ORNL/ER-155

**Phase 1 Data Summary Report for the Clinch River Remedial
Investigation: Health Risk and Ecological Risk
Screening Assessment**

ORNL/ER--155

DE93 006272

This page can be trimmed for use as a spine on a three-ring binder.

MASTER

JP

Energy Systems Environmental Restoration Program
Clinch River Environmental Restoration Program

**Phase 1 Data Summary Report for the Clinch River Remedial
Investigation: Health Risk and Ecological
Risk Screening Assessment**

R. B. Cook	S. K. Holladay
S. M. Adams	L. A. Hook
J. J. Beauchamp	D. A. Levine
M. S. Bevelhimer	R. C. Longman
B. G. Blaylock	C. W. McGinn
C. C. Brandt	J. L. Skiles
C. J. Ford	G. W. Suter
M. L. Frank	L. F. Williams
M. J. Gentry	

Date Issued—December 1992

Prepared by
Environmental Sciences Division
Oak Ridge National Laboratory
ESD Publication 4021

Prepared for
U.S. Department of Energy
Office of Environmental Restoration and Waste Management
under budget and reporting code EW 20

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6285
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400

Author Affiliations

R. B. Cook (Program Manager), S. M. Adams, B. G. Blaylock, C. J. Ford, M. L. Frank, S. K. Holladay, L. A. Hook, and G. W. Suter are members of the Environmental Sciences Division; J. J. Beauchamp and C. C. Brandt are members of the Engineering Physics and Mathematics Division; and C. W. McGinn and J. L. Skiles are members of the Health and Safety Research Division. All of the aforementioned divisions are part of Oak Ridge National Laboratory, Martin Marietta Energy Systems, Inc. M. S. Bevelhimer and D. A. Levine are with Automated Sciences Group, Inc., in Oak Ridge, Tennessee. M. J. Gentry, R. C. Longman, and L. F. Williams are with The University of Tennessee, Knoxville.

CONTENTS

FIGURES	v
TABLES	vii
ABBREVIATIONS	xiii
EXECUTIVE SUMMARY	xv
1. INTRODUCTION	1-1
1.1 OBJECTIVES AND APPROACH OF THE CLINCH RIVER REMEDIAL INVESTIGATION	1-3
1.2 SUMMARY OF THE SITE CONCEPTUAL MODEL	1-5
1.3 REPORT ORGANIZATION	1-6
2. PHASE 1 SAMPLE COLLECTION AND ANALYSIS	2-1
2.1 WATER	2-1
2.2 SEDIMENT	2-8
2.3 FISH	2-9
2.4 QA/QC PROGRAM FOR THE PHASE 1 DATA COLLECTION	2-12
2.4.1 General	2-12
2.4.2 Field Quality Control	2-17
2.4.3 Laboratory Quality Control Program	2-18
2.4.4 Data Validation	2-18
2.4.5 Data Assessment	2-19
3. SITE CHARACTERIZATION	3-1
3.1 INTRODUCTION	3-1
3.2 WATER	3-1
3.2.1 Discussion of Contaminants for the System	3-1
3.2.2 Discussion of Contaminants by Study Reach	3-2
3.2.3 Summary	3-3
3.3 SEDIMENT	3-4
3.3.1 Discussion of Contaminants for the System	3-5
3.3.2 Discussion of Contaminants by Study Reach	3-6
3.3.3 Summary	3-16
3.4 FISH ANALYSIS	3-18
3.4.1 Discussion of Contaminants for the System	3-19
3.4.2 Discussion of Contaminants by Study Reach	3-19
3.4.3 Summary	3-25
4. SCREENING RISK ASSESSMENT	4-1
4.1 SCREENING ASSESSMENT FOR HUMAN EXPOSURE	4-1
4.1.1 Approach	4-1
4.1.2 Exposure Pathways	4-1
4.1.3 Usage and Exposure Factors	4-2

4.1.4	Data Bases	4-2
4.1.5	Screening Criteria	4-3
4.1.6	Contaminant Screening Results for Primary Exposure Pathways	4-5
4.1.7	Dredging Scenario	4-16
4.1.8	Comparison of the Phase 1 Screening Analysis with a Previous Screening Analysis of Historical Data	4-19
4.1.9	Nondetected Contaminants	4-20
4.1.10	Contaminants Having No Available Toxicity Values	4-20
4.1.11	Special Case—Lead	4-21
4.1.12	High Priority Contaminants and Contaminants Requiring Further Investigation	4-21
4.1.13	Summary for Screening Risk Assessment	4-22
4.2	ECOLOGICAL RISK	4-24
4.2.1	Hazard Definition	4-25
4.2.2	Exposure Assessment	4-25
4.2.3	Effects Assessment: Screening Benchmarks	4-27
4.2.4	Risk Characterization: Contaminant Screening Results	4-28
4.2.5	Comparison to Previous Results	4-33
5.	SUMMARY	5-1
6.	REFERENCES	6-1
Appendix A.	LISTING OF ALL ANALYTES WITH CAS NUMBERS	A-1
Appendix B.	SUMMARY STATISTICS FOR WATER SAMPLES	B-1
Appendix C.	SUMMARY STATISTICS FOR SEDIMENT SAMPLES	C-1
Appendix D.	STATISTICAL PROCEDURE FOR PAIRWISE COMPARISON OF FISH TISSUE RESULTS	D-1
Appendix E.	HUMAN HEALTH RISK METHODS	E-1
Appendix F.	ECOLOGICAL RISK METHODS	F-1
Appendix G.	BRADEN BRANCH	G-1

FIGURES

1.1	Map showing the location of the Clinch River, Watts Bar Reservoir, and the Department of Energy's Oak Ridge Reservation	1-2
1.2	The phased approach used in the Clinch River Remedial Investigation	1-4
2.1	Locations of reaches used in Phase 1 sampling, analysis, and risk assessment	2-2
2.2	Location of sample sites for Phase 1 water and sediment core collection in Clinch River/Watts Bar system	2-5
2.3	Location of sites for Phase 1 surface sediment grab sample collection in Clinch River/Watts Bar system	2-10
2.4	Location of sample sites for Phase 1 fish collection in Clinch River/Watts Bar Reservoir system	2-13
3.1	Average ^{137}Cs concentration in surface sediment grabs within each reach ...	3-7
3.2	Average ^{60}Co concentration in surface sediment grabs within each reach ...	3-9
3.3	Average concentrations of arsenic and zinc for cores sampling sites	3-10
3.4	Average concentrations of beryllium and cadmium for cores sampling sites	3-11
3.5	Average concentrations of chromium, copper, and lead for cores sampling sites	3-12
3.6	Average concentrations of mercury for cores sampling sites	3-13
3.7	Average concentrations of ^{137}Cs for cores sampling sites	3-14
3.8	Average concentrations of ^{60}Co for core sampling sites	3-17
3.9	Arithmetic mean concentrations of total chlordane in Clinch River/Watts Bar fish tissue for all sample sites	3-21
3.10	Arithmetic mean concentrations of total PCBs in Clinch River/Watts Bar fish tissue for each site sampled in Phase 1	3-22
3.11	Arithmetic mean concentrations of mercury in Clinch River/Watts Bar fish tissue samples from each Phase 1 site	3-23

3.12	Arithmetic mean concentrations of ^{137}Cs in Clinch River/Watts Bar fish tissue sampled from each Phase 1 site	3-24
4.1	Screening criteria for carcinogens	4-4
4.2	Screening criteria for noncarcinogens	4-6
E.1	Location of NPDES monitoring sites in Clinch River/Watts Bar System	E-5
G.1	Braden Branch near-shore sample sites	G-4
G.2	Cobalt-60 concentrations at Braden Branch	G-10
G.3	Cobalt concentrations in Braden Branch core (No. 18700G)	G-11

TABLES

2.1	Stream and river reaches used in the CRRI site characterization and screening risk assessment	2-3
2.2	Matrices sampled in each reach during Phase 1 of the CRRI (validated and accepted data)	2-4
2.3	Summary of surface water analysis for the contaminants of concern	2-6
2.4	Number of CRRI water samples, by site and type of analysis (validated and accepted data)	2-7
2.5	Number of CR ERP sediment samples by reach and type of analysis (validated and accepted data)	2-11
2.6	Sample sites for Phase 1 fish collection in Clinch River/Watts Bar Reservoir system	2-14
2.7	Number of regular samples (i.e., does not include QC samples) submitted for analysis for each analytical group by species and by sampling location (validated and accepted data)	2-15
2.8	Data qualifiers	2-20
2.9	Summary of the number of rejected samples as a result of the validation of laboratory analyses	2-21
3.1	Summary of study site-to-reference site comparisons for fish tissue results	3-25
4.1	Summary of adult screening results for carcinogens in the fish ingestion pathway	4-7
4.2	Summary of screening results for noncarcinogens in the fish ingestion pathways	4-8
4.3	Summary of adult screening results for radionuclides in near-shore surface sediment exposure	4-9
4.4	Summary of adult screening results for carcinogens in the water ingestion pathway	4-10
4.5	Summary of screening results for noncarcinogens in the water ingestion pathway	4-12

4.6	Summary by reach of contaminants in the dredging scenario identified by conservative screening as requiring further investigation before taking action or designating as low priority	4-17
4.7	Contaminants for which the use of detection limits produced screening indices $>10^{-4}$ for carcinogens or >1.0 for noncarcinogens by nonconservative screening	4-21
4.8	Summary of contaminants requiring further investigation before taking action or designating as low priority based on fish ingestion and dredging scenarios	4-23
4.9	Contaminants of concern for ecological risk assessment	4-30
A.1	Listing of all analytes with CAS numbers and matrices analyzed for each analyte	A-3
B.1	Summary statistics for water data summarized for all reaches sorted by type of analysis and decreasing percentage detected	B-3
B.2	Summary statistics for water data detected compounds summarized by reach sorted by type of analysis, analysis, and reach	B-7
C.1	Summary statistics for sediment core data summarized for all reaches sorted by type of analysis and decreasing percentage detected	C-3
C.2	Summary statistics for sediment core data detected compounds summarized by reach sorted by type of analysis, analysis, and reach	C-6
C.3	Summary statistics for sediment grab data summarized for all reaches sorted by type of analysis and decreasing percentage detected	C-13
C.4	Summary statistics for sediment grab data detected compounds summarized by reach sorted by type of analysis, analysis, and reach	C-14
D.1	Summary statistics for fish tissue data summarized for all reaches sorted by type of analysis and decreasing percentage detected	D-6
D.2	Summary statistics for fish tissue data detected compounds summarized by subreach sorted by type of analysis, reach, and subreach	D-9
E.1	Receiving reaches for Oak Ridge Reservation source monitoring and surface water environmental monitoring stations	E-6
E.2	Usage and exposure factors for conservative and nonconservative screening	E-8

E.3	Reference dose factors and slope (cancer potency) factors for inorganic compounds	E-10
E.4	Reference dose factors and slope (cancer potency) factors for organic compounds	E-11
E.5	Slope (cancer potency) factors for radionuclides	E-15
E.6	Summary of carcinogens in the secondary pathways identified as low priority for further investigation	E-16
E.7	Summary of noncarcinogens in the secondary pathways identified as low priority for further investigation	E-19
E.8	Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-22
E.9	Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-36
E.10	Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-45
E.11	Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-60
E.12	Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-69
E.13	Results of adult conservative screening for carcinogens in dredged sediments where at least one measurement in a reach was above detection limits	E-78
E.14	Results of adult conservative screening for noncarcinogens in dredged sediments where at least one measurement in a reach was above detection limits	E-83
E.15	Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits	E-87

E.16	Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits	E-99
E.17	Results of adult nonconservative screening for carcinogens in dredged sediments where no measurements in a reach were above detection limits	E-113
E.18	Results of adult nonconservative screening for noncarcinogens in dredged sediments where no measurements in a reach were above detection limits	E-120
E.19	Unclassified analytes	E-128
F.1	Mean and upper confidence limit (UCL) concentrations of chemicals detected in the dissolved phase of CRRRI water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-6
F.2	Limits of detection of chemicals not detected in the dissolved phase of the CRRRI water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-8
F.3	Mean and upper confidence limit (UCL) concentrations of chemicals detected in NPDES water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-22
F.4	Limits of detection of chemicals not detected in NPDES water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-35
F.5	Mean and upper confidence limit (UCL) concentrations of chemicals detected in fish ($\mu\text{g/L}$), bioconcentration factors, estimated exposure in water ($\mu\text{g/L}$), and quotients for exposure to that water concentration	F-54
F.6	Limits of detection of chemicals not detected in fish ($\mu\text{g/L}$), bioconcentration factors, estimated exposure in water ($\mu\text{g/L}$), and quotients for exposure to that water concentration	F-66
F.7	Concentrations of inorganic chemicals detected in sediment ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-86
F.8	Detection limits for inorganic chemicals not detected in sediment ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-91

F.9	Mean and upper confidence limit (UCL) concentrations of organic chemicals detected in sediment ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-92
F.10	Detection limits for organic chemicals not detected in sediment ($\mu\text{g/L}$), estimated pore water concentrations ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-95
F.11	Mean and upper confidence limit (UCL) concentrations of chemicals in fish fillets (mg/kg), toxicological benchmarks for wildlife (mg/kg), and quotients for hazards to piscivorous wildlife	F-140
F.12	Minimum and upper confidence limit (UCL) detection limits for chemicals not detected in fish fillets (mg/kg), toxicological benchmarks for wildlife (mg/kg), and quotients for hazards to piscivorous wildlife	F-143
G.1	Data listing for CR-ERP Braden Branch sediment grab sample radiological analyses	G-6
G.2	Summary statistics for Braden Branch sediment core data	G-7
G.3	Data listing for CR-ERP Braden Branch sediment core samples radiological analyses	G-8
G.4	Summary statistics for Braden Branch sediment grab data	G-9
G.5	Results of adult nonconservative screening for carcinogens in surface sediment of Braden Branch where at least one measurement was above detection limits	G-13
G.6	Results of adult conservative screening for carcinogens in surface sediment of Braden Branch where at least one measurement was above detection limits	G-14

ABBREVIATIONS

AA	atomic absorption
BB	Braden Branch
BCF	bioconcentration factor
BMAP	Biological Monitoring and Abatement Program
CAS	Chemical Abstract Service
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CR	Clinch River
CR-ERP	Clinch River Environmental Restoration Program
CRM	Clinch River mile
CRRFI	Clinch River RCRA Facility Investigation
CRRI	Clinch River Remedial Investigation
CR/WBR	Clinch River/Watts Bar Reservoir
DDD	dichlorodiphenyldichloroethane
DDE	dichlorodiphenyldichloroethene
DDT	dichlorodiphenyltrichloroethane
DL	detection limit
DOE	U.S. Department of Energy
EFPC	East Fork Poplar Creek
EPA	U.S. Environmental Protection Agency
ER-L	effects range—low
ER-M	effects range—median
ESD	Environmental Sciences Division
FDA	U.S. Food and Drug Administration
FFA	Federal Facilities Agreement
foc	fractional organic matter content
HDPE	high-density polyethylene
HSWA	Hazardous and Solid Wastes Amendments
ICP	inductively coupled plasma emission spectroscopy
Koc	organic matter/water partitioning coefficient
Kp	partitioning coefficient
LCV	lowest chronic value
LTV	lowest toxic value
NAWQC	National Ambient Water Quality Criteria
NOAA	National Oceanic and Atmospheric Administration
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NWQC	National Water Quality Criteria
ORAU	Oak Ridge Associated Universities
ORNL	Oak Ridge National Laboratory
ORR	Oak Ridge Reservation
PCB	polychlorinated biphenyl
PCK	Poplar Creek kilometer
QA	quality assurance
QA/QC	quality assurance/quality control
RCRA	Resource Conservation and Recovery Act of 1976

RfD	reference dose
RI	remedial investigation
SARA	Superfund Amendments and Reauthorization Act of 1986
SI	screening indices
SQL	sample quantitation limit
TDEC	Tennessee Department of Environment and Conservation
TRM	Tennessee River mile
TVA	Tennessee Valley Authority
UCL	upper confidence limit
WBR	Watts Bar Reservoir
WOCE	White Oak Creek Embayment

EXECUTIVE SUMMARY

The Clinch River Remedial Investigation (CRRI) is designed to address the transport, fate, and distribution of waterborne contaminants released from the U.S. Department of Energy's (DOE's) Oak Ridge Reservation (ORR) and to assess potential risks to human health and the environment associated with these contaminants. The contaminants released since the early 1940s include a variety of radionuclides, metals, and organic compounds. Primary areas of investigation are Melton Hill Reservoir, the Clinch River from Melton Hill Dam to its confluence with the Tennessee River, Poplar Creek, and Watts Bar Reservoir. The receiving river-reservoir system encompasses 140 river miles in length and 44,000 acres in surface area and is used for municipal water supply, sport fishing, navigation, boating, swimming, tourism, and residential development. The contaminants identified in the Clinch River/Watts Bar Reservoir (CR/WBR) downstream of the ORR are those associated with the water, suspended particles, deposited sediments, aquatic organisms, and wildlife feeding on aquatic organisms.

A phased remedial investigation of the CR/WBR system is underway to (1) define the nature and extent of the off-site contamination, (2) evaluate associated environmental and human health risks, and (3) preliminarily identify and evaluate potential remediation alternatives. The purpose of this report is to summarize the results of Phase 1 of the CRRI.

Phase 1 of the CRRI was a preliminary sampling and analysis of fish, sediment, and water in selected areas of the CR/WBR chosen to represent differing levels of media contamination. Specifically, Phase 1 was designed to (1) obtain high-quality data to confirm existing historical data for contaminant levels in fish, sediment, and water from the CR/WBR; (2) determine the range of contaminant concentrations present in the river-reservoir system; (3) identify specific contaminants of concern; and (4) establish the reference (background) concentrations for those contaminants. Previously, scoping studies were conducted to provide an initial determination of contaminant distribution, in the sediments of Watts Bar Reservoir (Olsen et al. 1992) and to preliminarily assess potential human health and ecological risk (Suter 1991; Hoffman et al. 1991).

For Phase 1 of the CRRI, the CR/WBR system was divided into ten reaches, six of which are potentially affected by releases from the ORR. The other four reaches serve as reference areas. Contaminant concentrations for fish, sediment, and water were summarized for each of the 10 reaches. Analytes detected at least once in a reach were evaluated separately from those never measured above their analytical detection limit (i.e., nondetected).

In water samples collected during Phase 1, inorganic and organic constituents with the potential to pose a risk to human health or the environment were within the ranges reported previously in monitoring reports (e.g., Kornegay et al. 1991). Each inorganic contaminant detected during the Phase 1 investigation is well below the water quality criteria promulgated by the Tennessee Department of Environment and Conservation (rules of the Tennessee Department of Environment and Conservation, Chapter 1200-4-3).

The majority of potential contaminants of concern in Phase 1 water samples were detected between White Oak Creek and Poplar Creek on the Clinch River (reach 2) and in

the Poplar Creek downstream of the confluence with East Fork Poplar Creek, which receive effluents from several ORR sources for these contaminants. Total copper and zinc were detected in reach 2 surface waters and had values in the range of those previously observed (Kornegay et al. 1991). Dissolved lead, observed in water samples from Poplar Creek upstream of the confluence with East Fork Poplar Creek, may originate from an upstream source.

The Oak Ridge Y-12 Plant, the city of Oak Ridge wastewater treatment facility, and Oak Ridge K-25 Site operations are among the possible sources for mercury, silver, and lead detected in water samples from Poplar Creek downstream of the confluence with East Fork Poplar Creek. Elevated levels of cadmium and lead in water samples from the Clinch River between Poplar Creek and the confluence with the Tennessee River may originate from several sources in and upstream from this reach.

Concentrations of all radiological constituents in Phase 1 water samples were within the ranges of those observed during environmental monitoring (Kornegay et al. 1991) and with previous scoping investigations (Olsen et al. 1992). Dissolved ^{137}Cs and particle-associated ^{137}Cs were detected in the Clinch River arm of Watts Bar Reservoir and Watts Bar Reservoir downstream of the confluence with the Clinch River, with values similar to those reported previously for these reaches. Dissolved ^{137}Cs was detected in Poplar Creek downstream of the confluence with East Fork Poplar Creek, possibly a result of backflow from the Clinch River. Cobalt-60 in Watts Bar Reservoir below the entry of the Clinch was detected at levels comparable with those reported previously. Average ^{90}Sr values for all reaches never exceeded drinking water standards. Each of these radionuclides is currently released in permitted discharges from the ORR to surface waters.

Data resulting from the measurement of inorganic, organic, and radiological contaminants from the Phase 1 water sampling, in combination with historical data, indicate that a focused characterization effort in Phase 2 is warranted for Poplar Creek downstream of the confluence with East Fork Poplar Creek and for the Clinch River arm of Watts Bar Reservoir. Additionally, a more thorough characterization of reference sites is needed to rigorously establish background concentrations in surface water.

Concentrations of ^{137}Cs in surface sediments (0 to 10 cm) in near-shore areas of Norris, Melton Hill Reservoirs, and the Tennessee River arm of Watts Bar Reservoir should represent background levels (<1.0 pCi/g) because they are upstream of ORR and only received fallout from atmospheric testing of nuclear weapons. Concentrations in Poplar Creek and the Emory River average 1 pCi/g, which is consistent with background concentrations. Cesium-137 concentrations in the near-shore surface sediments in the Clinch River arm of Watts Bar Reservoir are between 2 and 3 pCi/g. Concentrations in Watts Bar Reservoir below the confluence of the Tennessee and Clinch rivers are usually less than 1 pCi/g, which are equivalent to background concentrations. Three areas within the Clinch River arm of Watts Bar Reservoir had concentrations between 10 and 34 pCi/g in surface sediment and will be targeted for sampling and analysis during Phase 2 of the CRRI. Samples collected at these areas will be analyzed for the contaminants identified by the risk screening as requiring further research.

Patterns of inorganic contaminants in sediment core samples indicate that there are multiple sources for these contaminants. Arsenic, beryllium, cadmium, chromium, copper, and mercury in Poplar Creek and the downstream portion of the Clinch River arm of Watts Bar Reservoir appear to be originating, at least in part, from Poplar Creek. Mercury releases from the Oak Ridge Y-12 Plant into the Poplar Creek system are well documented (Olsen et al. 1992, Turner et al. 1985). The other metals may be derived from disposal of coal fly-ash material and photographic processing chemicals from the Oak Ridge K-25 Site or from the municipal wastewater treatment facility for the city of Oak Ridge. Concentrations of cadmium, lead, and zinc are higher in the sediments in Watts Bar Reservoir below the confluence of the Tennessee River and Clinch River than elsewhere in the system, suggesting that there are sources of these contaminants to the Tennessee River arm of the reservoir or to WBR below the confluence of the two rivers. Phase 2 sampling and analysis will focus on locating any areas where concentrations of these contaminants pose an elevated risk to human health and the environment.

The lack of detectable quantities in sediment samples of most of the 92 organic chemicals for which analyses were conducted allows significant reduction in the list of organic contaminants requiring further study. Flouranthene, phenanthrene, and pyrene were found in Poplar Creek downstream of the confluence with East Fork Poplar Creek and in the Clinch River arm of Watts Bar Reservoir. Aroclor 1254 was found only in sediments at Clinch River Mile 9.5, which is immediately downstream of Poplar Creek. Potential sources for these contaminants in sediment are ORNL, the Y-12 Plant, the K-25 Site, and the municipal wastewater treatment facility. Sampling and analysis for these contaminants in Poplar Creek downstream of the confluence with East Fork Poplar Creek and in the Clinch River arm of Watts Bar Reservoir will continue in Phase 2 with the objective of identifying and delineating any areas requiring remediation.

Alpha-emitting radionuclides were detected in every site sampled for sediments. However, concentrations were all at or near background concentrations, except in Poplar Creek downstream of the confluence with East Fork Poplar Creek. Concentrations of ^{234}U , ^{235}U , and ^{238}U in Poplar Creek sediments were above background levels. Poplar Creek downstream of the confluence with East Fork Poplar Creek will be the focus of study for these contaminants during Phase 2 activities.

In order to characterize the nature and extent of contamination in the biota of the CR/WBR system, 126 bluegill sunfish, 119 channel catfish, and 43 largemouth bass were sampled from 15 sites in 8 reaches. These fish species were chosen for sampling by virtue of their positions in the foodweb, their migration/mobility behavior, and relative availability. Of the 77 organic and 13 inorganic analytes, a majority (63 organics and 7 inorganics) were not detected in fish at any site. Of the radionuclides, only ^{137}Cs , ^{60}Co , and ^{90}Sr were detected in fish. Four analytes were rarely detected, occurring in less than 5% of the samples. Seventeen analytes were detected in more than 5% of the samples, with only three of those (^{90}Sr , mercury, and zinc) detectable in all samples.

In general, the Phase 1 maxima (individual maximum concentration of an analyte at each site) for fish are lower than those from previous sampling. Exceptions to this generality are mercury, selenium, and Aroclor 1260. Maximum values of mercury in fish collected in Phase 1 were higher than historical values only in lower Watts Bar Reservoir, and at no sites did the

maximum mercury concentration exceed the Food and Drug Administration's (FDA's) action level of 1.0 mg/kg. Selenium maxima in fish exceeded historical values in Poplar Creek downstream of the confluence with East Fork Poplar Creek and in the Clinch River upstream of Poplar Creek and below Melton Hill Reservoir. Aroclor 1260 (a PCB congener) maxima exceeded historical values in Poplar Creek downstream of the confluence with East Fork Poplar Creek and in the Clinch River below Poplar Creek. Average values of Aroclor 1260 in fish did not exceed 1.0 mg/kg at any site or within any reach, but Aroclor 1260 maxima in fish exceeded the FDA action level of 2.0 mg/kg at one site in Melton Hill Reservoir, one site in Poplar Creek, and three sites in the Clinch River arm of WBR.

Mean concentrations of some contaminants in fish at some CR/WBR sites were higher than the concentrations at the reference site (Norris Reservoir). Arsenic concentrations were significantly ($P < 0.05$) higher than concentrations at the reference site at only one location (CRM 0.5). In Poplar Creek below the confluence with East Fork Poplar Creek and at the first site downstream of the confluence of Poplar Creek and the Clinch River (Brashear Island), mercury concentrations in fish were predictably higher than those of the reference sites. Selenium concentrations in fish were higher than those in the reference site in main channel sites in the upper part of the system (Upper Melton Hill to Brashear Island). Zinc concentrations in fish at 5 of the 12 sites were higher than those at the reference site, but no obvious spatial distribution was evident. Chlordane concentrations in fish from Upper Melton Hill to the mouth of the Clinch River were higher than reference concentrations. Total PCB concentrations at all CR/WBR sites were higher than those in fish from the reference site (Norris Reservoir). Cesium-137 levels for all fish species were usually higher than the reference, particularly below the confluence of White Oak Creek and the Clinch River. Concentrations of ^{137}Cs and mercury in fish are both elevated at sites downstream of known sources of these contaminants, White Oak Creek and East Fork Poplar Creek, respectively. No such patterns of downstream gradients of chlordane or PCBs in fish are evident.

A screening assessment of ecological effects in the CR/WBR was conducted concurrently with the health screening assessment. The ecological screening assessment considered potential toxic effects on aquatic biota due to exposure to contaminants in water, potential toxic effects to benthic and epibenthic organisms due to contaminants in sediment, and risks to piscivorous wildlife due to ingestion of contaminants in fish flesh. The assessment was performed by comparing chemical concentrations in media to screening benchmarks that are either regulatory criteria or concentrations that are estimated to have minimal effects on the abundance or production of exposed populations. The screening ecological risk assessment for Phase 1 identified 14 contaminants of concern among the chemicals that were analyzed for and detected in water, sediment, or fish fillets. In addition, 19 chemicals that were not detected were identified as potential contaminants of concern because their limits of detection exceed screening benchmarks.

Comparison of media concentrations to toxicological benchmarks produced ambiguous results because of the large number of chemicals that were not detected but had limits of detection higher than potentially toxic concentrations. However, the number of potential contaminants of concern is much lower than in previous screening assessments because of generally lower contaminant concentrations than in the historical data, use of bioavailable (dissolved) concentrations of metals in water rather than total concentrations, and use of better screening benchmarks for metals in sediment.

A screening analysis using human health risk methodology similar to the approach used in Hoffman et al. (1991) to screen historical data was conducted with data collected during Phase 1 of the CRRI. The purpose of this screening exercise, along with the results obtained in Hoffman et al. (1991), is to help guide the development of the Phase 2 sampling plan, which will focus the collection of data on contaminants of concern. A future baseline risk assessment will be completed when data from the Phase 2 sampling are available.

Conservatively biased risk estimates were used to identify contaminants, exposure pathways, and river and reservoir reaches that have a low priority for further investigations. It is highly unlikely that these conservative estimates will underestimate the potential maximum exposure to an individual residing in the vicinity of the Clinch River where it receives releases of effluents from the ORR. Nonconservative estimates of risk were used to identify contaminants, pathways, and reaches that have a high priority for further investigation. These nonconservative estimates are more realistic than the conservatively biased estimates and should not substantially overestimate maximum exposures to local populations or subgroups.

The primary pathways examined were exposure to three media: fish, near-shore surface sediments, and surface water. Currently, there is not a realistic direct exposure pathway to deep sediment, where the highest concentrations of particle-reactive contaminants have been observed (Olsen et al. 1992). However, a dredging scenario was included in the screening analysis to evaluate potential exposure pathways if the sediment should be dredged and placed up on the shore. In addition to the direct exposure pathways from exposed sediment (ingestion and external exposure to radionuclides) included in the dredging scenario, secondary pathways were also considered (ingestion of meat, milk, and vegetables produced on dredged sediments). The dredging scenario depended on mathematical models to determine the concentration of contaminants in the agricultural products. Because the models are designed to increase the conservatism of the results, the dredging scenario was used only to identify contaminants of low priority for further investigation.

Nonconservative screening of data for the fish ingestion pathway identified Aroclor 1260, a carcinogenic PCB, as a high-priority contaminant requiring immediate consideration for further investigation and potential remedial action for all reaches except reference reach 10 (Norris Reservoir). This contaminant was identified previously in Hoffman et al. (1991) as a high-priority contaminant. Other investigations have shown that PCBs contaminate most eastern Tennessee rivers and lakes, and although ORR is a probable source, it is not the only source of PCBs for the Clinch River and Tennessee River system. The Tennessee Department of Environment and Conservation has issued fish consumption advisories for several eastern Tennessee reservoirs (e.g., Fort Loudoun, Watts Bar, Melton Hill) as a result of PCB contamination.

None of the noncarcinogens that were analyzed in fish were identified as high priority contaminants. Conservative screening indicated that radionuclides (^{60}Co , ^{137}Cs , and ^{90}Sr) found in fish sampled during Phase 1 could be assigned a low priority for future consideration in all reaches except for ^{137}Cs in the Clinch River arm of Watts Bar Reservoir. Conservative screening identified several carcinogens and noncarcinogens that had been identified previously (Hoffman et al. 1991) as requiring further investigation in fish (i.e., Aroclor 1254, chlordane, arsenic, beryllium, mercury, and selenium).

Nonconservative screening of the ^{137}Cs and ^{60}Co data from surface-sediment samples from near-shore areas did not identify these radionuclides as high priority contaminants. Conservative screening identified ^{60}Co and ^{137}Cs , the only contaminants for which the samples were analyzed, as low priority in all reaches for the ingestion and inhalation pathways. In the external exposure pathway, ^{137}Cs was identified in Poplar Creek downstream of the mouth of East Fork Poplar Creek and in Melton Hill and Watts Bar reservoirs, and ^{60}Co in Poplar Creek and the Clinch River arm of Watts Bar Reservoir as contaminants requiring further investigation. However, Melton Hill Reservoir and the Tennessee River arm of WBR do not receive effluents from the ORR; therefore, the levels of ^{137}Cs found in the surface sediments of these reaches are background levels. A known source of ^{60}Co exists on the Clinch River upstream of the ORR, and it probably influences the screening indices for ^{60}Co in the Clinch River (Appendix G).

The results of the screening analysis for the water ingestion pathway should be viewed with caution because of the limited numbers of samples and because source monitoring data were used as estimates of concentrations in the reservoir. Thorium-228 was tentatively designated as a high priority contaminant in Poplar Creek downstream of the confluence with East Fork Poplar Creek in the water ingestion pathway. However, this designation results from including data from source monitoring sites. The monitoring data should be evaluated further before ^{228}Th , which is a daughter of ^{232}U and naturally occurring ^{232}Th , is designated a high priority contaminant. Screening of the water ingestion pathway showed that Poplar Creek, which receives effluents from the K-25 Site and from the Y-12 Plant and the city of Oak Ridge west-end sewage treatment plant via East Fork Poplar Creek, has the greatest number of contaminants that require further investigation. In the other reaches, most of the contaminant values were below detection limits or are considered low priority for further investigation.

In the dredging scenario, conservative screening was conducted on two direct pathways—external exposure and ingestion—and three secondary pathways—ingestion of meat, milk, and vegetables. Conservative screening of the secondary pathways identified many organic and inorganic contaminants in all reaches that could be assigned low priority for further investigation. Except for ^{137}Cs and to a lesser extent ^{60}Co and ^{90}Sr , most radionuclides for which data were available were assigned low priority for the secondary pathways in all reaches.

Nonconservative screening was conducted for contaminants that had concentrations below analytical detection limits to identify contaminants for which either better analytical methodology is needed or for which source-term data are needed.

Results from Phase 1 of the CRRI indicate that Poplar Creek and the Clinch River arm of Watts Bar Reservoir are the reaches of greatest concern based on the number of contaminants that need further evaluation during Phase 2 of the remedial investigation.

The screening assessment of risk to the environment for Phase 1 indicates that cadmium, mercury and silver in water need to be further investigated to determine the risk to aquatic biota in the CR/WBR system. Arsenic, lead, mercury, and zinc in sediments of the CR/WBR system, which may pose a risk to benthic dwelling organisms, need to be further assessed

during Phase 2. More detailed sampling, analysis, and assessment are needed during Phase 2 of the CRRI to conduct the baseline assessment for ecological risk.

For assessment of the risk to human health, the Phase 1 fish data are the most complete. The Phase 2 sampling for fish should concentrate on providing statistically valid data for a future baseline assessment of human health risk. Additional samples of water need to be collected and analyzed in surface water areas proximate to the ORR outfalls and sources to provide data for the baseline assessment of human health risk. Organic and inorganic contaminants common to eastern Tennessee and other metropolitan areas should be identified through further analysis of the available data for downstream and reference reaches. Analysis of organic contaminants in water and sediment should concentrate on those contaminants that are routinely released or have been released in the past from the ORR. The Phase 2 sampling and analysis should concentrate on previously identified contaminants in Poplar Creek downstream of the mouth of East Fork Poplar Creek and in the Clinch River arm of Watts Bar Reservoir to provide statistically valid data for a baseline risk assessment.

REFERENCES FOR EXECUTIVE SUMMARY

Hoffman, F. O., B. G. Blaylock, M. L. Frank, L. A. Hook, E. L. Etnier, and S. S. Talmage. 1991. *Preliminary Screening of Contaminants in the Off-Site Surface Water Environment Downstream of the U.S. Department of Energy Oak Ridge Reservation*. ORNL/ER-9. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Kornegay, F. C., D. C. West, R. A. Evans, S. T. Goodpasture, M. F. Tardiff, and A. R. Wilson. 1991. *Oak Ridge Reservation Environmental Report for 1990. Volume 2: Data Presentation*. ES/ESH-18/V2. Martin Marietta Energy Systems, Inc., Oak Ridge, Tennessee.

Olsen, C. R., I. L. Larsen, P. D. Lowry, C. R. Moriones, C. J. Ford, K. C. Dearstone, R. R. Turner, B. L. Kimmel, and C. C. Brandt. 1992. *Transport and Accumulation of Cesium-137 and Mercury in the Clinch River and Watts Bar Reservoir System*. ORNL/ER-7. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Turner, R. R., C. R. Olsen, and W. J. Wilcox, Jr. 1985. Environmental fate of Hg and ¹³⁷Cs discharged from Oak Ridge facilities. pp. 329-338. In D. D. Hemphill (ed.), *Trace Substances in Environmental Health—XVIII*. University of Missouri, Columbia.

1. INTRODUCTION

The purpose of this report is to present and discuss the results of Phase 1 of the Department of Energy's (DOE) remedial investigation (RI) of the Clinch River/Watts Bar Reservoir (CR/WBR) system. A screening risk assessment for human and ecological risk is used to identify contaminants, environmental pathways, and sites in the river and reservoir that require further investigation during Phase 2 of the RI.

The Clinch River RI (CRRI) focuses on the portions of the Clinch and Tennessee rivers that may have been adversely affected by contaminants that were released from DOE's Oak Ridge Reservation (ORR) from the mid-1940s to present. The contaminants released from the ORR to the Clinch and Tennessee rivers originated from research, industrial, and waste disposal activities conducted on the ORR at Oak Ridge National Laboratory (ORNL), the Oak Ridge Gaseous Diffusion Plant (now known as the K-25 Site), and the Y-12 Plant. The contaminants released from these facilities into the Clinch River include a variety of radionuclides, metals, and organic compounds.

Melton Hill Reservoir and the Clinch River, downstream from Melton Hill Dam, form the southern and eastern boundaries of the ORR (Fig. 1.1). All surface water draining the ORR eventually enters the Clinch River system. The Clinch River flows into the Tennessee River system of multiple-purpose impoundments near Kingston, Tennessee, 34 km downstream from the ORR (Fig. 1.1). Watts Bar Reservoir extends from Watts Bar Dam at Tennessee River Mile (TRM) 529.9 to Fort Loudoun Dam at TRM 602.3, and up the Clinch River to Melton Hill Dam at Clinch River Mile (CRM) 23.1. Chickamauga Reservoir is the next mainstream impoundment downstream of Watts Bar Dam. Watts Bar Reservoir is large, encompassing 140 river miles in length and 44,000 acres in surface area, and is used for municipal water supply, sport fishing, boating, swimming, tourism, and residential development. The CR/WBR system receives effluents from DOE facilities on the ORR, as well as effluents from municipal and industrial water treatment facilities and runoff from agricultural and urban areas.

DOE's ORR was added to the National Priority List (NPL) in December 1989. A Federal Facilities Agreement (FFA) [under Sect. 120 of the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), as amended by the Superfund Amendment and Reauthorization Act (SARA) and Sect. 6001 of the Resource Conservation and Recovery Act (RCRA) of 1976] between DOE, the U.S. Environmental Protection Agency (EPA) Region IV, and the Tennessee Department of Environment and Conservation (TDEC) became effective on January 1, 1992. These parties intend to coordinate DOE's CERCLA/RCRA response obligation with corrective measures required and conducted by DOE under its current permit under the Hazardous and Solid Waste Amendments (HSWA) to RCRA of 1984. Section 3004(v) of the HSWA of RCRA specifically addresses requirements for corrective actions for releases of hazardous wastes or constituents beyond the boundaries of RCRA-permitted sites. In addition TDEC, DOE, and EPA expect that the response actions under the FFA, together with corrective measures, will achieve comprehensive remediation of releases and potential releases of contaminants from the ORR. The CRRI is being conducted in compliance with the FFA and with Sect. 3004(v) of RCRA.

Fig. 1.1. Map showing the location of the Clinch

Clinch River Watts Bar Reservoir

Environmental Restoration Program

Clinch River study area is
shown by river boundary.

Department of Energy Reservation
shown by shaded area

Tennessee State Plane
Datum NAD83
Units Meters
November 1992

iver, Watts Bar Reservoir, and DOE's Oak Ridge Reservation.

1.1 OBJECTIVES AND APPROACH OF THE CLINCH RIVER REMEDIAL INVESTIGATION

The primary objectives of the CRRRI (Martin Marietta Energy Systems 1990) are to

- define the nature and extent of contamination of the off-site surface water and sediment environment downstream from the ORR, which resulted from operations and waste disposal activities at DOE's facilities on the ORR;
- quantify any risk to human health and to the environment resulting from the existing off-site contamination; and
- preliminarily identify and evaluate potential corrective measures and remedial action alternatives.

A phased approach is being employed in the CRRRI because a large number of samples potentially may be required to characterize such a large and complex river and reservoir system. The phased approach relies heavily on iterative screening-level risk analysis for estimating human and ecological health and is based on obtaining the data needed for risk analyses and for evaluating remediation alternatives (Fig. 1.2). Primary components of the phased approach include scoping, Phase 1, and Phase 2 studies.

The scoping studies were performed before the work plan for the CRRRI was developed and approved. In the scoping studies, extant data were synthesized and used in a screening-level risk analysis, which served to (1) preliminarily identify contaminants of concern in the CR/WBR system and (2) identify areas of the river and reservoir in which additional data collection was necessary for site characterization, risk assessment, and evaluation of remediation alternatives. The scoping studies were summarized in Olsen et al. (1992), Suter (1992), and Hoffman et al. (1991).

The Phase 1 study followed the scoping study and consisted of a preliminary sampling and analysis in areas selected to represent differing levels of contamination (including reference sites located upstream from the ORR). The objectives of the Phase 1 study were to (1) obtain high-quality data to confirm and modify preliminary results from the scoping study, (2) determine the range of contaminant concentrations present in the off-site environment, (3) identify the contaminants of concern, and (4) establish the reference concentrations for those contaminants. The Phase 1 study has been completed and is summarized in this report.

The objectives of the Phase 2 study are to (1) perform iterative risk analyses as additional data become available, (2) quantitatively estimate potential risks to human health and the environment, and (3) focus additional sampling efforts. These risk analyses will guide more extensive sampling and analysis and risk assessment, ultimately leading to a baseline risk assessment presented in a final RI report. In addition, the Phase 2 study will include a preliminary evaluation of potential remediation alternatives and identification of effective and acceptable corrective measures. The Phase 2 study is presently in the planning stages.

After the Phase 1 sampling and analysis began, a task was initiated to characterize the near-shore sediment in the CR/WBR. The primary objective of the near-shore sediment task is to provide information on the risks posed to human health from surface sediments (between 0 and 10 cm depth in the sediments) in shallow waters of the near-shore areas.

Fig. 1.2. The phased approach used in the Clinch River Remedial Investigation.

These sediments are exposed to the air during the winter season, when the water levels in these reservoirs are lowered as part of the water management activities of the Tennessee Valley Authority (TVA). This task supports the efforts of the TVA and the U.S. Army Corps of Engineers to evaluate permit applications for sediment-disturbing activities (e.g., dredging for private boat access channels, installation of boat docks, shoreline stabilization). An interagency working group composed of the TVA, the Corps of Engineers, DOE, TDEC, EPA, and staff from the CRRI preliminarily evaluates the levels of contamination prior to TVA and the Corps performing their evaluation. Almost 600 surface sediment samples in near-shore areas of Watts Bar Reservoir and the Clinch River were collected as part of this task and are discussed in Sects. 3 and 4.

1.2 SUMMARY OF THE SITE CONCEPTUAL MODEL

A site conceptual model was formulated in Martin Marietta Energy Systems (1990) to identify and summarize the current knowledge and understanding of contaminant sources, the potential environmental pathways of contaminant exposure (including transport media and exposure routes), and the potential receptors for contaminants that may have been released from the ORR to downstream surface waters.

A complicating factor in identifying sources and media for the CRRI site conceptual model is that there also may be non-DOE sources of contaminants to the CR/WBR. The Clinch and Tennessee rivers receive inputs from municipalities, agricultural activities, and industries that may contribute contaminants to the media, pathways, and endpoints of concern.

Operations and waste disposal activities at the Y-12 Plant, ORNL, and the K-25 Site on the ORR have been sources of a variety of airborne, liquid, and solid contaminants. Some of these contaminants have affected local streams that ultimately drain into the CR/WBR. For the CRRI, the contaminants identified in the CR/WBR were transported by surface water from the point of release on the ORR to the location being investigated. Therefore, the media of concern in the CRRI are those associated with the surface water environment downstream from the ORR: the water itself, suspended particles and sediments, deposited sediments, aquatic organisms, and wildlife feeding on aquatic organisms.

The deposited sediments also serve as a secondary source of contaminants. For many contaminants found in deposited sediments (i.e., particle-reactive contaminants), the largest releases occurred 20 to 40 years ago (Martin Marietta Energy Systems 1990). These contaminated sediments were buried by sediments that contain much lower levels of contaminants (Turner et al. 1985; Olsen et al. 1992). Unusual storm events or dredging activities may resuspend these deeply buried contaminated sediments and reintroduce contaminants into the water column or into the aquatic food chain.

In the CR/WBR system, the highest potential exposure to contaminants released from the ORR occurs at several spatial scales. The highest potential exposure to dissolved contaminants in off-site surface waters should occur at the ORR boundary, where surface water flows from the reservation to the public domain. Exposure points with lower predicted chemical concentrations, but larger potentially exposed receptor populations, are locations of municipal water supply intakes. Points of high potential exposure to particle-associated

contaminants occur at points of sediment deposition and accumulation. A major effort of the planned Phase 2 study in the CRRI is to locate areas of high rates of sediment and contaminant accumulation.

The primary exposure routes for the environment (aquatic biota and wildlife feeding on aquatic biota) are uptake and metabolism of dissolved contaminants, direct contact or exposure, absorption, and ingestion of contaminated food organisms or particulate matter. The primary exposure routes of greatest potential risk to human health are ingestion of contaminated water, ingestion of contaminated food (e.g., fish, waterfowl, or vegetables grown on contaminated sediment or soil), direct ingestion of contaminated sediment or soil, and direct exposure to contaminated sediment or soil.

The results of the scoping studies conducted by the CRRI confirm this conceptual site model and further indicate that ingestion of contaminated biota and direct exposure to contaminated sediments are the greatest potential exposure pathways.

1.3 REPORT ORGANIZATION

The CRRI Phase 1 Data Summary Report is organized to provide the flow of information from the objectives and phased approach of the CRRI, to the description of Phase 1 data, and to discussion of the screening risk assessment.

In the first volume, Sect. 1 introduces the objectives and approach of the study and provides the regulatory background. Section 2 presents (1) a description of the study area and sampling sites for fish, sediment, and water; (2) methods used for sample collection and analysis; and (3) a brief description of the quality assurance program for Phase 1.

The contaminant concentrations in fish, sediment, and water, based on the Phase 1 data, is summarized in Sect. 3. The risk screening analyses are presented in Sect. 4 for human and ecological health. Section 5 summarizes the Phase 1 results and describes how these results lead to program activities in Phase 2. Supporting information about Phase 1, including summary tables of the analytical data, are collected in Appendixes A through G.

2. PHASE 1 SAMPLE COLLECTION AND ANALYSIS

This section briefly summarizes the rationale and methods for the collection and analysis of water, sediment, and fish samples for Phase 1 of the CRRI. A complete description of the rationale and methods may be found in the Clinch River RCRA Facility Investigation Work Plan (Martin Marietta Energy Systems 1990).

The primary areas of investigation during Phase 1 of the CRRI were the Melton Hill and Watts Bar Reservoirs, with particular emphasis on the Clinch River arm of Watts Bar Reservoir (Melton Hill to the Clinch River—Tennessee River confluence) and Poplar Creek. Phase 1 sampling from these areas occurred from December 1989 to February 1990 for water, October 1989 to April 1990 for fish, and December 1989 to July 1990 for sediment. For the purposes of sampling and analysis, the CR/WBR system was partitioned into reaches (Fig. 2.1, Table 2.1), following those identified by Hoffman et al. (1991). The number of samples collected in each reach for each medium type (water, fish, and sediment) is listed in Table 2.2 and in the section corresponding to that medium. A list of analytes for each medium, with cross-reference to its Chemical Abstract Service (CAS) number, is found in Appendix A.

2.1 WATER

Surface water is the primary transport pathway for contaminants leaving the reservation. It is also the principal exposure medium for the initial introduction of dissolved (non-particle-associated) contaminants into the food web. The complex hydrodynamics and spatial heterogeneity of the CR/WBR system make it difficult to adequately estimate the temporal and spatial extent of surface water contamination, even with a large number of samples (Martin Marietta Energy Systems 1990). The monitoring of NPDES permitted discharges from the ORR for NPDES and DOE Order 5400.1 provides considerable additional information on contaminant concentrations near discharge points.

Phase 1 sampling of water was undertaken to confirm suspected low dissolved concentrations of contaminants that were identified in scoping studies (Suter 1991, Hoffman et al. 1991, Olsen et al. 1992). Accordingly, characterization of the nature and extent of surface water contamination in the CR/WBR system during Phase 1 of the CRRI was limited to a single sampling event for each station (Fig. 2.2). The focus of the first phase of CRRI sample collection was sediment and biota.

Water samples were analyzed for inorganic, radiochemical and organic constituents, as listed in Table 2.3. Field chain-of-custody was established for these samples at the time of collection and maintained throughout processing, handling, transport, and transfer to other analytical laboratories. All sample collection devices were cleaned following appropriate procedures identified in Martin Marietta Energy Systems 1990. Water samples were collected from Norris Reservoir; upper Melton Hill Reservoir; Solway; lower Melton Hill Reservoir; Jones Island; the mouth of Grassy Creek mbayment; Poplar Creek at the Hartland Bridge; four sites within Poplar Creek below the mouth of East Fork; Brashear Island; Kingston City park; and upper, mid and lower Watts Bar Reservoir (Table 2.4). Additional samples for contaminant screening from public water intake structures and beaches in the CR/WBR system were collected by the TVA in 1991 (TVA 1991). Samples were collected using clean

Fig. 2.1. Locations of reaches used

STUDY REACHES

- 1 Clinch River
from Melton Hill Dam
to Norris Dam
- 2 Clinch River
from Poplar Creek
to Melton Hill Dam
- 3 Poplar Creek
from East Fork Poplar
to Clinch River
- 4 Clinch River
from Tennessee River
to Poplar Creek
- 5 Tennessee River
from Watts Bar Dam
to Clinch River
- 6 Emory River
- 13 Poplar Creek
above confluence
with East Fork Poplar
- 15 Chickamauga Reservoir
- 18 Tennessee River
from Clinch River
to Loudon Dam

10 KM

Tennessee State Plane

Datum NAD83

Units Meters

November 1992

**Table 2.1. Stream and river reaches used in the
CRRRI site characterization and screening risk assessment**

Reach number	Reach name (description)	River mile/km^a
1	Melton Hill Reservoir (from the Oak Ridge Marina to Melton Hill Dam)	CRM 52.0-23.1
2	Clinch (Clinch River from Melton Hill Dam to Poplar Creek)	CRM 23.0-12.1
3	Poplar Creek embayment (Poplar Creek below the confluence of the East Fork)	PCK 8.9-0.0
4	Poplar Creek Clinch (the Clinch River from the mouth of Poplar Creek to the confluence with the Tennessee River)	CRM 12.0-0.0
5	Watts Bar Reservoir (the Tennessee River from the confluence of the Clinch River to Watts Bar Dam)	TRM 567.5-530.0
6	Emory River	ERM 14.0-0.0
10	Norris Reservoir (the Clinch and Powell Rivers above Norris Dam)	
13	Poplar Creek (Poplar Creek above the confluence of the East Fork)	PCK 11.0-9.0
15	Chickamauga Reservoir (below Watts Bar Dam)	TRM 529.9-515.0
18	Tennessee River arm of WBR (from Fort Loudon and Tellico Dams to the confluence of the Clinch River)	TRM 602.0-567.6

^aCRM = Clinch River mile, PCK = Poplar Creek kilometer, TRM = Tennessee River mile, and ERM = Emory River mile.

Table 2.2. Matrices sampled in each reach during Phase 1 of the CRRI (validated and accepted data). Entries with an "X" indicate that a particular reach was sampled for that matrix. Blanks indicate that no sample was collected from the reach for the particular matrix.

Reach	Matrix			
	Fish	Sediment core	Surface grab sediment	Water
1: Melton Hill Reservoir	X	X	X	X
2: Clinch R.: Poplar Ck -> Melton Hill Dam	X	X	X	X
3: Poplar Ck -> East Fork Confluence	X	X	X	X
4: Clinch R.: Mouth -> Poplar Ck Confluence	X	X	X	X
5: Watts Bar Reservoir	X	X	X	X
6: Emory River			X	
10: Norris Reservoir	X	X	X	X
13: Poplar Creek Above East Fork Confluence	X			X
15: Chickamauga Reservoir	X			
18: Tennessee R: Clinch R. Confl -> Ft. Loudon			X	
Total number of reaches sampled	8	6	8	7

Table 2.3. Summary of surface water analysis for the contaminants of concern. Analysis level from ESP-700 Section (Kimbrough et al. 1988).

Parameter	Analytical level	Container ^a and processing specifics	Analytical Procedure
³ H	IV	4 L Cubitainer, chill return to lab and acidify (HNO ₃) pH < 2	Method 906.0 ^b
⁹⁰ Sr	IV	4 L Cubitainer, chill return to lab and acidify (HNO ₃) pH < 2	Method 905.0 ^b
Gamma emitting radionuclides		21 L Cubitainer, chill to be used for:	
Particle-Associated	IV	filter all of Gamma, collect, air dry filters	
Dissolved	IV	filtrate from Gamma, evaporate to 90 mL	
Inorganics		4 L Cubitainer, chill, to be used for:	
Total metals	IV	1 L PPE bottle, HNO ₃ pH < 2, chill	CLP SOW ^c
Dissolved metals	IV	Filter ^d into 1 L PPE bottle, HNO ₃ pH < 2, chill	CLP SOW ^c
Organics (as listed in Table 4.1, Martin Marietta Energy Systems 1990)	IV	2.3 L Amber glass bottle, teflon lined, chill	CLP SOW ^e

^aAll containers precleaned to EPA specifications. Cubitainers are manufactured from polycarbonate materials.

^bEPA-600/4-80-032 1980.

^cEPA 1987.

^dFilter through nitric acid washed, 0.45 μm diameter effective pore size filters.

^eEPA 1988.

Fig. 2.2. Location of sample sites for Phase 1 w

Water and Sediment Core Collection Sites

Bold numbers are study reaches
Bold lines are reach boundaries

- ▲ water sample
- sediment sample
- water and sediment samples

10 KM

Tennessee State Plane
Datum NAD83
Units Meters
November 1992

er and sediment core collections in Clinch River/Watts Bar System.

Table 2.4. Number of Clinch River RI water samples, by site and type of analysis (validated & accepted data). Dissolved portion includes both inorganic filtered and gamma spec. evaporate samples. See text for details.

Reach	Site	Type of Analysis	Water Sample Matrix			Subtotal
			Dissolved	Residue	Total	
1	Upper Melton Hill	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
1	Solway	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
1	Lower Melton Hill	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
2	Jones Island	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
2	Grassy Creek	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
3	PC/Mouth of East Fork	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	0	1	2
3	K-1007	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
3	Lower Poplar Creek	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
4	Brashear Island	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
4	Kingston City Park	Inorganics	2	0	2	4
		Organics	0	0	2	2
		Radionuclides	1	1	0	2
5	Upper Watts Bar	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	0	2
5	Mid Watts Bar	Inorganics	2	0	2	4
		Organics	0	0	2	2
		Radionuclides	2	1	2	5
5	Lower Watts Bar	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
10	Norris Reservoir	Inorganics	2	0	2	4
		Organics	0	0	2	2
		Radionuclides	2	1	2	5
13	Hartland Bridge	Inorganics	1	0	1	2
		Organics	0	0	1	1
		Radionuclides	1	1	1	3
Total			37	15	54	106

sample devices, constructed from stainless steel for organic analytes and PVC plastic for inorganic and radiological analytes. Samples were collected using the following criteria. All samples were collected during winter conditions to ensure nearly uniform physical-chemical parameters (such as temperature, pH, dissolved oxygen, or water flow) throughout the water column. With the exception of Norris Reservoir, a single sample was collected from 1.0 m below the water's surface for each station. Samples were processed in a manner compatible with the requested analyses (Sect. 9.6.2, Martin Marietta Energy Systems 1990). Samples were collected from 1.5 and 16.5 m deep in Norris Reservoir (reach 10).

Water samples were apportioned for contaminant analysis as follows: (1) gamma-emitting radionuclide samples were filtered, and the filtrate was collected; both filter residue (particle-associated) and dissolved filtrate, evaporated to 90 mL) portions were analyzed for gamma spectrometry; (2) inorganic samples were split into two aliquots for total and dissolved trace metals; dissolved were passed through a 0.45 μm effective pore size filter, acidified following filtration, and chilled, while total were only acidified and chilled; and (3) organic samples were chilled upon collection (Table 2.3). The complete scheme for apportionment and analysis of samples, including water quality and nutrient analyses, is given in Sect. 9.6.3 and Fig. 9.7 of Martin Marietta Energy Systems (1990). Nineteen individual samples were collected from 16 stations in 7 reaches (Fig. 2.2), with the resulting verification, validation, and acceptance of 106 results. Not all analyses were completed for each individual sample. Additionally, field water-quality parameters were measured at each station using a Hydrolab Surveyor II multiparameter water quality monitoring device. All samples were stored, refrigerated over night, and delivered to the analytical laboratory the following morning.

The water concentration values used in the Phase 1 human health and ecological risk screening (Sect. 4) were supplemented with environmental monitoring samples, including routine NPDES monitoring and grab samples collected by environmental monitoring and compliance personnel at each ORR site. Additionally, these data are the current benchmark against which Phase 1 water data are compared in Sect. 3.2.

2.2 SEDIMENT

Sediments may be the short- and long-term repository for particle-associated contaminants. Contaminated surface sediments are also a direct source of particle-associated contaminants to aquatic biota. Through the continuous process of sediment accumulation, sediments preserve a record of the depositional history for particle-associated contaminants, which may be directly proportional to the contaminant release histories. Vertical profiles of contaminants in the sediments can be used to reconstruct past releases of particle-associated contaminants. Data from contaminant surveys of Watts Bar Reservoir (Turner et al. 1985; Olsen et al. 1992) show higher concentrations of ^{137}Cs and mercury at greater depth in the sediment than at the sediment surface. These higher values were associated with the relatively large releases of these contaminants from the ORR in the mid- to late 1950s and early 1960s.

Both surface sediment grab and sediment core samples were collected from all sites, following the criteria established in Sect. 9.5.2 of the CRRFI Plan (Fig. 2.2). Samples were collected from December 1989 through July 1990. Sediment core samples were collected from Norris Reservoir; Solway; lower Melton Hill Reservoir; Jones Island; the mouth of Grassy Creek Embayment; Poplar Creek at the mouth of Mitchell Branch; lower Poplar Creek;

Brashear Island; Kingston Steam Plant; Kingston City park; and upper, mid, and lower Watts Bar Reservoir. In addition to these sites, surface sediment grab samples were collected from Melton Hill Park, Walker Branch Embayment, Grubb Island, the K-770 area, Campbell Bend, Sugar Grove, the Emory River, and Webster Bluff. Samples were sent to several analytical laboratories for analysis of inorganic and organic chemicals and radionuclides. Surface sediment grab samples were collected using a ponar grab sampler. Sediment cores for contaminant profile determinations were collected using either a free-fall gravity corer (at stations with water depths greater than 5 m) or a vibracore device (for shallow water stations), using tubing appropriate to the requested analyses. Sediment cores were sectioned by removing the overlying water, extruding the core vertically, and slicing the core into segments as specified in Sect. 9.5.2 of the CRRFI Plan. All sections were placed in containers appropriate to the requested analyses (glass for organics, glass or plastic for metals, and aluminum cans for radiological analyses). Seventy-one sediment cores, split into 1027 samples (Fig. 2.2), and 487 surface sediment grab samples (Fig. 2.3) were collected from 16 stations in 7 reaches (Table 2.5).

An extensive project of surface sediment grab sample collections from near-shore areas of the CR/WBR system was initiated in July 1990 to assess the extent of surface sediment contamination. This sampling was initiated at the request of an interagency working group on Watts Bar Reservoir contaminated-sediment dredging. This sampling is limited to near-shore sediments in Watts Bar Reservoir, which may be (1) contaminated from past ORR operations or (2) susceptible to sediment disturbing activities. The results of this sampling have been reported regularly to the Watts Bar Interagency Working Group.

Sediment samples were also collected from Braden Branch and the White Oak Creek Embayment (WOCE) areas in conjunction with Phase 1 of the CRRI. Contamination observed in Braden Branch is associated with a state of Tennessee superfund site, not related to DOE ORR operations. Braden Branch was sampled in order to confirm a source of radiological contamination to Melton Hill, above the ORR. Braden Branch data are summarized in Appendix G for informational purposes only. They are not included in the risk screening for human and ecological health (Sect. 4), but are the focus of a separate risk assessment in Appendix G.

Sediment data from WOCE that were collected as part of Phase 1 sampling are summarized in Blaylock et al. (1992). These Phase 1 data led directly to the accelerated site characterization of this area, with the subsequent decision to construct a coffer cell-sediment retention structure at the mouth of the embayment to prevent erosion and transport from WOCE to the Clinch River. The rationale and the activities undertaken by the WOCE project are summarized in Leslie and Kimmel (1992) and Blaylock et al. (1992).

2.3 FISH

The objectives of the Phase 1 fish sampling and analysis were to

1. obtain high quality data to confirm or modify existing data for contaminant levels in fish from the CR/WBR system,
2. determine the range of contaminant concentrations in CR/WBR fish and the spatial distributions of these concentrations in reference to ORR,

Fig. 23. Location of sites for Phase 1 surface survey

ent grab sample collection in Clinch River/Watts Bar system.

**Table 2.5. Number of Clinch River ERP sediment
samples, by reach and type of analysis (validated & accepted data)**

Reach	Site	Type of Analysis	Sample Matrix		Subtotal
			Core Section	Surface Grab	
1	Melton Hill Reservoir	Inorganics	49	0	49
		Organics	25	0	25
		Radionuclides	28	13	41
2	Clinch River: Melton Hill Dam to Poplar Creek	Inorganics	59	0	59
		Organics	46	0	46
		Radionuclides	58	46	104
3	Poplar Creek: East Fork Confluence to Poplar Creek Mouth	Inorganics	79	0	79
		Organics	64	0	64
		Radionuclides	80	3	83
4	Clinch River: Poplar Creek Confluence to Clinch River Mouth	Inorganics	52	0	52
		Organics	59	0	59
		Radionuclides	110	93	203
5	Watts Bar Reservoir	Inorganics	70	0	70
		Organics	63	0	63
		Radionuclides	119	277	396
6	Emory River	Radionuclides	0	31	31
10	Norris Reservoir	Inorganics	23	0	23
		Organics	22	0	22
		Radionuclides	21	1	22
18	Tennessee River: Ft. Loudoun Dam to Clinch River Confluence	Radionuclides	0	23	23
Total			1027	487	1514

3. provide the necessary data required to perform human health and ecological screening risk assessments to refine the list of contaminants of concern, and
4. establish reference (background) concentrations for contaminants of concern for comparison to those sites impacted by releases from ORR.

Because fish accumulate contaminants from water, sediments, and food, they are integrators of many types of environmental conditions (Adams and McLean 1985). Characterization of the nature and extent of contamination in the aquatic biota in the CR/WBR system included the collection of 288 fish (representing three species) from 15 sites in 8 reaches. To characterize the spatial distribution of contaminants in fish, samples were collected from six sites on the Clinch River, three sites on the Tennessee River downstream of its confluence with the Clinch River, five sites on tributaries of the Clinch River, and one reference site (Norris Reservoir) also in the Clinch River system, but upstream of any influence of ORR (Fig. 2.4, Table 2.6). WOCE was also sampled as part of Phase 1 and these results are presented in Blaylock et al. (1992).

Three species representing different trophic levels were sampled with electrofishing gear and gillnets from August 1989 to April 1990. This sampling effort resulted in the collection of 126 bluegill sunfish (*Lepomis macrochirus*), 119 channel catfish (*Ictalurus punctatus*), and 43 largemouth bass (*Micropterus salmoides*), hereafter referred to as bluegill, catfish, and bass. The target sample size of 8 to 10 of each species at each site was achieved at nearly every site for bluegill and catfish, but rarely for bass because of their unavailability at some sites (Table 2.7). Catfish at four sites (CRM 24.0, CRM 20.6, Poplar Creek Mile (PCK) 7.4, and CRM 9.5) were originally collected as part of the Biological Monitoring and Abatement Plan (BMAP) sampling program on the Clinch River.

Field and laboratory processing of the fish followed procedures described in the CRRFI Plan. Fish fillets were analyzed for a total of 85 chemical constituents (Table A.1) including 13 metals, 2 PCB congeners, 5 pesticides (and derivatives), and 65 semivolatile compounds. Catfish fillets were submitted for analysis of the organic analytes and bass and bluegill for the inorganic analytes. Radionuclide analysis included measuring ^{90}Sr in catfish vertebrae and gamma emitters in the fillets of all three species. The inorganic and organic analyses were performed by two laboratories in the Analytical Chemistry Division of ORNL, and the ^{90}Sr and gamma emitters analyses were performed by Oak Ridge Associated Universities (ORAU) and the Environmental Sciences Division (ESD), respectively. Sample batches consisted of 10 to 20 samples and usually included one duplicate sample, one matrix blank sample, and one matrix spiked sample.

2.4 QA/QC PROGRAM FOR THE PHASE 1 DATA COLLECTION

2.4.1 General

The quality assurance program for Phase I was designed to comply with both the U.S. Environmental Protection Agency (EPA) QAMS-005/80 and the ANSI/ASME NQA-1 guidelines. The QA objectives for the data follow.

Fig. 2.4. Location of sample sites for Phase 1

Fish Collection Sites

Bold numbers are study reaches

Bold lines are reach boundaries

② Site identifier shown by
numbers within circles

10 KM

Tennessee State Plane

Datum NAD83

Units Meters

November 1992

collection in Clinch River/Watts Bar Reservoir system.

**Table 2.6. Sample sites for Phase I fish
collection in Clinch River/Watts Bar Reservoir system**

Site No.	Reach segment ^a	River mile or km	Name
1	10.1	CRM 105.0 ^b	Norris Reservoir (reference site)
2	1.1	CRM 51.0	Upper Melton Hill Reservoir
3	1.5	CRM 24.0	Lower Melton Hill Reservoir
5	2.3	CRM 20.6	Jones Island
6	2.5	CRM 14.7	Grassy Creek mouth ^c
7	13.1	PCK 9.6	Poplar Cr. at Hartland Bridge
8	3.1	PCK 8.6	Poplar Cr. at East Fork Poplar Cr. mouth
9	3.2	PCK 7.4	Poplar Cr. at Mitchell Branch mouth
10	3.3	PCK 2.2	Poplar Cr. at K-1007B pond outfall
11 ^d	3.4	PCK 0.8	Lower Poplar Cr. embayment
12	4.2	CRM 9.5	Brashear Island
13	4.5	CRM 0.5	Kingston city park
14	5.2	TRM 557.0	Mid Watts Bar Reservoir
15	5.3	TRM 530.5	Lower Watts Bar Reservoir
16	15.1	TRM 518.0	Upper Chickamauga Reservoir

^aReach segments are divisions of reaches. The number to the left of the decimal is the reach and the number to the right of the decimal indicates the segment.

^bChannel catfish were collected in the Powell River arm of Norris Reservoir at PRM 30.

^cOnly channel catfish were collected at the Grassy Creek site.

^dFish at sites 10 and 11 were sometimes combined because of small sample sizes.

Table 2.7. Number of regular samples (i.e., does not include QC samples) submitted for analysis for each analytical group by species (BLG, bluegill; LMB, largemouth bass; and CHC, channel catfish) and by sampling location

Site	Metals		Pest/ PCBs	Semi- vols.	Gamma rads			SR-90
	BLG	LMB	CHC	CHC	CHC	BLG	LMB	CHC
Norris Res.	10	5	8	8	8	5	4	8
Upper Melton Hill	10	5	8	8	8	2	5	8
Lower Melton Hill	10	3	8	8	8	2	3	
Jones Island	11	2	8	8	8	11	2	
Grassy Cr.	-	-	6	6	6	-	-	6
PC/Hartland Br.	10	-	4	4	4	2	-	4
PC/EFPC mouth	9	-	8	7	8	4	-	8
PC/Mitchell Br.	9	4	8	8	8	2	4	
PC-K-1007	2	3	10	10	10	5	3	10
PC Embayment	8	-	10	9	10	-	-	10
Brashear Island	9	3	8	8	8	9	3	
Kingston Park	10	5	10	10	10	5	5	10
Mid Watts Bar	10	2	9	9	9	2	2	9
Lower Watts Bar	10	3	10	10	10	2	3	10
Chickamauga Res.	8	8	4	4	4	2	8	4
Total	126	43	119	117	119	53	42	87

- Scientific data generated would withstand scientific scrutiny.
- Data would be gathered using appropriate procedures for field sampling, chain-of-custody, laboratory analyses, and data reporting.
- Data would be of known precision and accuracy.

Audits and surveillances were performed by DOE-OR, MMES Central QA, and others to review and evaluate the adequacy of field and laboratory performance, and to ascertain whether the QA/QC Plan was completely and uniformly implemented. Results of these audits and surveillances were documented and reported to management. Follow-up action was taken and corrective actions monitored by the Clinch River QAS.

The CRRRI samples were analyzed for potential contaminants of concern using established procedures.

Water samples were analyzed for metals, semivolatiles, and pesticide/PCBs by the K-25 ACD. All sediment metals analyses were subcontracted to IT laboratories by the K-25 ACD. The following inorganic elements were reported for sediment: Sb, As, Be, Cd, Cr, Cu, Pb, Hg, Ni, Se, Ag, Tl, and Zn. Inorganic elements that were reported for water were as follows: Ca, Fe, Mg, K, Na, Sb, As, Be, Cd, Cr, Cu, Pb, Hg, Ni, Se, Ag, Tl, and Zn. Organic sediment analyses were performed both by the K-25 ACD and by IT. Metals were analyzed by Contract Laboratory Program (CLP) Statement of Work (SOW) 7/87 or 7/88. Organics were analyzed by CLP SOW 2/88, with the exception that K-25 did not use a second column confirmation for the pesticides/PCBs analyses.

Fish samples were analyzed for semivolatiles and pesticides/PCBs by the Organic Analysis Laboratory (OAL) at ORNL. Semivolatiles were analyzed using a modified GC/MS procedure with no acid surrogates or matrix spike/matrix spike duplicate samples. Laboratory quality control elements that were utilized were method blanks, internal standardization, initial calibration, continuing calibration, and GC/MS tuning. Pesticide/PCB samples were extracted by sonication in hexane with sodium sulfate. Samples were spiked with tetrachloro-m-xylene and decachlorobiphenyl. Extracts were concentrated using Kuderna-Danish concentrators and nitrogen-blowdown. Pesticide extracts were acid washed using concentrated sulfuric acid. Percent lipids were determined on all samples. A method blank (sodium sulfate) was extracted with each group of samples. One sample in each batch was spiked with a matrix spike solution containing 0.5 ppm alpha-chlordane, 0.5 ppm gamma-chlordane and 5 ppm Aroclor 1260. The extracts were analyzed on a DB-5 capillary column and a DB-1701 megabore column using electron capture detection.

Additional PCB analyses, as well as metals analyses were performed by the ORNL Inorganic Analysis Group (IAG). An internal procedure was used for sample preparation and analysis for PCBs and lipids. An International Atomic Energy Agency (IAEA) fish homogenate was used as a laboratory control sample. Mercury was analyzed by the ORNL $\text{HNO}_3/\text{HClO}_4$ method (AC-MM10915) for sample preparation and a cold vapor atomic absorption method (AC-MM1214922) for the analysis of Hg. The cold vapor method measures mercury using a standard addition technique. A National Institute of Standards and Technology (NIST) standard was used for the standard addition. For analysis of metals, ICP/MS was used to measure As, Be, Tl, and ICP/OES was used to measure Sb, Cd, Cr, Cu, Pb, Ni, Se, Ag, and Zn. An internal method was used for sample preparation. A VG Plasma Quad P02 was used for the ICP/MS analyses and a JY-48 instrument was used for the

ICP/OES analyses. The ICP/MS measurements were made using the standard addition technique.

The Oak Ridge Associated University (ORAU) Radiological Laboratory provided radiochemical analytical services for the alpha-beta radionuclide analysis of all samples. QA activities were identified in Weaver (1988) and standard laboratory procedures were described in Weaver et al. (1988). The Oak Ridge National Laboratory-Environmental Sciences Division (ORNL-ESD) was responsible for analysis of gamma emitting radionuclides (gamma spectroscopy) of biota, sediment, and water.

The QA objectives for precision and accuracy in the laboratory were defined by control limits established for the EPA Contract Laboratory Program. Acceptance criteria for the radiological analyses were generated by each laboratory.

The CRRRI program required the independent review and validation of the analytical laboratory data for all Phase I analyses. The review and validation was conducted by a subcontractor following EPA and/or project-specified guidelines to ensure uniformity and comprehensiveness of the review.

2.4.2 Field Quality Control

Quality Control in field operations, including selection of sampling locations, field data recording and sample collection, were implemented to meeting the Phase 1 project objectives (outlined in Sect. 1.1). Sampling sites for each matrix (fish, sediment or water) and QC sample collection frequencies were specified by the CRRFI Plan, with final determination of each site, and QC sample collection the responsibility of the sample team leader. Methodologies followed for the major categories of field activities, including record keeping, establishment and maintenance of sample custody, instrument calibration, and sample identification were specified in standard operating procedures (SOPs); sample collection procedures followed those of Kimbrough et al. (1990).

All field activities followed standard record keeping, chain-of-custody procedures. These included recording of site specific information in water-proof notebooks, with routine reviews of the notebooks. Sample custody was established by the sampling team upon collection, through the use of standard chain-of-custody forms, and was maintained throughout sample processing, and delivery to analytical services. The goal for submittal of quality control samples was once for every 20 samples, as appropriate. Quality control samples include field duplicates, matrix spikes, equipment cleaning rinse samples and field rinse water blanks. Matrix specific field quality control activities are discussed below.

Fish. Site selection was dependent upon the species targeted for collection. Sampling occurred during the fall and spring, when fish were available. Duplicate samples from different halves of the fish were submitted for the requested analysis. Matrix spikes, fish flesh spiked with known quantities of analytes of interest were submitted for organic analysis approximately once every 20 routine samples. Replicate samples were submitted for organic and inorganic analysis approximately once every twenty samples.

Sediment. Criteria for selection of sample collection sites included those that were representative of sediment deposition zones, as defined in Martin Marietta Energy

Systems (1990). Reference material, consisting either of homogenized sediment, stored in a deep freeze, or NBS standard material, was submitted for each sediment core, for inorganic and organic analyses. Approximately one set of duplicate sections for each sediment core was submitted for each requested analysis group.

Water. Samples were collected from mid-channel in the same vicinity as other media. Due to the small number of samples collected, one field duplicate sample, and one equipment cleaning blank was collected per week of sampling. Field blanks and trip blanks were collected approximately once every twenty samples.

2.4.3 Laboratory Quality Control Program

The laboratory quality control program must ensure that all data generated and reported are scientifically valid, consistent with accepted methods, and of known accuracy. An accurate method will produce precise and unbiased results. Results from quality control samples are used to document data quality, verify that the analytical system is functioning for a given matrix/analyte, indicate that an instrument is functioning properly, identify when additional corrections need to be made to the analytical system, and determine the effect of these corrections. The QC data provide an important indication of how well the data quality objectives were met. The types of quality control samples used during Phase I are summarized below.

Radiological check sources were used to determine general performance of the counting system and to ensure that the efficiency of the system had not changed.

2.4.4 Data Validation

Analytical results from the laboratories were received by the project database manager in either electronic or hard copy format and merged into a large working database. Analytical results included the concentration of the analyte in the sample if it was present and measurable. When an analyte was not detected in a sample, the sample quantitation limit (SQL) for that particular sample was reported as the result accompanied by a "U" prefix. When spectral data indicated the presence of a compound that met the identification criteria but the result was less than the sample quantitation limit, a "J" flag was assigned indicating that the result was an estimated value. In data analyses, these "J" values were treated as normal results.

The primary goal of the Clinch River Environmental Restoration Quality Assurance (QA) Program is to ensure that the analysis of all environmental samples produces data of known quality. According to EPA, "the quality of data is known when all components associated with its derivation are thoroughly documented, with such documentation being verifiable and defensible" (USEPA, 1990). In fulfillment of this goal, a subcontractor, independent of the analytical laboratory that generated the data, was employed by the CR-ERP to validate the analytical results from Phase 1. Organic data packages were assessed by "Laboratory Data Validation Functional Guidelines for Evaluating Organics Analyses" (USEPA), and inorganic data packages were assessed by "Laboratory Data Validation Functional Guidelines for Evaluating Inorganics Analyses" (USEPA), as applicable. Radiological parameters were assessed by SAIC's "Laboratory Data Validation Guidelines for Evaluating Radionuclides Analysis," revision 3. The validation process included an assessment

of a variety of Quality Assurance/Quality Control (QA/QC) practices, including holding times, matrix spike recovery rates, surrogate recovery rates, calibration standards, and method blanks. Validation flags (or qualifiers) were assigned by the reviewer to each result depending on the quality of the data as determined by the reviewer. A total of 268 data packages, containing approximately 1700 inorganic samples, 1500 organic samples, and 2200 radiological samples were validated. As a result of the external validation some data were rejected as unusable and others were flagged as estimated values. Table 2.8 summarizes the various data qualifier combinations which were attached to each result by the analytical laboratory and/or the validators. Additional data checking was performed as the data were summarized and evaluated for the health and ecological screening assessment.

2.4.5 Data Assessment

All data deficiencies uncovered in the data validation process were reported to the CR-ERP by the subcontractor's nonconformance process. The CR-ERP reported all nonconformances to laboratory or program personnel for resolution. A nonconformance database was established to track resolution and trends. Two major areas for nonconformances were method use and missing data information. Other areas for nonconformances were illegibility, missing quality control information, matrix spikes missing or recoveries outside limits, and logbooks or worksheets missing or incorrect.

The number of rejected samples is summarized in Table 2.9. All pesticide/PCB and semivolatile analysis of sediment samples that were analyzed by IT were rejected due to missed holding times. Approximately 50% of the ORNL semivolatile data was rejected because of multiple problems affecting analytical performance, including surrogate recoveries outside of limits, surrogates not added, and inadequate quality control. The QA completeness objective for the project was to obtain valid analytical results for at least 95% of the samples collected. Overall completeness for all Phase 1 media and analyses combined was 87.8% valid data. If all the data is considered except for the analysis of semivolatiles in fish (which had the highest rejection rate, 48%), then the overall completeness is 96%.

Table 2.8. Data qualifiers

Qualifier	Definition
U	Compound was analyzed for but not detected. Sample quantitation limit was reporter as the result.
J	Indicates an estimated value either because of laboratory estimation of concentration or because of validation deficiency.
UJ	Compound was analyzed for but not detected, and detection limit was qualified as an estimate due to validation deficiency.
NJ	Tentative compound identification only and estimated concentration. No second column confirmation of pesticides.
X	Fish tissue which was held frozen while awaiting extraction exceeded allowable holding time for water samples. However, these criteria are not applicable to biological samples, and results are considered valid.
UX	Compound was analyzed for but not detected, and water holding times were exceeded (see X definition).
V/V	Indicates that the result has been verified and is a valid result.

Note: Data qualifiers applied to the sample analysis result by the analytical laboratories and/or independent data validator. In cases where there was no data qualifier required, "V/V" has been inserted in the data qualifier field.

**Table 2.9. Summary of the number of rejected
samples as a result of the validation of laboratory analyses**

Analysis class	Analysis type	Laboratory	Total no. of analyses	No. of accepted analyses	No. of rejected analyses	Percent rejected of total
INORGANICS	METALS	K25	664	660	4	0.6%
		K25/ IT	4313	3818	495	11.5%
		ORNL/ CPA	2027	1933	94	4.6%
	MERCURY	ORNL/ EAL	169	169	0	0.0%
ORGANICS	PEST/ PCB	K25	7830	7205	625	8.0%
		K25/ IT	27	0	27	100%
		ORNL/ CPA	238	225	13	5.5%
		ORNL/ OAL	1160	1123	37	3.2%
	SEMI- VOLS	K25	19306	19234	72	0.4%
		K25/ IT	65	0	65	100%
		ORNL/ OAL	7539	3930	3609	47.9%
		OAU	941	904	37	3.9%
RADIO- NUCLIDES		ORNL/ ESD	2339	2336	3	0.1%
TOTAL NUMBER OF SAMPLES			46618	41537	5081	12.2%

3. SITE CHARACTERIZATION

3.1 INTRODUCTION

The purpose of this section is to discuss contaminant concentrations observed in Phase 1 of the CRRI across the system and in each reach. While data tables referred to in this section include all contaminants for which analyses were performed and accepted through the verification and validation process, discussions are limited to the contaminants identified by the risk screening analysis as high priority contaminants or those contaminants requiring further study. This information is presented in individual sections for each medium type (water, sediment, and fish).

All of the summary tables referred to in this report include data qualifier flags (Table 2.8). It is important to recognize that the derivation of the reported data is dependent on understanding these qualifiers. Contaminants that were not detected at all, or that were detected but not quantifiable, have reported concentrations that are estimates, based upon the detection limits of the instruments used. Assigning the value of the "limit of detection" to compounds that were not detected yields a conservatively high "estimated concentration" for those contaminants.

3.2 WATER

Surface water, the primary transport pathway for all contaminants leaving the reservation, is also the principal exposure medium for the initial introduction of dissolved (non-particle-associated) contaminants into the food web. The complex hydrodynamics and spatial heterogeneity of the CR/WBR system make it exceedingly difficult to adequately estimate the temporal and spatial extent of surface water contamination, even with a large number of samples. The NPDES monitoring of permitted discharges provides considerable information on contaminant concentrations near discharge points.

Phase 1 sampling of water was undertaken to (1) obtain high-quality data to confirm or modify suspected low concentrations of contaminants that were identified in scoping studies (Suter 1991, Hoffman et al. 1991, Olsen et al. 1992), (2) determine the range of contaminant concentrations present in the river-reservoir system, (3) provide data for the identification of contaminants of concern, and (4) establish reference concentrations for those contaminants. Accordingly, characterization of the nature and extent of surface water contamination in the CR/WBR system during Phase 1 of the CRRI was limited to a single sampling event for each station. This single water sample from each station was collected during winter conditions of uniform physical-chemical parameters. Samples were analyzed for inorganic, radiochemical, and organic constituents. These data were collected to confirm the results of scoping studies and monitoring reports and they will be supplemented with NPDES monitoring data to evaluate human and ecological risk on a screening level. The intent is to focus sampling and analyses on specific areas and contaminants during Phase 2 of the CRRI.

3.2.1 Discussion of Contaminants for the System

The following results reflect the outcome of analyses of total dissolved (filtered) water samples for metals, PCB/pesticides, semivolatile organic compounds, and radionuclides. Few

contaminants were detected in Phase 1 water samples. Organic, inorganic, and radiological analytes that were observed, verified, and validated as acceptable values are included in the data tables, the text, and this discussion. The detected analytes are listed by descending frequency of occurrence in Appendix B.

The most prevalent analytes, observed in both total and dissolved portions of all samples submitted, are the cations calcium, magnesium, potassium, and sodium. Total and dissolved iron were observed less frequently.

Constituents with the potential to pose a risk to human health or the environment rarely met all of the quality criteria for acceptance into the database. Potential contaminants of concern include total and dissolved lead, total copper, dissolved cadmium, total and dissolved zinc, dissolved mercury, and total and dissolved silver (Appendix B). The distribution of these contaminants is discussed in Sect. 3.2.2. Inorganic contaminant values that were averaged over the whole CR/WBR system, with one exception (zinc), were in the ranges expected based on previous monitoring reports (Kornegay et al. 1991). Each inorganic contaminant detected during the Phase 1 investigation is well below the Tennessee Water Quality Criteria for Designated Water Uses (rules of the TDEC, Chapter 1200-4-3). The observed values for zinc (1 total and 1 filtered) are greater than the maximum reported for environmental monitoring of the Clinch River. Bis-(2-ethylhexyl)-phthalate was the only organic contaminant observed in any of the samples collected for Phase 1 site characterization (Appendix B).

Radiological constituents observed include ^{90}Sr , particle-associated and dissolved (evaporate) ^{137}Cs , and dissolved ^{60}Co (Appendix B). Radionuclides that have been observed, verified and validated were more common than organic or most inorganic contaminants. For the dissolved and total water phase radionuclides, observed values were within and below the range of values reported for environmental monitoring data (Kornegay et al. 1991). Analogous values for particle-associated ^{137}Cs are not present in the environmental monitoring report; however, values from Appendix B are within the range of those reported by Olsen et al. (1992).

3.2.2 Discussion of Contaminants by Study Reach

The discussion of patterns in the distribution of organic, inorganic, and radiological analytes is restricted to listing the contaminants found in particular reaches; there are too few observations to adequately address variability of the surface water environment. The data for this section are compiled in Appendix B.

Dissolved zinc was detected only at the Norris Reservoir reference site (reach 10). This observation should be investigated further, since no other values for dissolved zinc were detected in Phase 1 data. Dissolved cadmium and total and dissolved lead were observed in upper Melton Hill Reservoir (reach 1), upstream from all active ORR inputs to the CR/WBR system. The majority of the potential contaminants of concern for which observations have been verified, validated, and accepted were located downstream from ORNL operations (reach 2, between White Oak Creek and Poplar Creek on the Clinch River) and in the vicinity of the K-25 Site (reach 3, Poplar Creek Embayment). Total copper and zinc were detected in reach 2 surface waters, with values in the range previously detected for environmental monitoring (Kornegay et al. 1991).

Values for dissolved lead, observed in Poplar Creek upstream from the East Fork confluence (reach 13), were comparable to those reported previously (Kornegay et al. 1991); however, dissolved lead was not detected in any Phase 1 sample from reach 3, downstream in Poplar Creek. Dissolved mercury, total and dissolved silver, and total lead were all observed in upper portions of reach 3 (Appendix B). Mercury and lead values are within the range previously reported for this reach, though silver values are greater than previously reported (Kornegay et al. 1991). Dissolved cadmium and dissolved and total lead values in the Clinch River above the confluence of the Clinch and Tennessee rivers (reach 4) are greater than those observed for environmental monitoring.

Gamma-emitting radionuclides were not observed in water samples from any of the reference reaches, including Norris Reservoir and Poplar Creek. Additionally, none were detected in Melton Hill Reservoir. Both dissolved and particle-associated ^{137}Cs were detected in reaches 2, 4, and 5 (Appendix B); all values were similar to those reported by Olsen et al. (Table 5 in Olsen et al. 1992) and Kornegay et al. (1991). Only dissolved ^{137}Cs was detected in Poplar Creek (Appendix B). Cobalt-60 was detected in a single filtered sample from reach 5 (Appendix B) at levels comparable with those reported by Olsen et al. (Table 5). Though ^{90}Sr was reported as present in all reaches, the average values for each reach were less than the standard deviation of the measurements (Appendix B). Concentrations of radionuclides in surface waters are comparable with values for both surface sediments and current releases from the ORR (Olsen et al. 1992; Kornegay et al. 1991).

3.2.3 Summary

The cations calcium, magnesium, potassium, and sodium—the most prevalent analytes—are constituents of weathered rock and are commonly observed in surface waters. Values for the CR/WBR system are comparable with those observed in routine monitoring near the ORR and in other freshwater ecosystems (Kornegay et al. 1991; Table 10.1, Goldman and Horne 1983). The concentrations of these cations and shifts in their relative abundance within the system affect the geochemistry of the surface water, influencing contaminant availability and the ability of the ecosystem to respond to perturbation.

Inorganic and organic constituents with the potential to pose a risk to human health or the environment were in the ranges expected, based upon previous monitoring reports (eg. Kornegay et al. 1991). The majority of potential contaminants of concern were detected between White Oak Creek and Poplar Creek on the Clinch River (reach 2) and in the Poplar Creek Embayment (reach 3), both of which drain several ORR contaminant sources. Total copper and zinc were detected in reach 2 surface waters, with values in the range previously observed for environmental monitoring (Kornegay et al. 1991). Dissolved lead, observed in samples from the Poplar Creek reference reach (reach 13), may originate from an upstream contamination source.

The Y-12 Plant, the city of Oak Ridge waste water treatment facility, and the K-25 Site operations are among the possible sources for mercury, silver, and lead in detected in reach 3. The elevated levels of cadmium and lead in the Clinch River between Poplar Creek and the Tennessee River confluence may come from several sources in and upstream from this reach.

Phase 1 values for all radiological constituents were comparable with the mean values reported for environmental monitoring (Kornegay et al. 1991) and previous field

investigations (Olsen et al. 1992). These values should be used with caution as they may exaggerate the ^{137}Cs concentration observed for surface water systems. The methodology used (Sect 2.2) is sensitive to low concentrations of suspended solids and tends to overrepresent any ^{137}Cs detected on the filters. Concentrations of ^{137}Cs on suspended particles should be compared only generally with those observed for surface sediments, as the latter is a much more accurate measurement.

Dissolved and particle-associated ^{137}Cs were detected in reaches 2, 4, and 5, with values similar to those reported previously for these reaches. The dissolved ^{137}Cs detected in Poplar Creek, was possibly due to backflow from the Clinch River. Cobalt-60 in reach 5 was detected at levels comparable with those reported previously. Average ^{90}Sr values for all reaches are below levels of concern. Currently, each of these radionuclides is released from the ORR to surface waters. Though comparable with surface sediment values, these surface water radionuclides likely do not result from resuspension of surface sediments in the CR/WBR system.

High quality data confirming the contaminant concentrations observed in the scoping studies, the first objective of the Phase 1 sampling, was achieved for the limited sampling performed. Additionally the range of contaminant concentrations present in the CR/WBR system, the second objective of the Phase 1 sampling was determined, relative to the sampling conducted. The bulk of observations were comparable with those observed previously, and they did demonstrate the range of contaminant concentrations observed within the system. However, more high quality information will be collected as part of Phase 2 of the CRRI in order to determine the spatial-temporal patterns in contaminant concentrations.

Phase 1 water sampling did provide data, augmented with source monitoring data from NPDES and DOE Order 5400.1, for identification of contaminants of concern. In order to more fully support the identification of contaminants of concern, the second sampling phase must characterize the relationship between routine monitoring results and ambient surface water conditions, as well as refining the temporal-spatial concentration range for possible contaminants of concern. Contaminant concentrations from reference stations must also be assessed over time in order to provide relevance to values for contaminants of concern.

The inorganic, organic, and radiological results from these Phase 1 water samples, in combination with historic data, indicate that a greater effort must be made to characterize the Clinch River, Poplar Creek Embayment, and Kingston areas. The sampling and analysis plan for Phase 2 will have a much greater emphasis on the investigation of surface waters.

3.3 SEDIMENT

Many of the potential contaminants of concern identified in the screening phase of the CRRI are particle reactive and thus are associated with sediment particles; therefore, the scoping phase of the CRRI focused on identifying sediment deposition areas below the ORR. Because mercury and ^{137}Cs had known release histories and ^{137}Cs is easily analyzed, they were the only constituents analyzed in sediment during the scoping phase. The goal of the scoping phase was to identify areas of contaminated sediment accumulation in the CR/WBR. The results from the scoping phase indicated that sediment with contaminants originating from the ORR were predominantly found in the deep channel sediments of Watts Bar Reservoir. Previous studies (Oakes et al. 1982, Turner et al. 1985) also indicated that contaminated

sediment accumulated in deep channels of the Clinch River arm of Watts Bar Reservoir. Sediment core samples during Phase 1 were collected from these deep channel areas and other suspected sediment deposition zones and analyzed for potential contaminants of concern. Because sampling occurred in areas of known contamination, the Phase 1 sampling was biased towards "hot spots" and yielded conservatively high estimates of average concentrations for reaches. These data provide the information needed to evaluate human and ecological risk on a screening level. The intent was to further focus sampling and analyses during Phase 2 to specific areas and specific contaminants.

Near-shore surface sediment characterization sampling was initiated during Phase 1 to provide the data necessary to evaluate human health risks from direct exposure to sediment. This was a concern because near-shore sediment in Watts Bar Reservoir is exposed for parts of each year, and most of the sediment disturbing activities (e.g., dredging and dock construction) occur in this zone of the reservoir. Therefore, as part of the permitting process conducted by the U.S. Army Corps of Engineers and TVA for (sediment disturbing activities), samples were collected to evaluate human health risk on a permit-by-permit basis. Near-shore sediment samples were only analyzed for gamma-emitting radionuclides. The rationale for this approach follows that of the phased approach described for the sediment sampling and analysis. If concentrations of ^{137}Cs or ^{60}Co were high enough to indicate a human health risk resulting from deposition of contaminants originating from the ORR, additional analyses would be performed to determine concentrations of the other contaminants.

3.3.1 Discussion of Contaminants for the System

3.3.1.1 Near-shore sediment samples

Cesium-137 was present in detectable quantities in 99% of the 487 samples that were analyzed. Cobalt-60 was present in detectable quantities in 58% of the 488 samples that were analyzed (Appendix C.3). Of all the analyses performed on these samples, less than 4% were rejected (Table 2.9). This satisfies the completeness requirement of 95% defined in the Work Plan. One limitation with this data set is that only one grab sample was collected in Norris Reservoir (reach 10); therefore, a reference concentration average was calculated from the single grab sample and the top layer (3 cm) of the sediment core taken in reach 10. Data from 23 grab samples from the Tennessee River above the confluence with the Clinch River (reach 18) also provide a reference for surface sediment concentrations of gamma-emitting radionuclides (Appendix C.4).

3.3.1.2 Sediment cores

Sediment cores were collected to provide data for evaluating secondary pathways of human health risk and ecological risk. The exposure pathways resulted from a dredging scenario where deep sediment is dredged and spread on the nearby land. Cores were sectioned in 3- to 6-cm intervals and analyses were performed on each section. The summary tables report core averages and/or reach averages when more than one core was collected in a reach (Appendix C.2).

Detectable concentrations were reported for all 13 inorganic contaminants for which analyses were conducted; however, selenium, antimony, and thallium were detected in less than 30% of the samples (Appendix C.1). Some maximum values are estimates based on

detection limits; thus, in some cases, concentrations used in the risk screening analyses may be conservatively high. The number of inorganic samples rejected was less than 12% (Table 2.9). This does not satisfy the completeness requirement defined in the Work Plan. However 9 of the 13 inorganic contaminants had less than 1% rejected samples. Antimony, mercury, selenium, and silver all had greater than 10% but less than 50% of the samples rejected.

Twenty-two of the 92 organic analytes were detected in at least one sample. Eighteen of these were detected in less than 7% of the samples. The highest percentage (36%) of detected organics in core samples was for bis(2-ethylhexyl)phthalate, which is a plasticizer found in many environmental samples as a common artifact. Twenty percent of the samples analyzed had detectable quantities of Aroclor 1254. Fifteen percent had detectable levels of fluoranthene, 11% had detectable quantities of pyrene, and less than 7% had detectable levels of phenanthrene (Appendix C.1). Because of the small percentages of detected organic contaminants reported, minimum, maximum, and average values are almost always estimated values. These are indicated by the "U" or "U/J" codes in the data qualifier fields in the tables. Less than 1.0% of the semi-volatile organic analyses were rejected during the validation procedure (Table 2.9). This is well within the completeness requirement of 95% as outlined in the Work Plan. Eight percent of the PCB/pesticide organic analyses were rejected.

Samples were analyzed for alpha-emitting and gamma-emitting radionuclides. Every sample for which analysis for alpha-emitters was performed had detectable, though not always quantifiable, levels. Because of the low values observed, many of the concentrations listed in the tables are estimated from detection limits. Therefore, the risk screening analysis is based on estimated concentrations that are conservatively high. For gamma-emitters, ^{137}Cs was present in detectable quantities in 93% of the samples analyzed, and ^{60}Co was detected in 36% of the samples analyzed (Appendix C.1). Only 3% of the radionuclide analyses were rejected during the validation procedure (Table 2.9). This is within the completeness requirements defined in the Work Plan.

3.3.2 Discussion of Contaminants by Study Reach

3.3.2.1 Near-shore sediment samples

Cesium-137 concentrations in the near-shore sediments follow an expected geographic trend (Fig. 3.1). Norris Reservoir (reach 10) and Melton Hill Reservoir (reach 1) show background concentrations (0.33 and 0.26 pCi/g, respectively). The Clinch River between the White Oak Creek confluence and Poplar Creek confluence (reach 2) has the highest average concentrations and the highest individual values in the CR/WBR system. The Clinch River below Poplar Creek and above the confluence with the Tennessee River (reach 4) has a slightly lower average ^{137}Cs concentration than reach 2. Poplar Creek (reach 3) and the Emory River (reach 6), both of which are tributaries to the Clinch River, have concentrations above background but approximately 2.5 times lower than reaches 2 and 4. Slightly elevated levels of ^{137}Cs in these reaches are not unexpected due to the reversing flows into both Poplar Creek and the Emory River resulting from TVA reservoir operations for peak electrical power. Concentrations of ^{137}Cs in the Tennessee River arm of WBR (reach 18) are at background levels. The near-shore surface sediments in the main body of Watts Bar Reservoir (reach 5) have an average concentration below 1 pCi/g, which approaches background concentration levels (NCRP 1978).

Reach

Fig. 3.1. Average ^{137}Cs concentration in surface sediment grabs within each reach. Value for reach 10 is not an average. It is the concentration of the single sample collected in this reach during Phase 1 sampling. Reaches are arranged from upstream on the top to downstream on the bottom.

The average ^{60}Co concentrations within the study reaches show the same geographic trend as ^{137}Cs , with two exceptions (Fig. 3.2). First, ^{60}Co was not detected in Norris Reservoir. Second, the average ^{60}Co concentration in reach 1 is higher than background levels. This can be explained by releases from a closed medical supply facility in Braden Branch that acts as a source within reach 1 but was not part of the ORR operations. This source is discussed in further detail in Appendix G.

3.3.2.2 Core sediment samples

Summary results for sediment cores are presented in Appendix C.2. The arithmetic averages for each reach, reported in Appendix C.2, were calculated using all sections of all cores within a reach and detection limits as actual values when concentrations were below detection limits. For the sake of comparison, if a contaminant was detected in any reach, the average concentration for that contaminant is reported for all reaches even if that contaminant was not detected in all reaches. Again, the data qualifier flags indicate if reported concentrations are estimates. Average concentrations are discussed, rather than minimum and/or maximum concentrations, because in a dredging scenario, exposure would be to well-mixed sediment. The averaging of concentrations across core sections and within reaches mathematically approximates the mixing of sediment as a result of dredging.

Inorganics

There are several spatial patterns in the distributions of average core concentrations of inorganic contaminants. These patterns are shown in Figs. 3.3 through 3.7, which portray average contaminant concentrations for sediment cores arranged along hydrologic continuum. Presenting the averages by site rather than by reach, but still along a downstream continuum, provides more insight to possible sources of contaminants and will help determine where to focus sampling during Phase 2. Arsenic (Fig. 3.3), beryllium (Fig. 3.4), chromium and copper (Fig. 3.5), and mercury (Fig. 3.6) all show a peak concentration in Poplar Creek (reach 3) or at Brashear Island immediately downstream Poplar Creek (first bar in reach 4), suggesting that Poplar Creek is a source of these contaminants into Watts Bar Reservoir. Concentrations of thallium were detected in Poplar Creek (reach 3) and in McCoy Branch (reach 7), both of which are downstream of a coal ash-pile disposal areas. Average concentrations of arsenic, beryllium, and mercury decrease in downstream sediment core samples, with some concentration averages approaching reference levels in reach 5. Zinc (Fig. 3.3); cadmium (Fig. 3.4); and chromium, copper, and lead (Fig. 3.5) all have peak average concentrations in reach 5 (Watts Bar Reservoir). The location of these peaks suggests that there is a source of these contaminants downstream of the ORR either on the Tennessee River arm or below the confluence of the Tennessee and Clinch rivers. There are two superfund sites in Rockwood that drain into Caney Creek above the three sampling sites in reach 5. Additionally, historic metal processing operations based in Rockwood may explain the elevated levels of these metals in Watts Bar Reservoir. Industrial activities on the Tennessee River arm of Watts Bar Reservoir could also be responsible for these contaminants. Future sampling will focus on determining the source of these metals. There are no distinctive patterns in the vertical profiles of the inorganic contaminants with the exception of mercury. Concentration peaks of mercury between 40 and 80 cm deep are apparent in sediment cores from sites in and downstream of Poplar Creek. These peaks coincide with releases of mercury during the mid-to late 1950s and have been well documented by Olsen et al. (1992) and Turner et al. (1985).

Reach

Fig. 3.2. Average ⁶⁰Co concentration in surface sediment grabs within each reach. Value for reach 10 is not an average. It is the concentration of the single sample collected in this reach during Phase 1 sampling. Reaches are arranged from upstream on the top to downstream on the bottom.

Fig. 3.3. Average concentrations of arsenic and zinc for cores sampling sites. Reaches and cores sampling sites within reaches are arranged from upstream on the top to downstream on the bottom.

Fig. 3.4. Average concentrations of beryllium and cadmium for cores sampling sites. Reaches and cores within reaches are arranged from upstream on the top to downstream on the bottom.

Fig. 3.5. Average concentrations of chromium, copper, and lead for cores sampling sites. Reaches and cores within reaches are arranged from upstream on the top to downstream on the bottom.

Fig. 3.6. Average concentrations of mercury for cores sampling sites. Reaches and cores within reaches are arranged from upstream on the top to downstream on the bottom.

Fig. 3.7. Average concentrations of ^{137}Cs for cores sampling sites. Reaches and cores within reaches are arranged from upstream on the top to downstream on the bottom.

Organics

It is somewhat difficult to interpret the concentrations of organic compounds in the sediment cores because the variability associated with individual estimated averages are often larger than the averages themselves (Appendix C.2). These large variances are a result of ambient concentrations at or below analytical detection limits and indicate generally low concentrations of organics in sediments throughout the system.

Aroclor 1254 was detected only in reaches 2 (once), 3 (twice), and 4 (51 times). Aroclor 1254 was found almost exclusively in reach 4, specifically in the core at CRM 9.5. This suggests an upstream source and could include the ORR from Poplar Creek or the municipal wastewater treatment facilities for the City of Oak Ridge or Oliver Springs. Aroclor 1260 was detected in reach 2 once.

Bis(2-ethylhexyl)phthalate was observed in detectable quantities in reaches 2, 3, 4, 5, and 10. This compound is a plasticizer and is commonly found in sediments in industrial countries. Fluoranthene, phenanthrene, and pyrene in the CR/WBR system were detected in reach 2 and 3 and to a lesser degree in reach 4. This implies that these compounds originated from the ORR or another upstream source. These polycyclic aromatic hydrocarbons are contaminants commonly found in streams receiving industrial effluent.

Radionuclides

All of the alpha-emitting radionuclide values for sediment cores were estimates because they were reported as being present but not in high enough concentrations to be quantifiable. Whereas average concentrations of ^{234}U , ^{235}U , and ^{238}U are higher in reach 3, the ratios of $^{235}\text{U}/^{238}\text{U}$ (0.04) and $^{234}\text{U}/^{238}\text{U}$ (1.10) are close to what is expected in undisturbed regions ($^{235}\text{U}/^{238}\text{U} = 0.046$ and $^{234}\text{U}/^{238}\text{U} = 1.0$) (NCRP 1978, Olsen et al. 1989). Average ^{90}Sr concentrations are highest (0.396 pCi/g) in reach 3 also but are well below soil concentrations expected from fallout (1 pCi/g) (NCRP 1978). Higher concentrations of these radionuclides would be expected in Poplar Creek given the proximity to the Y-12 Plant and the K-25 Site. Sampling and analysis during Phase 2 will focus on these contaminants in Poplar Creek.

The gamma-emitting radionuclides show expected spatial trends in core average concentrations (Fig. 3.7). Background concentrations of ^{137}Cs are found in reaches 10 and 1 which are both upstream of White Oak Creek, the primary source of this contaminant. Immediately downstream of the Clinch River and White Oak Creek confluence, average core concentrations of ^{137}Cs are above background concentrations. Poplar Creek (reach 3) shows background levels at the site just downstream East Fork Poplar Creek and slightly elevated concentrations at the lower site, which is influenced by backflow from the Clinch River and the K-25 coal-fired power plant which took Clinch River water for cooling and discharged it into Poplar Creek. The highest average concentration and individual concentration are found at the Brashear Island core. This area is the first major depositional zone downstream of White Oak Creek and Poplar Creek. Average ^{137}Cs concentrations drop downstream but remain above background levels. It is important to recognize that the concentrations in the surface layers for these samples are relatively low and that the averages are dominated by large peak ^{137}Cs concentrations that are typically 50 to 80 cm deep in the sediment (Olsen et al. 1992). These peaks are associated with relatively large releases of ^{137}Cs from White Oak Lake in the mid- to late 1950s (DOE 1988).

Average concentrations of ^{60}Co follow a similar trend as ^{137}Cs concentrations except for slightly higher levels in reach 1 (Fig. 3.8). However, because ^{60}Co was detected in only 36% of all samples, many of the concentrations reported in Appendix C are estimates based on detection limits. These higher concentrations probably result from releases from the off-site medical equipment productions facility discussed in Appendix G.

3.3.3 Summary

Concentrations of ^{137}Cs in near-shore surface sediments in Norris, Melton Hill reservoirs, and the Tennessee River arm of Watts Bar Reservoir represent background levels (<1.0) because they are upstream of ORR and only received radioactive fallout from atmospheric testing of nuclear weapons. Concentrations in Poplar Creek and the Emory River average 1 pCi/g, which is consistent with background concentrations. Cesium-137 concentrations in the near-shore surface sediments in the Clinch River arm of Watts Bar reservoir are between 2.7 and 3 pCi/g. Concentrations in Watts Bar Reservoir below the confluence of the Tennessee and Clinch rivers are below 1 pCi/g, again within background concentrations. To date, no sediment disturbance permit that was evaluated by TVA, ACOE, and DOE were rejected on the basis of radiological contamination. Three sites within the Clinch River arm of Watts Bar Reservoir had concentrations between 10 and 34 pCi/g and will be targeted for sampling and analysis during Phase 2 of the CRRI. Samples collected at these areas will be analyzed for the contaminants identified by the risk screening as requiring further research.

Patterns of inorganic contaminants in sediment core samples indicate that there are multiple sources for these contaminants. Arsenic, beryllium, cadmium, chromium, copper, and mercury in Poplar Creek and the downstream portion of the Clinch River arm of Watts Bar Reservoir appear to be originating, at least in part, from Poplar Creek. Mercury releases from the Y-12 processing plant into Poplar Creek are well documented (Olsen et al. 1992, Turner et al. 1985). Other inorganic contaminants likely result from disposal of coal fly-ash material and photographic processing chemicals from the K-25 Site or the municipal wastewater treatment facilities for the cities of Oak Ridge and Oliver Springs. Concentrations of cadmium, lead, and zinc are higher in the sediments in Watts Bar Reservoir below the confluence of the Tennessee River and Clinch River arms, suggesting that there is a source of these contaminants either within the Tennessee River arm of the reservoir or below the confluence of the two rivers. Phase 2 sampling and analysis will focus on locating areas where concentrations of these contaminants are above acceptable levels.

The lack of detectable quantities of most of the 92 organic chemicals for which analyses were conducted allows significant reduction of the list of organic contaminants to be studied. Fluoranthene, phenanthrene, and pyrene were found almost exclusively in the Clinch River arm of Watts Bar Reservoir. Aroclor 1254 was only found at CRM 9.5, which is immediately downstream of Poplar Creek. Likely sources for these contaminants are the Y-12 Plant, the K-25 site, or the municipal wastewater treatment facility. Sampling and analysis for these contaminants in Poplar Creek and the Clinch River arm of Watts Bar Reservoir will continue in Phase 2 with the objective of identifying and delineating any areas requiring remediation.

Quantities of alpha-emitting radionuclides were at or near background concentrations for almost every site sampled. Concentrations of ^{234}U , ^{235}U , and ^{238}U in Poplar Creek were above background levels. Poplar Creek will be the focus of study for these contaminants during Phase 2 activities.

Fig. 3.8. Average concentrations of ^{60}Co for core sampling sites. Reaches and cores within reaches are arranged from upstream on the top to downstream on the bottom.

Phase 1 sediment sampling and analysis was conducted to:

- Obtain high quality data to confirm or modify existing data for contaminant levels in sediment.

This goal was met. Results from Phase 1 are consistent with previous studies (Olsen et al. 1992, Ashwood et al. 1986, and Oakes et al. 1982). Patterns of organics and radionuclides concentrations were similar to previous studies. Inorganic concentrations in Poplar Creek were found to be somewhat lower than previous studies indicated (Ashwood et al. 1986).

- Determine the range of contaminant concentrations present in the system.

This goal was met. We now have a working knowledge of concentrations levels of contaminants throughout the system.

- Identify contaminants of concern.

This goal was met. The Risk Analysis sections (Sect. 4.1 and 4.2) discuss this in detail. However, Phase 1 results allowed the list of contaminants of concern to be reduced significantly. Phase 1 data also strengthens level of confidence for establishing low- and high-priority contaminants.

- Establish reference (background) concentrations for contaminants.

This goal was partially met. Whereas Phase 1 sampling and analysis results provide some insight to reference concentrations, statistical comparability is not yet possible for sediment core data. Additional sampling and analysis during Phase 2 to provide the necessary reference data is the highest priority.

3.4 FISH ANALYSIS

In order to characterize the nature and extent of contamination in the principal biota of the CR/WBR system, three species of fish representing different trophic levels were sampled during the fall of 1989 and spring of 1990. Sampling included 126 bluegill sunfish, 119 channel catfish, and 43 largemouth bass. Sampling was conducted at 15 sites throughout the CR/WBR system in order to (1) bracket the entire geographic extent of this system into the sampling design and (2) focus on certain areas of the CR/WBR system such as the upper Clinch River below Melton Hill Dam and Poplar Creek where certain contaminants were known to be elevated. Results of the contaminant analysis in these three species of fish are summarized for all sites combined and individually by site.

The data validation process documented the quality of the data and determined that most of it (>95% of the metals, PCBs/pesticides, and radionuclides) met the quality standards established by the CR-ERP (see Table 2.9). One exception was the analysis of semivolatiles which was found to be deficient a large portion of the time, resulting in the rejection of nearly 50% of those results. The proportion of semivolatiles that was judged acceptable included very few detectable results, and is probably not of sufficient quantity to meet the program objectives without additional sampling.

3.4.1 Discussion of Contaminants for the System

Appendix D (Table D.1) summarizes the results of the fish analyses for all sites combined. Of the 77 organic and 13 inorganic analytes, a majority (63 organics and 7 inorganics) were not detected at any site. Of the radionuclides, only ^{137}Cs , ^{60}Co , and ^{90}Sr were detected. Four analytes were rarely detected, occurring in less than 5% of the samples. Seventeen analytes were detected in more than 5% of the samples, with only three of those (^{90}Sr , mercury, and zinc) detectable in all samples.

If either of the Aroclors was not detected, the detection limit value was included in the sum. Total PCB was calculated by summing the reported values of Aroclor 1254 and Aroclor 1260. Total chlordane was calculated by summing the concentrations of the detectable chlordane congeners. The six congeners measured in this study were α -chlordane, γ -chlordane, α -chlordene, γ -chlordene, chlordene, and oxychlordane. At least one chlordane congener and one Aroclor were detected in every sample submitted for analysis; therefore, total chlordane and total PCB were considered to be detected 100% of the time. Both total PCB and the individual Aroclors were included in further analysis because of possible point source differences for the two congeners.

Because a preliminary risk assessment was performed on historical contaminant data from the CR/WBR (Hoffman et al. 1991), a comparison of the Phase 1 data with the historical data provides a preliminary indication of any new areas of concern regarding contaminant concentrations and human health risk. Of the 24 detected analytes in Phase 1, four were not reported in the historical database (Hoffman et al. 1991): bis(2-ethylhexyl)phthalate, di-n-octylphthalate, aldrin, and 4,4'-DDE. Bis(2-ethylhexyl)phthalate and di-n-octylphthalate were detected in only one sample each. Although bis(2-ethylhexyl)phthalate was not previously detected in fish, it was detected in the sediment in past studies. According to the analytical chemists who performed the analyses, aldrin, 4,4'-DDE, and 4,4'-DDD values are suspect because of the difficulty involved in measuring these compounds when high levels of PCB and chlordane are present (personal communication, L. Wachter, May 31, 1990). Four of the analytes that were reported at detectable levels in the historical data but that were not detected in the Phase 1 sampling are antimony, chromium, silver, and thallium.

3.4.2 Discussion of Contaminants by Study Reach

A summary of the concentrations of those analytes that were detected at least once is found in Appendix D (Table D.2). For the most part, the Phase 1 maxima are lower than those from the historical data which were summarized by reach in Hoffman et al. (1991). Maximum values of mercury in fish collected in Phase 1 were higher than historical values only at reach 5, and at no sites did the maximum mercury concentration exceed the Food and Drug Administration's (FDA's) action level of 1.0 mg/kg or EPA's acceptable fish tissue concentration which is also 1.0 mg/kg. Selenium maxima exceeded historical values in reaches 2 and 3; Aroclor 1260 maxima exceeded historical values in reaches 3 and 4. Aroclor 1260 maxima exceeded the FDA action level of 2.0 mg/kg at five sites in reaches 2, 3, and 4, but average values did not exceed 1.0 mg/kg at any site or within any reach. Maximum and average values of both Aroclors 1260 and 1254 exceeded the EPA's 10^{-6} cancer risk concentration of 0.0014 mg/kg at all sites.

Because fish are mobile organisms (some species more than others), the degree to which their contaminant body burden is representative of the site in which they were collected from may be highly dependent on how long they have been residents of that area and the heterogeneity in contaminant concentrations in the surrounding environment. This along with other factors (e.g., such as size and diet) can contribute to a high degree of variability in a sample of several fish from a single site. A large amount of variation within sites is evident in both the mercury and the Aroclor 1260 results in the Phase 1 data (see standard deviations in Table D.2). The greatest variation in mercury concentrations within a site occurred in Poplar Creek at PCK 9.6 (Hartland Bridge upstream of the influence of ORR) and PCK 8.6 (mouth of EFPC). Because of the short distance between Hartland Bridge and EFPC (the source of mercury to Poplar Creek) the occurrence of a few fish with high mercury concentrations at Hartland Bridge can probably be explained by the movement of fish from the lower contaminated reaches of Poplar Creek into the upstream areas. The Hartland Bridge site was originally chosen as a potential reference site, but given this information, it should not be considered as a reference outside of the influence of releases from ORR.

The spatial distributions of total chlordane, total PCBs, mercury, and ^{137}Cs in fish are illustrated in Figs. 3.9 through 3.12, respectively. Concentrations of ^{137}Cs and mercury in fish are elevated at sites downstream of known sources of these contaminants, White Oak Creek and East Fork Poplar Creek, respectively. No such patterns of downstream gradients of chlordane or PCBs in fish are evident.

There are only a few analytes for which mean concentrations at any of the study sites exceeded those of the reference site (Norris Reservoir) by more than a factor of 2: mean mercury concentrations at two of the Poplar Creek sites and at Brashear Island, and mean copper levels at mid Watts Bar. Elevated levels of mercury at Poplar Creek and Brashear Island are not unexpected since the main source of mercury to the CR/WBR is the Y-12 facility via East Fork Poplar Creek (EFPC). Copper values are somewhat suspect for two reasons. First, elevated copper was detected at only one site in the entire system, and one fish had an abnormal concentration of 170 mg/kg. Second, copper is a well-regulated nutrient element which does not accumulate in muscle tissue even when exposure concentrations are high (Benoit 1975). Total chlordane concentrations were more than twice the reference values at nearly every site, while Aroclor 1254 and Aroclor 1260 concentrations at the affected sites ranged from 3 to 33 times higher than reference concentrations.

Using pairwise statistical tests, we compared the mean concentrations of detected analytes from each CR/WBR site with the reference site. When the data are from censored distributions, the usual analysis-of-variance techniques for comparing equality of the means of k distributions do not apply. However, if there are some *a priori* comparisons of particular sites that are of interest (e.g., each affected site with the reference site), a z -test can be used. Because this comparison was made with the means of two lognormal distributions, a statistical procedure was developed especially for this analysis (Appendix D).

The results of these site-to-reference comparisons are summarized in Table 3.1. Arsenic concentrations were significantly ($P < 0.05$) higher than concentrations at the reference site at only one location (Kingston Park). Mercury concentrations were predictably higher than those of the reference sites in Poplar Creek below the confluence with EFPC and at the first site downstream of the confluence of Poplar Creek and the Clinch River (Brashear Island).

Fig. 3.9. Arithmetic mean concentrations of total chlordane in Clinch River/Watts Bar fish tissue for all sample sites.

Fig. 3.10. Arithmetic mean concentrations of total PCBs in Clinch River/Watts Bar fish tissue for each site sampled in Phase 1.

Fig. 3.11. Arithmetic mean concentrations of mercury in Clinch River/Watts Bar fish tissue samples from each Phase 1 site.

Fig. 3.12. Arithmetic mean concentrations of ^{137}Cs in Clinch River/Watts Bar fish tissue sampled from each Phase 1 site. Fish analyzed for ^{137}Cs from reach 10 were collected from the Powell River arm of Norris Reservoir (PRM 30) as well as the Clinch River arm.

Table 3.1. Summary of study site-to-reference site comparisons for fish tissue results^a

Affected site	As	Hg	Se	Zn	Aroclor 1254	Aroclor 1260	Total PCB	Total chlordane	¹³⁷ Cs
2 Upper Melton Hill			**	**		**	**	**	*
3 Lower Melton Hill			**	**	**	**	**	**	
5 Jones Island			**		**	**	**	**	**
6 Grassy Creek	-	-	-	-		**	**		**
7 PC/Hartland Br.				**		**	**		**
8 PC/EFPC mouth		**			**	**	**	**	**
9 PC/Mitchell Br.		**			**	**	**	**	**
10 PC/K-1007		**		**	*	**	**	**	**
11 PC embayment	-	-	-	-	**	**	**	**	a
12 Brashear Island		**	**			**	**	**	**
13 Kingston Park	*				**	**	**	**	**
14 Mid Watts Bar					*	**	**		**
15 Lower Watts Bar				**	**	**	**	*	**
16 Chickamauga Res.					**	**	**		

^aAnalysis type was either z-test or confidence interval (CI) comparison as described in Appendix D (Pairwise Statistical Comparison). Final result was either (*) - concentrations from affected site significantly higher than reference site at $0.01 < P < 0.05$, (**) - concentrations from affected site significantly higher than reference site at $P < 0.01$, (a) - concentrations from study site apparently higher than reference site based on CI comparison, or (no symbol) no apparent difference in concentrations from study site and reference site. A dash (-) means no fish were analyzed for that analyte at that site. As, Hg, Se, and Zn values were from bluegill and bass only; Aroclor 1254, Aroclor 1260, total PCB, total chlordane were from catfish only; and ¹³⁷Cs was from all three species.

Selenium concentrations were higher than the reference site concentrations in main channel sites in the upper part of the CR/WBR (Upper Melton Hill to Brashear Island). Zinc concentrations at 5 of the 12 sites were higher than the those at the reference site, but no obvious spatial distribution was evident. Chlordane concentrations were higher than reference concentrations from Upper Melton Hill to Kingston Park. Total PCB concentrations at all affected sites were higher than the reference site concentrations. At nearly all sites, ¹³⁷Cs levels for all species were higher than the reference site levels and especially below the White Oak Creek confluence with the Clinch River.

3.4.3 Summary

The sampling plan for fish in Phase 1 of the CRRI was designed to characterize the extent of contamination in some major fish species in the CR/WBR system. The extent to which Phase 1 objectives as described at the beginning of this section were achieved is summarized below.

1. Data validation/verification exercises indicated that a majority of the data was acceptable for site characterization. At least 95% of the analyses performed for inorganic, PCBs/pesticides, and radionuclide analyses were judged acceptable by the validators. Analyses of semivolatiles were less reliable; nearly 50% were rejected due to validation deficiencies. Based on the acceptable data, a majority of the 90 plus analytes were never or rarely

detected. Except for aldrin, 4,4'-DDE (which are possible analytical artifacts) and the two organic compounds detected once each, the Phase 1 study did not identify any contaminants in fish tissue that were not already known from previous studies. In addition, of those analytes that were detected during Phase 1, concentrations were not markedly higher than those reported historically. The Phase 1 data set is markedly more complete and accurate than the historical data, however, and allows a much more comprehensive characterization, particularly on a spatial basis, of biota contamination in the CR/WBR system. Since the semivolatile data set was not complete, more samples should be collected for analysis of those compounds to support the finding that semivolatiles are rarely detected in fish tissue.

2. The data provided preliminary information on the spatial distribution of several contaminants in fish. Some of the contaminants (PCBs and chlordane) are widespread throughout the system and have no apparent spatial relationship to ORR. While others (mercury and ^{137}Cs) show a distinct relationship between downstream concentrations and distance from known contaminant sources. This information suggests that future sampling should be concentrated in those areas nearest ORR, Poplar Creek and the Clinch River arm of Watts Bar Reservoir.

3. Most of the data resulting from the analysis of fish samples was of sufficient quantity and quality for the screening risk assessment. The quality of the analysis of some pesticides (aldrin, DDT, and DDE) and copper are questionable, and these should be reinvestigated in future sampling. A discussion of the refinement of the contaminants of concern list will be reported in the risk assessment section of this document.

4. Fish samples collected from Norris Reservoir, the primary reference site, provided enough quality data for a preliminary characterization of that area. A pair-wise statistical analysis that compared the Norris Reservoir results to those of the study sites found that mercury, selenium, zinc, total chlordane, PCBs, and ^{137}Cs occur at some of the CR/WBR sites at significantly higher concentrations than at the reference site. Samples were also collected from a site in Reach 13 (Poplar Creek upstream of the influence of ORR, PCK 9.8) for evaluation as an additional reference site. After evaluation of the data, the usefulness of this site as a reference site for fish is questionable. Although the site is upstream of any ORR contamination sources, it is close enough to the mouth of EFPC that fish could move into the upstream site after residing in contaminated areas downstream. Other reference areas should be sampled in Phase 2 to better characterize background concentrations of contaminants of concern.

4. SCREENING RISK ASSESSMENT

4.1 SCREENING ASSESSMENT FOR HUMAN EXPOSURE

4.1.1 Approach

The objective of the screening assessment, which is based on human health risk methodology, is to utilize data collected in Phase 1 of the CRRI to (1) identify reaches and contaminants that are high priority for further evaluation and (2) identify reaches and contaminants that have a low priority for further investigation.

The screening approaches used in this assessment are similar to those employed in Hoffman et al. (1991), Blaylock et al. (1992a), and Blaylock et al. (1992b) reports. The first approach is the use of conservatively biased calculations to identify contaminants, exposure pathways, and reaches that have a low priority for further investigation. It is unlikely that these conservative calculations will underestimate the potential maximum exposures to an individual residing in the vicinity of the CR/WBR, which receives releases from the ORR. The second approach uses nonconservative calculations and assumptions to identify contaminants, pathways, and reaches that have a high priority for further investigation. These nonconservative risk estimates are more realistic than the conservative estimates and should not substantially overestimate maximum exposures to local population subgroups. Results of this screening assessment do not constitute a baseline risk assessment and are not intended to address issues regarding compliance with regulatory limits. The purpose of this screening exercise, along with the results obtained in Hoffman et al. (1991), is to help guide the development of the Phase 2 sampling plan, which will focus the collection of data on contaminants of concern. A future baseline risk assessment will be completed when data from the Phase 2 field sampling are available.

4.1.2 Exposure Pathways

The screening assessment was conducted on data from five reaches on the Clinch and Tennessee River system, which receives contaminant releases from the ORR (Fig. 2.4) and two reference reaches—Norris Reservoir and Poplar Creek above the confluence with East Fork Poplar Creek. Conservative and nonconservative screening analyses are performed for selected media in both primary and secondary exposure pathways.

Primary exposure pathways are the most realistic and currently the most likely to be encountered by a potential receptor. The primary pathways represent exposure to three media: fish, near-shore surface sediment, and surface water. The primary exposure pathways are: (1) ingestion of fish, (2) external exposure to radionuclides in surface sediment, (3) inadvertent ingestion of surface sediment, (4) inhalation of contaminated air resulting from resuspension of surface sediment, and (5) ingestion of surface water.

Secondary pathways are less likely scenarios and represent potential or future pathways of exposure. The scenario for secondary exposure pathways assumes that sediment from the Clinch and Tennessee River system is dredged and that the dredged material is placed on land where it supports agricultural activities including growing pasture as forage for cows and garden vegetables for human consumption. The pasture-to-cow pathway results in human

exposure via consumption of milk and meat. Direct analytical measurements of contaminants are not available for the dredging scenario except for the concentrations of contaminants in the dredged sediments. Environmental pathway models that incorporate generic transfer coefficients from sediment are used to predict the concentrations of contaminants in the other exposure media.

Model-predicted concentrations of contaminants have a much higher degree of uncertainty than concentrations based on actual measurements. An irrigation scenario was not considered in this screening analysis because the number of Phase 1 water samples was too limited to reliably extrapolate from water to other media. For the same reason, the consumption of surface water by cows was not included in the cow-meat-milk pathway. Pathways for the dredging scenario include (1) external exposure to dredged sediments, (2) inadvertent ingestion of dredged sediment, (3) inhalation of air contaminated by resuspension of sediments, and (4) ingestion of agricultural products (meat, milk, and vegetables).

4.1.3 Usage and Exposure Factors

Usage and exposure factors for conservative and nonconservative screening are listed in Appendix E, Table E.2. These factors are similar to the ones used by Hoffman et al. (1991) and Blaylock et al. (1992a, 1992b); however, various intake rates, exposure times, toxicity values, and other parameters for both carcinogens and noncarcinogens have been modified since the publication of these reports. For example, EPA has reconsidered the maximum exposure duration and now recommends a residential exposure period of 350 d/year for 30 years instead of the previous recommendation of 365 d/year for 70 years (OSWER 1991).

4.1.4 Data Bases

All data used in this screening analysis were collected in the Phase 1 sampling effort except for the surface water. Because of the limited number of Phase 1 water samples, data from source monitoring sites were also included in the data base (Table E.1). The source monitoring program collects and analyzes water from discharge points, before the effluent readies the surface water body, for NPDES and DOE Order 5400.1 permits. See Appendices B, C, and D for summaries of the Phase 1 data.

More than 500 near-shore grab samples of surface-sediment (Sect. 2) were taken to aid several agencies, including the regulators, TVA, and the U.S. Army Corps of Engineers, in evaluating permit applications for activities that would disturb near-shore sediment. The majority of these samples were taken in shallow water where the sediment would be exposed during winter months when the water level in Watts Bar Reservoir is low. These samples were analyzed for gamma-emitting radionuclides only, primarily ^{60}Co and ^{137}Cs . Data from these samples were used in this screening analysis for primary sediment exposure pathways (i.e., external exposure, inadvertent ingestion of sediment, and inhalation of resuspended material).

Currently there is not a realistic direct pathway for human exposure to deep sediment where the highest concentrations of contaminants have been observed (Olsen et al. 1992). However, a dredging scenario was included in the screening analysis to consider potential exposure pathways if the sediment should be placed upon shore. This scenario would insure

that elevated concentrations of contaminants in deep sediments are considered in the screening analysis (Sect. 3.3). Data from the Phase 1 sediment cores that were used in this analysis are summarized in Appendix C.

The data were divided into two categories—detected and nondetected contaminants. Contaminants for which at least one measurement was above the level of detection for a medium and reach were included in the detected data base. Contaminants for which all measurements in a reach and medium were below the detection limit were included in the nondetected data base. For conservative screening, the upper 95% confidence limit of the arithmetic mean (UCL) for each contaminant in the detectable data set was computed. The UCL was compared with the actual maximum detected value, and the smaller of the two values was used in the risk calculations, as directed in *Risk Assessment Guidance for Superfund* (RAGS) (EPA 1989). For nonconservative screening, the geometric mean of each contaminant in the detected data set and the lowest detection limit in the nondetected data set were used as the contaminant concentration in the environmental medium.

4.1.5 Screening Criteria

4.1.5.1 Screening indices

Screening indices (SIs) for carcinogens and noncarcinogens are calculated for contaminants of potential concern to human health. These indices are based upon EPA-approved or -suggested cancer slope factors and toxicity reference doses (RfDs). The slope factor is based upon an estimate of the lifetime risk of an incremental cancer incidence per unit of exposure. The RfD is an exposure level (threshold) below which no adverse health effects should occur. Tables E.3 and E.5 list the slope factors and RfDs for contaminants evaluated in this report.

The screening index for carcinogens is calculated by multiplying an estimate of exposure to a contaminant via external exposure, ingestion, or inhalation by an EPA slope factor. This index indicates the potential lifetime risk of developing additional cancer (EPA 1992, 1990a) above and beyond the normal incidence of cancer. The screening index for noncarcinogens is the ratio of the estimated daily ingestion or inhalation of a contaminant divided by the contaminant-specific RfD.

To estimate the potential risk from all contaminants in a particular reach, the screening indices are summed for all contaminants in the reach. This summation has been termed a "total reach screening index" throughout this report. Summation is conducted separately for carcinogens and noncarcinogens.

4.1.5.2 Screening criteria for carcinogens

Conservative and nonconservative screening criteria are applied to the EPA action level (*Federal Register* March 8, 1990) to categorize carcinogenic contaminants as low priority, high priority, or requiring further investigation before designating high or low priority. *Low priority* indicates that a carcinogen is of little concern for further investigation; *high priority* requires either immediate consideration for remedial action or further analysis; and contaminants identified as neither low nor high priority must be evaluated further before this determination can be made. Figure 4.1 summarizes the screening criteria for carcinogens used in this report.

**Conservative Estimate
of Exposure** **Nonconservative Estimate
of Exposure**

Screening index (SI) = exposure multiplied by a lifetime cancer slope factor

Fig. 4.1. Screening criteria for carcinogens.

For conservative screening, carcinogens having screening indices $<10^{-6}$ are considered low priority contaminants for further consideration. Carcinogens with screening indices between 10^{-6} and 10^{-4} require further investigation before either taking action or designating these substances as high or low priority. Carcinogens with conservative screening indices $\geq 10^{-4}$ require further investigation before taking action.

For nonconservative screening of carcinogens, substances having screening indices $\geq 10^{-4}$ are high priority carcinogens that require immediate consideration for remedial action. Carcinogens with screening indices between 10^{-4} and 10^{-6} require further investigation before action is taken (i.e., re-examination of the data base, checking of parameter values, or recalculation of screening indices, etc.). Because nonconservative screening employs parameter values that should not greatly overestimate maximum exposures to a contaminant (and in some cases could underestimate maximum exposure), nonconservative screening is not used to identify contaminants with low priority for further consideration.

4.1.5.3 Screening criteria for noncarcinogens

The screening criteria used for noncarcinogens are summarized in Fig. 4.2. For conservative screening, contaminants with screening indices <0.1 are low priority for further consideration, and contaminants with screening indices between 0.1 and 1.0 require further investigation before designating these substances as high or low priority. Noncarcinogens with conservative screening indices ≥ 1.0 require further investigation before action is taken.

For nonconservative screening of noncarcinogens, contaminants with screening indices ≥ 1.0 are high priority substances requiring immediate consideration for remedial action, and contaminants with screening indices between 1.0 and 0.1 require further investigation before taking action. Nonconservative screening indices <0.1 are not used to designate low priority contaminants because nonconservative screening employs parameter values that, under some circumstances, may underestimate maximum exposures to a contaminant.

4.1.6 Contaminant Screening Results for Primary Exposure Pathways

Screening results for individual contaminants for each primary exposure pathway and each reach are given in Appendix E, Tables E.8 through E.12. The pathways considered to be primary pathways for this screening exercise are: (1) ingestion of fish, (2) external exposure to gamma-emitting radionuclides present in surface sediment, (3) inadvertent ingestion of surface sediment, (4) inhalation of contaminated air (via either dust resuspension, volatilization, or vaporization of contaminants), and (5) ingestion of surface water. Tables 4.1 through 4.5 summarize the results for carcinogens and noncarcinogens in each exposure pathway for both the conservative and nonconservative screening approaches. These tables identify the high priority contaminants, low priority contaminants, and contaminants requiring further investigation; they also show the reaches where no adverse health effects are expected and those where no data were available.

4.1.6.1 Fish ingestion pathway

Carcinogens. Listed in Table 4.1 are the results of the conservative and nonconservative screening analyses for carcinogens in the fish ingestion pathway. Nonconservative screening identified the PCB Aroclor 1260 in all reaches except reach 1 (Melton Hill Reservoir) and

Screening index (SI) = exposure divided by reference dose factor (RfD)

Fig. 4.2. Screening criteria for noncarcinogens.

Table 4.1. Summary of adult screening results for carcinogens in the fish ingestion pathway

Carcinogen	Reach						
	1	2	3	4	5	10	13
Organics							
4,4'-DDE							
4,4'-DDT	N		N			N	N
Aldrin	N		N			N	
Aroclor 1254						N	
Aroclor 1260		X	X	X	X		X
Bis(2-ethylhexyl) phthalate	N	N		N	S	S	S
Chlordane							
Inorganics							
Arsenic							
Beryllium		N		N			N
Radionuclides							
⁶⁰ Co		N		N	N		
¹³⁷ Cs							
⁹⁰ Sr

Carcinogens having nonconservative screening indices $> 10^{-4}$ that require immediate consideration.

Carcinogens with conservative or nonconservative screening indices between 10^{-6} and 10^{-4} that further investigation before either taking action or designating as low priority.

Carcinogens having conservative screening indices $< 10^{-6}$ and can be designated as low priority.

N

Nondetects - contaminant concentrations were below detection limits.

S

No data available.

Table 4.2. Summary of screening results for noncarcinogens in the fish ingestion pathways

Contaminant	Reach						
	1	2	3	4	5	10	13
Organics							
4,4'-DDT	N		N			N	N
Aldrin	N		N			N	
Bis(2-ethylhexyl phthalate)	N	N		N	S	S	S
Chlordane							
Di-n-octylphthalate	N	N		N	S	S	S
Inorganics							
Arsenic							
Beryllium		N		N			N
Copper	N	N	N	N		N	N
Mercury							
Selenium					N		N
Zinc							

X	Contaminants with adult nonconservative screening indices ≥ 1.0 that require immediate consideration.
	Contaminants with child conservative screening indices between 0.1 and 1.0 that require further investigation before either taking action or designating as low priority.
	Contaminants having child conservative screening indices < 0.1 and can be designated as low priority.
N	Nondetects - contaminant concentrations were below detection limits.
S	No data available.

Table 4.3. Summary of adult screening results for radionuclides in near-shore surface sediment exposure

	Reach							
	1	2	3	4	5	10	13	18
External Exposure								
⁶⁰ Co						S	S	
¹³⁷ Cs						S	S	
Ingestion Exposure								
⁶⁰ Co						S	S	
¹³⁷ Cs						S	S	
Inhalation Pathway								
⁶⁰ Co						S	S	
¹³⁷ Cs						S	S

Carcinogens with nonconservative screening indices $\geq 10^{-4}$ that require immediate consideration.

Carcinogens with screening indices between 10^{-6} and 10^{-4} that require further investigation before designating as low priority.

Carcinogens having conservative screening indices $< 10^{-6}$ and can be designated as low priority.

N

Nondetects - contaminant concentrations were below detection limits.

S

No data available.

Table 4.4. Summary of adult screening results for carcinogens in the water ingestion pathway

Contaminant	Reach						
	1	2	3	4	5	10	13
Organics							
1,2-Dichloroethane	S	N		N	S	S	N
1,2-Dichloropropane	S		N	N	S	S	N
1,4-Dichlorobenzene	N	N		N	N	N	N
Anthracene	N	N		N	N	N	N
Aroclor 1254	N	N		N	N	N	S
Benzene	S	N		N	S	S	N
Benzo(a)anthracene	N	N		N	N	N	N
Bis(2-ethylhexyl)phthalate	N	N		N	N		N
Bromoform	S	N		N	S	S	N
Chlorodibromomethane	S	N		N	S	S	N
Chloroform	S			N	S	S	N
Chrysene	N	N		N	N	N	N
Dichlorobromomethane	S	N		N	S	S	N
1,1-Dichloroethylene							
Fluoranthene	N	N		N	N	N	N
Methylene Chloride	S			N	S	S	N
N-nitro-di-n-propylamine	N	N		N	N	N	N
Phenanthrene	N	N		N	N	N	N
Pyrene	N	N		N	N	N	N
Tetrachloroethylene	S			N	S	S	N
Trans-1,3-dichloropropene	S	N		N	S	S	N
Trichloroethylene	S	N		N	S	S	N
Vinyl Chloride	S	N		N	S	S	N
Inorganics							
Arsenic		N		N	N	N	N
Beryllium		N		N	N	N	N

Table 4.4 (continued)

Contaminant	Reach						
	1	2	3	4	5	10	13
Radionuclides							
⁶⁰ Co			N		N	N	N
¹³⁷ Cs						N	
³ H	S		S		S	S	S
²³⁷ Np	S			S	S	S	S
¹⁹¹ Os	S		S	S	S	S	S
²³⁴ Pa	S				S	S	S
²³⁸ Pu	S				S	S	S
²³⁹ Pu		S			S	S	S
⁹⁰ Sr						S	S
⁹⁹ Tc	S				S	S	
²²⁸ Th	S		X		S	S	
²³⁴ Th	S				S	S	
²³⁴ U					S	S	S
²³⁵ U					S	S	S
²³⁶ U	S	S			S	S	S
²³⁸ U					S	S	S

Carcinogens having nonconservative screening indices $> 10^{-4}$ that require immediate consideration.

Carcinogens with conservative or nonconservative screening indices between 10^{-6} and 10^{-4} that further investigation before designating as low priority.

Carcinogens having conservative screening indices $< 10^{-6}$ and can be designated as low priority.

N Nondetects - contaminant concentrations were below detection limits.

S No data available.

Table 4.5. Summary of screening results for noncarcinogens in the water ingestion pathway

Contaminant	Reach						
	1	2	3	4	5	10	13
Organics							
1,1-Dichloroethane	S	N		N	S	S	N
1,1,1-Trichloroethane	S	N		N	S	S	N
2,4-Dichlorophenol	N	N		N	N	N	N
Acetone	S		N	S	S	S	S
Anthracene	N	N		N	N	N	N
Bis(2-ethylhexyl)phthalate	N	N		N	N		N
Bromoform	S	N		N	S	S	N
Butylbenzylphthalate	N	N		N	N	N	N
Carbon Disulfide	S		S	S	S	S	S
Chlorobenzene	S	N		N	S	S	N
Chlorodibromomethane	S	N		N	S	S	N
Chloroform	S			N	S	S	N
Di-n-butylphthalate	N	N		N	N	N	N
Dichlorobromomethane	S	N		N	S	S	N
Diethylphthalate	N	N		N	N	N	N
Dimethylphthalate	N	N		N	N	N	N
Ethylbenzene	S	N		N	S	S	N
Fluoranthene	N	N		N	N	N	N
Methylene Chloride	S			N	S	S	N
Phenol	N				N	N	N
Pyrene	N	N		N	N	N	N
Tetrachloroethylene	S			N	S	S	N
Toluene	S			N	S	S	N
Trans-1,2-Dichloroethene	S	N		N	S	S	N
Trans-1,3-dichloropropene	S	N		N	S	S	N
Xylene	S		N	S	S	S	S

Table 4.5 (continued)

Contaminant	Reach						
	1	2	3	4	5	10	13
Inorganics							
Antimony		N	N	N	N	N	N
Arsenic		N		N	N	N	N
Barium					S	S	S
Beryllium		N		N	N	N	N
Boron	N	N			S	S	S
Cadmium		N		N	N	N	N
Chromium		N			N	N	
Copper					N	N	N
Cyanide	S				S	S	
Manganese					S	S	
Mercury		N			N	N	N
Molybdenum	N	N		N	S	S	
Nickel					N	N	
Selenium				N	N	N	N
Silver	N	N		N	N	N	N
Strontium			S	S	S	S	S
Tin	N	N		N	S	S	S
Uranium	S				S	S	N
Vanadium		N	N	S	S	S	S
Zinc					N	N

Noncarcinogens having adult nonconservative screening indices ≥ 1.0 that require immediate consideration.

Noncarcinogens with child conservative screening indices between 0.1 and 1.0 that require further investigation before either taking action or designating as low priority.

Noncarcinogens having child conservative screening indices < 0.1 and can be designated as low priority.

N Nondetects - contaminant concentrations were below detection limits.

S No data available.

reach 10 (Norris Reservoir) as a high priority contaminant requiring immediate consideration for remedial action. In addition, Aroclor 1254 was identified as requiring further study in all reaches except reach 10. In a previous screening analysis of historical data (Hoffman et al. 1991), PCBs were identified as a potentially high priority contaminant (screening index between 10^{-4} and 10^{-3}). Other investigations have shown that PCBs are a common pollutant in East Tennessee rivers and reservoirs. Although the ORR is a known source of PCBs, it is not the only one contributing PCBs to the Clinch River/Watts Bar Reservoir system.

Several contaminants were identified as requiring further investigation (screening indices between 10^{-4} and 10^{-6}) before action is taken or they are designated as low priority (Table 4.1). Arsenic and chlordane in all reaches, including reference reach 10 (Norris Reservoir), were identified as requiring further investigation. These contaminants were identified in Hoffman et al. (1991) as potentially high priority contaminants. Arsenic is a common pollutant resulting from industrial operations and fossil fuel use in East Tennessee, especially metropolitan areas. A special study is being conducted under the sponsorship of the CRRRI to document the concentration of arsenic in water and sediment in areas receiving effluents from the ORR (Byrd 1991). Beryllium was identified in all reaches except 2, 4, and 13, and ^{137}Cs was identified in reaches 2 and 4 as requiring further study.

Several pesticides found in fish (4,4'-DDE, 4,4'-DDT, and Aldrin) were identified as requiring further investigation. These are common agricultural pesticides that were identified in only a few reaches but will require further investigation; however, the occurrence of these contaminants in fish may be the result of analytical procedures.

Noncarcinogens. Exposure parameters for a child were used to identify the low priority noncarcinogens in the fish ingestion pathway because the child scenario is more conservative than the adult and would eliminate fewer contaminants. Results of both conservative (child) and nonconservative (adult) screening analyses for noncarcinogens in the fish ingestion pathway are summarized in Table 4.2. None of the noncarcinogens were identified by nonconservative screening as high priority contaminants that would require immediate consideration for remedial action. Also, none of the sums of the conservative screening indices for the reaches were <0.1 , including reference reach 10 (Norris Reservoir) (see Tables E.9 and E.12 in Appendix E), thus, no reach could be eliminated from further consideration. Contaminants that were classified as requiring further investigation in most reaches are chlordane, arsenic, mercury, and selenium. These contaminants are ubiquitous in East Tennessee aquatic systems, and, with the exception of mercury, which was released from the Y-12 Plant in large quantities in the past, are common industrial pollutants and not unique to the ORR.

4.1.6.2 Near-shore surface sediments

Results of conservative and nonconservative screening analyses for the near-shore surface sediment for the ingestion, inhalation, and exposure pathways are given in Table 4.3. Nonconservative screening indices were $<10^{-4}$ for all pathways and reaches. Conservative screening indices for the ingestion and inhalation pathways were $<10^{-6}$ for all reaches and can be assigned low priority for consideration for further investigation. Therefore, ^{60}Co and ^{137}Cs , the only contaminants for which the near-shore surface sediment samples were analyzed, are assigned low priority for further consideration in ingestion and inhalation pathways for all reaches.

For the external exposure pathway, ^{60}Co in reaches 1-4 and ^{137}Cs in reaches 1-5 and 18 were identified as contaminants with screening indices of $>10^{-6}$. These radionuclides require further investigation before action is taken or they are assigned low priority. In most reaches, the concentrations of ^{60}Co and ^{137}Cs were <1 and 4 pCi/g, respectively, in the near-shore surface sediment. Three samples had concentrations >10 pCi/g (Section 3.3.1.1); however, these samples came from relatively deep water.

4.1.6.3 Water ingestion pathway

Carcinogens. Because of the limited number of Phase 1 water samples and the use of source monitoring data, which included NPDES data, the screening results for water should be used primarily to guide future sampling and analysis of the surface water. Although source monitoring data are useful in identifying contaminants that are released into the surface water, it is unlikely that these data represent the actual contaminant concentrations in the receiving body of water. Results of the screening analysis for the ingestion of carcinogens in surface water are summarized in Table 4.4. This table shows that reach 3 (Poplar Creek), which receives effluents from the K-25 Site, East Fork Poplar Creek, and the City of Oak Ridge west-end sewage treatment plant, has a large number of contaminants that require consideration for further investigation. Thorium-228, a daughter of ^{232}U and naturally occurring ^{232}Th , was reported in reach 3 at concentrations that produced a screening index $>10^{-3}$. By the current screening criteria, ^{228}Th would be classified as a high priority contaminant requiring immediate consideration for remedial action; however, the ^{228}Th was measured at only surface water monitoring sites, and these measurements are representative of a point source rather than of the downstream surface water. In the other six reaches shown in Table 4.4, either most of the organic and inorganic contaminants in water were below detection limits, or the contaminants were not measured.

In addition to reach 3, reaches 2 and 4, which are immediately downstream from ORNL and the K-25 Site, respectively, have concentrations of radionuclides that require further investigation. Many of these radionuclides are isotopes of uranium or their daughter products that occur in nature. In reach 2, ^{191}Os was detected once but probably will not be detected again because it is an isotope that is used in nuclear medicine and is not routinely released from ORNL. The tritium (^3H) concentration for reach 2 is from surface water monitoring data taken at White Oak Dam and is not representative of water in the Clinch River. Although ^{99}Tc is identified in reaches 3 and 13 as being a contaminant of concern, the analytical methodology may be questionable when applied to environmental samples. The ^{99}Tc data probably represent gross beta activity rather than the actual activity of ^{99}Tc .

Noncarcinogens. Only two organic noncarcinogens in water in reach 3 are identified in Table 4.5 as contaminants that would require further consideration for investigation. However, a relatively large number of inorganic noncarcinogens in reaches 1 and 3 were identified as requiring further investigation. Because the water data are composed of a limited number of Phase 1 samples and are supplemented by NPDES monitoring data that represent point sources of contaminants, results of the screening analyses performed on these limited data are inconclusive. Unless a methodology can be developed to convert the available monitoring data to concentrations representative of the concentrations in the receiving water, additional data for all contaminants in water will be needed for future assessment of the water ingestion pathway.

4.1.7 Dredging Scenario

Conservative screening was used in the dredging scenario to identify contaminants found in buried sediments that require further investigation before designation as high priority contaminants, i.e., carcinogens with screening indices between 10^{-4} and 10^{-6} and noncarcinogens with a screening indices between 1.0 and 0.1. Screening indices for the carcinogens are given in Table E.13 and for the noncarcinogens in Table E.14. Screening indices for some inorganics in all pathways were $>10^{-6}$, which indicates that they require further investigation.

Carcinogens and noncarcinogens in the direct and secondary exposure pathways that had screening indices $>10^{-6}$ or >0.1 , respectively, are summarized in Table 4.6. Reach 3 had the largest number of organic contaminants that require further investigation in both the direct and secondary exposure pathways. Arsenic, beryllium, and chromium in all reaches were identified as requiring further investigation. Mercury and nickel were identified in the screening analysis of the secondary pathways as requiring further investigation in all pathways. Cobalt-60 and ^{137}Cs were identified in the screening analysis of the direct pathways as requiring further investigation in all reaches. Most of the inorganics in the direct and indirect exposure pathways listed in Table 4.6 require further investigation for reach 5 (Watts Bar Reservoir). Further investigation should be conducted to determine the original source of the metals that have relatively high concentrations in reach 5 sediment.

Although no contaminant was identified in the dredging scenario as a high priority contaminant requiring immediate consideration, a relatively large number of organic and inorganic contaminants were identified in most reaches as requiring further investigation. Therefore, dredging should be strictly limited, if not prohibited, in these reaches.

Conservative screening was used in the dredging scenario to identify low priority contaminants, i.e., carcinogens with a screening index of $<10^{-6}$ and noncarcinogens with a screening index of <0.1 . Screening results for carcinogens in the secondary pathways are summarized in Table E.6 in Appendix E. Eleven organic and five inorganic carcinogens had screening indices that were $<10^{-6}$ in at least one pathway; but only bis(2-ethylhexyl)phthalate and cadmium can be designated low priority for all reaches. The remaining carcinogens should be considered candidates for future investigation. Ten radionuclides had screening indices that were $<10^{-6}$ in one or more pathways and five of these, ^{241}Am , $^{243,244}\text{Cm}$, ^{238}Pu , $^{239,240}\text{Pu}$, and ^{235}U , can be eliminated from further consideration based on the dredging scenario.

A summary of the noncarcinogens in the dredging scenario that were identified as low priority in one or more pathways is presented in Table E.7 in Appendix E. All nine of the organics had screening indices that were <0.1 and can be designated low priority for all pathways and reaches. None of the ten inorganics can be designated low priority because their screening indices for one or more pathways was >0.1 . Total screening indices for the inhalation pathway for organic and inorganic noncarcinogens were <0.1 for all reaches, indicating that this pathway can be assigned low priority; however, only mercury has an inhalation RfD (Table E.14 in Appendix E). Total screening indices for the ingestion of meat, milk, and vegetable pathways were >0.1 and cannot be considered low priority.

Table 4.6. Summary by reach of contaminants in the dredging scenario identified by conservative screening as requiring further investigation before taking action or designating as low priority

Contaminants	Reaches					
	1	2	3	4	5	10
Sediment exposure via direct pathways (Ingestion, inhalation, and external exposure)						
Organic						
Anthracene			X			
Aroclor-1254		X	X			
Aroclor-1260		X				
Benzo(a)pyrene		X	X			
Fluoranthene	X	X	X	X		X
N-Nitro-di-n-propylamine				X		
Phenanthrene		X	X	X		X
Pyrene	X	X	X	X		X
Toxaphene		X				
Inorganic						
Arsenic	X	X	X	X	X	X
Beryllium	X	X	X	X	X	X
Chromium	X		X	X	X	X
Radionuclides						
⁶⁰ Co	X	X	X	X	X	
¹³⁷ Cs	X	X	X	X	X	X
²³⁸ U			X			
Additional contaminants identified from sediment exposure via secondary pathways (Ingestion of milk, meat, and vegetables produced on sediment)						
Organic						
Aroclor-1254				X		
4,4'-DDD	X					
4,4'-DDT	X					
Benzo(a)anthracene	X	X	X			
Benzo(b)fluoranthene		X	X			

Table 4.6 (continued)

	Reaches					
Contaminants	1	2	3	4	5	10
Organic (continued)						
Benzo(k)fluoranthene	X		X			
Chrysene						X
Indeno(1,2,2-cd)pyrene		X				
Inorganic						
Cadmium	X	X	X	X	X	X
Copper	X	X	X	X	X	X
Mercury	X	X	X	X	X	X
Nickel	X	X	X	X	X	X
Selenium	X	X	X	X		
Silver			X	X	X	
Thallium			X			
Uranium			X			
Zinc	X	X	X	X	X	X
Radionuclides						
⁹⁰ Sr		X	X	X	X	X
²³⁴ U			X			

4.1.8 Comparison of the Phase 1 Screening Analysis with a Previous Screening Analysis of Historical Data

The results of the screening analysis of the historical data base (Hoffman et al. 1991) were helpful in identifying areas and contaminants that required immediate consideration for remedial action. The screening analysis also helped guide the Phase 1 sampling, which was designed to confirm the areas and contaminants that will require more comprehensive sampling in Phase 2. White Oak Creek Embayment, previously included in reach 2, was identified in Hoffman et al. (1991) as an area requiring immediate attention. Results of intense sampling efforts by the CRRI staff led to the implementation of remedial measures to abate the release of contaminants from the embayment. The White Oak Creek Embayment remediation measures have been completed (Leslie and Kimmel 1992), and this area is no longer considered part of the CRRI. As a result of this remediation, ^{137}Cs , which was a high priority contaminant in the previous screening assessment, has been given a lower priority in reach 2 in the current screening assessment.

Generally, the results of the Phase 1 screening analysis are in agreement with results obtained from the earlier human health screening efforts. The screening methodologies used in the two reports are similar; however, modification of several exposure parameters, changes in some EPA-approved risk assessment factors and criteria, and the use of maximum instead of mean concentrations for some media have contributed to the differences in the screening results for the two reports. The identification of ^{228}Th as a high priority contaminant in water from Poplar Creek (reach 3) in the current screening assessments was the most notable difference between the two screening analyses. This result is probably biased and not representative of the entire reach because of the use of source monitoring data to supplement the Phase 1 water data. The source monitoring data should be investigated to document the concentrations of ^{228}Th reported in these data.

PCBs were identified in the fish ingestion pathway as potentially high priority pollutants (screening index between 10^{-4} and 10^{-3}) in Hoffman et al. (1991). The PCB Aroclor 1260 was identified in the current screening assessment as a high priority contaminant in fish (screening index $\geq 10^{-4}$) requiring immediate consideration for remedial action. A large number of inorganic and organic contaminants were identified as potentially high priority contaminants in the secondary pathways in the dredging scenario in Hoffman et al. (1991). Similar results were obtained for the secondary pathways in the dredging scenario in the current screening assessment, but the number of problem contaminants was smaller. For the dredging scenario, the maximum concentrations for the contaminants found in the sediment cores were used as the sediment concentration in Hoffman et al. (1991) whereas the mean concentration, which is a more realistic value, was used in the current screening assessment. Because of improved analytical procedures, more pesticides and polycyclic aromatic hydrocarbons were detected in the Phase 1 samples than were reported for samples in the historical data base. Several of the contaminants that were not reported in the historical data base are classified in the current screening assessment as requiring further evaluation. However, these contaminants are found in most streams that receive industrial and domestic waste and are not unique to streams that receive effluents from ORR.

4.1.9 Nondetected Contaminants

The nondetected contaminant data base contained several times as many contaminants as the detected contaminant data base. The screening analysis would not be complete without including the nondetected contaminants; however, caution should be exercised in interpreting the results because of the uncertainty associated with the reported contaminant concentrations.

Only the nonconservative screening approach was employed for the nondetected contaminants. This approach was used to identify contaminants that have a high priority for action to improve (lower) the detection limits (DLs) or, alternatively, those for which source-term data are needed to determine whether even extremely small quantities of these contaminants have been released from the ORR in the past. Results of the adult nonconservative screening for carcinogens and noncarcinogens in the nondetectable data base are given in Tables E.15-E.18 in Appendix E. Screening indices for individual contaminants are listed according to exposure pathway and reach. Concentrations used in the screening calculations were the minimum DL recorded for a medium across each reach. Screening indices were not summed for contaminant type, pathway, or reach because the uncertainty resulting from using DLs would make the sums overly conservative.

4.1.9.1 Carcinogens

For nondetected carcinogens, a nonconservative screening index $\geq 10^{-4}$ indicated the need to improve the DL of the contaminant for future samples. For organics in fish, the DLs for the same 11 contaminants in reaches 1, 2, and 4 produced screening indices $\geq 10^{-4}$, and four of these contaminants had screening indices of $> 10^{-4}$ in reach 3 (Table 4.7). Screening indices for six organic contaminants in water indicated a need for lower detection limits. No radionuclides in water or fish and no carcinogens in the deep core sediments had DLs that produced high priority screening indices (Tables E.15 and E.17 in Appendix E). Near-shore surface sediment grab samples had no problems with DLs.

4.1.9.2 Noncarcinogens

A noncarcinogen in the nondetected data base was assigned a high priority for improvement in analytical methodology if the screening index was ≥ 1.0 . In the fish and water exposure pathways, the detection limit for 4,6-dinitro-2-methylphenol produced screening indices > 1.0 in most reaches (Table 4.7). None of the DLs for noncarcinogens in sediment produced high priority screening indices (Table E.18 in Appendix E).

4.1.10 Contaminants Having No Available Toxicity Values

The number of contaminants that can be quantitatively evaluated in this risk assessment was limited by the availability of chemical-specific, EPA-approved dose/response information. Contaminants for which no slope or reference dose factors are currently available and were therefore not included in the screening assessment are listed in Table E.19 of Appendix E. Of these contaminants, some are essential nutrients that should have little or no toxic effects at the reported concentrations.

Table 4.7. Contaminants for which the use of detection limits produced screening indices $>10^{-4}$ for carcinogens or >1.0 for noncarcinogens by nonconservative screening

Contaminants		Reaches
Carcinogens	Fish pathway	Water pathway
Acenaphthene	1, 2, 3, 4	1,2,3,4,5,10,13
Acenaphthylene	1, 2, 3, 4	1,2,3,4,5,10,13
Anthracene	1, 2, 4	
Benzo(a)pyrene	1, 2, 4	
Benzidine		2,3,4,13
Benzo(g,h,i)perylene	1, 2, 4	
Dibenz(a,h)anthracene	1, 2, 3, 4	1,2,3,4,5,10,13
Fluoranthene	1, 2, 4	
Fluorene	1, 2, 3, 4	1,2,3,4,5,10,13
N-Nitroso-di-n-propylamine	1, 2, 4	
N-Nitrosodimethylamine		2,3,4,13
Phenanthrene	1, 2, 4	
Pyrene	1, 2, 4	
Noncarcinogens		
4,6-Dinitro-2-methylphenol	1,2,4,5,13	1,2,3,4,5,10,13

4.1.11 Special Case—Lead

Although an action level of 0.015 mg/L for lead has been adopted by EPA (EPA 1991), specific toxicity values are not presently available for risk assessment; instead, an EPA Uptake/Biokinetic Model (Version 0.50) designated for a target population of children (0 to 6 years old) was used to evaluate the risk from lead. The critical value for lead in the blood of young children is 10.0 $\mu\text{g/dL}$. Lead concentrations in the various media for the conservative exposure scenario were input to the model for each reach, and the percentage of the target population estimated to have lead levels greater than the critical value was calculated. EPA Region IV considers an area to be at risk if more than 5% of the population exceeds the critical value. Only reach 1 (Melton Hill Reservoir), with 6.4% of the population calculated to exceed the critical value, should be considered of marginal concern for lead exposure. Water concentrations appear to have the most influence on the model results and the model used the direct ingestion of surface water for drinking purposes.

4.1.12 High Priority Contaminants and Contaminants Requiring Further Investigation

The PCB Aroclor 1260 was identified as a high priority contaminant in the fish ingestion pathway (Section 4.1.6.1). Total PCB concentrations (i.e., Aroclor 1260, Aroclor 1254, and other formulations) should be considered when determining risk to human health; therefore, PCBs in the Clinch River/Watts Bar Reservoir should be considered high priority contaminants. The average concentration for PCBs in fish in the reaches of the Clinch and Tennessee River system used in this screening analysis (Appendix E) is less than the FDA action level of 2.0 ppm (mg/kg) in fish tissue.

Thorium-228 was identified as a high priority contaminant in the water ingestion pathway in reach 3, Poplar Creek embayment. Because of the limited water data and because the ^{228}Th

concentrations came from source monitoring data (see Sect. 4.1.6.3), ^{228}Th is not designated as a high priority contaminant in this screening analysis. The concentration of ^{228}Th from the source monitoring data which resulted in a screening index of $>10^{-4}$, for the water ingestion pathway, point out the need for collecting additional water samples and suggest that the source data should be investigated to document the concentrations of ^{228}Th reported in these data.

A listing of the contaminants requiring further investigation is provided in Table 4.8. Results of the water ingestion pathway are not included in this list because of the limited data available for analyses. The contaminants found in the sediments used for the dredging scenario are listed by direct and secondary exposure pathways. The uncertainty associated with results of the screening analysis increases in order from the fish ingestion pathway, to the direct exposure to sediment pathways, to the secondary (modeled) pathways. However, copper is an artifact in the fish ingestion pathway (see Sect. 2.4) and can be designated low priority. In addition, many of the contaminants were detected in less than 5% of the samples analyzed [i.e., bis(2-ethylhexyl)phthalate in the fish ingestion pathway]. For the organics listed in the secondary pathways, only Aroclor 1260, phenanthrene, and pyrene were detected in more than 5% of the samples analyzed. Therefore, additional analyses of these data and the methodology used in detecting these contaminants in environmental samples can undoubtedly reduce the number of contaminants listed in Table 4.8.

4.1.13 Summary for Screening Risk Assessment

The human health screening analysis of the Phase 1 data, in general, confirms the results of the previous screening of historical data (Hoffman et al. 1991). Interim remedial action has been completed on the White Oak Creek Embayment, which excludes it from the present study area; therefore, data from the embayment were not included in this report except for NPDES water monitoring data collected at White Oak Dam. The Phase 1 conclusions for the screening analysis for human health are listed here.

- Only two contaminants were identified by nonconservative screening as high priority contaminants requiring immediate consideration for remedial action. The PCB Aroclor 1260 was found in fish from the mouth of Poplar Creek to above the confluence with East Fork Poplar Creek, the Clinch River arm of Watts Bar Reservoir, Watts Bar Reservoir below the mouth of the Clinch River, and Chicamauga Reservoir at concentrations that classified it as a high priority contaminant; however, PCBs are found in most Tennessee rivers and lakes. Although the ORR is a probable source of PCBs, it is not the only one for the CR/WBR system.
- Thorium-228 was tentatively identified in Poplar Creek as a high priority contaminant in the water ingestion pathway. Thorium-228 was measured only at source monitoring sites at K-25, and the concentrations probably do not represent concentrations in the main body of water. Additionally, ^{228}Th is a daughter of ^{232}U or naturally occurring ^{232}Th , which were not reported in the monitoring data. The absence of either parent radionuclides indicate that the data or analytical methodology used for sample analyses are in need of reevaluation.

Table 4.8. Summary of contaminants requiring further investigation before taking action or designating as low priority based on fish ingestion and dredging scenarios

Contaminants		Pathways ^a	
Organic contaminants		Fish	Direct Secondary
4,4'-DDD ^b	G		M, L
4,4'-DDE ^b	G		
4,4'-DDT ^b	G		M, L
Aldrin ^b	G		
Anthracene ^c		G	M, L, V
Aroclor 1254	G	G	M, L, V
Aroclor 1260		G	M, L, V
Benzo(a)anthracene ^c			M, L, V
Benzo(a)pyrene ^c		G	M, L, V
Benzo(b)fluoranthene ^c			M, L, V
Benzo(k)fluoranthene ^c			M, L
Bis(2-ethylhexyl)phthalate	G		
Chlordane ^b	G		
Chrysene			M, L, V
Fluoranthene		G	M, L, V
Indeno(1,2,3-cd)pyrene			M, L
N-nitro-di-n-propylamine ^c		G	M, L, V
Phenanthrene ^c		G	M, L, V
Pyrene ^c		G	M, L, V
Toxaphene ^b		G	M, L, V
Inorganic contaminants			
Arsenic	G	G,H	M, V
Beryllium	G	G	M, V
Cadmium			M, L, V
Chromium		H	M, L, V
Copper	G		M, L, V
Mercury	G		M, L, V
Nickel			V
Selenium	G		M, L, V
Silver			L, V
Thallium			M, L, V
Uranium			M
Zinc			M, L, V
Radionuclides			
⁶⁰ Co		X	
¹³⁷ Cs	G	X	M, L, V
⁹⁰ Sr			L, V
²³⁴ U			M
²³⁸ U		X	M

^aExposure Pathways:

G=Ingestion

H=Inhalation

X=External Exposure

M=Meat Ingestion

L=Milk Ingestion

V=Vegetable Ingestion

^bInsecticides

^cPolycyclic Aromatic Hydrocarbons

- Results of the screening of the surface water data are inconclusive because of the limited number of Phase 1 samples and the uncertainty associated with using the source monitoring data to predict the concentrations of the contaminants in the different reaches. The water results should be viewed in terms of general patterns rather than results for individual contaminants. The results of the screening analysis indicate that a large number of organic contaminants in Poplar Creek would require further investigation; however, most of these contaminants are common industrial or laboratory chemicals or pesticides that are not unique to the ORR facilities.
- Screening of the contaminants in the fish ingestion pathway identified the PCB Aroclor 1260 as a high priority contaminant. Aroclor 1254, several pesticides (4,4'-DDE, aldrin, and chlordane) that are ubiquitous contaminants in East Tennessee aquatic environments, arsenic, beryllium, mercury, and selenium were identified as contaminants in fish that needed further evaluation. The only radionuclide in the fish ingestion pathway that was identified as needing additional evaluation was ^{137}Cs in the Clinch River arm of Watts Bar Reservoir.
- A conservative screening of near-shore surface-sediment data identified ^{60}Co and ^{137}Cs as low priority contaminants in the ingestion and inhalation pathways. In the external exposure pathway, ^{60}Co and ^{137}Cs were identified as contaminants that would require further investigation. However, because the concentrations at all sites are relatively low and the screening index from all pathways, including external exposure, is $<10^{-4}$, further evaluation of the data will probably classify these contaminants as low priority.
- Screening based on the dredging scenario identified a relatively large number of organic contaminants in Poplar Creek and the Clinch River arm of Watts Bar Reservoir above the mouth of Poplar Creek which would require further investigation. Many of these organic contaminants are common industrial and laboratory chemicals, or by-products of fossil fuel burning, and are not unique to the ORR.
- Conservative screening of Phase 1 sediment data in the dredging scenario eliminated all radionuclides that were identified from all pathways from further consideration except ^{60}Co , ^{137}Cs , and ^{238}U in the external exposure pathway; ^{90}Sr in the vegetation and milk ingestion pathways; ^{234}U and ^{238}U in the meat ingestion pathways; and ^{137}Cs in the milk, meat, and vegetable ingestion pathways.

4.2 ECOLOGICAL RISK

This preliminary screening ecological risk assessment is structured in terms of the standard paradigm for risk assessment (National Research Council, 1983, Risk Assessment in the Federal Government: Managing the Process, Washington, DC: National Academy Press, Suter 1992, Risk Assessment Forum 1992). It begins with a hazard definition phase (also termed hazard identification or problem formulation) that defines the contaminant sources, the receiving environment, and the assessment endpoints. It follows with an analytical phase consisting of exposure assessment and effects assessment. The results of these analyses are then combined in a risk characterization. Finally, the results of this screening ecological risk assessment are compared to a prior screening ecological risk assessment performed for the off-site RI (Suter 1991). This assessment constitutes initial screening of the Phase 1 data. The

assessment is intended to provide an interpretive summary of the data with respect to ecological risks and to guide additional sampling and analysis during the initial stages of Phase 2. Additional risk assessments of the Phase 1 data set will be conducted later.

4.2.1 Hazard Definition

The purpose of this screening assessment is to determine which of the contaminants detected in the off-site environment occur in forms and concentrations that constitute a hazard to the nonhuman biota of the Clinch River downstream of contaminant releases from the ORR. Those contaminants will be the focus of attention in the Phase 2 sampling and analysis program.

The ultimate sources of the contaminants considered in this assessment are the DOE's operations of the ORR, including waste disposal, spills, and use of chemicals such as pesticides. The intermediate sources are the streams that drain the ORR, all of which are tributaries of the Clinch River. The proximate sources are the contaminated media in the Clinch River: water, sediment, and fish tissues. Other contaminant sources occur on the Clinch and Tennessee rivers and their tributaries, which contribute to ambient concentrations; however, they are not objects of this assessment.

The assessment endpoints for this assessment are

1. a 20% or greater reduction in the abundance or production of any species of fish, and
2. a 20% or greater reduction in the abundance or production of any species of aquatic wildlife.

These endpoints are taken from the strategy for ecological risk assessment on the ORR (Suter et al. 1992a). The 20% level of effect is based on EPA regulatory precedents and practical limitations on the detection of ecological effects. Fish are chosen as the aquatic endpoint species because of their societal importance and because fish integrate direct toxic effects and effects on lower levels in the aquatic food web. All fish are included as potential endpoint species, but special emphasis is placed on the largemouth bass (*Micropterus salmoides*) because of its recreational importance and because it is a top carnivore (Suter et al. 1992b). The endpoint wildlife species are mink (*Mustela vison*) and great blue heron (*Ardea herodias*). These species were chosen on the basis of the prior screening ecological risk assessment, which showed the major risk to wildlife to be due to consumption of contaminated fish and showed that mink are particularly sensitive to the most hazardous contaminants and that great blue herons are the most abundant piscivorous bird in the Clinch River near the ORR (Suter 1991).

4.2.2 Exposure Assessment

Exposure assessment estimates the concentrations of contaminants to which organisms are exposed, taking into consideration (as far as possible) the spatial and temporal distributions of those concentrations. For detected chemicals, mean and upper 95% confidence limits on concentrations are considered. The mean concentration is assumed to represent typical chronic exposures. Upper 95% confidence limits may represent local extreme chronic concentrations or acute exposures; the data are not sufficient to determine which. For undetected chemicals, the minimum detection limit and the upper 95% confidence limits on

the detection limits are considered. The derivation of these exposure concentrations and the actual values are presented in Appendix F.

4.2.2.1 Aqueous exposures

The aquatic biota are assumed to be exposed to the dissolved fraction of the chemicals in water, because that is the bioavailable form (HECD 1992). Dissolved (i.e., 0.45 μm filtered) aqueous concentrations are available for inorganic chemicals, but only total concentrations are available for organic chemicals in the CRRI samples. Because only a few CRRI samples are available, NPDES samples are also included and are analyzed and discussed separately. The NPDES analyses are all from total (i.e., unfiltered) samples. In addition, fish tissue concentrations are used to estimate the concentrations of contaminants to which fish have been exposed. This is done by dividing the reported concentrations in fish filets by the bioconcentration factors (BCFs) provided by the EPA in the water quality criteria support documents.

4.2.2.2 Sediment exposures

Two different measures of exposure are used for sediments. First, whole sediment concentrations are used for comparison to screening benchmarks expressed as sediment concentrations. Second, for organic chemicals concentrations of chemicals in pore water are estimated by using the equilibrium partitioning approach. Pore-water concentrations of neutral (nonionic) organic chemicals can be calculated by assuming equilibrium partitioning between the pore water and the organic matter fraction of the sediment (OWRS 1989). The partitioning coefficient (K_p) is the product of the organic matter/water partitioning coefficient (K_{oc}) and the fractional organic matter content of the sediment (f_{oc}). Because sediment-dwelling organisms are approximately as sensitive to chemicals in water as the population of species used to derive the National Water Quality Criteria (NWQC) (OWRS 1989), the same screening criteria can be used as for water but corrected for partitioning. The formula is

$$C_s = K_p C_w$$

or

$$C_s = K_{oc} f_{oc} C_w$$

where C_s and C_w are equivalent concentrations in sediment and water, respectively. K_{oc} is seldom available, but it is quite accurately approximated by the octanol/water partitioning coefficient (K_{ow}) (DiToro 1985). This approach is being used by EPA to derive sediment quality criteria (OWRS 1988). This approach does not produce accurate predictions of pore water concentrations for ionic organic chemicals. However, it is used for all organic chemicals in this assessment because it is conservative for ionic chemicals (i.e., it somewhat overestimates pore water concentrations) and because alternative methods are not available for ionic organic chemicals.

4.2.2.3 Wildlife dietary exposures

Piscivorous wildlife are exposed through consumption of contaminated fish. The fish flesh concentrations reported are for filets rather than the whole fish consumed by a heron or mink. Because many contaminants concentrate in fat or in organs other than muscle, these concentrations are likely to be underestimates of the exposure experienced by piscivorous

wildlife. The error due to this factor is approximately a factor of 1.5 to 2 for metals and 2 to 3 for organic chemicals (Suter 1991). Therefore, the wildlife screening benchmarks (Sect. 4.2.3) include a factor of 2 for metals and 3 for organic chemicals to conservatively correct for this extrapolation.

4.2.3 Effects Assessment: Screening Benchmarks

In this screening assessment, effects are expressed as toxicological benchmarks: concentrations in media which are indicative of acceptable effects (lower screening benchmarks) or unacceptable effects (upper screening benchmarks). Only lower benchmarks were needed for this assessment, and they are presented in Appendix F.

4.2.3.1 Aquatic biota

The screening benchmarks for aquatic biota are taken from Suter et al. (1992b). Because there are no standard screening benchmarks, sets of alternative benchmarks were calculated. For the upper screening benchmark, they are the acute National Ambient Water Quality Criterion (ANAWQC) or, for chemicals with no criterion value, the lowest reported acute LC50 or EC50. The alternative chronic benchmarks are the chronic NAWQC (NAWQC ACTUAL); the chronic pesticide screening value, a concentration that is estimated with 95% confidence not to exceed the unknown chronic NAWQC for those chemicals with no NAWQC (NAWQC ADVISORY); the lowest chronic values for fish (LCV FISH) and daphnids (LCV DAPHNIDS); the lowest EC20 (concentration causing a 20% reduction in survival, growth, or fecundity) for fish (LTV EC20 FISH) and daphnids (LTV EC20 DAPHNID) from chronic toxicity tests; the estimated EC20 for a sensitive species (SS EC20); and the concentration estimated to cause a 20% reduction in the recruit abundance of largemouth bass (EC20 POPULATION). The methods of derivation and source data for these benchmarks are provided in Suter et al. (1992b).

4.2.3.2 Sediments

Two types of benchmarks were used with sediment concentrations. First, NOAA's Effects Range—Low (ER-L) and Effects Range—Median (ER-M) concentrations were used for the 43 chemicals for which they are available (Long and Morgan 1991). The ER-L and ER-M are concentrations in whole sediment that appeared to cause toxic effects in no more than 10% and 50% of sediments, respectively.

The second type of sediment screening benchmark is based on equilibrium partitioning of organic chemicals between sediments and pore water (Sect. 4.2.2). Because it is assumed that the exposure is due to exposure to the pore water, the aqueous screening benchmarks discussed above are used.

4.2.3.3 Wildlife

The screening criteria for wildlife are the same as the prior off-site screening ecological risk assessment (Suter 1991). Because there is not a set of consistent toxicity test data for wildlife as there is for aquatic biota, the screening benchmarks for wildlife are based on expert judgement concerning the best available test endpoint for wildlife found in the literature for each chemical. Wildlife toxicity data were preferred, but data for laboratory rodents and other

domestic animals were included in the review. Data for ruminants were excluded because of the influence of their unique digestive system on uptake. Data on cancer induction were also excluded because low frequencies of tumors have no effects on population endpoints. Toxicological benchmarks expressed as dietary concentrations are preferred and can be used without conversions. When benchmarks are expressed as acute doses, equivalent dietary concentrations are estimated by assuming that an acute dose is equivalent to one day's consumption and that the consumption rate is 8% of body weight per day.

Unlike the screening criteria that are based on NWQC, these criteria do not reflect the diversity of organisms that are exposed by this route. If chronic sublethal effects data are available for either mammals or birds but not both, a correction factor of 10 for differences in sensitivity is applied. This factor is based on limited studies that indicate that birds and mammals can differ in sensitivity by more than an order of magnitude and either can be more sensitive (Sigal and Suter 1989). If there are data for both birds and mammals, but fewer than five species including at least two avian species, we assume that the variance in sensitivity of the test species approximates the variance in all avian and mammalian species and apply no factor for differences among species. If the most sensitive test end point is death or another severe effect (e.g., reproductive failure), a factor of 5 is applied to estimate the threshold for effects on individuals that could result in population reductions (Tucker and Lietzke 1979). These factors are not applied to the upper criterion, which is intended to be a concentration at which effects are certain. The derivation of wildlife benchmarks for individual chemicals is presented in Suter (1991).

4.2.4 Risk Characterization: Contaminant Screening Results

The quotient method is used to integrate exposure and effects in screening assessments. That is, the environmental concentrations are divided by the screening benchmarks and the decision is based on the magnitude of the quotient. If the quotient is <1 for a lower screening benchmark, that indicates a nonhazardous exposure level. If the quotient is >1 for an upper screening benchmark, that indicates a clearly hazardous exposure. Chemical concentrations falling between those benchmarks indicate that a hazard worthy of additional investigation exists. All chemicals that have concentrations greater than lower screening benchmarks are potential "contaminants of concern."

Although the use of multiple upper and lower screening benchmarks increases confidence in the results of the screening assessment, it complicates interpretation of results. Because the alternative screening benchmarks are based on different assumptions, and none of the benchmarks has been shown to consistently estimate thresholds for significant effects, greater confidence in the screening results can be obtained by considering multiple benchmarks. However, interpretation of conflicting results requires expert judgement. The following interpretation rules are applied to the aqueous concentrations, including the estimated pore water concentrations for organics in sediment.

1. If the mean water concentration for a chemical exceeds its chronic NAWQC, then that chemical is a contaminant of concern.
2. If the maximum water concentration for a chemical exceeds its acute NAWQC, then that chemical is a contaminant of concern.
3. If two or more lower screening benchmarks are exceeded by the mean concentration, then that chemical is a contaminant of concern.

4. If a mean concentration exceeds a single lower screening criterion or a maximum concentration exceeds a single upper screening criterion, then the magnitude of the quotient and the conservatism of the benchmark will be considered in deciding whether to include the chemical as a contaminant of concern. However, a chemical is not a contaminant of concern when only the advisory concentration is exceeded.
5. Because of the uncertainty concerning the applicability of the EPA's bioconcentration factors (BCFs) to the field, aqueous concentrations estimated from fish concentrations are used only as supporting evidence.

Metals in sediment are considered to be contaminants of concern if the mean concentration exceeds the ERL or ERM or if the upper confidence limit concentration exceeds the ERM. Chemicals in fish fillets are considered to be contaminants of concern if their mean concentrations exceed lower screening benchmarks for piscivorous wildlife. The screening calculations and their results are presented in Appendix F. The potential contaminants of concern are discussed below, and the selected contaminants of concern are listed in Table 4.9.

4.2.4.1 Potential contaminants of concern in water

Aluminum. Aluminum concentrations in NPDES samples from reaches 1, 2, 3, and 4 exceed NAWQC and one or more other benchmarks. However, these concentrations are almost certainly due to particulate forms that are not bioavailable. Aluminum toxicity is not known to occur in circumneutral ambient waters (Baker et al. 1990).

Cadmium. In reaches 1 and 4, dissolved cadmium was detected in one of three samples. The dissolved concentrations in each reach were greater than the lowest chronic value for daphnids and the sensitive species EC20. These benchmarks were also exceeded by the limits of detection in the reaches where it was not detected. Mean cadmium concentrations in NPDES samples (i.e., total rather than bioavailable concentrations) from reach 1 exceeded all seven benchmarks, including the NAWQC, and in reach 3 exceeded all but the population EC20. Cadmium was not detected in fish fillets, but the detection limits were high relative to toxic concentrations.

Copper. Copper concentrations in fish tissues from reach 5 suggest exposure to extremely high aqueous concentrations. The high mean and upper confidence limit values are due to one extremely high value that is likely to be in error (Sect. 3.4.2). Mean copper concentrations in NPDES samples (i.e., total rather than bioavailable concentrations) exceed seven benchmarks in reach 3 (including the NAWQC), five benchmarks in reaches 1 and 2, and three benchmarks in reach 4. Copper was not detected in dissolved fraction analyses but detection limits exceeded four benchmarks in all reaches.

Cyanide. Mean cyanide concentrations in NPDES samples (i.e., total rather than bioavailable concentrations) exceed all five benchmarks in reaches 2, 3, 4, and 13 (including the NAWQC). Cyanide was not included among the dissolved water or fish analysis results for Phase 1.

Iron. Mean iron concentrations in NPDES samples (i.e., total rather than bioavailable concentrations) exceed two benchmarks in reaches 1, 2, 3, and 4 and the NAWQC in reach 3. Iron was detected in dissolved water analyses from reaches 2, and 5 but not 1, 3, 4, or 10.

Table 4.9. Contaminants of concern for ecological risk assessment

Detected	Undetected
Arsenic	Acenaphthene
Bis(2-ethylhexyl)phthalate	Anthracene
Cadmium	Aroclor 1248
Copper	Aroclor 1254
Cyanide	Benzo(a)anthracene
DDT	Benzo(a)pyrene
Iron	Boron
Lead	Chlordane
Mercury	Chromium
Nickel	Cobalt
PCBs (Aroclors)	Vanadium
Selenium	2-methylphenol
Silver	Benzoic acid
Zinc	Benzyl alcohol
	Lindane
	Heptachlor
	n-Nitrosodiphenylamine
	Phenol
	2,4-Dinitrophenol
	2-Methylphenol
	2-Nitrophenol

Both the detected concentrations and the detection limits were low. Iron was not measured in fish tissues because it is a macronutrient.

Lead. Mean lead concentrations in NPDES samples (i.e., total rather than bioavailable concentrations) exceed five benchmarks in reach 1 and two benchmarks in reaches 2 and 3 including the NAWQC in all three. Lead was detected in dissolved water analyses from reaches 1, 4, and 13 but not 2, 3, 5, or 10. Both the detected concentrations and the detection limits were low. Lead was not detected in fish fillets, and the detection limits were low.

Mercury. Mercury was detected in one dissolved-phase sample from reach 3. The mean concentration exceeded four of six benchmarks. Mean total mercury concentrations in NPDES samples from reaches 1 and 3 also exceed those benchmarks, and in reach 4 it exceeds three benchmarks. Mean aqueous detection limits in reaches 1, 2, 4, 5, 10, and 13 exceed three benchmarks. However, water concentrations estimated from fish fillet concentrations and EPA BCFs suggest aqueous exposure levels much lower than measured values in water. The benchmarks for methyl mercury were used for all of these calculations for conservatism. The relative abundance of organic and inorganic mercury in these samples is unknown.

Nickel. Mean nickel concentrations in NPDES samples (i.e., total rather than bioavailable concentrations) exceed five benchmarks in reach 3 and two benchmarks in reaches 1, 2 and 4. Nickel was not detected in dissolved water samples at a detection limit that was 12% of the NAWQC but above two out of seven benchmarks in all reaches. Nickel was not detected in fish tissues, and the detection limit was lower than all benchmarks.

Selenium. Mean selenium concentrations in fish tissues suggest that the fish were exposed to aqueous concentrations that were higher than two or more benchmarks in all reaches. In reaches 1, 2, 4, 10, and 15, the estimated mean aqueous concentration was greater than the NAWQC. However, selenium was not detected in the dissolved phase of water from any reach even though the detection limit was below the NAWQC and all other benchmarks, and total concentrations in NPDES samples were low. This discrepancy is likely to result from use of the EPA's BCF for this metal. There is now considerable evidence that, in the field, selenium is bioaccumulated through diet, which would make an aqueous concentration estimated using a BCF too high (Lemly 1985).

Silver. Dissolved silver was detected in one of four samples from reach 3. The concentration was greater than all seven lower benchmarks for that metal. In all other reaches silver was undetected, but the limits of detection exceeded all but one of the lower benchmarks. Mean concentrations of total silver in NPDES samples from reach 3 were greater than all seven benchmarks.

4.2.4.2 Potential contaminants of concern in sediment

Arsenic. The upper confidence limit arsenic concentration in sediment exceeds the ERM in reach 3 and the ERL in reaches 1 and 4.

Bis(2-ethylhexyl)phthalate. The estimated mean pore water concentration of this phthalate ester exceeds the lowest chronic value and the lowest EC20 for daphnids as well

as the advisory concentration in reach 5. The upper confidence limit estimated pore water concentration exceeds those three benchmarks in reaches 2, 3, and 5.

Lead. The mean sediment concentration exceeds the ER-L in reaches 5 and 10 and the upper confidence limit concentration exceeds the ER-M in reach 4 and the ER-L in reaches 1, 2, and 3.

Mercury. The mean mercury concentration in sediment exceeds the ER-M in reaches 3 and 4 and exceeds the ER-L in reaches 2 and 5. The upper confidence limit mercury concentration in sediment exceeds the ER-L in reach 1. The ER-L and ER-M values are based on naturally occurring mixtures of organic and inorganic mercury.

Zinc. The mean zinc concentration in sediment exceeds the ER-L in reach 5. The upper confidence limit zinc concentration in sediment exceeds the ER-M in reaches 1 and 4 and the ER-L in reach 3.

4.2.4.3 Potential contaminants of concern for wildlife

Aroclors. Mean concentrations of Aroclor 1254 in reaches 2, 3, 4, 5, 13, and 15 and of Aroclor 1260 in reaches 1, 2, 3, 4, 5, 13, and 15 exceeded the lower piscivore benchmark for PCBs. In addition, upper confidence limit concentrations of Aroclor 1254 in reach 1 and of Aroclor 1260 in reach 10 exceed the lower benchmark. That benchmark is a dietary concentration causing significant reductions in mink reproduction divided by a factor of 5 for an effects threshold and of 3 for fillet to whole fish conversion.

Copper. The upper confidence limit copper concentration in reach 5 exceeds the lower wildlife benchmark. However, this value is suspect (see Sect. 3.4).

4,4'-DDT and Metabolites. Mean DDE in reach 4 exceeded the lower piscivore benchmark, which is the lowest dietary concentration causing significant reductions in avian reproduction divided by a factor of 5 for an effects threshold and of 3 for fillet to whole fish conversion. Upper confidence limit fish exceeded that benchmark for DDE in reaches 1, 2, 4, and 5 for DDD in reaches 3 and 15, and for DDT in reach 4 and exceeded concentrations observed to cause reproductive effects in reaches 4 and 15. DDT is not known to be associated with ORR effluents or wastes, but the highest concentrations are found in fish from the Clinch River downstream from Poplar Creek (reach 4).

Mercury. Mean mercury concentrations in fish from all reaches exceeded the lower piscivore benchmark, which is the lowest dietary concentration causing significant reductions in avian reproduction divided by a factor of 2 for fillet to whole fish conversion. Mean fish filets from reaches 3, 4, 10, 13, and 15 exceeded the 0.1 mg/kg concentration observed to cause reduced reproduction in mallard ducks. The highest mean concentrations in fish were found in lower Poplar Creek (reach 3), but the second highest concentrations were found in Norris Reservoir (reach 10).

Selenium. Mean selenium concentrations in fish from reaches 1, 2, 4, 5, 10, 13, and 15 barely exceeded the lower piscivore benchmark, which is the lowest dietary concentration causing significant reductions in avian reproduction divided by a factor of 2 for fillet to whole fish conversion. None of the mean or upper confidence limit fish exceeded concentrations

observed to cause toxic effects in toxicity tests. The highest mean concentrations in fish were found in Melton Hill Reservoir (reach 1), and the second highest concentrations were found immediately below the reservoir but above Poplar Creek in reach 2.

4.2.4.4 Undetected potential contaminants of concern

Some contaminants were not detected in water, but the detection limits were high enough that they would be contaminants of concern if the actual concentrations were just below the detection limit. In a few cases, such as chlordane and lead, detection limits were above NAWQC.

4.2.5 Comparison to Previous Results

Because of declines in reported concentrations of most chemicals in most media, the number of potential contaminants of concern decreased greatly compared to the screening assessment based on historic data (Suter 1991). Mercury in fish tissues is an exception, having declined only slightly.

Data needs identified in the previous assessment have been met in part (Suter 1991). Concentrations of dissolved metals, which were not available before, were available for this assessment; the data are now consistent across reaches; and monitoring of contaminants in great blue heron eggs and nestlings has begun. However, some data needs identified in the previous assessment are still not filled. These include analyses of whole fish, of sediment pore water, and of certain species for arsenic, mercury, and chromium. In addition, indicators of response to chemical exposures other than body burdens have been monitored in herons but not fish.

5. SUMMARY

This report presents and discusses the results of Phase 1 of DOE's remedial investigation of the CR/WBR system. The approach in this report is to use a screening assessment for human health and ecological risk to identify contaminants, environmental pathways, and sites in the river-reservoir system that will require further investigation during Phase 2 of the CRRI. In addition, the screening assessment will identify contaminants that can be assigned a low priority for further investigation.

The CRRI is designed to address the transport, fate, and distribution of waterborne contaminants released from the ORR and to assess potential risks to human health and the environment associated with these contaminants. The contaminants released since the early 1940s include a variety of radionuclides, metals, and organic compounds. Primary areas of investigation are Melton Hill Reservoir, the Clinch River from Melton Hill Dam to its confluence with the Tennessee River, Poplar Creek, and Watts Bar Reservoir. The contaminants identified in the CR/WBR downstream of the ORR are those associated with the water, suspended particles, deposited sediments, aquatic organisms, and wildlife feeding on aquatic organisms.

A phased remedial investigation of the CR/WBR system is under way to (1) define the nature and extent of the off-site contamination, (2) evaluate associated environmental and human health risks, and (3) preliminarily identify and evaluate potential remediation alternatives.

Phase 1 of the CRRI was an initial sampling and analysis of fish, sediment, and water in selected areas of the CR/WBR chosen to represent differing levels of media contamination. Specifically, Phase 1 was designed to (1) obtain high-quality data to confirm existing historical data for contaminant levels in fish, sediment, and water from the CR/WBR; (2) determine the range of contaminant concentrations present in the river-reservoir system; (3) identify specific contaminants of concern; and (4) establish the reference (background) concentrations for those contaminants. Previously, scoping studies were conducted to provide an initial determination of contaminant distribution in the sediments of WBR (Olsen et al. 1992) and to preliminarily assess potential human health and ecological risk (Suter 1991; Hoffman et al. 1991) in the CR/WBR system.

For Phase 1 of the CRRI, the CR/WBR system was divided into ten reaches, six of which are potentially affected by releases from the ORR. The other four reaches serve as reference areas. Contaminant concentrations for fish, sediment, and water were summarized for each of the ten reaches. Analytes detected at least once in a reach were evaluated separately from those never measured above their analytical detection limit (i.e., nondetected).

Surface water samples were collected for Phase 1 during winter conditions and analyzed for inorganic, radiochemical, and organic constituents to confirm suspected low concentrations of contaminants in the CR/WBR system. Characterization of the nature and extent of surface water contamination in the CR/WBR system during Phase 1 of the CRRI was limited to a single sampling event for each station. The primary conclusions from the collection and analysis of surface water during Phase 1 are listed here.

- Inorganic and organic constituents that may pose a risk to human health or the environment had concentrations comparable to those reported previously and were well below the Tennessee Water Quality Criteria.
- Values for all radiological constituents in surface water were comparable with the mean values found in other investigations. Each of these radionuclides has been or is currently being released from the ORR to surface waters.
- The results from Phase 1 water samples, in combination with source monitoring and historical data, indicate that additional work is needed to characterize Poplar Creek below the mouth of East Fork Poplar Creek and the Clinch River arm of Watts Bar Reservoir.

The primary conclusions from the sediments collected and analyzed during Phase 1 and the near-shore sediment task are listed here.

- Concentrations of ^{137}Cs in the near-shore surface sediment of Watts Bar Reservoir below the mouth of the Clinch River are comparable to background levels. However, concentrations within the Clinch River arm of the reservoir are above background and warrant further study.
- Patterns of ^{137}Cs and ^{60}Co in sediment cores confirm previous studies and indicate that these contaminants are deep in the sediment throughout the system with highest concentrations in the Clinch River arm of the reservoir.
- Concentrations of ^{234}U , ^{235}U , and ^{238}U in sediment cores from Poplar Creek are about 5 times higher than background.
- Concentrations of arsenic, beryllium, cadmium, chromium, copper, and mercury in sediments are above background levels in the Clinch River arm of Watts Bar Reservoir and in Poplar Creek.
- Concentrations of cadmium, lead, and zinc in sediments are higher in the main body of Watts Bar Reservoir, below the mouth of the Clinch River, than in upstream areas.
- Detectable quantities of Aroclor 1254 were only found in sediments at Clinch River Mile 9.5, which is downstream of the confluence of the Clinch River and Poplar Creek.
- Detectable quantities of flouranthene, phenanthrene, and pyrene were found in sediments in Poplar Creek and in the Clinch River between Melton Hill Dam and the mouth of Poplar Creek.

To characterize the nature and extent of contamination in fish in the CR/WBR system, contaminants were analyzed in three species (channel catfish, largemouth bass, and bluegill sunfish) representing three trophic levels. The primary conclusions from the fish sampling and analyses conducted during Phase 1 are listed here.

- Of the 77 organic and 13 inorganic contaminants that were analyzed in fish, a majority (63 organics and 7 inorganics) were not detected from any CR/WBR site.
- Three of the contaminants (⁹⁰Sr, mercury, and zinc) were detected in all fish samples; 17 analytes were detected in greater than 5% of all fish samples. PCBs and chlordane were detected in fish at all CR/WBR sites, and ¹³⁷Cs was detected in 60% of the sites.
- Mercury and ¹³⁷Cs in fish were elevated at sites downstream of known sources of contaminant inputs, East Fork Poplar Creek and White Oak Creek, respectively. No pattern of downstream gradients of PCBs or chlordane was evident in fish collected in Phase 1.
- Phase 1 data for fish are not markedly different from previous historical data on contaminant levels in fish from the CR/WBR system.
- Mean mercury concentrations in fish were significantly higher in Poplar Creek below the confluence with East Fork Poplar Creek and at the first site downstream of the confluence of Poplar Creek and the Clinch River than at the reference site (Norris Reservoir). Mean selenium concentrations in Melton Hill Reservoir and the Clinch River arm of Watts Bar Reservoir were higher than those at the reference sites.
- Mean total PCB concentrations in fish at all CR/WBR sites were significantly higher than those at the reference, whereas mean chlordane concentrations were higher than reference concentrations at sites in Melton Hill Reservoir and the Clinch River arm of Watts Bar Reservoir.

The conclusions of the screening assessment for ecological risk for Phase 1 are listed here.

- The assessment of new data confirms the finding of the assessment of historical data that mercury and PCBs in fish constitute a significant risk to piscivorous wildlife.
- Phase 1 analytical data for dissolved (i.e., bioavailable) concentrations of inorganic chemicals in water greatly reduced the number of metals that appear to be hazardous to aquatic life relative to historical data or to source monitoring (e.g., National Pollutant Discharge Elimination System data).
- Cadmium, mercury, and silver appear to be hazardous to aquatic life in water from some reaches, but the number of samples in which they were detected is extremely small. In addition, the assessment of mercury conservatively assumed that all mercury in water was present as methyl mercury.
- Arsenic, lead, mercury, and zinc in sediments from some reaches exceed concentrations that have been found by the U.S. National Oceanic and Atmospheric Administration to cause effects on benthic communities at some sites.
- PCBs and several other organic chemicals that appeared to be hazardous to aquatic life because of their historical concentrations in sediments and water were not detected in those media in the Phase 1 investigation.

- Because this is a preliminary screening assessment of ecological risks, more detailed analyses will follow.

The human health screening analysis of the Phase 1 data, in general, confirms the results of the previous screening of historical data (Hoffman et al. 1991). The Phase 1 conclusions for the screening analysis for human health are listed here.

- Only two contaminants were identified by nonconservative screening as high priority contaminants requiring further investigation. The PCB Aroclor 1260 was found in fish from the mouth of Poplar Creek to above the confluence with East Fork Poplar Creek, the Clinch River arm of Watts Bar Reservoir, Watts Bar Reservoir below the mouth of the Clinch River, and Chicamauga Reservoir at concentrations that classified it as a high priority contaminant; however, PCBs are found in most Tennessee rivers and lakes. Although the ORR is a probable source of PCBs, it is not the only one for the CR/WBR system.
- Thorium-228 was tentatively identified in Poplar Creek as a high priority contaminant in the water ingestion pathway. Thorium-228 was detected only at source monitoring sites at K-25, and the concentrations probably do not represent concentrations in the main body of water. Additionally, ^{228}Th is a daughter of ^{232}U or naturally occurring ^{232}Th , which were not reported in the monitoring data. The absence of either parent radionuclides along with the ^{232}Th indicate that the data or analytical methodology used for sample analyses are in need of reevaluation.
- Results of the screening of the surface water data are inconclusive because of the limited number of Phase 1 samples and the uncertainty associated with using the source monitoring data to predict the concentrations of the contaminants in the river/reservoir system. The water results should be viewed in terms of general patterns rather than results for individual contaminants. The results of the screening analysis indicate that a large number of organic contaminants in Poplar Creek would require further investigation; however, most of these contaminants are common industrial or laboratory chemicals or pesticides that are not unique to the ORR facilities.
- Screening of the contaminants in the fish ingestion pathway identified the PCB Aroclor 1260 as a high priority contaminant. The Tennessee Department of Environment and Conservation has issued fish consumption advisories for several eastern Tennessee reservoirs (e.g., Fort Loudoun, Watts Bar, Melton Hill) as a result of PCB contamination. Aroclor 1254, several pesticides (4,4'-DDE, aldrin, and chlordane) that are ubiquitous contaminants in eastern Tennessee aquatic environments, arsenic, beryllium, mercury, and selenium were identified as contaminants in fish that needed further evaluation. The only radionuclide in the fish ingestion pathway that was identified as needing additional evaluation was ^{137}Cs in the Clinch River arm of Watts Bar Reservoir.
- A conservative screening of near-shore surface-sediment data identified ^{60}Co and ^{137}Cs as low priority contaminants in the ingestion and inhalation pathways. In the external exposure pathway, ^{60}Co and ^{137}Cs were identified as contaminants that would require further investigation. However, because the concentrations at all sites are relatively low

and the screening index from all pathways, including external exposure, is $<10^{-4}$, further evaluation of the data will probably classify these contaminants as low priority.

- Screening based on the dredging scenario identified a relatively large number of organic contaminants in Poplar Creek and the Clinch River arm of Watts Bar Reservoir above the mouth of Poplar Creek which would require further investigation. Many of these organic contaminants are common industrial and laboratory chemicals, or by-products of fossil fuel burning, and are not unique to the ORR.
- Conservative screening of Phase 1 sediment data in the dredging scenario eliminated all radionuclides that were identified from all pathways from further consideration except ^{60}Co , ^{137}Cs , and ^{238}U in the external exposure pathway; ^{90}Sr in the vegetation and milk ingestion pathways; ^{234}U and ^{238}U in the meat ingestion pathways; and ^{137}Cs in the milk, meat, and vegetable ingestion pathways.

Results from Phase 1 of the CRRI provide the following guidance for the collection and analysis of samples during Phase 2:

- Poplar Creek below the mouth of East Fork Poplar Creek and the Clinch River arm of Watts Bar Reservoir are the reaches of greatest concern based on the number of contaminants that need further evaluation.
- Additional data are required on bioavailable inorganic contaminants (i.e., cadmium, copper, cyanide, lead, mercury, nickel, selenium, and silver) in water to determine their risk to aquatic biota.
- Further sampling and analysis is required for bis(2-ethylhexyl)phthalate, arsenic, lead, mercury, and zinc in sediments. These contaminants may pose a risk to benthic dwelling organisms.
- Potential contaminants of concern for wildlife, identified from the Phase 1 data are Aroclor 1254, Aroclor 1260, 4,4'-DDT and its metabolites (4,4'-DDD and 4,4'-DDE), mercury and selenium.
- Data for the fish ingestion pathway are the most complete. Phase 2 sampling should concentrate on providing statistically valid data for a future baseline risk assessment for human health. Fish tissue analyses should concentrate on the evaluation of contaminants that are routinely released or have been released in the past from the ORR.
- The available data from the Phase 1 water sampling had to be augmented with source monitoring data in order to perform a screening assessment for human health. Additional water data for contaminants will be required, particularly in surface water areas proximate to ORR outfalls and sources, to provide data for the baseline risk assessment. Data from existing monitoring programs will be evaluated for use in Phase 2 of the CRRI.
- Organic and inorganic contaminants common to eastern Tennessee and other metropolitan areas should be identified through further analysis of the available data for downstream and reference reaches. Analyses of organic contaminants in water and

sediment should concentrate on those contaminants that are routinely released or have been released in the past from the ORR. In addition, analytical methods providing lower detection limits need to be evaluated and used for the analysis of organic contaminants.

- Phase 2 sediment sampling should concentrate on Poplar Creek below the mouth of East Fork Poplar Creek and on the Clinch River arm of Watts Bar Reservoir to provide statistically valid data for a baseline risk assessment. Analyses of sediment samples should concentrate on the contaminants identified in the screening analysis as needing further evaluation.

6. REFERENCES

- Adams, S. M., and R. B. McLean. 1985. Estimation of largemouth bass, *Micropterus salmoides* Lacepede, growth using the liver somatic index and physiological variables. *J. Fish Biol.* 26:111-126.
- Ashwood, T. L., C. R. Olsen, I. L. Larsen, and P. D. Lowry. 1986. *Sediment Contamination in Streams Surrounding the Oak Ridge Gaseous Diffusion Plant*. ORNL/TM-9791. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Baker, J. P., D. P. Bernard, S. W. Christensen, M. J. Sale, J. Freda, K. J. Heltcher, D. R. Marmorek, L. Rowe, P. F. Scanlon, G. W. Suter II, W. J. Warren-Hicks, and P. M. Welbourn. 1990. *Biological Effects of Changes in Surface Water Acid-Base Chemistry*. NAPAP Report 13. National Acid Precipitation Assessment Program, Washington, D.C.
- Benoit, D. A. 1975. Chronic effects of copper on survival, growth, and reproduction of the bluegill (*Lepomis macrochirus*). *Trans. Am. Fish. Soc.* 104:353-358.
- Blaylock, B. G., M. L. Frank, F. O. Hoffman, L. A. Hook, G. W. Suter, and J. A. Watts. 1992a. *Screening of Contaminants in Waste Area Grouping 2 at Oak Ridge National Laboratory, Oak Ridge, Tennessee*. ORNL/ER-62/R1. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Blaylock, B. G., M. L. Frank, L. A. Hook, F. O. Hoffman, and C. J. Ford. 1992b. *White Oak Creek Embayment Site Characterization and Contaminant Screening Report*. ORNL/ER-81. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Byrd, J. T. 1991. *Work Plan for Continuation of Arsenic Biogeochemistry in the Clinch River and Watts Bar Reservoir II*. Submitted to Oak Ridge National Laboratory. Skidaway Institute of Oceanography, Georgia.
- DiToro, D. M. 1985. A particle interaction model of reversible organic chemical sorption. *Chemosphere* 14:1503-1538.
- EPA (U.S. Environmental Protection Agency). 1990. *Health Effects Assessment Summary Tables*. NTIS No. PB90-921104.
- EPA (U.S. Environmental Protection Agency). 1992. *Health Effects Assessment Summary Tables, Annual Update*. OHEA, ECAO-CIN-821.
- Goldman, C. R., and A. J. Horne. 1983. *Limnology*. McGraw-Hill, New York.
- HECD (Health and Criteria Division). 1992. *Interim Guidance on Interpretation and Implementation of Aquatic Life Criteria for Metals*. U.S. Environmental Protection Agency, Washington, D.C.
- Hoffman, F. O., B. G. Blaylock, M. L. Frank, L. A. Hook, E. L. Etnier, and S. S. Talmage.

1991. *Preliminary Screening of Contaminants in the Off-Site Surface Water Environment Downstream of the U.S. Department of Energy Oak Ridge Reservation*. ORNL/ER-9. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Martin Marietta Energy Systems. 1990. *Clinch River RCRA Facility Investigation Plan*. ES/ER-1/D1. Martin Marietta Energy Systems, Inc., Oak Ridge, Tennessee.

Kornegay, F. C., D. C. West, R. A. Evans, S. T. Goodpasture, M. F. Tardiff, and A. R. Wilson. 1991. *Oak Ridge Reservation Environmental Report for 1990. Volume 2: Data Presentation*. ES/ESH-18/V2. Martin Marietta Energy Systems, Inc., Oak Ridge, Tennessee.

Lemly, A. D. 1985. Toxicology of selenium in a freshwater reservoir: Implications for environmental hazard evaluation and safety. *Ecotoxicol. Environ. Safety* 10:314-338.

Leslie, M., and B. L. Kimmel. 1992. *White Oak Creek Embayment Time-Critical CERCLA Removal Action Regulatory Compliance Study*. ORNL/ER/Sub-91-KA931/3. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Long, E. R., and L. G. Morgan. 1991. *The Potential for Biological Effects of Sediment-Sorbed Contaminants Tested in the National Status and Trends Program*. NOAA Technical Memo. NOS OMA 52. NOAA, Seattle, Washington.

National Research Council. 1983. *Risk Assessment in the Federal Government: Managing the Process*. National Academy of Science, Washington, D.C.

NCRP (National Council on Radiation Protection and Measurements). 1978. *A Handbook of Radioactivity Measurements Procedures*. Report 58. National Council on Radiation Protection and Measurements. Washington, D.C. 506 pp.

NUS Corporation. 1990. *Analytical Laboratory Guidebook for Environmental Professionals*. NUS Corporation, Pittsburgh, Pennsylvania.

Oakes, T. W., W. F. Ohnesorge, J. S. Eldridge, T. G. Scott, D. W. Parsons, H. M. Hubbard, O. M. Sealand, K. E. Shank, and L. D. Eyman. 1982. *Technical Background Information for the Environmental and Safety Report, Vol. 5: The 1977 Clinch River Sediment Survey-Data Presentation*. ORNL-5878. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Olsen, C. R., I. L. Larsen, P. D. Lowry, C. R. Moriones, C. J. Ford, K. C. Dearstone, R. R. Turner, B. L. Kimmel, and C. C. Brandt. 1992. *Transport and Accumulation of Cesium-137 and Mercury in the Clinch River and Watts Bar Reservoir System*. ORNL/ER-7. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Olsen, C. R., M. Thein, I. L. Larsen, P. D. Lowry, P. J. Mulholland, N. H. Cutshall, J. T. Byrd, and H. L. Windom. 1989. Plutonium, lead-210, and carbon isotopes in the Savannah Estuary: Riverine versus marine sources. *Environ. Sci. Technol.* 23(12):1474-1481.

OSWER (Office of Solid Waste and Emergency Response). 1991. OSWER Directive 9285.6-03, March 25, 1991. Office of Energy and Remedial Response Toxics Integration Branch, U.S. Environmental Protection Agency, Washington, D.C.

OWRS (Office of Water Regulations and Standards). 1989. *Briefing Report to the EPA Science Advisory Board on the Equilibrium Partitioning Approach to Generating Sediment Quality Criteria*. EPA 440/5-89-002. Environmental Protection Agency, Washington, D.C.

Risk Assessment Forum. 1992. *Framework for Ecological Risk Assessment*. EPA/630/R-92/001. U.S. Environmental Protection Agency, Washington, D.C.

Sigal, L. L., and G. W. Suter II. 1989. Potential effects of chemical agents on terrestrial resources. *Environ. Prof.* 11:376-384.

Suter, G. W., II. 1991. *Screening Level Risk Assessment for Off-Site Ecological Effects in Surface Waters Downstream from the U.S. Department of Energy Oak Ridge Reservation*. ORNL/ER-8. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Suter, G. W., II (ed.). 1992. *Ecological Risk Assessment*. Lewis Pub., Chelsea, Michigan.

Suter, G. W., II, A. Refearn, R. K. White, and R. A. Shaw. 1992a. *Approach and Strategy for Performing Ecological Risk Assessments for the Department of Energy Oak Ridge Field Office Environmental Restoration Program*. ES/ER/TM-33. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Suter, G. W., II, M. A. Futrell, and J. A. Kerchner. 1992b. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota on the Oak Ridge Reservation, Oak Ridge, Tennessee*. ORNL/ER-139. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Tucker, R. K., and J. S. Lietzke. 1979. Comparative toxicity of insecticides for vertebrate wildlife and fish. *Pharmacol. Ther.* 6:167-220.

Turner, R. R., C. R. Olsen, and W. J. Wilcox, Jr. 1985. Environmental fate of Hg and ¹³⁷Cs discharged from Oak Ridge facilities. pp. 329-338. In D. D. Hemphill (ed.), *Trace Substances in Environmental Health—XVIII*. University of Missouri, Columbia.

TVA (Tennessee Valley Authority). 1991. *Results of Sediment and Water Sampling for Inorganic, Organic, and Radionuclide Analysis at Recreation Areas and Water Intakes—Norris, Melton Hill, and Watts Bar Lakes*. Data Report. TVA Resource Development, River Basin Operations, Water Resources, Water Quality Department, Chattanooga, Tennessee.

USEPA (U.S. Environmental Protection Agency). 1989. *Interim Final: Risk Assessment Guidance for Superfund. Vol. I: Human Health Evaluation Manual*. OSWER Directive 9285.7-01a. EPA Office of Emergency and Remedial Response, Washington, D.C.

USEPA (U.S. Environmental Protection Agency). 1990. *Quality Assurance/Quality Control Guidance for Removal Activities, Sampling QA/QC Plan and Data Validation Procedures*. Interim Final. EPA/540/G-90/004. Environmental Protection Agency, Washington, D.C.

Appendix A

LISTING OF ALL ANALYTES WITH CAS NUMBERS

CONTENTS OF APPENDIX A

Table A.1	Listing of all analytes with CAS numbers and matrices analyzed for each analyte	A-3
------------------	--	------------

TABLE A.1
LISTING OF ALL ANALYTES WITH CAS NUMBERS AND MATRICIES ANALYZED FOR EACH ANALYTE

TYPE OF ANALYSIS	ANALYTE	CHEMICAL ABSTRACT SERVICE NUMBER	ANALYZED IN (FISH/SEDIMENT/WATER)
INORGANICS	ANTIMONY	7440-36-0	F/S/W
	ARSENIC	7440-38-2	F/S/W
	BERYLLIUM	7440-41-7	F/S/W
	CADMIUM	7440-43-9	F/S/W
	CALCIUM	7440-70-2	./S/W
	CHROMIUM	7440-47-3	F/S/W
	COPPER	7440-50-8	F/S/W
	IRON	7439-89-6	./S/W
	LEAD	7439-92-1	F/S/W
	MAGNESIUM	7439-95-4	./S/W
	MERCURY	7439-97-6	F/S/W
	NICKEL	7440-02-0	F/S/W
	POTASSIUM	7440-09-7	./S/W
	SELENIUM	7782-49-2	F/S/W
	SILVER	7440-22-4	F/S/W
	SODIUM	7440-23-5	./S/W
	THALLIUM	7440-28-0	F/S/W
	ZINC	7440-66-6	F/S/W
ORGANICS	ACENAPHTHENE	83-32-9	F/S/W
	ACENAPHTHYLENE	208-96-8	F/S/W
	ALDRIN	309-00-2	F/S/W
	ALPHA-BHC	319-84-6	./S/W
	ALPHA-CHLORDANE	5103-71-9	F/S/W
	ALPHA-CHLORDENE	56534-02-2	F/. /.
	ANTHRACENE	120-12-7	F/S/W
	AROCLOR-1016	12674-11-2	./S/W
	AROCLOR-1221	11104-28-2	./S/W
	AROCLOR-1232	11141-16-5	./S/W
	AROCLOR-1242	53469-21-9	./S/W
	AROCLOR-1248	12672-29-6	./S/W
	AROCLOR-1254	11097-69-1	F/S/W
	AROCLOR-1260	11096-82-5	F/S/W
	BENZO(A)ANTHRACENE	56-55-3	F/S/W
	BENZO(A)PYRENE	50-32-8	F/S/W
	BENZO(B)FLUORANTHENE	205-99-2	F/S/W
	BENZO(G,H,I)PERYLENE	191-24-2	F/S/W
	BENZO(K)FLUORANTHENE	207-08-9	F/S/W
	BENZOIC ACID	65-85-0	F/S/W
	BENZYL ALCOHOL	100-51-6	F/S/W
	BETA-BHC	319-85-7	./S/W
	BIS(2-CHLOROETHOXY)METHANE	111-91-1	F/S/W
	BIS(2-CHLOROETHYL)ETHER	111-44-4	F/S/W
	BIS(2-CHLOROISOPROPYL)ETHER	108-60-1	F/S/W
	BIS(2-ETHYLHEXYL)PHTHALATE	117-81-7	F/S/W
	BUTYLBENZYLPHTHALATE	85-68-7	F/S/W
	CHLORDENE	3734-48-3	F/. /.
	CHRYSENE	218-01-9	F/S/W
	DELTA-BHC	319-86-8	./S/W
	DI-N-BUTYLPHTHALATE	84-74-2	F/S/W
	DI-N-OCTYLPHTHALATE	117-84-0	F/S/W
	DIBENZ(A,H)ANTHRACENE	53-70-3	F/S/W
	DIBENZOPURAN	132-64-9	F/S/W
	DIELDRIN	60-57-1	./S/W
	DIETHYLPHTHALATE	84-66-2	F/S/W
	DIMETHYLPHTHALATE	131-11-3	F/S/W
	ENDOSULFAN I	959-98-8	./S/W
	ENDOSULFAN II	33213-65-9	./S/W

TABLE A.1
LISTING OF ALL ANALYTES WITH CAS NUMBERS AND MATRICIES ANALYZED FOR EACH ANALYTE

TYPE OF ANALYSIS	ANALYTE	CHEMICAL ABSTRACT SERVICE NUMBER	ANALYZED IN (FISH/SEDIMENT/WATER)
ORGANICS	ENDOSULFAN SULFATE	1031-07-8	./S/W
	ENDRIN	72-20-8	./S/W
	ENDRIN KETONE	53494-70-5	./S/W
	FLUORANTHENE	206-44-0	F/S/W
	FLUORENE	86-73-7	F/S/W
	GAMMA-BHC (LINDANE)	58-89-9	./S/W
	GAMMA-CHLORDANE	5103-74-2	F/S/W
	GAMMA-CHLORDENE	56641-38-4	F/. /.
	HEPTACHLOR	76-44-8	./S/W
	HEPTACHLOR EPOXIDE	1024-57-3	./S/W
	HEXACHLOROBENZENE	118-74-1	F/S/W
	HEXACHLOROBUTADIENE	87-68-3	F/S/W
	HEXACHLOROCYCLOPENTADIENE	77-47-4	F/S/W
	HEXACHLOROETHANE	67-72-1	F/S/W
	INDENO(1,2,3-CD)PYRENE	193-39-5	F/S/W
	ISOPHORONE	78-59-1	F/S/W
	METHOXYCHLOR	72-43-5	./S/W
	N-NITROSO-DI-N-PROPYLAMINE	621-64-7	F/S/W
	N-NITROSODIPHENYLAMINE	86-30-6	F/S/W
	NAPHTHALENE	91-20-3	F/S/W
	NITROBENZENE	98-95-3	F/S/W
	OXYCHLORDANE	27304-13-8	F/. /.
	PENTACHLOROPHENOL	87-86-5	F/S/W
	PHENANTHRENE	85-01-8	F/S/W
	PHENOL	108-95-2	F/S/W
	PYRENE	129-00-0	F/S/W
	TOXAPHENE	8001-35-2	./S/W
	1,2-DICHLOROBENZENE	95-50-1	F/S/W
	1,2,4-TRICHLOROBENZENE	120-82-1	F/S/W
	1,3-DICHLOROBENZENE	541-73-1	F/S/W
	1,4-DICHLOROBENZENE	106-46-7	F/S/W
	2-CHLORONAPHTHALENE	91-58-7	F/S/W
	2-CHLOROPHENOL	95-57-8	F/S/W
	2-METHYLNAPHTHALENE	91-57-6	F/S/W
	2-METHYLPHENOL	95-48-7	F/S/W
	2-NITROANILINE	88-74-4	F/S/W
	2-NITROPHENOL	88-75-5	F/S/W
	2,4-DICHLOROPHENOL	120-83-2	F/S/W
	2,4-DIMETHYLPHENOL	105-67-9	F/S/W
	2,4-DINITROPHENOL	51-28-5	F/S/W
	2,4-DINITROTOLUENE	121-14-2	F/S/W
	2,4,5-TRICHLOROPHENOL	95-95-4	F/S/W
	2,4,6-TRICHLOROPHENOL	88-06-2	F/S/W
	2,6-DINITROTOLUENE	606-20-2	F/S/W
	3-NITROANILINE	99-09-2	F/S/W
	3,3'-DICHLOROBENZIDINE	91-94-1	F/S/W
	4-BROMOPHENYL-PHENYLETHER	101-55-3	F/S/W
	4-CHLORO-3-METHYLPHENOL	59-50-7	F/S/W
	4-CHLOROANILINE	106-47-8	F/S/W
	4-CHLOROPHENYL-PHENYLETHER	7005-72-3	F/S/W
	4-METHYLPHENOL	106-44-5	F/S/W
	4-NITROANILINE	100-01-6	F/S/W
	4-NITROPHENOL	100-02-7	F/S/W
	4,4'-DDD	72-54-8	F/S/W
	4,4'-DDE	72-55-9	F/S/W
	4,4'-DDT	50-29-3	F/S/W
	4,6-DINITRO-2-METHYLPHENOL	534-52-1	F/S/W

TABLE A.1
LISTING OF ALL ANALYTES WITH CAS NUMBERS AND MATRICIES ANALYZED FOR EACH ANALYTE

TYPE OF ANALYSIS	ANALYTE	CHEMICAL ABSTRACT SERVICE NUMBER	ANALYZED IN (FISH/SEDIMENT/WATER)
RAD	AM-241	14596-10-2	./S/.
	CM-243,244	15757-87-6	./S/.
	CM-245,246	15621-76-8	./S/.
	CM-248	15758-33-5	./S/.
	CO-60	10198-40-0	F/S/W
	CS-137	10045-97-3	F/S/W
	H-3	10028-17-8	././W
	PU-238	13981-16-3	./S/.
	PU-239,240		./S/.
	SR-90	10098-97-2	F/S/W
	U-234	13966-29-5	./S/.
	U-235	15117-96-1	./S/.
	U-238	7440-61-1	./S/.

Appendix B

SUMMARY STATISTICS FOR WATER SAMPLES

CONTENTS OF APPENDIX B

Table B.1	Summary statistics for water data summarized for all reaches sorted by type of analysis and decreasing percentage detected	B-3
Table B.2	Summary statistics for water data detected compounds summarized by reach sorted by type of analysis, analysis, and reach	B-7

TABLE B.1
SUMMARY STATISTICS FOR WATER DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

TYPE OF ANALYSIS-INORGANICS											
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
CALCIUM, DISS.	FILTRATE	18	18	100.00	17700.00	V/V	34800.00	V/V	27318.67	7150.83	UG/L
CALCIUM, TOTAL	WATER	18	18	100.00	17600.00	V/V	35400.00	V/V	27305.53	7258.44	UG/L
MAGNESIUM, DISS.	FILTRATE	18	18	100.00	4550.000	V/V	10500.00	V/V	7986.67	2308.53	UG/L
MAGNESIUM, TOTAL	WATER	18	18	100.00	4520.000	V/V	10500.00	V/V	8018.89	2332.05	UG/L
POTASSIUM, DISS.	FILTRATE	18	18	100.00	1300.000	V/V	1840.00	V/V	1542.22	178.87	UG/L
POTASSIUM, TOTAL	WATER	18	18	100.00	1250.000	V/V	1840.00	V/V	1540.58	180.85	UG/L
SODIUM, DISS.	FILTRATE	18	18	100.00	2840.000	V/V	5280.00	V/V	4370.00	822.88	UG/L
SODIUM, TOTAL	WATER	18	18	100.00	2440.000	V/V	5240.00	V/V	4320.58	846.88	UG/L
IRON, TOTAL	WATER	18	18	88.89	15.000	U	745.00	J	288.39	208.95	UG/L
LEAD, TOTAL	WATER	19	4	21.05	0.360	U	11.00	U	1.90	2.49	UG/L
LEAD, DISS.	FILTRATE	19	3	15.79	0.360	U	2.50	U	1.25	0.80	UG/L
IRON, DISS.	FILTRATE	15	2	13.33	5.800	U	80.00	U	17.39	21.55	UG/L
COPPER, TOTAL	WATER	18	2	11.11	4.400	U	8.50	V/V	4.52	0.49	UG/L
COPPER, DISS.	FILTRATE	19	2	10.53	0.210	U	0.39	V/V	0.22	0.05	UG/L
BERYLLIUM, TOTAL	WATER	18	1	5.56	3.900	U	3.90	V/V	8.68	5.53	UG/L
ZINC, DISS.	FILTRATE	18	1	5.56	3.400	U	21.00	V/V	11.22	11.08	UG/L
ZINC, TOTAL	WATER	18	1	5.56	3.400	U	52.00	V/V	0.22	0.07	UG/L
MERCURY, DISS.	FILTRATE	19	1	5.26	0.200	U	0.52	V/V	1.39	2.81	UG/L
SILVER, DISS.	FILTRATE	19	1	5.26	0.740	U	13.00	V/V	0.95	0.93	UG/L
SILVER, TOTAL	WATER	19	1	5.26	0.740	U	4.80	V/V	0.95	0.93	UG/L
ANTIMONY, DISS.	FILTRATE	19	0	0.00	1.900	U	1.90	UJ	1.90	0.00	UG/L
ANTIMONY, TOTAL	WATER	19	0	0.00	1.900	U	1.30	UJ	1.30	0.00	UG/L
ARSENIC, DISS.	FILTRATE	19	0	0.00	1.300	U	1.30	UJ	1.30	0.00	UG/L
ARSENIC, TOTAL	WATER	19	0	0.00	1.300	U	1.30	UJ	1.30	0.00	UG/L
BERYLLIUM, DISS.	FILTRATE	18	0	0.00	3.900	U	3.90	U	3.90	0.00	UG/L
BERYLLIUM, TOTAL	WATER	18	0	0.00	0.210	U	0.31	U	0.22	0.02	UG/L
CHROMIUM, DISS.	FILTRATE	18	0	0.00	9.800	U	9.80	U	9.80	0.00	UG/L
CHROMIUM, TOTAL	WATER	18	0	0.00	9.800	U	9.80	U	9.80	0.00	UG/L
COPPER, DISS.	FILTRATE	18	0	0.00	4.400	U	4.40	U	4.40	0.00	UG/L
COPPER, TOTAL	WATER	19	0	0.00	0.200	U	0.20	U	0.20	0.00	UG/L
NICKEL, DISS.	FILTRATE	18	0	0.00	19.000	U	19.00	U	19.00	0.00	UG/L
NICKEL, TOTAL	WATER	18	0	0.00	19.000	U	19.00	U	19.00	0.00	UG/L
SELENIUM, DISS.	FILTRATE	19	0	0.00	0.970	U	0.97	UJ	0.97	0.00	UG/L
SELENIUM, TOTAL	WATER	19	0	0.00	0.970	U	0.97	UJ	0.97	0.00	UG/L
THALLIUM, DISS.	FILTRATE	18	0	0.00	2.700	U	2.70	U	2.70	0.00	UG/L
THALLIUM, TOTAL	WATER	19	0	0.00	2.700	U	2.70	U	2.70	0.00	UG/L

TYPE OF ANALYSIS-ORGANICS

ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
BIS(2-ETHYLHEXYL)PHTHALATE	WATER	18	1	5.56	3.000	U	10.00	UJ	8.94	2.46	UG/L
ACENAPHTHENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
ACENAPHTHYLENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
ALDRIN	WATER	12	0	0.00	0.050	U	0.05	U	0.05	0.00	UG/L

TABLE B.1
SUMMARY STATISTICS FOR WATER DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

TYPE OF ANALYSIS-ORGANICS (CONTINUED)											
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
ALPHA-BHC	WATER	12	0	0.00	0.050	U	0.05	U	0.05	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L	
WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00		

TABLE B.1
SUMMARY STATISTICS FOR WATER DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	%- DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
INDENO(1,2,3-CD)PYRENE	WATER	16	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
ISOPHORONE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
METHOXYCHLOR	WATER	12	0	0.00	0.500	U	0.50	U	0.50	0.00	UG/L
N-NITROSO-DI-N-PROPYLAMINE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
N-NITROSO-DIPHENYLAMINE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
NAPHTHALENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
NITROBENZENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
PENTACHLOROPHENOL	WATER	17	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
PHENANTHRENE	WATER	19	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
PHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
PYRENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
TOXAPENE	WATER	12	0	0.00	1.000	U	1.00	U	1.00	0.00	UG/L
1,2-DICHLOROBENZENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
1,2,4-TRICHLOROBENZENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
1,3-DICHLOROBENZENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
1,4-DICHLOROBENZENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2-CHLORONAPHTHALENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2-CHLOROPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2-METHYLNAPHTHALENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2-METHYLPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2-NITROANILINE	WATER	18	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
2-NITROPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2,4-DICHLOROPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2,4-DIMETHYLPHENOL	WATER	17	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
2,4-DINITROPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2,4-DINITROTOLUENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2,4,5-TRICHLOROPHENOL	WATER	15	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
2,4,6-TRICHLOROPHENOL	WATER	15	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
2,6-DINITROTOLUENE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
3-NITROANILINE	WATER	18	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
3,3'-DICHLOROBENZIDINE	WATER	18	0	0.00	20.000	U	20.00	UJ	20.00	0.00	UG/L
4-BROMOPHENYL-PHENYLETHER	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
4-CHLORO-3-METHYLPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
4-CHLOROANILINE	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
4-CHLOROPHENYL-PHENYLETHER	WATER	18	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
4-METHYLPHENOL	WATER	17	0	0.00	10.000	U	10.00	UJ	10.00	0.00	UG/L
4-NITROANILINE	WATER	18	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
4-NITROPHENOL	WATER	17	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L
4,4'-DDD	WATER	12	0	0.00	0.100	U	0.10	U	0.10	0.00	UG/L
4,4'-DDE	WATER	12	0	0.00	0.100	U	0.10	U	0.10	0.00	UG/L
4,4'-DDT	WATER	12	0	0.00	0.100	U	0.10	U	0.10	0.00	UG/L
4,6-DINITRO-2-METHYLPHENOL	WATER	17	0	0.00	50.000	U	50.00	UJ	50.00	0.00	UG/L

TABLE B.1
SUMMARY STATISTICS FOR WATER DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM		MAXIMUM		AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
					VALUE	QUAL.	VALUE	QUAL.			
SR-80	WATER	16	16	100.00	-0.500	J	1.50	J	-0.01	0.53	PCI/L
CS-137	EVAPORATE	18	7	38.89	0.009	V/V	0.22	V/V	0.12	0.05	PCI/L
CS-137	RESIDUE	15	3	20.00	2.578	UJ	57.10	UJ	14.12	12.82	PCI/G
CO-60	EVAPORATE	18	1	5.56	0.087	UJ	0.29	UJ	0.15	0.06	PCI/L
CO-60	RESIDUE	15	0	0.00	3.284	UJ	87.55	UJ	21.25	16.45	PCI/G

TABLE B.2
SUMMARY STATISTICS FOR WATER DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS-INORGANICS											
ANALYSIS	SAMPLE MATRIX	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. SAMPLE DEV. UNITS
BERYLLIUM, TOTAL	WATER	1	3	0	0.00	3.900	U	3.90	U	3.90	0.00 UG/L
	WATER	2	2	0	0.00	3.900	U	3.90	U	3.90	0.00 UG/L
	WATER	3	4	1	25.00	3.900	U	3.90	V/V	3.90	0.00 UG/L
	WATER	4	2	0	0.00	3.900	U	3.90	U	3.90	0.00 UG/L
	WATER	5	4	0	0.00	3.900	U	3.90	U	3.90	0.00 UG/L
	WATER	10	2	0	0.00	3.900	U	3.90	U	3.90	0.00 UG/L
	WATER	13	1	0	0.00	3.900	U	3.90	U	3.90	0.00 UG/L
	FILTRATE	1	3	1	33.33	0.210	U	0.38	V/V	0.27	0.10 UG/L
	FILTRATE	2	2	0	0.00	0.210	U	0.21	U	0.21	0.00 UG/L
	FILTRATE	3	4	0	0.00	0.210	U	0.21	U	0.21	0.00 UG/L
CADMIUM, DISS.	FILTRATE	4	3	1	33.33	0.210	U	0.31	V/V	0.24	0.08 UG/L
	FILTRATE	5	4	0	0.00	0.210	U	0.21	U	0.21	0.00 UG/L
	FILTRATE	10	2	0	0.00	0.210	U	0.21	U	0.21	0.00 UG/L
	FILTRATE	13	1	0	0.00	0.210	U	0.21	U	0.21	0.00 UG/L
	FILTRATE	1	3	3	100.00	33900.00	V/V	34300.00	V/V	34166.67	230.94 UG/L
	FILTRATE	2	2	2	100.00	33600.00	V/V	34500.00	V/V	34050.00	638.40 UG/L
	FILTRATE	3	4	4	100.00	21100.00	V/V	24600.00	V/V	22075.00	1685.97 UG/L
	FILTRATE	4	2	2	100.00	24600.00	V/V	34000.00	V/V	29300.00	8646.80 UG/L
	FILTRATE	5	4	4	100.00	18000.00	V/V	33000.00	V/V	21875.00	7353.17 UG/L
	FILTRATE	10	2	2	100.00	33900.00	V/V	34800.00	V/V	34300.00	707.11 UG/L
CALCIUM, DISS.	FILTRATE	13	1	1	100.00	17700.00	V/V	17700.00	V/V	17700.00	0.00 UG/L
	WATER	1	3	3	100.00	34100.00	V/V	35400.00	V/V	34566.67	723.42 UG/L
	WATER	2	2	2	100.00	33600.00	V/V	33800.00	V/V	33700.00	141.42 UG/L
	WATER	3	4	4	100.00	21000.00	V/V	24900.00	V/V	22175.00	1855.40 UG/L
	WATER	4	2	2	100.00	23500.00	V/V	33800.00	V/V	28650.00	7283.20 UG/L
	WATER	5	4	4	100.00	17800.00	V/V	33100.00	V/V	21925.00	7455.81 UG/L
	WATER	10	2	2	100.00	34300.00	V/V	34800.00	V/V	34550.00	353.55 UG/L
	WATER	13	1	1	100.00	17800.00	V/V	17800.00	V/V	17800.00	0.00 UG/L
	WATER	1	3	1	33.33	4.400	U	4.40	V/V	4.40	0.00 UG/L
	WATER	2	2	1	50.00	4.400	U	6.50	V/V	5.45	1.48 UG/L
COPPER, TOTAL	WATER	3	4	0	0.00	4.400	U	4.40	U	4.40	0.00 UG/L
	WATER	4	2	0	0.00	4.400	U	4.40	U	4.40	0.00 UG/L
	WATER	5	4	0	0.00	4.400	U	4.40	U	4.40	0.00 UG/L
	WATER	10	2	0	0.00	4.400	U	4.40	U	4.40	0.00 UG/L
	WATER	13	1	0	0.00	4.400	U	4.40	U	4.40	0.00 UG/L
	FILTRATE	1	2	0	0.00	8.000	U	80.00	U	44.50	50.20 UG/L
	FILTRATE	2	2	1	50.00	5.600	U	54.00	V/V	29.80	34.22 UG/L
	FILTRATE	3	4	0	0.00	6.700	U	10.00	U	7.82	1.48 UG/L
	FILTRATE	4	2	0	0.00	5.600	U	9.20	U	7.40	2.55 UG/L
	FILTRATE	5	3	1	33.33	12.000	U	29.00	V/V	18.33	9.29 UG/L
IRON, DISS.	FILTRATE	10	2	0	0.00	5.600	U	5.60	U	5.60	0.00 UG/L
	WATER	1	3	3	100.00	100.000	V/V	295.00	V/V	170.33	108.26 UG/L
	WATER	2	2	2	100.00	120.000	V/V	150.00	V/V	135.00	21.21 UG/L
	WATER	3	4	4	100.00	328.000	J	745.00	J	485.25	190.38 UG/L
	WATER	4	2	2	100.00	202.000	V/V	218.00	V/V	210.00	11.31 UG/L
	WATER	5	4	4	100.00	121.000	V/V	575.00	J	277.75	211.03 UG/L
	WATER	10	2	0	0.00	15.000	U	22.00	U	18.50	4.95 UG/L
	WATER	13	1	0	0.00	15.000	U	22.00	U	18.50	4.95 UG/L
	WATER	1	3	3	100.00	100.000	V/V	295.00	V/V	170.33	108.26 UG/L
	WATER	2	2	2	100.00	120.000	V/V	150.00	V/V	135.00	21.21 UG/L
WATER	3	4	4	100.00	328.000	J	745.00	J	485.25	190.38 UG/L	
WATER	4	2	2	100.00	202.000	V/V	218.00	V/V	210.00	11.31 UG/L	
WATER	5	4	4	100.00	121.000	V/V	575.00	J	277.75	211.03 UG/L	
WATER	10	2	0	0.00	15.000	U	22.00	U	18.50	4.95 UG/L	

TABLE B.2
SUMMARY STATISTICS FOR WATER DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH
TYPE OF ANALYSIS-INORGANICS
(CONTINUED)

ANALYSIS	SAMPLE MATRIX	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV. UNITS
IRON, TOTAL	WATER	13	1	1	100.00	821.000	J	821.00	J	821.00	UG/L
LEAD, DISS.	FILTRATE	1	3	1	33.33	0.700	U	2.40	V/V	1.80	0.95 UG/L
LEAD, DISS.	FILTRATE	2	2	0	0.00	2.000	U	2.50	U	2.25	0.35 UG/L
LEAD, DISS.	FILTRATE	3	4	0	0.00	0.360	U	1.80	U	1.34	0.70 UG/L
LEAD, DISS.	FILTRATE	4	3	1	33.33	0.600	U	1.50	U	0.97	0.47 UG/L
LEAD, DISS.	FILTRATE	5	4	0	0.00	0.360	U	2.40	U	0.98	0.98 UG/L
LEAD, DISS.	FILTRATE	10	2	0	0.00	0.500	U	0.70	U	0.60	0.14 UG/L
LEAD, DISS.	FILTRATE	13	1	1	100.00	0.600	V/V	0.60	V/V	0.60	UG/L
LEAD, TOTAL	WATER	1	3	1	33.33	0.900	U	2.30	U	1.73	0.74 UG/L
LEAD, TOTAL	WATER	2	2	0	0.00	0.360	U	0.40	U	0.38	0.03 UG/L
LEAD, TOTAL	WATER	3	4	2	50.00	0.360	U	1.80	UJ	1.08	0.53 UG/L
LEAD, TOTAL	WATER	4	3	1	33.33	1.000	V/V	1.50	U	1.30	0.28 UG/L
LEAD, TOTAL	WATER	5	4	0	0.00	0.360	U	11.00	U	3.77	4.96 UG/L
LEAD, TOTAL	WATER	10	2	0	0.00	0.600	U	1.00	U	0.80	0.28 UG/L
LEAD, TOTAL	WATER	13	1	0	0.00	5.200	U	5.20	U	5.20	UG/L
MAGNESIUM, DISS.	FILTRATE	1	3	3	100.00	9550.000	V/V	10400.00	V/V	10050.00	444.41 UG/L
MAGNESIUM, DISS.	FILTRATE	2	2	2	100.00	10200.00	V/V	10500.00	V/V	10350.00	212.13 UG/L
MAGNESIUM, DISS.	FILTRATE	3	4	4	100.00	6170.000	V/V	6310.00	V/V	6255.00	60.28 UG/L
MAGNESIUM, DISS.	FILTRATE	4	2	2	100.00	7030.000	V/V	10300.00	V/V	8865.00	2312.24 UG/L
MAGNESIUM, DISS.	FILTRATE	5	4	4	100.00	4550.000	V/V	9580.00	V/V	5930.00	2440.26 UG/L
MAGNESIUM, DISS.	FILTRATE	10	2	2	100.00	10100.00	V/V	10400.00	V/V	10250.00	212.13 UG/L
MAGNESIUM, DISS.	FILTRATE	13	1	1	100.00	8340.000	V/V	8340.00	V/V	8340.00	UG/L
MAGNESIUM, TOTAL	WATER	1	3	3	100.00	10000.00	V/V	10500.00	V/V	10233.33	251.66 UG/L
MAGNESIUM, TOTAL	WATER	2	2	2	100.00	10200.00	V/V	10300.00	V/V	10250.00	70.71 UG/L
MAGNESIUM, TOTAL	WATER	3	4	4	100.00	6160.000	V/V	6450.00	V/V	6335.00	134.29 UG/L
MAGNESIUM, TOTAL	WATER	4	2	2	100.00	6790.000	V/V	10300.00	V/V	8545.00	2481.94 UG/L
MAGNESIUM, TOTAL	WATER	5	4	4	100.00	4520.000	V/V	9600.00	V/V	5920.00	2460.31 UG/L
MAGNESIUM, TOTAL	WATER	10	2	2	100.00	10200.00	V/V	10500.00	V/V	10350.00	212.13 UG/L
MAGNESIUM, TOTAL	WATER	13	1	1	100.00	6330.000	V/V	6330.00	V/V	6330.00	UG/L
MERCURY, DISS.	FILTRATE	1	3	0	0.00	0.200	U	0.20	U	0.20	0.00 UG/L
MERCURY, DISS.	FILTRATE	2	2	0	0.00	0.200	U	0.20	U	0.20	0.00 UG/L
MERCURY, DISS.	FILTRATE	3	4	1	25.00	0.200	U	0.52	V/V	0.28	0.16 UG/L
MERCURY, DISS.	FILTRATE	4	3	0	0.00	0.200	U	0.20	U	0.20	0.00 UG/L
MERCURY, DISS.	FILTRATE	5	4	0	0.00	0.200	U	0.20	U	0.20	0.00 UG/L
MERCURY, DISS.	FILTRATE	10	2	0	0.00	0.200	U	0.20	U	0.20	0.00 UG/L
MERCURY, DISS.	FILTRATE	13	1	0	0.00	0.200	U	0.20	U	0.20	0.00 UG/L
POTASSIUM, DISS.	FILTRATE	1	3	3	100.00	1680.000	V/V	1710.00	V/V	1690.00	17.32 UG/L
POTASSIUM, DISS.	FILTRATE	2	2	2	100.00	1550.000	V/V	1750.00	V/V	1650.00	141.42 UG/L
POTASSIUM, DISS.	FILTRATE	3	4	4	100.00	1340.000	V/V	1380.00	V/V	1355.00	10.00 UG/L
POTASSIUM, DISS.	FILTRATE	4	2	2	100.00	1530.000	V/V	1730.00	V/V	1630.00	141.42 UG/L
POTASSIUM, DISS.	FILTRATE	5	4	4	100.00	1340.000	V/V	1660.00	V/V	1485.00	140.59 UG/L
POTASSIUM, DISS.	FILTRATE	10	2	2	100.00	1710.000	V/V	1840.00	V/V	1775.00	91.92 UG/L
POTASSIUM, DISS.	FILTRATE	13	1	1	100.00	1300.000	V/V	1300.00	V/V	1300.00	UG/L
POTASSIUM, TOTAL	WATER	1	3	3	100.00	1730.000	V/V	1840.00	V/V	1773.33	58.59 UG/L
POTASSIUM, TOTAL	WATER	2	2	2	100.00	1560.000	V/V	1770.00	V/V	1665.00	148.48 UG/L
POTASSIUM, TOTAL	WATER	3	4	4	100.00	1250.000	V/V	1510.00	V/V	1385.00	108.09 UG/L

TABLE B.2
SUMMARY STATISTICS FOR WATER DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS=INORGANICS
(CONTINUED)

ANALYSIS	SAMPLE MATRIX	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. SAMPLE DEV. UNITS
POTASSIUM, TOTAL	WATER	4	2	2	100.00	1380.000	V/V	1830.00	V/V	1555.00	108.07 UG/L
POTASSIUM, TOTAL	WATER	5	4	4	100.00	1250.000	V/V	1790.00	V/V	1485.00	238.54 UG/L
POTASSIUM, TOTAL	WATER	10	2	2	100.00	1600.000	V/V	1840.00	V/V	1620.00	28.28 UG/L
POTASSIUM, TOTAL	WATER	13	1	1	100.00	1330.000	V/V	1330.00	V/V	1330.00	0.00 UG/L
SILVER, DISS.	FILTRATE	1	3	0	0.00	0.740	U	0.74	UJ	0.74	0.00 UG/L
SILVER, DISS.	FILTRATE	2	2	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SILVER, DISS.	FILTRATE	3	4	1	25.00	0.740	U	13.00	V/V	3.81	6.13 UG/L
SILVER, DISS.	FILTRATE	4	3	0	0.00	0.740	U	0.74	UJ	0.74	0.00 UG/L
SILVER, DISS.	FILTRATE	5	4	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SILVER, DISS.	FILTRATE	10	2	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SILVER, DISS.	FILTRATE	13	1	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SILVER, TOTAL	WATER	1	3	0	0.00	0.740	UJ	0.74	UJ	0.74	0.00 UG/L
SILVER, TOTAL	WATER	2	2	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SILVER, TOTAL	WATER	3	4	1	25.00	0.740	U	4.80	V/V	1.76	2.03 UG/L
SILVER, TOTAL	WATER	4	3	0	0.00	0.740	U	0.74	UJ	0.74	0.00 UG/L
SILVER, TOTAL	WATER	5	4	0	0.00	0.740	U	0.74	UJ	0.74	0.00 UG/L
SILVER, TOTAL	WATER	10	2	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SILVER, TOTAL	WATER	13	1	0	0.00	0.740	U	0.74	U	0.74	0.00 UG/L
SODIUM, DISS.	FILTRATE	1	3	3	100.00	4480.000	V/V	5260.00	V/V	4910.00	389.74 UG/L
SODIUM, DISS.	FILTRATE	2	2	2	100.00	4760.000	V/V	5220.00	V/V	5000.00	311.13 UG/L
SODIUM, DISS.	FILTRATE	3	4	4	100.00	3060.000	V/V	4150.00	V/V	3440.00	484.22 UG/L
SODIUM, DISS.	FILTRATE	4	2	2	100.00	3780.000	V/V	4910.00	V/V	4335.00	813.17 UG/L
SODIUM, DISS.	FILTRATE	5	4	4	100.00	4310.000	V/V	5140.00	V/V	4715.00	339.26 UG/L
SODIUM, DISS.	FILTRATE	10	2	2	100.00	4940.000	V/V	5080.00	V/V	5000.00	84.85 UG/L
SODIUM, DISS.	FILTRATE	13	1	1	100.00	2640.000	V/V	2640.00	V/V	2640.00	0.00 UG/L
SODIUM, TOTAL	WATER	1	3	3	100.00	4720.000	V/V	5240.00	V/V	4928.67	275.92 UG/L
SODIUM, TOTAL	WATER	2	2	2	100.00	4760.000	V/V	5130.00	V/V	4945.00	261.83 UG/L
SODIUM, TOTAL	WATER	3	4	4	100.00	3050.000	V/V	4180.00	V/V	3420.00	516.07 UG/L
SODIUM, TOTAL	WATER	4	2	2	100.00	3820.000	V/V	4830.00	V/V	4225.00	855.60 UG/L
SODIUM, TOTAL	WATER	5	4	4	100.00	4120.000	V/V	5100.00	V/V	4850.00	402.82 UG/L
SODIUM, TOTAL	WATER	10	2	2	100.00	4800.000	V/V	5030.00	V/V	4985.00	81.92 UG/L
SODIUM, TOTAL	WATER	13	1	1	100.00	2440.000	V/V	2440.00	V/V	2440.00	0.00 UG/L
ZINC, DISS.	FILTRATE	1	3	0	0.00	3.400	U	14.00	U	9.13	5.35 UG/L
ZINC, DISS.	FILTRATE	2	2	0	0.00	3.400	U	12.00	U	7.70	6.08 UG/L
ZINC, DISS.	FILTRATE	3	4	0	0.00	3.400	U	15.00	U	8.63	5.35 UG/L
ZINC, DISS.	FILTRATE	4	2	0	0.00	3.400	U	3.40	U	3.40	0.00 UG/L
ZINC, DISS.	FILTRATE	5	4	0	0.00	3.400	U	18.00	U	7.55	5.84 UG/L
ZINC, DISS.	FILTRATE	10	2	1	50.00	7.900	V/V	21.00	V/V	14.45	9.26 UG/L
ZINC, DISS.	FILTRATE	13	1	0	0.00	13.000	U	13.00	U	13.00	0.00 UG/L
ZINC, TOTAL	WATER	1	3	0	0.00	3.400	U	11.00	U	6.37	4.06 UG/L
ZINC, TOTAL	WATER	2	2	1	50.00	8.400	U	52.00	V/V	30.20	30.83 UG/L
ZINC, TOTAL	WATER	3	4	0	0.00	5.300	U	15.00	U	10.50	5.23 UG/L
ZINC, TOTAL	WATER	4	2	0	0.00	3.400	U	5.90	U	4.65	1.77 UG/L
ZINC, TOTAL	WATER	5	4	0	0.00	3.400	U	13.00	U	8.03	4.05 UG/L
ZINC, TOTAL	WATER	10	2	0	0.00	10.000	U	11.00	U	10.50	0.71 UG/L
ZINC, TOTAL	WATER	13	1	0	0.00	18.000	U	18.00	U	18.00	0.00 UG/L

TABLE B.2
SUMMARY STATISTICS FOR WATER DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS--ORGANICS											
ANALYSIS	SAMPLE MATRIX	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. SAMPLE DEV. UNITS
BIS(2-ETHYLNEXYL)PHTHALATE WATER		1	3	0	0.00	10.000	U	10.00	U	10.00	0.00 UG/L
BIS(2-ETHYLNEXYL)PHTHALATE WATER		2	2	0	0.00	10.000	U	10.00	U	10.00	0.00 UG/L
BIS(2-ETHYLNEXYL)PHTHALATE WATER		3	4	0	0.00	10.000	U	10.00	U	10.00	0.00 UG/L
BIS(2-ETHYLNEXYL)PHTHALATE WATER		4	2	0	0.00	10.000	U	10.00	U	10.00	0.00 UG/L
BIS(2-ETHYLNEXYL)PHTHALATE WATER		5	4	0	0.00	3.000	U	10.00	U	6.50	4.04 UG/L
BIS(2-ETHYLNEXYL)PHTHALATE WATER		10	2	1	50.00	5.000	J	10.00	UJ	7.50	3.54 UG/L
BIS(2-ETHYLNEXYL)PHTHALATE WATER		13	1	0	0.00	10.000	U	10.00	U	10.00	UG/L
TYPE OF ANALYSIS--RAD											
ANALYSIS	SAMPLE MATRIX	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. SAMPLE DEV. UNITS
CO-80	EVAPORATE	1	3	0	0.00	0.083	UJ	0.15	UJ	0.12	0.04 PCI/L
CO-80	EVAPORATE	2	2	0	0.00	0.152	UJ	0.29	UJ	0.22	0.10 PCI/L
CO-80	EVAPORATE	3	4	0	0.00	0.109	UJ	0.24	UJ	0.16	0.06 PCI/L
CO-80	EVAPORATE	4	2	0	0.00	0.152	UJ	0.16	UJ	0.16	0.01 PCI/L
CO-80	EVAPORATE	5	4	1	25.00	0.080	UJ	0.18	V/V	0.14	0.04 PCI/L
CO-80	EVAPORATE	10	2	0	0.00	0.087	UJ	0.10	UJ	0.09	0.03 PCI/L
CO-80	EVAPORATE	13	1	0	0.00	0.217	UJ	0.22	UJ	0.22	PCI/L
CS-137	EVAPORATE	1	3	0	0.00	0.083	UJ	0.13	UJ	0.11	0.02 PCI/L
CS-137	RESIDUE	1	3	0	0.00	6.375	UJ	14.42	UJ	10.62	4.04 PCI/G
CS-137	EVAPORATE	2	2	1	50.00	0.009	V/V	0.16	UJ	0.08	0.11 PCI/L
CS-137	EVAPORATE	2	2	1	50.00	11.943	V/V	15.72	UJ	13.83	2.67 PCI/G
CS-137	RESIDUE	3	4	3	75.00	0.095	J	0.22	V/V	0.15	0.05 PCI/L
CS-137	RESIDUE	3	3	0	0.00	2.578	UJ	5.63	UJ	4.06	1.53 PCI/G
CS-137	EVAPORATE	4	2	2	100.00	0.077	J	0.13	J	0.10	0.04 PCI/L
CS-137	RESIDUE	4	2	1	50.00	12.990	UJ	18.87	V/V	16.33	4.72 PCI/G
CS-137	EVAPORATE	5	4	1	25.00	0.050	V/V	0.14	UJ	0.10	0.04 PCI/L
CS-137	RESIDUE	5	3	1	33.33	11.338	UJ	17.34	UJ	14.57	3.03 PCI/G
CS-137	EVAPORATE	10	2	0	0.00	0.075	UJ	0.12	UJ	0.10	0.03 PCI/L
CS-137	RESIDUE	10	1	0	0.00	57.102	UJ	57.10	UJ	57.10	PCI/G
CS-137	EVAPORATE	13	1	0	0.00	0.205	UJ	0.21	UJ	0.21	PCI/L
CS-137	RESIDUE	13	1	0	0.00	6.669	UJ	6.67	UJ	6.67	PCI/G
SR-90	WATER	1	3	3	100.00	0.200	J	1.50	J	0.70	0.70 PCI/L
SR-90	WATER	2	2	2	100.00	-0.500	J	-0.10	J	-0.30	0.28 PCI/L
SR-90	WATER	3	4	4	100.00	-0.400	J	0.20	J	-0.13	0.28 PCI/L
SR-90	WATER	4	1	1	100.00	-0.400	J	-0.40	J	-0.40	PCI/L
SR-90	WATER	5	3	3	100.00	-0.400	J	0.70	J	-0.00	0.61 PCI/L
SR-90	WATER	10	2	2	100.00	-0.400	J	-0.30	J	-0.35	0.07 PCI/L
SR-90	WATER	13	1	1	100.00	-0.100	J	-0.10	J	-0.10	PCI/L

Appendix C

SUMMARY STATISTICS FOR SEDIMENT SAMPLES

CONTENTS OF APPENDIX C

Table C.1	Summary statistics for sediment core data summarized for all reaches sorted by type of analysis and decreasing percentage detected	C-3
Table C.2	Summary statistics for sediment core data detected compounds summarized by reach sorted by type of analysis, analysis, and reach	C-6
Table C.3	Summary statistics for sediment grab data summarized for all reaches sorted by type of analysis and decreasing percentage detected	C-13
Table C.4	Summary statistics for sediment grab data detected compounds summarized by reach sorted by type of analysis, analysis, and reach . . .	C-14

TABLE C.1
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

TYPE OF ANALYSIS--INORGANICS										
ANALYSIS	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
CHROMIUM, TOTAL	332	332	100.00	3.000	J	74.30	V/V	21.55	10.44	MG/KG
LEAD, TOTAL	332	332	100.00	0.720	J	92.80	V/V	27.40	14.31	MG/KG
NICKEL, TOTAL	332	332	100.00	2.400	V/V	84.00	V/V	18.10	9.23	MG/KG
ZINC, TOTAL	332	332	100.00	14.500	J	725.00	J	110.86	89.18	MG/KG
ARSENIC, TOTAL	331	328	99.09	0.400	UJ	202.00	V/V	18.60	39.29	MG/KG
BERYLLIUM, TOTAL	332	321	96.69	0.200	U	2.20	V/V	0.82	0.45	MG/KG
COPPER, TOTAL	332	318	95.78	6.000	J	61.50	V/V	24.12	11.78	MG/KG
MERCURY, TOTAL	217	189	87.10	0.030	V/V	180.00	V/V	2.86	11.95	MG/KG
SILVER, TOTAL	174	136	78.18	0.020	U	3.00	J	0.31	0.37	MG/KG
CADMIUM, TOTAL	332	248	74.70	0.200	U	8.50	J	0.83	1.09	MG/KG
SELENIUM, TOTAL	273	76	27.84	0.200	UJ	14.30	J	0.83	1.49	MG/KG
ANTIMONY, TOTAL	169	21	12.43	0.040	U	1.50	J	0.39	0.13	MG/KG
THALLIUM, TOTAL	330	28	8.48	0.200	U	2.60	V/V	0.48	0.40	MG/KG

TYPE OF ANALYSIS--ORGANICS

ANALYSIS	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
BIS(2-ETHYLHEXYL)PHTHALATE	279	101	36.20	0.180	J	10.00	U	1.28	1.38	MG/KG
AROCLOR-1254	275	54	19.64	0.033	UJ	43.00	UJ	5.16	7.51	MG/KG
FLUORANTHENE	279	41	14.70	0.060	UJ	10.00	U	1.23	1.60	MG/KG
PYRENE	279	32	11.47	0.094	J	10.00	U	1.23	1.59	MG/KG
PHENANTHRENE	279	19	6.81	0.059	J	10.00	U	1.24	1.59	MG/KG
BENZO(B)FLUORANTHENE	279	5	1.79	0.170	UJ	10.00	U	1.27	1.60	MG/KG
CHRYSENE	279	5	1.43	0.160	UJ	10.00	U	1.25	1.58	MG/KG
BENZO(K)FLUORANTHENE	279	4	1.43	0.058	J	10.00	U	1.27	1.60	MG/KG
DI-N-BUTYLPHTHALATE	279	4	1.43	0.042	J	10.00	U	1.27	1.60	MG/KG
2-METHYLNAPHTHALENE	279	4	1.43	0.016	UJ	10.00	U	1.27	1.60	MG/KG
4,4'-DDD	273	3	1.10	0.170	UJ	4.30	UJ	0.62	0.75	MG/KG
BENZO(A)ANTHRAcene	278	3	1.08	0.170	UJ	10.00	U	1.28	1.60	MG/KG
NAPHTHALENE	279	3	1.08	0.038	J	10.00	U	1.27	1.60	MG/KG
4,4'-DBT	273	2	0.73	0.018	UJ	4.30	UJ	0.62	0.75	MG/KG
ANTHRAcene	279	2	0.72	0.059	UJ	10.00	U	1.26	1.58	MG/KG
BENZO(A)PYRENE	279	2	0.72	0.170	UJ	10.00	U	1.27	1.60	MG/KG
TOXAPHENE	261	1	0.38	0.190	UJ	43.00	UJ	5.68	6.99	MG/KG
AROCLOR-1280	273	1	0.37	0.190	UJ	43.00	UJ	6.20	7.52	MG/KG
DIETHYLPHTHALATE	279	1	0.36	0.076	U	10.00	U	1.27	1.60	MG/KG
DIMETHYLPHTHALATE	279	1	0.36	0.170	UJ	10.00	U	1.27	1.60	MG/KG
INDENO(1,2,3-CD)PYRENE	279	1	0.36	0.170	UJ	10.00	U	1.27	1.60	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
ACENAPHTHENE	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
ACENAPHTHYLENE	279	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
ALDRIN	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
ALPHA-BHC	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
ALPHA-CHLORDANE	241	0	0.00	0.009	UJ	1.80	UJ	2.38	3.08	MG/KG

TABLE C.1
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

----- TYPE OF ANALYSIS-ORGANICS
(CONTINUED)

ANALYSIS	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MTN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
AROCOR-1018	241	0	0.00	0.085	UJ	18.00	UJ	2.38	3.08	MG/KG
AROCOR-1221	241	0	0.00	0.095	UJ	18.00	UJ	2.38	3.08	MG/KG
AROCOR-1232	241	0	0.00	0.085	UJ	18.00	UJ	2.38	3.08	MG/KG
AROCOR-1242	241	0	0.00	0.085	UJ	18.00	UJ	2.38	3.08	MG/KG
AROCOR-1248	241	0	0.00	0.085	UJ	18.00	UJ	2.38	3.08	MG/KG
BENZO(O,H,1)PERYLENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
BENZOIC ACID	278	0	0.00	0.400	UJ	51.00	U	6.17	7.78	MG/KG
BENZYL ALCOHOL	271	0	0.00	0.170	UJ	10.00	U	1.29	1.62	MG/KG
BETA-BHC	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
BIS(2-CHLOROETHOXY)METHANE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
BIS(2-CHLOROETHYL)ETHER	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
BIS(2-CHLOROISOPROPYL)ETHE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
BUTYLBENZYLPHthalATE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
DELTA-BHC	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
DI-N-OCTYLPHthalATE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
DIBENZ(A,H)ANTHRACENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
DIBENZOFURAN	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
DIELDRIN	273	0	0.00	0.018	UJ	4.30	UJ	0.62	0.75	MG/KG
ENDOSULFAN I	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
ENDOSULFAN II	273	0	0.00	0.018	UJ	4.30	UJ	0.62	0.75	MG/KG
ENDOSULFAN SULFATE	273	0	0.00	0.018	UJ	4.30	UJ	0.62	0.75	MG/KG
ENDRIN	273	0	0.00	0.018	UJ	4.30	UJ	0.62	0.75	MG/KG
ENDRIN KETONE	273	0	0.00	0.018	UJ	4.30	UJ	0.62	0.75	MG/KG
FLUORENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
GAMMA-BHC (LINDANE)	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
GAMMA-CHLORDANE	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
HEPTACHLOR	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
HEPTACHLOR EPOXIDE	241	0	0.00	0.009	UJ	1.80	UJ	0.24	0.31	MG/KG
HEXACHLOROBENZENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
HEXACHLOROBUTADIENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
HEXACHLOROCYCLOPENTADIENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
HEXACHLOROETHANE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
ISOPHORONE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
METHOXYCHLOR	273	0	0.00	0.009	UJ	21.00	UJ	3.08	3.72	MG/KG
N-NITROSODIPHENYLAMINE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
NITROBENZENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
PENTACHLOROPHENOL	278	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG
PHENOL	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
1,2-DICHLOROBENZENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
1,2,4-TRICHLOROBENZENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
1,3-DICHLOROBENZENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
1,4-DICHLOROBENZENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2-CHLORONAPHTHALENE	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2-CHLOROPHENOL	278	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2-METHYLPHENOL	271	0	0.00	0.170	UJ	10.00	U	1.29	1.62	MG/KG
2-NITROANILINE	278	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG

TABLE C.1
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED
TYPE OF ANALYSIS=ORGANICS
(CONTINUED)

ANALYSIS	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
2-NITROPHENOL	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2,4-DICHLOROPHENOL	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2,4-DIMETHYLPHENOL	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2,4-DINITROPHENOL	256	0	0.00	0.830	UJ	51.00	U	6.48	8.05	MG/KG
2,4-DINITROTOLUENE	279	0	0.00	0.170	UJ	10.00	U	1.28	1.58	MG/KG
2,4,5-TRICHLOROPHENOL	279	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG
2,4,8-TRICHLOROPHENOL	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
2,6-DINITROTOLUENE	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
3-NITROANILINE	279	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG
3,3'-DICHLOROBENZIDINE	279	0	0.00	0.330	UJ	21.00	U	2.55	3.20	MG/KG
4-BROMOPHENYL-PHENYLETHYR	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
4-CHLORO-3-METHYLPHENOL	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
4-CHLOROANILINE	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
4-CHLOROPHENYL-PHENYLETHYR	279	0	0.00	0.170	UJ	10.00	U	1.27	1.60	MG/KG
4-METHYLPHENOL	279	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG
4-NITROANILINE	279	0	0.00	0.830	UJ	51.00	U	6.24	7.83	MG/KG
4-NITROPHENOL	266	0	0.00	0.830	UJ	4.30	UJ	0.62	0.75	MG/KG
4,4'-DDE	273	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG
4,6-DINITRO-2-METHYLPHENOL	279	0	0.00	0.830	UJ	51.00	U	6.18	7.77	MG/KG

ANALYSIS	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
AM-241	92	92	100.00	0.010	J	0.88	J	0.14	0.13	PCI/G (DRY)
CM-243,244	92	92	100.00	0.000	J	0.11	J	0.02	0.01	PCI/G (DRY)
CM-245,246	35	35	100.00	0.013	J	0.08	J	0.02	0.02	PCI/G (DRY)
CM-248	34	34	100.00	0.013	J	0.08	J	0.03	0.02	PCI/G (DRY)
PU-238	91	91	100.00	0.010	J	0.23	J	0.02	0.03	PCI/G (DRY)
PU-239,240	91	91	100.00	0.010	J	1.57	J	0.14	0.22	PCI/G (DRY)
SR-90	93	93	100.00	-0.060	J	1.25	J	0.25	0.28	PCI/G (DRY)
U-234	91	91	100.00	0.290	J	8.91	J	2.43	1.85	PCI/G (DRY)
U-235	91	91	100.00	0.010	J	0.68	J	0.10	0.10	PCI/G (DRY)
U-238	91	91	100.00	0.330	J	6.92	J	2.16	1.64	PCI/G (DRY)
CS-137	416	385	92.55	0.003	V/V	423.75	V/V	12.37	30.83	PCI/G (DRY)
CO-60	366	130	35.52	0.014	UJ	2.44	V/V	0.23	0.23	PCI/G (DRY)

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS=INORGANICS											
ANALYSIS	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
ANTIMONY, TOTAL	1	14	0	0.00	0.300	U	0.70	U	0.48	0.12	MG/KG
ANTIMONY, TOTAL	3	29	11	37.93	0.300	U	0.40	V/V	0.31	0.04	MG/KG
ANTIMONY, TOTAL	4	35	1	2.86	0.270	UJ	0.60	UJ	0.35	0.07	MG/KG
ANTIMONY, TOTAL	5	40	2	5.00	0.260	UJ	1.50	J	0.48	0.20	MG/KG
ANTIMONY, TOTAL	7	28	6	21.43	0.190	V/V	0.61	J	0.32	0.10	MG/KG
ANTIMONY, TOTAL	10	23	1	4.35	0.040	U	0.70	UJ	0.43	0.11	MG/KG
ARSENIC, TOTAL	1	21	20	95.24	3.400	V/V	20.60	J	9.14	3.99	MG/KG
ARSENIC, TOTAL	2	58	58	100.00	2.400	V/V	10.60	J	5.88	2.05	MG/KG
ARSENIC, TOTAL	3	78	79	100.00	0.700	J	202.00	V/V	51.21	71.22	MG/KG
ARSENIC, TOTAL	4	52	52	100.00	2.800	J	20.30	J	10.26	3.90	MG/KG
ARSENIC, TOTAL	5	70	70	100.00	3.100	J	15.30	J	6.62	2.38	MG/KG
ARSENIC, TOTAL	7	28	28	100.00	4.200	V/V	20.10	V/V	11.48	4.26	MG/KG
ARSENIC, TOTAL	10	23	21	91.30	0.400	UJ	8.80	V/V	5.20	2.12	MG/KG
BERYLLIUM, TOTAL	1	21	21	100.00	0.320	V/V	1.00	V/V	0.74	0.15	MG/KG
BERYLLIUM, TOTAL	2	59	48	81.36	0.200	U	1.60	V/V	0.70	0.30	MG/KG
BERYLLIUM, TOTAL	3	78	79	100.00	0.200	V/V	2.20	V/V	0.78	0.65	MG/KG
BERYLLIUM, TOTAL	4	52	52	100.00	0.300	V/V	1.60	V/V	0.80	0.29	MG/KG
BERYLLIUM, TOTAL	5	70	70	100.00	0.430	V/V	2.10	V/V	1.25	0.28	MG/KG
BERYLLIUM, TOTAL	7	28	28	100.00	0.640	V/V	1.70	V/V	1.04	0.26	MG/KG
BERYLLIUM, TOTAL	10	23	23	100.00	0.880	V/V	1.80	V/V	1.29	0.21	MG/KG
CADMIUM, TOTAL	1	21	13	61.90	0.300	U	1.30	V/V	0.63	0.35	MG/KG
CADMIUM, TOTAL	2	59	35	59.32	0.200	U	0.70	V/V	0.38	0.13	MG/KG
CADMIUM, TOTAL	3	78	65	82.28	0.200	U	3.60	J	0.76	0.63	MG/KG
CADMIUM, TOTAL	4	52	48	92.31	0.240	V/V	2.20	V/V	0.86	0.49	MG/KG
CADMIUM, TOTAL	5	70	69	98.57	0.430	V/V	8.50	J	2.16	1.75	MG/KG
CADMIUM, TOTAL	7	28	4	14.29	0.260	U	1.80	J	0.60	0.33	MG/KG
CADMIUM, TOTAL	10	23	14	60.87	0.400	U	1.40	V/V	0.64	0.23	MG/KG
CHROMIUM, TOTAL	1	21	21	100.00	7.100	V/V	24.00	V/V	18.14	5.33	MG/KG
CHROMIUM, TOTAL	2	59	59	100.00	5.000	V/V	28.00	V/V	14.34	4.12	MG/KG
CHROMIUM, TOTAL	3	78	79	100.00	3.000	J	40.00	J	17.48	10.48	MG/KG
CHROMIUM, TOTAL	4	52	52	100.00	5.800	V/V	47.70	V/V	21.08	10.08	MG/KG
CHROMIUM, TOTAL	5	70	70	100.00	16.000	V/V	51.90	V/V	31.70	7.20	MG/KG
CHROMIUM, TOTAL	7	28	28	100.00	17.300	V/V	74.30	V/V	27.30	11.54	MG/KG
CHROMIUM, TOTAL	10	23	23	100.00	12.600	V/V	33.60	V/V	22.24	4.69	MG/KG
COPPER, TOTAL	1	21	21	100.00	15.000	V/V	52.00	J	28.43	8.01	MG/KG
COPPER, TOTAL	2	59	59	100.00	6.000	J	32.00	V/V	15.24	5.13	MG/KG
COPPER, TOTAL	3	78	79	100.00	6.000	J	47.00	J	19.52	12.87	MG/KG
COPPER, TOTAL	4	52	52	100.00	9.400	V/V	53.80	V/V	25.08	9.79	MG/KG
COPPER, TOTAL	5	70	70	100.00	13.000	J	57.50	V/V	34.36	8.23	MG/KG
COPPER, TOTAL	7	28	28	100.00	10.400	V/V	48.40	V/V	28.31	10.31	MG/KG
COPPER, TOTAL	10	23	9	39.13	11.700	U	61.50	J	20.37	11.35	MG/KG
LEAD, TOTAL	1	21	21	100.00	18.600	J	47.80	J	36.42	7.02	MG/KG
LEAD, TOTAL	2	59	59	100.00	8.600	V/V	44.30	V/V	27.81	9.11	MG/KG
LEAD, TOTAL	3	78	79	100.00	0.800	J	69.90	J	14.83	10.50	MG/KG
LEAD, TOTAL	4	52	52	100.00	5.700	V/V	37.60	J	20.74	8.80	MG/KG
LEAD, TOTAL	5	70	70	100.00	0.720	J	92.80	V/V	43.95	12.08	MG/KG
LEAD, TOTAL	7	28	28	100.00	3.600	J	35.90	J	21.39	7.05	MG/KG

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS-INORGANICS
(CONTINUED)

ANALYSIS	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
LEAD, TOTAL	10	23	23	100.00	24.300	J	45.30	J	33.24	5.75	MG/KG
MERCURY, TOTAL	1	21	21	100.00	0.040	V/V	0.17	V/V	0.10	0.04	MG/KG
MERCURY, TOTAL	2	14	14	100.00	0.040	J	0.79	J	0.25	0.28	MG/KG
MERCURY, TOTAL	3	28	28	100.00	0.030	V/V	4.59	V/V	0.38	1.00	MG/KG
MERCURY, TOTAL	4	52	52	100.00	0.081	V/V	180.00	V/V	9.66	23.12	MG/KG
MERCURY, TOTAL	5	51	51	100.00	0.080	V/V	10.93	V/V	1.88	2.43	MG/KG
MERCURY, TOTAL	7	28	0	0.00	0.050	U	0.34	U	0.16	0.09	MG/KG
MERCURY, TOTAL	10	23	23	100.00	0.050	V/V	0.14	V/V	0.08	0.02	MG/KG
NICKEL, TOTAL	1	21	21	100.00	10.000	V/V	32.00	V/V	20.14	8.02	MG/KG
NICKEL, TOTAL	2	59	59	100.00	4.000	V/V	28.00	V/V	13.73	4.82	MG/KG
NICKEL, TOTAL	3	79	79	100.00	3.000	V/V	64.00	V/V	14.29	11.11	MG/KG
NICKEL, TOTAL	4	52	52	100.00	2.400	V/V	57.70	V/V	22.55	10.83	MG/KG
NICKEL, TOTAL	5	70	70	100.00	13.000	V/V	51.00	V/V	22.90	6.80	MG/KG
NICKEL, TOTAL	7	28	28	100.00	14.800	V/V	48.70	V/V	25.77	6.25	MG/KG
NICKEL, TOTAL	10	23	23	100.00	14.100	V/V	30.30	V/V	20.90	3.58	MG/KG
SELENIUM, TOTAL	1	21	12	57.14	0.300	UJ	1.70	J	0.63	0.38	MG/KG
SELENIUM, TOTAL	2	59	5	8.47	0.300	U	1.30	UJ	0.33	0.13	MG/KG
SELENIUM, TOTAL	3	79	31	39.24	0.200	UJ	14.30	J	1.34	2.50	MG/KG
SELENIUM, TOTAL	4	29	8	27.59	0.270	UJ	2.00	J	0.63	0.51	MG/KG
SELENIUM, TOTAL	5	40	0	0.00	0.300	UJ	0.82	UJ	0.47	0.11	MG/KG
SELENIUM, TOTAL	7	28	20	71.43	0.270	UJ	5.50	J	1.59	1.30	MG/KG
SELENIUM, TOTAL	10	17	0	0.00	0.400	UJ	0.70	UJ	0.46	0.08	MG/KG
SILVER, TOTAL	1	21	21	100.00	0.180	V/V	0.42	V/V	0.25	0.07	MG/KG
SILVER, TOTAL	2	59	43	72.88	0.030	U	3.00	J	0.31	0.43	MG/KG
SILVER, TOTAL	3	14	12	85.71	0.020	UJ	1.60	J	0.30	0.46	MG/KG
SILVER, TOTAL	4	2	2	100.00	1.800	J	1.80	J	1.80	0.00	MG/KG
SILVER, TOTAL	5	50	39	78.00	0.140	UJ	0.93	V/V	0.39	0.26	MG/KG
SILVER, TOTAL	7	28	19	67.86	0.020	U	0.70	U	0.14	0.22	MG/KG
SILVER, TOTAL	1	21	0	0.00	0.300	U	0.40	U	0.41	0.12	MG/KG
THALLIUM, TOTAL	2	59	0	0.00	0.300	U	0.40	U	0.31	0.03	MG/KG
THALLIUM, TOTAL	3	79	27	34.18	0.200	U	2.60	V/V	0.75	0.74	MG/KG
THALLIUM, TOTAL	4	52	0	0.00	0.270	U	0.60	U	0.34	0.06	MG/KG
THALLIUM, TOTAL	5	88	0	0.00	0.300	U	0.82	UJ	0.47	0.11	MG/KG
THALLIUM, TOTAL	7	28	1	3.57	0.210	U	0.83	V/V	0.42	0.13	MG/KG
THALLIUM, TOTAL	10	23	0	0.00	0.400	U	0.70	U	0.44	0.07	MG/KG
ZINC, TOTAL	1	21	21	100.00	59.300	V/V	183.00	J	104.60	31.98	MG/KG
ZINC, TOTAL	2	59	59	100.00	28.300	J	167.00	J	73.88	22.08	MG/KG
ZINC, TOTAL	3	79	79	100.00	14.500	J	725.00	J	54.92	78.77	MG/KG
ZINC, TOTAL	4	52	52	100.00	34.100	V/V	185.00	J	82.98	32.88	MG/KG
ZINC, TOTAL	5	70	70	100.00	94.700	V/V	437.00	J	246.51	69.15	MG/KG
ZINC, TOTAL	7	28	28	100.00	41.000	V/V	116.00	J	80.33	20.85	MG/KG
ZINC, TOTAL	10	23	23	100.00	55.500	V/V	134.00	V/V	87.75	17.01	MG/KG

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS=ORGANICS											
ANALYSIS	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
ANTHRACENE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
ANTHRACENE	2	46	0	0.00	0.330	UJ	0.63	UJ	0.51	0.06	MG/KG
ANTHRACENE	3	64	1	1.56	0.059	UJ	0.69	UJ	0.44	0.07	MG/KG
ANTHRACENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
ANTHRACENE	5	63	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
ANTHRACENE	10	22	1	4.55	0.460	V/V	10.00	U	4.43	1.88	MG/KG
AROCLOR-1254	1	23	0	0.00	0.230	UJ	10.00	UJ	2.75	2.24	MG/KG
AROCLOR-1254	2	47	1	2.13	0.220	UJ	10.00	UJ	2.70	2.17	MG/KG
AROCLOR-1254	3	63	2	3.17	0.190	UJ	9.70	UJ	1.29	1.44	MG/KG
AROCLOR-1254	4	57	51	89.47	0.033	NJ	14.00	U	0.69	2.14	MG/KG
AROCLOR-1254	5	63	0	0.00	0.560	UJ	43.00	UJ	11.64	9.36	MG/KG
AROCLOR-1254	10	22	0	0.00	1.600	UJ	41.00	UJ	17.12	6.94	MG/KG
AROCLOR-1260	1	23	0	0.00	0.230	UJ	10.00	UJ	2.75	2.24	MG/KG
AROCLOR-1260	2	47	1	2.13	0.220	UJ	10.00	UJ	2.70	2.17	MG/KG
AROCLOR-1260	3	63	0	0.00	0.190	UJ	9.70	UJ	1.35	1.46	MG/KG
AROCLOR-1260	4	55	0	0.00	0.230	UJ	22.00	U	5.61	5.50	MG/KG
AROCLOR-1260	5	63	0	0.00	0.560	UJ	43.00	UJ	11.64	9.36	MG/KG
AROCLOR-1260	10	22	0	0.00	1.600	UJ	41.00	UJ	17.12	6.94	MG/KG
BENZO(A)ANTHRACENE	1	25	1	4.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
BENZO(A)ANTHRACENE	2	46	1	2.17	0.330	UJ	0.63	UJ	0.50	0.06	MG/KG
BENZO(A)ANTHRACENE	3	63	1	1.59	0.210	UJ	0.69	UJ	0.44	0.06	MG/KG
BENZO(A)ANTHRACENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
BENZO(A)ANTHRACENE	5	63	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
BENZO(A)ANTHRACENE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.66	MG/KG
BENZO(A)PYRENE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
BENZO(A)PYRENE	2	46	1	2.17	0.330	UJ	0.63	UJ	0.50	0.06	MG/KG
BENZO(A)PYRENE	3	64	1	1.56	0.180	UJ	0.69	UJ	0.44	0.06	MG/KG
BENZO(A)PYRENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
BENZO(A)PYRENE	5	63	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
BENZO(A)PYRENE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.66	MG/KG
BENZO(B)FLUORANTHENE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
BENZO(B)FLUORANTHENE	2	46	3	6.52	0.190	J	0.63	UJ	0.49	0.07	MG/KG
BENZO(B)FLUORANTHENE	3	64	2	3.13	0.180	J	0.69	UJ	0.44	0.07	MG/KG
BENZO(B)FLUORANTHENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
BENZO(B)FLUORANTHENE	5	63	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
BENZO(B)FLUORANTHENE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.66	MG/KG
BENZO(K)FLUORANTHENE	1	25	1	4.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
BENZO(K)FLUORANTHENE	2	46	1	2.17	0.330	UJ	0.63	UJ	0.51	0.06	MG/KG
BENZO(K)FLUORANTHENE	3	64	2	3.13	0.180	UJ	0.69	UJ	0.44	0.07	MG/KG
BENZO(K)FLUORANTHENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
BENZO(K)FLUORANTHENE	5	63	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
BENZO(K)FLUORANTHENE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.66	MG/KG
BIS(2-ETHYLHEXYL)PHTHALATE	1	25	0	0.00	0.390	UJ	3.40	U	1.23	0.79	MG/KG
BIS(2-ETHYLHEXYL)PHTHALATE	2	46	30	65.22	0.180	J	1.10	U	0.48	0.20	MG/KG
BIS(2-ETHYLHEXYL)PHTHALATE	3	64	16	25.00	0.180	J	2.60	U	0.63	0.43	MG/KG
BIS(2-ETHYLHEXYL)PHTHALATE	4	59	29	49.15	0.320	U	8.00	U	1.31	1.46	MG/KG
BIS(2-ETHYLHEXYL)PHTHALATE	5	63	24	38.10	0.590	UJ	7.50	J	1.66	1.23	MG/KG

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS-ORGANICS (CONTINUED)											
ANALYSIS	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
BIS(2-ETHYLNXYL)PHTHALATE	10	22	2	9.09	0.810	J	10.00	U	3.49	2.32	MG/KG
CHRYSENE	1	25	1	4.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
CHRYSENE	2	48	2	4.35	0.330	UJ	0.63	UJ	0.50	0.06	MG/KG
CHRYSENE	3	64	1	1.56	0.210	UJ	0.69	UJ	0.44	0.06	MG/KG
CHRYSENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
CHRYSENE	5	83	0	0.00	0.250	UJ	8.60	U	1.40	1.55	MG/KG
CHRYSENE	10	22	1	4.55	0.740	V/V	10.00	U	4.45	1.85	MG/KG
DI-N-BUTYLPHTHALATE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
DI-N-BUTYLPHTHALATE	2	46	3	6.52	0.060	J	0.63	UJ	0.47	0.12	MG/KG
DI-N-BUTYLPHTHALATE	3	64	1	1.56	0.058	J	0.69	UJ	0.44	0.07	MG/KG
DI-N-BUTYLPHTHALATE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
DI-N-BUTYLPHTHALATE	5	83	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
DI-N-BUTYLPHTHALATE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.66	MG/KG
DIETHYLPHTHALATE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
DIETHYLPHTHALATE	2	46	1	2.17	0.140	J	0.63	UJ	0.50	0.08	MG/KG
DIETHYLPHTHALATE	3	64	0	0.00	0.078	U	0.70	UJ	0.44	0.08	MG/KG
DIETHYLPHTHALATE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
DIETHYLPHTHALATE	5	83	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
DIETHYLPHTHALATE	10	22	0	0.00	2.300	U	10.00	U	4.85	1.66	MG/KG
DIMETHYLPHTHALATE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
DIMETHYLPHTHALATE	2	46	0	0.00	0.330	UJ	0.63	UJ	0.51	0.06	MG/KG
DIMETHYLPHTHALATE	3	64	1	1.56	0.390	UJ	0.69	UJ	0.44	0.05	MG/KG
DIMETHYLPHTHALATE	4	58	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
DIMETHYLPHTHALATE	5	83	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
DIMETHYLPHTHALATE	10	22	0	0.00	2.300	U	10.00	U	4.85	1.66	MG/KG
FLUORANTHENE	1	25	1	4.00	0.470	UJ	2.10	UJ	0.72	0.33	MG/KG
FLUORANTHENE	2	46	21	45.65	0.085	J	0.58	UJ	0.39	0.16	MG/KG
FLUORANTHENE	3	64	11	17.19	0.060	UJ	0.90	J	0.41	0.13	MG/KG
FLUORANTHENE	4	59	7	11.86	0.170	UJ	8.00	U	1.60	1.68	MG/KG
FLUORANTHENE	5	83	0	0.00	0.480	UJ	8.60	U	1.41	1.54	MG/KG
FLUORANTHENE	10	22	1	4.55	0.610	V/V	10.00	U	4.44	1.66	MG/KG
INDENO(1,2,3-CD)PYRENE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
INDENO(1,2,3-CD)PYRENE	2	48	1	2.17	0.280	J	0.63	UJ	0.50	0.06	MG/KG
INDENO(1,2,3-CD)PYRENE	3	64	0	0.00	0.390	UJ	0.69	UJ	0.44	0.05	MG/KG
INDENO(1,2,3-CD)PYRENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
INDENO(1,2,3-CD)PYRENE	5	83	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
INDENO(1,2,3-CD)PYRENE	10	22	0	0.00	2.300	U	10.00	U	4.85	1.66	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	2	46	0	0.00	0.330	UJ	0.63	UJ	0.51	0.06	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	3	64	0	0.00	0.390	UJ	0.69	UJ	0.44	0.05	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	4	59	1	1.69	0.170	UJ	8.00	U	1.61	1.67	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	5	83	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	10	22	0	0.00	2.300	U	10.00	U	4.85	1.66	MG/KG
N-NITROSO-DI-N-PROPYLAMINE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
NAPHTHALENE	2	46	2	4.35	0.039	J	0.63	UJ	0.49	0.09	MG/KG
NAPHTHALENE	3	64	1	1.56	0.047	J	0.69	UJ	0.44	0.07	MG/KG

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

TYPE OF ANALYSIS-ORGANICS (CONTINUED)											
ANALYSIS	REACH	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
NAPHTHALENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
NAPHTHALENE	5	63	0	0.00	0.490	UJ	6.00	U	1.41	1.54	MG/KG
NAPHTHALENE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.86	MG/KG
PHENANTHRENE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
PHENANTHRENE	2	46	9	19.57	0.059	J	0.63	UJ	0.44	0.15	MG/KG
PHENANTHRENE	3	64	8	12.50	0.072	J	0.51	J	0.41	0.10	MG/KG
PHENANTHRENE	4	59	1	1.69	0.170	UJ	8.00	U	1.61	1.67	MG/KG
PHENANTHRENE	5	63	0	0.00	0.490	UJ	6.00	U	1.41	1.54	MG/KG
PHENANTHRENE	10	22	1	4.55	0.640	V/V	10.00	U	4.44	1.86	MG/KG
PYRENE	1	25	1	4.00	0.470	UJ	2.10	UJ	0.72	0.33	MG/KG
PYRENE	2	46	17	36.96	0.094	J	0.58	UJ	0.41	0.15	MG/KG
PYRENE	3	64	10	15.63	0.094	J	0.88	J	0.41	0.09	MG/KG
PYRENE	4	59	3	5.08	0.170	UJ	8.00	U	1.60	1.68	MG/KG
PYRENE	5	63	0	0.00	0.490	UJ	6.00	U	1.41	1.54	MG/KG
PYRENE	10	22	1	4.55	0.650	V/V	10.00	U	4.44	1.86	MG/KG
TOXAPHENE	1	23	0	0.00	0.230	UJ	10.00	UJ	2.75	2.24	MG/KG
TOXAPHENE	2	47	1	2.13	0.220	UJ	10.00	UJ	2.70	2.17	MG/KG
TOXAPHENE	3	63	0	0.00	0.180	UJ	9.70	UJ	1.35	1.46	MG/KG
TOXAPHENE	4	54	0	0.00	0.230	UJ	22.00	U	5.57	5.55	MG/KG
TOXAPHENE	5	63	0	0.00	0.580	UJ	43.00	UJ	11.64	9.36	MG/KG
TOXAPHENE	10	11	0	0.00	13.000	UJ	21.00	UJ	15.84	2.58	MG/KG
2-METHYLNAPHTHALENE	1	25	0	0.00	0.470	UJ	2.10	UJ	0.71	0.33	MG/KG
2-METHYLNAPHTHALENE	2	46	2	4.35	0.042	J	0.63	UJ	0.49	0.11	MG/KG
2-METHYLNAPHTHALENE	3	64	2	3.13	0.074	J	0.69	UJ	0.43	0.08	MG/KG
2-METHYLNAPHTHALENE	4	59	0	0.00	0.170	UJ	8.00	U	1.61	1.67	MG/KG
2-METHYLNAPHTHALENE	5	63	0	0.00	0.490	UJ	8.60	U	1.41	1.54	MG/KG
2-METHYLNAPHTHALENE	10	22	0	0.00	2.300	U	10.00	U	4.65	1.86	MG/KG
4,4'-DDD	1	23	3	13.04	0.016	NJ	1.00	UJ	0.28	0.22	MG/KG
4,4'-DDD	2	47	0	0.00	0.022	UJ	1.00	UJ	0.27	0.22	MG/KG
4,4'-DDD	3	63	0	0.00	0.019	UJ	0.97	UJ	0.14	0.15	MG/KG
4,4'-DDD	4	55	0	0.00	0.023	UJ	2.20	U	0.56	0.55	MG/KG
4,4'-DDD	5	63	0	0.00	0.056	UJ	4.30	UJ	1.16	0.94	MG/KG
4,4'-DDD	10	22	0	0.00	0.160	UJ	4.10	UJ	1.71	0.89	MG/KG
4,4'-DDT	1	23	2	8.70	0.023	UJ	1.00	UJ	0.28	0.22	MG/KG
4,4'-DDT	2	47	0	0.00	0.022	UJ	1.00	UJ	0.27	0.22	MG/KG
4,4'-DDT	3	63	0	0.00	0.019	UJ	0.97	UJ	0.14	0.15	MG/KG
4,4'-DDT	4	55	0	0.00	0.023	UJ	2.20	U	0.56	0.55	MG/KG
4,4'-DDT	5	63	0	0.00	0.056	UJ	4.30	UJ	1.16	0.94	MG/KG
4,4'-DDT	10	22	0	0.00	0.160	UJ	4.10	UJ	1.71	0.89	MG/KG

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

ANALYSIS	REACH	NO. OF OBS.	NO. DETECTED	%	TYPE OF ANALYSIS=RAD				AVG. VALUE	STD. DEV.	SAMPLE UNITS
					MIN VALUE	MIN	MAX VALUE	MAX QUAL.			
AM-241	2	28	28	100.00	0.020	J	0.45	J	0.16	0.12	PCI/G (DRY)
AM-241	3	17	17	100.00	0.010	J	0.25	J	0.11	0.07	PCI/G (DRY)
AM-241	4	18	18	100.00	0.020	J	0.88	J	0.20	0.23	PCI/G (DRY)
AM-241	5	23	23	100.00	0.010	V/V	0.30	J	0.10	0.07	PCI/G (DRY)
AM-241	10	6	6	100.00	0.010	J	0.08	J	0.08	0.03	PCI/G (DRY)
AM-243	2	28	28	100.00	0.013	J	0.03	V/V	0.01	0.00	PCI/G (DRY)
AM-243	3	17	17	100.00	0.013	J	0.01	V/V	0.01	0.00	PCI/G (DRY)
AM-243	4	18	18	100.00	0.013	J	0.11	J	0.02	0.02	PCI/G (DRY)
AM-243	5	23	23	100.00	0.013	J	0.04	J	0.02	0.01	PCI/G (DRY)
AM-243	10	6	6	100.00	0.000	J	0.04	J	0.01	0.01	PCI/G (DRY)
AM-245	2	7	7	100.00	0.013	V/V	0.03	V/V	0.01	0.00	PCI/G (DRY)
AM-245	3	6	6	100.00	0.013	V/V	0.03	V/V	0.02	0.01	PCI/G (DRY)
AM-245	4	8	8	100.00	0.013	J	0.08	J	0.03	0.02	PCI/G (DRY)
AM-245	5	8	8	100.00	0.013	J	0.06	V/V	0.04	0.02	PCI/G (DRY)
AM-245	10	6	6	100.00	0.013	J	0.03	J	0.02	0.01	PCI/G (DRY)
AM-248	2	7	7	100.00	0.013	V/V	0.08	V/V	0.03	0.02	PCI/G (DRY)
AM-248	3	7	7	100.00	0.013	J	0.04	V/V	0.02	0.01	PCI/G (DRY)
AM-248	4	8	8	100.00	0.013	J	0.08	J	0.03	0.02	PCI/G (DRY)
AM-248	5	6	6	100.00	0.013	J	0.08	V/V	0.03	0.02	PCI/G (DRY)
AM-248	10	6	6	100.00	0.013	J	0.08	V/V	0.03	0.02	PCI/G (DRY)
CO-60	1	20	4	20.00	0.016	UJ	0.68	V/V	0.24	0.16	PCI/G (DRY)
CO-60	2	51	19	37.25	0.021	UJ	0.75	V/V	0.16	0.15	PCI/G (DRY)
CO-60	3	55	3	5.45	0.014	UJ	1.21	UJ	0.23	0.03	PCI/G (DRY)
CO-60	4	110	68	61.82	0.014	V/V	2.44	V/V	0.34	0.34	PCI/G (DRY)
CO-60	5	101	31	30.69	0.016	V/V	1.21	UJ	0.20	0.14	PCI/G (DRY)
CO-60	7	8	2	25.00	0.055	UJ	0.65	V/V	0.18	0.05	PCI/G (DRY)
CO-60	10	21	3	14.29	0.081	UJ	0.26	UJ	0.68	0.59	PCI/G (DRY)
CS-137	1	20	18	90.00	0.056	V/V	2.01	V/V	16.39	25.31	PCI/G (DRY)
CS-137	2	58	58	100.00	0.073	V/V	106.01	V/V	3.52	4.67	PCI/G (DRY)
CS-137	3	80	65	81.25	0.003	V/V	17.22	V/V	25.78	53.95	PCI/G (DRY)
CS-137	4	110	103	93.64	0.009	V/V	423.75	V/V	8.71	7.42	PCI/G (DRY)
CS-137	5	119	116	97.48	0.004	J	32.64	V/V	0.69	0.44	PCI/G (DRY)
CS-137	7	8	8	100.00	0.319	V/V	1.47	V/V	1.06	1.04	PCI/G (DRY)
CS-137	10	21	17	80.95	0.068	UJ	3.58	V/V	0.02	0.03	PCI/G (DRY)
PU-238	2	28	28	100.00	0.010	J	0.15	V/V	0.02	0.01	PCI/G (DRY)
PU-238	3	19	19	100.00	0.010	J	0.04	V/V	0.02	0.02	PCI/G (DRY)
PU-238	4	18	18	100.00	0.010	J	0.07	J	0.02	0.05	PCI/G (DRY)
PU-238	5	22	22	100.00	0.010	J	0.23	J	0.03	0.00	PCI/G (DRY)
PU-238	10	4	4	100.00	0.010	J	0.01	J	0.01	0.19	PCI/G (DRY)
PU-239	2	28	28	100.00	0.010	J	0.69	V/V	0.20	0.01	PCI/G (DRY)
PU-239	3	19	19	100.00	0.010	J	0.03	J	0.02	0.01	PCI/G (DRY)

TABLE C.2
SUMMARY STATISTICS FOR SEDIMENT CORE DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

ANALYSIS	TYPE OF ANALYSIS-RAD (CONTINUED)										
	REACH	NO. OF OBS.	NO. DETECTED	%	MIN		MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
					VALUE	QUAL.					
PU-239,240	4	18	18	100.00	0.010	J	1.57	J	0.26	0.38	PCI/G (DRY)
	5	22	22	100.00	0.010	J	0.45	J	0.10	0.13	PCI/G (DRY)
	10	4	4	100.00	0.010	J	0.04	J	0.03	0.02	PCI/G (DRY)
PU-239,240	2	28	28	100.00	0.010	J	1.25	J	0.38	0.37	PCI/G (DRY)
	3	19	19	100.00	0.020	J	0.16	J	0.08	0.04	PCI/G (DRY)
	4	18	18	100.00	0.070	J	1.09	J	0.40	0.31	PCI/G (DRY)
SR-90	5	23	23	100.00	-0.060	J	0.39	J	0.15	0.11	PCI/G (DRY)
	10	5	5	100.00	0.040	J	0.34	J	0.12	0.12	PCI/G (DRY)
	2	28	28	100.00	0.290	J	1.99	J	1.13	0.42	PCI/G (DRY)
U-234	3	17	17	100.00	2.850	J	8.91	J	5.82	1.28	PCI/G (DRY)
	4	18	18	100.00	1.050	J	5.47	J	2.21	0.98	PCI/G (DRY)
	5	23	23	100.00	0.890	J	3.08	J	1.96	0.49	PCI/G (DRY)
U-234	10	5	5	100.00	1.050	J	1.27	J	1.19	0.09	PCI/G (DRY)
	2	26	26	100.00	0.010	J	0.11	J	0.05	0.03	PCI/G (DRY)
	3	17	17	100.00	0.060	J	0.30	J	0.20	0.08	PCI/G (DRY)
U-235	4	18	18	100.00	0.010	J	0.69	J	0.12	0.16	PCI/G (DRY)
	5	23	23	100.00	0.010	J	0.37	J	0.08	0.08	PCI/G (DRY)
	10	5	5	100.00	0.010	J	0.15	J	0.06	0.05	PCI/G (DRY)
U-235	2	28	28	100.00	0.330	J	1.83	J	1.03	0.41	PCI/G (DRY)
	3	17	17	100.00	2.910	J	8.92	J	5.26	0.91	PCI/G (DRY)
	4	18	18	100.00	0.940	J	4.03	J	1.88	0.75	PCI/G (DRY)
U-238	5	23	23	100.00	0.590	J	2.45	J	1.66	0.46	PCI/G (DRY)
	10	5	5	100.00	1.020	J	1.79	J	1.32	0.29	PCI/G (DRY)

TABLE C.3
SUMMARY STATISTICS FOR SEDIMENT GRAB DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

ANALYSIS	NO. OF OBS.	NO. DETECTED	% DETECTED	TYPE OF ANALYSIS-RAD				AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
				MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.			
CS-137	487	482	98.97	0.005	V/V	34.13	V/V	1.23	2.31	PCI/G (1967)
CO-60	488	283	57.99	0.001	J	0.52	J	0.07	0.08	PCI/G (1967)

TABLE C.4
SUMMARY STATISTICS FOR SEDIMENT GRAB DATA
DETECTED COMPOUNDS SUMMARIZED BY REACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS & REACH

ANALYSIS	TYPE OF ANALYSIS=RAD											
	REACH	NO. OF OBS.	NO. DETECTED	%	DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
CO-80	1	4	4	100.00	0.023	V/V	0.24	J	0.10	0.10	0.10	PCI/G (DRY)
CO-80	2	46	44	95.85	0.018	J	0.30	J	0.11	0.11	0.08	PCI/G (DRY)
CO-80	3	3	3	100.00	0.013	V/V	0.14	V/V	0.07	0.07	0.06	PCI/G (DRY)
CO-80	4	95	93	97.89	0.023	V/V	0.52	J	0.15	0.15	0.11	PCI/G (DRY)
CO-80	5	277	116	41.88	0.007	UJ	0.27	UJ	0.04	0.04	0.04	PCI/G (DRY)
CO-80	6	30	15	50.00	0.001	J	0.26	J	0.05	0.05	0.07	PCI/G (DRY)
CO-80	7	9	7	77.78	0.027	UJ	0.12	V/V	0.06	0.06	0.03	PCI/G (DRY)
CO-80	10	1	0	0.00	0.020	UJ	0.02	UJ	0.02	0.02	0.01	PCI/G (DRY)
CO-80	18	23	1	4.35	0.005	UJ	0.04	J	0.02	0.02	0.14	PCI/G (DRY)
CS-137	1	4	4	100.00	0.133	V/V	0.40	V/V	0.26	0.26	5.11	PCI/G (DRY)
CS-137	2	46	46	100.00	0.256	V/V	34.13	V/V	2.92	2.92	1.08	PCI/G (DRY)
CS-137	3	3	3	100.00	0.346	V/V	2.23	V/V	1.00	1.00	2.52	PCI/G (DRY)
CS-137	4	93	83	100.00	0.214	V/V	9.49	V/V	2.72	2.72	1.00	PCI/G (DRY)
CS-137	5	277	274	98.92	0.016	V/V	10.32	V/V	0.62	0.62	1.58	PCI/G (DRY)
CS-137	6	31	31	100.00	0.005	V/V	6.88	V/V	1.02	1.02	0.14	PCI/G (DRY)
CS-137	7	9	8	88.89	0.029	UJ	0.47	V/V	0.30	0.30	0.14	PCI/G (DRY)
CS-137	10	1	1	100.00	0.241	V/V	0.24	V/V	0.24	0.24	0.13	PCI/G (DRY)
CS-137	18	23	22	95.65	0.026	V/V	0.51	V/V	0.15	0.15	0.13	PCI/G (DRY)

Appendix D

STATISTICAL PROCEDURE FOR PAIRWISE COMPARISON OF FISH TISSUE RESULTS

CONTENTS OF APPENDIX D

D.1	DATA DISTRIBUTION AND TRANSFORMATION	D-3
D.2	ESTIMATING MEANS FROM CENSORED DATA SETS	D-3
D.3	PAIR-WISE COMPARISON OF CONTAMINANT CONCENTRATIONS	D-4
D.4	REFERENCES FOR APPENDIX D	D-5

TABLES

Table D.1	Summary statistics for fish tissue data summarized for all reaches sorted by type of analysis and decreasing percentage detected	D-6
Table D.2	Summary statistics for fish tissue data detected compounds summarized by subreach sorted by type of analysis, reach, and subreach	D-9

D.1 DATA DISTRIBUTION AND TRANSFORMATION

Before appropriate analytical methods and statistical tests can be selected and performed on the contaminant concentration data, there should be some basis for assuming a particular form of data distribution (e.g., normal or lognormal). One possible choice for the distribution is the lognormal, because the observations are bounded on the lower side by zero and their distribution tends to be positively skewed with a tail to the right—both properties that are compatible with the lognormal distribution. Because tests for normality or lognormality (e.g., the Shapiro-Wilk or Kolmogorov tests) are not formulated to manage censored observations, we restricted the analysis to combinations of site, analyte, and species for which censored observations were not present. This restriction resulted in our analyzing only Aroclor 1260 for catfish and mercury for bluegill. The data distribution of Aroclor 1260 was thus assumed to be representative of the organic analytes and mercury of the inorganic analytes. That is, if the distribution for PCB or mercury was a member of a particular family of distributions (e.g., lognormal), then the other analytes within that group of analytes were assumed to be of the same family. In order to increase the sample size for evaluating the distribution types, sites were grouped together based on expectations of similar results due to their respective proximity to each other and contaminant point sources. Tests for normality and lognormality were performed on data from each group of sites. Except for one data set, the test for normality on the untransformed data was significant ($P < 0.01$), indicating a non-normal distribution. When the data were log transformed, the normality test statistic increased in all cases to the point at which the lognormal distribution seemed appropriate, except for the single case mentioned previously. Based on these results, the assumption of lognormality for the distribution of the organic and inorganic analytes seemed appropriate, and this distribution was assumed for all subsequent analyses.

D.2 ESTIMATING MEANS FROM CENSORED DATA SETS

Censored data sets are those which contain some observations reported only as greater or less than some value (less than an SQL in this case). Although various methods are available to obtain estimates of the mean and variance of censored data sets (such as replacing the censored observations with a value of 0, half the SQL, or the SQL itself, and then proceeding as if the observations were uncensored), many of these methods make inefficient use of the available data. When the distribution of the censored data set can be assumed to be of a particular form (e.g., normal, lognormal, etc.), the maximum likelihood (ML) method is known to be an efficient method of obtaining these estimates. ML estimates of the mean and variance of the analyte concentrations can be obtained with a method similar to that developed by Cohen (1963) for progressively censored samples. The results from an analysis of the PCB and mercury data distributions form the basis for assuming that the fish body burden data measured in Phase 1 are lognormally distributed. This assumption makes it possible to specify the distribution function needed in the likelihood function. The ML estimates of the unknown parameters needed to specify the lognormal distribution were obtained using the LIFEREG procedure from SAS (Version 6.06).

As the proportion of censoring increases and/or the sample size decreases, the confidence in the estimates of the distribution parameters decreases. This relationship is reflected in the general increase in the standard errors for the different parameter estimates as either of these occur. A plot of the relationship between the proportion of censoring and the relative

standard error (std. error/mean) from the Phase 1 data suggests that the reliability of the ML method begins to decrease for those data sets where the censoring exceeds 60 to 70%. The number of detectable observations in a sample is a simple measure which can also indicate a high degree of censoring or a small sample size (or both), either of which can produce less reliable ML estimates. Given this information, we considered the ML method inappropriate for calculating useful standard errors when the proportion of censoring exceeded 60% or when the number of detectable observations was less than three. When these criteria were exceeded, we deemed the estimation of a mean and variance by any method for the purpose of statistical analyses to be unsuitable.

D.3 PAIR-WISE COMPARISON OF CONTAMINANT CONCENTRATIONS

We adapted current statistical procedures to evaluate the difference in concentration between each affected site and the reference site on a pair-wise basis. A description of this method follows. Let $\hat{m}_i$, the estimates of the means of the two lognormal distributions, be

$$\hat{m}_i = \exp(\hat{\mu}_i + \frac{\hat{\sigma}_i^2}{2}), \quad (1)$$

where

$$i = 1, 2,$$

$\hat{\mu}_i$ = estimated mean for the transformed (log) data at site i ,

and

$\hat{\sigma}_i^2$ = estimated variance for the transformed log data at site i .

The two means of interest correspond to the affected and reference sites. Since differences in the $\hat{m}_i$'s would be reflected by differences in the terms $\hat{\mu}_i + \hat{\sigma}_i^2/2$, the comparison of the two distributions will be done in terms of these latter quantities. An asymptotic z-test was used to compare the two means with the test statistic calculated as

$$z = \frac{\ln(\hat{m}_1) - \ln(\hat{m}_2)}{\sqrt{(s_1)^2 + (s_2)^2}} \quad (2)$$

where

$$s_i = s.e.(\hat{\mu}_i + \frac{\hat{\sigma}_i^2}{2}) \quad (3)$$

The z value was then compared with a standard normal distribution. This test was adequate only when a reliable mean and s.e. could be calculated for both sites i and j. That is, either by standard procedures if there was no censoring, or by the maximum likelihood method for censored data if censoring did not exceed 60% and the number of detectable results was greater than 2.

If a reliable mean and s.e. could only be calculated for either site i or j (but not both), then a second method of site comparison was required. First, a 95% confidence interval on the mean of the untransformed data was calculated for the site with the reliable mean. If none of the individual values for the other site fell within the confidence interval, we concluded that there was a strong indication that the two sites were different. Otherwise we concluded that the data was insufficient to determine any significant difference between the two sites.

Lastly, if reliable means and s.e.'s were not attainable for both sites which were being compared, we considered the data to be insufficient to draw any conclusions about the similarity of the two sites.

D.4 REFERENCE FOR APPENDIX D

Cohen, A.C. 1963. Progressively censored samples in life testing. *Technometrics* 5:327-339.

TABLE D.1
SUMMARY STATISTICS FOR FISH TISSUE DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

TYPE OF ANALYSIS-INORGANICS											
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
MERCURY	FISH	169	169	100.00	0.019	J	0.88	J	0.21	0.21	MG/KG
ZINC	FISH	168	168	100.00	2.200	J	71.00	J	8.54	5.28	MG/KG
ARSENIC	FISH	169	117	69.23	0.050	UJ	0.24	J	0.08	0.04	MG/KG
SELENIUM	FISH	169	69	40.83	0.390	UJ	1.10	J	0.62	0.15	MG/KG
BERYLLIUM	FISH	154	19	12.34	0.003	UJ	0.04	J	0.00	0.00	MG/KG
COPPER	FISH	169	6	3.55	0.280	UJ	170.00	J	1.55	13.04	MG/KG
ANTIMONY	FISH	169	0	0.00	0.280	UJ	0.63	UJ	0.50	0.06	MG/KG
CADMIUM	FISH	169	0	0.00	0.110	UJ	0.28	UJ	0.20	0.03	MG/KG
CHROMIUM	FISH	169	0	0.00	0.280	UJ	0.63	UJ	0.51	0.07	MG/KG
LEAD	FISH	119	0	0.00	0.280	UJ	0.61	UJ	0.50	0.07	MG/KG
NICKEL	FISH	149	0	0.00	0.280	UJ	1.10	UJ	0.50	0.08	MG/KG
SILVER	FISH	169	0	0.00	0.110	UJ	0.25	UJ	0.20	0.03	MG/KG
THALLIUM	FISH	160	0	0.00	0.020	UJ	0.02	UJ	0.02	0.00	MG/KG
TYPE OF ANALYSIS-ORGANICS											
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS
AROCLOL-1260	FISH	119	117	98.32	0.010	NJ	3.60	NJ	0.55	0.60	MG/KG
ALPHA-CHLORDANE	FISH	109	102	93.58	0.001	NJ	0.13	J	0.01	0.02	MG/KG
GAMMA-CHLORDANE	FISH	110	94	85.45	0.000	NJ	0.07	J	0.01	0.01	MG/KG
AROCLOL-1254	FISH	106	88	83.02	0.010	U	0.77	NJ	0.11	0.11	MG/KG
4,4'-DDE	FISH	113	73	64.60	0.000	J	0.20	UX	0.03	0.05	MG/KG
OXYCHLORDANE	FISH	110	95	59.09	0.000	NJ	0.20	UX	0.02	0.04	MG/KG
GAMMA-CHLORDENE	FISH	113	61	53.98	0.000	NJ	0.20	J	0.02	0.04	MG/KG
ALPHA-CHLORDENE	FISH	113	34	30.09	0.000	NJ	0.20	UX	0.02	0.05	MG/KG
4,4'-DDD	FISH	114	31	27.19	0.000	J	0.20	UX	0.03	0.05	MG/KG
ALDRIN	FISH	111	8	7.21	0.000	J	0.20	UX	0.03	0.05	MG/KG
4,4'-DDT	FISH	114	8	7.02	0.000	J	0.20	UX	0.03	0.05	MG/KG
CHLORDENE	FISH	116	8	6.90	0.000	NJ	0.20	UX	0.03	0.05	MG/KG
BIS(2-ETHYLHEXYL)PHTHALATE	FISH	56	1	1.79	0.047	J	1.90	UX	0.57	0.34	MG/KG
DI-N-OCTYLPHTHALATE	FISH	56	1	1.79	0.024	J	1.90	UX	0.55	0.35	MG/KG
ACENAPHTHENE	FISH	57	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
ACENAPHTHYLENE	FISH	57	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
ANTHRACENE	FISH	57	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BENZO(A)ANTHRACENE	FISH	57	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BENZO(A)PYRENE	FISH	56	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BENZO(B)FLUORANTHENE	FISH	56	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BENZO(G,H,I)PERYLENE	FISH	55	0	0.00	0.180	UX	1.90	UX	0.58	0.34	MG/KG
BENZO(K)FLUORANTHENE	FISH	56	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BENZOIC ACID	FISH	50	0	0.00	0.930	UX	9.30	UX	2.75	1.76	MG/KG
BENZYL ALCOHOL	FISH	50	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BIS(2-CHLOROETHOXY)METHANE	FISH	57	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG
BIS(2-CHLOROETHYL)ETHER	FISH	57	0	0.00	0.180	UX	1.90	UX	0.57	0.33	MG/KG
BIS(2-CHLOROISOPROPYL)ETHE	FISH	58	0	0.00	0.180	UX	1.90	UX	0.57	0.34	MG/KG

TABLE D.1
SUMMARY STATISTICS FOR FISH TISSUE DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

TYPE OF ANALYSIS-ORGANICS (CONTINUED)												
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS	
BUTYLBENZYLPHthalATE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
CHRYSENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
DI-N-BUTYLPHthalATE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
DIBENZ(A,H)ANTHRACENE	FISH	56	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
DIBENZOFURAN	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
DIETHYLPHthalATE	FISH	57	0	0.00	0.080	UX	1.90	UX	0.55	0.33	MG/KG	
DIMETHYLPHthalATE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
FLUORANTHENE	FISH	58	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
FLUORENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
HEXACHLOROBENZENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
HEXACHLOROBUTADIENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
HEXACHLOROCYCLOPENTADIENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
HEXACHLOROTHANE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
INDENO(1,2,3-CD)PYRENE	FISH	56	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
ISOPHORONE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
N-NITROSO-DI-N-PROPYLAMINE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
N-NITROSDIPHENYLAMINE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
NAPHTHALENE	FISH	58	0	0.00	0.057	UJ	1.90	UX	0.56	0.34	MG/KG	
NITROBENZENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
PENTACHLOROPHENOL	FISH	76	0	0.00	0.930	UJ	9.30	UX	3.04	1.48	MG/KG	
PHENANTHRENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
PHENOL	FISH	76	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
PYRENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
1,2-DICHLOROBENZENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
1,2,4-TRICHLOROBENZENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
1,3-DICHLOROBENZENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
1,4-DICHLOROBENZENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
2-CHLORONAPHTHALENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
2-CHLOROPHENOL	FISH	77	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
2-METHYLNAPHTHALENE	FISH	80	0	0.00	0.035	UJ	1.90	UX	0.54	0.35	MG/KG	
2-METHYLPHENOL	FISH	76	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
2-NITROANILINE	FISH	56	0	0.00	0.930	UX	9.30	UX	2.83	1.67	MG/KG	
2-NITROPHENOL	FISH	76	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
2,4-DICHLOROPHENOL	FISH	77	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
2,4-DIMETHYLPHENOL	FISH	77	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
2,4-DINITROPHENOL	FISH	76	0	0.00	0.930	UJ	9.30	UX	3.04	1.48	MG/KG	
2,4-DINITROTOLUENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
2,4,5-TRICHLOROPHENOL	FISH	76	0	0.00	0.930	UJ	9.30	UX	3.04	1.48	MG/KG	
2,4,6-TRICHLOROPHENOL	FISH	76	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
2,6-DINITROTOLUENE	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
3-NITROANILINE	FISH	35	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
3,3'-DICHLOROBENZIDINE	FISH	57	0	0.00	0.370	UX	3.70	UX	1.14	0.66	MG/KG	
4-BROMOPHENYL-PHENYLETHYER	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	
4-CHLORO-3-METHYLPHENOL	FISH	78	0	0.00	0.190	UJ	1.90	UX	0.61	0.30	MG/KG	
4-CHLORANILINE	FISH	51	0	0.00	0.190	UX	1.90	UJ	0.50	0.25	MG/KG	
4-CHLOROPHENYL-PHENYLETHYER	FISH	57	0	0.00	0.190	UX	1.90	UX	0.57	0.34	MG/KG	

TABLE D.1
SUMMARY STATISTICS FOR FISH TISSUE DATA
SUMMARIZED FOR ALL REACHES
SORTED BY TYPE OF ANALYSIS & DECREASING %-DETECTED

TYPE OF ANALYSIS-ORGANICS (CONTINUED)												
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS	
4-METHYLPHENOL	FISH	77	0	0.00	0.190	UJ	1.90	UX	0.81	0.30	MG/KG	
4-NITROANILINE	FISH	50	0	0.00	0.930	UX	9.30	UX	2.75	1.76	MG/KG	
4-NITROPHENOL	FISH	76	0	0.00	0.930	UJ	9.30	UX	3.04	1.48	MG/KG	
4,6-DINITRO-2-METHYLPHENOL	FISH	75	0	0.00	0.930	UJ	9.30	UX	3.03	1.49	MG/KG	
TYPE OF ANALYSIS-RAD												
ANALYSIS	SAMPLE MATRIX	NO. OF OBS.	NO. DETECTED	% DETECTED	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	STD. DEV.	SAMPLE UNITS	
SR-90	FISH	87	87	100.00	-0.300	J	1.00	J	0.19	0.21	PCI/G (BONE)	
CS-137	FISH	214	127	59.35	0.005	V/V	2.31	V/V	0.14	0.22	PCI/G (WET)	
CO-60	FISH	211	4	1.90	0.005	UJ	1.21	UJ	0.14	0.15	PCI/G (WET)	

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH
----- TYPE OF ANALYSIS-INORGANICS -----

ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
ARSENIC	FISH	1.1	15	9	60.00	0.050	UJ	0.17	J	0.08	0.04	MG/KG
ARSENIC	FISH	1.5	13	8	61.54	0.050	UJ	0.18	J	0.08	0.04	MG/KG
ARSENIC	FISH	2.3	13	5	38.46	0.050	UJ	0.20	J	0.07	0.04	MG/KG
ARSENIC	FISH	3.1	9	5	55.56	0.050	UJ	0.07	J	0.06	0.01	MG/KG
ARSENIC	FISH	3.2	13	3	23.08	0.050	UJ	0.14	J	0.07	0.03	MG/KG
ARSENIC	FISH	3.3	13	11	84.62	0.050	UJ	0.12	J	0.08	0.03	MG/KG
ARSENIC	FISH	4.2	12	10	83.33	0.050	UJ	0.15	J	0.09	0.03	MG/KG
ARSENIC	FISH	4.5	15	15	100.00	0.050	J	0.14	J	0.10	0.02	MG/KG
ARSENIC	FISH	5.2	12	12	100.00	0.060	J	0.17	J	0.08	0.04	MG/KG
ARSENIC	FISH	5.3	13	11	84.62	0.050	UJ	0.13	J	0.08	0.03	MG/KG
ARSENIC	FISH	10.1	15	14	93.33	0.050	UJ	0.24	J	0.08	0.05	MG/KG
ARSENIC	FISH	13.1	10	4	40.00	0.050	UJ	0.08	J	0.05	0.01	MG/KG
ARSENIC	FISH	15.1	18	10	62.50	0.050	UJ	0.17	J	0.08	0.04	MG/KG
ARSENIC	FISH	1.1	15	3	20.00	0.003	UJ	0.01	J	0.00	0.00	MG/KG
BERYLLIUM	FISH	1.5	10	1	10.00	0.003	UJ	0.01	J	0.00	0.00	MG/KG
BERYLLIUM	FISH	2.3	13	0	0.00	0.003	UJ	0.00	UJ	0.00	0.00	MG/KG
BERYLLIUM	FISH	3.1	9	0	0.00	0.003	UJ	0.00	UJ	0.00	0.00	MG/KG
BERYLLIUM	FISH	3.2	13	4	30.77	0.003	UJ	0.01	J	0.00	0.00	MG/KG
BERYLLIUM	FISH	3.3	13	3	23.08	0.003	UJ	0.02	J	0.00	0.00	MG/KG
BERYLLIUM	FISH	4.2	12	0	0.00	0.003	UJ	0.00	UJ	0.00	0.00	MG/KG
BERYLLIUM	FISH	4.5	15	0	0.00	0.003	UJ	0.00	UJ	0.00	0.00	MG/KG
BERYLLIUM	FISH	5.2	12	0	0.00	0.003	UJ	0.00	UJ	0.00	0.00	MG/KG
BERYLLIUM	FISH	5.3	13	2	15.38	0.003	UJ	0.03	J	0.01	0.01	MG/KG
BERYLLIUM	FISH	10.1	10	1	10.00	0.003	UJ	0.04	J	0.01	0.00	MG/KG
BERYLLIUM	FISH	13.1	3	0	0.00	0.003	UJ	0.02	J	0.01	0.01	MG/KG
BERYLLIUM	FISH	15.1	18	5	31.25	0.003	UJ	0.55	UJ	0.44	0.09	MG/KG
COPPER	FISH	1.1	15	0	0.00	0.280	UJ	0.60	UJ	0.52	0.04	MG/KG
COPPER	FISH	1.5	13	0	0.00	0.460	UJ	0.59	UJ	0.51	0.08	MG/KG
COPPER	FISH	2.3	13	0	0.00	0.460	UJ	0.59	UJ	0.53	0.03	MG/KG
COPPER	FISH	3.1	9	0	0.00	0.300	UJ	0.58	UJ	0.51	0.04	MG/KG
COPPER	FISH	3.2	13	0	0.00	0.440	UJ	0.57	UJ	0.48	0.05	MG/KG
COPPER	FISH	3.3	13	0	0.00	0.480	UJ	0.61	UJ	0.55	0.05	MG/KG
COPPER	FISH	4.2	12	0	0.00	0.480	UJ	0.58	UJ	0.53	0.03	MG/KG
COPPER	FISH	4.5	15	0	0.00	0.480	UJ	0.68	J	0.55	0.06	MG/KG
COPPER	FISH	5.2	12	2	16.67	0.460	UJ	170.00	J	14.05	46.87	MG/KG
COPPER	FISH	5.3	13	4	30.77	0.460	UJ	0.61	UJ	0.48	0.07	MG/KG
COPPER	FISH	10.1	15	0	0.00	0.350	UJ	0.59	UJ	0.52	0.05	MG/KG
COPPER	FISH	13.1	10	0	0.00	0.450	UJ	0.57	UJ	0.48	0.05	MG/KG
COPPER	FISH	15.1	18	0	0.00	0.350	UJ	0.20	J	0.08	0.05	MG/KG
MERCURY	FISH	1.1	15	15	100.00	0.029	J	0.37	J	0.09	0.02	MG/KG
MERCURY	FISH	1.5	13	13	100.00	0.019	J	0.07	J	0.04	0.10	MG/KG
MERCURY	FISH	2.3	13	13	100.00	0.037	J	0.88	J	0.57	0.22	MG/KG
MERCURY	FISH	3.1	9	9	100.00	0.202	J	0.75	J	0.55	0.19	MG/KG
MERCURY	FISH	3.2	13	13	100.00	0.086	J	0.56	J	0.34	0.12	MG/KG
MERCURY	FISH	3.3	13	13	100.00	0.072	J	0.77	J	0.43	0.18	MG/KG
MERCURY	FISH	4.2	12	12	100.00	0.186	J	0.22	J	0.11	0.06	MG/KG
MERCURY	FISH	4.5	15	15	100.00	0.044	J					

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

TYPE OF ANALYSIS-INORGANICS (CONTINUED)											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV. UNITS
MERCURY	FISH	5.2	12	12	100.00	0.033	J	0.16	J	0.06	0.04 MG/KG
MERCURY	FISH	5.3	13	13	100.00	0.032	J	0.25	J	0.10	0.08 MG/KG
MERCURY	FISH	10.1	15	15	100.00	0.107	J	0.39	J	0.21	0.10 MG/KG
MERCURY	FISH	13.1	10	10	100.00	0.040	J	0.62	J	0.17	0.22 MG/KG
MERCURY	FISH	15.1	16	16	100.00	0.061	J	0.38	J	0.15	0.08 MG/KG
SELENIUM	FISH	1.1	15	14	93.33	0.500	J	1.00	J	0.89	0.15 MG/KG
SELENIUM	FISH	1.5	13	12	92.31	0.670	J	1.10	J	0.87	0.12 MG/KG
SELENIUM	FISH	2.3	13	11	84.62	0.540	UJ	1.10	J	0.75	0.15 MG/KG
SELENIUM	FISH	3.1	8	0	0.00	0.480	UJ	0.59	UJ	0.53	0.03 MG/KG
SELENIUM	FISH	3.2	13	1	7.69	0.390	UJ	1.10	U	0.59	0.17 MG/KG
SELENIUM	FISH	3.3	13	1	7.69	0.440	UJ	0.65	J	0.49	0.06 MG/KG
SELENIUM	FISH	4.2	12	11	91.67	0.540	UJ	0.97	J	0.77	0.11 MG/KG
SELENIUM	FISH	4.5	15	3	20.00	0.480	UJ	0.64	J	0.54	0.05 MG/KG
SELENIUM	FISH	5.2	12	0	0.00	0.480	UJ	0.83	UJ	0.53	0.04 MG/KG
SELENIUM	FISH	5.3	13	0	0.00	0.480	UJ	0.62	UJ	0.54	0.04 MG/KG
SELENIUM	FISH	10.1	15	8	53.33	0.480	UJ	0.72	J	0.56	0.08 MG/KG
SELENIUM	FISH	13.1	10	0	0.00	0.450	UJ	0.59	UJ	0.52	0.05 MG/KG
SELENIUM	FISH	15.1	16	8	50.00	0.450	UJ	0.80	J	0.81	0.12 MG/KG
ZINC	FISH	1.1	15	15	100.00	4.500	J	11.00	J	6.92	2.12 MG/KG
ZINC	FISH	1.5	13	13	100.00	4.900	J	11.00	J	6.69	1.63 MG/KG
ZINC	FISH	2.3	13	13	100.00	3.900	J	8.10	J	5.84	1.11 MG/KG
ZINC	FISH	3.1	8	8	100.00	2.200	J	7.10	J	5.22	1.34 MG/KG
ZINC	FISH	3.2	13	13	100.00	3.900	J	7.60	J	5.58	1.14 MG/KG
ZINC	FISH	3.3	13	13	100.00	4.100	J	8.60	J	6.82	1.31 MG/KG
ZINC	FISH	4.2	12	12	100.00	4.200	J	7.40	J	5.92	1.03 MG/KG
ZINC	FISH	4.5	15	15	100.00	4.000	J	9.50	J	5.81	1.78 MG/KG
ZINC	FISH	5.2	11	11	100.00	3.900	J	8.10	J	6.19	1.13 MG/KG
ZINC	FISH	5.3	13	13	100.00	3.100	J	71.00	J	12.05	18.01 MG/KG
ZINC	FISH	10.1	15	15	100.00	3.500	J	7.30	J	5.32	1.05 MG/KG
ZINC	FISH	13.1	10	10	100.00	5.200	J	7.60	J	6.36	0.82 MG/KG
ZINC	FISH	15.1	16	16	100.00	4.000	J	11.00	J	5.97	1.74 MG/KG

----- TYPE OF ANALYSIS-ORGANICS -----											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV. UNITS
ALDRIN	FISH	1.1	8	0	0.00	0.010	UX	0.10	UX	0.04	0.05 MG/KG
ALDRIN	FISH	1.5	8	0	0.00	0.010	UJ	0.20	UJ	0.13	0.08 MG/KG
ALDRIN	FISH	2.3	8	0	0.00	0.010	UX	0.10	UX	0.09	0.03 MG/KG
ALDRIN	FISH	2.5	6	2	33.33	0.000	J	0.01	UX	0.01	0.01 MG/KG
ALDRIN	FISH	3.1	7	0	0.00	0.008	UJ	0.01	UJ	0.01	0.00 MG/KG
ALDRIN	FISH	3.2	8	0	0.00	0.008	UX	0.01	UJ	0.01	0.00 MG/KG
ALDRIN	FISH	3.3	10	0	0.00	0.008	UX	0.01	UX	0.01	0.00 MG/KG
ALDRIN	FISH	3.4	9	0	0.00	0.008	UX	0.01	UX	0.01	0.00 MG/KG

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

-----	TYPE OF ANALYSIS-ORGANICS	-----
	(CONTINUED)	

ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	%	MIN VALUE	MAX VALUE	AVG. VALUE	STD. DEV.	SAMPLE UNITS
ALDRIN	FISH	4.2	4	2	50.00	0.001	0.01	0.01	0.01	MG/KG
	FISH	4.5	10	0	0.00	0.010	0.20	0.04	0.06	MG/KG
	FISH	5.2	9	0	0.00	0.010	0.10	0.02	0.03	MG/KG
	FISH	5.3	9	2	22.22	0.000	0.01	0.01	0.00	MG/KG
	FISH	10.1	8	0	0.00	0.010	0.01	0.00	0.00	MG/KG
	FISH	13.1	3	2	66.67	0.001	0.01	0.00	0.01	MG/KG
	FISH	15.1	4	0	0.00	0.050	0.20	0.14	0.08	MG/KG
	FISH	1.1	7	7	100.00	0.003	0.05	0.02	0.02	MG/KG
	FISH	1.5	8	3	37.50	0.003	0.01	0.01	0.00	MG/KG
	FISH	2.3	8	7	87.50	0.004	0.05	0.02	0.02	MG/KG
	FISH	2.5	2	2	100.00	0.002	0.00	0.00	0.00	MG/KG
	FISH	3.1	7	7	100.00	0.006	0.03	0.01	0.01	MG/KG
	FISH	3.2	8	8	100.00	0.007	0.03	0.02	0.01	MG/KG
	FISH	3.3	10	10	100.00	0.002	0.01	0.01	0.00	MG/KG
	FISH	3.4	9	9	100.00	0.003	0.01	0.01	0.00	MG/KG
	FISH	4.2	8	7	87.50	0.003	0.05	0.02	0.01	MG/KG
	FISH	4.5	10	10	100.00	0.001	0.06	0.02	0.02	MG/KG
	FISH	5.2	9	9	100.00	0.001	0.13	0.02	0.04	MG/KG
	FISH	5.3	9	9	100.00	0.003	0.03	0.01	0.01	MG/KG
	FISH	10.1	8	8	100.00	0.001	0.01	0.00	0.00	MG/KG
	FISH	13.1	2	2	100.00	0.005	0.01	0.01	0.00	MG/KG
	FISH	15.1	4	4	100.00	0.004	0.01	0.01	0.00	MG/KG
	FISH	1.1	7	3	42.86	0.001	0.10	0.02	0.04	MG/KG
	FISH	1.5	8	3	37.50	0.000	0.20	0.09	0.13	MG/KG
	FISH	2.3	8	2	25.00	0.002	0.10	0.06	0.05	MG/KG
	FISH	2.5	5	0	0.00	0.010	0.01	0.01	0.00	MG/KG
	FISH	3.1	7	1	14.29	0.010	0.02	0.01	0.01	MG/KG
	FISH	3.2	8	1	12.50	0.006	0.01	0.01	0.00	MG/KG
	FISH	3.3	10	0	0.00	0.002	0.01	0.01	0.00	MG/KG
	FISH	3.4	9	0	0.00	0.010	0.01	0.01	0.00	MG/KG
	FISH	4.2	8	2	25.00	0.001	0.01	0.01	0.00	MG/KG
	FISH	4.5	10	6	60.00	0.001	0.20	0.03	0.06	MG/KG
FISH	5.2	9	8	88.89	0.000	0.01	0.01	0.00	MG/KG	
FISH	5.3	8	0	0.00	0.010	0.01	0.01	0.00	MG/KG	
FISH	10.1	8	7	87.50	0.000	0.01	0.01	0.00	MG/KG	
FISH	13.1	4	0	0.00	0.010	0.01	0.01	0.00	MG/KG	
FISH	15.1	4	1	25.00	0.002	0.20	0.09	0.08	MG/KG	
AROCLOR-1254	FISH	1.1	8	5	62.50	0.010	0.14	0.07	0.13	MG/KG
AROCLOR-1254	FISH	1.5	8	7	87.50	0.010	0.77	0.22	0.24	MG/KG
AROCLOR-1254	FISH	2.3	8	2	100.00	0.090	0.19	0.14	0.07	MG/KG
AROCLOR-1254	FISH	2.5	2	2	100.00	0.020	0.14	0.08	0.04	MG/KG
AROCLOR-1254	FISH	3.1	7	7	100.00	0.060	0.33	0.22	0.10	MG/KG
AROCLOR-1254	FISH	3.2	8	8	100.00	0.060	0.37	0.13	0.11	MG/KG
AROCLOR-1254	FISH	3.3	7	7	100.00	0.060	0.19	0.09	0.07	MG/KG
AROCLOR-1254	FISH	3.4	9	9	100.00	0.020	0.19	0.08	0.10	MG/KG
AROCLOR-1254	FISH	4.2	8	5	62.50	0.010	0.27	0.09	0.10	MG/KG

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

TYPE OF ANALYSIS-ORGANICS (CONTINUED)											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
AROCLOR-1254	FISH	4.5	10	10	100.00	0.030 NJ	0.24 NJ		0.12	0.09	MG/KG
	FISH	5.2	9	8	88.87	0.010 U	0.18 NJ		0.07	0.06	MG/KG
	FISH	5.3	8	8	100.00	0.010 J	0.32 J		0.15	0.11	MG/KG
	FISH	10.1	8	0	0.00	0.010 U	0.05 U		0.02	0.02	MG/KG
	FISH	13.1	2	2	100.00	0.040 J	0.19 J		0.11	0.11	MG/KG
	FISH	15.1	4	4	100.00	0.080 NJ	0.20 NJ		0.13	0.05	MG/KG
	FISH	1.1	8	8	100.00	0.050 NJ	2.55 NJ		0.74	0.95	MG/KG
	FISH	1.5	8	8	100.00	0.100 NJ	0.41 NJ		0.21	0.10	MG/KG
	FISH	2.3	8	8	100.00	0.320 NJ	2.29 NJ		0.99	0.70	MG/KG
	FISH	2.5	8	8	100.00	0.230 J	1.42 J		0.68	0.51	MG/KG
	FISH	3.1	8	8	100.00	0.030 NJ	0.93 NJ		0.33	0.28	MG/KG
	FISH	3.2	8	8	100.00	0.280 NJ	3.00 NJ		0.94	0.87	MG/KG
	FISH	3.3	10	10	100.00	0.150 NJ	1.54 NJ		0.58	0.42	MG/KG
	FISH	3.4	10	10	100.00	0.210 NJ	0.55 NJ		0.38	0.12	MG/KG
	FISH	4.2	8	7	87.50	0.010 UX	2.10 NJ		0.70	0.71	MG/KG
	FISH	4.5	10	10	100.00	0.140 NJ	3.60 NJ		0.88	1.03	MG/KG
	FISH	5.2	9	8	100.00	0.030 NJ	0.45 NJ		0.26	0.15	MG/KG
	FISH	5.3	10	10	100.00	0.190 J	1.24 J		0.67	0.34	MG/KG
	FISH	10.1	8	7	87.50	0.010 NJ	0.08 NJ		0.03	0.03	MG/KG
	FISH	13.1	4	4	100.00	0.190 J	0.39 J		0.30	0.08	MG/KG
FISH	15.1	4	4	100.00	0.100 NJ	0.65 NJ		0.39	0.25	MG/KG	
BIS(2-ETHYLHEXYL)PHTHALATE	FISH	1.5	8	0	0.00	0.680 UX	0.87 UX		1.18	0.45	MG/KG
	FISH	2.3	8	0	0.00	0.700 UX	1.90 UX		0.37	0.01	MG/KG
	FISH	3.1	7	0	0.00	0.350 UX	0.39 UX		0.37	0.01	MG/KG
	FISH	3.2	8	0	0.00	0.380 UX	0.39 UX		0.37	0.01	MG/KG
	FISH	3.3	9	0	0.00	0.380 UX	0.41 UX		0.38	0.02	MG/KG
	FISH	3.4	9	1	11.11	0.047 J	0.41 UX		0.29	0.13	MG/KG
	FISH	4.2	1	0	0.00	0.680 UX	0.66 UX		0.66	0.02	MG/KG
	FISH	4.5	5	0	0.00	0.670 UX	0.72 UX		0.69	0.02	MG/KG
	FISH	15.1	3	0	0.00	0.670 UX	0.71 UX		0.69	0.05	MG/KG
	FISH	1.1	8	0	0.00	0.010 UX	0.10 UX		0.13	0.08	MG/KG
	FISH	1.5	8	0	0.00	0.010 UX	0.10 UX		0.09	0.03	MG/KG
	FISH	2.3	8	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG
	FISH	2.5	6	0	0.00	0.010 UX	0.01 J		0.01	0.00	MG/KG
	FISH	3.1	7	2	28.57	0.010 UX	0.01 UX		0.01	0.00	MG/KG
	FISH	3.2	8	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG
	FISH	3.3	10	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG
	FISH	3.4	9	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG
	FISH	4.2	8	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG
	FISH	4.5	10	0	0.00	0.010 UX	0.20 UX		0.04	0.06	MG/KG
	FISH	5.2	9	5	55.56	0.000 NJ	0.10 UX		0.01	0.03	MG/KG
FISH	5.3	9	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG	
CHLORDENE	FISH	10.1	8	1	12.50	0.002 NJ	0.01 UX		0.01	0.00	MG/KG
CHLORDENE	FISH	13.1	4	0	0.00	0.010 UX	0.01 UX		0.01	0.00	MG/KG
CHLORDENE	FISH	15.1	4	0	0.00	0.050 UX	0.20 UX		0.14	0.08	MG/KG
CHLORDENE	FISH	1.5	8	0	0.00	0.680 UX	0.87 UX		0.75	0.08	MG/KG

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	%	TYPE OF ANALYSIS-ORGANICS		MIN VALUE	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
						DETECTED	VALUE						
DI-N-OCTYLPHTHALATE	FISH	2.3	8	0	0.00	0.00	0.082	UX	1.90	UX	1.10	0.58	MG/KG
DI-N-OCTYLPHTHALATE	FISH	3.1	7	0	0.00	0.00	0.350	UX	0.39	UX	0.37	0.01	MG/KG
DI-N-OCTYLPHTHALATE	FISH	3.2	8	1	12.50	0.024	J	0.024	0.38	UJ	0.33	0.12	MG/KG
DI-N-OCTYLPHTHALATE	FISH	3.3	9	0	0.00	0.360	UJ	0.41	0.41	UJ	0.38	0.02	MG/KG
DI-N-OCTYLPHTHALATE	FISH	3.4	9	0	0.00	0.190	UX	0.190	0.41	UJ	0.32	0.10	MG/KG
DI-N-OCTYLPHTHALATE	FISH	4.2	1	0	0.00	0.860	UX	0.860	0.66	UX	0.66		MG/KG
DI-N-OCTYLPHTHALATE	FISH	4.5	5	0	0.00	0.670	UX	0.72	0.71	UX	0.69	0.02	MG/KG
DI-N-OCTYLPHTHALATE	FISH	15.1	3	0	0.00	0.670	UJ	0.670	0.71	UX	0.69	0.02	MG/KG
DI-N-OCTYLPHTHALATE	FISH	1.1	7	3	42.86	0.002	U	0.04	0.04	U	0.01	0.01	MG/KG
GAMMA-CHLORDANE	FISH	1.5	8	0	0.00	0.003	U	0.01	0.01	U	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	2.3	8	6	75.00	0.002	U	0.02	0.02	J	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	2.5	3	2	66.67	0.001	J	0.01	0.01	UX	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	3.1	7	7	100.00	0.016	J	0.05	0.05	J	0.03	0.01	MG/KG
GAMMA-CHLORDANE	FISH	3.2	8	8	100.00	0.018	J	0.02	0.07	J	0.04	0.02	MG/KG
GAMMA-CHLORDANE	FISH	3.3	10	10	100.00	0.008	J	0.02	0.02	J	0.02	0.01	MG/KG
GAMMA-CHLORDANE	FISH	3.4	9	8	100.00	0.001	J	0.03	0.03	J	0.02	0.01	MG/KG
GAMMA-CHLORDANE	FISH	4.2	8	8	100.00	0.002	J	0.02	0.02	J	0.01	0.01	MG/KG
GAMMA-CHLORDANE	FISH	4.5	10	10	100.00	0.001	J	0.03	0.03	J	0.01	0.01	MG/KG
GAMMA-CHLORDANE	FISH	5.2	9	9	100.00	0.000	NJ	0.00	0.01	J	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	5.3	9	8	88.89	0.002	J	0.06	0.06	J	0.02	0.02	MG/KG
GAMMA-CHLORDANE	FISH	10.1	8	8	100.00	0.000	J	0.00	0.00	J	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	13.1	2	2	100.00	0.003	J	0.01	0.01	J	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	15.1	4	4	100.00	0.002	J	0.01	0.01	J	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	1.1	7	6	85.71	0.000	J	0.01	0.01	UX	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	1.5	8	2	25.00	0.001	J	0.20	0.20	J	0.13	0.08	MG/KG
GAMMA-CHLORDANE	FISH	2.3	8	2	25.00	0.008	NJ	0.10	0.10	UX	0.08	0.04	MG/KG
GAMMA-CHLORDANE	FISH	2.5	6	0	0.00	0.010	UX	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	3.1	7	6	85.71	0.003	J	0.03	0.03	J	0.02	0.01	MG/KG
GAMMA-CHLORDANE	FISH	3.2	8	8	100.00	0.007	X	0.04	0.04	J	0.02	0.01	MG/KG
GAMMA-CHLORDANE	FISH	3.3	10	3	30.00	0.003	J	0.01	0.01	UJ	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	3.4	9	7	77.78	0.003	J	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	4.2	8	2	25.00	0.003	J	0.01	0.01	UJ	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	4.5	9	7	77.78	0.002	J	0.02	0.02	J	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	5.2	9	8	88.89	0.000	NJ	0.01	0.01	UX	0.00	0.00	MG/KG
GAMMA-CHLORDANE	FISH	5.3	9	1	11.11	0.000	J	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	10.1	8	7	87.50	0.000	NJ	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	13.1	3	0	0.00	0.010	UJ	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	15.1	4	2	50.00	0.003	J	0.10	0.10	UX	0.04	0.05	MG/KG
GAMMA-CHLORDANE	FISH	1.1	7	7	100.00	0.001	J	0.03	0.03	J	0.01	0.01	MG/KG
GAMMA-CHLORDANE	FISH	1.5	8	2	25.00	0.000	U	0.20	0.20	UJ	0.05	0.09	MG/KG
GAMMA-CHLORDANE	FISH	2.3	8	6	75.00	0.002	J	0.10	0.10	UX	0.03	0.04	MG/KG
GAMMA-CHLORDANE	FISH	2.5	3	0	0.00	0.010	UX	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	3.1	7	5	71.43	0.003	J	0.01	0.01	J	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	3.2	8	8	100.00	0.001	J	0.07	0.07	J	0.01	0.02	MG/KG
GAMMA-CHLORDANE	FISH	3.3	10	4	40.00	0.001	J	0.01	0.01	UX	0.01	0.00	MG/KG
GAMMA-CHLORDANE	FISH	3.4	9	7	77.78	0.001	J	0.01	0.01	UX	0.00	0.00	MG/KG

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

TYPE OF ANALYSIS-ORGANICS (CONTINUED)											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV.	SAMPLE UNITS
OXYCHLORDANE	FISH	4.2	8	7	87.50	0.008 J	0.09 J	J	0.03	0.03	MG/KG
	FISH	4.5	10	3	30.00	0.000 J	0.01 J	J	0.01	0.00	MG/KG
	FISH	5.2	9	3	33.33	0.000 NJ	0.10 UX	UX	0.02	0.03	MG/KG
	FISH	5.3	9	3	33.33	0.001 J	0.01 UX	UX	0.01	0.00	MG/KG
	FISH	10.1	8	7	87.50	0.000 NJ	0.01 UX	UX	0.00	0.00	MG/KG
	FISH	13.1	2	2	100.00	0.051 J	0.06 J	J	0.06	0.01	MG/KG
	FISH	15.1	4	1	25.00	0.004 J	0.20 UX	UX	0.11	0.10	MG/KG
	FISH	1.1	7	5	71.43	0.000 J	0.10 UX	UX	0.02	0.04	MG/KG
	FISH	1.5	8	0	0.00	0.003 U	0.20 UJ	UJ	0.12	0.10	MG/KG
	FISH	2.3	8	0	0.00	0.010 UX	0.10 UX	UX	0.08	0.03	MG/KG
	FISH	2.5	8	0	0.00	0.010 UX	0.01 UX	UX	0.01	0.00	MG/KG
	FISH	3.1	7	0	0.00	0.015 UJ	0.02 UJ	UJ	0.02	0.00	MG/KG
	FISH	3.2	8	2	25.00	0.000 J	0.02 UJ	UJ	0.01	0.01	MG/KG
	FISH	3.3	10	0	0.00	0.016 UX	0.02 UX	UX	0.02	0.00	MG/KG
	FISH	3.4	9	0	0.00	0.016 UX	0.01 UJ	UJ	0.01	0.00	MG/KG
	FISH	4.2	8	0	0.00	0.010 UJ	0.01 UJ	UJ	0.01	0.06	MG/KG
	FISH	4.5	10	4	40.00	0.003 J	0.20 UX	UX	0.03	0.06	MG/KG
	FISH	5.2	9	9	100.00	0.000 NJ	0.01 J	J	0.01	0.00	MG/KG
	FISH	5.3	8	3	37.50	0.002 J	0.02 J	J	0.01	0.01	MG/KG
	FISH	10.1	8	7	87.50	0.000 NJ	0.01 UX	UX	0.01	0.00	MG/KG
	FISH	13.1	4	0	0.00	0.010 UJ	0.01 UX	UX	0.13	0.09	MG/KG
	FISH	15.1	4	1	25.00	0.005 J	0.20 UX	UX	0.01	0.01	MG/KG
	FISH	1.1	8	6	75.00	0.001 J	0.03 J	J	0.01	0.01	MG/KG
	FISH	1.5	8	2	25.00	0.010 UJ	0.20 UJ	UJ	0.13	0.08	MG/KG
	FISH	2.3	8	4	50.00	0.005 J	0.10 UX	UX	0.05	0.05	MG/KG
	FISH	2.5	5	2	40.00	0.002 J	0.01 UX	UX	0.01	0.00	MG/KG
	FISH	3.1	7	7	100.00	0.005 J	0.04 J	J	0.01	0.01	MG/KG
	FISH	3.2	8	8	100.00	0.000 J	0.02 J	J	0.01	0.01	MG/KG
	FISH	3.3	10	10	100.00	0.000 J	0.00 J	J	0.00	0.00	MG/KG
	FISH	3.4	9	9	100.00	0.000 J	0.00 J	J	0.00	0.00	MG/KG
	FISH	4.2	8	6	75.00	0.004 J	0.02 J	J	0.01	0.01	MG/KG
	FISH	4.5	10	6	60.00	0.010 UX	0.20 UX	UX	0.04	0.06	MG/KG
	FISH	5.2	9	3	33.33	0.010 J	0.02 J	J	0.01	0.01	MG/KG
	FISH	5.3	9	6	66.67	0.005 J	0.03 J	J	0.01	0.01	MG/KG
	FISH	10.1	8	2	25.00	0.002 J	0.01 UX	UX	0.01	0.00	MG/KG
	FISH	13.1	2	2	100.00	0.002 J	0.01 J	J	0.00	0.00	MG/KG
	FISH	15.1	4	0	0.00	0.050 UX	0.20 UX	UX	0.14	0.08	MG/KG
	FISH	1.1	8	0	0.00	0.010 UX	0.10 UX	UX	0.04	0.05	MG/KG
	FISH	1.5	8	0	0.00	0.010 UJ	0.20 UJ	UJ	0.13	0.08	MG/KG
	FISH	2.3	8	0	0.00	0.010 UX	0.10 UX	UX	0.09	0.03	MG/KG
FISH	2.5	5	2	40.00	0.000 J	0.01 UX	UX	0.01	0.01	MG/KG	
FISH	3.1	7	0	0.00	0.015 UJ	0.02 UJ	UJ	0.02	0.00	MG/KG	
FISH	3.2	8	0	0.00	0.016 UX	0.02 UX	UX	0.02	0.00	MG/KG	
FISH	3.3	10	0	0.00	0.016 UX	0.02 UX	UX	0.02	0.00	MG/KG	
FISH	3.4	9	0	0.00	0.010 UX	0.02 UX	UX	0.02	0.00	MG/KG	
FISH	4.2	8	1	12.50	0.002 J	0.01 UJ	UJ	0.01	0.00	MG/KG	

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

TYPE OF ANALYSIS=ORGANICS (CONTINUED)											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. SAMPLE DEV. UNITS
4,4'-DDT	FISH	4.5	10	0	0.00	0.010	UX	0.20	UX	0.04	0.06 MG/KG
4,4'-DDT	FISH	5.2	9	0	0.00	0.010	UX	0.01	UX	0.01	0.00 MG/KG
4,4'-DDT	FISH	5.3	9	5	55.56	0.000	J	0.01	UX	0.00	0.00 MG/KG
4,4'-DDT	FISH	10.1	8	0	0.00	0.010	UX	0.01	UX	0.01	0.00 MG/KG
4,4'-DDT	FISH	13.1	3	0	0.00	0.010	UX	0.01	UX	0.01	0.00 MG/KG
4,4'-DDT	FISH	15.1	4	0	0.00	0.050	UX	0.20	UX	0.14	0.06 MG/KG
TYPE OF ANALYSIS=RAD											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. SAMPLE DEV. UNITS
CO-80	FISH	1.1	15	1	6.67	0.031	UJ	0.16	UJ	0.09	0.05 PCI/G (WET)
CO-80	FISH	1.5	13	0	0.00	0.029	UJ	0.19	UJ	0.10	0.05 PCI/G (WET)
CO-80	FISH	2.3	21	0	0.00	0.042	U	0.70	UJ	0.31	0.23 PCI/G (WET)
CO-80	FISH	2.5	6	0	0.00	0.076	UJ	0.43	UJ	0.25	0.17 PCI/G (WET)
CO-80	FISH	3.1	12	0	0.00	0.055	U	0.35	UJ	0.13	0.08 PCI/G (WET)
CO-80	FISH	3.2	14	0	0.00	0.032	UJ	0.18	UJ	0.09	0.04 PCI/G (WET)
CO-80	FISH	3.3	18	1	5.56	0.092	UJ	0.18	UJ	0.13	0.03 PCI/G (WET)
CO-80	FISH	3.4	10	0	0.00	0.068	UJ	0.38	UJ	0.18	0.10 PCI/G (WET)
CO-80	FISH	4.2	19	0	0.00	0.031	U	1.21	UJ	0.28	0.31 PCI/G (WET)
CO-80	FISH	4.5	19	0	0.00	0.006	UJ	0.24	UJ	0.13	0.05 PCI/G (WET)
CO-80	FISH	5.2	13	0	0.00	0.040	U	0.24	UJ	0.12	0.05 PCI/G (WET)
CO-80	FISH	5.3	15	0	0.00	0.029	UJ	0.19	UJ	0.10	0.05 PCI/G (WET)
CO-80	FISH	10.1	16	1	6.25	0.005	UJ	0.14	UJ	0.05	0.05 PCI/G (WET)
CO-80	FISH	13.1	8	1	18.67	0.051	UJ	0.17	UJ	0.08	0.04 PCI/G (WET)
CO-80	FISH	15.1	14	0	0.00	0.009	UJ	0.15	UJ	0.07	0.06 PCI/G (WET)
CS-137	FISH	1.1	15	7	46.67	0.010	J	0.12	UJ	0.06	0.04 PCI/G (WET)
CS-137	FISH	1.5	13	7	53.85	0.009	V/V	0.12	V/V	0.06	0.04 PCI/G (WET)
CS-137	FISH	2.3	21	18	85.71	0.095	J	2.31	V/V	0.49	0.52 PCI/G (WET)
CS-137	FISH	2.5	6	5	83.33	0.072	J	0.32	UJ	0.19	0.10 PCI/G (WET)
CS-137	FISH	3.1	12	5	41.67	0.036	V/V	0.29	UJ	0.10	0.07 PCI/G (WET)
CS-137	FISH	3.2	14	8	57.14	0.021	V/V	0.14	UJ	0.09	0.03 PCI/G (WET)
CS-137	FISH	3.3	18	10	55.56	0.005	V/V	0.17	V/V	0.08	0.04 PCI/G (WET)
CS-137	FISH	3.4	10	3	30.00	0.055	V/V	0.34	UJ	0.17	0.09 PCI/G (WET)
CS-137	FISH	4.2	20	14	70.00	0.035	V/V	1.03	UJ	0.20	0.25 PCI/G (WET)
CS-137	FISH	4.5	20	14	70.00	0.043	J	0.25	UJ	0.13	0.08 PCI/G (WET)
CS-137	FISH	5.2	13	10	76.92	0.020	V/V	0.16	UJ	0.09	0.05 PCI/G (WET)
CS-137	FISH	5.3	15	9	60.00	0.014	J	0.14	UJ	0.07	0.05 PCI/G (WET)
CS-137	FISH	10.1	17	9	52.94	0.013	J	0.17	UJ	0.07	0.06 PCI/G (WET)
CS-137	FISH	13.1	6	3	50.00	0.034	J	0.15	UJ	0.08	0.04 PCI/G (WET)
CS-137	FISH	15.1	14	5	35.71	0.016	J	0.16	UJ	0.08	0.05 PCI/G (WET)
SR-90	FISH	1.1	8	6	100.00	-0.100	J	0.10	J	0.01	0.10 PCI/G (BONE)
SR-90	FISH	2.5	6	6	100.00	0.200	V/V	0.70	V/V	0.45	0.19 PCI/G (BONE)
SR-90	FISH	3.1	8	8	100.00	-0.300	J	0.20	V/V	0.06	0.17 PCI/G (BONE)

TABLE D.2
SUMMARY STATISTICS FOR FISH TISSUE DATA
DETECTED COMPOUNDS SUMMARIZED BY SUBREACH
SORTED BY TYPE OF ANALYSIS, ANALYSIS, REACH & SUBREACH

TYPE OF ANALYSIS-RAD (CONTINUED)											
ANALYSIS	SAMPLE MATRIX	REACH SEGMENT	NO. OF OBS.	NO. DETECTED	% DETECTED	MIN VALUE	MIN QUAL.	MAX VALUE	MAX QUAL.	AVG. VALUE	STD. DEV. SAMPLE UNITS
SR-90	FISH	3.3	10	10	100.00	0.000	J	0.80	J	0.39	0.23 PC1/G (BONE)
SR-90	FISH	3.4	10	10	100.00	0.000	J	0.50	J	0.16	0.18 PC1/G (BONE)
SR-90	FISH	4.5	10	10	100.00	0.000	J	0.40	J	0.15	0.15 PC1/G (BONE)
SR-90	FISH	5.2	8	8	100.00	-0.100	J	1.00	J	0.24	0.30 PC1/G (BONE)
SR-90	FISH	5.3	10	10	100.00	0.000	J	0.80	J	0.17	0.17 PC1/G (BONE)
SR-90	FISH	10.1	8	8	100.00	-0.100	J	0.30	J	0.11	0.12 PC1/G (BONE)
SR-90	FISH	13.1	4	4	100.00	0.000	J	0.30	J	0.15	0.13 PC1/G (BONE)
SR-90	FISH	15.1	4	4	100.00	0.100	J	0.20	J	0.15	0.08 PC1/G (BONE)

Appendix E

HUMAN HEALTH RISK METHODS

CONTENTS OF APPENDIX E

E.1 HUMAN HEALTH SCREENING RISK ASSESSMENTS	E-3
E.2 FISH	E-3
E.3 SEDIMENT	E-4
E.3.1 Surface Sediments	E-4
E.3.2 Cores	E-4
E.4 WATER	E-4

TABLES AND FIGURES

Table E.1	Receiving reaches for Oak Ridge Reservation source monitoring and surface water environmental monitoring stations	E-6
Table E.2	Usage and exposure factors for conservative and nonconservative screening	E-8
Table E.3	Reference dose factors and slope (cancer potency) factors for inorganic compounds	E-10
Table E.4	Reference dose factors and slope (cancer potency) factors for organic compounds	E-11
Table E.5	Slope (cancer potency) factors for radionuclides	E-15
Table E.6	Summary of carcinogens in the secondary pathways identified as low priority for further investigation ...	E-16
Table E.7	Summary of noncarcinogens in the secondary pathways identified as low priority for further investigation	E-19
Table E.8	Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-22
Table E.9	Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-36
Table E.10	Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-45
Table E.11	Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-60
Table E.12	Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits	E-69
Table E.13	Results of adult conservative screening for carcinogens in dredged sediments where at least one measurement in a reach was above detection limits	E-78
Table E.14	Results of adult conservative screening for noncarcinogens in dredged sediments where at least one measurement in a reach was above detection limits	E-83
Table E.15	Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits	E-87
Table E.16	Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits	E-99
Table E.17	Results of adult nonconservative screening for carcinogens in dredged sediments where no measurements in a reach were above detection limits	E-113
Table E.18	Results of adult nonconservative screening for noncarcinogens in dredged sediments where no measurements in a reach were above detection limits	E-120
Table E.19	Unclassified analytes	E-128
Figure E.1	Location of NPDES monitoring sites in Clinch River/Watts Bar System	E-5

E.1 HUMAN HEALTH SCREENING RISK ASSESSMENTS

Conservative and nonconservative screening risk analyses were performed for selected media and primary and secondary exposure pathways. Primary pathways are those most realistic and likely to be encountered. Secondary pathways are less likely scenarios or potential and/or future pathways of exposure. A listing of risk assessment scenarios, usage, and exposure factors used for the human health screening is given in Table E.2. Tables E.3 through E.5 give toxicity reference doses and cancer factors for those contaminants considered in the risk analyses. Screening results are presented in Tables E.6 through E.18.

Media and Primary Exposure Pathways

1. Fish:
 - ingestion by adult and child.
2. Surface Sediment:
 - external exposure to sediments for adult,
 - ingestion of sediments for adult and child, and
 - inhalation of sediments for adult.
3. Water:
 - ingestion by adult and child.

Media and Secondary Exposure Pathway

Buried Sediment (dredging of deep sediments):

- external exposure to dredged sediments for adult;
- ingestion of dredged sediments for adult;
- inhalation of dredged sediments for adult; and
- ingestion of milk, meat, and vegetables produced on dredged sediments for adult.

The results of the analyses for the selected analytes (see Table A.1) provided a comprehensive view of the extent of contamination in the selected media. Actually, only 10% of the determinations exceeded their method detection limits. This is a common occurrence for regulatory-driven environmental sampling and analysis activities. For the screening risk analyses, a distinction was made between *detected contaminants*, contaminants with at least one detected value in a given reach, and *nondetected contaminants*, contaminants with no detected values in a reach. Separate analyses and interpretations of these results were done for detected and the nondetected contaminants.

The following sections describe the specific data manipulations performed for each of the selected media.

E.2 FISH

Except for PCBs data were combined for all fish species within a reach. Only catfish results were used for PCBs because catfish values are typically much greater than bass and are useful for the conservative screening. Only detected chlordane isomer values were summed. The average detection limit was used when all chlordane isomers were nondetects.

Given the potential for means of distributions with numerous non-detect values (censored data) to be positively biased, the upper 95% confidence limit of the arithmetic mean and the geometric mean were calculated using a maximum likelihood estimator (SAS PROC LIFEREG). The UCL was compared with the actual maximum value, and whichever is smaller was used in the conservative risk calculation as detected contaminants. The geometric mean was used for nonconservative calculations. Only a nonconservative calculation using the minimum detection limit was done for nondetected contaminants.

E3 SEDIMENT

E3.1 Surface Sediments

All grabs are summarized for each reach.

Surface sediment samples in a reach were used on the screening analysis. The upper 95% confidence limit of the arithmetic mean and the geometric mean were calculated using a maximum likelihood estimator (SAS PROC LIFEREG). The UCL was compared with the actual maximum value and which ever is smaller was used in the conservative risk calculations: for detected contaminants the geometric mean will be used for nonconservative calculations.

Only a nonconservative calculation using the minimum detection limit was done for the nondetected contaminants.

E3.2 Cores

A mass-weighted average concentration was calculated to a depth of 1 m. For cores shorter than 1 m, the mass-weighted average of what was collected was calculated and was assumed to represent the concentration in the top 1 m of sediment. When there was more than one core per reach, the mass-weighted average for each core was calculated and the upper 95% confidence limit of the average of the core averages, or the maximum core average, (whichever is smaller) was used as the reach value for conservative screening. The arithmetic mean of the core averages was used for nonconservative screening. Only a nonconservative calculation using the minimum detection limit of the core averages was done for the nondetected contaminants.

E4 WATER

The description of the water data used in the screening analyses requires some additional explanation. CR-ERP sampling consisted of a single grab sample collected at the same times (late 1989 and early 1990) and sites as the initial sediment and fish samples. To better represent the water quality of the CR/WBR system, additional data were obtained from ORR compliance and environmental monitoring programs. Table E.1 lists the sources of these data for each reach. The sampling locations relative to DOE facilities are shown in Fig. E.1. The source monitoring data are from samples collected at permitted outfalls to determine regulatory compliance with NPDES and DOE Order 5400.1. These samples were collected and analyzed at varying frequencies for the analytes specified in each outfall permit. Environmental monitoring samples were generally collected quarterly from locations upstream and downstream from DOE facilities. The quality of the analytical data is good, but not as well documented as the CR-ERP. Standard EPA sampling and analytical

Fig. E.1. Location of NPDES monitoring sites in Clinch River/Watts Bar System.

Table E.1. Receiving reaches for Oak Ridge Reservation source monitoring and surface water environmental monitoring stations. Source monitoring data are collected in support of NPDES permits and DOE Order 5400.1. (Data from these locations for years 1990, 1991, and part of 1992, were used in the human health and ecological risk screening analyses).

Reach	Location	Source Type
1	Kerr Hollow Quarry	Source Monitoring
1	Rogers Quarry	Source Monitoring
1	Melton Hill Dam	Environmental Monitoring
2	White Oak Dam	Source Monitoring
2	K-1513	Environmental Monitoring
2	K-1515C	Source Monitoring
2	Gallaher Bridge	Environmental Monitoring
3	K-1710	Environmental Monitoring
3	K-1700	Source Monitoring
3	K-1407E	Source Monitoring
3	K-1407F	Source Monitoring
3	K-1407J	Source Monitoring
3	K-1203	Source Monitoring
3	K-1007B	Source Monitoring
3	K-716	Environmental Monitoring
4	K-901A	Source Monitoring
4	K-1770	Environmental Monitoring
4	Clinch River (Brashar Island)	Environmental Monitoring
4	Kingston	Environmental Monitoring
13	West Fork Poplar Creek	Environmental Monitoring

methods were used, but the results were not validated against the EPA functional guidelines for CERCLA as was done for CR-ERP. These data were integrated with CR-ERP results and all of the data used in the human health risk screening analyses.

Combining data from various organizations, collected and analyzed for different purposes, makes interpretation of the summarized data more difficult. For example, the CR-ERP gamma spectroscopy results which were less than detection limits were reported as a minimum detectable activity along with a data qualifier for the value. Similar source monitoring gamma results were reported as background corrected values and associated counting uncertainty. Therefore, negative and very small values were common, and distinguishing non-detected values was not possible. These results were used as is and included as detected contaminants in the screening analyses. Resulting risk summary tables, therefore, list numerous radionuclides with very low risk values and, more importantly, indicate that these radionuclides were detected in all samples (See ratio of detected to total analyses in Tables E.8 thru E.12.).

Results of analyses performed on raw/unfiltered water samples (e.g., total aluminum) were used when available. The data for the water pathways included samples collected by both the CR-ERP and by the ORR compliance and monitoring organizations.

The values used for the risk screening analyses for a given contaminant in a reach were calculated by first averaging all results by location and then determining the upper 95% confidence limit of the arithmetic mean of the location averages. The UCL or the maximum location average (whichever is smaller) was used as the value for conservative screening for each reach. The arithmetic mean of the location averages was used for nonconservative screening.

Table E.2. Usage and exposure factors for conservative and nonconservative screening

Risk	Route	Media	Estimate	Age	Body weight (kg)		Exposure duration (yr)	Exposure frequency	Averaging time (yr)	Ingestion Rate	Intake Rate	Units
Cancer	Exposure	Sed.	C	A			30	0.2282				
Cancer	Exposure	Sed.	NC	A			30	0.0114				
Cancer	Ingestion	Fish	C	A	70		30	0.1370	70	2.30E-01	1.35E-02	kg/d
Cancer	Ingestion	Fish	NC	A	70		30	0.0329	70	2.30E-01	3.24E-03	kg/d
Cancer	Ingestion	Meat	C	A	70		30	1.0000	70	2.00E-01	8.57E-02	kg/d
Cancer	Ingestion	Milk	C	A	70		30	1.0000	70	5.00E-01	2.14E-01	L/d
Cancer	Ingestion	Sed.	C	A	70		30	1.0000	70	1.00E-04	4.29E-05	kg/d
Cancer	Ingestion	Sed.	NC	A	70		30	1.0000	70	1.00E-05	4.29E-06	kg/d
Cancer	Ingestion	Veg.	C	A	70		30	1.0000	70	2.00E-01	8.57E-02	kg/d
Cancer	Ingestion	Water	C	A	70		30	1.0000	70	2.00E+00	8.57E-01	L/d
Cancer	Ingestion	Water	NC	A	70		30	1.0000	70	2.00E-01	8.57E-02	L/d
Cancer	Inhalation	Sed.	C	A	70		30	1.0000	70	2.30E+01	9.86E+00	m3/d
Toxic	Ingestion	Fish	C	A	70		1	0.2740	1	2.30E-01	6.30E-02	kg/d
				C	20		1	0.2740	1	1.15E-01	3.15E-02	kg/d
Toxic	Ingestion	Fish	NC	A	70		1	0.0274	1	2.30E-01	6.30E-03	kg/d
Toxic	Ingestion	Meat	C	A	70		1	1.0000	1	2.00E-01	2.00E-01	kg/d
				C	20		1	1.0000	1	1.00E-01	1.00E-01	kg/d
Toxic	Ingestion	Milk	C	A	70		1	1.0000	1	5.00E-01	5.00E-01	L/d
				C	10		1	1.0000	1	1.00E+00	1.00E+00	L/d
Toxic	Ingestion	Sed.	C	A	70		1	1.0000	1	1.00E-04	1.00E-04	kg/d
				C	15		1	1.0000	1	2.00E-04	2.00E-04	kg/d

Table E.2 (continued)

Risk	Route	Media	Estimate	Age	Body weight (kg)	Exposure duration (yr)	Exposure frequency	Averaging time (yr)	Ingestion Rate	Intake Rate	Units
Toxic	Ingestion	Sed.	NC	A	70	1	1.0000	1	1.00E-05	1.00E-05	kg/d
Toxic	Ingestion	Veg.	C	C	70	1	1.0000	1	2.00E-01	2.00E-01	kg/d
									1.00E-01	1.00E-01	kg/d
Toxic	Ingestion	Water	C	C	70	1	1.0000	1	2.00E+00	2.00E+00	L/d
									1.00E+00	1.00E+00	L/d
Toxic	Ingestion	Water	NC	A	70	1	1.0000	1	2.00E-01	2.00E-01	L/d
Toxic	Inhalation	Sed.	C	A	70	1	1.0000	1	2.30E+01	2.30E+01	m3/d

Table E.3. Reference doses and slope (cancer potency) factors for inorganic compounds

CAS No.	Compound	Oral reference dose (mg/kg/d)	Inhalation reference concentration (mg/m ³)	Oral slope factor (mg/kg/d) ⁻¹	Inhalation slope factor (mg/kg/d) ⁻¹
7429-90-5	Aluminum	4.00E-04 ^a			
7440-36-0	Antimony	3.00E-04 ^a			
7440-38-2	Arsenic	7.00E-02 ^a	5.00E-04 ^b	1.75E+00 ^a	5.00E+01 ^b
7440-39-3	Barium	5.00E-03 ^a			
7440-41-7	Beryllium	9.00E-02 ^a	2.00E-02 ^b	4.30E+00 ^a	8.40E+00 ^b
7440-42-8	Boron	5.00E-04 ^a			
7440-43-9	Cadmium	5.00E-03 ^a			
18540-29-9	Chromium	3.70E-02 ^c			
7440-50-8	Copper	2.00E-02 ^a			
57-12-5	Cyanide				
7439-89-6	Iron				
7439-92-1	Lead				
7439-95-4	Magnesium				
7439-96-5	Manganese				
7439-97-6	Mercury				
7439-98-7	Molybdenum	1.00E-01 ^a	4.00E-04 ^a		
7440-02-0	Nickel	3.00E-04 ^b	3.00E-04 ^b		
7782-49-2	Selenium	5.00E-03 ^b			8.40E-01 ^a
7440-22-4	Silver	2.00E-02 ^a			
7440-24-6	Strontium	5.00E-03 ^a			
7440-28-0	Thallium	8.80E-01 ^b			
7440-31-5	Tin	7.00E-05 ^c			
7440-61-1	Uranium	6.00E-01 ^c			
7440-62-2	Vanadium	3.00E-03 ^a			
7440-66-6	Zinc	7.00E-03 ^b			
		2.00E-01 ^b			

^aUSEPA 1992. Health Effects Assessment Summary Tables FY-1992. OHEA ECAO-CIN-821. EPA Office of Research and Development and Office of Emergency and Remedial Response, Washington, D.C.

^bORNL 1992. Toxicity Values from the U.S. Environmental Protection Agency Integrated Risk Information System and Health Effects Assessment Summary Table. Health and Safety Research Division, Oak Ridge, TN.

^cEstimated value

Table E.4. Reference doses and slope (cancer potency) factors for organic compounds

CAS No.	Compound	Oral reference dose (mg/kg/d)	Inhalation reference concentration (mg/m ³)	Oral slope factor (mg/kg/d) ⁻¹	Inhalation slope factor (mg/kg/d) ⁻¹
71-55-6	1,1,1-Trichloroethane	9.00E-02 ^b	1.00E+00 ^b	2.00E-01 ^a	2.00E-01 ^b
79-34-5	1,1,2,2-Tetrachloroethane	4.00E-03 ^a		5.70E-02 ^a	5.70E-02 ^b
79-00-5	1,1,2-Trichloroethane	1.00E-01 ^b	5.00E-01 ^b	6.00E-01 ^a	1.20E+00 ^b
75-34-3	1,1-Dichloroethane	9.00E-03 ^a	9.00E-03 ^b		
75-35-4	1,1-Dichloroethylene	1.00E-02 ^b	2.00E-01 ^b		
120-82-1	1,2,4-Trichlorobenzene	9.00E-02 ^b			
95-50-1	1,2-Dichlorobenzene	1.00E-02 ^b			
107-06-2	1,2-Dichloroethane	1.00E-02 ^b			
156-59-2	1,2-Dichloroethene (total)	1.00E-02 ^b			
78-87-5	1,2-Dichloropropane		4.00E-03 ^a	6.80E-02 ^b	
541-73-1	1,3-Dichlorobenzene		7.00E-01 ^b	2.40E-02 ^b	
106-46-7	1,4-Dichlorobenzene				
95-95-4	2,4,5-Trichlorophenol	1.00E-01 ^a		1.10E-02 ^a	1.00E-02 ^b
88-06-2	2,4,6-Trichlorophenol				
120-83-2	2,4-Dichlorophenol	3.00E-03 ^a			
105-67-9	2,4-Dimethylphenol	2.00E-02 ^a			
51-28-5	2,4-Dinitrophenol	2.00E-03 ^a			
121-14-2	2,4-Dinitrotoluene				
606-20-2	2,6-Dinitrotoluene				
91-58-7	2-Chloronaphthalene	8.00E-02 ^a		6.80E-01 ^b	
95-57-8	2-Chlorophenol	5.00E-03 ^a		6.80E-01 ^b	
591-78-6	2-Hexanone				
91-57-6	2-Methylnaphthalene				
95-48-7	2-Methylphenol	5.00E-02 ^a			
88-74-4	2-Nitroaniline				
88-75-5	2-Nitrophenol				
91-94-1	3,3'-Dichlorobenzidine				
99-09-2	3-Nitroaniline			4.50E-01 ^a	
72-54-8	4,4'-DDD				
72-55-9	4,4'-DDE				
50-29-3	4,4'-DDT				
534-52-1	4,6-Dinitro-2-methylphenol	5.00E-04 ^a		2.40E-01 ^a	
101-55-3	4-Bromophenyl-phenylether	1.00E-04 ^c		3.40E-01 ^a	3.40E-01 ^b

Table B.4 (continued)

CAS No.	Compound	Oral reference dose (mg/kg/d)	Inhalation reference concentration (mg/m ³)	Oral slope factor (mg/kg/d) ⁻¹	Inhalation slope factor (mg/kg/d) ⁻¹
59-50-7	4-Chloro-3-methylphenol				
106-47-8	4-Chloroaniline	4.00E-03 ^a		3.50E-02 ^c	
7005-72-3	4-Chlorophenyl-phenylether	5.00E-02 ^b	8.00E-02 ^b		
108-10-1	4-Methyl-2-pentanone	5.00E-02 ^b			
106-44-5	4-Methylphenol				
100-01-6	4-Nitroaniline	6.00E-02 ^a		1.15E+01 ^c	
100-02-7	4-Nitrophenol	1.00E-01 ^a		1.15E+01 ^c	
83-32-9	Acenaphthene	3.00E-05 ^b			1.70E+01 ^b
208-96-8	Acenaphthylene				6.30E+00 ^b
67-64-1	Acetone				
309-00-2	Aldrin	3.00E-01 ^a		1.70E+01 ^a	
319-84-6	Alpha-BHC			6.30E+00 ^a	
120-12-7	Anthracene	3.00E-01 ^a		7.30E+00 ^c	
11097-69-1	Aroclor-1254			7.70E+00 ^c	
11096-82-5	Aroclor-1260			7.70E+00 ^c	
71-43-2	Benzene			2.90E-02 ^a	2.90E-02 ^b
92-87-5	Benzidine	3.00E-03 ^a		2.30E+02 ^a	2.30E+02 ^a
56-55-3	Benzo(a)anthracene			7.30E-01 ^c	6.10E+00 ^b
50-32-8	Benzo(a)pyrene			7.30E+00 ^a	
205-99-2	Benzo(b)fluoranthene			7.30E-01 ^c	
191-24-2	Benzo(g,h,i)perylene			7.30E+00 ^c	
207-08-9	Benzo(k)fluoranthene			7.30E+00 ^c	
65-85-0	Benzoic acid	4.00E+00 ^a		7.30E-01 ^c	
100-51-6	Benzyl alcohol	3.00E-01 ^b			
319-85-7	Beta-BHC			1.80E+00 ^a	1.80E+00 ^b
111-91-1	Bis(2-chloroethoxy)methane				
111-44-4	Bis(2-chloroethyl)ether	4.00E-02 ^a		1.10E+00 ^a	1.10E+00 ^b
108-60-1	Bis(2-chloroisopropyl)ether	2.00E-02 ^a		7.00E-02 ^b	3.50E-02 ^b
117-81-7	Bis(2-ethylhexyl)phthalate	2.00E-02 ^a		1.40E-02 ^a	
75-25-2	Bromoform	2.00E-02 ^a		7.90E-03 ^a	3.90E-03 ^b
85-68-7	Butylbenzylphthalate	2.00E-01 ^a			
75-15-0	Carbon disulfide	1.00E-01 ^a	1.00E-02 ^b	1.30E-01 ^a	5.30E-02 ^b
56-23-5	Carbon tetrachloride	7.00E-04 ^a			

Table E.4 (continued)

CAS No.	Compound	Oral reference dose (mg/kg/d)	Inhalation reference concentration (mg/m ³)	Oral slope factor (mg/kg/d) ⁻¹	Inhalation slope factor (mg/kg/d) ⁻¹
108-90-7	Chlorobenzene	2.00E-02 ^a	2.00E-02 ^b	1.30E+00 ^a	1.30E+00 ^b
57-73-9	Chloroethane	6.00E-05 ^a		9.40E-02 ^a	
124-48-1	Chlorodibromomethane	2.00E-02 ^a			
75-00-3	Chloroethane		1.00E+01 ^a	6.10E-03 ^a	8.10E-02 ^b
67-66-3	Chloroform	1.00E-02 ^a		1.30E-02 ^b	6.30E-03 ^b
74-87-3	Chloromethane			7.30E-02 ^c	
218-01-9	Chrysene	3.00E-04 ^c		1.80E-01 ^c	1.30E-01 ^c
10061-01-5	Cis-1,3-dichloropropene				
319-86-8	Delta-BHC	1.00E-01 ^b		1.15E+01 ^c	
84-74-2	Di-n-butylphthalate	2.00E-02 ^b			
117-84-0	Di-n-octylphthalate				
53-70-3	Dibenz(a,h)anthracene			1.30E-01 ^a	1.60E+01 ^b
132-64-9	Dibenzofuran	2.00E-02 ^a			
75-27-4	Dichlorobromomethane	5.00E-05 ^a		1.60E+01 ^a	
60-57-1	Dieldrin	8.00E-01 ^a			
84-66-2	Diethylphthalate	1.00E+00 ^b			
131-11-3	Dimethylphthalate	5.00E-05 ^c			
959-98-8	Endosulfan I	5.00E-05 ^c			
33213-65-9	Endosulfan II				
1031-07-8	Endosulfan sulfate	3.00E-04 ^a	1.00E+00 ^a		
72-20-8	Endrin	1.00E-01 ^a			
100-41-4	Ethylbenzene	4.00E-02 ^a		7.30E+00 ^c	
206-44-0	Fluoranthene	4.00E-02 ^a		1.15E+01 ^c	
86-73-7	Fluorene	3.00E-04 ^a		1.30E+00 ^b	
58-89-9	Gamma-BHC (Lindane)	5.00E-04 ^a		4.50E+00 ^a	4.50E+00 ^b
76-44-8	Heptachlor	1.30E-05 ^a		9.10E+00 ^b	9.10E+00 ^b
1024-57-3	Heptachlor epoxide	8.00E-04 ^a		1.60E+00 ^a	1.60E+00 ^b
118-74-1	Hexachlorobenzene	2.00E-03 ^a		7.80E-02 ^a	7.80E-02 ^b
87-68-3	Hexachlorobutadiene	7.00E-03 ^a	7.00E-05 ^b		
77-47-4	Hexachlorocyclopentadiene	1.00E-03 ^a		1.40E-02 ^a	1.40E-02 ^b
67-72-1	Hexachloroethane			7.30E-01 ^c	
193-39-5	Indeno(1,2,3-cd)pyrene	2.00E-01 ^a		4.10E-03 ^a	
78-59-1	Isophorone				

Table E.4 (continued)

CAS No.	Compound	Oral reference dose (mg/kg/d)	Inhalation reference concentration (mg/m ³)	Oral slope factor (mg/kg/d) ⁻¹	Inhalation slope factor (mg/kg/d) ⁻¹
72-43-5	Methoxychlor	5.00E-03 ^a			
74-83-9	Methyl bromide	1.40E-03 ^a	5.00E-03 ^a		
75-09-2	Methylene chloride	6.00E-02 ^a	3.00E+00 ^b	7.50E-03 ^a	
621-64-7	N-Nitroso-di-n-propylamine			7.00E+00 ^a	5.10E+01 ^b
62-75-9	N-Nitrosodimethylamine			5.10E+01 ^a	
86-30-6	N-Nitrosodiphenylamine			4.90E-03 ^a	
91-20-3	Naphthalene	4.00E-02 ^b			
98-95-3	Nitrobenzene	5.00E-04 ^a	2.00E-03 ^b		
87-86-5	Pentachlorophenol	3.00E-02 ^a		1.20E-01 ^a	
85-01-8	Phenanthrene			7.30E+00 ^c	
108-95-2	Phenol	6.00E-01 ^a			
129-00-0	Pyrene	3.00E-02 ^a		7.30E+00 ^c	
100-42-5	Styrene	2.00E-01 ^a		3.00E-02 ^c	2.00E-03 ^c
127-18-4	Tetrachloroethylene	1.00E-02 ^a		5.10E-02 ^c	
108-88-3	Toluene	2.00E-01 ^a	4.00E-01 ^b		
8001-35-2	Toxaphene			1.10E+00 ^a	1.10E+00 ^b
156-60-5	Trans-1,2-dichloroethene	2.00E-02 ^a			
10061-02-6	Trans-1,3-dichloropropene	3.00E-04 ^c		1.80E-01 ^c	1.30E-01 ^c
79-01-6	Trichloroethylene			1.10E-02 ^c	1.70E-02 ^c
75-69-4	Trichlorofluoromethane	3.00E-01 ^a	7.00E-01 ^b		
108-05-4	Vinyl acetate	1.00E+00 ^b	2.00E-01 ^a		
75-01-4	Vinyl chloride			1.90E+00 ^b	3.00E-01 ^b
1330-20-7	Xylene	2.00E+00 ^a			

^aUSEPA 1992. Health Effects Assessment Summary Tables FY-1992. OHEA ECAO-CIN-821. EPA Office of Research and Development and Office of Emergency and Remedial Response, Washington, D.C.

^bORNL 1992. Toxicity Values from the U.S. Environmental Protection Agency Integrated Risk Information System and Health Effects Assessment Summary Table. Health and Safety Research Division, Oak Ridge, TN.

^cEstimated value

Table E.5. Slope (cancer potency) factors for radionuclides^a

CAS No.	Radionuclide	External exposure		
		slope factor (kg/Bq)	Oral slope factor (1/Bq)	Inhalation slope factor (1/Bq)
14596-10-2	Am-241	1.30E-10	6.50E-09	8.60E-07
13981-15-2	Cm-244	8.10E-13	4.30E-09	5.90E-07
10198-40-0	Co-60	2.30E-07	4.10E-10	4.10E-09
10045-97-3	Cs-137	5.40E-08	7.60E-10	5.10E-10
10028-17-8	H-3		1.50E-12	2.10E-12
13994-20-2	Mp-237	1.20E-08	5.90E-09	7.80E-07
14119-24-5	Os-191	2.30E-09	1.80E-11	9.70E-11
15100-28-4	Pa-234	1.60E-07	1.80E-11	1.50E-11
13981-16-3	Pu-238	7.60E-13	5.90E-09	1.10E-06
15117-48-3	Pu-239	4.60E-13	6.20E-09	1.00E-06
10098-97-2	Sr-90		9.70E-10	1.70E-09
14133-76-7	Tc-99	1.60E-13	3.50E-11	2.20E-10
14274-82-9	Th-228	1.50E-07	1.50E-09	2.10E-06
15065-10-8	Th-234	9.50E-11	1.10E-10	8.60E-10
13966-29-5	U-234	8.10E-13	4.30E-10	7.00E-07
15117-96-1	U-235	6.50E-09	4.30E-10	6.80E-07
13982-70-2	U-236	6.50E-13	4.10E-10	6.80E-07
7440-61-1	U-238	9.70E-10	7.60E-10	1.40E-06

^aUSEPA 1992. Health Effects Assessment Summary Tables FY-1992. OHEA ECAO-CIN-821.

EPA Office of Research and Development and Office of Emergency and Remedial Response, Washington, D.C.

Table E.6 Summary of carcinogens in the secondary pathways identified as low priority for further investigation

Contaminant	Type	Contaminant	Reach					
			1	2	3	4	5	10
Organics								
		4,4'-DDD	G V	N	N	N	N	N
		4,4'-DDT	G H V	N	N	N	N	N
		Aroclor-1254	N			G	N	N
		Benzo(a)anthracene	G	G	G	N	N	N
		Benzo(a)pyrene	N	H	H	N	N	N
		Benzo(b)fluoranthene	N	G	G V	N	N	N
		Benzo(k)fluoranthene	G V	N	G V	N	N	N
		Bis(2-ethylhexyl)phthalate	N	G M L V	G M L V	G M L V	G M L V	G M L V
		Chrysene	G M L V	G M L V	G M L V	N	N	G
		Indeno(1,2,3-cd)pyrene	N	G V	N	N	N	N
		Toxaphene	N	H	N	N	N	N
Inorganics								
		Arsenic	H M L	H L	L	H M L	H M L	H M L
		Beryllium	H L	H L	H L	H L	H L	H L
		Cadmium	H	H	H	H	H	H

Table E.6 (continued)

Contaminant	Type	Contaminant	Reach					
			1	2	3	4	5	10
Inorganics (cont.)								
		Chromium		H				
		Nickel	H	H	H	H	H	H
Radionuclides								
		Am-241	S	X G H M L V	X G H M L V	X G H M L V	X G H M L V	X G H M L V
		Cm-243,244	S	X G H M L V	X G H M L V	X G H M L V	X G H M L V	X G H M L V
		Co-60	G H M L V	G H M L V	G H M L V	G H M L V	G H M L V	N N N N N
		Cs-137	G H	G H	G H	H	G H	G H
		Pu-238	S	X G H M L V	X G H M L V	X G H M L V	X G H M L V	X G H M L V
		Pu-239,240	S	X G H M L V	X G H M L V	X G H M L V	X G H M L V	X G H M L V

Table E.6 (continued)

Contaminant	Type	Contaminant	Reach					
			1	2	3	4	5	10
		Sr-90	S	G H M	G H M L	G H M	G H M	G H M L
		U-234	S	X G H M L V	X G H L V	X G H M L V	X G H M L V	X G H M L V
		U-235	S	X G H M L V	X G H M L V	X G H M L V	X G H M L V	X G H M L V
		U-238	S	X G H M L V	G H L V	X G H M L V	X G H M L V	X G H M L V

X	Carcinogens having conservative screening indices $< 10^{-4}$ for direct external exposure to radionuclides in dredged sediment.
G	Carcinogens having conservative screening indices $< 10^{-4}$ for direct ingestion of dredged sediment.
H	Carcinogens having conservative screening indices $< 10^{-4}$ for inhalation of air contaminated by dredged sediment.
M	Carcinogens having conservative screening indices $< 10^{-4}$ for ingestion of meat.
L	Carcinogens having conservative screening indices $< 10^{-4}$ for ingestion of milk.
V	Carcinogens having conservative screening indices $< 10^{-4}$ for ingestion of vegetables grown in dredged sediment.
N	Nondetects - contaminant concentrations were below detection limits.
S	No data available.

Table E.7 Summary of noncarcinogens in the secondary pathways identified as low priority for further investigation

Contaminant Type	Contaminant	Reach					
		1	2	3	4	5	10
Organics							
	4,4'-DDT	G M L V	N	N	N	N	N
	Anthracene	N	N	G M L V	N	N	G M L V
	Bis(2-ethylhexyl)phthalate	N	G M L V	G M L V	G M L V	G M L V	G M L V
	Di-n-butylphthalate	N	G M L V	G M L V	N	N	N
	Diethylphthalate	N	G M L V	N	N	N	N
	Dimethylphthalate	N	N	G M L V	N	N	N
	Fluoranthene	G M L V	G M L V	G M L V	G M L V	N	G M L V
	Naphthalene	N	G M L V	G M L V	N	N	N
	Pyrene	G M L V	G M L V	G M L V	G M L V	N	G M L V

Table E.7 (continued)

Contaminant Type	Contaminant	Reach					
		1	2	3	4	5	10
Inorganics	Antimony	N	S	G M L V	G M L V	G M L V	G M L V
	Arsenic	G M L	G M L	M L	G M L	G M L V	G M L V
	Beryllium	G M L V	G M L V	G M L V	G M L V	G M L V	G M L V
	Cadmium	G M	G M	G M	G	G	G M
	Chromium	G L	G L V	G	G L	G	G L
	Copper	G	L G V	G	G		G V
	Mercury	G H L	G H L	G H L	G H	G H	G H L M
	Nickel	G M L	G M L	G M L	G M L	G M L	G M L
	Selenium	G L V	G I V	G	G V	N	N
	Silver	G M L V	G M L V	G M V	G M	G M V	
	Thallium	N	N	G	N	N	N

Table E.7 (continued)

Contaminant Type	Contaminant	Reach					
		1	2	3	4	5	10
	Uranium	S	G M L V	G L V	G M L V	M L V	G M L V
	Zinc	G	G	G	G	G	G

X	Noncarcinogens having conservative screening indices < 0.1 for direct external exposure to radionuclides in dredged sediment.
G	Noncarcinogens having conservative screening indices < 0.1 for direct ingestion of dredged sediment.
H	Noncarcinogens having conservative screening indices < 0.1 for inhalation of air contaminated by dredged sediment.
M	Noncarcinogens having conservative screening indices < 0.1 for ingestion of meat.
L	Noncarcinogens having conservative screening indices < 0.1 for ingestion of milk.
V	Noncarcinogens having conservative screening indices < 0.1 for ingestion of vegetables grown in dredged sediment.
N	Nondetects - contaminant concentrations were below detection limits.
S	No data available.

[illegible]

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
1	Sed. Ingestion	Radionuclides	Cs-137 Co-60 TOTAL	4/4 4/4	8.49E+00 (G M) 2.34E+00 (G M)	7.6E-10 4.1E-10	9.3E-01 2.6E-01	7.1E-10 1.0E-10 8.1E-10
1	3ed. Ingestion							8.1E-10
	Water Ingestion	Radionuclides	Sr-90 U-234 Pu-239 Cs-137 U-238 Co-60 U-235 TOTAL	4/4 1/1 1/1 1/4 1/1 1/4 1/1	2.64E-02 (G M) 1.50E-02 (G M) 1.00E-03 (G M) 7.12E-03 (G M) 6.00E-03 (G M) 9.57E-03 (G M) 3.00E-03 (G M)	9.7E-10 4.3E-10 6.2E-09 7.6E-10 7.6E-10 4.1E-10 4.3E-10	Bq/Life 5.8E+01 3.3E+01 2.2E+00 1.6E+01 1.3E+01 2.1E+01 6.6E+00	5.6E-08 1.4E-08 1.4E-08 1.2E-08 1.0E-08 8.6E-09 2.8E-09 1.2E-07
		Inorganics	Beryllium Arsenic TOTAL	1/4 1/5	2.68E-03 (G M) 5.04E-03 (G M)	4.3E+00 1.8E+00	3.3E-06 6.2E-06	1.4E-05 1.1E-05 2.5E-05
1	Water Ingestion							2.5E-05
1								1.3E-04
2	Fish Ingestion	Radionuclides	Cs-137 Sr-90	23/27 6/6	9.13E+00 (G M) 1.53E-01 (G M)	7.6E-10 9.7E-10	7.6E+02 1.3E+01	5.7E-07 1.2E-08

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index			
2	Fish Ingestion	Inorganics	TOTAL		mg/kg	1/(mg/kg/d)	mg/kg/d	5.9E-07			
			Arsenic TOTAL	5/13	3.80E-02 (G M)	1.8E+00	1.8E-06	3.1E-06 3.1E-06			
		Organics	Aroclor-1280	14/14	6.69E-01 (G M)	7.7E+00	3.1E-05	2.4E-04			
			Aroclor-1254	9/10	1.21E-01 (G M)	7.7E+00	5.6E-06	4.3E-05			
			Chlordane	9/14	1.14E-02 (G M)	1.3E+00	5.3E-07	6.9E-07			
			Aldrin	2/14	1.28E-04 (G M)	1.7E+01	6.0E-09	1.0E-07			
			4,4'-DDE	6/13	5.29E-03 (G M)	3.4E-01	2.5E-07	8.3E-08			
			4,4'-DDT	2/13	4.67E-04 (G M)	3.4E-01	2.2E-08	7.4E-09 2.8E-04			
			TOTAL								
		2	Fish Ingestion							2.9E-04	
Radionuclides	Cs-137			46/46	6.64E+01 (G M)	5.4E-08	1.4E+01	7.8E-07			
	Co-60			44/46	3.21E+00 (G M)	2.3E-07	2.5E-01	5.8E-08 8.4E-07			
	TOTAL										
2	Sed. Exposure									8.4E-07	
2	Sed. Ingestion	Radionuclides	Cs-137	46/46	6.64E+01 (G M)	7.6E-10	7.3E+00	5.5E-09			
			Co-60	44/46	3.21E+00 (G M)	4.1E-10	3.5E-01	1.4E-10 5.7E-09			
		TOTAL									

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
3	Fish Ingestion	Radionuclides	Cs-137	26/54	Bq/kg	1/8q	Bq/life	
			Sr-90	28/28	1.78E+00 (G M)	7.6E-10	1.5E+02	1.1E-07
			Co-60	1/54	8.97E-02 (G M)	9.7E-10	7.4E+00	7.2E-09
			TOTAL		1.42E-01 (G M)	4.1E-10	1.2E+01	4.8E-09 1.2E-07
		Inorganics	Arsenic	19/35	mg/kg	1/(mg/kg/d)	mg/kg/d	
			Beryllium	7/35	5.31E-02 (G M)	1.8E+00	2.5E-06	4.3E-06
			TOTAL		9.22E-04 (G M)	4.3E+00	4.3E-08	1.8E-07 4.5E-06
		Organics	Aroclor-1260	36/36	mg/kg	1/(mg/kg/d)	mg/kg/d	
			Aroclor-1254	31/31	3.91E-01 (G M)	7.7E+00	1.8E-05	1.4E-04
			Chlordane	34/34	8.90E-02 (G M)	7.7E+00	4.6E-06	3.5E-05
			Bis(2-ethylhexyl)phthalate	1/33	3.99E-02 (G M)	1.3E+00	1.8E-06	2.4E-06
			4,4'-DDE	34/34	4.71E-02 (G M)	1.4E-02	2.2E-06	3.1E-08
			4,4'-DDT	2/34	1.04E-03 (G M)	3.4E-01	4.8E-08	1.6E-08
			TOTAL		3.00E-04 (G M)	2.4E-01	1.4E-08	3.3E-09 1.8E-04
3	Fish Ingestion							1.8E-04
	Sed. Exposure	Radionuclides	Cs-137	3/3	Bq/kg	1/Bq	Bq/yr	
			Co-60	3/3	2.55E+01 (G M)	5.4E-08	5.5E+00	3.0E-07
			TOTAL		1.76E+00 (G M)	2.3E-07	1.4E-01	3.2E-08 3.3E-07
3	Sed. Exposure							3.3E-07

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
<hr/>								
3	Sed. Ingestion	Radionuclides	Cs-137	3/3	2.55E+01 (G M)	1/Bq	Bq/life	2.1E-09
			Co-60	3/3	1.76E+00 (G M)	7.6E-10	2.8E+00	7.9E-11
			TOTAL			4.1E-10	1.9E-01	2.2E-09
<hr/>								
3	Sed. Ingestion							2.2E-09
<hr/>								
3	Water Ingestion	Radionuclides	Th-228	3/3	4.91E+01 (G M)	1/Bq	Bq/Life	1.6E-04
			Th-234	7/7	5.87E+00 (G M)	1.5E-09	1.1E+05	1.4E-06
			Np-237	7/7	4.84E-02 (G M)	1.1E-10	1.3E+04	6.3E-07
			Pu-238	7/7	2.42E-02 (G M)	5.9E-09	1.1E+02	3.1E-07
			Pa-234	7/7	7.61E+00 (G M)	5.9E-09	5.3E+01	3.0E-07
			U-234	6/6	2.59E-01 (G M)	1.8E-11	1.7E+04	2.4E-07
			Tc-99	8/8	3.04E+00 (G M)	4.3E-10	5.7E+02	2.3E-07
			U-238	6/6	1.38E-01 (G M)	3.5E-11	6.7E+03	2.3E-07
			Pu-239	7/7	4.36E-03 (G M)	7.6E-10	3.0E+02	5.9E-08
			Cs-137	8/11	2.07E-02 (G M)	6.2E-09	9.5E+00	3.4E-08
			Sr-90	4/4	7.40E-03 (G M)	7.6E-10	4.5E+01	1.6E-08
			U-235	7/7	1.42E-02 (G M)	9.7E-10	1.6E+01	1.3E-08
			U-236	6/6	3.00E-03 (G M)	4.3E-10	3.1E+01	2.7E-09
			TOTAL			4.1E-10	6.6E+00	1.6E-04
					mg/L	1/(mg/kg/d)	mg/kg/d	
		Inorganics	Arsenic	3/10	3.47E-03 (G M)	1.8E+00	4.2E-06	7.4E-06
			Beryllium	2/10	1.37E-03 (G M)	4.3E+00	1.7E-06	7.2E-06
			TOTAL					1.5E-05
		Organics	Pyrene	1/8	1.00E-02 (G M)	1/(mg/kg/d)	mg/kg/d	9.0E-05
			Fluoranthene	1/8	1.00E-02 (G M)	7.3E+00	1.2E-05	9.0E-05
			Anthracene	1/8	1.00E-02 (G M)	7.3E+00	1.2E-05	9.0E-05

=====

8.5E-04

=====

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
4	Water Ingestion	Radionuclides	Pu-239	4/4	Bq/L	1/Bq	Bq/life	8.8E-07
			Pu-238	4/4	6.50E-02 (G M)	6.2E-09	1.4E+02	6.1E-07
			Th-228	1/1	4.69E-02 (G M)	5.9E-09	1.0E+02	1.7E-07
			Th-234	3/3	5.18E-02 (G M)	1.5E-09	1.1E+02	1.2E-07
			Tc-98	3/3	4.84E-01 (G M)	1.1E-10	1.1E+03	8.4E-08
			U-234	1/1	1.09E+00 (G M)	3.5E-11	2.4E+03	7.8E-08
			U-238	1/1	8.24E-02 (G M)	4.3E-10	1.8E+02	7.3E-07
			H-3	1/1	4.41E-02 (G M)	7.6E-10	9.7E+01	5.5E-08
			Sr-90	2/2	1.68E+01 (G M)	1.5E-12	3.7E+04	2.5E-08
			Cs-137	5/6	1.19E-02 (G M)	9.7E-10	2.6E+01	2.0E-08
			Pa-234	3/3	1.20E-02 (G M)	7.6E-10	2.6E+01	1.7E-08
			Co-60	1/3	4.33E-01 (G M)	1.8E-11	9.5E+02	6.4E-09
			U-235	1/1	7.14E-03 (G M)	4.1E-10	1.6E+01	1.9E-09
			U-236	1/1	2.04E-03 (G M)	4.3E-10	4.5E+00	7.6E-10
			TOTAL	1/1	8.50E-04 (G M)	4.1E-10	1.9E+00	2.1E-06
								2.1E-06
								1.9E-04
5	Fish Ingestion	Radionuclides	Cs-137	19/28	Bq/kg	1/Bq	Bq/life	9.8E-08
			Sr-90	19/19	1.55E+00 (G M)	7.6E-10	1.3E+02	5.9E-09
			TOTAL	19/19	7.37E-02 (G M)	9.7E-10	6.1E+00	1.0E-07
		Inorganics	Arsenic	23/25	mg/kg	1/(mg/kg/d)	mg/kg/d	6.0E-06
			Beryllium	2/25	7.46E-02 (G M)	1.8E+00	3.5E-06	3.2E-09
			TOTAL		1.61E-05 (G M)	4.3E+00	7.4E-10	6.1E-06

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
					mg/kg	1/(mg/kg/d)	mg/kg/d	
5	Fish Ingestion	Organics	Aroclor-1260	19/19	3.48E-01 (G M)	7.7E+00	1.6E-05	1.2E-04
			Aroclor-1254	14/17	5.83E-02 (G M)	7.7E+00	2.7E-06	2.1E-05
			Chlordane	18/18	1.06E-02 (G M)	1.3E+00	4.9E-07	6.4E-07
			Aldrin	2/18	4.64E-04 (G M)	1.7E+01	2.2E-08	3.7E-07
			4,4'-DDE	9/18	8.03E-03 (G M)	3.4E-01	3.7E-07	1.3E-07
			4,4'-DDD	12/17	1.55E-03 (G M)	2.4E-01	7.2E-08	1.7E-08
			4,4'-DDT	5/18	7.03E-04 (G M)	3.4E-01	3.3E-08	1.1E-08
			TOTAL					1.5E-04

5	Fish Ingestion							1.5E-04

Sed. Exposure	Radionuclides	Cs-137 Co-60 TOTAL		274/277 116/277	Bq/kg	1/Bq	Bq/yr	
					1.02E+01 (G M) 5.35E-01 (G M)	5.4E-08 2.3E-07	2.2E+00 4.2E-02	1.2E-07 9.7E-09 1.3E-07

5	Sed. Exposure							1.3E-07

Sed. Ingestion	Radionuclides	Cs-137 Co-60 TOTAL		274/277 116/277	Bq/kg	1/Bq	Bq/life	
					1.02E+01 (G M) 5.35E-01 (G M)	7.6E-10 4.1E-10	1.1E+00 5.9E-02	8.5E-10 2.4E-11 8.7E-10

5	Sed. Ingestion							8.7E-10

Water Ingestion	Radionuclides	Sr-90		3/3	Bq/L	1/Bq	Bq/life	
					2.59E-02 (G M)	9.7E-10	5.7E+01	5.5E-08

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
5	Water Ingestion	Radionuclides	Cs-137 TOTAL	1/3	2.83E-03 (G M)	7.6E-10	6.2E+00	4.7E-09 6.0E-08
5	Water Ingestion							6.0E-08
5								1.5E-04
10	Fish Ingestion	Radionuclides	Cs-137 Sr-90 Co-60 TOTAL	9/17 8/8 1/16	Bq/kg 9.34E-01 (G M) 5.60E-02 (G M) 7.60E-02 (G M)	1/Bq 7.6E-10 9.7E-10 4.1E-10	Bq/life 7.7E+01 4.6E+00 6.3E+00	5.9E-08 4.5E-09 2.6E-09 6.6E-08
		Inorganics	Arsenic Beryllium TOTAL	14/15 1/10	mg/kg 7.14E-02 (G M) 9.70E-06 (G M)	1/(mg/kg/d) 1.8E+00 4.3E+00	mg/kg/d 3.3E-06 4.5E-10	5.8E-06 1.9E-09 5.8E-06
		Organics	Aroclor-1260 Chlordane 4,4'-DDE 4,4'-DDD TOTAL	7/8 8/8 2/8 7/8	mg/kg 2.20E-02 (G M) 8.32E-03 (G M) 2.56E-03 (G M) 6.87E-04 (G M)	1/(mg/kg/d) 7.7E+00 1.3E+00 3.4E-01 2.4E-01	mg/kg/d 1.0E-06 3.9E-07 1.2E-07 3.2E-08	7.8E-06 5.0E-07 4.0E-08 7.6E-09 8.4E-06
10	Fish Ingestion				mg/L	1/(mg/kg/d)	mg/kg/d	1.4E-05

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
10	Water Ingestion	Organics	Bis(2-ethylhexyl)phthalate TOTAL	1/1	7.50E-03 (G M)	1.4E-02	9.2E-06	1.3E-07 1.3E-07
10	Water Ingestion							1.3E-07
10								1.4E-05
13	Fish Ingestion	Radionuclides	Cs-137 Co-60 Sr-90 TOTAL	3/6 1/6 4/4	1.91E+00 (G M) 1.82E+00 (G M) 6.72E-02 (G M)	7.6E-10 4.1E-10 9.7E-10	1.6E+02 1.5E+02 5.6E+00	1.2E-07 6.2E-08 5.4E-09 1.9E-07
		Inorganics	Arsenic TOTAL	4/10	mg/kg 4.51E-02 (G M)	1/(mg/kg/d) 1.8E+00	mg/kg/d 2.1E-06	3.7E-06 3.7E-06
		Organics	Aroclor-1260 Aroclor-1254 Chlordane Aldrin 4,4'-DDE TOTAL	4/4 2/2 2/4 2/3 2/2	mg/kg 2.88E-01 (G M) 8.72E-02 (G M) 1.37E-02 (G M) 9.60E-04 (G M) 3.87E-03 (G M)	1/(mg/kg/d) 7.7E+00 7.7E+00 1.3E+00 1.7E+01 3.4E-01	mg/kg/d 1.3E-05 4.0E-06 6.3E-07 4.4E-08 1.8E-07	1.0E-04 3.1E-05 8.2E-07 7.6E-07 6.1E-08 1.4E-04
13	Fish Ingestion				Bq/L	1/Bq	Bq/Life	1.4E-04

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
13	Water Ingestion	Radionuclides	Th-228 Tc-99 Th-234 Cs-137 Pa-234 TOTAL	1/1 1/1 1/1 1/2 1/1	4.44E-02 (G M) 1.49E+00 (G M) 6.17E-02 (G M) 7.03E-03 (G M) 1.76E-01 (G M)	1.5E-09 3.5E-11 1.1E-10 7.6E-10 1.8E-11	9.7E+01 3.3E+03 1.4E+02 1.5E+01 3.8E+02	1.5E-07 1.1E-07 1.5E-08 1.2E-08 6.9E-09 2.9E-07
13	Water Ingestion							2.9E-07
13								1.4E-04
18	Sed. Exposure	Radionuclides	Cs-137 Co-60 TOTAL	22/23 1/23	3.66E+00 (G M) 8.94E-03 (G M)	5.4E-08 2.3E-07	7.9E-01 7.1E-04	4.3E-08 1.6E-10 4.3E-08
18	Sed. Exposure							4.3E-08
	Sed. Ingestion	Radionuclides	Cs-137 Co-60 TOTAL	22/23 1/23	3.66E+00 (G M) 8.94E-03 (G M)	7.6E-10 4.1E-10	4.0E-01 9.8E-04	3.0E-10 4.0E-13 3.0E-10
18	Sed. Ingestion							3.0E-10

Table E.8 Results of adult nonconservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
=====								=====
18								4.3E-08
=====								=====

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
1	Fish Ingestion	Inorganics	Arsenic	17/28	6.16E-02 (G M)	3.0E-04	mg/kg/d	1.8E-02
			Mercury	28/28	5.37E-02 (G M)	3.0E-04	5.5E-06	1.6E-02
			Selenium	28/28	7.48E-01 (G M)	5.0E-03	6.7E-05	1.3E-02
			Zinc	28/28	6.59E+00 (G M)	2.0E-01	5.9E-04	3.0E-03
			Beryllium TOTAL	4/25	1.47E-03 (G M)	5.0E-03	1.3E-07	2.6E-05
	Organics	Chlordane TOTAL		11/16	1.65E-02 (G M)	6.0E-05	mg/kg/d	2.5E-02
							1.5E-06	2.5E-02
1	Fish Ingestion	Inorganics	Arsenic	1/5	5.04E-03 (G M)	3.0E-04	mg/kg/d	4.8E-02
			Antimony	1/4	4.32E-03 (G M)	4.0E-04	1.2E-05	3.1E-02
			Chromium	3/6	7.86E-03 (G M)	5.0E-03	2.2E-05	4.5E-03
			Manganese	1/1	1.44E-01 (G M)	1.0E-01	4.1E-04	4.1E-03
			Cadmium	1/5	7.11E-04 (G M)	5.0E-04	2.0E-06	4.1E-03
	Organics	Chlordane TOTAL	Mercury	2/5	2.39E-04 (G M)	3.0E-04	6.8E-07	2.3E-03
			Nickel	3/6	1.59E-02 (G M)	2.0E-02	4.5E-05	2.3E-03
			Beryllium	1/4	2.68E-03 (G M)	5.0E-03	7.6E-06	1.5E-03
			Barium	1/1	3.30E-02 (G M)	7.0E-02	9.4E-05	1.3E-03
			Selenium	3/6	2.33E-03 (G M)	5.0E-03	6.7E-06	1.3E-03
Water Ingestion	Inorganics	Vanadium	1/1	2.48E-03 (G M)	7.0E-03	7.1E-06	1.0E-03	
		Copper	2/6	5.13E-03 (G M)	3.7E-02	1.5E-05	4.0E-04	
		Strontium	1/1	8.73E-02 (G M)	8.8E-01	2.5E-04	2.8E-04	
		Zinc	3/6	7.81E-03 (G M)	2.0E-01	2.2E-05	1.1E-04	
		TOTAL					1.0E-01	

Table E.8 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
1	Water Ingestion							1.0E-01
1								1.8E-01
2	Fish Ingestion	Inorganics	Mercury	13/13	6.38E-02 (G M)	3.0E-04	5.7E-06	1.9E-02
			Selenium	11/13	7.28E-01 (G M)	5.0E-03	6.5E-05	1.3E-02
			Arsenic	5/13	3.80E-02 (G M)	3.0E-04	3.4E-06	1.1E-02
			Zinc	13/13	5.74E+00 (G M)	2.0E-01	5.2E-04	2.6E-03
			TOTAL					4.6E-02
		Organics	Chlordane	9/14	1.14E-02 (G M)	6.0E-05	1.0E-06	1.7E-02
			Aldrin	2/14	1.28E-04 (G M)	3.0E-05	1.2E-08	3.9E-04
			4,4'-DDT	2/13	4.67E-04 (G M)	5.0E-04	4.2E-08	8.4E-05
			TOTAL					1.8E-02
2	Fish Ingestion							6.4E-02
	Water Ingestion	Inorganics	Cyanide	1/1	1.06E-01 (G M)	2.0E-02	3.0E-04	1.5E-02
			Nickel	1/3	2.00E-02 (G M)	2.0E-02	5.7E-05	2.9E-03
			Selenium	1/4	2.37E-03 (G M)	5.0E-03	6.8E-06	1.4E-03
			Barium	2/2	3.04E-02 (G M)	7.0E-02	8.7E-05	1.2E-03
			Manganese	1/1	3.51E-02 (G M)	1.0E-01	1.0E-04	1.0E-03
			Uranium	2/2	7.49E-04 (G M)	3.0E-03	2.1E-06	7.1E-04
			Copper	2/3	5.09E-03 (G M)	3.7E-02	1.5E-05	3.9E-04
			Strontium	1/1	8.90E-02 (G M)	8.8E-01	2.5E-04	2.9E-04
			Zinc	2/3	1.30E-02 (G M)	2.0E-01	3.7E-05	1.9E-04

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

2	Water Ingestion	Organics	TOTAL		mg/L	1/(mg/kg/d)	mg/kg/d	2.3E-02
			Tetrachloroethylene	1/3	4.92E-03 (G M)	1.0E-02	1.4E-05	1.4E-03
			Chloroform	1/2	4.47E-03 (G M)	1.0E-02	1.3E-05	1.3E-03
			Acetone	1/1	9.45E-03 (G M)	1.0E-01	2.7E-05	2.7E-04
			Carbon disulfide	1/1	4.69E-03 (G M)	1.0E-01	1.3E-05	1.3E-04
			Methylene chloride	2/3	2.79E-03 (G M)	6.0E-02	8.0E-06	1.3E-04
			Toluene	1/3	4.86E-03 (G M)	2.0E-01	1.4E-05	6.9E-05
			Phenol	1/2	1.11E-02 (G M)	6.0E-01	3.2E-05	5.3E-05
			Xylene	1/1	4.75E-03 (G M)	2.0E+00	1.4E-05	6.8E-06
			TOTAL					3.3E-03

2	Water Ingestion							2.6E-02

=====								
2								9.0E-02
=====								
3	Fish Ingestion	Inorganics	Mercury	35/35	4.23E-01 (G M)	3.0E-04	3.8E-05	1.3E-01
			Arsenic	19/35	5.31E-02 (G M)	3.0E-04	4.8E-06	1.6E-02
			Selenium	2/35	2.31E-01 (G M)	5.0E-03	2.1E-05	4.2E-03
			Zinc	35/35	5.76E+00 (G M)	2.0E-01	5.2E-04	2.6E-03
			Beryllium	7/35	9.22E-04 (G M)	5.0E-03	8.3E-08	1.7E-05
			TOTAL					1.5E-01
	Organics	Chlordane Bis(2-ethylhexyl)phthalate Di-n-octylphthalate TOTAL	34/34	3.99E-02 (G M)	6.0E-05	3.6E-06	6.0E-02	
			1/33	4.71E-02 (G M)	2.0E-02	4.2E-06	2.1E-04	
			1/33	2.40E-02 (G M)	2.0E-02	2.2E-06	1.1E-04	
							6.0E-02	

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
3	Fish Ingestion							2.1E-01
3	Water Ingestion	Inorganics	Arsenic	3/10	3.47E-03 (G M)	3.0E-04	9.9E-06	3.3E-02
			Cyanide	3/3	9.77E-02 (G M)	2.0E-02	2.8E-04	1.4E-02
			Uranium	8/8	7.18E-03 (G M)	3.0E-03	2.1E-05	6.8E-03
			Molybdenum	1/4	1.08E-02 (G M)	5.0E-03	3.1E-05	6.2E-03
			Chromium	7/12	1.05E-02 (G M)	5.0E-03	3.0E-05	6.0E-03
			Nickel	5/10	3.76E-02 (G M)	2.0E-02	1.1E-04	5.4E-03
			Cadmium	3/11	9.16E-04 (G M)	5.0E-04	2.6E-06	5.2E-03
			Manganese	8/6	1.27E-01 (G M)	1.0E-01	3.6E-04	2.6E-03
			Barium	3/4	6.46E-02 (G M)	7.0E-02	1.8E-04	3.6E-03
			Mercury	5/10	2.67E-04 (G M)	3.0E-04	7.6E-07	2.5E-03
			Silver	4/11	4.27E-03 (G M)	5.0E-03	1.2E-05	2.4E-03
			Selenium	2/11	3.34E-03 (G M)	5.0E-03	9.6E-06	1.9E-03
			Boron	4/4	2.56E-02 (G M)	9.0E-02	7.3E-05	8.1E-04
			Beryllium	2/10	1.37E-03 (G M)	5.0E-03	3.9E-06	7.8E-04
			Copper	6/10	8.10E-03 (G M)	3.7E-02	1.7E-05	4.7E-04
			Zinc	7/11	1.47E-02 (G M)	2.0E-01	4.2E-05	2.1E-04
			Tin	1/3	1.14E-02 (G M)	6.0E-01	3.3E-05	5.4E-05
			TOTAL					9.2E-02
		Organics	Trans-1,3-dichloropropene	1/5	5.00E-03 (G M)	3.0E-04	1.4E-05	4.8E-02
			2,4-Dichlorophenol	1/8	1.00E-02 (G M)	3.0E-03	2.9E-05	9.5E-03
			1,1-Dichloroethylene	2/5	4.91E-03 (G M)	9.0E-03	1.4E-05	1.6E-03
			Bis(2-ethylhexyl)phthalate	2/8	1.01E-02 (G M)	2.0E-02	2.9E-05	1.4E-03
			Tetrachloroethylene	3/5	3.87E-03 (G M)	1.0E-02	1.1E-05	1.1E-03
			Chloroform	3/5	3.46E-03 (G M)	1.0E-02	9.9E-06	9.9E-04
			Pyrene	1/8	1.00E-02 (G M)	3.0E-02	2.9E-05	9.6E-04
			Trans-1,2-dichloroethylene	3/5	5.14E-03 (G M)	2.0E-02	1.5E-05	7.3E-04
			Fluoranthene	1/8	1.00E-02 (G M)	4.0E-02	2.9E-05	7.2E-04
			Bromoform	1/5	5.01E-03 (G M)	2.0E-02	1.4E-05	7.2E-04
			Chlorobenzene	1/5	5.00E-03 (G M)	2.0E-02	1.4E-05	7.1E-04

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
3	Water Ingestion	Organics	Chlorodibromomethane	1/5	5.00E-03 (G M)	2.0E-02	1.4E-05	7.1E-04
			Dichlorobromomethane	2/5	4.88E-03 (G M)	2.0E-02	1.4E-05	6.9E-04
			Di-n-butylphthalate	1/8	9.88E-03 (G M)	1.0E-01	2.8E-05	2.8E-04
			Methylene chloride	3/5	4.84E-03 (G M)	6.0E-02	1.4E-05	2.3E-04
			Butylbenzylphthalate	1/8	1.00E-02 (G M)	2.0E-01	2.9E-05	1.4E-04
			Ethylbenzene	1/5	5.00E-03 (G M)	1.0E-01	1.4E-05	1.4E-04
			1,1-Dichloroethane	2/5	4.87E-03 (G M)	1.0E-01	1.4E-05	1.4E-04
			1,1,1-Trichloroethane	2/5	4.29E-03 (G M)	9.0E-02	1.2E-05	1.4E-04
			Anthracene	1/8	1.00E-02 (G M)	3.0E-01	2.9E-05	9.5E-05
			Toluene	2/5	4.95E-03 (G M)	2.0E-01	1.4E-05	7.1E-05
			Phenol	4/8	1.10E-02 (G M)	6.0E-01	3.2E-05	5.3E-05
			Diethylphthalate	1/8	1.00E-02 (G M)	8.0E-01	2.9E-05	3.6E-05
			Dimethylphthalate	1/8	1.00E-02 (G M)	1.0E+00	2.9E-05	2.9E-05
			TOTAL					6.9E-02
								1.6E-01
								3.7E-01
4	Fish Ingestion	Inorganics	Mercury	27/27	1.85E-01 (G M)	3.0E-04	1.7E-05	5.6E-02
			Arsenic	25/27	8.79E-02 (G M)	3.0E-04	7.9E-06	2.6E-02
			Selenium	14/27	5.57E-01 (G M)	5.0E-03	5.0E-05	1.0E-02
			Zinc	27/27	5.75E+00 (G M)	2.0E-01	5.2E-04	2.6E-03
			TOTAL					9.5E-02
		Organics	Chlordane	18/18	2.84E-02 (G M)	6.0E-05	2.6E-06	4.3E-02
			Aldrin	2/14	4.9E-04 (G M)	3.0E-05	7.6E-08	2.5E-03
			4,4'-DDT	1/18	3.8E-03 (G M)	5.0E-04	1.4E-07	2.8E-04
			TOTAL					4.5E-02

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
4	Fish Ingestion				mg/L	1/(mg/kg/d)	mg/kg/d	1.4E-01
4	Water Ingestion	Inorganics	Cyanide	1/2	9.17E-02 (G M)	2.0E-02	2.6E-04	1.3E-02
			Chromium	3/5	1.08E-02 (G M)	5.0E-03	3.1E-05	6.1E-03
			Nickel	2/4	1.82E-02 (G M)	2.0E-02	4.6E-05	2.3E-03
			Mercury	1/5	2.00E-04 (G M)	3.0E-04	5.7E-07	1.9E-03
			Manganese	2/2	5.12E-02 (G M)	1.0E-01	1.5E-04	1.5E-03
			Barium	1/1	3.30E-02 (G M)	7.0E-02	9.4E-05	1.3E-03
			Boron	1/1	2.30E-02 (G M)	9.0E-02	6.6E-05	7.3E-04
			Uranium	4/4	7.48E-04 (G M)	3.0E-03	2.1E-06	7.1E-04
			Copper	2/4	4.44E-03 (G M)	3.7E-02	1.3E-05	3.4E-04
			Zinc	2/4	5.71E-03 (G M)	2.0E-01	1.6E-05	8.2E-05
			TOTAL					2.8E-02
		Organics	Phenol	1/4	1.07E-02 (G M)	6.0E-01	3.0E-05	5.1E-05
			TOTAL					5.1E-05
4	Water Ingestion				mg/kg	1/(mg/kg/d)	mg/kg/d	2.8E-02
4								1.7E-01
5	Fish Ingestion	Inorganics	Arsenic	23/25	7.46E-02 (G M)	3.0E-04	6.7E-06	2.2E-02
			Mercury	25/25	6.60E-02 (G M)	3.0E-04	5.9E-06	2.0E-02
			Zinc	24/24	6.98E+00 (G M)	2.0E-01	6.3E-04	3.1E-03
			Copper	6/25	4.65E-02 (G M)	3.7E-02	4.2E-06	1.1E-04
			Beryllium	2/25	1.61E-05 (G M)	5.0E-03	1.4E-09	2.9E-07

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

5								
Fish Ingestion								

TOTAL					mg/kg	1/(mg/kg/d)	mg/kg/d	4.5E-02
Organics					1.06E-02 (G M)	6.0E-05	9.6E-07	1.6E-02
Chlordane					4.84E-04 (G M)	3.0E-05	4.2E-08	1.4E-03
Aldrin					7.03E-04 (G M)	5.0E-04	6.3E-08	1.3E-04
4,4'-DDT								1.7E-02
TOTAL								

5								
Fish Ingestion								

TOTAL					mg/kg	1/(mg/kg/d)	mg/kg/d	6.3E-02
=====								
5								
Fish Ingestion								

TOTAL					mg/kg	1/(mg/kg/d)	mg/kg/d	5.7E-02
Mercury					1.90E-01 (G M)	3.0E-04	1.7E-05	2.1E-02
Arsenic					7.14E-02 (G M)	3.0E-04	6.4E-06	9.4E-03
Selenium					5.22E-01 (G M)	5.0E-03	4.7E-05	2.4E-03
Zinc					5.22E+00 (G M)	2.0E-01	4.7E-04	1.7E-07
Beryllium					9.70E-06 (G M)	5.0E-03	8.7E-10	9.0E-02
TOTAL								

10								
Fish Ingestion								

TOTAL					mg/kg	1/(mg/kg/d)	mg/kg/d	1.2E-02
Chlordane					8.32E-03 (G M)	6.0E-05	7.5E-07	1.2E-02

10								
Fish Ingestion								

TOTAL					mg/L	1/(mg/kg/d)	mg/kg/d	1.0E-01

10								
Water Ingestion								

TOTAL					mg/L	1/(mg/kg/d)	mg/kg/d	1.1E-03
Organics					7.50E-03 (G M)	2.0E-02	2.1E-05	1.1E-03
Bis(2-ethylhexyl)phthalate								

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

TOTAL								1.1E-03

10	Water Ingestion							1.1E-03

=====								
10								1.0E-01
=====								
13	Fish Ingestion	Inorganics	Mercury	10/10	1.06E-01 (G M)	3.0E-04	9.5E-06	3.2E-02
			Arsenic	4/10	4.51E-02 (G M)	3.0E-04	4.1E-06	1.4E-02
			Zinc	10/10	6.31E+00 (G M)	2.0E-01	5.7E-04	2.8E-03
			TOTAL					4.8E-02

		Organics	Chlordane	2/4	1.37E-02 (G M)	6.0E-05	1.2E-06	2.1E-02
			Aldrin	2/3	9.60E-04 (G M)	3.0E-05	8.6E-08	2.9E-03
			TOTAL					2.3E-02

13	Fish Ingestion							7.1E-02

	Water Ingestion	Inorganics	Cyanide	1/1	8.79E-02 (G M)	2.0E-02	2.5E-04	1.3E-02
			Chromium	1/2	1.04E-02 (G M)	5.0E-03	3.0E-05	5.9E-03
			Manganese	1/1	1.54E-01 (G M)	1.0E-01	4.4E-04	4.4E-03
			Nickel	1/2	1.44E-02 (G M)	2.0E-02	4.1E-05	2.1E-03
			Zinc	1/2	1.04E-02 (G M)	2.0E-01	3.0E-05	1.5E-04
			TOTAL					2.5E-02

Table E.9 Results of adult nonconservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
13	Water Ingestion							2.5E-02
13								9.7E-02

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
					Bq/kg	1/Bq	Bq/life	
1	Fish Ingestion	Radionuclides	Cs-137	14/28	1.54E+00 (UCL)	7.6E-10	5.3E+02	4.0E-07
			Co-60	1/28	2.38E+00 (UCL)	4.1E-10	8.2E+02	3.4E-07
			Sr-90	8/8	3.70E-02 (Max)	9.7E-10	1.3E+01	1.2E-08
			TOTAL					7.5E-07
		Inorganics	Arsenic	17/28	mg/kg	1/(mg/kg/d)	mg/kg/d	3.3E-05
			Beryllium	4/25	9.82E-02 (UCL)	1.8E+00	1.9E-05	2.7E-06
			TOTAL		3.26E-03 (UCL)	4.3E+00	6.3E-07	3.6E-05
		Organics	Aroclor-1260	16/16	mg/kg	1/(mg/kg/d)	mg/kg/d	1.3E-03
			Aroclor-1254	13/16	8.97E-01 (UCL)	7.7E+00	1.7E-04	3.0E-04
			Chlordane	11/16	1.99E-01 (UCL)	7.7E+00	3.8E-05	2.7E-05
			4,4'-DDE	8/16	1.09E-01 (UCL)	1.3E+00	2.1E-05	4.2E-06
			4,4'-DDD	5/15	6.43E-02 (UCL)	3.4E-01	1.2E-05	2.7E-07
			TOTAL		5.90E-03 (UCL)	2.4E-01	1.1E-06	1.7E-03
1	Fish Ingestion							1.7E-03
	Sed. Exposure	Radionuclides	Cs-137	4/4	Bq/kg	1/Bq	Bq/yr	3.5E-06
			Co-60	4/4	1.48E+01 (Max)	5.4E-08	6.4E+01	3.2E-06
			TOTAL		8.91E+00 (Max)	2.3E-07	1.4E+01	6.7E-06
1	Sed. Exposure							6.7E-06
					Bq/kg	1/Bq	Bq/life	

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
1	Sed. Ingestion	Radionuclides	Cs-137 Co-60 TOTAL	4/4 4/4	1.48E+01 (Max) 8.91E+00 (Max)	7.6E-10 4.1E-10	1.6E+01 9.8E+00	1.2E-08 4.0E-09 1.6E-08
1	Sed. Ingestion							1.6E-08
	Sed. Inhalation	Radionuclides	Co-60 Cs-137 TOTAL	4/4 4/4	Bq/kg 8.91E+00 (Max) 1.48E+01 (Max)	1/Bq 4.1E-09 5.1E-10	Bq/Life 6.2E-03 2.8E-02	2.6E-11 1.4E-11 4.0E-11
1	Sed. Inhalation							4.0E-11
	Water Ingestion	Radionuclides	Cs-137 Co-60 Sr-90 U-234 Pu-239 U-238 U-235 TOTAL	1/4 1/4 4/4 1/1 1/1 1/1 1/1	Bq/L 1.49E-01 (Max) 1.90E-01 (Max) 8.00E-02 (Max) 1.50E-02 (Max) 1.00E-03 (Max) 6.00E-03 (Max) 3.00E-03 (Max)	1/Bq 7.6E-10 4.1E-10 9.7E-10 4.3E-10 6.2E-09 7.6E-10 4.3E-10	Bq/Life 3.3E+03 4.2E+03 1.8E+03 3.3E+02 2.2E+01 1.3E+02 6.6E+01	2.5E-06 1.7E-06 1.7E-06 1.4E-07 1.4E-07 1.0E-07 2.8E-08 6.3E-06
	Inorganics	Arsenic Beryllium TOTAL		1/5 1/4	mg/L 4.28E-02 (UCL) 3.90E-03 (Max)	1/(mg/kg/d) 1.8E+00 4.3E+00	mg/kg/d 5.2E-04 4.8E-05	9.2E-04 2.1E-04 1.1E-03

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
1	Water Ingestion	Radionuclides	Cs-137 Sr-90 TOTAL	23/27 6/6	Bq/kg	1/Bq	Bq/life	1.1E-03
					2.19E+01 (UCL)	7.6E-10	7.6E+03	5.8E-06
					2.59E-01 (Max)	9.7E-10	8.9E+01	8.7E-08
2	Fish Ingestion	Inorganics	Arsenic TOTAL	5/13	mg/kg	1/(mg/kg/d)	mg/kg/d	2.8E-03
					9.80E-02 (UCL)	1.8E+00	1.9E-05	3.3E-05
					mg/kg	1/(mg/kg/d)	mg/kg/d	3.3E-05
	Fish Ingestion	Organics	Aroclor-1260 Aroclor-1254 Chlordane 4,4'-DDE Aldrin 4,4'-DDT TOTAL	14/14 9/10 9/14 6/13 2/14 2/13	mg/kg	1/(mg/kg/d)	mg/kg/d	2.1E-03
					1.41E+00 (UCL)	7.7E+00	2.7E-04	1.1E-03
					7.53E-01 (UCL)	7.7E+00	1.5E-04	2.0E-05
2	Sed. Exposure	Radionuclides	Cs-137 Co-60	46/46 44/46	Bq/kg	1/Bq	Bq/yr	1.2E-06
					1.27E+02 (UCL)	5.4E-08	5.5E+02	3.0E-05
					5.22E+00 (UCL)	2.3E-07	8.3E+00	1.9E-06

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index

TOTAL								
2	Sed. Exposure							3.2E-05

2	Sed. Ingestion	Radionuclides	Cs-137 Co-60 TOTAL	46/46 44/46	Bq/kg 1.27E+02 (UCL) 5.22E+00 (UCL)	1/Bq 7.6E-10 4.1E-10	Bq/Life 1.4E+02 5.7E+00	1.1E-07 2.3E-09 1.1E-07

2	Sed. Ingestion							1.1E-07

	Sed. Inhalation	Radionuclides	Cs-137 Co-60 TOTAL	46/46 44/46	Bq/kg 1.27E+02 (UCL) 5.22E+00 (UCL)	1/Bq 5.1E-10 4.1E-09	Bq/Life 2.4E-01 3.7E-03	1.2E-10 1.5E-11 1.4E-10

2	Sed. Inhalation							1.4E-10

	Water Ingestion	Radionuclides	H-3 Sr-90 Cs-137 Np-237 Th-228 Co-60 U-234 Os-191 Th-234 Tc-99	2/2 4/4 4/5 1/1 1/1 2/4 1/1 1/1 1/1 1/1	Bq/L 6.80E+03 (Max) 8.42E+00 (Max) 2.14E+00 (UCL) 1.18E-01 (Max) 3.55E-01 (Max) 2.77E-01 (Max) 1.90E-01 (Max) 3.30E+00 (Max) 1.04E+00 (Max)	1/Bq 1.5E-12 9.7E-10 7.6E-10 5.9E-09 1.5E-09 4.1E-10 4.3E-10 1.8E-11 3.5E-11	Bq/Life 1.5E+08 1.8E+05 4.7E+04 2.6E+03 7.8E+03 6.1E+03 4.2E+03 7.2E+04 8.7E+03 2.3E+04	2.2E-04 1.8E-04 3.6E-05 1.5E-05 1.2E-05 2.5E-06 1.8E-06 1.3E-06 9.6E-07 8.0E-07

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
2	Water Ingestion	Radionuclides	U-235	2/2	6.40E-02 (Max)	4.3E-10	1.4E+03	6.0E-07
			U-238	1/1	3.40E-02 (Max)	7.6E-10	7.4E+02	5.7E-07
			Pa-234	1/1	1.51E-01 (Max)	1.8E-11	3.3E+03	6.0E-08
			Pu-238	3/3	1.75E-05 (Max)	5.9E-09	3.8E-01	2.3E-09
			TOTAL					4.7E-04
		Organics	1,2-Dichloropropane	1/3	5.00E-03 (Max)	6.8E-02	6.1E-05	4.2E-06
			Tetrachloroethylene	1/3	5.00E-03 (Max)	5.1E-02	6.1E-05	3.1E-06
			Methylene chloride	2/3	5.00E-03 (Max)	7.5E-03	6.1E-05	4.6E-07
			Chloroform	1/2	5.00E-03 (Max)	6.1E-03	6.1E-05	3.7E-07
			TOTAL					8.1E-06

2	Water Ingestion							

=====								
2								
=====								
3	Fish Ingestion	Radionuclides	Cs-137	26/54	2.98E+00 (UCL)	7.6E-10	1.0E+03	7.8E-07
			Co-60	1/54	2.88E+00 (UCL)	4.1E-10	9.9E+02	4.1E-07
			Sr-90	28/28	1.51E-01 (UCL)	9.7E-10	5.2E+01	5.1E-08
			TOTAL					1.2E-06
		Inorganics	Arsenic	19/35	7.50E-02 (UCL)	1.8E+00	1.4E-05	2.5E-05
			Beryllium	7/35	5.45E-03 (UCL)	4.3E+00	1.1E-06	4.5E-06
			TOTAL					3.0E-05
		Organics	Aroclor-1280	36/36	7.71E-01 (UCL)	7.7E+00	1.5E-04	1.1E-03

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
3	Fish Ingestion	Organics	Aroclor-1254	31/31	1.86E-01 (UCL)	7.7E+00	3.6E-05	2.8E-04
			Chlordane	34/34	7.45E-02 (UCL)	1.3E+00	1.4E-05	1.9E-05
			Bis(2-ethylhexyl)phthalate	1/33	4.10E-01 (Max)	1.4E-02	7.9E-05	1.1E-06
			4,4'-DDD	2/34	1.71E-02 (Max)	2.4E-01	3.3E-06	7.9E-07
			4,4'-DDE	34/34	9.10E-03 (UCL)	3.4E-01	1.8E-06	6.0E-07
			TOTAL				1.4E-03	

3	Fish Ingestion							1.5E-03

3	Sed. Exposure	Radionuclides	Cs-137	3/3	8.24E+01 (Max)	5.4E-08	3.6E+02	1.9E-05
			Co-60	3/3	5.00E+00 (Max)	2.3E-07	7.9E+00	1.8E-06
			TOTAL					2.1E-05
							2.1E-05	

3	Sed. Exposure							

3	Sed. Ingestion	Radionuclides	Cs-137	3/3	8.24E+01 (Max)	7.6E-10	9.0E+01	6.9E-08
			Co-60	3/3	5.00E+00 (Max)	4.1E-10	5.5E+00	2.2E-09
			TOTAL					7.1E-08
							7.1E-08	

3	Sed. Ingestion							

3	Sed. Inhalation	Radionuclides	Cs-137	3/3	8.24E+01 (Max)	5.1E-10	1.6E-01	8.1E-11
			Co-60	3/3	5.00E+00 (Max)	4.1E-09	3.5E-03	1.4E-11
			TOTAL					9.5E-11

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
3	Sed. Inhalation							9.5E-11
3	Water Ingestion	Radionuclides	Th-228	3/3	5.99E+01 (Max)	1.5E-09	1.3E+06	2.0E-03
			U-234	6/6	9.38E+00 (UCL)	4.3E-10	2.1E+05	8.8E-05
			U-238	6/6	5.04E+00 (UCL)	7.6E-10	1.1E+05	8.4E-05
			Pa-234	7/7	1.27E+02 (UCL)	1.8E-11	2.8E+06	5.0E-05
			Th-234	7/7	1.74E+01 (UCL)	1.1E-10	3.8E+05	4.2E-05
			Np-237	7/7	1.90E-01 (UCL)	5.9E-09	4.2E+03	2.5E-05
			Tc-99	8/8	1.02E+01 (UCL)	3.5E-11	2.2E+05	7.8E-06
			Pu-238	7/7	3.84E-02 (UCL)	5.9E-09	8.4E+02	5.0E-06
			Pu-239	7/7	1.90E-02 (UCL)	6.2E-09	4.2E+02	2.6E-06
			Cs-137	8/11	1.41E-01 (UCL)	7.6E-10	3.1E+03	2.4E-06
			U-235	7/7	2.08E-01 (UCL)	4.3E-10	4.6E+03	2.0E-06
			U-236	6/6	9.70E-02 (UCL)	4.1E-10	2.1E+03	8.7E-07
			Sr-90	4/4	7.40E-03 (Max)	9.7E-10	1.6E+02	1.6E-07
			TOTAL					2.3E-03
		Inorganics	Arsenic	3/10	mg/L	1/(mg/kg/d)	mg/kg/d	
			Beryllium	2/10	1.14E-02 (UCL)	1.8E+00	1.4E-04	2.4E-04
			TOTAL		3.20E-03 (UCL)	4.3E+00	3.9E-05	1.7E-04
								4.1E-04
		Organics	Pyrene	1/8	mg/L	1/(mg/kg/d)	mg/kg/d	
			Fluoranthene	1/8	1.01E-02 (UCL)	7.3E+00	1.2E-04	9.0E-04
			Anthracene	1/8	1.01E-02 (UCL)	7.3E+00	1.2E-04	9.0E-04
			Phenanthrene	1/8	1.01E-02 (UCL)	7.3E+00	1.2E-04	9.0E-04
			N-Nitroso-di-n-propylamine	1/8	1.01E-02 (UCL)	7.3E+00	1.2E-04	8.6E-04
			Vinyl chloride	2/5	1.00E-02 (Max)	1.9E+00	1.2E-04	2.3E-04
			Aroclor-1254	2/3	1.04E-03 (Max)	7.7E+00	1.3E-05	9.8E-05
			Benzo(a)anthracene	1/8	1.01E-02 (UCL)	7.3E-01	1.2E-04	9.0E-05
			1,1-Dichloroethylene	2/5	5.01E-03 (Max)	6.0E-01	6.1E-05	3.7E-05

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
3	Water Ingestion	Organics	Trans-1,3-dichloropropene	1/5	5.01E-03 (UCL)	1.8E-01	6.1E-05	1.1E-05
			Chrysene	1/8	1.01E-02 (UCL)	7.3E-02	1.2E-04	9.0E-06
			Dichlorobromomethane	2/5	5.00E-03 (Max)	1.3E-01	6.1E-05	8.0E-06
			1,2-Dichloroethane	1/5	5.31E-03 (UCL)	9.1E-02	6.5E-05	5.9E-06
			Chlorodibromomethane	1/5	5.00E-03 (Max)	8.4E-02	6.1E-05	5.1E-06
			Trichloroethylene	3/5	3.20E-02 (UCL)	1.1E-02	3.9E-04	4.3E-06
			Tetrachloroethylene	3/5	5.03E-03 (UCL)	5.1E-02	6.2E-05	3.1E-06
			1,4-Dichlorobenzene	1/9	1.00E-02 (Max)	2.4E-02	1.2E-04	2.9E-06
			Benzene	2/5	5.00E-03 (Max)	2.9E-02	1.1E-05	1.8E-06
			Bis(2-ethylhexyl)phthalate	2/8	1.03E-02 (UCL)	1.4E-02	1.3E-04	1.8E-06
			Methylene chloride	3/5	5.53E-03 (UCL)	7.5E-03	6.8E-05	5.1E-07
			Bromoform	1/5	5.03E-03 (UCL)	7.9E-03	6.2E-05	4.9E-07
			Chloroform	3/5	5.10E-03 (UCL)	6.1E-03	6.2E-05	3.8E-07
		TOTAL					5.0E-03	
3	Water Ingestion							7.7E-03
3								9.2E-03
4	Fish Ingestion	Radionuclides	Cs-137	28/40	4.69E+00 (UCL)	7.6E-10	1.6E+03	1.2E-06
			Sr-90	10/10	1.34E-01 (UCL)	9.7E-10	4.6E+01	4.5E-08
			TOTAL					1.3E-06
					mg/kg	1/(mg/kg/d)	mg/kg/d	
		Inorganics	Arsenic	25/27	1.07E-01 (UCL)	1.8E+00	2.1E-05	3.6E-05
			TOTAL					3.6E-05
		Organics	Aroclor-1260	17/18	2.79E+00 (UCL)	7.7E+00	5.4E-04	4.1E-03
			Aroclor-1254	15/18	2.70E-01 (Max)	7.7E+00	5.2E-05	4.0E-04

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
4	Fish Ingestion	Organics	Chlordane	18/18	9.72E-02 (UCL)	1.3E+00	1.9E-05	2.4E-05
			4,4'-DDT	1/18	2.00E-01 (Max)	3.4E-01	3.9E-05	1.3E-05
			Aldrin	2/14	1.33E-03 (UCL)	1.7E+01	2.6E-07	4.4E-06
			4,4'-DDE	12/18	2.44E-02 (UCL)	3.4E-01	4.7E-06	1.6E-06
			4,4'-DDD	4/18	9.04E-03 (UCL)	2.4E-01	1.7E-06	4.2E-07
			TOTAL				4.6E-03	

4	Fish Ingestion				Bq/kg	1/Bq	Bq/yr	4.6E-03

4	Sed. Exposure	Radionuclides	Cs-137	93/93	1.33E+02 (UCL)	5.4E-08	5.8E+02	3.1E-05
			Co-60	93/95	6.43E+00 (UCL)	2.3E-07	1.0E+01	2.3E-06
			TOTAL				3.3E-05	

4	Sed. Exposure				Bq/kg	1/Bq	Bq/life	3.3E-05

4	Sed. Ingestion	Radionuclides	Cs-137	93/93	1.33E+02 (UCL)	7.6E-10	1.5E+02	1.1E-07
			Co-60	93/95	6.43E+00 (UCL)	4.1E-10	7.0E+00	2.9E-09
			TOTAL				1.1E-07	

4	Sed. Ingestion				Bq/kg	1/Bq	Bq/life	1.1E-07

4	Sed. Inhalation	Radionuclides	Cs-137	93/93	1.33E+02 (UCL)	5.1E-10	2.5E-01	1.3E-10
			Co-60	93/95	6.43E+00 (UCL)	4.1E-09	4.5E-03	1.8E-11
			TOTAL				1.5E-10	

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
4	Sed. Inhalation							1.5E-10
4	Water Ingestion	Radionuclides	Pa-234 Pu-239 Pu-238 Th-228 Th-234 Cs-137 Tc-99 U-234 U-238 H-3 Sr-90 Co-60 U-235 U-236 TOTAL	3/3 4/4 4/4 1/1 3/3 5/8 3/3 1/1 1/1 1/1 2/2 1/3 1/1 1/1	7.17E+01 (Max) 6.50E-02 (Max) 4.69E-02 (Max) 5.18E-02 (Max) 6.91E-01 (Max) 7.18E-02 (UCL) 1.09E+00 (Max) 8.24E-02 (Max) 4.41E-02 (Max) 1.68E+01 (Max) 1.19E-02 (Max) 1.88E-02 (Max) 2.04E-03 (Max) 8.50E-04 (Max)	1/Bq	Bq/Life	2.8E-05 8.8E-06 6.1E-06 1.7E-06 1.7E-06 1.2E-06 8.4E-07 7.8E-07 7.3E-07 5.5E-07 2.5E-07 1.8E-07 1.9E-08 7.6E-09 5.1E-05
4	Water Ingestion							5.1E-05
4								4.7E-03
5	Fish Ingestion	Radionuclides	Cs-137 Sr-90 TOTAL	19/28 19/19	2.64E+00 (UCL) 1.23E-01 (UCL)	1/Bq	Bq/Life	6.9E-07 4.1E-08 7.3E-07
					mg/kg	1/(mg/kg/d)	mg/kg/d	

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index		
5	Fish Ingestion	Inorganics	Arsenic Beryllium TOTAL	23/25	9.23E-02 (UCL)	1.8E+00	1.8E-05	3.1E-05		
				2/25	2.70E-02 (Max)	4.3E+00	5.2E-06	2.2E-05 5.4E-05		
		Organics	Aroclor-1260 Aroclor-1254 Aldrin Chlordane 4,4'-DDE 4,4'-DDD 4,4'-DDT TOTAL	19/19	8.90E-01 (UCL)	7.7E+00	1.7E-04	1.3E-03		
				14/17	3.20E-01 (Max)	7.7E+00	6.2E-05	4.8E-04		
				2/18	5.45E-03 (UCL)	1.7E+01	1.1E-06	1.8E-05		
				18/18	4.55E-02 (UCL)	1.3E+00	8.8E-06	1.1E-05		
				9/18	1.32E-02 (UCL)	3.4E-01	2.9E-06	8.7E-07		
				12/17	5.83E-03 (UCL)	2.4E-01	1.1E-06	2.7E-07		
				5/18	1.29E-03 (UCL)	3.4E-01	2.5E-07	8.5E-08 1.8E-03		
				5	Fish Ingestion	Radionuclides	Cs-137 Co-60 TOTAL	274/277	2.70E+01 (UCL)	5.4E-08
116/277	1.46E+00 (UCL)	2.3E-07	2.3E+00					5.3E-07 6.9E-06		
Sed. Exposure			Bq/kg			1/Bq	Bq/yr		6.9E-06	
5	Sed. Exposure	Radionuclides	Cs-137 Co-60 TOTAL			274/277	2.70E+01 (UCL)	7.6E-10	3.0E+01	2.2E-08
						116/277	1.46E+00 (UCL)	4.1E-10	1.6E+00	6.6E-10 2.3E-08

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
5	Sed. Ingestion							2.3E-08
5	Sed. Inhalation	Radionuclides	Cs-137 Co-60 TOTAL	274/277 116/277	Bq/kg 2.70E+01 (UCL) 1.46E+00 (UCL)	1/Bq 5.1E-10 4.1E-09	Bq/life 5.2E-02 1.0E-03	2.6E-11 4.2E-12 3.1E-11
5	Sed. Inhalation							3.1E-11
	Water Ingestion	Radionuclides	Sr-90 Cs-137 TOTAL	3/3 1/3	Bq/L 2.59E-02 (Max) 3.27E-03 (Max)	1/Bq 9.7E-10 7.6E-10	Bq/life 5.7E+02 7.2E+01	5.5E-07 5.4E-08 6.0E-07
5	Water Ingestion							6.0E-07
5								1.9E-03
10	Fish Ingestion	Radionuclides	Cs-137 Co-60 Sr-90 TOTAL	9/17 1/16 8/8	Bq/kg 1.58E+00 (UCL) 8.02E-01 (UCL) 9.38E-02 (UCL)	1/Bq 7.6E-10 4.1E-10 9.7E-10	Bq/life 5.4E+02 2.8E+02 3.2E+01	4.1E-07 1.1E-07 3.1E-08 5.6E-07
	Inorganics	Arsenic		14/15	mg/kg 9.68E-02 (UCL)	1/(mg/kg/d) 1.8E+00	mg/kg/d 1.9E-05	3.3E-05

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
10	Fish Ingestion	Inorganics	Beryllium TOTAL	1/10	3.70E-02 (Max)	4.3E+00	7.1E-06	3.1E-05 6.3E-05
		Organics	Aroclor-1260 Chlordane 4,4'-DDE 4,4'-DDD TOTAL	7/8 8/8 2/8 7/8	mg/kg 7.31E-02 (UCL) 1.57E-02 (UCL) 2.95E-03 (UCL) 1.22E-03 (UCL)	1/(mg/kg/d) 7.7E+00 1.3E+00 3.4E-01 2.4E-01	mg/kg/d 1.4E-05 3.0E-06 5.7E-07 2.4E-07	1.1E-04 3.9E-06 1.9E-07 5.6E-08 1.1E-04
10	Fish Ingestion				mg/L	1/(mg/kg/d)	mg/kg/d	1.8E-04
	Water Ingestion	Organics	Bis(2-ethylhexyl)phthalate TOTAL	1/1	7.50E-03 (Max)	1.4E-02	9.2E-05	1.3E-06 1.3E-06
10	Water Ingestion				Bq/kg	1/Bq	Bq/life	1.3E-06
10		Radionuclides	Cs-137 Co-60 Sr-90 TOTAL	3/6 1/6 4/4	Bq/kg 3.45E+00 (UCL) 2.68E+00 (UCL) 1.11E-01 (Max)	7.6E-10 4.1E-10 9.7E-10	1.2E+03 9.2E+02 3.8E+01	9.0E-07 3.8E-07 3.7E-08 1.3E-06
		Inorganics	Arsenic	4/10	mg/kg 5.90E-02 (UCL)	1/(mg/kg/d) 1.8E+00	mg/kg/d 1.1E-05	2.0E-05

Table E.10 Results of adult conservative screening for carcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
13 Fish Ingestion								
		Organics	Aroclor-1260	4/4	3.90E-01 (Max)	7.7E+00	7.5E-05	2.0E-05
			Aroclor-1254	2/2	1.90E-01 (Max)	7.7E+00	3.7E-05	5.8E-04
			Chlordane	2/4	7.29E-02 (Max)	1.3E+00	1.4E-05	2.8E-04
			Aldrin	2/3	1.30E-03 (UCL)	1.7E+01	2.5E-07	1.8E-05
			4,4'-DDE	2/2	6.82E-03 (Max)	3.4E-01	1.3E-06	4.3E-06
			TOTAL					4.5E-07
								8.8E-04
13 Fish Ingestion								
		Radionuclides	Th-228	1/1	4.44E-02 (Max)	1.5E-09	9.7E+02	1.5E-06
			Tc-99	1/1	1.49E+00 (Max)	3.5E-11	3.3E+04	1.1E-06
			Cs-137	1/2	1.24E-02 (Max)	7.6E-10	2.7E+02	2.1E-07
			Th-234	1/1	6.17E-02 (Max)	1.1E-10	1.4E+03	1.5E-07
			Pa-234	1/1	1.76E-01 (Max)	1.8E-11	3.8E+03	6.9E-08
			TOTAL					3.0E-06
13 Water Ingestion								
		Radionuclides	Cs-137	22/23	8.66E+00 (UCL)	5.4E-08	3.8E+01	2.0E-06
			Co-60	1/23	1.39E+00 (Max)	2.3E-07	2.2E+00	5.1E-07
			TOTAL					2.5E-06
18 Sed. Exposure								
		Radionuclides	Cs-137	22/23	8.66E+00 (UCL)	5.4E-08	3.8E+01	2.0E-06
			Co-60	1/23	1.39E+00 (Max)	2.3E-07	2.2E+00	5.1E-07
			TOTAL					2.5E-06

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
18	Sed. Exposure							2.5E-06
18	Sed. Ingestion	Radionuclides	Cs-137 Co-60 TOTAL	22/23 1/23	8.66E+00 (UCL) 1.39E+00 (Max)	1/Bq	Bq/life	7.2E-09 6.3E-10 7.8E-09
18	Sed. Ingestion							7.8E-09
	Sed. Inhalation	Radionuclides	Cs-137 Co-60 TOTAL	22/23 1/23	8.66E+00 (UCL) 1.39E+00 (Max)	1/Bq	Bq/life	8.5E-12 4.0E-12 1.2E-11
18	Sed. Inhalation							1.2E-11
18								2.5E-06

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
					mg/kg	1/(mg/kg/d)	mg/kg/d	
1	Fish Ingestion	Inorganics	Arsenic	17/28	9.82E-02 (UCL)	3.0E-04	8.8E-05	2.9E-01
			Mercury	28/28	8.19E-02 (UCL)	3.0E-04	7.4E-05	2.5E-01
			Selenium	28/28	8.34E-01 (UCL)	5.0E-03	7.5E-04	1.5E-01
			Zinc	28/28	7.55E+00 (UCL)	2.0E-01	6.8E-03	3.4E-02
			Beryllium	4/25	3.26E-03 (UCL)	5.0E-03	2.9E-06	5.9E-04
			TOTAL					7.3E-01
		Organics	Chlordane	11/16	1.09E-01 (UCL)	6.0E-05	9.9E-05	1.6E+00
			TOTAL					1.6E+00
1	Fish Ingestion							2.4E+00
1	Water Ingestion	Inorganics	Arsenic	1/5	4.28E-02 (UCL)	3.0E-04	1.2E-03	4.1E+00
			Antimony	1/4	5.09E-02 (Max)	4.0E-04	1.5E-03	3.6E+00
			Cadmium	1/5	4.95E-03 (UCL)	5.0E-04	1.4E-04	2.8E-01
			Selenium	3/6	1.84E-02 (UCL)	5.0E-03	5.3E-04	1.1E-01
			Chromium	3/6	9.80E-03 (Max)	5.0E-03	2.8E-04	5.6E-02
			Nickel	3/6	3.02E-02 (UCL)	2.0E-02	8.6E-04	4.3E-02
			Manganese	1/1	1.44E-01 (Max)	1.0E-01	4.1E-03	4.1E-02
			Mercury	2/5	3.50E-04 (UCL)	3.0E-04	1.0E-05	3.3E-02
			Beryllium	1/4	3.90E-03 (Max)	5.0E-03	1.1E-04	2.2E-02
			Barium	1/1	3.30E-02 (Max)	7.0E-02	9.4E-04	1.3E-02
			Vanadium	1/1	2.48E-03 (Max)	7.0E-03	7.1E-05	1.0E-02
			Copper	2/6	7.44E-03 (UCL)	3.7E-02	2.1E-04	5.7E-03
			Strontium	1/1	8.73E-02 (Max)	8.8E-01	2.5E-03	2.8E-03
			Zinc	3/6	1.40E-02 (UCL)	2.0E-01	4.0E-04	2.0E-03
			TOTAL					8.3E+00

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
1	Water Ingestion	Inorganics	Mercury	13/13	1.35E-01 (UCL)	3.0E-04	1.2E-04	4.1E-01
			Arsenic	5/13	9.80E-02 (UCL)	3.0E-04	8.8E-05	2.9E-01
1	Water Ingestion	Inorganics	Selenium	11/13	8.51E-01 (UCL)	5.0E-03	7.7E-04	1.5E-01
			Zinc	13/13	6.57E+00 (UCL)	2.0E-01	5.9E-03	3.0E-02
			TOTAL					8.8E-01
2	Fish Ingestion	Organics	Chlordane	9/14	8.16E-02 (UCL)	6.0E-05	7.3E-05	1.2E+00
			Aldrin	2/14	1.65E-04 (UCL)	3.0E-05	1.5E-07	5.0E-03
			4,4'-DDT	2/13	8.59E-04 (UCL)	5.0E-04	7.7E-07	1.5E-03
			TOTAL					1.2E+00
2	Fish Ingestion	Inorganics	Cyanide	1/1	1.06E-01 (Max)	2.0E-02	3.0E-03	1.5E-01
			Selenium	1/4	6.72E-03 (Max)	5.0E-03	1.9E-04	3.8E-02
			Nickel	1/3	2.21E-02 (Max)	2.0E-02	6.3E-04	3.2E-02
			Barium	2/2	3.30E-02 (Max)	7.0E-02	9.4E-04	1.3E-02
			Manganese	1/1	3.51E-02 (Max)	1.0E-01	1.0E-03	1.0E-02
			Uranium	2/2	8.42E-04 (Max)	3.0E-03	2.4E-05	8.0E-03
			Zinc	2/3	5.20E-02 (Max)	2.0E-01	1.5E-03	7.4E-03
			Copper	2/3	6.50E-03 (Max)	3.7E-02	1.9E-04	5.0E-03
			Strontium	1/1	8.90E-02 (Max)	8.8E-01	2.5E-03	2.9E-03

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
2	Water Ingestion	Organics	TOTAL		mg/L	1/(mg/kg/d)	mg/kg/d	2.7E-01
			Tetrachloroethylene	1/3	5.00E-03 (Max)	1.0E-02	1.4E-04	1.4E-02
			Chloroform	1/2	5.00E-03 (Max)	1.0E-02	1.4E-04	1.4E-02
			Acetone	1/1	9.45E-03 (Max)	1.0E-01	2.7E-04	2.7E-03
			Methylene chloride	2/3	5.00E-03 (Max)	6.0E-02	1.4E-04	2.4E-03
			Carbon disulfide	1/1	4.69E-03 (Max)	1.0E-01	1.3E-04	1.3E-03
			Toluene	1/3	5.00E-03 (Max)	2.0E-01	1.4E-04	7.1E-04
			Phenol	1/2	1.23E-02 (Max)	6.0E-01	3.5E-04	5.8E-04
			Xylene	1/1	4.75E-03 (Max)	2.0E+00	1.4E-04	6.8E-05
			TOTAL					3.6E-02

2	Water Ingestion							3.0E-01

=====								
2								2.4E+00
=====								
3	Fish Ingestion	Inorganics	Mercury	35/35	6.16E-01 (UCL)	3.0E-04	5.5E-04	1.8E+00
			Arsenic	19/35	7.50E-02 (UCL)	3.0E-04	6.8E-05	2.3E-01
			Selenium	2/35	5.77E-01 (UCL)	5.0E-03	5.2E-04	1.0E-01
			Zinc	35/35	6.55E+00 (UCL)	2.0E-01	5.9E-03	2.9E-02
			Beryllium	7/35	5.45E-03 (UCL)	5.0E-03	4.9E-06	9.8E-04
			TOTAL					2.2E+00
		Organics	Chlordane	34/34	7.45E-02 (UCL)	6.0E-05	6.7E-05	1.1E+00
			Bis(2-ethylhexyl)phthalate	1/33	4.10E-01 (Max)	2.0E-02	3.7E-04	1.8E-02
			Di-n-octylphthalate	1/33	4.10E-01 (Max)	2.0E-02	3.7E-04	1.8E-02
			TOTAL					1.2E+00

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
3	Fish Ingestion				mg/L	1/(mg/kg/d)	mg/kg/d	3.4E+00
3	Water Ingestion	Inorganics	Uranium	8/8	2.91E-01 (UCL)	3.0E-03	8.3E-03	2.8E+00
			Arsenic	3/10	1.14E-02 (UCL)	3.0E-04	3.3E-04	1.1E+00
			Nickel	5/10	1.29E-01 (UCL)	2.0E-02	3.7E-03	1.8E-01
			Cyanide	3/3	1.01E-01 (Max)	2.0E-02	2.9E-03	1.4E-01
			Cadmium	3/11	2.13E-03 (UCL)	5.0E-04	6.1E-05	1.2E-01
			Molybdenum	1/4	1.38E-02 (Max)	5.0E-03	3.9E-04	7.9E-02
			Boron	4/4	2.31E-01 (Max)	9.0E-02	6.6E-03	7.3E-02
			Chromium	7/12	1.13E-02 (UCL)	5.0E-03	3.2E-04	6.5E-02
			Manganese	6/6	2.19E-01 (UCL)	1.0E-01	6.2E-03	6.2E-02
			Selenium	2/11	1.08E-02 (UCL)	5.0E-03	3.1E-04	6.2E-02
			Silver	4/11	9.36E-03 (UCL)	5.0E-03	2.7E-04	5.3E-02
			Barium	3/4	1.18E-01 (Max)	7.0E-02	3.4E-03	4.8E-02
			Mercury	5/10	5.03E-04 (UCL)	3.0E-04	1.4E-05	4.8E-02
			Beryllium	2/10	3.20E-03 (UCL)	5.0E-03	9.1E-05	1.8E-02
			Copper	6/10	1.54E-02 (UCL)	3.7E-02	4.4E-04	1.2E-02
			Zinc	7/11	2.98E-02 (UCL)	2.0E-01	8.5E-04	4.2E-03
			Tin	1/3	1.48E-02 (Max)	6.0E-01	4.2E-04	7.0E-04
			TOTAL					4.8E+00
		Organics	Trans-1,3-dichloropropene	1/5	5.01E-03 (UCL)	3.0E-04	1.4E-04	4.8E-01
			2,4-Dichlorophenol	1/8	1.01E-02 (UCL)	3.0E-03	2.9E-04	9.6E-02
			Trans-1,2-dichloroethylene	3/5	2.23E-02 (UCL)	2.0E-02	6.4E-04	3.2E-02
			1,1-Dichloroethylene	2/5	5.01E-03 (Max)	9.0E-03	1.4E-04	1.6E-02
			Bis(2-ethylhexyl)phthalate	2/8	1.03E-02 (UCL)	2.0E-02	2.9E-04	1.5E-02
			Chloroform	3/5	5.10E-03 (UCL)	1.0E-02	1.5E-04	1.5E-02
			Tetrachloroethylene	3/5	5.03E-03 (UCL)	1.0E-02	1.4E-04	1.4E-02
			Pyrene	1/8	1.01E-02 (UCL)	3.0E-02	2.9E-04	9.6E-03
			Fluoranthene	1/8	1.01E-02 (UCL)	4.0E-02	2.9E-04	7.2E-03
			Bromoforn	1/5	5.03E-03 (UCL)	2.0E-02	1.4E-04	7.2E-03
			Dichlorobromomethane	2/5	5.00E-03 (Max)	2.0E-02	1.4E-04	7.1E-03

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
3	Water Ingestion	Organics	Chlorobenzene	1/5	5.00E-03 (Max)	2.0E-02	1.4E-04	7.1E-03
			Chlorodibromomethane	1/5	5.00E-03 (Max)	2.0E-02	1.4E-04	7.1E-03
			Di-n-butylphthalate	1/8	1.00E-02 (Max)	1.0E-01	2.9E-04	2.9E-03
			Methylene chloride	3/5	5.53E-03 (UCL)	6.0E-02	1.6E-04	2.6E-03
			1,1,1-Trichloroethane	2/5	5.02E-03 (Max)	9.0E-02	1.4E-04	1.6E-03
			Butylbenzylphthalate	1/8	1.00E-02 (UCL)	2.0E-01	2.9E-04	1.4E-03
			1,1-Dichloroethane	2/5	5.00E-03 (Max)	1.0E-01	1.4E-04	1.4E-03
			Ethylbenzene	1/5	5.00E-03 (Max)	1.0E-01	1.4E-04	1.4E-03
			Anthracene	1/8	1.01E-02 (UCL)	3.0E-01	2.9E-04	9.6E-04
			Toluene	2/5	5.00E-03 (Max)	2.0E-01	1.4E-04	7.1E-04
			Phenol	4/8	1.24E-02 (UCL)	6.0E-01	3.5E-04	5.9E-04
			Diethylphthalate	1/8	1.01E-02 (UCL)	8.0E-01	2.9E-04	3.6E-04
			Dimethylphthalate	1/8	1.01E-02 (UCL)	1.0E+00	2.9E-04	7.2E-04
			TOTAL					
								5.6E+00
								8.9E+00
4	Fish Ingestion	Inorganics	Mercury	27/27	3.91E-01 (UCL)	3.0E-04	3.5E-04	1.2E+00
			Arsenic	25/27	1.07E-01 (UCL)	3.0E-04	9.7E-05	3.2E-01
			Selenium	14/27	6.86E-01 (UCL)	5.0E-03	6.2E-04	1.2E-01
			Zinc	27/27	6.55E+00 (UCL)	2.0E-01	5.9E-03	2.9E-02
			TOTAL					1.6E+00
		Organics	Chlordane	18/18	9.72E-02 (UCL)	6.0E-05	8.7E-05	1.5E+00
			4,4'-DDT	1/18	2.00E-01 (Max)	5.0E-04	1.8E-04	3.6E-01
			Aldrin	2/14	1.33E-03 (UCL)	3.0E-05	1.2E-06	4.0E-02
			TOTAL					1.9E+00

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
4	Fish Ingestion							3.5E+00
4	Water Ingestion	Inorganics	Cyanide Chromium Uranium Nickel Mercury Manganese Barium Boron Copper Zinc TOTAL	1/2 3/5 4/4 2/4 1/5 2/2 1/1 1/1 2/4 2/4	9.59E-02 (Max) 1.62E-02 (Max) 3.58E-03 (UCL) 1.90E-02 (Max) 2.01E-04 (UCL) 6.17E-02 (Max) 3.30E-02 (Max) 2.30E-02 (Max) 4.53E-03 (Max) 8.12E-03 (Max)	2.0E-02 5.0E-03 3.0E-03 2.0E-02 3.0E-04 1.0E-01 7.0E-02 9.0E-02 3.7E-02 2.0E-01	2.7E-03 4.6E-04 1.0E-04 5.4E-04 5.8E-06 1.8E-03 9.4E-04 6.6E-04 1.3E-04 2.3E-04	1.4E-01 9.3E-02 3.4E-02 2.7E-02 1.9E-02 1.8E-02 1.3E-02 7.3E-03 3.5E-03 1.2E-03 3.5E-01
4	Water Ingestion	Organics	Phenol TOTAL	1/4	1.29E-02 (Max)	6.0E-01	3.7E-04	6.1E-04 6.1E-04
4	Water Ingestion							3.5E-01
4	Water Ingestion							3.9E+00
5	Fish Ingestion	Inorganics	Copper Mercury Arsenic Zinc Beryllium	6/25 25/25 23/25 24/24 2/25	1.70E+02 (Max) 1.08E-01 (UCL) 9.23E-02 (UCL) 1.09E+01 (UCL) 2.70E-02 (Max)	3.7E-02 3.0E-04 3.0E-04 2.0E-01 5.0E-03	1.5E-01 9.7E-05 8.3E-05 9.8E-03 2.4E-05	4.1E+00 3.2E-01 2.8E-01 4.9E-02 4.9E-03

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

5	Fish Ingestion	Organics	Chlordane Aldrin 4,4'-DDT TOTAL	18/18 2/18 5/18	mg/kg 4.55E-02 (UCL) 5.45E-03 (UCL) 1.29E-03 (UCL)	1/(mg/kg/d) 6.0E-05 3.0E-05 5.0E-04	mg/kg/d 4.1E-05 4.9E-06 1.2E-06	4.8E+00 6.8E-01 1.6E-01 2.3E-03 8.5E-01

5	Fish Ingestion							5.6E+00

=====								
5								5.6E+00
=====								
10	Fish Ingestion	Inorganics	Mercury Arsenic Selenium Zinc Beryllium TOTAL	15/15 14/15 8/15 15/15 1/10	mg/kg 2.80E-01 (UCL) 9.68E-02 (UCL) 5.82E-01 (UCL) 5.97E+00 (UCL) 3.70E-02 (Max)	1/(mg/kg/d) 3.0E-04 3.0E-04 5.0E-03 2.0E-01 5.0E-03	mg/kg/d 2.5E-04 8.7E-05 5.2E-04 5.4E-03 3.3E-05	8.4E-01 2.9E-01 1.0E-01 2.7E-02 6.7E-03 1.3E+00

10	Fish Ingestion	Organics	Chlordane TOTAL	8/8	mg/kg 1.57E-02 (UCL)	1/(mg/kg/d) 6.0E-05	mg/kg/d 1.4E-05	2.4E-01 2.4E-01

10	Fish Ingestion							1.5E+00

	Water Ingestion	Organics	Bis(2-ethylhexyl)phthalate	1/1	mg/L 7.50E-03 (Max)	1/(mg/kg/d) 2.0E-02	mg/kg/d 2.1E-04	1.1E-02

Table E.11 Results of adult conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

10								
Water Ingestion								

TOTAL								
								1.1E-02

10								
Water Ingestion								

TOTAL								
								1.1E-02

13								
Fish Ingestion								

		Inorganics	Mercury	10/10	3.67E-01 (UCL)	3.0E-04	3.3E-04	1.1E+00
			Arsenic	4/10	5.90E-02 (UCL)	3.0E-04	5.3E-05	1.8E-01
			Zinc	10/10	6.98E+00 (UCL)	2.0E-01	6.3E-03	3.1E-02
			TOTAL					1.3E+00

		Organics	Chlordane	2/4	7.29E-02 (Max)	6.0E-05	6.6E-05	1.1E+00
			Aldrin	2/3	1.30E-03 (UCL)	3.0E-05	1.2E-06	3.9E-02
			TOTAL					1.1E+00

13								
Fish Ingestion								

TOTAL								
								2.4E+00

13								
Water Ingestion								

		Inorganics	Cyanide	1/1	8.79E-02 (Max)	2.0E-02	2.5E-03	1.3E-01
			Chromium	1/2	1.0E-02 (Max)	5.0E-03	3.1E-04	6.3E-02
			Manganese	1/1	1.54E-01 (Max)	1.0E-01	4.4E-03	4.4E-02
			Nickel	1/2	1.90E-02 (Max)	2.0E-02	5.4E-04	2.7E-02
			Zinc	1/2	1.80E-02 (Max)	2.0E-01	5.1E-04	2.6E-03
			TOTAL					2.6E-01

[illegible]

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
					mg/kg	1/(mg/kg/d)	mg/kg/d	
1	Fish Ingestion	Inorganics	Arsenic	17/28	9.82E-02 (UCL)	3.0E-04	1.5E-04	5.2E-01
			Mercury	28/28	8.19E-02 (UCL)	3.0E-04	1.3E-04	4.3E-01
			Selenium	28/28	8.34E-01 (UCL)	5.0E-03	1.3E-03	2.6E-01
			Zinc	28/28	7.55E+00 (UCL)	2.0E-01	1.2E-02	5.9E-02
			Beryllium TOTAL	4/25	3.28E-03 (UCL)	5.0E-03	5.1E-06	1.0E-03 1.3E+00
1	Fish Ingestion	Organics	Chlordane TOTAL	11/16	1.09E-01 (UCL)	6.0E-05	1.7E-04	2.9E+00 2.9E+00
					mg/kg	1/(mg/kg/d)	mg/kg/d	4.1E+00
1	Water Ingestion	Inorganics	Arsenic	1/5	4.28E-02 (UCL)	3.0E-04	2.9E-03	9.5E+00
			Antimony	1/4	5.09E-02 (Max)	4.0E-04	3.4E-03	8.5E+00
			Cadmium	1/5	4.95E-03 (UCL)	5.0E-04	3.3E-04	6.6E-01
			Selenium	3/6	1.84E-02 (UCL)	5.0E-03	1.2E-03	2.5E-01
			Chromium	3/6	9.80E-03 (Max)	5.0E-03	6.5E-04	1.3E-01
			Nickel	3/6	3.02E-02 (UCL)	2.0E-02	2.0E-03	1.0E-01
			Manganese	1/1	1.44E-01 (Max)	1.0E-01	9.6E-03	9.6E-02
			Mercury	2/5	3.50E-04 (UCL)	3.0E-04	2.3E-05	7.8E-02
			Beryllium	1/4	3.90E-03 (Max)	5.0E-03	2.6E-04	5.2E-02
			Barium	1/1	3.30E-02 (Max)	7.0E-02	3.1E-03	3.1E-02
			Vanadium	1/1	2.48E-03 (Max)	7.0E-03	1.7E-04	2.4E-02
			Copper	2/6	7.44E-03 (UCL)	3.7E-02	5.0E-04	1.3E-02
			Strontium	1/1	8.73E-02 (Max)	8.8E-01	5.8E-03	6.6E-03
			Zinc	3/6	1.40E-02 (UCL)	2.0E-01	9.3E-04	4.7E-03
			TOTAL					1.9E+01

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
1	Water Ingestion							1.9E+01
2	Fish Ingestion	Inorganics	Mercury	13/13	1.35E-01 (UCL)	3.0E-04	2.1E-04	7.1E-01
			Arsenic	5/13	9.80E-02 (UCL)	3.0E-04	1.5E-04	5.1E-01
			Selenium	11/13	8.51E-01 (UCL)	5.0E-03	1.3E-03	2.7E-01
			Zinc	13/13	6.57E+00 (UCL)	2.0E-01	1.0E-02	5.2E-02
			TOTAL					1.5E+00
		Organics	Chlordane	9/14	8.16E-02 (UCL)	6.0E-05	1.3E-04	2.1E+00
			Aldrin	2/14	1.65E-04 (UCL)	3.0E-05	2.6E-07	8.7E-03
			4,4'-DDT	2/13	8.59E-04 (UCL)	5.0E-04	1.4E-06	2.7E-03
			TOTAL					2.2E+00
								3.7E+00
2	Fish Ingestion							
	Water Ingestion	Inorganics	Cyanide	1/1	1.08E-01 (Max)	2.0E-02	7.0E-03	3.5E-01
			Selenium	1/4	6.72E-03 (Max)	5.0E-03	4.5E-04	9.0E-02
			Nickel	1/3	2.21E-02 (Max)	2.0E-02	1.5E-03	7.4E-02
			Barium	2/2	3.30E-02 (Max)	7.0E-02	2.2E-03	3.1E-02
			Manganese	1/1	3.51E-02 (Max)	1.0E-01	2.3E-03	2.3E-02
			Uranium	2/2	8.42E-04 (Max)	3.0E-03	5.6E-05	1.9E-02
			Zinc	2/3	5.20E-02 (Max)	2.0E-01	3.5E-03	1.7E-02
			Copper	2/3	6.50E-03 (Max)	3.7E-02	4.3E-04	1.2E-02
			Strontium	1/1	8.90E-02 (Max)	8.8E-01	5.9E-03	6.7E-03

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index		
2	Water Ingestion	Organics	TOTAL			1/(mg/kg/d)	mg/kg/d	6.2E-01		
			tetrachloroethylene	1/3	5.00E-03 (Max)	1.0E-02	3.3E-04	3.3E-02		
			Chloroform	1/2	5.00E-03 (Max)	1.0E-02	3.3E-04	3.3E-02		
			Acetone	1/1	9.45E-03 (Max)	1.0E-01	6.3E-04	6.3E-03		
			Methylene chloride	2/3	5.00E-03 (Max)	6.0E-02	3.3E-04	5.6E-03		
			Carbon disulfide	1/1	4.69E-03 (Max)	1.0E-01	3.1E-04	3.1E-03		
			Toluene	1/3	5.00E-03 (Max)	2.0E-01	3.3E-04	1.7E-03		
			Phenol	1/2	1.23E-02 (Max)	6.0E-01	8.2E-04	1.4E-03		
			Xylene	1/1	4.75E-03 (Max)	2.0E+00	3.2E-04	1.6E-04		
			TOTAL					8.5E-02		
									7.1E-01	
										Water Ingestion
3	Fish Ingestion	Inorganics	Mercury	35/35	6.18E-01 (UCL)	3.0E-04	9.7E-04	3.2E+00		
			Arsenic	19/35	7.50E-02 (UCL)	3.0E-04	1.2E-04	3.9E-01		
			Selenium	2/35	5.77E-01 (UCL)	5.0E-03	9.1E-04	1.8E-01		
			Zinc	35/35	6.55E+00 (UCL)	2.0E-01	1.0E-02	5.2E-02		
			Beryllium	7/35	5.45E-03 (UCL)	5.0E-03	8.6E-06	1.7E-03		
			TOTAL					3.9E+00		
				Organics	Chlordane	34/34	7.45E-02 (UCL)	6.0E-05	1.2E-04	2.0E+00
					Bis(2-ethylhexyl)phthalate	1/33	4.10E-01 (Max)	2.0E-02	6.5E-04	3.2E-02
					Di-n-octylphthalate	1/33	4.10E-01 (Max)	2.0E-02	6.5E-04	3.2E-02
	TOTAL							2.0E+00		

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
3	Fish Ingestion							5.9E+00
3	Water Ingestion	Inorganics	Uranium	8/8	2.91E-01 (UCL)	3.0E-03	1.9E-02	6.5E+00
			Arsenic	3/10	1.14E-02 (UCL)	3.0E-04	7.6E-04	2.5E+00
			Nickel	5/10	1.29E-01 (UCL)	2.0E-02	8.8E-03	4.3E-01
			Cyanide	3/3	1.01E-01 (Max)	2.0E-02	6.8E-03	3.4E-01
			Cadmium	3/11	2.13E-03 (UCL)	5.0E-04	1.4E-04	2.8E-01
			Molybdenum	1/4	1.38E-02 (Max)	5.0E-03	9.2E-04	1.8E-01
			Boron	4/4	2.31E-01 (Max)	9.0E-02	1.5E-02	1.7E-01
			Chromium	7/12	1.13E-02 (UCL)	5.0E-03	7.5E-04	1.5E-01
			Manganese	6/6	2.19E-01 (UCL)	1.0E-01	1.5E-02	1.5E-01
			Selenium	2/11	1.08E-02 (UCL)	5.0E-03	7.2E-04	1.4E-01
			Silver	4/11	9.36E-03 (UCL)	5.0E-03	6.2E-04	1.2E-01
			Barium	3/4	1.18E-01 (Max)	7.0E-02	7.9E-03	1.1E-01
			Mercury	5/10	5.03E-04 (UCL)	3.0E-04	3.4E-05	1.1E-01
			Beryllium	2/10	3.20E-03 (UCL)	5.0E-03	2.1E-04	4.3E-02
			Copper	6/10	1.54E-02 (UCL)	3.7E-02	1.0E-03	2.8E-02
			Zinc	7/11	2.96E-02 (UCL)	2.0E-01	2.0E-03	9.9E-03
			Tin	1/3	1.48E-02 (Max)	6.0E-01	9.9E-04	1.6E-03
			TOTAL					1.1E+01
		Organics	Trans-1,3-dichloropropene	1/5	5.01E-03 (UCL)	3.0E-04	3.3E-04	1.1E+00
			2,4-Dichlorophenol	1/8	1.01E-02 (UCL)	3.0E-03	6.7E-04	2.2E-01
			Trans-1,2-dichloroethylene	3/5	2.23E-02 (UCL)	2.0E-02	1.5E-03	7.4E-02
			1,1-Dichloroethylene	2/5	5.01E-03 (Max)	9.0E-03	3.3E-04	3.7E-02
			Bis(2-ethylhexyl)phthalate	2/8	1.03E-02 (UCL)	2.0E-02	6.9E-04	3.4E-02
			Chloroform	3/5	5.10E-03 (UCL)	1.0E-02	3.4E-04	3.4E-02
			Tetrachloroethylene	3/5	5.03E-03 (UCL)	1.0E-02	3.4E-04	3.4E-02
			Pyrene	1/8	1.01E-02 (UCL)	3.0E-02	6.7E-04	2.2E-02
			Fluoranthene	1/8	1.01E-02 (UCL)	4.0E-02	6.7E-04	1.7E-02
			Bromoform	1/5	5.03E-03 (UCL)	2.0E-02	3.4E-04	1.7E-02
			Dichlorobromomethane	2/5	5.00E-03 (Max)	2.0E-02	3.3E-04	1.7E-02

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index		
3	Water Ingestion	Organics	Chlorobenzene	1/5	5.00E-03 (Max)	2.0E-02	3.3E-04	1.7E-02		
			Chlorodibromomethane	1/5	5.00E-03 (Max)	2.0E-02	3.3E-04	1.7E-02		
			Di-n-butylphthalate	1/8	1.00E-02 (Max)	1.0E-01	6.7E-04	6.7E-03		
			Methylene chloride	3/5	5.53E-03 (UCL)	6.0E-02	3.7E-04	6.1E-03		
			1,1,1-Trichloroethane	2/5	5.02E-03 (Max)	9.0E-02	3.3E-04	3.7E-03		
			Butylbenzylphthalate	1/8	1.00E-02 (UCL)	2.0E-01	6.7E-04	3.3E-03		
			1,1-Dichloroethane	2/5	5.00E-03 (Max)	1.0E-01	3.3E-04	3.3E-03		
			Ethylbenzene	1/5	5.00E-03 (Max)	1.0E-01	3.3E-04	3.3E-03		
			Anthracene	1/8	1.01E-02 (UCL)	3.0E-01	6.7E-04	2.2E-03		
			Toluene	2/5	5.00E-03 (Max)	2.0E-01	3.3E-04	1.7E-03		
			Phenol	4/8	1.24E-02 (UCL)	6.0E-01	1.4E-03	1.4E-03		
			Diethylphthalate	1/8	1.01E-02 (UCL)	8.0E-01	6.7E-04	8.4E-04		
			Dimethylphthalate	1/8	1.01E-02 (UCL)	1.0E+00	6.7E-04	6.7E-04		
TOTAL						1.7E+00				
3	Water Ingestion						1.3E+01			
3							1.9E+01			
4	Fish Ingestion	Inorganics	Mercury	27/27	3.91E-01 (UCL)	3.0E-04	6.2E-04	2.1E+00		
			Arsenic	25/27	1.07E-01 (UCL)	3.0E-04	1.7E-04	5.6E-01		
			Selenium	14/27	6.86E-01 (UCL)	5.0E-03	1.1E-03	2.2E-01		
			Zinc	27/27	5.55E+00 (UCL)	2.0E-01	1.0E-02	5.2E-02		
			TOTAL						2.9E+00	
			Organics	Chlordane	18/18	9.72E-02 (UCL)	6.0E-05	1.5E-04	2.6E+00	
				4,4'-DDT	1/18	2.00E-01 (Max)	5.0E-04	3.2E-04	6.3E-01	
				Aldrin	2/14	1.33E-03 (UCL)	3.0E-05	2.1E-06	7.0E-02	
				TOTAL						3.3E+00

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
4	Fish Ingestion				mg/L	1/(mg/kg/d)	mg/kg/d	6.1E+00
4	Water Ingestion	Inorganics	Cyanide	1/2	9.59E-02 (Max)	2.0E-02	6.4E-03	3.2E-01
			Chromium	3/5	1.62E-02 (Max)	5.0E-03	1.1E-03	2.2E-01
			Uranium	4/4	3.58E-03 (UCL)	3.0E-03	2.4E-04	8.0E-02
			Nickel	2/4	1.90E-02 (Max)	2.0E-02	1.3E-03	6.3E-02
			Mercury	1/5	2.01E-04 (UCL)	3.0E-04	1.3E-05	4.5E-02
			Manganese	2/2	6.17E-02 (Max)	1.0E-01	4.1E-03	4.1E-02
			Barium	1/1	3.30E-02 (Max)	7.0E-02	2.2E-03	3.1E-02
			Boron	1/1	2.30E-02 (Max)	9.0E-02	1.5E-03	1.7E-02
			Copper	2/4	4.53E-03 (Max)	3.7E-02	3.0E-04	8.2E-03
			Zinc	2/4	8.12E-03 (Max)	2.0E-01	5.4E-04	2.7E-03
			TOTAL					8.2E-01
		Organics	Phenol	1/4	1.29E-02 (Max)	6.0E-01	8.6E-04	1.4E-03
			TOTAL					1.4E-03
4	Water Ingestion				mg/kg	1/(mg/kg/d)	mg/kg/d	8.3E-01
4								7.0E+00
5	Fish Ingestion	Inorganics	Copper	6/25	1.70E+02 (Max)	3.7E-02	2.7E-01	7.2E+00
			Mercury	25/25	1.08E-01 (UCL)	3.0E-04	1.7E-04	5.7E-01
			Arsenic	23/25	9.23E-02 (UCL)	2.0E-04	1.5E-04	4.8E-01
			Zinc	24/24	1.03E+01 (UCL)	2.0E-01	1.7E-02	8.6E-02
			Beryllium	2/25	2.70E-02 (Max)	5.0E-03	4.3E-05	8.5E-03

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

TOTAL					mg/kg	1/(mg/kg/d)	mg/kg/d	8.4E+00
5	Fish Ingestion	Organics	Chlordane	18/18	4.55E-02 (UCL)	6.0E-05	7.2E-05	1.2E+00
			Aldrin	2/18	5.45E-03 (UCL)	3.0E-05	8.6E-06	2.9E-01
			4,4'-DDT	5/18	1.29E-03 (UCL)	5.0E-04	2.0E-06	4.1E-03
			TOTAL					1.5E+00

5	Fish Ingestion							9.9E+00
=====								
5								9.9E+00
=====								
10	Fish Ingestion	Inorganics	Mercury	15/15	2.80E-01 (UCL)	3.0E-04	4.4E-04	1.5E+00
			Arsenic	14/15	9.68E-02 (UCL)	3.0E-04	1.5E-04	5.1E-01
			Selenium	8/15	5.82E-01 (UCL)	5.0E-03	9.2E-04	1.8E-01
			Zinc	15/15	5.97E+00 (UCL)	2.0E-01	9.4E-03	4.7E-02
			Beryllium	1/10	3.70E-02 (Max)	5.0E-03	5.8E-05	1.2E-02
			TOTAL					2.2E+00

		Organics	Chlordane	8/8	1.57E-02 (UCL)	6.0E-05	2.5E-05	4.1E-01
			TOTAL					4.1E-01

10	Fish Ingestion							2.6E+00

	Water Ingestion	Organics	Bis(2-ethylhexyl)phthalate	1/1	7.50E-03 (Max)	2.0E-02	5.0E-04	2.5E-02

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of detects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index

10	Water Ingestion		TOTAL					2.5E-02

10								2.5E-02

13	Fish Ingestion	Inorganics	Mercury	10/10	3.87E-01 (UCL)	3.0E-04	5.8E-04	1.9E+00
			Arsenic	4/10	5.90E-02 (UCL)	3.0E-04	9.3E-05	3.1E-01
			Zinc	10/10	6.98E+00 (UCL)	2.0E-01	1.1E-02	5.5E-02
			TOTAL					2.3E+00

		Organics	Chlordane	2/4	7.29E-02 (Max)	6.0E-05	1.1E-04	1.9E+00
			Aldrin	2/3	1.30E-03 (UCL)	3.0E-05	2.1E-06	6.8E-02
			TOTAL					2.0E+00

13	Fish Ingestion							4.3E+00

	Water Ingestion	Inorganics	Cyanide	1/1	8.79E-02 (Max)	2.0E-02	5.9E-03	2.9E-01
			Chromium	1/2	1.10E-02 (Max)	5.0E-03	7.3E-04	1.5E-01
			Manganese	1/1	1.54E-01 (Max)	1.0E-01	1.0E-02	1.0E-01
			Nickel	1/2	1.90E-02 (Max)	2.0E-02	1.3E-03	6.3E-02
			Zinc	1/2	1.80E-02 (Max)	2.0E-01	1.2E-03	6.0E-03
			TOTAL					6.1E-01

Table E.12 Results of child conservative screening for noncarcinogens in fish, surface sediment, and water where at least one measurement in a reach was above detection limits.

Reach	Media	Contaminant type	Contaminant	Ratio of defects to total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
13	Water Ingestion							6.1E-01
13								4.9E+00

Table E.13 Results of adult conservative screening for carcinogens in dredged sediments
where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Sediment exposure	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
1	Inorganics	Arsenic	1.2E-05	7.9E-07		1.4E-08	2.1E-07	2.1E-05	3.3E-05
		Beryllium	2.1E-06	9.9E-09		1.8E-08	1.3E-05	1.5E-05	3.0E-05
		Chromium		1.2E-06					1.2E-06
		Nickel		2.9E-08					2.9E-08
		Cadmium		6.2E-09					6.2E-09
	Organics	TOTAL	1.4E-05	2.0E-06		3.2E-08	1.4E-05	3.6E-05	6.5E-05
		Pyrene	3.7E-06			1.5E-05	1.4E-05	9.3E-05	1.3E-04
		Fluoranthene	3.7E-06			1.6E-05	1.5E-05	9.1E-05	1.3E-04
		Benzo(k)fluoranthene	3.6E-07			6.0E-05	5.7E-05	6.9E-07	1.2E-04
		Benzo(a)anthracene	3.6E-07			5.0E-06	4.7E-06	3.4E-06	1.4E-05
		4,4'-DDT	8.1E-08	1.9E-10		5.1E-06	4.9E-06	2.8E-07	1.0E-05
		4,4'-DDD	5.7E-08			1.7E-06	1.7E-06	3.2E-07	3.8E-06
		Chrysene	3.6E-08			6.8E-07	8.4E-07	2.3E-07	2.0E-06
		TOTAL	8.3E-06	1.9E-10		1.0E-04	9.9E-05	1.9E-04	4.0E-04
2	Radionuclides	Cs-137	3.2E-08	3.8E-11	9.0E-06	1.8E-06	2.0E-06	1.2E-06	1.4E-05
		Co-60	5.2E-09	3.3E-11	4.7E-06	8.5E-08	3.8E-07	7.3E-08	4.8E-06
		TOTAL	3.7E-08	7.1E-11	1.3E-05	1.9E-06	2.4E-06	1.3E-06	1.9E-05
	Inorganics	Beryllium	2.1E-06	9.7E-09		1.7E-08	1.3E-05	1.5E-05	3.0E-05
		Arsenic	6.5E-06	4.4E-07		7.9E-09	1.2E-07	1.2E-05	1.9E-05
		Chromium		9.7E-07					9.7E-07
		Nickel		1.9E-08					1.9E-08
		Cadmium		4.2E-09					4.2E-09
	Organics	TOTAL	8.6E-06	1.4E-06		2.5E-08	1.3E-05	2.8E-05	4.9E-05
		Aroclor-1260	1.7E-05			5.6E-03	5.3E-03	2.2E-05	1.1E-02
		Aroclor-1254	1.7E-05			5.3E-04	5.1E-04	9.5E-05	1.1E-03
		Toxaphene	2.4E-06			1.4E-06	1.3E-06	5.1E-04	5.1E-04
		Benzo(a)pyrene	2.2E-06	5.8E-09		7.1E-05	6.7E-05	1.2E-05	1.5E-04
		Phenanthrene	2.2E-06	4.5E-09		5.7E-06	5.4E-06	8.3E-05	9.6E-05
		Pyrene	1.9E-06			7.9E-06	7.5E-06	4.9E-05	6.6E-05
		Fluoranthene	1.9E-06			8.0E-06	7.6E-06	4.7E-05	6.4E-05
		Indeno(1,2,3-cd)pyrene	2.2E-07			2.2E-05	2.1E-05	6.0E-07	4.3E-05

Table E.13 Results of adult conservative screening for carcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Sediment exposure	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
2	Organics	Benzo(b)fluoranthene	2.2E-07			8.4E-06	8.0E-06	1.1E-06	1.8E-05
		Benzo(a)anthracene	2.2E-07			3.1E-06	3.0E-06	2.2E-06	8.5E-06
		Chrysene	2.2E-08			5.5E-07	5.2E-07	1.4E-07	1.2E-06
		Bis(2-ethylhexyl)phthalate	5.1E-08			2.1E-08	2.0E-08	1.3E-07	1.7E-07
		TOTAL	4.6E-05	1.0E-08		6.3E-03	5.9E-03	8.2E-04	1.3E-02
2	Radionuclides	Cs-137	8.6E-07	1.0E-09	2.4E-04	4.9E-05	5.5E-05	3.3E-05	3.8E-04
		Sr-90	1.4E-08	4.3E-11		2.9E-06	8.6E-07	5.3E-06	9.1E-06
		Co-60	5.8E-09	3.7E-11	4.7E-06	9.5E-08	4.3E-07	8.1E-08	5.3E-06
		U-238	3.8E-08	1.7E-07	2.6E-07	7.9E-08	3.9E-07	1.3E-07	1.1E-06
		U-234	2.4E-08	9.2E-08	2.4E-10	4.9E-08	2.5E-07	8.3E-08	4.9E-07
		Am-241	4.9E-08	1.5E-08	5.2E-08	5.9E-11	8.8E-10	8.6E-08	1.6E-07
		U-235	1.1E-09	4.1E-09	8.8E-08	2.3E-09	1.1E-08	3.8E-09	1.1E-07
		Pu-239, 240	4.6E-08	1.8E-08	1.8E-11	1.3E-11	3.9E-10	4.2E-08	1.1E-07
		Pu-238	7.1E-09	2.8E-09	4.4E-12	1.8E-12	5.4E-11	5.8E-09	1.6E-08
		Ca-243, 244	2.7E-09	5.4E-10	1.7E-12	2.0E-12	3.0E-11	2.9E-09	6.2E-09
3	Inorganics	TOTAL	1.0E-06	3.0E-07	2.5E-04	5.2E-05	5.7E-05	3.9E-05	4.0E-04
3	Organics	Arsenic	1.5E-04	1.0E-05		1.8E-07	2.7E-06	2.7E-04	4.3E-04
		Beryllium	4.1E-06	1.9E-08		3.4E-08	2.6E-05	2.9E-05	5.9E-05
		Chromium		1.8E-06					1.8E-06
		Nickel		2.8E-08					2.9E-08
		Cadmium		1.2E-08					1.2E-08
		TOTAL	1.5E-04	1.2E-05		2.2E-07	2.8E-05	3.0E-04	4.9E-04
3	Organics	Aroclor-1254	1.0E-05			3.3E-04	3.1E-04	5.8E-05	7.0E-04
		Benzo(e)pyrene	2.1E-06			6.7E-05	6.3E-05	1.1E-05	1.4E-04
		Anthracene	2.1E-06	4.2E-09		4.6E-06	4.4E-06	9.4E-05	1.0E-04
		Phenanthrene	1.9E-06			4.9E-06	4.6E-06	7.1E-05	8.3E-05
		Benzo(k)fluoranthene	2.1E-07			3.6E-05	3.4E-05	4.0E-07	7.0E-05
		Fluoranthene	1.9E-06			8.1E-06	7.6E-06	4.7E-05	6.5E-05
		Pyrene	1.9E-06			7.6E-06	7.2E-06	4.7E-05	6.4E-05
		Benzo(b)fluoranthene	2.0E-07			7.6E-06	7.2E-06	9.9E-07	1.6E-05
		Benzo(a)anthracene	2.1E-07			2.9E-06	2.8E-06	2.0E-06	8.0E-06
		Chrysene	2.1E-08			5.2E-07	4.9E-07	1.4E-07	1.2E-06

Table E.13 Results of adult conservative screening for carcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Sediment exposure	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
3	Organics	Bis(2-ethylhexyl)phthalate	8.4E-08			3.5E-08	3.3E-08	2.1E-07	2.9E-07
		TOTAL	2.1E-05	4.2E-09		4.6E-04	4.4E-04	3.3E-04	1.3E-03
	Radionuclides	Cs-137	2.5E-07	2.9E-10	6.9E-05	1.4E-05	1.6E-05	9.4E-06	1.1E-04
		U-238	1.6E-07	7.0E-07	1.1E-06	3.3E-07	1.7E-06	5.6E-07	4.5E-06
		U-234	1.0E-07	3.9E-07	1.0E-09	2.1E-07	1.0E-06	3.5E-07	2.1E-06
		Sr-90	3.0E-09	6.0E-12		6.0E-07	1.8E-07	1.1E-06	1.9E-06
		Co-60	1.5E-09	9.8E-12	1.2E-06	2.5E-08	1.1E-07	2.1E-08	1.4E-06
		U-235	3.3E-08	1.3E-08	2.7E-07	7.0E-09	3.5E-08	1.2E-08	3.4E-07
		Am-241	2.8E-08	8.8E-09	3.0E-09	3.3E-11	5.0E-10	4.9E-08	8.9E-08
		Pu-238, 240	3.9E-09	1.5E-09	1.5E-12	1.1E-12	3.3E-11	3.5E-09	8.9E-09
		Pu-238	3.5E-09	1.4E-09	2.2E-12	8.9E-13	2.7E-11	2.9E-09	7.8E-09
		Cm-243, 244	2.2E-09	4.3E-10	1.3E-12	1.6E-12	2.4E-11	2.3E-09	5.0E-09
		TOTAL	5.5E-07	1.1E-08	7.2E-05	1.5E-05	1.9E-05	1.1E-05	1.2E-04
			=====	=====	=====	=====	=====	=====	=====
			1.8E-04	1.3E-05	7.2E-05	4.8E-04	4.9E-04	6.4E-04	1.9E-03
4	Inorganics	Beryllium	3.1E-08	1.4E-08		2.6E-08	1.9E-05	2.1E-05	4.4E-05
		Arsenic	1.2E-05	8.0E-07		1.4E-08	2.1E-07		3.4E-05
		Chromium		1.6E-06					1.6E-06
		Nickel		3.9E-08					3.6E-08
		Cadmium		1.3E-08					1.3E-08
		TOTAL	1.5E-05	2.4E-06		4.0E-08	1.9E-05	4.2E-05	7.9E-05
	Organics	N-Nitroso-di-n-propylamine	1.5E-05			1.2E-06	1.2E-06	4.2E-02	4.2E-02
		Phenanthrene	1.6E-05			4.2E-05	4.0E-05	6.2E-04	7.1E-04
		Pyrene	1.6E-05			6.6E-05	6.3E-05	4.1E-04	5.5E-04
		Fluoranthene	1.6E-05			6.8E-05	6.5E-05	4.0E-04	5.5E-04
		Aroclor-1254	8.3E-07			2.6E-05	2.5E-05	4.6E-06	5.6E-05
		Bis(2-ethylhexyl)phthalate	1.7E-08			7.0E-08	6.6E-08	4.2E-07	5.7E-07
		TOTAL	6.5E-05			2.0E-04	1.9E-04	4.4E-02	4.4E-02
	Radionuclides	Cs-137	1.4E-06	1.6E-09	3.9E-04	7.9E-05	8.9E-05	5.3E-05	6.2E-04
		Sr-90	1.6E-08	4.8E-11		3.2E-06	9.6E-07	5.9E-06	1.0E-05
		Co-60	9.7E-09	6.2E-11	7.9E-06	1.6E-07	7.1E-07	1.4E-07	8.9E-06
		U-238	5.8E-08	2.5E-07	4.0E-07	1.2E-07	6.0E-07	2.0E-07	1.6E-06
		U-234	3.9E-03	1.5E-07	3.9E-10	8.0E-08	4.0E-07	1.4E-07	8.1E-07

Table E.13 Results of adult conservative screening for carcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Sediment exposure	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
4	Radionuclides	U-235	2.1E-09	8.0E-09	1.7E-07	4.4E-09	2.2E-08	7.4E-09	2.2E-07
		Am-241	5.2E-08	1.8E-08	5.5E-08	6.2E-11	9.4E-10	9.1E-08	1.7E-07
		Pu-239,240	6.4E-08	2.5E-08	2.6E-11	1.8E-11	5.5E-10	5.9E-08	1.5E-07
		Pu-238	5.3E-09	2.1E-09	3.3E-12	1.3E-12	4.0E-11	4.3E-09	1.2E-08
		Cm-243,244	3.9E-09	7.7E-10	2.4E-12	2.9E-12	4.3E-11	4.2E-09	8.9E-09
		TOTAL	1.6E-08	4.8E-07	4.0E-04	8.3E-05	9.2E-05	6.0E-05	6.4E-04
4	Inorganics	Beryllium	3.9E-06	1.8E-08	4.0E-04	2.9E-04	3.0E-04	4.4E-02	4.5E-02
		Arsenic	9.7E-06	6.6E-07	4.0E-04	2.9E-04	3.0E-04	4.4E-02	4.5E-02
		Chromium		2.1E-08					
		Cadmium		3.8E-08					
		Nickel	1.4E-05	3.4E-08					
		TOTAL		2.9E-06					
5	Organics	Bis(2-ethylhexyl)phthalate	2.0E-08						
		TOTAL	2.0E-08						
	Radionuclides	Cs-137	3.7E-07	4.3E-10	1.0E-04	2.1E-05	2.3E-05	1.4E-05	1.6E-04
		Co-60	4.6E-09	2.9E-11	3.7E-06	7.4E-08	3.4E-07	6.3E-08	4.2E-06
		Sr-90	5.6E-09	1.7E-11		1.1E-08	3.4E-07	2.1E-06	3.6E-06
		U-238	5.2E-08	2.3E-07	3.6E-07	1.1E-07	5.4E-07	1.8E-07	1.5E-06
		U-234	3.3E-08	1.3E-07	3.4E-10	6.9E-08	3.5E-07	1.2E-07	7.0E-07
		U-235	1.6E-09	5.9E-09	1.3E-07	3.2E-09	1.6E-08	5.5E-09	1.6E-07
10	Inorganics	Am-241	2.7E-08	8.2E-09	2.8E-09	3.2E-11	4.8E-10	4.7E-08	8.5E-08
		Pu-239,240	2.4E-08	9.3E-09	8.7E-12	6.9E-12	2.1E-10	2.2E-08	5.6E-08
		Pu-238	7.5E-09	3.0E-09	4.6E-12	1.9E-12	5.7E-11	6.1E-09	1.7E-08
		Ca-243,244	3.1E-09	6.2E-10	1.9E-12	2.3E-12	3.5E-11	3.4E-09	7.2E-09
		TOTAL	5.3E-07	3.9E-07	1.1E-04	2.2E-05	2.5E-05	1.7E-05	1.7E-04
5	Inorganics	Beryllium	3.4E-06	1.6E-08	1.1E-04	2.2E-05	4.9E-05	6.1E-05	2.6E-04
		Arsenic	5.5E-06	3.8E-07					

Table E.13 Results of adult conservative screening for carcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Sediment exposure	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
10	Inorganics	Chromium		1.3E-06					1.3E-06
		Nickel		2.6E-08					2.6E-08
		Cadmium		5.0E-09					5.0E-09
		TOTAL	8.9E-06	1.8E-06		3.5E-08	2.1E-05	3.4E-05	8.6E-05
	Organics	Anthracene	1.9E-05			4.2E-05	4.0E-05	8.5E-04	9.5E-04
		Phenanthrene	1.8E-05			5.0E-05	4.7E-05	7.2E-04	8.4E-04
		Pyrene	1.9E-05			7.8E-05	7.4E-05	4.8E-04	6.5E-04
		Fluoranthene	1.9E-05			8.0E-05	7.6E-05	4.7E-04	6.4E-04
		Chrysene	1.8E-07			4.6E-08	4.4E-08	1.2E-06	1.0E-05
		Bis(2-ethylhexyl)phthalate	2.8E-08			1.2E-07	1.1E-07	7.1E-07	9.8E-07
		TOTAL	7.6E-05			2.5E-04	2.4E-04	2.5E-03	3.1E-03
10	Radionuclides	Cs-137	2.7E-08	3.2E-11	7.7E-06	1.6E-06	1.8E-08	1.0E-06	1.2E-05
		Sr-90	4.6E-09	1.4E-11		9.3E-07	2.8E-07	1.7E-06	2.9E-06
		U-238	4.1E-08	1.6E-07	2.8E-07	8.5E-08	4.2E-07	1.4E-07	1.1E-06
		U-234	2.1E-08	8.1E-08	2.1E-10	4.3E-08	2.2E-07	7.3E-08	4.4E-07
		U-235	1.0E-08	3.9E-09	8.5E-08	2.2E-09	1.1E-08	3.9E-09	1.1E-07
		As-241	1.4E-08	4.4E-09	1.5E-09	1.7E-11	2.6E-10	2.5E-08	4.5E-08
		Pu-238, 240	7.3E-09	2.8E-09	2.9E-12	2.1E-12	6.2E-11	6.7E-09	1.7E-08
		Cs-243, 244	2.4E-09	4.7E-10	1.5E-12	1.7E-12	2.6E-11	2.5E-09	5.4E-09
		Pu-238	2.4E-09	9.5E-10	1.5E-12	6.1E-13	1.8E-11	2.0E-09	5.3E-09
		TOTAL	1.2E-07	2.7E-07	8.1E-06	2.6E-06	2.7E-06	3.0E-06	1.7E-05
10			8.5E-05	2.0E-06	8.1E-06	2.6E-04	2.7E-04	2.6E-03	3.2E-03

Table E.14 Results of adult conservative screening for noncarcinogens in dredged sediments
where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
1	Inorganics	Zinc	8.7E-04		7.2E-01	2.2E+00	6.9E-01	3.6E+00
		Cadmium	2.0E-03		3.3E-01	5.0E-02	2.0E+00	2.4E+00
		Mercury	5.8E-04	4.9E-08	2.4E-02	1.5E-01	3.5E-01	5.2E-01
		Chromium	5.6E-03		7.1E-02	3.2E-01	1.1E-01	5.1E-01
		Copper	1.1E-03		1.5E-01	2.3E-01	1.1E-01	4.9E-01
		Selenium	2.0E-04		8.8E-02	2.6E-01	4.0E-02	3.8E-01
		Nickel	1.7E-03		3.2E-02	1.9E-02	1.7E-01	2.2E-01
		Arsenic	5.2E-02		7.2E-05	1.1E-03	1.1E-01	1.6E-01
		Silver	9.4E-05		7.8E-02	1.2E-02	3.7E-02	1.3E-01
		Beryllium	2.3E-04		2.2E-06	1.7E-03	1.9E-03	3.8E-03
		TOTAL	6.4E-02	4.9E-08	1.5E+00	3.2E+00	3.8E+00	8.4E+00
		4,4'-DDT	1.1E-03		8.1E-02	7.7E-02	4.5E-03	1.6E-01
		Pyrene	3.8E-05		1.9E-04	1.8E-04	1.1E-03	1.5E-03
		Fluoranthene	3.0E-05		1.4E-04	1.4E-04	8.4E-04	1.2E-03
		TOTAL	1.2E-03		8.2E-02	7.8E-02	6.5E-03	1.7E-01
2	Inorganics	Zinc	5.1E-04		4.2E-01	1.3E+00	4.0E-01	2.1E+00
		Cadmium	1.4E-03		2.2E-01	3.4E-02	1.3E+00	1.6E+00
		Mercury	1.2E-03	9.6E-08	4.8E-02	2.9E-01	6.9E-01	1.0E+00
		Chromium	4.8E-03		5.9E-02	2.7E-01	9.3E-02	4.2E-01
		Copper	5.6E-04		7.5E-02	1.1E-01	5.6E-02	2.4E-01
		Selenium	9.7E-05		4.2E-02	1.2E-01	1.9E-02	1.9E-01
		Nickel	1.1E-03		2.1E-02	1.3E-02	1.1E-01	1.5E-01
		Arsenic	2.9E-02		4.1E-05	6.1E-04	5.9E-02	8.9E-02
		Silver	3.9E-05		3.2E-02	4.8E-03	1.5E-02	5.2E-02
		Uranium	2.1E-03		4.9E-03	2.5E-02	8.3E-03	4.0E-02
		Beryllium	2.3E-04		2.2E-06	1.6E-03	1.8E-03	3.7E-03
		TOTAL	4.1E-02	9.6E-08	9.3E-01	2.1E+00	2.8E+00	5.9E+00
		Naphthalene	1.8E-05		1.6E-05	1.5E-05	2.9E-03	2.9E-03
		Bis(2-ethylhexyl)phthalate	4.2E-05		2.0E-04	1.9E-04	1.2E-03	1.7E-03
		Pyrene	2.0E-05		9.7E-05	9.2E-05	6.0E-04	8.1E-04
		Diethylphthalate	8.9E-07		2.5E-07	2.4E-07	6.4E-04	6.5E-04
		Fluoranthene	1.5E-05		7.4E-05	7.0E-05	4.3E-04	5.9E-04

Table E.14 Results of adult conservative screening for noncarcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
2	Organics	Di-n-butylphthalate	7.0E-06		3.4E-05	3.2E-05	2.0E-04	2.7E-04
		TOTAL	1.0E-04		4.2E-04	4.0E-04	6.0E-03	6.9E-03
2			4.1E-02	9.8E-06	9.3E-01	2.1E+00	2.8E+00	5.9E+00
3	Inorganics	Thallium	3.0E-02		4.6E+00	9.2E+00	1.2E+01	2.8E+01
		Cadmium	3.9E-03		6.3E-01	9.5E-02	3.8E+00	4.5E+00
		Mercury	2.3E-03	1.9E-05	9.6E-02	5.7E-01	1.4E+00	2.1E+00
		Arsenic	6.7E-01		9.4E-04	1.4E-02	1.4E+00	2.0E+00
		Selenium	8.9E-04		3.8E-01	1.1E+00	1.8E-01	1.7E+00
		Zinc	4.0E-04		3.3E-01	9.9E-01	3.2E-01	1.6E+00
		Chromium	8.5E-03		1.1E-01	4.9E-01	1.7E-01	7.7E-01
		Copper	1.3E-03		1.8E-01	2.7E-01	1.3E-01	5.8E-01
		Silver	2.2E-04		1.8E-01	2.7E-02	8.7E-02	3.0E-01
		Nickel	1.7E-03		3.3E-02	2.0E-02	1.7E-01	2.2E-01
		Uranium	8.7E-03		2.1E-02	1.0E-01	3.5E-02	1.7E-01
		Antimony	1.1E-03		8.1E-04	2.4E-03	2.3E-02	2.7E-02
		Beryllium	4.5E-04		4.3E-06	3.2E-03	3.6E-03	7.3E-03
		TOTAL	7.3E-01	1.9E-05	6.5E+00	1.3E+01	2.0E+01	4.0E+01
	Organics	Bis(2-ethylhexyl)phthalate	7.0E-05		3.4E-04	3.2E-04	2.0E-03	2.7E-03
		Naphthalene	1.6E-05		1.5E-05	1.4E-05	2.6E-03	2.7E-03
		Dimethylphthalate	6.9E-07		1.3E-07	1.3E-07	7.9E-04	7.9E-04
		Pyrene	2.0E-05		9.4E-05	8.9E-05	5.8E-04	7.8E-04
		Fluoranthene	1.5E-05		7.4E-05	7.0E-05	4.3E-04	5.9E-04
		Di-n-butylphthalate	6.6E-06		3.2E-05	3.1E-05	1.9E-04	2.6E-04
		Anthracene	2.2E-06		5.7E-06	5.4E-06	1.1E-04	1.3E-04
		TOTAL	1.3E-04		5.6E-04	5.3E-04	6.7E-03	8.0E-03
3			7.3E-01	1.9E-05	6.5E+00	1.3E+01	2.0E+01	4.0E+01
4	Inorganics	Mercury	7.6E-02	6.3E-04	3.1E+00	1.9E-01	4.5E+01	6.7E+01
		Cadmium	4.0E-03		6.7E-01	1.0E-01	4.0E+00	4.8E+00
		Zinc	8.9E-04		7.3E-01	2.2E+00	7.1E-01	3.6E+00
		Silver	5.1E-04		4.3E-01	6.4E-02	2.0E-01	6.9E-01

Table E.14 Results of adult conservative screening for noncarcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total
4	Inorganics	Chromium	7.6E-03		9.7E-02	4.4E-01	1.5E-01	6.9E-01
		Selenium	3.5E-04		1.5E-01	4.5E-01	7.0E-02	6.7E-01
		Copper	1.3E-03		1.8E-01	2.7E-01	1.3E-01	5.9E-01
		Nickel	2.1E-03		4.0E-02	2.4E-02	2.1E-01	2.7E-01
		Arsenic	5.2E-02		7.3E-05	1.1E-03	1.1E-01	1.6E-01
		Uranium	3.2E-03		7.6E-03	3.8E-02	1.3E-02	6.1E-02
		Antimony	1.5E-03		1.1E-03	3.2E-03	2.9E-02	3.5E-02
		Beryllium	3.3E-04		3.2E-06	2.4E-03	2.7E-03	5.4E-03
		TOTAL	1.5E-01	6.3E-04	5.4E+00	2.2E+01	5.1E+01	7.9E+01
		Pyrene	1.7E-04		8.1E-04	7.7E-04	5.0E-03	6.8E-03
		Bis(2-ethylhexyl)phthalate	1.4E-04		6.7E-04	6.3E-04	4.0E-03	5.4E-03
		Fluoranthene	1.3E-04		6.3E-04	6.0E-04	3.7E-03	5.0E-03
		TOTAL	4.4E-04		2.1E-03	2.0E-03	1.3E-02	1.7E-02
			1.5E-01	6.3E-04	5.4E+00	2.2E+01	5.1E+01	7.9E+01
5	Inorganics	Cadmium	1.2E-02		2.0E+00	3.0E-01	1.2E+01	1.4E+01
		Mercury	1.4E-02		5.8E-01	3.5E+00	8.4E+00	1.3E+01
		Zinc	1.9E-03	1.2E-04	1.6E+00	4.7E+00	1.5E+00	7.8E+00
		Chromium	1.0E-02		1.3E-01	5.8E-01	2.0E-01	9.2E-01
		Copper	1.5E-03		2.0E-01	3.0E-01	1.5E-01	6.5E-01
		Nickel	2.0E-03		3.8E-02	2.3E-02	2.0E-01	2.6E-01
		Silver	1.8E-04		1.5E-01	2.2E-02	7.0E-02	2.4E-01
		Arsenic	4.3E-02		6.1E-05	9.1E-04	8.9E-02	1.3E-01
		Uranium	2.8E-03		6.8E-03	3.4E-02	1.1E-02	5.5E-02
		Antimony	1.7E-03		1.2E-03	3.7E-03	3.4E-02	4.1E-02
		Beryllium	4.1E-04		3.9E-06	2.9E-03	3.3E-03	6.6E-03
		TOTAL	9.0E-02	1.2E-04	4.7E+00	9.5E+00	2.3E+01	3.7E+01
		Bis(2-ethylhexyl)phthalate	1.8E-04		7.9E-04	7.5E-04	4.7E-03	6.4E-03
		TOTAL	1.6E-04		7.9E-04	7.5E-04	4.7E-03	6.4E-03
			9.0E-02	1.2E-04	4.7E+00	9.5E+00	2.3E+01	3.7E+01

Table E.14 Results of adult conservative screening for noncarcinogens in dredged sediments where at least one measurement in a reach was above detection limits.

Resch	Contaminant type	Contaminant	Sediment ingestion	Sediment inhalation	Milk ingestion	Meat ingestion	Vegetable ingestion	Multiple pathway total	
10	Inorganics	Zinc	6.3E-04		5.2E-01	1.6E+00	5.0E-01	2.6E+00	
		Cadmium	1.6E-03		2.6E-01	3.9E-02	1.6E+00	1.9E+00	
		Chromium	6.3E-03		8.1E-02	3.6E-01	1.3E-01	5.8E-01	
		Mercury	3.9E-04	3.2E-06	1.6E-02	9.6E-02	2.3E-01	3.4E-01	
		Copper	7.7E-04		1.0E-01	1.5E-01	7.7E-02	3.4E-01	
		Nickel	1.5E-03		2.9E-02	1.7E-02	1.5E-01	2.0E-01	
		Arsenic	2.4E-02		3.4E-05	5.2E-04	5.0E-02	7.5E-02	
		Uranium	2.1E-03		5.1E-03	2.6E-02	8.7E-03	4.2E-02	
		Antimony	1.5E-03		1.1E-03	3.2E-03	3.0E-02	3.5E-02	
		Beryllium	3.7E-04		3.5E-06	2.7E-03	3.0E-03	6.0E-03	
		TOTAL	4.0E-02	3.2E-06	1.0E+00	2.3E+00	2.8E+00	6.1E+00	
		Organics	Bis(2-ethylhexyl)phthalate	2.4E-04		1.1E-03	1.1E-03	6.9E-03	9.3E-03
			Pyrene	2.0E-04		9.5E-04	9.0E-04	5.9E-03	7.9E-03
Fluoranthene	1.5E-04			7.4E-04	7.0E-04	4.3E-03	5.9E-03		
Anthracene	2.0E-05			5.2E-05	4.9E-05	1.0E-03	1.2E-03		
TOTAL	6.1E-04			2.9E-03	2.7E-03	1.8E-02	2.4E-02		
10			4.0E-02	3.2E-06	1.0E+00	2.3E+00	2.8E+00	6.1E+00	

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
				mg/kg			mg/L	
1	Organics	1,4-Dichlorobenzene	6	6.80E-01 (Min)	7.6E-07	3	1.00E-02 (Min)	2.9E-07
		2,4,6-Trichlorophenol	6	6.80E-01 (Min)	3.5E-07	3	1.00E-02 (Min)	1.3E-07
		2,4-Dinitrotoluene	6	6.80E-01 (Min)	2.1E-05	3	1.00E-02 (Min)	8.3E-06
		2,6-Dinitrotoluene	6	6.80E-01 (Min)	2.1E-05	3	1.00E-02 (Min)	8.3E-06
		3,3'-Dichlorobenzidine	6	1.40E+00 (Min)	2.9E-05	3	2.00E-02 (Min)	1.1E-05
		4,4'-DDD				2	1.00E-04 (Min)	2.9E-08
		4,4'-DDE				2	1.00E-04 (Min)	4.2E-08
		4,4'-DDT				2	1.00E-04 (Min)	4.2E-08
		4-Chloroaniline	18	1.00E-02 (Min)	1.6E-07	2	1.00E-04 (Min)	4.3E-07
		Acenaphthene	6	6.80E-01 (Min)	1.1E-06	3	1.00E-02 (Min)	1.4E-04
		Acenaphthylene	6	6.80E-01 (Min)	3.6E-04	3	1.00E-02 (Min)	1.4E-04
		Aldrin	18	1.00E-02 (Min)	7.9E-06	2	5.00E-05 (Min)	1.0E-06
		Alpha-BHC				2	5.00E-05 (Min)	3.9E-07
		Anthracene	6	6.80E-01 (Min)	2.3E-04	3	1.00E-02 (Min)	8.9E-05
		Aroclor-1254				2	1.00E-03 (Min)	9.4E-06
		Aroclor-1260				2	1.00E-03 (Min)	9.4E-06
		Benzo(a)anthracene	6	6.80E-01 (Min)	2.3E-05	3	1.00E-02 (Min)	8.9E-06
		Benzo(a)pyrene	6	6.80E-01 (Min)	2.3E-04	3	1.00E-02 (Min)	8.9E-05
		Benzo(b)fluoranthene	6	6.80E-01 (Min)	2.3E-05	3	1.00E-02 (Min)	8.9E-06
		Benzo(g,h,i)perylene	6	6.80E-01 (Min)	2.3E-04	3	1.00E-02 (Min)	8.9E-05
		Benzo(k)fluoranthene	6	6.80E-01 (Min)	2.3E-05	3	1.00E-02 (Min)	8.9E-06
		Beta-BHC				2	5.00E-05 (Min)	1.1E-07
		Bis(2-chloroethyl)ether	6	6.80E-01 (Min)	3.5E-05	3	1.00E-02 (Min)	1.3E-05
		Bis(2-chloroisopropyl)ether	6	6.80E-01 (Min)	2.2E-06	3	1.00E-02 (Min)	8.6E-07
		Bis(2-ethylhexyl)phthalate	6	6.80E-01 (Min)	4.4E-07	3	1.00E-02 (Min)	1.7E-07
		Chlordane				4	5.00E-04 (Min)	8.0E-07
		Chrysene	6	6.80E-01 (Min)	2.3E-06	3	1.00E-02 (Min)	8.9E-07
		Dibenz(a,h)anthracene	6	6.80E-01 (Min)	3.6E-04	3	1.00E-02 (Min)	1.4E-04
		Dieldrin				2	1.00E-04 (Min)	2.0E-06
		Fluoranthene	6	6.80E-01 (Min)	2.3E-04	3	1.00E-02 (Min)	8.9E-05
		Fluorene	6	6.80E-01 (Min)	3.6E-04	3	1.00E-02 (Min)	1.4E-04
		Gamma-BHC (Lindane)				2	5.00E-05 (Min)	8.0E-08
		Heptachlor				2	5.00E-05 (Min)	2.8E-07
		Heptachlor epoxide				2	5.00E-05 (Min)	5.6E-07
		Hexachlorobenzene	6	6.80E-01 (Min)	5.0E-05	3	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene	6	6.80E-01 (Min)	2.5E-06	3	1.00E-02 (Min)	9.6E-07

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
1	Organics	Hexachloroethane	6	6.80E-01 (Min)	4.4E-07	3	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene	6	6.80E-01 (Min)	2.3E-05	3	1.00E-02 (Min)	8.9E-06
		Isophorone	6	6.80E-01 (Min)	1.3E-07	3	1.00E-02 (Min)	5.0E-08
		N-Nitroso-di-n-propylamine	6	6.80E-01 (Min)	2.2E-04	3	1.00E-02 (Min)	8.6E-05
		N-Nitrosodiphenylamine	6	6.80E-01 (Min)	1.5E-07	3	1.00E-02 (Min)	6.0E-08
		Pentachlorophenol	6	3.40E+00 (Min)	1.9E-05	3	5.00E-02 (Min)	7.3E-06
		Phenanthrene	6	6.80E-01 (Min)	2.3E-04	3	1.00E-02 (Min)	8.9E-05
		Pyrene	6	6.80E-01 (Min)	2.3E-04	3	1.00E-02 (Min)	8.9E-05
		Toxaphene	6	6.80E-01 (Min)	2.3E-04	2	1.00E-03 (Min)	1.3E-06
2	Inorganics	Arsenic	13	3.00E-03 (Min)	6.0E-07	3	1.30E-03 (Min)	2.8E-06
		Beryllium				4	2.00E-04 (Min)	1.1E-06
Organics	Organics	1,1,2,2-Tetrachloroethane				3	4.84E-03 (Min)	1.2E-06
		1,1,2-Trichloroethane				3	4.84E-03 (Min)	3.4E-07
		1,1-Dichloroethylene				3	4.84E-03 (Min)	3.6E-06
		1,2-Dichloroethane	8	7.00E-01 (Min)	7.8E-07	3	4.84E-03 (Min)	5.4E-07
		1,4-Dichlorobenzene	8	7.00E-01 (Min)	3.6E-07	3	9.50E-03 (Min)	2.8E-07
		2,4,6-Trichlorophenol	8	7.00E-01 (Min)	3.6E-07	2	1.00E-02 (Min)	1.3E-07
		2,4-Dinitrotoluene	8	7.00E-01 (Min)	2.2E-05	3	1.00E-02 (Min)	8.3E-06
		2,6-Dinitrotoluene	8	7.00E-01 (Min)	2.2E-05	3	1.00E-02 (Min)	8.3E-06
		3,3'-Dichlorobenzidine	8	1.40E+00 (Min)	2.9E-05	3	2.00E-02 (Min)	1.1E-05
		4,4'-DDD	14	1.00E-02 (Min)	1.1E-07	2	1.00E-04 (Min)	2.9E-08
		4,4'-DDE				2	1.00E-04 (Min)	4.2E-08
		4,4'-DDT				2	1.00E-04 (Min)	4.2E-08
		4-Chloroaniline	2	7.00E-01 (Min)	1.1E-06	2	1.00E-02 (Min)	4.3E-07
		Acenaphthene	8	7.00E-01 (Min)	3.7E-04	3	1.00E-02 (Min)	1.4E-04
		Acenaphthylene	8	7.00E-01 (Min)	3.7E-04	3	1.00E-02 (Min)	1.4E-04
		Aldrin				2	5.00E-05 (Min)	1.0E-06
		Alpha-BHC				2	5.00E-05 (Min)	3.9E-07
		Anthracene	8	7.00E-01 (Min)	2.4E-04	3	1.00E-02 (Min)	8.9E-05
		Aroclor-1254				2	1.00E-02 (Min)	9.4E-06
		Aroclor-1260				2	1.00E-03 (Min)	9.4E-06

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
2	Organics	Benzene				3	4.84E-03 (Min)	1.7E-07
		Benztidine				1	2.00E-02 (Min)	5.6E-03
		Benzo(a)anthracene	8	7.00E-01 (Min)	2.4E-05	3	1.00E-02 (Min)	8.9E-06
		Benzo(a)pyrene	8	7.00E-01 (Min)	2.4E-04	3	1.00E-02 (Min)	8.9E-05
		Benzo(b)fluoranthene	8	7.00E-01 (Min)	2.4E-05	3	1.00E-02 (Min)	8.9E-06
		Benzo(g,h,i)perylene	8	7.00E-01 (Min)	2.4E-04	3	1.00E-02 (Min)	8.9E-05
		Benzo(k)fluoranthene	8	7.00E-01 (Min)	2.4E-05	3	1.00E-02 (Min)	8.9E-06
		Beta-BHC				2	5.00E-05 (Min)	1.1E-07
		Bis(2-chloroethyl)ether	8	7.00E-01 (Min)	3.8E-05	3	1.00E-02 (Min)	1.3E-05
		Bis(2-chloroisopropyl)ether	8	7.00E-01 (Min)	2.3E-06	3	1.00E-02 (Min)	8.6E-07
		Bis(2-ethylhexyl)phthalate	8	7.00E-01 (Min)	4.5E-07	3	1.00E-02 (Min)	1.7E-07
		Bromoform				3	4.84E-03 (Min)	4.7E-08
		Carbon tetrachloride				3	4.84E-03 (Min)	7.7E-07
		Chlordane				4	5.00E-04 (Min)	8.0E-07
		Chlorodibromomethane				3	4.84E-03 (Min)	5.0E-07
		Chloroethane				3	9.69E-03 (Min)	1.5E-07
		Chrysene	8	7.00E-01 (Min)	2.4E-06	3	1.00E-02 (Min)	8.9E-07
		Cis-1,3-Dichloropropene				3	4.84E-03 (Min)	1.1E-06
		Dibenz(a,h)anthracene	8	7.00E-01 (Min)	3.7E-04	3	1.00E-02 (Min)	1.4E-04
		Dichlorobromomethane				3	4.84E-03 (Min)	7.7E-07
		Dieldrin				2	1.00E-04 (Min)	2.0E-06
		Fluoranthene	8	7.00E-01 (Min)	2.4E-04	3	1.00E-02 (Min)	8.9E-05
		Fluorene	8	7.00E-01 (Min)	3.7E-04	3	1.00E-02 (Min)	1.4E-04
		Gamma-BHC (Lindane)				2	5.00E-05 (Min)	8.0E-08
		Heptachlor				2	5.00E-05 (Min)	2.8E-07
		Heptachlor epoxide				2	5.00E-05 (Min)	5.6E-07
		Hexachlorobenzene	8	7.00E-01 (Min)	5.2E-05	3	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene	8	7.00E-01 (Min)	2.5E-06	3	1.00E-02 (Min)	9.6E-07
		Hexachloroethane	8	7.00E-01 (Min)	4.5E-07	3	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene	8	7.00E-01 (Min)	2.4E-05	3	1.00E-02 (Min)	8.9E-06
		Isophorone	8	7.00E-01 (Min)	1.3E-07	3	1.00E-02 (Min)	5.0E-08
		N-Nitroso-di-n-propylamine				3	1.00E-02 (Min)	8.6E-05
		N-Nitrosodimethylamine	8	7.00E-01 (Min)	2.3E-04	1	1.00E-02 (Min)	6.2E-04
		N-Nitrosodiphenylamine	8	7.00E-01 (Min)	1.6E-07	3	1.00E-02 (Min)	6.0E-08
		Nitrophenol	8	3.50E-00 (Min)	1.9E-05	2	5.00E-02 (Min)	7.3E-06
		Phenanthrene	8	7.00E-01 (Min)	2.4E-04	3	1.00E-02 (Min)	8.9E-05
		Pyrene	8	7.00E-01 (Min)	2.4E-04	3	1.00E-02 (Min)	8.9E-05
		Styrene				1	4.84E-03 (Min)	1.8E-07

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water
where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
<hr/>								
2	Organics	Toxaphene				2	1.00E-03 (Min)	1.3E-06
		Trans-1,3-dichloropropene				3	4.84E-03 (Min)	1.1E-06
		Trichloroethylene				2	5.00E-03 (Min)	6.7E-08
		Vinyl chloride				3	9.69E-03 (Min)	2.3E-05
<hr/>								
				Bq/kg			Bq/L	
	Radionuclides	Co-60	27	1.58E+00 (Min)	5.3E-08			
<hr/>								
3	Organics	1,1,2,2-Tetrachloroethane				5	5.00E-03 (Min)	1.2E-06
		1,1,2-Trichloroethane				5	5.00E-03 (Min)	3.5E-07
		1,2-Dichloropropane				5	5.00E-03 (Min)	4.2E-07
		1,4-Dichlorobenzene	34	1.90E-01 (Min)	2.1E-07	8	1.00E-02 (Min)	1.3E-07
		2,4,6-Trichlorophenol	34	1.90E-01 (Min)	9.7E-08	8	1.00E-02 (Min)	8.3E-06
		2,4-Dinitrotoluene	34	1.90E-01 (Min)	6.0E-06	8	1.00E-02 (Min)	8.3E-06
		2,6-Dinitrotoluene	34	1.90E-01 (Min)	6.0E-06	8	2.00E-02 (Min)	1.1E-05
		3,3'-Dichlorobenzidine	34	3.70E-01 (Min)	7.7E-06	1	1.00E-04 (Min)	2.9E-08
		4,4'-DDD				1	1.00E-04 (Min)	4.2E-08
		4,4'-DDE				1	1.00E-04 (Min)	4.2E-08
		4,4'-DDT				1	1.00E-04 (Min)	4.3E-07
		4-Chloroaniline	34	1.00E-02 (Min)	1.6E-07	5	1.00E-02 (Min)	4.3E-07
		Acenaphthene	34	1.90E-01 (Min)	3.1E-07	8	1.00E-02 (Min)	1.4E-04
		Acenaphthylene	34	1.90E-01 (Min)	1.0E-04	8	1.00E-02 (Min)	1.4E-04
		Aldrin	34	7.72E-03 (Min)	1.0E-04	1	5.00E-05 (Min)	1.0E-06
		Alpha-BHC				1	5.00E-05 (Min)	3.9E-07
		Anthracene	34	1.90E-01 (Min)	6.4E-05	3	1.00E-03 (Min)	9.4E-06
		Aroclor-1260				4	1.00E-02 (Min)	2.8E-03
		Benzidine						
		Benzo(a)anthracene	34	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-05
		Benzo(a)pyrene	33	1.90E-01 (Min)	6.4E-05	8	1.00E-02 (Min)	8.9E-06
		Benzo(b)fluoranthene	33	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-05
		Benzo(g,h,i)perylene	32	1.90E-01 (Min)	6.4E-05	8	1.00E-02 (Min)	8.9E-06
		Benzo(k)fluoranthene	33	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-06
		Beta-BHC				1	5.00E-05 (Min)	1.1E-07
		Bis(2-chloroethyl)ether	34	1.90E-01 (Min)	9.7E-06	8	1.00E-02 (Min)	1.3E-05
<hr/>								
				mg/kg			mg/L	
		1,1,2,2-Tetrachloroethane				5	5.00E-03 (Min)	1.2E-06
		1,1,2-Trichloroethane				5	5.00E-03 (Min)	3.5E-07
		1,2-Dichloropropane				5	5.00E-03 (Min)	4.2E-07
		1,4-Dichlorobenzene	34	1.90E-01 (Min)	2.1E-07	8	1.00E-02 (Min)	1.3E-07
		2,4,6-Trichlorophenol	34	1.90E-01 (Min)	9.7E-08	8	1.00E-02 (Min)	8.3E-06
		2,4-Dinitrotoluene	34	1.90E-01 (Min)	6.0E-06	8	1.00E-02 (Min)	8.3E-06
		2,6-Dinitrotoluene	34	1.90E-01 (Min)	6.0E-06	8	2.00E-02 (Min)	1.1E-05
		3,3'-Dichlorobenzidine	34	3.70E-01 (Min)	7.7E-06	1	1.00E-04 (Min)	2.9E-08
		4,4'-DDD				1	1.00E-04 (Min)	4.2E-08
		4,4'-DDE				1	1.00E-04 (Min)	4.2E-08
		4,4'-DDT				1	1.00E-04 (Min)	4.3E-07
		4-Chloroaniline	34	1.00E-02 (Min)	1.6E-07	5	1.00E-02 (Min)	4.3E-07
		Acenaphthene	34	1.90E-01 (Min)	3.1E-07	8	1.00E-02 (Min)	1.4E-04
		Acenaphthylene	34	1.90E-01 (Min)	1.0E-04	8	1.00E-02 (Min)	1.4E-04
		Aldrin	34	7.72E-03 (Min)	1.0E-04	1	5.00E-05 (Min)	1.0E-06
		Alpha-BHC				1	5.00E-05 (Min)	3.9E-07
		Anthracene	34	1.90E-01 (Min)	6.4E-05	3	1.00E-03 (Min)	9.4E-06
		Aroclor-1260				4	1.00E-02 (Min)	2.8E-03
		Benzidine						
		Benzo(a)anthracene	34	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-05
		Benzo(a)pyrene	33	1.90E-01 (Min)	6.4E-05	8	1.00E-02 (Min)	8.9E-06
		Benzo(b)fluoranthene	33	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-05
		Benzo(g,h,i)perylene	32	1.90E-01 (Min)	6.4E-05	8	1.00E-02 (Min)	8.9E-06
		Benzo(k)fluoranthene	33	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-06
		Beta-BHC				1	5.00E-05 (Min)	1.1E-07
		Bis(2-chloroethyl)ether	34	1.90E-01 (Min)	9.7E-06	8	1.00E-02 (Min)	1.3E-05

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
3	Organics	Bis(2-chloroisopropyl)ether	34	1.90E-01 (Min)	6.2E-07	8	1.00E-02 (Min)	8.6E-07
		Carbon tetrachloride				5	5.00E-03 (Min)	8.0E-07
		Chlordane				2	5.00E-04 (Min)	8.0E-07
		Chloromethane				5	1.00E-02 (Min)	1.6E-07
		Chrysene	34	1.90E-01 (Min)	6.4E-07			
		Cis-1,3-Dichloropropene				5	5.00E-03 (Min)	1.1E-06
		Dibenz(a,h)anthracene	33	1.90E-01 (Min)	1.0E-04	8	1.00E-02 (Min)	1.4E-04
		Dieldrin				1	1.00E-04 (Min)	2.0E-06
		Fluoranthene	34	1.90E-01 (Min)	6.4E-05			
		Fluorene	34	1.90E-01 (Min)	1.0E-04	8	1.00E-02 (Min)	1.4E-04
		Gamma-BHC (Lindane)				1	5.00E-05 (Min)	8.0E-08
		Heptachlor				1	5.00E-05 (Min)	2.8E-07
		Heptachlor epoxide				1	5.00E-05 (Min)	5.6E-07
		Hexachlorobenzene	34	1.90E-01 (Min)	1.4E-05	8	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene	34	1.90E-01 (Min)	6.9E-07	8	1.00E-02 (Min)	9.6E-07
		Hexachloroethane	34	1.90E-01 (Min)	1.2E-07	8	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene	33	1.90E-01 (Min)	6.4E-06	8	1.00E-02 (Min)	8.9E-06
		Isophorone	34	1.90E-01 (Min)	3.6E-08	8	1.00E-02 (Min)	5.0E-08
		N-Nitroso-di-n-propylamine	34	1.90E-01 (Min)	6.2E-05			
		N-Nitrosodimethylamine				4	1.00E-02 (Min)	6.2E-04
4	Inorganics	N-Nitrosodiphenylamine	34	1.90E-01 (Min)	4.3E-08	8	1.00E-02 (Min)	6.0E-08
		Pentachlorophenol	34	1.90E-01 (Min)	5.2E-06	8	5.00E-02 (Min)	7.3E-06
		Phenanthrene	34	1.90E-01 (Min)	6.4E-05			
		Pyrene	34	1.90E-01 (Min)	6.4E-05	1	5.00E-03 (Min)	1.8E-07
4	Radionuclides	Styrene				1	1.00E-03 (Min)	1.3E-06
		Toxaphene						
		Co-60		Bq/kg		3	2.19E-03 (Min)	2.0E-09
4	Inorganics	Arsenic		mg/kg				
		Beryllium	27	3.00E-03 (Min)	6.0E-07	5	1.30E-03 (Min)	2.8E-06
						3	3.00E-04 (Min)	1.6E-06

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
				mg/kg			mg/L	
4	Organics	1,1,2,2-Tetrachloroethane				2	5.00E-03 (Min)	1.2E-06
		1,1,2-Trichloroethane				2	5.00E-03 (Min)	3.5E-07
		1,1-Dichloroethylene				2	5.00E-03 (Min)	3.7E-06
		1,2-Dichloroethane				2	5.00E-03 (Min)	5.6E-07
		1,2-Dichloropropane				2	5.00E-03 (Min)	4.2E-07
		1,4-Dichlorobenzene	6	6.60E-01 (Min)	7.3E-07	4	9.50E-03 (Min)	2.8E-07
		2,4,6-Trichlorophenol	13	6.60E-01 (Min)	3.4E-07	4	1.00E-02 (Min)	1.3E-07
		2,4-Dinitrotoluene	6	6.60E-01 (Min)	2.1E-05	4	1.00E-02 (Min)	8.3E-06
		2,6-Dinitrotoluene	6	6.60E-01 (Min)	2.1E-05	4	1.00E-02 (Min)	8.3E-06
		3,3'-Dichlorobenzidine	6	1.30E+00 (Min)	2.7E-05	4	2.00E-02 (Min)	1.1E-05
		4,4'-DDD				3	1.00E-04 (Min)	2.9E-08
		4,4'-DDE				3	1.00E-04 (Min)	4.2E-08
		4,4'-DDT				3	1.00E-04 (Min)	4.2E-08
		4-Chloroaniline	6	6.60E-01 (Min)	1.1E-06	2	1.00E-02 (Min)	4.3E-07
		Acenaphthene	6	6.60E-01 (Min)	3.5E-04	4	1.00E-02 (Min)	1.4E-04
		Acenaphthylene	6	6.60E-01 (Min)	3.5E-04	4	1.00E-02 (Min)	1.4E-04
		Aldrin				3	5.00E-05 (Min)	1.0E-06
		Alpha-BHC				3	5.00E-05 (Min)	3.9E-07
		Anthracene	6	6.60E-01 (Min)	2.2E-04	4	1.00E-02 (Min)	8.9E-05
		Aroclor-1254				4	1.00E-03 (Min)	9.4E-06
		Aroclor-1260				4	1.00E-03 (Min)	9.4E-06
		Benzene				2	5.00E-03 (Min)	1.8E-07
		Benzidine				2	1.00E-02 (Min)	2.8E-03
		Benzo(a)anthracene	6	6.60E-01 (Min)	2.2E-05	4	1.00E-02 (Min)	8.9E-06
		Benzo(a)pyrene	6	6.60E-01 (Min)	2.2E-04	4	1.00E-02 (Min)	8.9E-05
		Benzo(b)fluoranthene	6	6.60E-01 (Min)	2.2E-05	4	1.00E-02 (Min)	8.9E-06
		Benzo(g,h,i)perylene	6	6.60E-01 (Min)	2.2E-04	4	1.00E-02 (Min)	8.9E-05
		Benzo(k)fluoranthene	6	6.60E-01 (Min)	2.2E-05	4	1.00E-02 (Min)	8.9E-06
		Beta-BHC				3	5.00E-05 (Min)	1.1E-07
		Bis(2-chloroethyl)ether	7	6.60E-01 (Min)	3.4E-05	4	1.00E-02 (Min)	1.3E-05
		Bis(2-chloroisopropyl)ether	6	6.60E-01 (Min)	2.1E-06	4	1.00E-02 (Min)	8.6E-07
		Bis(2-ethylhexyl)phthalate	6	6.60E-01 (Min)	4.3E-07	4	1.00E-02 (Min)	1.7E-07
		Bromoform				2	5.00E-03 (Min)	4.8E-08
		Carbon tetrachloride				2	5.00E-03 (Min)	8.0E-07
		Chlordane	6			6	5.00E-04 (Min)	8.0E-07
		Chlorodibromomethane				2	5.00E-03 (Min)	5.1E-07

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water
where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
4	Organics	Chloroform				2	5.00E-03 (Min)	3.7E-08
		Chloromethane				2	1.00E-02 (Min)	1.6E-07
		Chrysene	6	6.60E-01 (Min)	2.2E-06	4	1.00E-02 (Min)	8.9E-07
		Cis-1,3-Dichloropropene				2	5.00E-03 (Min)	1.1E-06
		Dibenz(a,h)anthracene	6	6.60E-01 (Min)	3.5E-04	4	1.00E-02 (Min)	1.4E-04
		Dichlorobromomethane				2	5.00E-03 (Min)	8.0E-07
		Dieldrin				3	1.00E-04 (Min)	2.0E-06
		Fluoranthene	7	6.60E-01 (Min)	2.2E-04	4	1.00E-02 (Min)	8.9E-05
		Fluorene	6	6.60E-01 (Min)	3.5E-04	4	1.00E-02 (Min)	1.4E-04
		Gamma-BHC (Lindane)				3	5.00E-05 (Min)	8.0E-08
		Heptachlor				3	5.00E-05 (Min)	2.8E-07
		Heptachlor epoxide				3	5.00E-05 (Min)	5.6E-07
		Hexachlorobenzene	6	6.60E-01 (Min)	4.9E-05	4	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene	6	6.60E-01 (Min)	2.4E-06	4	1.00E-02 (Min)	9.6E-07
		Hexachloroethane	6	6.60E-01 (Min)	4.3E-07	4	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene	6	6.60E-01 (Min)	2.2E-05	4	1.00E-02 (Min)	8.9E-06
		Isophorone	6	6.60E-01 (Min)	1.3E-07	4	1.00E-02 (Min)	5.0E-08
		Methylene chloride				2	5.00E-03 (Min)	4.6E-08
		N-Nitroso-di-n-propylamine	6	6.60E-01 (Min)	2.1E-04	4	1.00E-02 (Min)	8.6E-05
		N-Nitrosodimethylamine				2	1.00E-02 (Min)	6.2E-04
		N-Nitrosodiphenylamine	6	6.60E-01 (Min)	1.5E-07	4	1.00E-02 (Min)	6.0E-08
		Pentachlorophenol	13	3.30E+00 (Min)	1.8E-05	4	5.00E-02 (Min)	7.3E-06
		Phenanthrene	6	6.60E-01 (Min)	2.2E-04	4	1.00E-02 (Min)	8.9E-05
		Pyrene	6	6.60E-01 (Min)	2.2E-04	4	1.00E-02 (Min)	8.9E-05
		Tetrachloroethylene				2	5.00E-03 (Min)	3.1E-07
		Toxaphene				3	1.00E-03 (Min)	1.3E-06
		Trans-1,3-dichloropropene				2	5.00E-03 (Min)	1.1E-06
		Trichloroethylene				2	5.00E-03 (Min)	6.7E-08
		Vinyl chloride				2	1.00E-02 (Min)	2.3E-05
						Bq/L		
	Radionuclides	Co-60	38	2.33E-01 (Min)	7.9E-09			
							mg/L	
5	Inorganics	Arsenic	4	1.30E-03 (Min)	2.8E-06			

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
5	Inorganics	Beryllium	4	3.90E-03 (Min)	2.1E-05			
	Organics	1,4-Dichlorobenzene	4	1.00E-02 (Min)	2.9E-07			
		2,4,6-Trichlorophenol	4	1.00E-02 (Min)	1.3E-07			
		2,4-Dinitrotoluene	4	1.00E-02 (Min)	8.3E-06			
		2,6-Dinitrotoluene	4	1.00E-02 (Min)	8.3E-06			
		3,3'-Dichlorobenzidine	4	2.00E-02 (Min)	1.1E-05			
		4,4'-DDD	3	1.00E-04 (Min)	2.9E-08			
		4,4'-DDE	3	1.00E-04 (Min)	4.2E-08			
		4,4'-DDT	3	1.00E-04 (Min)	4.2E-08			
		4-Chloroaniline	4	1.00E-02 (Min)	4.3E-07			
		Acenaphthene	4	1.00E-02 (Min)	1.4E-04			
		Acenaphthylene	4	1.00E-02 (Min)	1.4E-04			
		Aldrin	3	5.00E-05 (Min)	1.0E-06			
		Alpha-BHC	3	5.00E-05 (Min)	3.9E-07			
		Anthracene	4	1.00E-02 (Min)	8.9E-05			
		Aroclor-1254	3	1.00E-03 (Min)	9.4E-06			
		Aroclor-1260	3	1.00E-03 (Min)	9.4E-06			
		Benzo(a)anthracene	4	1.00E-02 (Min)	8.9E-06			
		Benzo(a)pyrene	4	1.00E-02 (Min)	8.9E-05			
		Benzo(b)fluoranthene	4	1.00E-02 (Min)	8.9E-06			
		Benzo(g,h,i)perylene	4	1.00E-02 (Min)	8.9E-05			
		Benzo(k)fluoranthene	4	1.00E-02 (Min)	8.9E-06			
		Beta-BHC	3	5.00E-05 (Min)	1.1E-07			
		Bis(2-chloroethyl)ether	4	1.00E-02 (Min)	1.3E-05			
		Bis(2-chloroisopropyl)ether	4	1.00E-02 (Min)	8.6E-07			
		Bis(2-ethylhexyl)phthalate	4	3.00E-03 (Min)	5.1E-08			
		Chlordane	6	5.00E-04 (Min)	8.0E-07			
		Chrysene	4	1.00E-02 (Min)	8.9E-07			
		Dibenz(a,h)anthracene	4	1.00E-02 (Min)	1.4E-04			
		Dieldrin	3	1.00E-04 (Min)	2.0E-06			
		Fluoranthene	4	1.00E-02 (Min)	8.9E-05			
		Fluorene	4	1.00E-02 (Min)	1.4E-04			
		Gamma-BHC (Lindane)	3	5.00E-05 (Min)	8.0E-08			
		Heptachlor	3	5.00E-05 (Min)	2.8E-07			
		Heptachlor epoxide	3	5.00E-05 (Min)	5.6E-07			

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
5	Organics	Hexachlorobenzene				4	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene				4	1.00E-02 (Min)	9.6E-07
		Hexachloroethane				4	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene				4	1.00E-02 (Min)	8.9E-06
		Isophorone				4	1.00E-02 (Min)	5.0E-08
		N-Nitroso-di-n-propylamine				4	1.00E-02 (Min)	8.6E-05
		N-Nitrosodiphenylamine				4	1.00E-02 (Min)	6.0E-08
		Pentachlorophenol				4	5.00E-02 (Min)	7.3E-06
		Phenanthrene	9	3.30E+00 (Min)	1.8E-05	4	1.00E-02 (Min)	8.9E-05
		Pyrene				4	1.00E-02 (Min)	8.9E-05
		Toxaphene				3	1.00E-03 (Min)	1.3E-06
				Bq/kg			Bq/L	
		Radionuclides Co-60	28	1.08E+00 (Min)	3.7E-08	3	3.18E-03 (Min)	2.9E-09
10	Inorganics	Arsenic		mg/kg		1	1.30E-03 (Min)	2.8E-06
		Beryllium				1	3.90E-03 (Min)	2.1E-05
	Organics			mg/kg			mg/L	
		1,4-Dichlorobenzene				1	1.00E-02 (Min)	2.9E-07
		2,4-Dinitrotoluene				1	1.00E-02 (Min)	8.3E-06
		2,6-Dinitrotoluene				1	1.00E-02 (Min)	8.3E-06
		3,3'-Dichlorobenzidine				1	2.00E-02 (Min)	1.1E-05
		4,4'-DDD				1	1.00E-04 (Min)	2.9E-08
		4,4'-DDE				1	1.00E-04 (Min)	4.2E-08
		4,4'-DDT				1	1.00E-04 (Min)	4.2E-08
		4-Chloroaniline	8	1.00E-02 (Min)	1.6E-07	1	1.00E-02 (Min)	4.3E-07
		Acenaphthene				1	1.00E-02 (Min)	1.4E-04
		Acenaphthylene				1	1.00E-02 (Min)	1.0E-06
		Aldrin	8	1.00E-02 (Min)	7.9E-06	1	5.00E-05 (Min)	1.0E-06
		Alpha-BHC				1	5.00E-05 (Min)	3.9E-07
		Anthracene				1	1.00E-02 (Min)	8.9E-05
		Aroclor-1254	8	1.00E-02 (Min)	3.8E-06	1	1.00E-03 (Min)	9.4E-06

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
10	Organics	Aroclor-1260	1			1	1.00E-03 (Min)	9.4E-06
		Benzo(a)anthracene	1			1	1.00E-02 (Min)	8.9E-06
		Benzo(a)pyrene	1			1	1.00E-02 (Min)	8.9E-05
		Benzo(b)fluoranthene	1			1	1.00E-02 (Min)	8.9E-06
		Benzo(g,h,i)perylene	1			1	1.00E-02 (Min)	8.9E-05
		Benzo(k)fluoranthene	1			1	1.00E-02 (Min)	8.9E-06
		Beta-BHC	1			1	5.00E-05 (Min)	1.1E-07
		Bis(2-chloroethyl)ether	1			1	1.00E-02 (Min)	1.3E-05
		Bis(2-chloroisopropyl)ether	1			1	1.00E-02 (Min)	8.6E-07
		Chlordane	2			2	5.00E-04 (Min)	8.0E-07
		Chrysene	1			1	1.00E-02 (Min)	8.9E-07
		Dibenz(a,h)anthracene	1			1	1.00E-02 (Min)	1.4E-04
		Dieldrin	1			1	1.00E-04 (Min)	2.0E-06
		Fluoranthene	1			1	1.00E-02 (Min)	8.9E-05
		Fluorene	1			1	1.00E-02 (Min)	1.4E-04
		Gamma-BHC (Lindane)	1			1	5.00E-05 (Min)	8.0E-08
		Heptachlor	1			1	5.00E-05 (Min)	2.8E-07
		Heptachlor epoxide	1			1	5.00E-05 (Min)	5.6E-07
		Hexachlorobenzene	1			1	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene	1			1	1.00E-02 (Min)	9.6E-07
		Hexachloroethane	1			1	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene	1			1	1.00E-02 (Min)	8.9E-06
		Isophorone	1			1	1.00E-02 (Min)	5.0E-08
		N-Nitroso-di-n-propylamine	1			1	1.00E-02 (Min)	8.6E-05
		N-Nitrosodiphenylamine	1			1	1.00E-02 (Min)	6.0E-08
		Pentachlorophenol	1			1	5.00E-02 (Min)	7.3E-06
		Phenanthrene	1			1	1.00E-02 (Min)	8.9E-05
		Pyrene	1			1	1.00E-02 (Min)	8.9E-05
		Toxaphene	1			1	1.00E-03 (Min)	1.3E-06
				Bq/kg			Bq/L	
	Radionuclides	Cs-137	1			1	2.37E-03 (Min)	3.9E-09
		Co-60	1			1	2.80E-03 (Min)	2.5E-09
				mg/kg			mg/L	

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index	
13	Inorganics	Arsenic	3	3.00E-03 (Min)	8.0E-07	2	1.30E-03 (Min)	2.8E-08	
		Beryllium				1	3.90E-03 (Min)	2.1E-05	
	Organics	1,1,2,2-Tetrachloroethane				1	5.00E-03 (Min)	1.2E-06	
		1,1,2-Trichloroethane				1	5.00E-03 (Min)	3.5E-07	
		1,1-Dichloroethylene				1	5.00E-03 (Min)	3.7E-06	
		1,2-Dichloroethane				1	5.00E-03 (Min)	5.6E-07	
		1,2-Dichloropropane				1	5.00E-03 (Min)	4.2E-07	
		1,4-Dichlorobenzene				2	1.00E-02 (Min)	2.9E-07	
		2,4,6-Trichlorophenol				2	1.00E-02 (Min)	1.3E-07	
		2,4-Dinitrotoluene				2	1.00E-02 (Min)	8.3E-06	
		2,6-Dinitrotoluene				2	1.00E-02 (Min)	8.3E-06	
		3,3'-Dichlorobenzidine				2	2.00E-02 (Min)	1.1E-05	
		4,4'-DDD			4	1.00E-02 (Min)	1.1E-07		
		4,4'-DDE			3	1.00E-02 (Min)	1.6E-07		
		4-Chloroaniline					1	1.00E-02 (Min)	4.3E-07
		Acenaphthene					2	1.00E-02 (Min)	1.4E-04
		Acenaphthylene					2	1.00E-02 (Min)	1.4E-04
		Anthracene					2	1.00E-02 (Min)	8.9E-05
		Benzene					1	5.00E-03 (Min)	1.8E-07
		Benzidine					1	1.00E-02 (Min)	2.8E-03
		Benzo(a)anthracene					2	1.00E-02 (Min)	8.9E-06
		Benzo(a)pyrene					2	1.00E-02 (Min)	8.9E-05
		Benzo(b)fluoranthene					2	1.00E-02 (Min)	8.9E-05
		Benzo(g,h,i)perylene					2	1.00E-02 (Min)	8.9E-06
		Benzo(k)fluoranthene					2	1.00E-02 (Min)	1.3E-05
		Bis(2-chloroethyl)ether					2	1.00E-02 (Min)	8.6E-07
		Bis(2-chloroisopropyl)ether					2	1.00E-02 (Min)	1.7E-07
		Bis(2-ethylhexyl)phthalate					2	1.00E-02 (Min)	4.8E-08
		Bromoform					1	5.00E-03 (Min)	8.0E-07
		Carbon tetrachloride					1	5.00E-03 (Min)	5.1E-07
		Chlorodibromomethane					1	5.00E-03 (Min)	3.7E-08
		Chloroform					1	1.00E-02 (Min)	1.6E-07
		Chloromethane					2	1.00E-02 (Min)	8.9E-07
		Chrysene					1	5.00E-03 (Min)	1.1E-06
		Cis-1,3-Dichloropropene							

Table E.15 Results of adult nonconservative screening for carcinogens in fish and water
where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
13	Organics	Dibenz(a,h)anthracene	2			2	1.00E-02 (Min)	1.4E-04
		Dichlorobromomethane	1			1	5.00E-03 (Min)	8.0E-07
		Fluoranthene	2			2	1.00E-02 (Min)	8.9E-05
		Fluorene	2			2	1.00E-02 (Min)	1.4E-04
		Hexachlorobenzene	2			2	1.00E-02 (Min)	2.0E-05
		Hexachlorobutadiene	2			2	1.00E-02 (Min)	9.6E-07
		Hexachloroethane	2			2	1.00E-02 (Min)	1.7E-07
		Indeno(1,2,3-cd)pyrene	2			2	1.00E-02 (Min)	8.9E-08
		Isophorone	2			2	1.00E-02 (Min)	5.0E-08
		Methylene chloride	1			1	5.00E-03 (Min)	4.6E-08
		N-Nitroso-di-n-propylamine	2			2	1.00E-02 (Min)	8.6E-05
		N-Nitrosodimethylamine	1			1	1.00E-02 (Min)	6.2E-04
		N-Nitrosodiphenylamine	2			2	1.00E-02 (Min)	6.0E-08
		Pentachlorophenol	2	3.40E+00 (Min)	1.9E-05	2	5.00E-02 (Min)	7.3E-06
		Phenanthrene	2			2	1.00E-02 (Min)	8.9E-05
		Pyrene	2			2	1.00E-02 (Min)	8.9E-05
		Tetrachloroethylene	1			1	5.00E-03 (Min)	3.1E-07
		Trans-1,3-dichloropropene	1			1	5.00E-03 (Min)	1.1E-06
		Trichloroethylene	1			1	5.00E-03 (Min)	6.7E-08
		Vinyl chloride	1			1	1.00E-02 (Min)	2.3E-05
				Bq/kg		Bq/L		
			1			8.37E-03 (Min)	7.5E-09	

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water
where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
				mg/kg			mg/L	
1	Inorganics	Antimony	28	2.80E-01 (Min)	6.3E-02	1	8.00E-02 (Min)	2.5E-03
		Boron						
		Cadmium	28	1.10E-01 (Min)	2.0E-02			
		Chromium	28	2.80E-01 (Min)	5.0E-03			
		Copper	28	2.80E-01 (Min)	6.8E-04			
		Molybdenum				1	4.00E-02 (Min)	2.3E-02
		Nickel	28	2.80E-01 (Min)	1.3E-03			
		Silver	28	1.10E-01 (Min)	2.0E-03	4	7.40E-04 (Min)	4.2E-04
		Thallium	28	2.00E-02 (Min)	2.6E-02	3	2.70E-03 (Min)	1.1E-01
		Tin				1	5.00E-02 (Min)	2.4E-04
	Organics	1,2,4-Trichlorobenzene	6	6.80E-01 (Min)	6.1E-03	3	1.00E-02 (Min)	2.9E-03
		1,2-Dichlorobenzene	6	6.80E-01 (Min)	6.8E-04	3	1.00E-02 (Min)	3.2E-04
		2,4,5-Trichlorophenol	6	3.40E+00 (Min)	3.1E-03	3	5.00E-02 (Min)	1.4E-03
		2,4-Dichlorophenol	6	6.80E-01 (Min)	2.0E-02	3	1.00E-02 (Min)	9.5E-03
		2,4-Dimethylphenol	6	6.80E-01 (Min)	3.1E-03	3	1.00E-02 (Min)	1.4E-03
		2,4-Dinitrophenol	6	3.40E+00 (Min)	1.5E-01	3	5.00E-02 (Min)	7.1E-02
		2-Chloronaphthalene	6	6.80E-01 (Min)	7.7E-04	3	1.00E-02 (Min)	3.6E-04
		2-Chlorophenol	6	6.80E-01 (Min)	1.2E-02	3	1.00E-02 (Min)	5.7E-03
		2-Methylphenol	6	6.80E-01 (Min)	1.2E-03	3	1.00E-02 (Min)	5.7E-04
		4,4'-DDT	16	1.00E-02 (Min)	1.8E-03	2	1.00E-04 (Min)	5.7E-04
		4,6-Dinitro-2-methylphenol	6	3.40E+00 (Min)	3.1E+00	3	5.00E-02 (Min)	1.4E+00
		4-Chloroaniline	6	6.80E-01 (Min)	1.5E-02	3	1.00E-02 (Min)	7.1E-03
		4-Methylphenol	6	6.80E-01 (Min)	1.2E-03	3	1.00E-02 (Min)	5.7E-04
		Acenaphthene	6	6.80E-01 (Min)	1.0E-03	3	1.00E-02 (Min)	4.8E-04
		Aldrin	16	1.00E-02 (Min)	3.0E-02	2	5.00E-05 (Min)	4.8E-03
		Anthracene	6	6.80E-01 (Min)	2.0E-04	3	1.00E-02 (Min)	9.5E-05
		Benzoic acid	6	3.40E+00 (Min)	7.7E-05	3	5.00E-02 (Min)	3.6E-05
		Benzyl alcohol	6	6.80E-01 (Min)	2.0E-04	3	1.00E-02 (Min)	9.5E-05
		Bis(2-chloroisopropyl)ether	6	6.80E-01 (Min)	1.5E-03	3	1.00E-02 (Min)	7.1E-04
		Bis(2-ethylhexyl)phthalate	6	6.80E-01 (Min)	3.1E-03	3	1.00E-02 (Min)	1.4E-03
		Butylbenzyl phthalate	6	6.80E-01 (Min)	3.1E-04	3	1.00E-02 (Min)	1.4E-04
		Chlordane				4	5.00E-04 (Min)	2.4E-02
		Di-n-butyl phthalate	6	6.80E-01 (Min)	6.1E-04	3	1.00E-02 (Min)	2.9E-04

Table E.18 Results of adult nonconservative screening for noncarcinogens in fish and water
where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
1	Organics	Di-n-octylphthalate	6	6.80E-01 (Min)	3.1E-03	3	1.00E-02 (Min)	1.4E-03
		Dieldrin				2	1.00E-04 (Min)	5.7E-03
		Diethylphthalate	6	6.80E-01 (Min)	7.7E-05	3	1.00E-02 (Min)	3.6E-05
		Dimethylphthalate	6	6.80E-01 (Min)	6.1E-05	3	1.00E-02 (Min)	2.9E-05
		Endosulfan I				2	5.00E-05 (Min)	2.9E-03
		Endosulfan II				2	1.00E-04 (Min)	5.7E-03
		Endrin				2	1.00E-04 (Min)	9.5E-04
		Fluoranthene	6	6.80E-01 (Min)	1.5E-03	3	1.00E-02 (Min)	7.1E-04
		Fluorene	6	6.80E-01 (Min)	1.5E-03	3	1.00E-02 (Min)	7.1E-04
		Gamma-BHC (Lindane)				2	5.00E-05 (Min)	4.8E-04
		Heptachlor				2	5.00E-05 (Min)	2.9E-04
		Heptachlor epoxide				2	5.00E-05 (Min)	1.1E-02
		Hexachlorobenzene	6	6.80E-01 (Min)	7.7E-02	3	1.00E-02 (Min)	3.6E-02
		Hexachlorobutadiene	6	6.80E-01 (Min)	3.1E-02	3	1.00E-02 (Min)	1.4E-02
		Hexachlorocyclopentadiene	6	6.80E-01 (Min)	8.7E-03	3	1.00E-02 (Min)	4.1E-03
		Hexachloroethane	6	6.80E-01 (Min)	6.1E-02	3	1.00E-02 (Min)	2.9E-02
		Isophorone	6	6.80E-01 (Min)	3.1E-04	3	1.00E-02 (Min)	1.4E-04
		Methoxychlor				2	5.00E-04 (Min)	2.9E-04
		Naphthalene	6	6.80E-01 (Min)	1.5E-03	3	1.00E-02 (Min)	7.1E-04
		Nitrobenzene	6	6.80E-01 (Min)	1.2E-01	3	1.00E-02 (Min)	5.7E-02
		Pentachlorophenol	6	3.40E+00 (Min)	1.0E-02	3	5.00E-02 (Min)	4.8E-03
		Phenol	6	6.80E-01 (Min)	1.0E-04	3	1.00E-02 (Min)	4.8E-05
		Pyrene	6	6.80E-01 (Min)	2.0E-03	3	1.00E-02 (Min)	9.5E-04
2	Inorganics	Antimony						
		Arsenic	13	4.60E-01 (Min)	1.0E-01	4	1.90E-03 (Min)	1.4E-02
		Beryllium				3	1.30E-03 (Min)	1.2E-02
		Boron	13	3.00E-03 (Min)	5.4E-05	4	2.00E-04 (Min)	1.1E-04
		Cadmium	13	1.80E-01 (Min)	3.2E-02	2	4.00E-03 (Min)	1.3E-04
		Chromium	13	4.60E-01 (Min)	8.3E-03	3	2.10E-04 (Min)	1.2E-03
		Copper	13	4.60E-01 (Min)	1.1E-03	3	9.80E-03 (Min)	5.6E-03
		Mercury				3	2.00E-04 (Min)	1.9E-03
		Molybdenum	13	4.60E-01 (Min)	2.1E-03	2	1.00E-02 (Min)	5.7E-03
		Nickel	13	1.80E-01 (Min)	3.2E-03	3	7.40E-04 (Min)	4.2E-04
		Silver						

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water
where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
2	Inorganics	Thallium	13	2.00E-02 (Min)	2.6E-02	3	2.70E-03 (Min)	1.1E-01
		Tin				2	1.00E-02 (Min)	4.8E-05
		Vanadium				1	2.00E-03 (Min)	8.2E-04
	Organics			mg/kg			mg/L	
1,1,1-Trichloroethane					3	4.84E-03 (Min)	1.5E-04	
1,1,2-Trichloroethane					3	4.84E-03 (Min)	3.5E-03	
1,1-Dichloroethane					3	4.84E-03 (Min)	1.4E-04	
1,1-Dichloroethylene					3	4.84E-03 (Min)	1.5E-03	
1,2,4-Trichlorobenzene		8	7.00E-01 (Min)	6.3E-03	3	1.00E-02 (Min)	2.9E-03	
1,2-Dichlorobenzene		8	7.00E-01 (Min)	7.0E-04	3	9.50E-03 (Min)	3.0E-04	
1,2-Dichloroethylene (total)					1	4.84E-03 (Min)	1.4E-03	
2,4,5-Trichlorophenol		8	3.50E+00 (Min)	3.2E-03	1	5.00E-02 (Min)	1.4E-03	
2,4-Dichlorophenol		8	7.00E-01 (Min)	2.1E-02	2	1.00E-02 (Min)	9.5E-03	
2,4-Dimethylphenol		8	7.00E-01 (Min)	3.2E-03	2	1.00E-02 (Min)	1.4E-03	
2,4-Dinitrophenol		8	3.50E+00 (Min)	1.6E-01	1	5.00E-02 (Min)	7.1E-02	
2-Chloronaphthalene		8	7.00E-01 (Min)	7.9E-04	3	1.00E-02 (Min)	3.6E-04	
2-Chlorophenol		8	7.00E-01 (Min)	1.3E-02	2	1.00E-02 (Min)	5.7E-03	
2-Methylphenol		8	7.00E-01 (Min)	1.3E-03	1	1.00E-02 (Min)	5.7E-04	
4,4'-DDT					2	1.00E-04 (Min)	5.7E-04	
4,6-Dinitro-2-methylphenol		8	3.50E+00 (Min)	3.2E+00	2	5.00E-02 (Min)	1.4E+00	
4-Chloroaniline		2	7.00E-01 (Min)	1.6E-02	2	1.00E-02 (Min)	7.1E-03	
4-Methyl-2-pentanone					1	9.69E-03 (Min)	5.5E-04	
4-Methylphenol		8	7.00E-01 (Min)	1.3E-03	1	1.00E-02 (Min)	5.7E-04	
Acenaphthene		8	7.00E-01 (Min)	1.1E-03	3	1.00E-02 (Min)	4.8E-04	
Aldrin					2	5.00E-05 (Min)	4.8E-03	
Anthracene		8	7.00E-01 (Min)	2.1E-04	3	1.00E-02 (Min)	9.5E-05	
Benzo(a)pyrene					1	2.00E-02 (Min)	1.9E-02	
Benzoic acid		8	3.50E+00 (Min)	7.9E-05	2	5.00E-02 (Min)	3.6E-05	
Benzyl alcohol		8	7.00E-01 (Min)	2.1E-04	2	1.00E-02 (Min)	9.5E-05	
Bis(2-chloroisopropyl)ether		8	7.00E-01 (Min)	1.6E-03	3	1.00E-02 (Min)	7.1E-04	
Bis(2-ethylhexyl)phthalate		8	7.00E-01 (Min)	3.2E-03	3	1.00E-02 (Min)	1.4E-03	
Bromoforn					3	4.84E-03 (Min)	6.9E-04	
Butylbenzylphthalate		8	7.00E-01 (Min)	3.2E-04	3	1.00E-02 (Min)	1.4E-04	
Carbon tetrachloride				3	4.84E-03 (Min)	2.0E-02		
Chlordane				4	5.00E-04 (Min)	2.4E-02		
Chlorobenzene				3	4.84E-03 (Min)	6.9E-04		

Table E.18 Results of adult nonconservative screening for nencarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
2	Organics	Chlorodibromomethane				3	4.84E-03 (Min)	6.9E-04
		Cis-1,3-Dichloropropene				3	4.84E-03 (Min)	4.8E-02
		Di-n-butylphthalate	8	7.00E-01 (Min)	6.3E-04	3	1.00E-02 (Min)	2.9E-04
		Di-n-octylphthalate	8	8.20E-02 (Min)	3.7E-04	3	1.00E-02 (Min)	1.4E-03
		Dichlorobromomethane				3	4.84E-03 (Min)	6.9E-04
		Dieldrin				2	1.00E-04 (Min)	5.7E-03
		Diethylphthalate	8	8.00E-02 (Min)	9.0E-06	3	1.00E-02 (Min)	3.6E-05
		Dimethylphthalate	8	7.00E-01 (Min)	6.3E-05	3	1.00E-02 (Min)	2.9E-05
		Endosulfan I				2	5.00E-05 (Min)	2.9E-03
		Endosulfan II				2	1.00E-04 (Min)	5.7E-03
		Endrin				2	1.00E-04 (Min)	9.5E-04
		Ethylbenzene	8	7.00E-01 (Min)	1.6E-03	3	4.84E-03 (Min)	1.4E-04
		Fluoranthene	8	7.00E-01 (Min)	1.6E-03	3	1.00E-02 (Min)	7.1E-04
		Fluorene				3	1.00E-02 (Min)	7.1E-04
		Gamma-BHC (Lindane)				2	5.00E-05 (Min)	4.8E-04
		Heptachlor				2	5.00E-05 (Min)	2.9E-04
		Heptachlor epoxide				2	5.00E-05 (Min)	1.1E-02
		Hexachlorobenzene	8	7.00E-01 (Min)	7.9E-02	3	1.00E-02 (Min)	3.6E-02
		Hexachlorobutadiene	8	7.00E-01 (Min)	3.2E-02	3	1.00E-02 (Min)	1.4E-02
		Hexachlorocyclopentadiene	8	7.00E-01 (Min)	9.0E-03	3	1.00E-02 (Min)	4.1E-03
		Hexachloroethane	8	7.00E-01 (Min)	6.3E-02	3	1.00E-02 (Min)	2.9E-02
		Isophorone	8	7.00E-01 (Min)	3.2E-04	3	1.00E-02 (Min)	1.4E-04
		Isophorone				2	5.00E-04 (Min)	2.9E-04
		Methoxychlor				2	5.00E-04 (Min)	2.9E-04
		Methyl bromide	8	7.00E-01 (Min)	1.6E-03	3	9.69E-03 (Min)	2.0E-02
		Naphthalene	8	7.00E-01 (Min)	1.3E-01	3	1.00E-02 (Min)	7.1E-04
		Nitrobenzene	8	3.50E+09 (Min)	1.1E-02	3	1.00E-02 (Min)	5.7E-02
		Pentachlorophenol	8	7.00E-01 (Min)	1.1E-04	2	5.00E-02 (Min)	4.8E-03
		Phenol						
		Pyrene	8	7.00E-01 (Min)	1.1E-03	3	1.00E-02 (Min)	9.5E-04
		Styrene				1	4.84E-03 (Min)	6.9E-05
		Trans-1,2-dichloroethylene	2			2	5.00E-03 (Min)	7.1E-04
		Trans-1,3-dichloropropene	3			3	4.84E-03 (Min)	4.6E-02
		Trichlorofluoromethane	1			1	5.00E-03 (Min)	4.8E-05
		Vinyl acetate	1			1	9.69E-03 (Min)	2.8E-05

mg/L

mg/kg

Reach	Contaminant type	Contaminant	Fish ingest.			Water ingest.		
			Fish samples	Fish concentration	Fish carcinogen screening index	Water samples	Water concentration	Water carcinogen screening index
3	Inorganics	Antimony	35	3.00E-01 (Min)	6.8E-02	9	1.90E-03 (Min)	1.4E-02
		Cadmium	35	1.20E-01 (Min)	2.2E-02			
		Chromium	35	3.00E-01 (Min)	5.4E-03			
		Copper	35	3.00E-01 (Min)	7.3E-04			
		Nickel	35	3.00E-01 (Min)	1.4E-03			
		Silver	35	1.20E-01 (Min)	2.2E-03	8	2.70E-03 (Min)	1.1E-01
		Thallium	35	2.00E-02 (Min)	2.6E-02	1	5.00E-01 (Min)	2.0E-01
		Vanadium						
	Organics	1,1,2-Trichloroethane				5	5.00E-03 (Min)	3.6E-03
		1,2,4-Trichlorobenzene	34	1.90E-01 (Min)	1.7E-03	8	1.00E-02 (Min)	2.9E-03
		1,2-Dichlorobenzene	34	1.90E-01 (Min)	1.9E-04	9	4.86E-03 (Min)	1.5E-04
		1,2-Dichloroethylene (total)				1	5.00E-03 (Min)	1.4E-03
		2,4,5-Trichlorophenol	34	9.30E-01 (Min)	8.4E-04	5	5.00E-02 (Min)	1.4E-03
		2,4-Dichlorophenol	34	1.90E-01 (Min)	5.7E-03			
		2,4-Dimethylphenol	34	1.90E-01 (Min)	8.6E-04	8	1.00E-02 (Min)	1.4E-03
		2,4-Dinitrophenol	34	9.30E-01 (Min)	4.2E-02	4	5.00E-02 (Min)	7.1E-02
		2-Chloronaphthalene	34	1.90E-01 (Min)	2.1E-04	8	1.00E-02 (Min)	3.6E-04
		2-Chlorophenol	34	1.90E-01 (Min)	3.4E-03	8	1.00E-02 (Min)	5.7E-03
		2-Methylphenol	34	1.90E-01 (Min)	3.4E-04	5	1.00E-02 (Min)	5.7E-04
		4,4'-DDT	34	1.00E-02 (Min)	1.8E-03	1	1.00E-04 (Min)	5.7E-04
		4,6-Dinitro-2-methylphenol	34	9.30E-01 (Min)	8.4E-01	8	5.00E-02 (Min)	1.4E+00
		4-Chloroaniline	34	1.90E-01 (Min)	4.3E-03	5	1.00E-02 (Min)	7.1E-03
		4-Methyl-2-pentanone				1	1.00E-02 (Min)	5.7E-04
		4-Methylphenol	34	1.90E-01 (Min)	3.4E-04	5	1.00E-02 (Min)	5.7E-04
		Acenaphthene	34	1.90E-01 (Min)	2.9E-04	8	1.00E-02 (Min)	4.8E-04
		Acetone				1	1.00E-02 (Min)	2.9E-04
		Aldrin	34	7.72E-03 (Min)	2.3E-02	1	5.00E-05 (Min)	4.8E-03
		Anthracene	34	1.90E-01 (Min)	5.7E-05			
		Benzidine				4	1.00E-02 (Min)	9.5E-03
		Benzoic acid	34	9.30E-01 (Min)	2.1E-05	5	5.00E-02 (Min)	3.6E-05
		Benzyl alcohol	34	1.90E-01 (Min)	5.7E-05	5	1.00E-02 (Min)	9.5E-05
		Bis(2-chloroisopropyl)ether	34	1.90E-01 (Min)	4.3E-04	8	1.00E-02 (Min)	7.1E-04
		Butylbenzylphthalate						
		Carbon tetrachloride	34	1.90E-01 (Min)	8.6E-05			
		Chlordane				5	5.00E-03 (Min)	2.0E-02
				2	5.00E-04 (Min)	2.4E-02		

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
3	Organics	Cis-1,3-Dichloropropene	34	1.90E-01 (Min)	1.7E-04	5	5.00E-03 (Min)	4.8E-02
		Di-n-butylphthalate	34	1.90E-01 (Min)	1.7E-04	8	1.00E-02 (Min)	1.4E-03
		Di-n-octylphthalate	34	1.90E-01 (Min)	1.7E-05	1	1.00E-04 (Min)	5.7E-03
		Dieldrin	34	1.90E-01 (Min)	2.1E-05	1	5.00E-05 (Min)	2.9E-03
		Diethylphthalate	34	1.90E-01 (Min)	1.7E-05	1	1.00E-04 (Min)	5.7E-03
		Dimethylphthalate	34	1.90E-01 (Min)	1.7E-05	1	1.00E-04 (Min)	9.5E-04
		Endosulfan I	34	1.90E-01 (Min)	1.7E-05	1	1.00E-04 (Min)	9.5E-04
		Endosulfan II	34	1.90E-01 (Min)	1.7E-05	1	1.00E-04 (Min)	9.5E-04
		Endrin	34	1.90E-01 (Min)	4.3E-04	8	1.00E-02 (Min)	7.1E-04
		Fluoranthene	34	1.90E-01 (Min)	4.3E-04	1	5.00E-05 (Min)	4.8E-04
		Fluorene	34	1.90E-01 (Min)	4.3E-04	1	5.00E-05 (Min)	2.9E-04
		Gamma-BHC (Lindane)	34	1.90E-01 (Min)	4.3E-04	1	5.00E-05 (Min)	1.1E-02
		Heptachlor	34	1.90E-01 (Min)	2.1E-02	8	1.00E-02 (Min)	3.6E-02
		Heptachlor epoxide	34	1.90E-01 (Min)	8.6E-03	8	1.00E-02 (Min)	1.4E-02
		Hexachlorobenzene	34	1.90E-01 (Min)	2.4E-03	8	1.00E-02 (Min)	4.1E-03
		Hexachlorobutadiene	34	1.90E-01 (Min)	1.7E-02	8	1.00E-02 (Min)	2.9E-02
		Hexachlorocyclopentadiene	34	1.90E-01 (Min)	8.6E-05	8	1.00E-02 (Min)	1.4E-04
		Hexachloroethane	34	1.90E-01 (Min)	8.6E-05	1	5.00E-04 (Min)	2.9E-04
		Isophorone	34	1.90E-01 (Min)	8.6E-05	5	9.99E-03 (Min)	2.0E-02
		Methoxychlor	34	1.90E-01 (Min)	4.3E-04	8	1.00E-02 (Min)	7.1E-04
		Methyl bromide	34	1.90E-01 (Min)	3.4E-02	8	1.00E-02 (Min)	5.7E-02
		Naphthalene	34	1.90E-01 (Min)	2.8E-03	8	1.00E-02 (Min)	4.8E-03
		Nitrobenzene	34	9.30E-01 (Min)	2.8E-03	8	1.00E-02 (Min)	4.8E-03
		Pentachlorophenol	34	1.90E-01 (Min)	2.9E-05	8	5.00E-02 (Min)	4.8E-03
		Phenol	34	1.90E-01 (Min)	5.7E-04	1	5.00E-03 (Min)	7.1E-05
		Pyrene	34	1.90E-01 (Min)	5.7E-04	4	5.00E-03 (Min)	4.8E-05
		Styrene	34	1.90E-01 (Min)	5.7E-04	1	1.00E-02 (Min)	2.9E-05
		Trichlorofluoromethane	34	1.90E-01 (Min)	5.7E-04	1	5.00E-03 (Min)	7.1E-06
		Vinyl acetate	34	1.90E-01 (Min)	5.7E-04	1	5.00E-03 (Min)	7.1E-06
		Xylene	34	1.90E-01 (Min)	5.7E-04	1	5.00E-03 (Min)	7.1E-06
4	Inorganics	Antimony	27	4.80E-01 (Min)	1.1E-01	4	1.90E-03 (Min)	1.4E-02
		Arsenic	27	3.00E-03 (Min)	5.4E-05	5	1.30E-03 (Min)	1.2E-02
		Beryllium	27	1.90E-01 (Min)	3.4E-02	3	3.00E-04 (Min)	1.7E-04
		Cadmium	27	1.90E-01 (Min)	3.4E-02	5	2.10E-04 (Min)	1.2E-03

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
4	Inorganics	Chromium	27	4.80E-01 (Min)	8.8E-03			
		Copper	27	4.80E-01 (Min)	1.2E-03	1	1.00E-02 (Min)	5.7E-03
		Molybdenum						
		Nickel	27	4.80E-01 (Min)	2.2E-03			
		Selenium	27	1.80E-01 (Min)	3.4E-03	4	9.70E-04 (Min)	5.5E-04
		Silver	27	2.00E-02 (Min)	2.6E-02	4	7.40E-04 (Min)	4.2E-04
		Thallium				4	2.70E-03 (Min)	1.1E-01
		Tin				1	1.00E-02 (Min)	4.8E-05
	Organics	1,1,1-Trichloroethane				2	5.00E-03 (Min)	1.6E-04
		1,1,2-Trichloroethane				2	5.00E-03 (Min)	3.6E-03
		1,1-Dichloroethane				2	5.00E-03 (Min)	1.4E-04
		1,1-Dichloroethylene				2	5.00E-03 (Min)	1.6E-03
		1,2,4-Trichlorobenzene	6	6.60E-01 (Min)	5.8E-03	4	1.00E-02 (Min)	2.9E-03
		1,2-Dichlorobenzene	6	6.60E-01 (Min)	6.6E-04	4	9.50E-03 (Min)	3.0E-04
		2,4,5-Trichlorophenol	13	3.30E+00 (Min)	3.0E-03	2	5.00E-02 (Min)	1.4E-03
		2,4-Dichlorophenol	14	6.60E-01 (Min)	2.0E-02	4	1.00E-02 (Min)	9.5E-03
		2,4-Dimethylphenol	14	6.60E-01 (Min)	3.0E-03	4	1.00E-02 (Min)	1.4E-03
		2,4-Dinitrophenol	13	3.30E+00 (Min)	1.5E-01	2	5.00E-02 (Min)	7.1E-02
		2-Chloronaphthalene	6	6.60E-01 (Min)	7.4E-04	4	1.00E-02 (Min)	3.6E-04
		2-Chlorophenol	14	6.60E-01 (Min)	1.2E-02	4	1.00E-02 (Min)	5.7E-03
		2-Methylphenol	13	6.60E-01 (Min)	1.2E-03	2	1.00E-02 (Min)	5.7E-04
		4,4'-DDT				3	1.00E-04 (Min)	5.7E-04
		4,6-Dinitro-2-methylphenol	13	3.30E+00 (Min)	3.0E+00	4	5.00E-02 (Min)	1.4E+00
		4-Chloroaniline	6	6.60E-01 (Min)	1.5E-02	2	1.00E-02 (Min)	7.1E-03
		4-Methylphenol	14	6.60E-01 (Min)	1.2E-03	2	1.00E-02 (Min)	5.7E-04
		Acenaphthene	6	6.60E-01 (Min)	9.9E-04	4	1.00E-02 (Min)	4.8E-04
		Aldrin				3	5.00E-05 (Min)	4.8E-03
		Anthracene	6	6.60E-01 (Min)	2.0E-04	4	1.00E-02 (Min)	9.5E-05
		Benzidine				2	1.00E-02 (Min)	9.5E-03
		Benzoic acid	2	3.30E+00 (Min)	7.4E-05	2	5.00E-02 (Min)	3.6E-05
		Benzyl alcohol	6	6.60E-01 (Min)	2.0E-04	2	1.00E-02 (Min)	9.5E-05
		Bis(2-chloroisopropyl)ether	6	6.60E-01 (Min)	1.5E-03	4	1.00E-02 (Min)	7.1E-04
		Bis(2-ethylhexyl)phthalate	6	6.60E-01 (Min)	3.0E-03	4	1.00E-02 (Min)	1.4E-03
		Bromoform				2	5.00E-03 (Min)	7.1E-04
		Butylbenzylphthalate	6	6.60E-01 (Min)	3.0E-04	4	1.00E-02 (Min)	1.4E-04

Table E.18 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
4	Organics	Carbon tetrachloride				2	5.00E-03 (Min)	2.0E-02
		Chlordane				6	5.00E-04 (Min)	2.4E-02
		Chlorobenzene				2	5.00E-03 (Min)	7.1E-04
		Chlorodibromomethane				2	5.00E-03 (Min)	7.1E-04
		Chloroform				2	5.00E-03 (Min)	1.4E-03
		Cis-1,3-Dichloropropene				2	5.00E-03 (Min)	4.8E-02
		Di-n-butylphthalate	6	6.80E-01 (Min)	5.9E-04	4	1.00E-02 (Min)	2.9E-04
		Di-n-octylphthalate	6	6.80E-01 (Min)	3.0E-03	4	1.00E-02 (Min)	1.4E-03
		Dichlorobromomethane				2	5.00E-03 (Min)	7.1E-04
		Dieldrin				3	1.00E-04 (Min)	5.7E-03
		Diethylphthalate	6	6.80E-01 (Min)	7.4E-05	4	1.00E-02 (Min)	3.6E-05
		Dimethylphthalate	6	6.80E-01 (Min)	5.9E-05	4	1.00E-02 (Min)	2.9E-05
		Endosulfan I				3	5.00E-05 (Min)	2.9E-03
		Endosulfan II				3	1.00E-04 (Min)	5.7E-03
		Endrin				3	1.00E-04 (Min)	9.5E-04
		Ethylbenzene				2	5.00E-03 (Min)	1.4E-04
		Fluoranthene	7	6.80E-01 (Min)	1.5E-03	4	1.00E-02 (Min)	7.1E-04
		Fluorene	6	6.80E-01 (Min)	1.5E-03	4	1.00E-02 (Min)	7.1E-04
		Gamma-BHC (Lindane)				3	5.00E-05 (Min)	4.8E-04
		Heptachlor				3	5.00E-05 (Min)	2.9E-04
		Heptachlor epoxide				3	5.00E-05 (Min)	1.1E-02
		Hexachlorobenzene	6	6.80E-01 (Min)	7.4E-02	4	1.00E-02 (Min)	3.6E-02
		Hexachlorobutadiene	6	6.80E-01 (Min)	3.0E-02	4	1.00E-02 (Min)	1.4E-02
		Hexachlorocyclopentadiene	6	6.80E-01 (Min)	8.5E-03	4	1.00E-02 (Min)	4.1E-03
		Hexachloroethane	6	6.80E-01 (Min)	5.9E-02	4	1.00E-02 (Min)	2.9E-02
		Isophorone	6	6.80E-01 (Min)	3.0E-04	4	1.00E-02 (Min)	1.4E-04
		Methoxychlor				3	5.00E-04 (Min)	2.9E-04
		Methyl bromide				2	1.00E-02 (Min)	2.0E-02
		Methylene chloride				2	5.00E-03 (Min)	2.4E-04
		Naphthalene	6	6.80E-01 (Min)	1.5E-03	4	1.00E-02 (Min)	7.1E-04
		Nitrobenzene	6	6.80E-01 (Min)	1.2E-01	4	1.00E-02 (Min)	5.7E-02
		Pentachlorophenol	13	3.30E+00 (Min)	9.9E-03	4	5.00E-02 (Min)	4.8E-03
		Phenol	13	6.80E-01 (Min)	9.9E-05	4	5.00E-02 (Min)	4.8E-03
		Pyrene	6	6.80E-01 (Min)	2.0E-03	4	1.00E-02 (Min)	9.5E-04
		Tetrachloroethylene				2	5.00E-03 (Min)	1.4E-03
		Toluene				2	5.00E-03 (Min)	7.1E-05
		Trans-1,2-dichloroethylene				2	5.00E-03 (Min)	7.1E-04
		Trans-1,3-dichloropropene				2	5.00E-03 (Min)	4.8E-02

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
4	Organics	Trichlorofluoromethane	1			1	5.00E-03 (Min)	4.8E-05
5	Inorganics	Antimony	25	4.80E-01 (Min)	1.0E-01	4	1.90E-03 (Min)	1.4E-02
		Arsenic				4	1.30E-03 (Min)	1.2E-02
		Beryllium				4	3.90E-03 (Min)	2.2E-03
		Cadmium	25	1.80E-01 (Min)	3.4E-02	4	2.10E-04 (Min)	1.2E-03
		Chromium	25	4.60E-01 (Min)	8.3E-03	4	9.80E-03 (Min)	5.6E-03
		Copper				4	4.40E-03 (Min)	3.4E-04
		Mercury	5	4.60E-01 (Min)	2.1E-03	4	2.00E-04 (Min)	1.9E-03
		Nickel	25	4.80E-01 (Min)	8.3E-03	4	1.90E-02 (Min)	2.7E-03
		Selenium	25	1.80E-01 (Min)	3.4E-03	4	9.70E-04 (Min)	5.5E-04
		Silver	25	2.00E-02 (Min)	2.6E-02	4	7.40E-04 (Min)	4.2E-04
		Thallium	18			4	2.70E-03 (Min)	1.1E-01
		Zinc				4	3.40E-03 (Min)	4.9E-05
	Organics	1,2,4-Trichlorobenzene	9	3.30E+00 (Min)	3.0E-03	4	1.00E-02 (Min)	2.9E-03
		1,2-Dichlorobenzene				4	1.00E-02 (Min)	3.2E-04
		2,4,5-Trichlorophenol	9	6.80E-01 (Min)	2.0E-02	4	5.00E-02 (Min)	1.4E-03
		2,4-Dichlorophenol	9	6.80E-01 (Min)	3.0E-03	4	1.00E-02 (Min)	9.5E-03
		2,4-Dimethylphenol	9	3.30E+00 (Min)	1.5E-01	4	1.00E-02 (Min)	1.4E-03
		2,4-Dinitrophenol				4	5.00E-02 (Min)	7.1E-02
		2-Chloronaphthalene	9	6.80E-01 (Min)	1.2E-02	4	1.00E-02 (Min)	3.6E-04
		2-Chlorophenol	9	6.80E-01 (Min)	1.2E-03	4	1.00E-02 (Min)	5.7E-03
		2-Methylphenol	8	3.30E+00 (Min)	3.0E+00	3	1.00E-04 (Min)	5.7E-04
		4,4'-DDT	9	6.80E-01 (Min)	1.2E-03	4	5.00E-02 (Min)	1.4E+00
		4,6-Dinitro-2-methylphenol				4	1.00E-02 (Min)	7.1E-03
		4-Chloroaniline	9	6.80E-01 (Min)	1.2E-03	4	1.00E-02 (Min)	5.7E-04
		4-Methylphenol				4	1.00E-02 (Min)	5.7E-04
		Acenaphthene				4	1.00E-05 (Min)	4.8E-04
		Aldrin	3			3	5.00E-05 (Min)	4.8E-03
		Anthracene	4			4	1.00E-02 (Min)	9.5E-05
		Benzoic acid	4			4	5.00E-02 (Min)	3.6E-05
		Benzyl alcohol	4			4	1.00E-02 (Min)	9.5E-05

Table E.18 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
5	Organics	Bis(2-chloroisopropyl)ether				4	1.00E-02 (Min)	7.1E-04
		Bis(2-ethylhexyl)phthalate				4	3.00E-03 (Min)	4.3E-04
		Butylbenzylphthalate				4	1.00E-02 (Min)	1.4E-04
		Chlordane				6	5.00E-04 (Min)	2.4E-02
		Di-n-butylphthalate				4	1.00E-02 (Min)	2.9E-04
		Di-n-octylphthalate				4	1.00E-02 (Min)	1.4E-03
		Dieldrin				3	1.00E-04 (Min)	5.7E-03
		Diethylphthalate				4	1.00E-02 (Min)	3.6E-05
		Dimethylphthalate				4	1.00E-02 (Min)	2.9E-05
		Endosulfan I				3	5.00E-05 (Min)	2.9E-03
		Endosulfan II				3	1.00E-04 (Min)	5.7E-03
		Endrin				3	1.00E-04 (Min)	9.5E-04
		Fluoranthene				4	1.00E-02 (Min)	7.1E-04
		Fluorene				4	1.00E-02 (Min)	7.1E-04
		Gamma-BHC (Lindane)				3	5.00E-05 (Min)	4.8E-04
		Heptachlor				3	5.00E-05 (Min)	2.9E-04
		Heptachlor epoxide				3	5.00E-05 (Min)	1.1E-02
		Hexachlorobenzene				3	1.00E-02 (Min)	3.6E-02
		Hexachlorobutadiene				4	1.00E-02 (Min)	1.4E-02
		Hexachlorocyclopentadiene				4	1.00E-02 (Min)	4.1E-03
		Hexachloroethane				4	1.00E-02 (Min)	2.9E-02
		Isophorone				4	1.00E-02 (Min)	1.4E-04
		Methoxychlor				3	5.00E-04 (Min)	2.9E-04
		Naphthalene				4	1.00E-02 (Min)	7.1E-04
		Nitrobenzene				4	1.00E-02 (Min)	5.7E-02
		Pentachlorophenol				4	5.00E-02 (Min)	4.8E-03
		Phenol				4	1.00E-02 (Min)	4.8E-05
		Pyrene				4	1.00E-02 (Min)	9.5E-04
			1	5.70E-02 (Min)	1.3E-04			
			9	3.30E+00 (Min)	9.9E-03			
			9	6.60E-01 (Min)	9.9E-05			
10	Inorganics			mg/kg			mg/L	
		Antimony	15	3.50E-01 (Min)	7.9E-02	1	1.90E-03 (Min)	1.4E-02
		Arsenic				1	1.30E-03 (Min)	1.2E-02
		Beryllium	15	1.40E-01 (Min)	2.5E-02	1	3.90E-03 (Min)	2.2E-03
		Cadmium	15	3.50E-01 (Min)	6.3E-03	1	2.10E-04 (Min)	1.2E-03
		Chromium	15	3.50E-01 (Min)	8.5E-04	1	9.80E-03 (Min)	5.6E-03
		Copper				1	4.40E-03 (Min)	3.4E-04

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
10	Inorganics	Mercury	15	3.50E-01 (Min)	1.6E-03	1	2.00E-04 (Min)	1.9E-03
		Nickel				1	1.90E-02 (Min)	2.7E-03
		Selenium				1	9.70E-04 (Min)	5.5E-04
		Silver	15	1.40E-01 (Min)	2.5E-03	1	7.40E-04 (Min)	4.2E-04
		Thallium	15	2.00E-02 (Min)	2.6E-02	1	2.70E-03 (Min)	1.1E-01
		Zinc				1	1.05E-02 (Min)	1.5E-04
<hr/>								
			ag/kg		ag/L			
Organics		1,2,4-Trichlorobenzene	8	1.00E-02 (Min)	1.8E-03	1	1.00E-02 (Min)	2.9E-03
		1,2-Dichlorobenzene				1	1.00E-02 (Min)	3.2E-04
		2,4-Dichlorophenol				1	1.00E-02 (Min)	9.5E-03
		2,4-Dimethylphenol				1	1.00E-02 (Min)	1.4E-03
		2,4-Dinitrophenol				1	5.00E-02 (Min)	7.1E-02
		2-Chloronaphthalene				1	1.00E-02 (Min)	3.6E-04
		2-Chlorophenol				1	1.00E-02 (Min)	5.7E-03
		2-Methylphenol				1	1.00E-02 (Min)	5.7E-04
		4,4'-DDT				1	1.00E-04 (Min)	5.7E-04
		4,6-Dinitro-2-methylphenol	8	1.00E-02 (Min)	1.8E-03	1	1.00E-02 (Min)	1.4E+00
		4-Chloroaniline				1	1.00E-02 (Min)	7.1E-03
		4-Methylphenol				1	1.00E-02 (Min)	5.7E-04
		Acenaphthene				1	1.00E-02 (Min)	4.8E-04
		Aldrin	8	1.00E-02 (Min)	3.0E-02	1	5.00E-05 (Min)	4.8E-03
		Anthracene				1	1.00E-02 (Min)	9.5E-05
		Benzoic acid				1	5.00E-02 (Min)	3.6E-05
		Benzyl alcohol				1	1.00E-02 (Min)	9.5E-05
		Bis(2-chloroisopropyl)ether				1	1.00E-02 (Min)	7.1E-04
		Butylbenzylphthalate				1	1.00E-02 (Min)	1.4E-04
		Chlordane				2	5.00E-04 (Min)	2.4E-02
		Di-n-butylphthalate				1	1.00E-02 (Min)	2.9E-04
		Di-n-octylphthalate				1	1.00E-02 (Min)	1.4E-03
		Dieldrin				1	1.00E-04 (Min)	5.7E-03
		Diethylphthalate				1	1.00E-02 (Min)	3.6E-05
		Diethylphthalate				1	1.00E-02 (Min)	2.9E-05
		Endosulfan I				1	5.00E-05 (Min)	2.9E-03
		Endosulfan II				1	1.00E-04 (Min)	5.7E-03
		Endrin				1	1.00E-04 (Min)	9.5E-04
		Fluoranthene				1	1.00E-02 (Min)	7.1E-04

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish ingest.			Water ingest.		
			Fish samples	Fish concentration	Fish carcinogen screening index	Water samples	Water concentration	Water carcinogen screening index
10	Organics	Fluorene				1	1.00E-02 (Min)	7.1E-04
		Gamma-BHC (Lindane)				1	5.00E-05 (Min)	4.8E-04
		Heptachlor				1	5.00E-05 (Min)	2.9E-04
		Heptachlor epoxide				1	5.00E-05 (Min)	1.1E-02
		Hexachlorobenzene				1	1.00E-02 (Min)	3.6E-02
		Hexachlorobutadiene				1	1.00E-02 (Min)	1.4E-02
		Hexachlorocyclopentadiene				1	1.00E-02 (Min)	4.1E-03
		Hexachloroethane				1	1.00E-02 (Min)	2.9E-02
		Isophorone				1	1.00E-02 (Min)	1.4E-04
		Methoxychlor				1	5.00E-04 (Min)	2.9E-04
		Naphthalene				1	1.00E-02 (Min)	7.1E-04
		Nitrobenzene				1	5.00E-02 (Min)	5.7E-02
		Pentachlorophenol				1	5.00E-02 (Min)	4.8E-03
		Phenol				1	1.00E-02 (Min)	4.8E-05
		Pyrene				1	1.00E-02 (Min)	9.5E-04
13	Inorganics			mg/kg			mg/L	
		Antimony	10	4.80E-01 (Min)	1.0E-01	1	1.90E-03 (Min)	1.4E-02
		Arsenic				2	1.30E-03 (Min)	1.2E-02
		Beryllium	3	3.00E-03 (Min)	5.4E-05	1	3.90E-03 (Min)	2.2E-03
		Cadmium	10	1.80E-01 (Min)	3.2E-02	2	2.10E-04 (Min)	1.2E-03
		Chromium	10	4.50E-01 (Min)	8.1E-03			
		Copper	10	4.50E-01 (Min)	1.1E-03	2	4.00E-03 (Min)	3.1E-04
		Mercury				2	2.00E-04 (Min)	1.9E-03
		Nickel	10	4.50E-01 (Min)	2.0E-03			
		Selenium	10	4.50E-01 (Min)	8.1E-03	1	9.70E-04 (Min)	5.5E-04
		Silver	10	1.80E-01 (Min)	3.2E-03	1	7.40E-04 (Min)	4.2E-04
		Thallium	10	2.00E-02 (Min)	2.6E-02	1	2.70E-03 (Min)	1.1E-01
		Uranium				1	6.67E-04 (Min)	6.3E-04
				mg/kg			mg/L	
		Organics	1,1,1-Trichloroethane	1			1	5.00E-03 (Min)
1,1,2-Trichloroethane	1				1	5.00E-03 (Min)	3.6E-03	
1,1-Dichloroethane	1				1	5.00E-03 (Min)	1.4E-04	
1,1-Dichloroethylene	1				1	5.00E-03 (Min)	1.6E-03	

Table E.18 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
13	Organics	1,2,4-Trichlorobenzene				2	1.00E-02 (Min)	2.9E-03
		1,2-Dichlorobenzene				2	1.00E-02 (Min)	3.2E-04
		2,4,5-Trichlorophenol				1	5.00E-02 (Min)	1.4E-03
		2,4-Dichlorophenol	2	3.40E+00 (Min)	3.1E-03	2	1.00E-02 (Min)	9.5E-03
		2,4-Dimethylphenol	2	6.90E-01 (Min)	2.1E-02	2	1.00E-02 (Min)	1.4E-03
		2,4-Dinitrophenol	2	6.90E-01 (Min)	3.1E-03	1	5.00E-02 (Min)	7.1E-02
		2-Chloronaphthalene	2	3.40E+00 (Min)	1.5E-01	2	1.00E-02 (Min)	3.6E-04
		2-Chlorophenol	2	6.90E-01 (Min)	1.2E-02	2	1.00E-02 (Min)	5.7E-03
		2-Methylphenol	2	6.90E-01 (Min)	1.2E-03	1	1.00E-02 (Min)	5.7E-04
		4,4'-DDT	3	1.00E-02 (Min)	1.8E-03			
		4,6-Dinitro-2-methylphenol	2	3.40E+00 (Min)	3.1E+00	2	5.00E-02 (Min)	1.4E+00
		4-Chloroaniline				1	1.00E-02 (Min)	7.1E-03
		4-Methylphenol				1	1.00E-02 (Min)	5.7E-04
		Acenaphthene	2	8.90E-01 (Min)	1.2E-03	2	1.00E-02 (Min)	4.8E-04
		Anthracene				2	1.00E-02 (Min)	9.5E-05
		Benzidine				1	1.00E-02 (Min)	9.5E-03
		Benzoic acid				1	5.00E-02 (Min)	3.6E-05
		Benzyl alcohol				1	1.00E-02 (Min)	9.5E-05
		Bis(2-chloroisopropyl)ether				2	1.00E-02 (Min)	7.1E-04
		Bis(2-ethylhexyl)phthalate				2	1.00E-02 (Min)	1.4E-03
		Bromoform				1	5.00E-03 (Min)	7.1E-04
		Butylbenzylphthalate				2	1.00E-02 (Min)	1.4E-04
		Carbon tetrachloride				1	5.00E-03 (Min)	2.0E-02
		Chlorobenzene				1	5.00E-03 (Min)	7.1E-04
		Chlorodibromomethane				1	5.00E-03 (Min)	1.4E-03
		Chloroform				1	5.00E-03 (Min)	4.8E-02
		Cis-1,3-Dichloropropene				2	1.00E-02 (Min)	2.9E-04
		Di-n-butylphthalate				2	1.00E-02 (Min)	1.4E-03
		Di-n-octylphthalate				2	1.00E-02 (Min)	7.1E-04
		Dichlorobromomethane				2	5.00E-03 (Min)	3.6E-05
		Diethylphthalate				2	1.00E-02 (Min)	2.9E-05
		Dimethylphthalate				1	5.00E-03 (Min)	1.4E-04
		Ethylbenzene				2	1.00E-02 (Min)	7.1E-04
		Fluoranthene				2	1.00E-02 (Min)	7.1E-04
		Fluorene				2	1.00E-02 (Min)	3.6E-02
		Hexachlorobenzene				2	1.00E-02 (Min)	1.4E-02
		Hexachlorobutadiene				2	1.00E-02 (Min)	4.1E-03
		Hexachlorocyclopentadiene				2	1.00E-02 (Min)	

Table E.16 Results of adult nonconservative screening for noncarcinogens in fish and water where no measurement in a reach was above detection limits.

Reach	Contaminant type	Contaminant	Fish samples	Fish concentration	Fish ingest. carcinogen screening index	Water samples	Water concentration	Water ingest. carcinogen screening index
13	Organics	Hexachloroethane				2	1.00E-02 (Min)	2.9E-02
		Isophorone				2	1.00E-02 (Min)	1.4E-04
		Methyl bromide				1	1.00E-02 (Min)	2.0E-02
		Methylene chloride				1	5.00E-03 (Min)	2.4E-04
		Naphthalene				2	1.00E-02 (Min)	7.1E-04
		Nitrobenzene				2	1.00E-02 (Min)	5.7E-02
		Pentachlorophenol	2	3.40E+00 (Min)	1.0E-02	2	5.00E-02 (Min)	4.8E-03
		Phenol	2	8.90E-01 (Min)	1.0E-04	2	1.00E-02 (Min)	4.8E-05
		Pyrene				2	1.00E-02 (Min)	9.5E-04
		Tetrachloroethylene				1	5.00E-03 (Min)	1.4E-03
		Toluene				1	5.00E-03 (Min)	7.1E-05
		Trans-1,2-dichloroethylene				1	5.00E-03 (Min)	7.1E-04
		Trans-1,3-dichloropropene				1	5.00E-03 (Min)	4.8E-02

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
1	Sej. Ingestion	Organics			mg/kg	1/(mg/kg/d)	mg/kg/d	
			Aroclor-1260	23	1.20E+00 (Min)	7.7E+00	7.3E-08	5.6E-07
			Aroclor-1254	23	1.20E+00 (Min)	7.7E+00	7.3E-08	5.6E-07
			Dibenz(a,h)anthracene	25	5.16E-01 (Min)	1.2E+01	3.2E-08	3.6E-07
			Acenaphthene	25	5.16E-01 (Min)	1.2E+01	3.2E-08	3.6E-07
			Fluorene	25	5.16E-01 (Min)	1.2E+01	3.2E-08	3.6E-07
			Acenaphthylene	25	5.16E-01 (Min)	1.2E+01	3.2E-08	3.6E-07
			Benzo(a)pyrene	25	5.16E-01 (Min)	7.3E+00	3.2E-08	2.3E-07
			Phenanthrene	25	5.16E-01 (Min)	7.3E+00	3.2E-08	2.3E-07
			Anthracene	25	5.16E-01 (Min)	7.3E+00	3.2E-08	2.3E-07
			Benzo(g,h,i)perylene	25	5.16E-01 (Min)	7.0E+00	3.2E-08	2.2E-07
			N-Nitroso-di-n-propylamine	25	5.16E-01 (Min)	1.6E+01	7.3E-09	1.2E-07
			Dieldrin	23	1.20E-01 (Min)	1.1E+00	7.3E-08	8.1E-08
			Toxaphene	23	1.20E+00 (Min)	1.7E+01	3.7E-09	6.2E-08
			Aldrin	23	5.98E-02 (Min)	1.6E+00	3.2E-08	5.1E-08
			Hexachlorobenzene	25	5.16E-01 (Min)	1.3E+00	3.7E-08	4.8E-08
			Chlordane	23	5.98E-01 (Min)	1.1E+00	3.2E-08	3.5E-08
			Bis(2-chloroethyl)ether	25	5.16E-01 (Min)	9.1E+00	3.7E-09	3.3E-08
			Heptachlor epoxide	25	5.98E-02 (Min)	4.5E-01	6.3E-08	2.9E-08
			3,3'-Dichlorobenzidine	25	1.04E+00 (Min)	7.3E-01	3.2E-08	2.3E-08
			Indeno(1,2,3-cd)pyrene	25	5.16E-01 (Min)	7.3E-01	3.2E-08	2.3E-08
			Benzo(b)fluoranthene	25	5.16E-01 (Min)	6.3E+00	3.7E-09	2.3E-08
			Alpha-BHC	23	5.98E-02 (Min)	6.8E-01	3.2E-08	2.1E-08
			2,4-Dinitrotoluene	25	5.16E-01 (Min)	6.8E-01	3.2E-08	2.1E-08
			2,6-Dinitrotoluene	25	5.98E-02 (Min)	1.2E-01	1.5E-07	1.8E-08
			Pentachlorophenol	23	5.98E-02 (Min)	4.5E+00	3.7E-09	1.6E-08
			Heptachlor	23	5.98E-02 (Min)	1.8E+00	3.7E-09	6.6E-09
			Beta-BHC	23	5.98E-02 (Min)	1.3E+00	3.7E-09	4.8E-09
			Gamma-BHC (Lindane)	23	5.98E-02 (Min)	3.4E-01	7.3E-09	2.5E-09
			4,4'-DDE	23	1.20E-01 (Min)	7.8E-02	3.2E-08	2.5E-09
			Hexachlorobutadiene	25	5.16E-01 (Min)	7.0E-02	3.2E-08	2.2E-09
			Bis(2-chloroisopropyl)ether	25	5.16E-01 (Min)	3.5E-02	3.2E-08	1.1E-09
			4-Chloroaniline	25	5.16E-01 (Min)	2.4E-02	3.2E-08	7.6E-10
			1,4-Dichlorobenzene	25	7.48E-01 (Min)	1.4E-02	4.6E-08	6.4E-10
			Bis(2-ethylhexyl)phthalate	25	5.16E-01 (Min)	1.4E-02	3.2E-08	4.4E-10
			Hexachloroethane	25	5.16E-01 (Min)	1.1E-02	3.2E-08	3.5E-10
			2,4,6-Trichlorophenol	25	5.16E-01 (Min)	4.9E-03	3.2E-08	1.5E-10
			N-Nitrosodiphenylamine	25	5.16E-01 (Min)			

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment
where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
1	Sed. Ingestion	Organics	Isophorone	25	5.18E-01 (Min)	4.1E-03	3.2E-08	1.3E-10
2	Sed. Ingestion	Organics	Dibenz(a,h)anthracene	40	5.00E-01 (Min)	1.2E+01	3.1E-08	3.5E-07
			Acenaphthene	40	5.00E-01 (Min)	1.2E+01	3.1E-08	3.5E-07
			Fluorene	40	5.00E-01 (Min)	1.2E+01	3.1E-08	3.5E-07
			Acenaphthylene	40	5.00E-01 (Min)	1.2E+01	3.1E-08	3.5E-07
			Anthracene	40	5.00E-01 (Min)	7.3E+00	3.1E-08	2.2E-07
			Benzo(g,h,i)perylene	40	5.00E-01 (Min)	7.0E+00	3.1E-08	2.1E-07
			N-Nitroso-di-n-propylamine	40	5.00E-01 (Min)	1.6E+01	1.2E-08	1.9E-07
			Dieldrin	41	1.90E-01 (Min)	1.7E+01	5.8E-09	9.8E-08
			Aldrin	41	9.42E-02 (Min)	1.3E+00	5.7E-08	7.5E-08
			Chlordane	41	9.39E-01 (Min)	1.3E+00	5.8E-09	5.2E-08
			Heptachlor epoxide	41	9.42E-02 (Min)	9.1E+00	3.1E-08	4.9E-08
			Hexachlorobenzene	40	5.00E-01 (Min)	1.6E+00	5.8E-09	3.6E-08
			Alpha-BHC	41	9.42E-02 (Min)	6.3E+00	3.1E-08	3.4E-08
			Bis(2-chloroethyl)ether	40	5.00E-01 (Min)	1.1E+00	6.1E-08	2.7E-08
			3,3'-Dichlorobenzidine	40	9.97E-01 (Min)	4.5E-01	5.8E-09	2.6E-08
			Heptachlor	41	9.42E-02 (Min)	4.5E+00	3.1E-08	2.2E-08
			Benzo(k)fluoranthene	40	5.00E-01 (Min)	7.3E-01	3.1E-08	2.1E-08
			2,4-Dinitrotoluene	40	5.00E-01 (Min)	6.8E-01	3.1E-08	2.1E-08
			2,6-Dinitrotoluene	40	5.00E-01 (Min)	6.8E-01	3.1E-08	2.1E-08
			Pentachlorophenol	40	2.43E+00 (Min)	1.2E-01	1.5E-07	1.8E-08
			Beta-BHC	41	9.42E-02 (Min)	1.8E+00	5.8E-09	1.0E-08
			Gamma-BHC (Lindane)	41	9.42E-02 (Min)	1.3E+00	5.8E-09	7.5E-09
			4,4'-DDT	41	1.90E-01 (Min)	3.4E-01	1.2E-08	3.9E-09
			4,4'-DDE	41	1.90E-01 (Min)	3.4E-01	1.2E-08	3.9E-09
			4,4'-DDD	41	1.90E-01 (Min)	2.4E-01	1.2E-08	2.8E-09
			Hexachlorobutadiene	40	5.00E-01 (Min)	7.8E-02	3.1E-08	2.4E-09
			Bis(2-chloroisopropyl)ether	40	5.00E-01 (Min)	7.0E-02	3.1E-08	2.1E-09
			4-Chloroaniline	40	5.00E-01 (Min)	3.5E-02	3.1E-08	1.1E-09
			1,4-Dichlorobenzene	40	5.00E-01 (Min)	2.4E-02	3.1E-08	7.3E-10
			Hexachloroethane	40	5.00E-01 (Min)	1.4E-02	3.1E-08	4.3E-10
			2,4,6-Trichlorophenol	40	5.00E-01 (Min)	1.1E-02	3.1E-08	3.4E-10
			N-Nitrosodiphenylamine	40	5.00E-01 (Min)	4.9E-03	3.1E-08	1.5E-10
			Isophorone	40	5.00E-01 (Min)	4.1E-03	3.1E-08	1.3E-10

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
3	Sed. Ingestion	Organics	Aroclor-1260	49	9.38E-01 (Min)	7.7E+00	5.7E-08	4.4E-07
			Dibenz(a,h)anthracene	50	4.17E-01 (Min)	1.2E+01	2.6E-08	2.9E-07
			Acenaphthene	50	4.17E-01 (Min)	1.2E+01	2.6E-08	2.9E-07
			Fluorene	50	4.17E-01 (Min)	1.2E+01	2.6E-08	2.9E-07
			Acenaphthylene	50	4.17E-01 (Min)	1.2E+01	2.6E-08	2.9E-07
			Benzo(g,h,i)perylene	50	4.17E-01 (Min)	7.3E+00	2.6E-08	1.9E-07
			N-Nitroso-dl-n-propylamine	50	4.17E-01 (Min)	7.0E+00	2.6E-08	1.8E-07
			Dieldrin	49	9.38E-02 (Min)	1.6E+01	5.7E-09	9.2E-08
			Toxaphene	49	9.38E-01 (Min)	1.1E+00	5.7E-08	6.3E-08
			Aldrin	49	4.62E-02 (Min)	1.7E+01	2.8E-09	4.8E-08
			Hexachlorobenzene	50	4.17E-01 (Min)	1.6E+00	2.6E-08	4.1E-08
			Chlordane	49	4.62E-01 (Min)	1.3E+00	2.8E-08	3.7E-08
			Bis(2-chloroethyl)ether	50	4.17E-01 (Min)	1.1E+00	2.6E-08	2.8E-08
			Heptachlor epoxide	49	4.62E-02 (Min)	9.1E+00	2.8E-09	2.6E-08
			3,3'-Dichlorobenzidine	50	8.34E-01 (Min)	4.5E-01	5.1E-08	2.3E-08
			Indeno(1,2,3-cd)pyrene	50	4.17E-01 (Min)	7.3E-01	2.6E-08	1.9E-08
			Alpha-BHC	48	4.62E-02 (Min)	6.3E+00	2.8E-09	1.8E-08
			2,4-Dinitrotoluene	50	4.17E-01 (Min)	6.8E-01	2.6E-08	1.7E-08
			2,6-Dinitrotoluene	50	2.02E+00 (Min)	1.2E-01	2.6E-08	1.5E-08
			Pentachlorophenol	49	4.62E-02 (Min)	4.5E+00	2.8E-09	1.3E-08
			Heptachlor	49	4.62E-02 (Min)	1.8E+00	2.8E-09	5.1E-09
			Beta-BHC	49	4.62E-02 (Min)	1.3E+00	2.8E-09	3.7E-09
			Gamma-BHC (Lindane)	49	4.62E-02 (Min)	7.8E-02	2.8E-08	2.0E-09
			Hexachlorobutadiene	50	4.17E-01 (Min)	3.4E-01	5.7E-09	2.0E-09
			3,4'-DDT	49	9.38E-02 (Min)	3.4E-01	5.7E-09	2.0E-09
			4,4'-DDE	50	4.17E-01 (Min)	7.0E-02	2.6E-08	1.8E-09
			Bis(2-chloroisopropyl)ether	49	9.38E-02 (Min)	2.4E-01	5.7E-09	1.4E-09
			4,4'-DDD	50	4.17E-01 (Min)	3.5E-02	2.6E-08	8.9E-10
			4-Chloroaniline	50	4.17E-01 (Min)	2.4E-02	2.6E-08	6.1E-10
			1,4-Dichlorobenzene	50	4.17E-01 (Min)	1.4E-02	2.6E-08	3.6E-10
			Hexachloroethane	50	4.17E-01 (Min)	1.1E-02	2.6E-08	2.8E-10
			2,4,6-Trichlorophenol	50	4.17E-01 (Min)	4.9E-03	2.6E-08	1.3E-10
			N-Nitrosodiphenylamine	50	4.17E-01 (Min)	4.1E-03	2.6E-08	1.0E-10
			Isophorone	50	4.17E-01 (Min)	4.1E-03	2.6E-08	1.0E-10

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
4	Sed. Ingestion	Organics	Aroclor-1260	45	1.91E+00 (Min)	7.7E+00	1.2E-07	9.0E-07
			Dibenz(a,h)anthracene	47	4.97E-01 (Min)	1.2E+01	3.0E-08	3.5E-07
			Acenaphthene	47	4.97E-01 (Min)	1.2E+01	3.0E-08	3.5E-07
			Fluorene	47	4.97E-01 (Min)	1.2E+01	3.0E-08	3.5E-07
			Acenaphthylene	47	4.97E-01 (Min)	1.2E+01	3.0E-08	3.5E-07
			Benzo(a)pyrene	47	4.97E-01 (Min)	7.3E+00	3.0E-08	2.2E-07
			Anthracene	47	4.97E-01 (Min)	7.3E+00	3.0E-08	2.2E-07
			Benzo(g,h,i)perylene	47	4.97E-01 (Min)	7.3E+00	3.0E-08	2.2E-07
			Dieldrin	45	1.91E-01 (Min)	1.6E+01	1.2E-08	1.9E-07
			Toxaphene	44	1.91E+00 (Min)	1.1E+00	1.2E-07	1.3E-07
			Aldrin	44	9.54E-02 (Min)	1.7E+01	5.8E-09	9.9E-08
			Chlordane	44	9.54E-01 (Min)	1.3E+00	5.8E-09	7.6E-08
			Heptachlor epoxide	44	9.54E-02 (Min)	9.1E+00	5.8E-09	5.3E-08
			Hexachlorobenzene	47	4.97E-01 (Min)	1.6E+00	3.0E-08	4.9E-08
			Alpha-BHC	44	9.54E-02 (Min)	6.3E+00	5.8E-09	3.7E-08
			Bis(2-chloroethyl)ether	47	4.97E-01 (Min)	1.1E+00	3.0E-08	3.3E-08
			3,3'-dichlorobenzidine	47	9.90E-01 (Min)	4.5E-01	6.1E-08	2.7E-08
			Heptachlor	44	4.97E-01 (Min)	4.5E+00	3.0E-08	2.6E-08
			Benzo(a)anthracene	47	4.97E-01 (Min)	7.3E-01	3.0E-08	2.2E-08
			Indeno(1,2,3-cd)pyrene	47	4.97E-01 (Min)	7.3E-01	3.0E-08	2.2E-08
			Benzo(b)fluoranthene	47	4.97E-01 (Min)	7.3E-01	3.0E-08	2.2E-08
			Benzo(k)fluoranthene	47	4.97E-01 (Min)	6.8E-01	3.0E-08	2.1E-08
			2,4-Dinitrotoluene	47	4.97E-01 (Min)	6.8E-01	3.0E-08	2.1E-08
			2,6-Dinitrotoluene	47	4.97E-01 (Min)	1.2E-01	1.5E-07	1.8E-08
			Pentachlorophenol	44	2.40E+00 (Min)	1.8E+00	5.8E-09	1.1E-08
			Beta-BHC	44	9.54E-02 (Min)	1.3E+00	5.8E-09	7.6E-09
			Gamma-BHC (Lindane)	44	9.54E-02 (Min)	1.3E+00	5.8E-09	7.6E-09
			4,4'-DDT	45	1.91E-01 (Min)	3.4E-01	1.2E-08	4.0E-09
			4,4'-DDE	45	1.91E-01 (Min)	3.4E-01	1.2E-08	4.0E-09
			4,4'-DDD	45	1.91E-01 (Min)	2.4E-01	1.2E-08	2.8E-09
			Hexachlorobutadiene	47	4.97E-01 (Min)	7.8E-02	3.0E-08	2.4E-09
			Chrysene	47	4.97E-01 (Min)	7.3E-02	3.0E-08	2.2E-09
			Bis(2-chloroisopropyl)ether	47	4.97E-01 (Min)	7.0E-02	3.0E-08	2.1E-09
			4-Chloroaniline	47	4.97E-01 (Min)	3.5E-02	3.0E-08	1.1E-09

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
4	Sed. Ingestion	Organics	1,4-Dichlorobenzene	47	4.97E-01 (Min)	2.4E-02	3.0E-08	7.3E-10
			Hexachloroethane	47	4.97E-01 (Min)	1.4E-02	3.0E-08	4.3E-10
			2,4,6-Trichlorophenol	47	4.97E-01 (Min)	1.1E-02	3.0E-08	3.3E-10
			N-Nitrosodiphenylamine	47	4.97E-01 (Min)	4.9E-03	3.0E-08	1.5E-10
			Isophorone	47	4.97E-01 (Min)	4.1E-03	3.0E-08	1.2E-10
5	Sed. Ingestion	Organics	Aroclor-1260	62	7.12E+00 (Min)	7.7E+00	4.4E-07	3.4E-06
			Aroclor-1254	62	7.12E+00 (Min)	7.7E+00	4.4E-07	3.4E-06
			Dieldrin	62	7.12E-01 (Min)	1.6E+01	4.4E-08	7.0E-07
			Dibenz(a,h)anthracene	62	7.25E-01 (Min)	1.2E+01	4.4E-08	5.1E-07
			Acenaphthene	62	7.25E-01 (Min)	1.2E+01	4.4E-08	5.1E-07
			Fluorene	62	7.25E-01 (Min)	1.2E+01	4.4E-08	5.1E-07
			Acenaphthylene	62	7.25E-01 (Min)	1.2E+01	4.4E-08	5.1E-07
			Toxaphene	62	7.12E+00 (Min)	1.1E+00	4.4E-07	4.8E-07
			Aldrin	42	3.55E-01 (Min)	1.7E+01	2.2E-08	3.7E-07
			Benzo(a)pyrene	62	7.25E-01 (Min)	7.3E+00	4.4E-08	3.2E-07
			Phenanthrene	62	7.25E-01 (Min)	7.3E+00	4.4E-08	3.2E-07
			Anthracene	62	7.25E-01 (Min)	7.3E+00	4.4E-08	3.2E-07
			Pyrene	62	7.25E-01 (Min)	7.3E+00	4.4E-08	3.2E-07
			Benzo(g,h,i)perylene	62	7.25E-01 (Min)	7.3E+00	4.4E-08	3.2E-07
			Fluoranthene	62	7.25E-01 (Min)	7.3E+00	4.4E-08	3.2E-07
			N-Nitroso-di-n-propylamine	62	7.25E-01 (Min)	7.0E+00	4.4E-08	3.1E-07
			Chlordane	42	3.55E+00 (Min)	1.3E+00	2.2E-07	2.8E-07
			Heptachlor epoxide	42	3.55E-01 (Min)	9.1E+00	2.2E-08	2.0E-07
			Alpha-BHC	42	3.55E-01 (Min)	6.3E+00	2.2E-08	1.4E-07
			Heptachlor	42	3.55E-01 (Min)	4.5E+00	2.2E-08	9.8E-08
			Hexachlorobenzene	62	7.25E-01 (Min)	1.6E+00	4.4E-08	7.1E-08
			Bis(2-chloroethyl)ether	62	7.25E-01 (Min)	1.1E+00	4.4E-08	4.9E-08
			3,3'-dichlorobenzidine	62	1.44E+00 (Min)	4.5E-01	8.8E-08	4.0E-08
			Beta-BHC	42	3.55E-01 (Min)	1.8E+00	2.2E-08	3.9E-08
			Benzo(a)anthracene	62	7.25E-01 (Min)	7.3E-01	4.4E-08	3.2E-08
			Indeno(1,2,3-cd)pyrene	62	7.25E-01 (Min)	7.3E-01	4.4E-08	3.2E-08
			Benzo(b)fluoranthene	62	7.25E-01 (Min)	7.3E-01	4.4E-08	3.2E-08
			Benzo(k)fluoranthene	62	7.25E-01 (Min)	7.3E-01	4.4E-08	3.2E-08
			2,4-Dinitrotoluene	62	7.25E-01 (Min)	6.8E-01	4.4E-08	3.0E-08

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
5	Sed. Ingestion	Organics	2,6-Dinitrotoluene	62	7.25E-01 (Min)	6.8E-01	4.4E-08	3.0E-08
			Gamma-BHC (Lindane)	42	3.55E-01 (Min)	1.3E+00	2.2E-08	2.8E-08
			Pentachlorophenol	62	3.54E+00 (Min)	1.2E-01	2.2E-07	2.6E-08
			4,4'-DDT	62	7.12E-01 (Min)	3.4E-01	4.4E-08	1.5E-08
			4,4'-DDE	62	7.12E-01 (Min)	3.4E-01	4.4E-08	1.5E-08
			4,4'-DDD	62	7.12E-01 (Min)	2.4E-01	4.4E-08	1.0E-08
			Hexachlorobutadiene	62	7.25E-01 (Min)	7.8E-02	4.4E-08	3.5E-09
			Chrysene	62	7.25E-01 (Min)	7.3E-02	4.4E-08	3.2E-09
			Bis(2-chloroisopropyl)ether	62	7.25E-01 (Min)	7.0E-02	4.4E-08	3.1E-09
			4-Chloroaniline	62	7.25E-01 (Min)	3.5E-02	4.4E-08	1.6E-09
			1,4-Dichlorobenzene	62	7.25E-01 (Min)	2.4E-02	4.4E-08	1.1E-09
			Hexachloroethane	62	7.25E-01 (Min)	1.4E-02	4.4E-08	6.2E-10
			2,4,6-Trichlorophenol	62	7.25E-01 (Min)	1.1E-02	4.4E-08	4.9E-10
			N-Nitrosodiphenylamine	62	7.25E-01 (Min)	4.9E-03	4.4E-08	2.2E-10
			Isophorone	62	7.25E-01 (Min)	4.1E-03	4.4E-08	1.8E-10
10	Sed. Exposure	Radionuclides	Co-60	21	6.35E+00 (Min)	1/Bq	Bq/yr	1.2E-07
	Sed. Ingestion	Radionuclides	Co-60	21	6.35E+00 (Min)	1/Bq	Bq/Life	2.9E-10
	Sed. Ingestion	Organics	Aroclor-1260	22	1.68E+01 (Min)	7.7E+00	1.0E-06	7.9E-06
			Aroclor-1254	22	1.68E+01 (Min)	7.7E+00	1.0E-06	7.9E-06
			Dibenz(a,h)anthracene	22	4.43E+00 (Min)	1.2E+01	2.7E-07	3.1E-06
			Acenaphthene	22	4.43E+00 (Min)	1.2E+01	2.7E-07	3.1E-06
			Fluorene	22	4.43E+00 (Min)	1.2E+01	2.7E-07	3.1E-06
			Acenaphthylene	22	4.43E+00 (Min)	1.2E+01	2.7E-07	3.1E-06
			Benzo(a)pyrene	22	4.43E+00 (Min)	7.3E+00	2.7E-07	2.0E-06
			Benzo(g,h,i)perylene	22	4.43E+00 (Min)	7.3E+00	2.7E-07	2.0E-06
			N-Nitroso-di-n-propylamine	22	4.43E+00 (Min)	7.0E+00	2.7E-07	1.9E-06
			Dieldrin	22	1.68E+00 (Min)	1.6E+01	1.0E-07	1.6E-06

Table E.17 Results of adult nonconservative screening for carcinogens in dredged sediment where no measurements in a reach were above detection limits.

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
10	Sed. Ingestion	Organics	Toxaphene	11	1.55E+01 (Min)	1.1E+00	9.5E-07	1.0E-06
			Aldrin	11	7.72E-01 (Min)	1.7E+01	4.7E-08	8.0E-07
			Chlordane	11	7.72E+00 (Min)	1.3E+00	4.7E-07	6.1E-07
			Hexachlorobenzene	22	4.43E+00 (Min)	1.6E+00	2.7E-07	4.3E-07
			Heptachlor epoxide	11	7.72E-01 (Min)	9.1E+00	4.7E-08	4.3E-07
			Bis(2-chloroethyl)ether	22	4.43E+00 (Min)	1.1E+00	2.7E-07	3.0E-07
			Alpha-BHC	11	7.72E-01 (Min)	6.3E+00	4.7E-08	3.0E-07
			3,3'-Dichlorobenzidine	22	8.91E+00 (Min)	4.5E-01	5.5E-07	2.5E-07
			Heptachlor	11	7.72E-01 (Min)	4.5E+00	4.7E-08	2.1E-07
			Benzo(a)anthracene	22	4.43E+00 (Min)	7.3E-01	2.7E-07	2.0E-07
			Indeno(1,2,3-cd)pyrene	22	4.43E+00 (Min)	7.3E-01	2.7E-07	2.0E-07
			Benzo(b)fluoranthene	22	4.43E+00 (Min)	7.3E-01	2.7E-07	2.0E-07
			Benzo(k)fluoranthene	22	4.43E+00 (Min)	6.8E-01	2.7E-07	1.8E-07
			2,4-Dinitrotoluene	22	4.43E+00 (Min)	6.8E-01	2.7E-07	1.8E-07
			2,6-Dinitrotoluene	22	2.16E+01 (Min)	1.2E-01	1.3E-06	1.6E-07
			Pentachlorophenol	22	4.43E+00 (Min)	1.8E+00	4.7E-08	8.5E-08
			Beta-BHC	11	7.72E-01 (Min)	1.3E+00	4.7E-08	6.1E-08
			Gamma-BHC (Lindane)	11	7.72E-01 (Min)	1.3E+00	4.7E-08	6.1E-08
			4,4'-DDT	22	1.68E+00 (Min)	3.4E-01	1.0E-07	3.5E-08
			4,4'-DDE	22	1.68E+00 (Min)	3.4E-01	1.0E-07	3.5E-08
			4,4'-DDD	22	1.68E+00 (Min)	2.4E-01	1.0E-07	2.5E-08
			Hexachlorobutadiene	22	4.43E+00 (Min)	7.8E-02	2.7E-07	2.1E-08
			Bis(2-chloroisopropyl)ether	22	4.43E+00 (Min)	7.0E-02	2.7E-07	1.9E-08
			4-Chloroaniline	22	4.43E+00 (Min)	3.5E-02	2.7E-07	9.5E-09
			1,4-Dichlorobenzene	22	4.43E+00 (Min)	2.4E-02	2.7E-07	6.5E-09
			Hexachloroethane	22	4.43E+00 (Min)	1.4E-02	2.7E-07	3.8E-09
			2,4,6-Trichlorophenol	22	4.43E+00 (Min)	1.1E-02	2.7E-07	3.0E-09
			N-Nitrosodiphenylamine	22	4.43E+00 (Min)	4.9E-03	2.7E-07	1.3E-09
			Isophorone	22	4.43E+00 (Min)	4.1E-03	2.7E-07	1.1E-09

Table E.18. Results of adult nonconservative screening for noncarcinogens in dredged sediment where no measurements in a reach were above detection limits

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
1	Sed. Ingestion	Inorganics	Thallium Antimony	21	mg/kg	1/(mg/kg/d)	mg/kg/d	
				14				
				21	3.17E-01 (Min)	7.0E-05	4.5E-08	6.5E-04
				14	4.50E-01 (Min)	4.0E-04	6.4E-08	1.6E-04
				25	mg/kg	1/(mg/kg/d)	mg/kg/d	
				23				
		Organics	4,6-Dinitro-2-methylphenol Chlordane Heptachlor epoxide Dieldrin Endosulfan II Aldrin 2,4-Dinitrophenol Endosulfan I Nitrobenzene Hexachlorobenzene Hexachloroethane Endrin Hexachlorobutadiene Gamma-BHC (Lindane) 2,4-Dichlorophenol 4-Chloroaniline Heptachlor Methoxychlor 2-Chlorophenol Pentachlorophenol Hexachlorocyclopentadiene 1,2,4-Trichlorobenzene Bis(2-ethylhexyl)phthalate 2,4-Dimethylphenol Di-n-octylphthalate 2,4,5-Trichlorophenol Fluorene Naphthalene Bis(2-chloroisopropyl) ether 2-Methylphenol 4-Methylphenol Acenaphthene	25	2.49E+00 (Min)	1.0E-04	3.6E-07	3.6E-03
				23	5.98E-01 (Min)	6.0E-05	8.5E-08	1.4E-03
				23	5.98E-02 (Min)	1.3E-05	8.5E-09	6.6E-04
				23	1.20E-01 (Min)	5.0E-05	1.7E-08	3.4E-04
				23	1.20E-01 (Min)	5.0E-05	1.7E-08	3.4E-04
				23	5.98E-02 (Min)	3.0E-05	8.5E-09	2.8E-04
				25	2.49E+00 (Min)	2.0E-03	3.6E-07	1.8E-04
				23	5.98E-02 (Min)	5.0E-05	8.5E-09	1.7E-04
				25	5.16E-01 (Min)	5.0E-04	7.4E-08	1.5E-04
				25	5.16E-01 (Min)	8.0E-04	7.4E-08	9.2E-05
				25	5.16E-01 (Min)	1.0E-03	7.4E-08	7.4E-05
				23	1.20E-01 (Min)	3.0E-04	1.7E-08	5.7E-05
				25	5.16E-01 (Min)	2.0E-03	7.4E-08	3.7E-05
				23	5.98E-02 (Min)	3.0E-04	8.5E-09	2.8E-05
				25	5.16E-01 (Min)	4.0E-03	7.4E-08	2.5E-05
				25	5.16E-01 (Min)	4.0E-03	7.4E-08	1.8E-05
				23	5.98E-02 (Min)	5.0E-04	8.5E-09	1.7E-05
				23	5.98E-01 (Min)	5.0E-03	8.5E-08	1.7E-05
				25	5.16E-01 (Min)	5.0E-03	7.4E-08	1.5E-05
				25	2.49E+00 (Min)	3.0E-02	3.6E-07	1.2E-05
				25	5.16E-01 (Min)	7.0E-03	7.4E-08	1.1E-05
				25	5.16E-01 (Min)	2.0E-02	7.4E-08	7.4E-06
				25	7.48E-01 (Min)	2.0E-02	7.4E-08	5.3E-06
				25	5.16E-01 (Min)	2.0E-02	7.4E-08	3.7E-06
				25	5.16E-01 (Min)	2.0E-02	7.4E-08	3.7E-06
				25	2.49E+00 (Min)	1.0E-01	3.6E-07	3.6E-06
				25	5.16E-01 (Min)	4.0E-02	7.4E-08	1.8E-06
				25	5.16E-01 (Min)	4.0E-02	7.4E-08	1.8E-06
				25	5.16E-01 (Min)	4.0E-02	7.4E-08	1.8E-06
				25	5.16E-01 (Min)	5.0E-02	7.4E-08	1.5E-06
				25	5.16E-01 (Min)	5.0E-02	7.4E-08	1.5E-06
				25	5.16E-01 (Min)	6.0E-02	7.4E-08	1.2E-06

Table E.18 (continued)

Resch	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
1	Sed. Ingestion	Organics	2-Chloronaphthalene 1,2-Dichlorobenzene Di-n-butylphthalate Isophorone Butylbenzylphthalate Benzyl alcohol Anthracene Phenol Diethylphthalate Benzoic acid Dimethylphthalate	25 25 25 25 25 25 25 25 25 25 25 25	5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 5.16E-01 (Min) 2.49E+00 (Min) 5.16E-01 (Min)	8.0E-02 9.0E-02 1.0E-01 2.0E-01 2.0E-01 3.0E-01 3.0E-01 6.0E-01 8.0E-01 4.0E+00 1.0E+00	7.4E-08 7.4E-08 7.4E-08 7.4E-08 7.4E-08 7.4E-08 7.4E-08 7.4E-08 7.4E-08 3.6E-07 7.4E-08	9.2E-07 8.2E-07 7.4E-07 3.7E-07 3.7E-07 2.5E-07 2.5E-07 1.2E-07 9.2E-08 8.9E-08 7.4E-08
2	Sed. Ingestion	Inorganics	Thallius	40	mg/kg 3.02E-01 (Min)	1/(mg/kg/d) 7.0E-05	mg/kg/d 4.3E-08	6.2E-04
		Organics	4,6-Dinitro-2-methylphenol Chlordane Heptachlor epoxide Dieldrin Endosulfan II Aldrin Endosulfan I 2,4-Dinitrophenol Nitrobenzene Endrin Hexachlorobenzene Hexachloroethane 4,4'-DDT Gamma-BHC (Lindane) Hexachlorobutadiene Heptachlor Methoxychlor 2,4-Dichlorophenol 4-Chloroaniline	40 41 41 41 41 41 41 40 40 41 40 41 41 41 40 41 41 40 40 40	2.43E+00 (Min) 9.39E-01 (Min) 9.42E-02 (Min) 1.90E-01 (Min) 1.90E-01 (Min) 9.42E-02 (Min) 9.42E-01 (Min) 2.43E+00 (Min) 5.00E-01 (Min) 1.90E-01 (Min) 5.00E-01 (Min) 1.90E-01 (Min) 9.42E-02 (Min) 5.00E-01 (Min) 9.42E-02 (Min) 9.39E-01 (Min) 5.00E-01 (Min) 5.00E-01 (Min)	1.0E-04 6.0E-05 1.3E-05 5.0E-05 5.0E-05 3.0E-05 5.0E-05 2.0E-03 3.0E-04 8.0E-04 1.0E-03 5.0E-04 3.0E-03 2.0E-03 5.0E-04 3.0E-03 3.0E-03 4.0E-03	3.5E-07 1.3E-07 1.3E-08 2.7E-08 2.7E-08 1.3E-08 1.3E-08 3.5E-07 7.1E-08 2.7E-08 7.1E-08 2.7E-08 1.3E-08 1.3E-08 7.1E-08 7.1E-08 7.1E-08 7.1E-08	3.5E-03 2.2E-03 1.0E-03 5.4E-04 5.4E-04 4.5E-04 2.7E-04 1.7E-04 1.4E-04 9.0E-05 8.9E-05 5.4E-05 4.5E-05 3.6E-05 2.7E-05 2.7E-05 2.4E-05 1.8E-05

Reech	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
2	Sed. Ingestion	Organics	2-Chlorophenol	40	5.00E-01 (Min)	5.0E-03	7.1E-08	1.4E-05
			Pentachlorophenol	40	2.43E+00 (Min)	3.0E-02	3.5E-07	1.2E-05
			Hexachlorocyclopentadiene	40	5.00E-01 (Min)	7.0E-03	7.1E-08	1.0E-05
			1,2,4-Trichlorobenzene	40	5.00E-01 (Min)	1.0E-02	7.1E-08	7.1E-06
			2,4-Dimethylphthalate	40	5.00E-01 (Min)	2.0E-02	7.1E-08	3.6E-06
			Di-n-octylphthalate	40	5.00E-01 (Min)	2.0E-02	7.1E-08	3.6E-06
			2,4,5-Trichlorophenol	40	2.43E+00 (Min)	1.0E-01	3.5E-07	3.5E-06
			Fluorene	40	5.00E-01 (Min)	4.0E-02	7.1E-08	1.8E-06
			Bis(2-chloroisopropyl) ether	40	5.00E-01 (Min)	4.0E-02	7.1E-08	1.8E-06
			2-Methylphenol	40	5.00E-01 (Min)	5.0E-02	7.1E-08	1.4E-06
			4-Methylphenol	40	5.00E-01 (Min)	5.0E-02	7.1E-08	1.4E-06
			Acenaphthene	40	5.00E-01 (Min)	6.0E-02	7.1E-08	1.2E-06
			2-Chloronaphthalene	40	5.00E-01 (Min)	8.0E-02	7.1E-08	8.9E-07
			1,2-Dichlorobenzene	40	5.00E-01 (Min)	9.0E-02	7.1E-08	7.9E-07
			Isophorone	40	5.00E-01 (Min)	2.0E-01	7.0E-08	3.5E-07
			Butylbenzylphthalate	40	4.89E-01 (Min)	2.0E-01	7.1E-08	2.4E-07
			Benzyl alcohol	40	5.00E-01 (Min)	3.0E-01	7.1E-08	2.4E-07
			Anthracene	40	5.00E-01 (Min)	6.0E-01	7.1E-08	1.2E-07
			Phenol	40	5.00E-01 (Min)	4.0E+00	3.5E-07	8.7E-08
			Benzoic acid	40	2.43E+00 (Min)	1.0E+00	7.1E-08	7.1E-08
			Dimethylphthalate	40	5.00E-01 (Min)			
					mg/kg	1/(mg/kg/d)	mg/kg/d	
3	Sed. Ingestion	Organics	4,6-Dinitro-2-methylphenol	50	2.02E+00 (Min)	1.0E-04	2.9E-07	2.9E-03
			Chlordane	49	4.62E-01 (Min)	6.0E-05	6.6E-08	1.1E-03
			Heptachlor epoxide	49	4.62E-02 (Min)	1.3E-05	6.6E-09	5.1E-04
			Dieldrin	49	9.38E-02 (Min)	5.0E-05	1.3E-08	2.7E-04
			Endosulfan II	49	9.38E-02 (Min)	5.0E-05	1.3E-08	2.7E-04
			Aldrin	49	4.62E-02 (Min)	3.0E-05	6.6E-09	2.2E-04
			2,4-Dinitrophenol	50	2.02E+00 (Min)	2.0E-03	2.9E-07	1.4E-04
			Endosulfan I	49	4.62E-02 (Min)	5.0E-05	6.6E-09	1.3E-04
			Nitrobenzene	50	4.17E-01 (Min)	5.0E-04	6.0E-08	1.2E-04
			Hexachlorobenzene	50	4.17E-01 (Min)	8.0E-04	6.0E-08	7.4E-05
			Hexachloroethane	50	4.17E-01 (Min)	1.0E-03	6.0E-08	6.0E-05
			Endrin	49	9.38E-02 (Min)	3.0E-04	1.3E-08	4.5E-05
			Hexachlorobutadiene	50	4.17E-01 (Min)	2.0E-03	6.0E-08	3.0E-05

Table E.18 (continued)

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
3	Sed. Ingestion	Organics	4,4'-DDT	49	9.30E-02 (Min)	5.0E-04	1.3E-08	2.7E-05
			Gamma-BHC (Lindene)	49	4.62E-02 (Min)	3.0E-04	6.6E-09	2.2E-05
			2,4-Dichlorophenol	50	4.17E-01 (Min)	3.0E-03	6.0E-08	2.0E-05
			4-Chloroaniline	50	4.17E-01 (Min)	4.0E-03	6.0E-08	1.5E-05
			Heptachlor	49	4.62E-02 (Min)	5.0E-04	6.6E-09	1.3E-05
			Methoxychlor	49	4.62E-01 (Min)	5.0E-03	6.6E-08	1.3E-05
			2-Chlorophenol	50	4.17E-01 (Min)	5.0E-03	6.0E-08	1.2E-05
			Pentachlorophenol	50	2.02E+00 (Min)	3.0E-02	2.9E-07	9.6E-06
			Hexachlorocyclopentadiene	50	4.17E-01 (Min)	7.0E-03	6.0E-08	8.5E-06
			1,2,4-Trichlorobenzene	50	4.17E-01 (Min)	1.0E-02	6.0E-08	6.0E-06
			2,4-Dimethylphenol	50	4.17E-01 (Min)	2.0E-02	6.0E-08	3.0E-06
			Di-n-octylphthalate	50	4.17E-01 (Min)	2.0E-02	6.0E-08	3.0E-06
			2,4,5-Trichlorophenol	50	2.02E+00 (Min)	1.0E-01	2.9E-07	2.9E-06
			Fluorene	50	4.17E-01 (Min)	4.0E-02	6.0E-08	1.5E-06
			Bis(2-chloroisopropyl) ether	50	4.17E-01 (Min)	4.0E-02	6.0E-08	1.5E-06
			2-Methylphenol	50	4.17E-01 (Min)	5.0E-02	6.0E-08	1.2E-06
			4-Methylphenol	50	4.17E-01 (Min)	5.0E-02	6.0E-08	1.2E-06
			Acenaphthene	50	4.17E-01 (Min)	6.0E-02	6.0E-08	9.9E-07
			2-Chloronaphthalene	50	4.17E-01 (Min)	8.0E-02	6.0E-08	7.4E-07
			1,2-Dichlorobenzene	50	4.17E-01 (Min)	9.0E-02	6.0E-08	6.6E-07
			Isophorone	50	4.17E-01 (Min)	2.0E-01	6.0E-08	3.0E-07
			Butylbenzylphthalate	50	4.17E-01 (Min)	2.0E-01	6.0E-08	3.0E-07
			Benzyl alcohol	50	4.17E-01 (Min)	3.0E-01	6.0E-08	2.0E-07
			Phenol	50	4.17E-01 (Min)	6.0E-01	6.0E-08	9.9E-08
			Diethylphthalate	50	4.28E-01 (Min)	8.0E-01	6.1E-08	7.6E-08
			Benzoic acid	50	2.02E+00 (Min)	4.0E+00	2.9E-07	7.2E-08
4	Sed. Ingestion	Inorganics	Thallium	40	3.12E-01 (Min)	1/(mg/kg/d)	mg/kg/d	6.4E-04
					mg/kg	1/(mg/kg/d)	mg/kg/d	
			Organics	44	47	2.40E+00 (Min)	1.0E-04	3.4E-07
			4,6-Dinitro-2-methylphenol	44	9.54E-01 (Min)	6.0E-05	1.4E-07	2.3E-03
			Chlordane	44	9.54E-02 (Min)	1.3E-05	1.4E-08	1.0E-03
			Heptachlor epoxide	45	1.91E-01 (Min)	5.0E-05	2.7E-08	5.4E-04
			Dieldrin					

Table E.18 (continued)

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
4	Sed. Ingestion	Organics	Endosulfan II	45	1.91E-01 (Min)	5.0E-05	2.7E-08	5.4E-04
			Aldrin	44	9.54E-02 (Min)	3.0E-05	1.4E-08	4.5E-04
			Endosulfan I	44	9.54E-02 (Min)	5.0E-05	1.4E-08	2.7E-04
			2,4-Dinitrophenol	44	2.41E+00 (Min)	2.0E-03	3.4E-07	1.7E-04
			Nitrobenzene	47	4.97E-01 (Min)	5.0E-04	7.1E-08	1.4E-04
			Endrin	45	1.91E-01 (Min)	3.0E-04	2.7E-08	9.1E-05
			Hexachlorobenzene	47	4.97E-01 (Min)	8.0E-04	7.1E-08	8.9E-05
			Hexachloroethane	47	4.97E-01 (Min)	1.0E-03	7.1E-08	7.1E-05
			4,4'-DDT	45	1.91E-01 (Min)	5.0E-04	2.7E-08	5.4E-05
			Gamma-BHC (Lindane)	44	9.54E-02 (Min)	3.0E-04	1.4E-08	4.5E-05
			Hexachlorobutadiene	47	4.97E-01 (Min)	2.0E-03	7.1E-08	3.5E-05
			Methoxychlor	45	9.54E-01 (Min)	5.0E-03	1.4E-07	2.7E-05
			Heptachlor	44	9.54E-02 (Min)	5.0E-04	1.4E-08	2.7E-05
			2,4-Dichlorophenol	47	4.97E-01 (Min)	3.0E-03	7.1E-08	2.4E-05
			4-Chloroaniline	47	4.97E-01 (Min)	4.0E-03	7.1E-08	1.8E-05
			2-Chlorophenol	47	4.97E-01 (Min)	5.0E-03	7.1E-08	1.4E-05
			Pentachlorophenol	47	2.40E+00 (Min)	3.0E-02	3.4E-07	1.1E-05
			Hexachlorocyclopentadiene	47	4.97E-01 (Min)	7.0E-03	7.1E-08	1.0E-05
			1,2,4-Trichlorobenzene	47	4.97E-01 (Min)	1.0E-02	7.1E-08	7.1E-06
			2,4-Dimethylphenol	47	4.97E-01 (Min)	2.0E-02	7.1E-08	3.5E-06
			Di-n-octylphthalate	47	4.97E-01 (Min)	2.0E-02	7.1E-08	3.5E-06
			2,4,5-Trichlorophenol	47	2.40E+00 (Min)	1.0E-01	3.4E-07	3.4E-06
			Fluorene	47	4.97E-01 (Min)	4.0E-02	7.1E-08	1.8E-06
			Naphthalene	47	4.97E-01 (Min)	4.0E-02	7.1E-08	1.8E-06
			Bis(2-chloroisopropyl)ether	47	4.97E-01 (Min)	4.0E-02	7.1E-08	1.8E-06
			2-Methylphenol	47	4.97E-01 (Min)	5.0E-02	7.1E-08	1.4E-06
			4-Methylphenol	47	4.97E-01 (Min)	5.0E-02	7.1E-08	1.4E-06
			Acenaphthene	47	4.97E-01 (Min)	6.0E-02	7.1E-08	1.2E-06
			2-Chloronaphthalene	47	4.97E-01 (Min)	8.0E-02	7.1E-08	8.9E-07
			1,2-Dichlorobenzene	47	4.97E-01 (Min)	9.0E-02	7.1E-08	7.9E-07
			Di-n-butylphthalate	47	4.97E-01 (Min)	1.0E-01	7.1E-08	7.1E-07
			Isophorone	47	4.97E-01 (Min)	2.0E-01	7.1E-08	3.5E-07
			Butylbenzylphthalate	47	4.97E-01 (Min)	2.0E-01	7.1E-08	3.5E-07
			Benzyl alcohol	47	4.97E-01 (Min)	3.0E-01	7.1E-08	2.4E-07
			Anthracene	47	4.97E-01 (Min)	3.0E-01	7.1E-08	2.4E-07
			Phenol	47	4.97E-01 (Min)	6.0E-01	7.1E-08	1.2E-07
			Diethylphthalate	47	4.97E-01 (Min)	8.0E-01	7.1E-08	8.9E-08
			Benzoic acid	47	2.40E+00 (Min)	4.0E+00	3.4E-07	8.6E-08
			Dimethylphthalate	47	4.97E-01 (Min)	1.0E+00	7.1E-08	7.1E-08

Table E.18 (continued)

Resch	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
5	Sed. Ingestion	Inorganics	Thallium Selenium	68 40	3.99E-01 (Min) 4.01E-01 (Min)	7.0E-05 5.0E-03	5.7E-08 5.7E-08	8.1E-04 1.1E-05
					mg/kg	1/(mg/kg/d)	mg/kg/d	
		Organics	Chlordane	42	3.55E+00 (Min)	6.0E-05	5.1E-07	8.4E-03
			4,6-Dinitro-2-methylphenol	62	3.54E+00 (Min)	1.0E-04	5.1E-07	5.1E-03
			Heptachlor epoxide	42	3.55E-01 (Min)	1.3E-05	5.1E-08	3.9E-03
			Dieldrin	62	7.12E-01 (Min)	5.0E-05	1.0E-07	2.0E-03
			Endosulfan II	62	7.12E-01 (Min)	5.0E-05	1.0E-07	2.0E-03
			Aldrin	42	3.55E-01 (Min)	3.0E-05	5.1E-08	1.7E-03
			Endosulfan I	42	3.55E-01 (Min)	5.0E-05	5.1E-08	1.0E-03
			Endrin	62	7.12E-01 (Min)	3.0E-04	1.0E-07	3.4E-04
			2,4-Dinitrophenol	51	3.54E+00 (Min)	2.0E-03	5.1E-07	2.5E-04
			Nitrobenzene	62	7.25E-01 (Min)	5.0E-04	1.0E-07	2.1E-04
			4,4'-DDT	62	7.12E-01 (Min)	5.0E-04	1.0E-07	2.0E-04
			Gamma-BHC (Lindane)	42	3.55E-01 (Min)	3.0E-04	5.1E-08	1.7E-04
			Hexachlorobenzene	62	7.25E-01 (Min)	8.0E-04	1.0E-07	1.3E-04
			Hexachloroethane	62	7.25E-01 (Min)	1.0E-03	1.0E-07	1.0E-04
			Methoxychlor	62	7.25E-01 (Min)	5.0E-04	5.1E-07	1.0E-04
			Heptachlor	42	3.55E-01 (Min)	5.0E-04	5.1E-08	1.0E-04
			Hexachlorobutadiene	62	7.25E-01 (Min)	2.0E-03	1.0E-07	5.2E-05
			2,4-Dichlorophenol	62	7.25E-01 (Min)	3.0E-03	1.0E-07	3.5E-05
			4-Chloroaniline	62	7.25E-01 (Min)	4.0E-03	1.0E-07	2.6E-05
			2-Chlorophenol	62	7.25E-01 (Min)	5.0E-03	1.0E-07	2.1E-05
			Pentachlorophenol	62	3.54E+00 (Min)	3.0E-02	5.1E-07	1.7E-05
			Hexachlorocyclopentadiene	62	7.25E-01 (Min)	7.0E-03	1.0E-07	1.5E-05
			1,2,4-Trichlorobenzene	62	7.25E-01 (Min)	1.0E-02	1.0E-07	1.0E-05
			2,4-Dimethylphenol	62	7.25E-01 (Min)	2.0E-02	1.0E-07	5.2E-06
			Di-n-octylphthalate	62	3.54E+00 (Min)	2.0E-02	1.0E-07	5.2E-06
			2,4,5-Trichlorophenol	62	7.25E-01 (Min)	1.0E-01	5.1E-07	5.1E-06
			Pyrene	62	7.25E-01 (Min)	3.0E-02	1.0E-07	3.5E-06
			Fluorene	62	7.25E-01 (Min)	4.0E-02	1.0E-07	2.6E-06
			Naphthalene	62	7.25E-01 (Min)	4.0E-02	1.0E-07	2.6E-06

Table E.18 (continued)

Reach	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
5	Sed. Ingestion	Organics	Bis(2-chloroisopropyl)ether	62	7.25E-01 (Min)	4.0E-02	1.0E-07	2.6E-06
			Fluoranthene	62	7.25E-01 (Min)	4.0E-02	1.0E-07	2.6E-06
			2-Methylphenol	54	7.25E-01 (Min)	5.0E-02	1.0E-07	2.1E-06
			4-Methylphenol	62	7.25E-01 (Min)	5.0E-02	1.0E-07	2.1E-06
			Acenaphthene	62	7.25E-01 (Min)	6.0E-02	1.0E-07	1.7E-06
			2-Chloronaphthalene	62	7.25E-01 (Min)	8.0E-02	1.0E-07	1.3E-06
			1,2-Dichlorobenzene	62	7.25E-01 (Min)	9.0E-02	1.0E-07	1.2E-06
			Di-n-butylphthalate	62	7.25E-01 (Min)	1.0E-01	1.0E-07	1.0E-06
			Isophorone	62	7.25E-01 (Min)	2.0E-01	1.0E-07	5.2E-07
			Butylbenzylphthalate	62	7.25E-01 (Min)	2.0E-01	1.0E-07	5.2E-07
			Benzyl alcohol	54	7.25E-01 (Min)	3.0E-01	1.0E-07	3.5E-07
			Anthracene	62	7.25E-01 (Min)	3.0E-01	1.0E-07	3.5E-07
			Phenol	62	7.25E-01 (Min)	6.0E-01	1.0E-07	1.7E-07
			Diethylphthalate	62	7.25E-01 (Min)	8.0E-01	1.0E-07	1.3E-07
			Benzoic acid	62	3.54E+00 (Min)	4.0E+00	5.1E-07	1.3E-07
			Dimethylphthalate	62	7.25E-01 (Min)	1.0E+00	1.0E-07	1.0E-07
10	Sed. Ingestion	Inorganics	Thallium	23	4.37E-01 (Min)	7.0E-05	6.2E-08	8.9E-04
			Selenium	17	4.51E-01 (Min)	5.0E-03	6.4E-08	1.3E-05
					mg/kg	1/(mg/kg/d)	mg/kg/d	
					mg/kg	1/(mg/kg/d)	mg/kg/d	
		Organics	4,6-Dinitro-2-methylphenol	22	2.16E+01 (Min)	1.0E-04	3.1E-06	3.1E-02
			Chlordane	11	7.72E+00 (Min)	6.0E-05	1.1E-06	1.8E-02
			Heptachlor epoxide	11	7.72E-01 (Min)	1.5E-05	1.1E-07	8.5E-03
			Dieldrin	22	1.68E+00 (Min)	5.0E-05	2.4E-07	4.8E-03
			Endosulfan II	22	1.68E+00 (Min)	5.0E-05	2.4E-07	4.8E-03
			Aldrin	11	7.72E-01 (Min)	3.0E-05	1.1E-07	3.7E-03
			Endosulfan I	11	7.72E-01 (Min)	5.0E-05	1.1E-07	2.2E-03
			2,4-Dinitrophenol	22	2.16E+01 (Min)	2.0E-03	3.1E-06	1.5E-03
			Nitrobenzene	22	4.43E+00 (Min)	5.0E-04	6.3E-07	1.3E-03
			Endrin	22	4.43E+00 (Min)	3.0E-04	2.4E-07	8.0E-04
			Hexachlorobenzene	22	4.43E+00 (Min)	8.0E-04	6.3E-07	7.9E-04
			Hexachloroethane	22	4.43E+00 (Min)	1.0E-03	6.3E-07	6.3E-04
			4,4'-DDT	22	1.68E+00 (Min)	5.0E-04	2.4E-07	4.8E-04

Table E.18 (continued)

Resch	Media	Contaminant type	Contaminant	Total samples	Concentration (type)	Reference dose	Intake rate	Noncarcinogen screening index
10	Sed. Ingestion	Organics	Gamma-BHC (Lindane)	11	7.72E-01 (Min)	3.0E-04	1.1E-07	3.7E-04
			Hexachlorobutadiene	22	4.43E+00 (Min)	2.0E-03	6.3E-07	3.2E-04
			Methoxychlor	22	8.32E+00 (Min)	5.0E-03	1.2E-06	2.4E-04
			Heptachlor	11	7.72E-01 (Min)	5.0E-04	1.1E-07	2.2E-04
			2,4-Dichlorophenol	22	4.43E+00 (Min)	3.0E-03	6.3E-07	2.1E-04
			4-Chloroaniline	22	4.43E+00 (Min)	4.0E-03	6.3E-07	1.6E-04
			2-Chlorophenol	22	4.43E+00 (Min)	5.0E-03	6.3E-07	1.3E-04
			Pentachlorophenol	22	2.16E+01 (Min)	3.0E-02	3.1E-06	1.0E-04
			Hexachlorocyclopentadiene	22	4.43E+00 (Min)	7.0E-03	6.3E-07	9.0E-05
			1,2,4-Trichlorobenzene	22	4.43E+00 (Min)	1.0E-02	6.3E-07	6.3E-05
			2,4-Dimethylphenol	22	4.43E+00 (Min)	2.0E-02	6.3E-07	3.2E-05
			Di-n-octylphthalate	22	4.43E+00 (Min)	2.0E-02	6.3E-07	3.1E-05
			2,4,5-Trichlorophenol	22	2.16E+01 (Min)	1.0E-01	3.1E-06	1.6E-05
			Fluorene	22	4.43E+00 (Min)	4.0E-02	6.3E-07	1.6E-05
			Naphthalene	22	4.43E+00 (Min)	4.0E-02	6.3E-07	1.6E-05
			Bis(2-chloroisopropyl)ether	22	4.43E+00 (Min)	4.0E-02	6.3E-07	1.6E-05
			2-Methylphenol	22	4.43E+00 (Min)	5.0E-02	6.3E-07	1.3E-05
			4-Methylphenol	22	4.43E+00 (Min)	5.0E-02	6.3E-07	1.3E-05
			Acenaphthene	22	4.43E+00 (Min)	6.0E-02	6.3E-07	1.1E-05
			2-Chloronaphthalene	22	4.43E+00 (Min)	8.0E-02	6.3E-07	7.9E-06
			1,2-Dichlorobenzene	22	4.43E+00 (Min)	9.0E-02	6.3E-07	7.0E-06
			Di-n-butylphthalate	22	4.43E+00 (Min)	1.0E-01	6.3E-07	6.3E-06
			Isophorone	22	4.43E+00 (Min)	2.0E-01	6.3E-07	3.2E-06
			Butylbenzylphthalate	22	4.43E+00 (Min)	2.0E-01	6.3E-07	3.2E-06
			Benzyl alcohol	22	4.43E+00 (Min)	3.0E-01	6.3E-07	2.1E-06
			Phenol	22	4.43E+00 (Min)	6.0E-01	6.3E-07	1.1E-06
			Diethylphthalate	22	4.43E+00 (Min)	8.0E-01	6.3E-07	7.9E-07
			Benzoic acid	22	2.16E+01 (Min)	4.0E+00	3.1E-06	7.7E-07
			Dimethylphthalate	22	4.43E+00 (Min)	1.0E+00	6.3E-07	6.3E-07

Table E.19. Analytes without reference doses and
slope (cancer potency) factors

Type	Analyte	CAS No.
Inorganics	Aluminum	7429-90-5
	Iron	7439-89-6
	Lead	7439-92-1
	Magnesium	7439-95-4
Organics	1,3-Dichlorobenzene	541-73-1
	2-Hexanone	591-78-6
	2-Methylnaphthalene	91-57-6
	2-Nitroaniline	88-74-4
	2-Nitrophenol	88-75-5
	3-Nitroaniline	99-09-2
	4-Bromophenyl-phenylether	101-55-3
	4-Chloro-3-methylphenol	59-50-7
	4-Chlorophenyl-phenylether	7005-72-3
	4-Nitroaniline	100-01-6
	4-Nitrophenol	100-02-7
	Bis(2-chloroethoxy)methane	111-91-1
	Chloroethane	75-00-3
	Delta-BHC	319-86-8
	Dibenzofuran	132-64-9
	Endosulfan sulfate	1031-07-8
	Endrin ketone	53494-70-5
Radionuclides	Cm-245	15621-76-8
	Cm-248	15758-33-5

Appendix F
ECOLOGICAL RISK METHODS

CONTENTS OF APPENDIX F

F.1	DATA SUMMARIZATION FOR ECOLOGICAL RISK ASSESSMENT	F-3
F.2	FISH	F-3
F.3	SEDIMENT CORES	F-4
F.4	WATER	F-4

TABLES

F.1	Mean and upper confidence limit (UCL) concentrations of chemicals detected in the dissolved phase of CRRI water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-6
F.2	Limits of detection of chemicals not detected in the dissolved phase of the CRRI water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-8
F.3	Mean and upper confidence limit (UCL) concentrations of chemicals detected in NPDES water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-22
F.4	Limits of detection of chemicals not detected in NPDES water samples ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for aquatic life	F-35
F.5	Mean and upper confidence limit (UCL) concentrations of chemicals detected in fish ($\mu\text{g/L}$), bioconcentration factors, estimated exposure in water ($\mu\text{g/L}$), and quotients for exposure to that water concentration	F-54
F.6	Limits of detection of chemicals not detected in fish ($\mu\text{g/L}$), bioconcentration factors, estimated exposure in water ($\mu\text{g/L}$), and quotients for exposure to that water concentration	F-66
F.7	Concentrations of inorganic chemicals detected in sediment ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-86
F.8	Detection limits for inorganic chemicals not detected in sediment ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-91
F.9	Mean and upper confidence limit (UCL) concentrations of organic chemicals detected in sediment ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-92
F.10	Detection limits for organic chemicals not detected in sediment ($\mu\text{g/L}$), estimated pore water concentrations ($\mu\text{g/L}$), toxicological benchmarks ($\mu\text{g/L}$), and quotients for benthic aquatic life	F-95
F.11	Mean and upper confidence limit (UCL) concentrations of chemicals in fish filets (mg/kg), toxicological benchmarks for wildlife (mg/kg), and quotients for hazards to piscivorous wildlife	F-140
F.12	Minimum and upper confidence limit (UCL) detection limits for chemicals not detected in fish filets (mg/kg), toxicological benchmarks for wildlife (mg/kg), and quotients for hazards to piscivorous wildlife	F-143

F.1 DATA SUMMARIZATION FOR ECOLOGICAL RISK ASSESSMENT

The presentation of results of the analyses for the selected analytes (Table A.1) provides a comprehensive view of the extent of contamination in the three media. However, for the risk assessment, these data must be summarized as typical and extreme exposure concentrations. For detected chemicals, mean concentrations and the upper 95% confidence limits of the mean concentrations were considered. The mean concentrations are assumed to represent typical chronic exposures. Upper 95% confidence limits may represent local extreme chronic concentrations or acute exposures. For undetected chemicals, minimum detection limits and the upper 95% confidence limits of the mean of the detection limits were the concentrations considered.

Only 10% of the determinations were above their respective method detection limit. This is a common observation for regulatory driven environmental sampling and analysis activities. It follows that 90% of the analyte determinations were not detected above their respective analytical method detection limits in CRRI samples. For the screening risk analyses, a distinction was made between *detected contaminants*, contaminants with at least one detected value in a given reach, and *non-detected contaminants*, contaminants with no detected values in a reach. Separate analyses and interpretations of these results were required.

The following sections describe the specific data manipulations performed for each of the selected media results.

F.2 FISH

F.2.1 Subsetting

Data were combined for all fish species within a reach for all chemicals except for PCBs. Only catfish results were used for PCBs. Only detected chlordane isomer values were summed. Only regular samples—no field duplicates—were used.

F.2.2 Summary Statistics

F.2.2.1 Detected contaminants

Given the potential for means of distributions with numerous non-detect values (censored data) to be positively biased, the upper 95% confidence limit of the arithmetic mean and the geometric mean were estimated using maximum likelihood estimator calculations (SAS PROC LIFEREG). The upper 95% confidence limit was compared with the actual maximum value and the lower value was used in the risk characterization. Detection limits were used as is.

F.2.2.2 Non-detected contaminants

The upper 95% confidence limit of the arithmetic mean was calculated and compared to the maximum analyte detection limit. The smaller of the two values, along with the minimum detection limit were used for the exposure assessment.

F.3 SEDIMENT CORES

F.3.1 Subsetting

Only analytical data for core segments from depths ≤ 6 cm were used.

F.3.2 Summary Statistics

F.3.2.1 Detected contaminants

The arithmetic mean concentration was calculated to a depth of 6 cm. When there was more than one core per reach, the average for each core was calculated and then the upper 95% confidence limit of the mean of the core averages, or the maximum core average, (whichever was smaller) was used as the value for the reach. The arithmetic mean of the core averages was used as the estimate of chronic exposure.

F.3.2.2 Non-detected contaminants

The upper 95% confidence limit of the arithmetic mean was calculated and compared to the maximum analyte detection limit. The smaller of the two values, along with the minimum detection limit was used for the exposure assessment.

F.4 WATER

F.4.1 Subsetting

F.4.1.1 Dissolved constituents

Dissolved metal concentrations and total organic concentrations from CRRRI samples were used for one set of exposure assessments. Organic analyses were performed on unfiltered CRRRI samples.

F.4.1.2 Total concentrations

Because dissolved concentrations were available for only the limited CRRRI samples, total metal concentrations from unfiltered CRRRI samples and all contaminant results from unfiltered ORR NPDES and environmental monitoring samples (Table E.1 and Fig. E.1) were included, but exposures were analyzed and discussed separately.

F.4.2 Summary Statistics

F.4.2.1 Detected contaminants

The values for the screening assessment for a given contaminant in a reach were calculated by first averaging all results at a location and then determining the upper 95% confidence limit of the mean of the location averages. Either the upper 95% confidence limit or the maximum location average, whichever was smaller, was used as the value for the reach.

F.4.2.2 Nondetected contaminants

The upper 95% confidence limit of the arithmetic mean was calculated and compared to the maximum analyte detection limit. The smaller of the two values, along with the minimum detection limit was used for the exposure assessment.

Table F.1. Mean and upper confidence limit (UCL) concentration of chemicals detected in the dissolved phase of CRR1 water samples (ug/L), toxicological benchmarks (ug/L), and quotients for aquatic life

Analyte	Reach	Number		Geo. mean ^a water conc.	UCL ^b water conc.	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
		analyzed in reach (water)	Number detects in reach (water)							
CADMIUM	1	3	1	0.27	0.39	20.769	30.00	0.01		SS EC20
	1	3	1	0.27	0.39	1.800	2.60	0.15		LCV DAPHNIDS
	1	3	1	0.27	0.39	0.360	0.52	0.75		LTV EC20 DAPHNID
	1	3	1	0.27	0.39	0.245	0.35	1.10		NAWQ ACTUAL
	1	3	1	0.27	0.39	0.159	0.23	1.70		LCV FISH
	1	3	1	0.27	0.39	0.150	0.22	1.80		LTV EC20 FISH
	1	3	1	0.27	0.39	0.063	0.09	4.30		EC20 POPULATION
	1	3	1	0.27	0.31	18.718	23.85	0.01		SS EC20
	4	3	1	0.24	0.31	1.622	2.07	0.15		LCV DAPHNIDS
	4	3	1	0.24	0.31	0.324	0.41	0.75		LTV EC20 DAPHNID
	4	3	1	0.24	0.31	0.221	0.28	1.10		NAWQ ACTUAL
	4	3	1	0.24	0.31	0.143	0.18	1.70		LCV FISH
	4	3	1	0.24	0.31	0.135	0.17	1.80		LTV EC20 FISH
	4	3	1	0.24	0.31	0.057	0.07	4.30		EC20 POPULATION
	2	2	1	29.80	54.03	1.862	3.37	16.00		LTV EC20 DAPHNID
	2	2	1	29.80	54.03	0.189	0.34	158.00		LCV DAPHNIDS
IRON	2	2	1	29.80	54.03	0.030	0.05	1000.00		NAWQ ACTUAL
	5	3	1	18.33	29.00	1.146	1.81	16.00		LTV EC20 DAPHNID
	5	3	1	18.33	29.00	0.116	0.18	158.00		LCV DAPHNIDS
	5	3	1	18.33	29.00	0.018	0.03	1000.00		NAWQ ACTUAL
	1	3	1	1.80	2.40	5.143	6.86	0.35		SS EC20
	1	3	1	1.80	2.40	0.562	0.75	3.20		NAWQ ACTUAL
LEAD	1	3	1	1.80	2.40	0.147	0.20	12.26		LCV DAPHNIDS
	1	3	1	1.80	2.40	0.095	0.13	18.88		LCV FISH
	1	3	1	1.80	2.40	0.082	0.11	22.00		LTV EC20 FISH
	1	3	1	1.80	2.40	0.025	0.03	71.00		EC20 POPULATION
	4	3	1	0.97	1.50	2.762	4.29	0.35		SS EC20
	4	3	1	0.97	1.50	0.302	0.47	3.20		NAWQ ACTUAL
	4	3	1	0.97	1.50	0.079	0.12	12.26		LCV DAPHNIDS
	4	3	1	0.97	1.50	0.051	0.08	18.88		LCV FISH
	4	3	1	0.97	1.50	0.044	0.07	22.00		LTV EC20 FISH
	4	3	1	0.97	1.50	0.014	0.02	71.00		EC20 POPULATION
	13	1	1	0.60	0.60	1.714	1.71	0.35		SS EC20
	13	1	1	0.60	0.60	0.187	0.19	3.20		NAWQ ACTUAL
	13	1	1	0.60	0.60	0.049	0.05	12.26		LCV DAPHNIDS
	13	1	1	0.60	0.60	0.032	0.03	18.88		LCV FISH
	13	1	1	0.60	0.60	0.027	0.03	22.00		LTV EC20 FISH
	13	1	1	0.60	0.60	0.008	0.01	71.00		EC20 POPULATION

Table F.1 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Geo. mean ^a water conc.	UCL ^b water conc.	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
MERCURY	3	4	1	0.28	0.52	936.667	1746.67	0.00	<	NAHQ ADVISORY
	3	4	1	0.28	0.52	9.367	17.47	0.03	<	LTV EC20 FISH
	3	4	1	0.28	0.52	7.025	13.10	0.04		LCV DAPHNIDS
	3	4	1	0.28	0.52	1.004	1.87	0.28		EC20 POPULATION
	3	4	1	0.28	0.52	0.540	1.01	0.52		LCV FISH
	3	4	1	0.28	0.52	0.323	0.60	0.87		LTV EC20 DAPHNID
SILVER	3	4	1	3.80	13.00	31.708	108.33	0.12		LCV FISH
	3	4	1	3.80	13.00	27.179	92.86	0.14		SS EC20
	3	4	1	3.80	13.00	19.025	65.00	0.20		LTV EC20 FISH
	3	4	1	3.80	13.00	15.220	52.00	0.25		NAHQ ADVISORY
	3	4	1	3.80	13.00	11.891	40.63	0.32		EC20 POPULATION
	3	4	1	3.80	13.00	6.795	23.21	0.56	<	LTV EC20 DAPHNID
SODIUM	3	4	1	3.80	13.00	1.463	5.00	2.60		LCV DAPHNIDS
	1	3	3	4910.00	5260.00	0.058	0.06	85049.00	*	LCV DAPHNIDS
	2	2	2	5000.00	5220.00	0.059	0.06	85049.00	*	LCV DAPHNIDS
	3	4	4	3440.00	4150.00	0.040	0.05	85049.00	*	LCV DAPHNIDS
	4	2	2	4335.00	4910.00	0.051	0.06	85049.00	*	LCV DAPHNIDS
	5	4	4	4715.00	5140.00	0.055	0.06	85049.00	*	LCV DAPHNIDS
	10	2	2	5000.00	5060.00	0.059	0.06	85049.00	*	LCV DAPHNIDS
	13	1	1	2640.00	2640.00	0.031	0.03	85049.00	*	LCV DAPHNIDS
	10	2	1	14.45	21.00	0.688	1.00	21.00		SS EC20
	10	2	1	14.45	21.00	0.397	0.58	36.41		LCV FISH
	10	2	1	14.45	21.00	0.309	0.45	46.73		LCV DAPHNIDS
	10	2	1	14.45	21.00	0.307	0.45	47.00		LTV EC20 FISH
ZINC	10	2	1	14.45	21.00	0.181	0.26	80.00		EC20 POPULATION
	10	2	1	14.45	21.00	0.131	0.19	110.00		NAHQ ACTUAL

^aIn cases of mixed detects and nondetects, statistics are calculated using all values, "as is." The mean concentration is the geometric mean.

^bUCL concentration was compared with the minimum value, and the smaller has been used as the "UCL Concentration."

Table F.2. Limits of detection of chemicals not detected in the dissolved phase of the CRRJ water samples (ug/L), toxicological benchmarks (ug/L), and quotients for aquatic life

analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
ANTIMONY	1	3	0	1.90	1.900	0.106	0.106	18.00	NAHQ ADVISORY
	1	3	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	1	3	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	1	3	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	1	3	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH
	1	3	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	1	3	0	1.90	1.900	0.106	0.106	18.00	NAHQ ADVISORY
	2	2	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	2	2	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	2	2	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	2	2	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH
	2	2	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	3	4	0	1.90	1.900	0.106	0.106	18.00	NAHQ ADVISORY
	3	4	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	3	4	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	3	4	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	3	4	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH
	3	4	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	4	3	0	1.90	1.900	0.106	0.106	18.00	NAHQ ADVISORY
	4	3	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	4	3	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	4	3	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	4	3	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH
	4	3	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	5	4	0	1.90	1.900	0.106	0.106	18.00	NAHQ ADVISORY
	5	4	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	5	4	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	5	4	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	5	4	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH
	5	4	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	5	4	0	1.90	1.900	0.106	0.106	18.00	NAHQ ADVISORY
	10	2	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	10	2	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	10	2	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	10	2	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH
	10	2	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	10	2	0	1.90	1.900	0.000	0.000	18.00	NAHQ ADVISORY
	13	1	0	1.90	1.900	0.024	0.024	79.00	EC20 POPULATION
	13	1	0	1.90	1.900	0.001	0.001	1600.00	LCV FISH
	13	1	0	1.90	1.900	0.001	0.001	1900.00	LTV EC20 DAPHNID
	13	1	0	1.90	1.900	0.001	0.001	2310.00	LTV EC20 FISH

Table F.2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	IDL quotient	Benchmark	Benchmark type
ARSENIC	13	1	0	1.90	1.900	0.000	0.000	5400.00	LCV DAPHNIDS
	1	3	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	1	3	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	1	3	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	1	3	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	1	3	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	1	3	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	1	3	0	1.30	1.300	0.000	0.000	2962.00	LCV FISH
	1	3	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	2	2	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	2	2	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	2	2	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	2	2	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	2	2	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	2	2	0	1.30	1.300	0.000	0.000	2962.00	LCV FISH
	2	2	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	3	4	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	3	4	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	3	4	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	3	4	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	3	4	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	3	4	0	1.30	1.300	0.000	0.000	2962.00	LCV FISH
	4	3	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	4	3	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	4	3	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	4	3	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	4	3	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	4	3	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	4	3	0	1.30	1.300	0.000	0.000	2962.00	LCV FISH
	5	4	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	5	4	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	5	4	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	5	4	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	5	4	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	5	4	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	5	4	0	1.30	1.300	0.000	0.000	2962.00	LCV FISH
	5	4	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	10	2	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	10	2	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	10	2	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	10	2	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	10	2	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	10	2	0	1.30	1.300	0.000	0.000	2962.00	LCV FISH

Table F.2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UPL quotient	Benchmark	Benchmark type
ARSENIC (continued)	13	1	0	1.30	1.300	0.024	0.024	55.00	SS EC20
	13	1	0	1.30	1.300	0.007	0.007	190.00	NAHQ ACTUAL
	13	1	0	1.30	1.300	0.002	0.002	633.00	LTV EC20 DAPHNID
	13	1	0	1.30	1.300	0.001	0.001	914.10	LCV DAPHNIDS
	13	1	0	1.30	1.300	0.001	0.001	1995.00	EC20 POPULATION
	13	1	0	1.30	1.300	0.001	0.001	2130.00	LTV EC20 FISH
	13	1	0	1.30	1.300	0.001	0.001	2962.00	LCV FISH
	13	1	0	1.30	1.300	0.000	0.000		
	1	3	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	1	3	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
BERYLLIUM	1	3	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	1	3	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	1	3	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	1	3	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	1	3	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	2	2	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	2	2	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	2	2	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	2	2	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	2	2	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	2	2	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	3	4	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	3	4	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	3	4	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	3	4	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	3	4	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	3	4	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	4	2	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	4	2	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	4	2	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	4	2	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	4	2	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	4	2	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	5	4	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	5	4	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	5	4	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	5	4	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	5	4	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	5	4	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	5	4	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	5	4	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	5	4	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	5	4	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	5	4	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	5	4	0	3.90	3.900	6.393	6.393	0.61	NAHQ ADVISORY
	10	2	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	10	2	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	10	2	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	10	2	0	3.90	3.900	0.068	0.068	57.00	LCV FISH

Table F-2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UOL quotient	Benchmark	Benchmark type
BERYLLIUM (continued)	10	2	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	13	1	0	3.90	3.900	6.393	6.393	0.61	NAWQ ADVISORY
	13	1	0	3.90	3.900	1.026	1.026	3.80	LTV EC20 DAPHNID
	13	1	0	3.90	3.900	0.736	0.736	5.30	LCV DAPHNIDS
	13	1	0	3.90	3.900	0.186	0.186	21.00	EC20 POPULATION
	13	1	0	3.90	3.900	0.068	0.068	57.00	LCV FISH
	13	1	0	3.90	3.900	0.026	0.026	148.00	LTV EC20 FISH
	2	2	0	0.21	0.210	16.154	16.154	0.01	SS EC20
	2	2	0	0.21	0.210	1.400	1.400	0.15	LCV DAPHNIDS
	2	2	0	0.21	0.210	0.280	0.280	0.75	LTV EC20 DAPHNID
CADMIUM	2	2	0	0.21	0.210	0.191	0.191	1.10	NAWQ ACTUAL
	2	2	0	0.21	0.210	0.124	0.124	1.70	LCV FISH
	2	2	0	0.21	0.210	0.117	0.117	1.80	LTV EC20 FISH
	2	2	0	0.21	0.210	0.049	0.049	4.30	EC20 POPULATION
	3	3	0	0.21	0.210	16.154	16.154	0.01	SS EC20
	3	3	0	0.21	0.210	1.400	1.400	0.15	LCV DAPHNIDS
	3	3	0	0.21	0.210	0.280	0.280	0.75	LTV EC20 DAPHNID
	3	3	0	0.21	0.210	0.191	0.191	1.10	NAWQ ACTUAL
	3	3	0	0.21	0.210	0.124	0.124	1.70	LCV FISH
	3	3	0	0.21	0.210	0.117	0.117	1.80	LTV EC20 FISH
	3	3	0	0.21	0.210	0.049	0.049	4.30	EC20 POPULATION
	4	4	0	0.21	0.210	16.154	16.154	0.01	SS EC20
	4	4	0	0.21	0.210	1.400	1.400	0.15	LCV DAPHNIDS
	4	4	0	0.21	0.210	0.280	0.280	0.75	LTV EC20 DAPHNID
	4	4	0	0.21	0.210	0.191	0.191	1.10	NAWQ ACTUAL
	4	4	0	0.21	0.210	0.124	0.124	1.70	LCV FISH
	4	4	0	0.21	0.210	0.117	0.117	1.80	LTV EC20 FISH
	4	4	0	0.21	0.210	0.049	0.049	4.30	EC20 POPULATION
	5	5	0	0.21	0.210	16.154	16.154	0.01	SS EC20
	5	5	0	0.21	0.210	1.400	1.400	0.15	LCV DAPHNIDS
	5	5	0	0.21	0.210	0.280	0.280	0.75	LTV EC20 DAPHNID
	5	5	0	0.21	0.210	0.191	0.191	1.10	NAWQ ACTUAL
	5	5	0	0.21	0.210	0.124	0.124	1.70	LCV FISH
	5	5	0	0.21	0.210	0.117	0.117	1.80	LTV EC20 FISH
	5	5	0	0.21	0.210	0.049	0.049	4.30	EC20 POPULATION
	10	2	0	0.21	0.210	16.154	16.154	0.01	SS EC20
	10	2	0	0.21	0.210	1.400	1.400	0.15	LCV DAPHNIDS
	10	2	0	0.21	0.210	0.280	0.280	0.75	LTV EC20 DAPHNID
	10	2	0	0.21	0.210	0.191	0.191	1.10	NAWQ ACTUAL
	10	2	0	0.21	0.210	0.124	0.124	1.70	LCV FISH
	10	2	0	0.21	0.210	0.117	0.117	1.80	LTV EC20 FISH
	10	2	0	0.21	0.210	0.049	0.049	4.30	EC20 POPULATION
	13	1	0	0.21	0.210	16.154	16.154	0.01	SS EC20
	13	1	0	0.21	0.210	1.400	1.400	0.15	LCV DAPHNIDS
	13	1	0	0.21	0.210	0.280	0.280	0.75	LTV EC20 DAPHNID
	13	1	0	0.21	0.210	0.191	0.191	1.10	NAWQ ACTUAL
	13	1	0	0.21	0.210	0.124	0.124	1.70	LCV FISH
	13	1	0	0.21	0.210	0.117	0.117	1.80	LTV EC20 FISH
	13	1	0	0.21	0.210	0.049	0.049	4.30	EC20 POPULATION

Table P.2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
CHROMIUM	1	3	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	1	3	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	1	3	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	1	3	0	9.80	9.800	0.891	0.891	11.00	NAMQ ACTUAL
	1	3	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	1	3	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	1	3	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION
	1	3	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	2	2	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	2	2	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	2	2	0	9.80	9.800	0.891	0.891	11.00	NAMQ ACTUAL
	2	2	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	2	2	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	2	2	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION
	2	2	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	3	4	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	3	4	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	3	4	0	9.80	9.800	0.891	0.891	11.00	NAMQ ACTUAL
	3	4	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	3	4	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	3	4	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION
	3	4	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	4	2	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	4	2	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	4	2	0	9.80	9.800	0.891	0.891	11.00	NAMQ ACTUAL
	4	2	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	4	2	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	4	2	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION
	4	2	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	5	4	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	5	4	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	5	4	0	9.80	9.800	0.891	0.891	11.00	NAMQ ACTUAL
	5	4	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	5	4	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	5	4	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION
	5	4	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	10	2	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	10	2	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	10	2	0	9.80	9.800	0.891	0.891	11.00	NAMQ ACTUAL
	10	2	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	10	2	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	10	2	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION

Table F.2 (continued)

Analyte	Reach	Number analyzed in rec (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
CHROMIUM (continued)	13	1	0	9.80	9.800	36.842	36.842	0.27	SS EC20
	13	1	0	9.80	9.800	19.600	19.600	0.50	LTV EC20 DAPHNID
	13	1	0	9.80	9.800	1.598	1.598	6.13	LCV DAPHNIDS
	13	1	0	9.80	9.800	0.891	0.891	11.00	MANQ ACTUAL
	13	1	0	9.80	9.800	0.192	0.192	51.00	LTV EC20 FISH
	13	1	0	9.80	9.800	0.134	0.134	73.18	LCV FISH
	13	1	0	9.80	9.800	0.031	0.031	316.00	EC20 POPULATION
	1	3	0	4.40	4.400	21.463	21.463	0.20	LTV EC20 DAPHNID
	1	3	0	4.40	4.400	19.130	19.130	0.23	LCV DAPHNIDS
	1	3	0	4.40	4.400	16.923	16.923	0.26	SS EC20
COPPER	1	3	0	4.40	4.400	1.136	1.136	3.87	LCV FISH
	1	3	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	1	3	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	1	3	0	4.40	4.400	0.367	0.367	12.00	MANQ ACTUAL
	1	3	0	4.40	4.400	0.367	0.367	0.20	LTV EC20 DAPHNID
	2	2	0	4.40	4.400	21.463	21.463	0.23	LCV DAPHNIDS
	2	2	0	4.40	4.400	19.130	19.130	0.26	SS EC20
	2	2	0	4.40	4.400	16.923	16.923	3.87	LCV FISH
	2	2	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	2	2	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	2	2	0	4.40	4.400	0.367	0.367	12.00	MANQ ACTUAL
	2	2	0	4.40	4.400	0.367	0.367	0.20	LTV EC20 DAPHNID
	3	4	0	4.40	4.400	21.463	21.463	0.23	LCV DAPHNIDS
	3	4	0	4.40	4.400	19.130	19.130	0.26	SS EC20
	3	4	0	4.40	4.400	16.923	16.923	3.87	LCV FISH
	3	4	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	3	4	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	3	4	0	4.40	4.400	0.367	0.367	12.00	MANQ ACTUAL
	3	4	0	4.40	4.400	0.367	0.367	0.20	LTV EC20 DAPHNID
	4	4	0	4.40	4.400	21.463	21.463	0.23	LCV DAPHNIDS
	4	4	0	4.40	4.400	19.130	19.130	0.26	SS EC20
	4	4	0	4.40	4.400	16.923	16.923	3.87	LCV FISH
	4	4	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	4	4	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	4	4	0	4.40	4.400	0.367	0.367	12.00	MANQ ACTUAL
	4	4	0	4.40	4.400	0.367	0.367	0.20	LTV EC20 DAPHNID
	5	5	0	4.40	4.400	21.463	21.463	0.23	LCV DAPHNIDS
	5	5	0	4.40	4.400	19.130	19.130	0.26	SS EC20
	5	5	0	4.40	4.400	16.923	16.923	3.87	LCV FISH
	5	5	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	5	5	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	5	5	0	4.40	4.400	0.367	0.367	12.00	MANQ ACTUAL

Table F.2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
COPPER (continued)	10	2	0	4.40	4.400	21.463	21.463	0.20	LTV EC20 DAPHNID
	10	2	0	4.40	4.400	19.130	19.130	0.23	LCV DAPHNIDS
	10	2	0	4.40	4.400	16.923	16.923	0.26	SS EC20
	10	2	0	4.40	4.400	1.136	1.136	3.87	LCV FISH
	10	2	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	10	2	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	10	2	0	4.40	4.400	0.367	0.367	12.00	MAWQ ACTUAL
	10	2	0	4.40	4.400	21.463	21.463	0.20	LTV EC20 DAPHNID
	13	1	0	4.40	4.400	19.130	19.130	0.23	LCV DAPHNIDS
	13	1	0	4.40	4.400	16.923	16.923	0.26	SS EC20
	13	1	0	4.40	4.400	1.136	1.136	3.87	LCV FISH
	13	1	0	4.40	4.400	0.880	0.880	5.00	LTV EC20 FISH
	13	1	0	4.40	4.400	0.512	0.512	8.60	EC20 POPULATION
	13	1	0	4.40	4.400	0.367	0.367	12.00	MAWQ ACTUAL
	1	2	0	9.00	495.570	0.562	30.973	16.00	LTV EC20 DAPHNID
	1	2	0	9.00	495.570	0.057	3.137	158.00	LCV DAPHNIDS
	1	2	0	9.00	495.570	0.009	0.496	1000.00	MAWQ ACTUAL
	3	2	0	6.70	10.176	0.419	0.636	16.00	LTV EC20 DAPHNID
	3	4	0	6.70	10.176	0.042	0.064	158.00	LCV DAPHNIDS
IRON	3	4	0	6.70	10.176	0.007	0.010	1000.00	MAWQ ACTUAL
	3	4	0	6.70	10.176	0.350	1.892	16.00	LTV EC20 DAPHNID
	4	2	0	5.60	30.271	0.035	0.192	158.00	LCV DAPHNIDS
	4	2	0	5.60	30.271	0.006	0.030	1000.00	MAWQ ACTUAL
	4	2	0	5.60	30.271	0.350	0.350	16.00	LTV EC20 DAPHNID
	10	2	0	5.60	5.600	0.035	0.035	158.00	LCV DAPHNIDS
	10	2	0	5.60	5.600	0.006	0.006	1000.00	MAWQ ACTUAL
	2	2	0	2.00	5.427	5.714	15.504	0.35	SS EC20
	2	2	0	2.00	5.427	0.625	1.696	3.20	MAWQ ACTUAL
	2	2	0	2.00	5.427	0.163	0.443	12.26	LCV DAPHNIDS
	2	2	0	2.00	5.427	0.106	0.287	18.88	LCV FISH
	2	2	0	2.00	5.427	0.091	0.247	22.00	LTV EC20 FISH
	2	2	0	2.00	5.427	0.028	0.076	71.00	EC20 POPULATION
	3	4	0	0.36	2.460	1.029	7.030	0.35	SS EC20
	3	4	0	0.36	2.460	0.112	0.769	3.20	MAWQ ACTUAL
	3	4	0	0.36	2.460	0.029	0.201	12.26	LCV DAPHNIDS
	3	4	0	0.36	2.460	0.019	0.130	18.88	LCV FISH
	3	4	0	0.36	2.460	0.016	0.112	22.00	LTV EC20 FISH
	3	4	0	0.36	2.460	0.005	0.035	71.00	EC20 POPULATION
LEAD	5	4	0	0.36	2.509	1.029	7.168	0.35	SS EC20
	5	4	0	0.36	2.509	0.112	0.784	3.20	MAWQ ACTUAL
	5	4	0	0.36	2.509	0.029	0.205	12.26	LCV DAPHNIDS

Table F-2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
LEAD (continued)	5	4	0	0.36	2.509	0.019	0.133	18.88	LCV FISH
	5	4	0	0.36	2.509	0.016	0.114	22.00	LTV EC20 FISH
	5	4	0	0.36	2.509	0.005	0.035	71.00	EC20 POPULATION
	10	2	0	0.50	1.871	1.429	5.345	0.35	SS EC20
	10	2	0	0.50	1.871	0.156	0.585	3.20	NAHQ ACTUAL
	10	2	0	0.50	1.871	0.041	0.153	12.26	LCV DAPHNIDS
	10	2	0	0.50	1.871	0.026	0.099	18.88	LCV FISH
	10	2	0	0.50	1.871	0.023	0.085	22.00	LTV EC20 FISH
	10	2	0	0.50	1.871	0.007	0.026	71.00	EC20 POPULATION
	1	3	0	0.20	0.200	666.667	666.667	0.00	NAHQ ADVISORY
	1	3	0	0.20	0.200	6.667	6.667	0.03	LTV EC20 FISH
	1	3	0	0.20	0.200	5.000	5.000	0.04	LCV DAPHNIDS
MERCURY	1	3	0	0.20	0.200	0.714	0.714	0.28	EC20 POPULATION
	1	3	0	0.20	0.200	0.385	0.385	0.52	LCV FISH
	1	3	0	0.20	0.200	0.230	0.230	0.87	LTV EC20 DAPHNID
	1	3	0	0.20	0.200	666.667	666.667	0.00	NAHQ ADVISORY
	2	2	0	0.20	0.200	6.667	6.667	0.03	LTV EC20 FISH
	2	2	0	0.20	0.200	5.000	5.000	0.04	LCV DAPHNIDS
	2	2	0	0.20	0.200	0.714	0.714	0.28	EC20 POPULATION
	2	2	0	0.20	0.200	0.385	0.385	0.52	LCV FISH
	2	2	0	0.20	0.200	0.230	0.230	0.87	LTV EC20 DAPHNID
	2	2	0	0.20	0.200	666.667	666.667	0.00	NAHQ ADVISORY
	4	3	0	0.20	0.200	6.667	6.667	0.03	LTV EC20 FISH
	4	3	0	0.20	0.200	5.000	5.000	0.04	LCV DAPHNIDS
	4	3	0	0.20	0.200	0.714	0.714	0.28	EC20 POPULATION
	4	3	0	0.20	0.200	0.385	0.385	0.52	LCV FISH
	4	3	0	0.20	0.200	0.230	0.230	0.87	LTV EC20 DAPHNID
	4	3	0	0.20	0.200	666.667	666.667	0.00	NAHQ ADVISORY
	5	4	0	0.20	0.200	6.667	6.667	0.03	LTV EC20 FISH
	5	4	0	0.20	0.200	5.000	5.000	0.04	LCV DAPHNIDS
	5	4	0	0.20	0.200	0.714	0.714	0.28	EC20 POPULATION
	5	4	0	0.20	0.200	0.385	0.385	0.52	LCV FISH
	5	4	0	0.20	0.200	0.230	0.230	0.87	LTV EC20 DAPHNID
	5	4	0	0.20	0.200	666.667	666.667	0.00	NAHQ ADVISORY
	10	2	0	0.20	0.200	6.667	6.667	0.03	LTV EC20 FISH
	10	2	0	0.20	0.200	5.000	5.000	0.04	LCV DAPHNIDS
	10	2	0	0.20	0.200	0.714	0.714	0.28	EC20 POPULATION
	10	2	0	0.20	0.200	0.385	0.385	0.52	LCV FISH
	10	2	0	0.20	0.200	0.230	0.230	0.87	LTV EC20 DAPHNID
	13	1	0	0.20	0.200	666.667	666.667	0.00	NAHQ ADVISORY
	13	1	0	0.20	0.200	6.667	6.667	0.03	LTV EC20 FISH
	13	1	0	0.20	0.200	5.000	5.000	0.04	LCV DAPHNIDS

Table F.2 (continued)

Analyte	Resch	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
MERCURY (continued)	13	1	0	0.20	0.200	0.714	0.714	0.28	EC20 POPULATION
	13	1	0	0.20	0.200	0.385	0.385	0.52	LCV FISH
	13	1	0	0.20	0.200	0.230	0.230	0.87	LTV EC20 DAPHNID
NICKEL	1	3	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	1	3	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	1	3	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	1	3	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID
	1	3	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	1	3	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	1	3	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	1	3	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	2	2	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	2	2	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	2	2	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID
	2	2	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	2	2	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	2	2	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	2	2	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	3	4	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	3	4	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	3	4	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID
	3	4	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	3	4	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	3	4	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	3	4	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	4	2	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	4	2	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	4	2	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID
	4	2	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	4	2	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	4	2	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	4	2	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	5	4	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	5	4	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	5	4	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID
	5	4	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	5	4	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	5	4	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	10	2	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	10	2	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	10	2	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	10	2	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID

Table F-2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
NICKEL (continued)	10	2	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	10	2	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	10	2	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	13	1	0	19.00	19.000	3.800	3.800	5.00	LCV DAPHNIDS
	13	1	0	19.00	19.000	1.727	1.727	11.00	SS EC20
	13	1	0	19.00	19.000	0.543	0.543	35.00	LCV FISH
	13	1	0	19.00	19.000	0.422	0.422	45.00	LTV EC20 DAPHNID
	13	1	0	19.00	19.000	0.306	0.306	62.00	LTV EC20 FISH
	13	1	0	19.00	19.000	0.119	0.119	160.00	NAHQ ACTUAL
	13	1	0	19.00	19.000	0.088	0.088	215.00	EC20 POPULATION
	1	3	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	1	3	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID
	1	3	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	1	3	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	1	3	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	1	3	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
SELENIUM	1	3	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	2	2	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID
	2	2	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	2	2	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	2	2	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	2	2	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
	3	4	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	3	4	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID
	3	4	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	3	4	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	3	4	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	3	4	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
	3	4	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	4	3	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID
	4	3	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	4	3	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	4	3	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	4	3	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
	4	3	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	5	4	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID
	5	4	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	5	4	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	5	4	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	5	4	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
	5	4	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	10	2	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID

Table F.2 (continued)

Analyte	Resch	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
SELENIUM (continued)	10	2	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	10	2	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	10	2	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	10	2	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
	13	1	0	0.97	0.970	0.373	0.373	2.60	SS EC20
	13	1	0	0.97	0.970	0.039	0.039	25.00	LTV EC20 DAPHNID
	13	1	0	0.97	0.970	0.028	0.028	35.00	NAHQ ACTUAL
	13	1	0	0.97	0.970	0.024	0.024	40.00	LTV EC20 FISH
	13	1	0	0.97	0.970	0.011	0.011	88.32	LCV FISH
	13	1	0	0.97	0.970	0.011	0.011	91.65	LCV DAPHNIDS
	1	3	0	0.74	0.740	6.167	6.167	0.12	LCV FISH
	1	3	0	0.74	0.740	5.286	5.286	0.14	SS EC20
	1	3	0	0.74	0.740	3.700	3.700	0.20	LTV EC20 FISH
	1	3	0	0.74	0.740	2.960	2.960	0.25	NAHQ ADVISORY
	1	3	0	0.74	0.740	2.313	2.313	0.32	EC20 POPULATION
	1	3	0	0.74	0.740	1.321	1.321	0.56	LTV EC20 DAPHNID
SILVER	1	3	0	0.74	0.740	0.285	0.285	2.60	LCV DAPHNIDS
	1	3	0	0.74	0.740	6.167	6.167	0.12	LCV FISH
	2	2	0	0.74	0.740	5.286	5.286	0.14	SS EC20
	2	2	0	0.74	0.740	3.700	3.700	0.20	LTV EC20 FISH
	2	2	0	0.74	0.740	2.960	2.960	0.25	NAHQ ADVISORY
	2	2	0	0.74	0.740	2.313	2.313	0.32	EC20 POPULATION
	2	2	0	0.74	0.740	1.321	1.321	0.56	LTV EC20 DAPHNID
	2	2	0	0.74	0.740	0.285	0.285	2.60	LCV DAPHNIDS
	4	3	0	0.74	0.740	6.167	6.167	0.12	LCV FISH
	4	3	0	0.74	0.740	5.286	5.286	0.14	SS EC20
	4	3	0	0.74	0.740	3.700	3.700	0.20	LTV EC20 FISH
	4	3	0	0.74	0.740	2.960	2.960	0.25	NAHQ ADVISORY
	4	3	0	0.74	0.740	2.313	2.313	0.32	EC20 POPULATION
	4	3	0	0.74	0.740	1.321	1.321	0.56	LTV EC20 DAPHNID
	4	3	0	0.74	0.740	0.285	0.285	2.60	LCV DAPHNIDS
	5	4	0	0.74	0.740	6.167	6.167	0.12	LCV FISH
	5	4	0	0.74	0.740	5.286	5.286	0.14	SS EC20
	5	4	0	0.74	0.740	3.700	3.700	0.20	LTV EC20 FISH
	5	4	0	0.74	0.740	2.960	2.960	0.25	NAHQ ADVISORY
	5	4	0	0.74	0.740	2.313	2.313	0.32	EC20 POPULATION
	5	4	0	0.74	0.740	1.321	1.321	0.56	LTV EC20 DAPHNID
	5	4	0	0.74	0.740	0.285	0.285	2.60	LCV DAPHNIDS
	10	2	0	0.74	0.740	6.167	6.167	0.12	LCV FISH
	10	2	0	0.74	0.740	5.286	5.286	0.14	SS EC20
	10	2	0	0.74	0.740	3.700	3.700	0.20	LTV EC20 FISH
	10	2	0	0.74	0.740	2.960	2.960	0.25	NAHQ ADVISORY

Table F-2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
SILVER (continued)	10	2	0	0.74	0.740	2.313	2.313	0.32	EC20 POPULATION
	10	2	0	0.74	0.740	1.321	1.321	0.56	LTV EC20 DAPHNID
	10	2	0	0.74	0.740	0.285	0.285	2.60	LCV DAPHNIDS
	13	1	0	0.74	0.740	6.167	6.167	0.12	LCV FISH
	13	1	0	0.74	0.740	5.286	5.286	0.14	SS EC20
	13	1	0	0.74	0.740	3.700	3.700	0.20	LTV EC20 FISH
	13	1	0	0.74	0.740	2.960	2.960	0.25	NAHQ ADVISORY
	13	1	0	0.74	0.740	2.313	2.313	0.32	EC20 POPULATION
	13	1	0	0.74	0.740	1.321	1.321	0.56	LTV EC20 DAPHNID
	13	1	0	0.74	0.740	0.285	0.285	2.60	LCV DAPHNIDS
	1	3	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	1	3	0	2.70	2.700	0.047	0.047	57.00	LCV FISH
	1	3	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
THALLIUM	1	3	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	1	3	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	1	3	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	1	3	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	2	2	0	2.70	2.700	0.047	0.047	57.00	LCV FISH
	2	2	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
	2	2	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	2	2	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	2	2	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	3	3	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	3	3	0	2.70	2.700	0.047	0.047	57.00	LCV FISH
	3	3	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
	3	3	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	3	3	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	3	3	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	4	4	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	4	4	0	2.70	2.700	0.047	0.047	57.00	LCV FISH
	4	4	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
	4	4	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	4	4	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	4	4	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	5	5	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	5	5	0	2.70	2.700	0.047	0.047	57.00	LCV FISH
	5	5	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
	5	5	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	5	5	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	5	5	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	10	2	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	10	2	0	2.70	2.700	0.047	0.047	57.00	LCV FISH

Table F.2 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
THALLIUM (continued)	10	2	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
	10	2	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	10	2	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	10	2	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	13	1	0	2.70	2.700	1.080	1.080	2.50	NAHQ ADVISORY
	13	1	0	2.70	2.700	0.047	0.047	57.00	LCV FISH
	13	1	0	2.70	2.700	0.042	0.042	64.00	LTV EC20 DAPHNID
	13	1	0	2.70	2.700	0.040	0.040	67.00	EC20 POPULATION
	13	1	0	2.70	2.700	0.033	0.033	81.00	LTV EC20 FISH
	13	1	0	2.70	2.700	0.020	0.020	135.00	LCV DAPHNIDS
	1	3	0	3.40	22.431	0.162	1.068	21.00	SS EC20
	1	3	0	3.40	22.431	0.093	0.616	36.41	LCV FISH
	1	3	0	3.40	22.431	0.073	0.480	46.73	LCV DAPHNIDS
ZINC	1	3	0	3.40	22.431	0.072	0.477	47.00	LTV EC20 FISH
	1	3	0	3.40	22.431	0.042	0.280	80.00	EC20 POPULATION
	1	3	0	3.40	22.431	0.031	0.204	110.00	NAHQ ACTUAL
	1	3	0	3.40	62.337	0.162	2.968	21.00	SS EC20
	2	2	0	3.40	62.337	0.093	1.712	36.41	LCV FISH
	2	2	0	3.40	62.337	0.073	1.334	46.73	LCV DAPHNIDS
	2	2	0	3.40	62.337	0.072	1.326	47.00	LTV EC20 FISH
	2	2	0	3.40	62.337	0.042	0.779	80.00	EC20 POPULATION
	2	2	0	3.40	62.337	0.031	0.567	110.00	NAHQ ACTUAL
	2	2	0	3.40	17.145	0.162	0.816	21.00	SS EC20
	3	4	0	3.40	17.145	0.093	0.471	36.41	LCV FISH
	3	4	0	3.40	17.145	0.073	0.367	46.73	LCV DAPHNIDS
	3	4	0	3.40	17.145	0.072	0.365	47.00	LTV EC20 FISH
	3	4	0	3.40	17.145	0.042	0.214	80.00	EC20 POPULATION
	3	4	0	3.40	17.145	0.031	0.156	110.00	NAHQ ACTUAL
	4	4	0	3.40	3.400	0.162	0.162	21.00	SS EC20
	4	2	0	3.40	3.400	0.093	0.093	36.41	LCV FISH
	4	2	0	3.40	3.400	0.073	0.073	46.73	LCV DAPHNIDS
	4	2	0	3.40	3.400	0.072	0.072	47.00	LTV EC20 FISH
	4	2	0	3.40	3.400	0.042	0.042	80.00	EC20 POPULATION
	4	2	0	3.40	3.400	0.031	0.031	110.00	NAHQ ACTUAL
	4	4	0	3.40	17.003	0.162	0.810	21.00	SS EC20
	5	4	0	3.40	17.003	0.093	0.467	36.41	LCV FISH
	5	4	0	3.40	17.003	0.073	0.364	46.73	LCV DAPHNIDS
	5	4	0	3.40	17.003	0.072	0.362	47.00	LTV EC20 FISH
	5	4	0	3.40	17.003	0.042	0.213	80.00	EC20 POPULATION
	5	4	0	3.40	17.003	0.031	0.155	110.00	NAHQ ACTUAL
	5	4	0	3.40	13.000	0.619	0.619	21.00	SS EC20
	13	1	0	13.00	13.000	0.357	0.357	36.41	LCV FISH

Table F-2 (continued)

Analyte	Reech	Number analyzed in reech (water)	Number detects in reech (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
ZINC (continued)	13	1	0	13.00	13.000	0.278	0.278	46.73	LCV DAPHNIDS
	13	1	0	13.00	13.000	0.277	0.277	47.00	LTV EC20 FISH
	13	1	0	13.00	13.000	0.163	0.163	80.00	EC20 POPULATION
	13	1	0	13.00	13.000	0.118	0.118	110.00	NAMQ ACTUAL

^aThe maximum detection limit was used in cases where the UCL was missing (i.e., where N = 1).

Table F.3. Mean and upper confidence limit (UCL) concentrations of chemicals detected in NPDES water samples (ug/L), toxicological benchmarks (ug/L), and quotients for aquatic life

Analyte	Resch	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
1,1,1-TRICHLOROETHANE	3	704	200	4.47	5.02	3.19	3.59	1.40		MAW ADVISORY
	3	704	200	4.47	5.02	0.02	0.02	251.00		EC20 POPULATION
	3	704	200	4.47	5.02	0.00	0.00	1300.00		LTV EC20 DAPHNID
	3	704	200	4.47	5.02	0.00	0.00	1770.00	*	LCV DAPHNIDS
	3	704	200	4.47	5.02	0.00	0.00	2457.00	*	LTV EC20 FISH
1,1-DICHLOROETHANE	3	705	41	4.87	5.00	9.55	9.80	0.51		MAW ADVISORY
	3	705	41	4.87	5.00	0.00	0.00	1585.00	*	EC20 POPULATION
	3	705	41	4.87	5.00	0.00	0.00	8219.00	*	LTV EC20 FISH
	3	705	41	4.87	5.00	0.00	0.00	14680.00	*	LCV FISH
	3	705	3	5.08	5.25	0.04	0.04	144.00		MAW ADVISORY
1,2-DICHLOROETHANE	3	705	3	5.08	5.25	0.00	0.00	1259.00		EC20 POPULATION
	3	705	3	5.08	5.25	0.00	0.00	11000.00	<	LTV EC20 DAPHNID
	3	705	3	5.08	5.25	0.00	0.00	15200.00		LCV DAPHNIDS
	3	705	3	5.08	5.25	0.00	0.00	29000.00		LTV EC20 FISH
	3	705	3	5.08	5.25	0.00	0.00	41364.00		LCV FISH
ACETONE	2	64	8	9.45	9.45	0.01	0.01	770.00		MAW ADVISORY
	2	64	8	9.45	9.45	0.00	0.00	23714.00	*	EC20 POPULATION
	2	64	8	9.45	9.45	0.00	0.00	161867.00	*	LTV EC20 FISH
	2	64	8	9.45	9.45	0.00	0.00	507640.00	*	LCV FISH
	2	64	8	9.45	9.45	0.00	0.00	3114182.00	*	LCV DAPHNIDS
ALUMINUM	1	64	52	618.34	618.34	8.24	8.24	75.00		SS EC20
	1	64	52	618.34	618.34	7.11	7.11	87.00		MAW ACTUAL
	1	64	52	618.34	618.34	1.15	1.15	540.00		LTV EC20 DAPHNID
	1	64	52	618.34	618.34	0.33	0.33	1900.00		LCV DAPHNIDS
	1	64	52	618.34	618.34	0.13	0.13	4700.00		LTV EC20 FISH
	1	64	52	618.34	618.34	0.11	0.11	5800.00		LCV FISH
	2	186	186	343.50	487.01	4.58	6.49	75.00		SS EC20
	2	186	186	343.50	487.01	3.95	5.60	87.00		MAW ACTUAL
	2	186	186	343.50	487.01	0.64	0.90	540.00		LTV EC20 DAPHNID
	2	186	186	343.50	487.01	0.18	0.26	1900.00		LCV DAPHNIDS
	2	186	186	343.50	487.01	0.07	0.10	4700.00		LTV EC20 FISH
	2	186	186	343.50	487.01	0.06	0.08	5800.00		LCV FISH
	2	186	186	343.50	487.01	0.06	0.08	75.00		SS EC20
	3	497	318	1774.17	3807.40	23.66	50.77	87.00		MAW ACTUAL
	3	497	318	1774.17	3807.40	20.39	43.76	540.00		LTV EC20 DAPHNID
	3	497	318	1774.17	3807.40	3.29	7.05	1900.00		LCV DAPHNIDS
	3	497	318	1774.17	3807.40	0.93	2.00	4700.00		LTV EC20 FISH
	3	497	318	1774.17	3807.40	0.38	0.81	5800.00		LCV FISH
	3	497	318	1774.17	3807.40	0.31	0.66	5800.00		LCV FISH

Table F.3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
ALUMINUM (continued)	4	1	1	540.00	540.00	7.20	7.20	75.00		SS EC20
	4	1	1	540.00	540.00	6.21	6.21	87.00		NAHQ ACTUAL
	4	1	1	540.00	540.00	1.00	1.00	540.00		LTV EC20 DAPHNID
	4	1	1	540.00	540.00	0.28	0.28	1900.00		LCV DAPHNIDS
	4	1	1	540.00	540.00	0.11	0.11	4700.00		LTV EC20 FISH
	4	1	1	540.00	540.00	0.09	0.09	5800.00		LCV FISH
ANTHRACENE	3	179	1	10.04	10.13	1004347.83	1012927.74	0.00		NAHQ ADVISORY
	3	179	1	10.04	10.13	111.59	112.55	0.09	*	LCV FISH
	3	179	1	10.04	10.13	28.94	28.94	0.35	*	LTV EC20 FISH
	3	179	1	10.04	10.13	4.78	4.82	2.10	<	LCV DAPHNIDS
	3	179	1	10.04	10.13	1.22	1.24	8.20	>	LTV EC20 DAPHNID
	1	64	2	50.87	50.87	2.83	2.83	18.00		NAHQ ADVISORY
	1	64	2	50.87	50.87	0.64	0.64	79.00		EC20 POPULATION
	1	64	2	50.87	50.87	0.03	0.03	1600.00		LCV FISH
	1	64	2	50.87	50.87	0.03	0.03	1900.00		LTV EC20 DAPHNID
	1	64	2	50.87	50.87	0.02	0.02	2310.00		LTV EC20 FISH
	1	64	2	50.87	50.87	0.01	0.01	5400.00		LCV DAPHNIDS
	3	210	5	1.01	1.04	50.45	51.91	0.02		NAHQ ADVISORY
	3	210	5	1.01	1.04	1.94	2.00	0.52		LTV EC20 FISH
	3	210	5	1.01	1.04	1.60	1.65	0.63		EC20 POPULATION
	3	210	5	1.01	1.04	0.84	0.87	1.00		LCV FISH
	3	210	5	1.01	1.04	0.48	0.49	1.20		LTV EC20 DAPHNID
	3	210	5	1.01	1.04	0.71	0.83	2.10		LCV DAPHNIDS
	1	219	56	39.09	45.71	0.21	0.24	55.00		SS EC20
	1	219	56	39.09	45.71	0.06	0.07	190.00		NAHQ ACTUAL
	1	219	56	39.09	45.71	0.04	0.05	633.00		LTV EC20 DAPHNID
	1	219	56	39.09	45.71	0.02	0.02	914.10		LCV DAPHNIDS
	1	219	56	39.09	45.71	0.02	0.02	1995.00		EC20 POPULATION
	1	219	56	39.09	45.71	0.02	0.02	2130.00		LTV EC20 FISH
	1	219	56	39.09	45.71	0.01	0.02	2982.00		LCV FISH
	3	412	26	8.78	16.14	0.16	0.29	55.00		SS EC20
	3	412	26	8.78	16.14	0.05	0.08	190.00		NAHQ ACTUAL
	3	412	26	8.78	16.14	0.01	0.03	633.00		LTV EC20 DAPHNID
	3	412	26	8.78	16.14	0.01	0.02	914.10		LCV DAPHNIDS
	3	412	26	8.78	16.14	0.00	0.01	1995.00		EC20 POPULATION
	3	412	26	8.78	16.14	0.00	0.01	2130.00		LTV EC20 FISH
	3	412	26	8.78	16.14	0.00	0.01	2982.00		LCV FISH

Table F3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
BARIUM	1	64	60	32.97	32.97	0.30	0.30	109.00	*	MAW ADVISORY
	1	64	60	32.97	32.97	0.00	0.00	20336.00		LCV DAPHNIDS
	2	2	2	30.50	33.00	0.28	0.30	109.00	*	MAW ADVISORY
	2	2	2	30.50	33.00	0.00	0.00	20336.00		LCV DAPHNIDS
	3	192	9	80.00	114.41	0.73	1.05	109.00	*	MAW ADVISORY
	3	192	9	80.00	114.41	0.00	0.01	20336.00		LCV DAPHNIDS
	4	1	1	33.00	33.00	0.30	0.30	109.00	*	MAW ADVISORY
	4	1	1	33.00	33.00	0.00	0.00	20336.00		LCV DAPHNIDS
	3	705	9	4.98	5.00	1.56	1.56	3.20		MAW ADVISORY
	3	705	9	4.98	5.00	0.24	0.24	21.00		LTV EC20 FISH
BENZENE	3	705	9	4.98	5.00	0.02	0.02	229.00	*	EC20 POPULATION
	3	705	9	4.98	5.00	0.02	0.02	299.00	*	LCV FISH
	3	705	9	4.98	5.00	0.00	0.00	98000.00	>	LCV DAPHNIDS
	3	179	1	10.05	10.15	3722.11	3758.42	0.00	*	MAW ADVISORY
	3	179	1	10.05	10.15	15.46	15.61	0.65	*	LCV DAPHNIDS
BERYLLIUM	1	64	4	0.87	0.87	1.42	1.42	0.61		MAW ADVISORY
	1	64	4	0.87	0.87	0.23	0.23	3.80		LTV EC20 DAPHNID
	1	64	4	0.87	0.87	0.16	0.16	5.30		LCV DAPHNIDS
	1	64	4	0.87	0.87	0.04	0.04	21.00	*	EC20 POPULATION
	1	64	4	0.87	0.87	0.02	0.02	57.00	*	LCV FISH
	1	64	4	0.87	0.87	0.01	0.01	148.00	*	LTV EC20 FISH
	3	398	13	0.79	1.10	1.30	1.81	0.61		MAW ADVISORY
	3	398	13	0.79	1.10	0.21	0.29	3.80		LTV EC20 DAPHNID
	3	398	13	0.79	1.10	0.15	0.21	5.30		LCV DAPHNIDS
	3	398	13	0.79	1.10	0.04	0.05	21.00	*	EC20 POPULATION
	3	398	13	0.79	1.10	0.01	0.02	57.00	*	LCV FISH
	3	398	13	0.79	1.10	0.01	0.01	148.00	*	LTV EC20 FISH
	3	179	28	10.19	10.56	1018.86	1055.79	0.01		MAW ADVISORY
	3	179	28	10.19	10.56	3.40	3.52	3.00	<	LCV DAPHNIDS
	3	179	28	10.19	10.56	3.40	3.52	3.00	<	LTV EC20 DAPHNID
BIS(2-ETHYLNEXYL)PHTHALATE	3	179	28	10.19	10.56	1.21	1.26	8.40		LCV FISH
	3	179	28	10.19	10.56	0.20	0.21	50.00		EC20 POPULATION
	3	179	28	10.19	10.56	0.19	0.20	54.00	>	LTV EC20 FISH
	3	192	191	69.67	176.38	9.95	25.20	7.00		LTV EC20 DAPHNID
	4	1	1	23.00	23.00	3.29	3.29	7.00		LTV EC20 DAPHNID
BCRON	3	192	191	69.67	176.38	9.95	25.20	7.00		LTV EC20 DAPHNID
	4	1	1	23.00	23.00	3.29	3.29	7.00		LTV EC20 DAPHNID

Table F-3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
CADMIUM	1	173	52	4.51	5.35	347.27	411.76	0.01		SS EC20
	1	173	52	4.51	5.35	30.10	35.69	0.15		LCV DAPHNIDS
	1	173	52	4.51	5.35	6.02	7.14	0.75		LTV EC20 DAPHNID
	1	173	52	4.51	5.35	4.10	4.87	1.10		NAHQ ACTUAL
	1	173	52	4.51	5.35	2.66	3.15	1.70		LCV FISH
	1	173	52	4.51	5.35	2.51	2.97	1.80		LTV EC20 FISH
	1	173	52	4.51	5.35	1.05	1.24	4.30		EC20 POPULATION
	1	173	52	4.51	5.35	165.35	187.09	0.01		SS EC20
	3	615	17	2.15	2.43	14.33	16.21	0.15		LCV DAPHNIDS
	3	615	17	2.15	2.43	2.87	3.24	0.75		LTV EC20 DAPHNID
	3	615	17	2.15	2.43	1.95	2.21	1.10		NAHQ ACTUAL
	3	615	17	2.15	2.43	1.26	1.43	1.70		LCV FISH
	3	615	17	2.15	2.43	1.19	1.35	1.80		LTV EC20 FISH
	3	615	17	2.15	2.43	0.50	0.57	4.30		EC20 POPULATION
CARBON DISULFIDE	2	32	2	4.69	4.69	5.15	5.15	0.91	*	NAHQ ADVISORY
	2	32	2	4.69	4.69	0.02	0.02	244.00	*	LCV DAPHNIDS
	2	32	2	4.69	4.69	0.00	0.00	1000.00	*	EC20 POPULATION
	2	32	2	4.69	4.69	0.00	0.00	5719.00	*	LTV EC20 FISH
	2	32	2	4.69	4.69	0.00	0.00	9538.00	*	LCV FISH
CHLOROBENZENE	3	705	1	5.00	5.00	0.43	0.43	11.60		NAHQ ADVISORY
	3	705	1	5.00	5.00	0.03	0.03	165.00		EC20 POPULATION
	3	705	1	5.00	5.00	0.00	0.00	1002.00	*	LTV EC20 FISH
	3	705	1	5.00	5.00	0.00	0.00	1203.00	*	LCV FISH
	3	705	1	5.00	5.00	0.00	0.00	15042.00	*	LCV DAPHNIDS
CHLOROFORM	2	9	1	4.50	5.00	0.32	0.36	14.00		NAHQ ADVISORY
	2	9	1	4.50	5.00	0.01	0.01	562.00	*	EC20 POPULATION
	2	9	1	4.50	5.00	0.00	0.00	3578.00	*	LCV FISH
	2	9	1	4.50	5.00	0.00	0.00	4483.00	*	LCV DAPHNIDS
	2	9	1	4.50	5.00	0.00	0.00	8400.00	*	LTV EC20 FISH
	2	9	1	4.50	5.00	0.30	0.38	14.00		NAHQ ADVISORY
	3	705	601	4.25	5.26	0.01	0.01	562.00	*	EC20 POPULATION
	3	705	601	4.25	5.26	0.00	0.00	3578.00	*	LCV FISH
	3	705	601	4.25	5.26	0.00	0.00	4483.00	*	LCV DAPHNIDS
	3	705	601	4.25	5.26	0.00	0.00	8400.00	*	LTV EC20 FISH
	3	705	601	4.25	5.26	0.00	0.00			

Table F-3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
CHROMIUM	1	211	96	6.32	6.96	23.77	26.16	0.27		SS EC20
	1	211	96	6.32	6.96	12.65	13.92	0.50		LTV EC20 DAPHNID
	1	211	96	6.32	6.96	1.03	1.13	6.13		LCV DAPHNIDS
	1	211	96	6.32	6.96	0.57	0.63	11.00		NAHQ ACTUAL
	1	211	96	6.32	6.96	0.12	0.14	51.00		LTV EC20 FISH
	1	211	96	6.32	6.96	0.09	0.10	73.18		LCV FISH
	1	211	96	6.32	6.96	0.02	0.02	316.00		EC20 POPULATION
	3	883	89	10.92	11.86	41.05	44.60	0.27		SS EC20
	3	883	89	10.92	11.86	21.84	23.72	0.50		LTV EC20 DAPHNID
	3	883	89	10.92	11.86	1.78	1.93	6.13		LCV DAPHNIDS
	3	883	89	10.92	11.86	0.99	1.08	11.00		NAHQ ACTUAL
	3	883	89	10.92	11.86	0.21	0.23	51.00		LTV EC20 FISH
	3	883	89	10.92	11.86	0.15	0.16	73.18		LCV FISH
	3	883	89	10.92	11.86	0.03	0.04	316.00		EC20 POPULATION
	4	184	39	13.54	16.20	50.90	60.90	0.27		SS EC20
	4	184	39	13.54	16.20	27.08	32.40	0.50		LTV EC20 DAPHNID
	4	184	39	13.54	16.20	2.21	2.64	6.13		LCV DAPHNIDS
	4	184	39	13.54	16.20	1.23	1.47	11.00		NAHQ ACTUAL
	4	184	39	13.54	16.20	0.27	0.32	51.00		LTV EC20 FISH
	4	184	39	13.54	16.20	0.19	0.22	73.18		LCV FISH
	4	184	39	13.54	16.20	0.04	0.05	316.00		EC20 POPULATION
	13	8	1	11.00	11.00	41.35	41.35	0.27		SS EC20
	13	8	1	11.00	11.00	22.00	22.00	0.50		LTV EC20 DAPHNID
	13	8	1	11.00	11.00	1.79	1.79	6.13		LCV DAPHNIDS
	13	8	1	11.00	11.00	1.00	1.00	11.00		NAHQ ACTUAL
	13	8	1	11.00	11.00	0.22	0.22	51.00		LTV EC20 FISH
	13	8	1	11.00	11.00	0.15	0.15	73.18		LCV FISH
	13	8	1	11.00	11.00	0.03	0.03	316.00		EC20 POPULATION
COPPER	1	211	66	6.31	9.34	30.79	45.56	0.20		LTV EC20 DAPHNID
	1	211	66	6.31	9.34	27.44	40.61	0.23		LCV DAPHNIDS
	1	211	66	6.31	9.34	24.27	35.92	0.26		SS EC20
	1	211	66	6.31	9.34	1.63	2.41	3.87		LCV FISH
	1	211	66	6.31	9.34	1.26	1.87	5.00		LTV EC20 FISH
	1	211	66	6.31	9.34	0.73	1.09	8.60		EC20 POPULATION
	1	211	66	6.31	9.34	0.53	0.78	12.00		NAHQ ACTUAL
	2	24	3	5.29	5.29	25.79	25.79	0.20		LTV EC20 DAPHNID
	2	24	3	5.29	5.29	22.99	22.99	0.23		LCV DAPHNIDS
	2	24	3	5.29	5.29	20.34	20.34	0.26		SS EC20
	2	24	3	5.29	5.29	1.37	1.37	3.87		LCV FISH
	2	24	3	5.29	5.29	1.06	1.06	5.00		LTV EC20 FISH
	2	24	3	5.29	5.29	0.61	0.61	8.60		EC20 POPULATION
	2	24	3	5.29	5.29	0.44	0.44	12.00		NAHQ ACTUAL

Table F-3 (continued)

Analyte	Reech	Number analyzed in reech (water)	Number detects in reech (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
COPPER (continued)	3	412	332	13.56	19.90	66.15	97.07	0.20		LTV EC20 DAPHNID
	3	412	332	13.56	19.90	58.96	86.52	0.23		LCV DAPHNIDS
	3	412	332	13.56	19.90	52.16	76.54	0.26		SS EC20
	3	412	332	13.56	19.90	3.50	5.14	3.87		LCV FISH
	3	412	332	13.56	19.90	2.71	3.98	5.00		LTV EC20 FISH
	3	412	332	13.56	19.90	1.58	2.31	8.60		EC20 POPULATION
	3	412	332	13.56	19.90	1.13	1.66	12.00		NAHQ ACTUAL
	3	412	332	13.56	19.90	21.83	22.07	0.20		LTV EC20 DAPHNID
	4	32	5	4.47	4.52	19.46	19.67	0.23		LCV DAPHNIDS
	4	32	5	4.47	4.52	17.21	17.40	0.26		SS EC20
	4	32	5	4.47	4.52	1.16	1.17	3.87		LCV FISH
	4	32	5	4.47	4.52	0.89	0.90	5.00		LTV EC20 FISH
	4	32	5	4.47	4.52	0.52	0.53	8.60		EC20 POPULATION
	4	32	5	4.47	4.52	0.37	0.38	12.00		NAHQ ACTUAL
	2	24	1	105.67	105.67	90.31	90.31	1.17		SS EC20
	2	24	1	105.67	105.67	20.32	20.32	5.20		NAHQ ACTUAL
	2	24	1	105.67	105.67	19.94	19.94	5.30		LTV EC20 FISH
	2	24	1	105.67	105.67	13.55	13.55	7.80		LCV FISH
	2	24	1	105.67	105.67	9.61	9.61	11.00		EC20 POPULATION
	3	191	16	97.77	101.29	83.56	86.57	1.17		SS EC20
CYANIDE	3	191	16	97.77	101.29	18.80	19.48	5.20		NAHQ ACTUAL
	3	191	16	97.77	101.29	18.45	19.11	5.30		LTV EC20 FISH
	3	191	16	97.77	101.29	12.53	12.99	7.80		LCV FISH
	3	191	16	97.77	101.29	8.89	9.21	11.00		EC20 POPULATION
	3	32	1	91.83	95.92	78.49	81.98	1.17		SS EC20
	4	32	1	91.83	95.92	17.66	18.45	5.20		NAHQ ACTUAL
	4	32	1	91.83	95.92	17.33	18.10	5.30		LTV EC20 FISH
	4	32	1	91.83	95.92	11.77	12.30	7.80		LCV FISH
	4	32	1	91.83	95.92	8.35	8.72	11.00		EC20 POPULATION
	4	32	1	87.87	87.87	75.11	75.11	1.17		SS EC20
	13	8	1	87.87	87.87	16.90	16.90	5.20		NAHQ ACTUAL
	13	8	1	87.87	87.87	16.58	16.58	5.30		LTV EC20 FISH
	13	8	1	87.87	87.87	11.27	11.27	7.80		LCV FISH
	13	8	1	87.87	87.87	7.99	7.99	11.00		EC20 POPULATION
	3	179	11	9.93	10.00	2.26	2.27	4.40		NAHQ ADVISORY
	3	179	11	9.93	10.00	0.04	0.04	251.00		EC20 POPULATION
	3	179	11	9.93	10.00	0.04	0.04	270.00		LTV EC20 FISH
	3	179	11	9.93	10.00	0.02	0.02	500.00		LTV EC20 DAPHNID
	3	179	11	9.93	10.00	0.01	0.01	717.00		LCV FISH
	3	179	11	9.93	10.00	0.01	0.01	717.00		LCV DAPHNIDS
D1-N-BUTYLPHTHALATE	3	179	11	9.93	10.00	2.26	2.27	4.40		NAHQ ADVISORY
	3	179	11	9.93	10.00	0.04	0.04	251.00		EC20 POPULATION
	3	179	11	9.93	10.00	0.04	0.04	270.00		LTV EC20 FISH
	3	179	11	9.93	10.00	0.02	0.02	500.00		LTV EC20 DAPHNID
	3	179	11	9.93	10.00	0.01	0.01	717.00		LCV FISH

Table P3 (continued)

Analyte	Resch	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^a water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
DIETHYLPHTHALATE	3	179	1	10.05	10.14	0.37	0.38	27.00		NAWQ ADVISORY EC20 POPULATION
	3	179	1	10.05	10.14	0.01	0.01	1000.00		
FLUORANTHENE	3	179	2	10.05	10.13	5.91	5.96	1.70		NAWQ ADVISORY EC20 POPULATION
	3	179	2	10.05	10.13	0.31	0.32	32.00		
	3	179	2	10.05	10.13	0.00	0.00	65795.00	*	LCV DAPHNIDS
IRON	1	211	140	265.29	621.36	16.58	38.84	16.00		LTV EC20 DAPHNID
	1	211	140	265.29	621.36	1.68	3.93	158.00		LCV DAPHNIDS
	1	211	140	265.29	621.36	0.27	0.62	1000.00		NAWQ ACTUAL
	2	1	1	220.00	220.00	13.75	13.75	16.00		LTV EC20 DAPHNID
	2	1	1	220.00	220.00	1.39	1.39	158.00		LCV DAPHNIDS
	2	1	1	220.00	220.00	0.22	0.22	1000.00		NAWQ ACTUAL
	3	363	352	1962.63	3835.87	122.66	239.74	16.00		LTV EC20 DAPHNID
	3	363	352	1962.63	3835.87	12.42	24.28	158.00		LCV DAPHNIDS
	3	363	352	1962.63	3835.87	1.96	3.84	1000.00		NAWQ ACTUAL
	4	1	1	660.00	660.00	41.25	41.25	16.00		LTV EC20 DAPHNID
	4	1	1	660.00	660.00	4.18	4.18	158.00		LCV DAPHNIDS
	4	1	1	660.00	660.00	0.66	0.66	1000.00		NAWQ ACTUAL
	1	215	16	29.89	46.44	85.41	132.69	0.35		SS EC20
	1	215	16	29.89	46.44	9.34	14.51	3.20		NAWQ ACTUAL
	1	215	16	29.89	46.44	2.44	3.79	12.26		LCV DAPHNIDS
LEAD	1	215	16	29.89	46.44	1.58	2.46	18.88		LCV FISH
	1	215	16	29.89	46.44	1.36	2.11	22.00		LTV EC20 FISH
	1	215	16	29.89	46.44	0.42	0.65	71.00		EC20 POPULATION
	1	215	16	29.89	46.44	11.98	11.98	0.35		SS EC20
	2	24	1	4.19	4.19	1.31	1.31	3.20		NAWQ ACTUAL
	2	24	1	4.19	4.19	0.34	0.34	12.26		LCV DAPHNIDS
	2	24	1	4.19	4.19	0.22	0.22	18.88		LCV FISH
	2	24	1	4.19	4.19	0.19	0.19	22.00		LTV EC20 FISH
	2	24	1	4.19	4.19	0.06	0.06	71.00		EC20 POPULATION
	2	24	1	4.19	4.19	22.63	40.53	0.35		SS EC20
	3	715	121	7.92	14.18	2.48	4.43	3.20		NAWQ ACTUAL
	3	715	121	7.92	14.18	0.65	1.16	12.26		LCV DAPHNIDS
	3	715	121	7.92	14.18	0.42	0.75	18.88		LCV FISH
	3	715	121	7.92	14.18	0.36	0.64	22.00		LTV EC20 FISH
	3	715	121	7.92	14.18	0.11	0.20	71.00		EC20 POPULATION
	3	715	121	7.92	14.18					

Table F.3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
MANGANESE	1	114	110	144.21	144.21	13.11	13.11	11.00		NAMQ ADVISORY
	1	114	110	144.21	144.21	1.29	1.29	112.00	<	EC20 POPULATION
	1	114	110	144.21	144.21	0.13	0.13	1100.00	<	LCV DAPHNIDS
	1	114	110	144.21	144.21	0.13	0.13	1100.00		LTV EC20 DAPHNID
	1	114	110	144.21	144.21	0.11	0.11	1270.00		LTV EC20 FISH
	1	114	110	144.21	144.21	0.08	0.08	1770.00		LCV FISH
	2	24	24	35.12	35.12	3.19	3.19	11.00		NAMQ ADVISORY
	2	24	24	35.12	35.12	0.31	0.31	112.00		EC20 POPULATION
	2	24	24	35.12	35.12	9.03	9.03	1100.00	<	LCV DAPHNIDS
	2	24	24	35.12	35.12	0.03	0.03	1100.00	<	LTV EC20 DAPHNID
	2	24	24	35.12	35.12	0.03	0.03	1270.00		LTV EC20 FISH
	2	24	24	35.12	35.12	0.02	0.02	1770.00		LCV FISH
	3	410	366	145.41	201.44	13.22	18.31	11.00		NAMQ ADVISORY
	3	410	366	145.41	201.44	1.30	1.80	112.00		EC20 POPULATION
	3	410	366	145.41	201.44	0.13	0.18	1100.00	<	LCV DAPHNIDS
	3	410	366	145.41	201.44	0.13	0.18	1100.00	<	LTV EC20 DAPHNID
	3	410	366	145.41	201.44	0.11	0.16	1270.00		LTV EC20 FISH
	3	410	366	145.41	201.44	0.08	0.11	1770.00		LCV FISH
	4	32	32	52.12	61.70	4.74	5.61	11.00		NAMQ ADVISORY
	4	32	32	52.12	61.70	0.47	0.55	112.00		EC20 POPULATION
	4	32	32	52.12	61.70	0.05	0.06	1100.00	<	LCV DAPHNIDS
	4	32	32	52.12	61.70	0.05	0.06	1100.00	<	LTV EC20 DAPHNID
	4	32	32	52.12	61.70	0.04	0.05	1270.00		LTV EC20 FISH
	4	32	32	52.12	61.70	0.03	0.03	1770.00		LCV FISH
	13	8	8	154.29	154.29	14.03	14.03	11.00		NAMQ ADVISORY
	13	8	8	154.29	154.29	1.38	1.38	112.00		EC20 POPULATION
	13	8	8	154.29	154.29	0.14	0.14	1100.00	<	LCV DAPHNIDS
	13	8	8	154.29	154.29	0.14	0.14	1100.00	<	LTV EC20 DAPHNID
	13	8	8	154.29	154.29	0.12	0.12	1270.00		LTV EC20 FISH
	13	8	8	154.29	154.29	0.09	0.09	1770.00		LCV FISH

Table F-3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
MERCURY	1	147	75	0.32	0.39	1065.08	1301.59	0.00		NAHQ ADVISORY
	1	147	75	0.32	0.39	10.65	13.02	0.03	<	LTV EC20 FISH
	1	147	75	0.32	0.39	7.99	9.76	0.04	<	LCV DAPHNIDS
	1	147	75	0.32	0.39	1.14	1.39	0.28		EC20 POPULATION
	1	147	75	0.32	0.39	0.61	0.75	0.52		LCV FISH
	1	147	75	0.32	0.39	0.37	0.45	0.87		LTV EC20 DAPHNID
	3	444	50	0.40	0.65	1326.54	2175.57	0.00		NAHQ ADVISORY
	3	444	50	0.40	0.65	13.27	21.76	0.03	<	LTV EC20 FISH
	3	444	50	0.40	0.65	9.95	16.32	0.04	<	LCV DAPHNIDS
	3	444	50	0.40	0.65	1.42	2.33	0.28		EC20 POPULATION
	3	444	50	0.40	0.65	0.77	1.26	0.52		LCV FISH
	3	444	50	0.40	0.65	0.46	0.75	0.87		LTV EC20 DAPHNID
	4	32	1	0.20	0.20	669.44	672.22	0.00		NAHQ ADVISORY
	4	32	1	0.20	0.20	6.69	6.72	0.03	<	LTV EC20 FISH
	4	32	1	0.20	0.20	5.02	5.04	0.04	<	LCV DAPHNIDS
	4	32	1	0.20	0.20	0.72	0.72	0.28		EC20 POPULATION
METHYLENE CHLORIDE	4	32	1	0.20	0.20	0.39	0.39	0.52		LCV FISH
	4	32	1	0.20	0.20	0.23	0.23	0.87		LTV EC20 DAPHNID
	2	41	9	3.67	5.00	0.02	0.02	220.00		NAHQ ADVISORY
	2	41	9	3.67	5.00	0.01	0.01	410.00		LTV EC20 FISH
	2	41	9	3.67	5.00	0.00	0.00	1259.00		EC20 POPULATION
	2	41	9	3.67	5.00	0.00	0.00	42667.00	*	LCV DAPHNIDS
	2	41	9	3.67	5.00	0.00	0.00	108000.00		LCV FISH
	3	705	70	5.04	5.41	0.02	0.02	220.00		NAHQ ADVISORY
	3	705	70	5.04	5.41	0.01	0.01	410.00		LTV EC20 FISH
	3	705	70	5.04	5.41	0.00	0.00	1259.00		EC20 POPULATION
	3	705	70	5.04	5.41	0.00	0.00	42667.00	*	LCV DAPHNIDS
	3	705	70	5.04	5.41	0.00	0.00	108000.00		LCV FISH
MOLYBDENUM	3	192	31	10.95	12.83	0.37	0.43	30.00		NAHQ ADVISORY
	3	192	31	10.95	12.83	0.03	0.04	360.00		LTV EC20 DAPHNID
	3	192	31	10.95	12.83	0.01	0.01	880.00		LCV DAPHNIDS

Table F.3 (continued)

Analyte	Reech	Number analyzed in reech (water)	Number detects in reech (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
NICKEL	1	211	95	18.03	38.01	3.61	7.60	5.00	<	LCV DAPHNIDS
	1	211	95	18.03	38.01	1.64	3.46	11.00	<	SS EC20
	1	211	95	18.03	38.01	0.52	1.09	35.00	<	LCV FISH
	1	211	95	18.03	38.01	0.40	0.84	45.00		LTV EC20 DAPHNID
	1	211	95	18.03	38.01	0.29	0.61	62.00		LTV EC20 FISH
	1	211	95	18.03	38.01	0.11	0.24	160.00		NAHQ ACTUAL
	1	211	95	18.03	38.01	0.38	0.18	215.00	<	EC20 POPULATION
	2	24	2	22.08	22.08	4.42	4.42	5.00	<	LCV DAPHNIDS
	2	24	2	22.08	22.08	2.01	2.01	11.00	<	SS EC20
	2	24	2	22.08	22.08	0.63	0.63	35.00	<	LCV FISH
	2	24	2	22.08	22.08	0.49	0.49	45.00		LTV EC20 DAPHNID
	2	24	2	22.08	22.08	0.36	0.36	62.00		LTV EC20 FISH
	2	24	2	22.08	22.08	0.14	0.14	160.00		NAHQ ACTUAL
	2	24	2	22.08	22.08	0.10	0.10	215.00	<	EC20 POPULATION
	3	412	88	95.90	181.14	19.18	36.23	5.00	<	LCV DAPHNIDS
	3	412	88	95.90	181.14	8.72	16.47	11.00	<	SS EC20
	3	412	88	95.90	181.14	2.74	5.18	35.00	<	LCV FISH
	3	412	88	95.90	181.14	2.13	4.03	45.00		LTV EC20 DAPHNID
	3	412	88	95.90	181.14	1.55	2.92	62.00		LTV EC20 FISH
	3	412	88	95.90	181.14	0.60	1.13	160.00		NAHQ ACTUAL
	3	412	88	95.90	181.14	0.45	0.84	215.00	<	EC20 POPULATION
	4	32	3	14.04	16.71	2.81	3.34	5.00	<	LCV DAPHNIDS
	4	32	3	14.04	16.71	1.28	1.52	11.00	<	SS EC20
	4	32	3	14.04	16.71	0.40	0.48	35.00	<	LCV FISH
	4	32	3	14.04	16.71	0.31	0.37	45.00		LTV EC20 DAPHNID
	4	32	3	14.04	16.71	0.23	0.27	62.00		LTV EC20 FISH
	4	32	3	14.04	16.71	0.09	0.10	160.00		NAHQ ACTUAL
	4	32	3	14.04	16.71	0.07	0.08	215.00	<	EC20 POPULATION
	13	8	2	10.87	10.87	2.17	2.17	5.00	<	LCV DAPHNIDS
	13	8	2	10.87	10.87	0.99	0.99	11.00	<	SS EC20
	13	8	2	10.87	10.87	0.31	0.31	35.00	<	LCV FISH
	13	8	2	10.87	10.87	0.24	0.24	45.00		LTV EC20 DAPHNID
	13	8	2	10.87	10.87	0.18	0.18	62.00		LTV EC20 FISH
	13	8	2	10.87	10.87	0.07	0.07	160.00		NAHQ ACTUAL
	13	8	2	10.87	10.87	0.05	0.05	215.00		EC20 POPULATION
PHENANTHRENE	3	179	1	10.04	10.12	23.35	23.55	0.43		NAHQ ADVISORY
	3	179	1	10.04	10.12	0.09	0.09	110.00		LTV EC20 DAPHNID
	3	179	1	10.04	10.12	0.05	0.05	200.00		LCV DAPHNIDS

Table F.3 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
PHENOL	2	8	1	12.25	12.25	0.58	0.58	21.00	<	NAHQ ADVISORY
	2	8	1	12.25	12.25	0.05	0.05	230.00		LTV EC20 FISH
	2	8	1	12.25	12.25	0.01	0.01	1369.00		LCV FISH
	2	8	1	12.25	12.25	0.01	0.01	2005.00	*	LCV DAPHNIDS
	2	8	1	12.25	12.25	0.00	0.00	4467.00		EC20 POPULATION
	3	179	20	12.23	12.23	0.58	0.64	21.00	<	NAHQ ADVISORY
	3	179	20	12.23	13.50	0.05	0.06	230.00		LTV EC20 FISH
	3	179	20	12.23	13.50	0.01	0.01	1369.00		LCV FISH
	3	179	20	12.23	13.50	0.01	0.01	2005.00	*	LCV DAPHNIDS
	3	179	20	12.23	13.50	0.00	0.00	4467.00		EC20 POPULATION
	4	10	2	11.44	12.87	0.54	0.61	21.00	<	NAHQ ADVISORY
	4	10	2	11.44	12.87	0.05	0.06	230.00		LTV EC20 FISH
	4	10	2	11.44	12.87	0.01	0.01	1369.00		LCV FISH
	4	10	2	11.44	12.87	0.01	0.01	2005.00	*	LCV DAPHNIDS
	4	10	2	11.44	12.87	0.00	0.00	4467.00		EC20 POPULATION
SELENIUM	1	260	106	11.60	29.50	4.46	11.35	2.60		SS EC20
	1	260	106	11.60	29.50	0.46	1.18	25.00		LTV EC20 DAPHNID
	1	260	106	11.60	29.50	0.33	0.84	35.00		NAHQ ACTUAL
	1	260	106	11.60	29.50	0.29	0.74	40.00		LTV EC20 FISH
	1	260	106	11.60	29.50	0.13	0.33	88.32		LCV FISH
	1	260	106	11.60	29.50	0.13	0.32	91.65		LCV DAPHNIDS
	2	26	2	5.86	6.72	2.25	2.58	2.60		SS EC20
	2	26	2	5.86	6.72	0.23	0.27	25.00		LTV EC20 DAPHNID
	2	26	2	5.86	6.72	0.17	0.19	35.00		NAHQ ACTUAL
	2	26	2	5.86	6.72	0.15	0.17	40.00		LTV EC20 FISH
	2	26	2	5.86	6.72	0.07	0.08	88.32		LCV FISH
	2	26	2	5.86	6.72	0.06	0.07	91.65		LCV DAPHNIDS
	3	571	2	8.55	14.76	3.29	5.68	2.60		SS EC20
	3	571	2	8.55	14.76	0.34	0.59	25.00		LTV EC20 DAPHNID
	3	571	2	8.55	14.76	0.24	0.42	35.00		NAHQ ACTUAL
	3	571	2	8.55	14.76	0.21	0.37	40.00		LTV EC20 FISH
	3	571	2	8.55	14.76	0.10	0.17	88.32		LCV FISH
	3	571	2	8.55	14.76	0.09	0.16	91.65		LCV DAPHNIDS
SILVER	3	570	6	9.17	10.88	76.41	90.70	0.12		LCV FISH
	3	570	6	9.17	10.88	65.50	77.75	0.14		SS EC20
	3	570	6	9.17	10.88	45.85	54.42	0.20		LTV EC20 FISH
	3	570	6	9.17	10.88	36.68	43.54	0.25		NAHQ ADVISORY
	3	570	6	9.17	10.88	28.66	34.01	0.32		EC20 POPULATION
	3	570	6	9.17	10.88	16.37	19.44	0.56	<	LTV EC20 DAPHNID
	3	570	6	9.17	10.88	3.53	4.19	2.60		LCV DAPHNIDS

Table F.3 (continued)

Analyte	Reech	Number analyzed in reech (water)	Number detects in reech (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
SODIUM	1	106	8	2836.41	4996.87	0.03	0.06	85049.00	*	LCV DAPHNIDS
	2	25	25	9143.96	14000.00	0.11	0.16	85049.00	*	LCV DAPHNIDS
	3	49	49	8215.83	13000.00	0.10	0.15	85049.00	*	LCV DAPHNIDS
	4	32	32	4317.29	4454.58	0.05	0.05	85049.00	*	LCV DAPHNIDS
	13	8	8	4265.00	4265.00	0.05	0.05	85049.00	*	LCV DAPHNIDS
TOLUENE	2	41	2	4.86	5.00	0.47	0.48	10.40	<	NAHQ ADVISORY
	2	41	2	4.86	5.00	0.19	0.19	26.00		LTV EC20 FISH
	2	41	2	4.86	5.00	0.02	0.02	200.00		EC20 POPULATION
	2	41	2	4.86	5.00	0.00	0.00	1269.00	*	LCV FISH
	2	41	2	4.86	5.00	0.00	0.00	25229.00	*	LCV DAPHNIDS
	2	41	2	4.86	5.00	0.48	0.48	10.40		NAHQ ADVISORY
	3	705	26	4.95	5.00	0.19	0.19	26.00	<	LTV EC20 FISH
	3	705	26	4.95	5.00	0.02	0.02	200.00		EC20 POPULATION
	3	705	26	4.95	5.00	0.00	0.00	1269.00	*	LCV FISH
	3	705	26	4.95	5.00	0.00	0.00	25229.00	*	LCV DAPHNIDS
	3	705	26	4.95	5.00	0.00	0.00	25229.00	*	LCV DAPHNIDS
	3	705	26	4.95	5.00	0.00	0.00	25229.00	*	LCV DAPHNIDS
URANIUM	2	46	11	0.75	0.84	107.73	120.23	0.01		NAHQ ADVISORY
	2	46	11	0.75	0.84	0.03	0.03	27.00		EC20 POPULATION
	2	46	11	0.75	0.84	0.01	0.01	142.00	*	LCV FISH
	2	46	11	0.75	0.84	0.00	0.00	455.00	*	LTV EC20 FISH
	2	46	11	0.75	0.84	0.00	0.00	455.00	*	NAHQ ADVISORY
	3	72	65	1.94	2.53	277.78	361.11	0.01		EC20 POPULATION
	3	72	65	1.94	2.53	0.07	0.09	27.00	*	LCV FISH
	3	72	65	1.94	2.53	0.01	0.02	142.00	*	LTV EC20 FISH
	3	72	65	1.94	2.53	0.00	0.01	455.00	*	NAHQ ADVISORY
	4	58	22	0.51	0.67	72.77	95.24	0.01		EC20 POPULATION
	4	58	22	0.51	0.67	0.02	0.02	27.00	*	LCV FISH
	4	58	22	0.51	0.67	0.00	0.00	142.00	*	NAHQ ADVISORY
	4	58	22	0.51	0.67	0.00	0.00	455.00	*	EC20 POPULATION
	4	58	22	0.51	0.67	0.00	0.00	455.00	*	LTV EC20 FISH
	4	58	22	0.51	0.67	0.00	0.00	455.00	*	LCV FISH
VANADIUM	1	64	12	2.47	2.47	0.82	0.82	3.00		NAHQ ADVISORY
	1	64	12	2.47	2.47	0.08	0.08	32.00		EC20 POPULATION
	1	64	12	2.47	2.47	0.06	0.06	41.00		LTV EC20 FISH
	1	64	12	2.47	2.47	0.03	0.03	80.00		LCV FISH
	1	64	12	2.47	2.47	0.01	0.01	430.00		LTV EC20 DAPHNID
	1	64	12	2.47	2.47	0.00	0.00	940.00	>	LCV DAPHNIDS
VINYL CHLORIDE	3	704	144	9.34	10.00	9.84	10.53	0.95	*	NAHQ ADVISORY
	3	704	144	9.34	10.00	0.00	0.00	14520.00	*	LTV EC20 FISH
	3	704	144	9.34	10.00	0.00	0.00	28879.00	*	LCV FISH
	3	704	144	9.34	10.00	0.00	0.00	28879.00	*	LCV FISH
XYLENE	2	32	1	4.75	4.75	3.25	3.25	1.46		NAHQ ADVISORY
	2	32	1	4.75	4.75	0.00	0.00	2680.00		LTV EC20 FISH
	2	32	1	4.75	4.75	0.00	0.00	62308.00	*	LCV FISH
	2	32	1	4.75	4.75	0.00	0.00	62308.00	*	LCV FISH

Table F.3 (continued)

Analyte	Reech	Number analyzed in reach (water)	Number detects in reach (water)	Mean ^a water concentration	UCL ^b water concentration	Mean quotient	UCL quotient	Benchmark	Flag	Benchmark type
ZINC	1	211	119	11.47	16.80	0.55	0.80	21.00		SS EC20
	1	211	119	11.47	16.80	0.31	0.46	36.41		LCV FISH
	1	211	119	11.47	16.80	0.25	0.36	46.73		LCV DAPHNIDS
	1	211	119	11.47	16.80	0.24	0.36	47.00		LTV EC20 FISH
	1	211	119	11.47	16.80	0.14	0.21	50.00		EC20 POPULATION
	1	211	119	11.47	16.80	0.10	0.15	110.00		MAWQ ACTUAL
	1	211	119	11.47	16.80	0.51	0.51	21.00		SS EC20
	2	24	15	10.61	10.61	0.29	0.29	36.41		LCV FISH
	2	24	15	10.61	10.61	0.23	0.23	46.73		LCV DAPHNIDS
	2	24	15	10.61	10.61	0.23	0.23	47.00		LTV EC20 FISH
	2	24	15	10.61	10.61	0.13	0.13	80.00		EC20 POPULATION
	2	24	15	10.61	10.61	0.10	0.10	110.00		MAWQ ACTUAL
	3	707	235	27.62	35.23	1.32	1.68	21.00		SS EC20
	3	707	235	27.62	35.23	0.76	0.97	36.41		LCV FISH
	3	707	235	27.62	35.23	0.59	0.75	46.73		LCV DAPHNIDS
	3	707	235	27.62	35.23	0.59	0.75	47.00		LTV EC20 FISH
	3	707	235	27.62	35.23	0.35	0.44	80.00		EC20 POPULATION
	3	707	235	27.62	35.23	0.25	0.32	110.00		MAWQ ACTUAL
	4	32	20	7.32	8.12	0.35	0.39	21.00		SS EC20
	4	32	20	7.32	8.12	0.20	0.22	36.41		LCV FISH
	4	32	20	7.32	8.12	0.16	0.17	46.73		LCV DAPHNIDS
	4	32	20	7.32	8.12	0.16	0.17	47.00		LTV EC20 FISH
	4	32	20	7.32	8.12	0.09	0.10	80.00		EC20 POPULATION
	4	32	20	7.32	8.12	0.07	0.07	110.00		MAWQ ACTUAL
	13	8	5	6.01	6.01	0.29	0.29	21.00		SS EC20
	13	8	5	6.01	6.01	0.17	0.17	36.41		LCV FISH
	13	8	5	6.01	6.01	0.13	0.13	46.73		LCV DAPHNIDS
	13	8	5	6.01	6.01	0.13	0.13	47.00		LTV EC20 FISH
	13	8	5	6.01	6.01	0.08	0.08	80.00		EC20 POPULATION
	13	8	5	6.01	6.01	0.05	0.05	110.00		MAWQ ACTUAL
ZIRCONIUM	1	106	28	11.98	20.00	56.23	93.90	0.21		MAWQ ADVISORY
	1	106	28	11.98	20.00	0.05	0.08	251.00	*	EC20 POPULATION
	1	106	28	11.98	20.00	0.02	0.04	548.00	*	LCV FISH
	1	106	28	11.98	20.00	0.00	0.01	2396.00	*	LTV EC20 FISH

^aIn cases of mixed detects and nondetects, statistics are calculated using all values, "as is." The mean concentration is the geometric mean.

^bUCL concentration was compared with the maximum value, and the smaller has been used as the "UCL concentration."

Table F-4. Limits of detection of chemicals not detected in NPDES water samples (ug/L), toxicological benchmarks (ug/L), and quotients for aquatic life

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
1,1,1-TRICHLOROETHANE	2	41	0	4.844	5.053	3.46	3.61	1.40	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.02	0.02	251.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.00	1300.00	LTV EC20 DAPHNID
	2	41	0	4.844	5.053	0.00	0.00	1770.00	LCV DAPHNIDS
	2	41	0	4.844	5.053	0.00	0.00	2457.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.00	0.00	3493.00	LCV FISH
	4	10	0	5.000	5.000	3.57	3.57	1.40	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.02	0.02	251.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	1300.00	LTV EC20 DAPHNID
	4	10	0	5.000	5.000	0.00	0.00	1770.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.00	0.00	2457.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	3493.00	LCV FISH
	13	2	0	5.000	5.000	3.57	3.57	1.40	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.02	0.02	251.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	1300.00	LTV EC20 DAPHNID
	13	2	0	5.000	5.000	0.00	0.00	1770.00	LCV DAPHNIDS
1,1,2,2-TETRACHLOROETHANE	13	2	0	5.000	5.000	0.00	0.00	2457.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	3493.00	LCV FISH
	2	41	0	4.844	5.053	0.16	0.16	31.00	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.01	0.01	420.00	LTV EC20 DAPHNID
	2	41	0	4.844	5.053	0.00	0.00	1400.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.00	0.00	1585.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.00	2400.00	LCV FISH
	2	41	0	4.844	5.053	0.00	0.00	9900.00	LCV DAPHNIDS
	3	705	0	5.000	5.000	0.16	0.16	31.00	NAHQ ADVISORY
	3	705	0	5.000	5.000	0.01	0.01	420.00	LTV EC20 DAPHNID
	3	705	0	5.000	5.000	0.00	0.00	1400.00	LTV EC20 FISH
	3	705	0	5.000	5.000	0.00	0.00	1585.00	EC20 POPULATION
	3	705	0	5.000	5.000	0.00	0.00	2400.00	LCV FISH
	3	705	0	5.000	5.000	0.00	0.00	9900.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.16	0.16	31.00	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.01	0.01	420.00	LTV EC20 DAPHNID
	4	10	0	5.000	5.000	0.00	0.00	1400.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	1585.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	2400.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	9900.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.16	0.16	31.00	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.01	0.01	420.00	LTV EC20 DAPHNID
	13	2	0	5.000	5.000	0.00	0.00	1400.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	1585.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	2400.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	9900.00	LCV DAPHNIDS

Table FA (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
1,1,2-TRICHLOROETHANE	2	41	0	4.844	5.053	0.05	0.05	92.00	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.00	0.00	9400.00	LCV FISH
	2	41	0	4.844	5.053	0.00	0.00	13000.00	LTV EC20 DAPHNID
	2	41	0	4.844	5.053	0.00	0.00	14800.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.00	0.00	15849.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.00	18400.00	LCV DAPHNIDS
	2	41	0	4.844	5.053	0.00	0.00	92.00	NAHQ ADVISORY
	3	705	0	5.000	5.000	0.05	0.05	9400.00	LCV FISH
	3	705	0	5.000	5.000	0.00	0.00	13000.00	LTV EC20 DAPHNID
	3	705	0	5.000	5.000	0.00	0.00	14800.00	LTV EC20 FISH
	3	705	0	5.000	5.000	0.00	0.00	15849.00	EC20 POPULATION
	3	705	0	5.000	5.000	0.00	0.00	18400.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.05	0.05	92.00	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.00	0.00	9400.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	13000.00	LTV EC20 DAPHNID
	4	10	0	5.000	5.000	0.00	0.00	14800.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	15849.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	18400.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.05	0.05	92.00	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.00	0.00	9400.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	13000.00	LTV EC20 DAPHNID
	13	2	0	5.000	5.000	0.00	0.00	14800.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	15849.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	18400.00	LCV DAPHNIDS
1,1-DICHLOROETHANE	2	41	0	4.844	5.053	9.50	9.91	0.51	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.00	0.00	1585.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.00	8219.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.00	0.00	14680.00	LCV FISH
	4	41	0	5.000	5.000	9.80	9.80	0.51	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.00	0.00	1585.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	8219.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	14680.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	0.51	NAHQ ADVISORY
	13	2	0	5.000	5.000	9.80	9.80	0.51	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	1585.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	8219.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	14680.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	0.51	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.00	0.00	1585.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	8219.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	14680.00	LCV FISH

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	HDL quotient	UDL quotient	Benchmark	Benchmark type
1,2-DICHLOROETHANE	2	41	0	4.844	5.053	0.03	0.04	144.00	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.00	0.00	1259.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.00	11000.00	LTV EC20 DAPHNID
	2	41	0	4.844	5.053	0.00	0.00	15200.00	LCV DAPHNIDS
	2	41	0	4.844	5.053	0.00	0.00	29000.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.00	0.00	41364.00	LCV FISH
	4	10	0	5.000	5.000	0.03	0.03	144.00	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.00	0.00	1259.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	11000.00	LTV EC20 DAPHNID
	4	10	0	5.000	5.000	0.00	0.00	15200.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.00	0.00	29000.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	41364.00	LCV FISH
2-HEXANONE	13	2	0	5.000	5.000	0.03	0.03	144.00	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.00	0.00	1259.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	11000.00	LTV EC20 DAPHNID
	13	2	0	5.000	5.000	0.00	0.00	15200.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.00	0.00	29000.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	41364.00	LCV FISH
	2	32	0	9.687	9.687	8.81	8.81	1.10	NAHQ ADVISORY
	2	32	0	9.687	9.687	0.01	0.01	1259.00	EC20 POPULATION
	2	32	0	9.687	9.687	0.00	0.00	16155.00	LTV EC20 FISH
	2	32	0	9.687	9.687	0.00	0.00	32783.00	LCV FISH
	3	2	0	10.000	10.000	9.09	9.09	1.10	NAHQ ADVISORY
	3	2	0	10.000	10.000	0.01	0.01	1259.00	EC20 POPULATION
2-METHYLPHENOL	3	2	0	10.000	10.000	0.00	0.00	16155.00	LTV EC20 FISH
	3	2	0	10.000	10.000	0.00	0.00	32783.00	LCV FISH
	3	1	0	10.000	10.000	1.30	1.30	7.70	NAHQ ADVISORY
	3	1	0	10.000	10.000	0.14	0.14	74.00	EC20 POPULATION
	3	1	0	10.000	10.000	0.02	0.02	470.00	LTV EC20 FISH
	3	1	0	10.000	10.000	0.02	0.02	489.00	LCV FISH
2-NITROPHENOL	3	1	0	10.000	10.000	0.01	0.01	1316.00	LCV DAPHNIDS
	2	8	0	10.000	10.000	0.07	0.07	150.00	NAHQ ACTUAL
	3	179	0	10.000	10.148	0.07	0.07	150.00	NAHQ ACTUAL
	4	10	0	10.000	10.000	0.07	0.07	150.00	NAHQ ACTUAL
	13	2	0	10.000	10.000	0.07	0.07	150.00	NAHQ ACTUAL
	3	1	0	0.100	0.100	238.10	238.10	0.00	NAHQ ADVISORY
4,4'-DDO	3	1	0	0.100	0.100	0.16	0.16	0.61	EC20 POPULATION
	3	1	0	0.100	0.100	0.06	0.06	1.69	LCV FISH
	3	1	0	0.100	0.100	0.03	0.03	3.99	LTV EC20 FISH
	3	1	0	0.100	0.100	0.03	0.03	3.99	LTV EC20 FISH

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
4,4'-DDE	3	1	0	0.100	0.100	12.50	12.50	0.01	SS EC20
	3	1	0	0.100	0.100	6.25	6.25	0.02	LCV DAPHNIDS
	3	1	0	0.100	0.100	5.00	5.00	0.02	NAHQ ADVISORY
	3	1	0	0.100	0.100	0.29	0.29	0.35	LTV EC20 FISH
	3	1	0	0.100	0.100	0.14	0.14	0.73	LCV FISH
4,4'-DDT	3	1	0	0.100	0.100	12.50	12.50	0.01	SS EC20
	3	1	0	0.100	0.100	6.25	6.25	0.02	LCV DAPHNIDS
	3	1	0	0.100	0.100	5.00	5.00	0.02	NAHQ ADVISORY
	3	1	0	0.100	0.100	0.29	0.29	0.35	LTV EC20 FISH
	3	1	0	0.100	0.100	0.14	0.14	0.73	LCV FISH
4-METHYL-2-PENTANONE	2	32	0	9.687	9.687	4.21	4.21	2.30	NAHQ ADVISORY
	2	32	0	9.687	9.687	0.01	0.01	1585.00	EC20 POPULATION
	2	32	0	9.687	9.687	0.00	0.00	77400.00	LCV FISH
	3	2	0	10.000	10.000	4.35	4.35	2.30	NAHQ ADVISORY
	3	2	0	10.000	10.000	0.01	0.01	1585.00	EC20 POPULATION
ACENAPHTHENE	3	2	0	10.000	10.000	0.00	0.00	77400.00	LCV FISH
	2	8	0	10.000	10.000	4.55	4.55	2.20	NAHQ ADVISORY
	2	8	0	10.000	10.000	0.05	0.05	197.00	LTV EC20 FISH
	2	8	0	10.000	10.000	0.02	0.02	413.00	LCV FISH
	3	8	0	10.000	10.000	0.00	0.00	6646.00	LCV DAPHNIDS
ACETONE	3	179	0	10.000	10.148	4.55	4.61	2.20	NAHQ ADVISORY
	3	179	0	10.000	10.148	0.05	0.05	197.00	LTV EC20 FISH
	3	179	0	10.000	10.148	0.02	0.02	413.00	LCV FISH
	3	179	0	10.000	10.148	0.00	0.00	6646.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	4.55	4.55	2.20	NAHQ ADVISORY
ALDRIN	4	10	0	10.000	10.000	0.05	0.05	197.00	LTV EC20 FISH
	4	10	0	10.000	10.000	0.02	0.02	413.00	LCV FISH
	4	10	0	10.000	10.000	0.00	0.00	6646.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	4.55	4.55	2.20	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.05	0.05	197.00	LTV EC20 FISH
ALDRIN	13	2	0	10.000	10.000	0.02	0.02	413.00	LCV FISH
	13	2	0	10.000	10.000	0.00	0.00	6646.00	LCV DAPHNIDS
	3	4	0	10.000	10.000	0.01	0.01	770.00	NAHQ ADVISORY
	3	4	0	10.000	10.000	0.00	0.00	23714.00	EC20 POPULATION
	3	4	0	10.000	10.000	0.00	0.00	161867.00	LTV EC20 FISH
ALDRIN	3	4	0	10.000	10.000	0.00	0.00	507640.00	LCV FISH
	3	4	0	10.000	10.000	0.00	0.00	3114182.00	LCV DAPHNIDS
	3	1	0	0.050	0.050	0.67	0.67	0.08	NAHQ ADVISORY

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UOL quotient	Benchmark	Benchmark type
ANTHRACENE	2	8	0	10.000	10.000	1000000.00	1000000.00	0.00	NAHQ ADVISORY
	2	8	0	10.000	10.000	111.11	111.11	0.09	LCV FISH
	2	8	0	10.000	10.000	28.57	28.57	0.35	LTV EC20 FISH
	2	8	0	10.000	10.000	4.76	4.76	2.10	LCV DAPHNIDS
	2	8	0	10.000	10.000	1.22	1.22	8.20	LTV EC20 DAPHNID
	4	10	0	10.000	10.000	1000000.00	1000000.00	0.00	NAHQ ADVISORY
	4	10	0	10.000	10.000	111.11	111.11	0.09	LCV FISH
	4	10	0	10.000	10.000	28.57	28.57	0.35	LTV EC20 FISH
	4	10	0	10.000	10.000	4.76	4.76	2.10	LCV DAPHNIDS
	4	10	0	10.000	10.000	1.22	1.22	8.20	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	1000000.00	1000000.00	0.00	NAHQ ADVISORY
	13	2	0	10.000	10.000	111.11	111.11	0.09	LCV FISH
	13	2	0	10.000	10.000	28.57	28.57	0.35	LTV EC20 FISH
	13	2	0	10.000	10.000	4.76	4.76	2.10	LCV DAPHNIDS
	13	2	0	10.000	10.000	1.22	1.22	8.20	LTV EC20 DAPHNID
ANTIMONY	2	2	0	50.000	50.000	2.78	2.78	18.00	NAHQ ADVISORY
	2	2	0	50.000	50.000	0.63	0.63	79.00	EC20 POPULATION
	2	2	0	50.000	50.000	0.03	0.03	1600.00	LCV FISH
	2	2	0	50.000	50.000	0.03	0.03	1900.00	LTV EC20 DAPHNID
	2	2	0	50.000	50.000	0.02	0.02	2310.00	LTV EC20 FISH
	2	2	0	50.000	50.000	0.01	0.01	5400.00	LCV DAPHNIDS
	3	194	0	50.000	50.000	2.78	3.17	18.00	NAHQ ADVISORY
	3	194	0	50.000	50.000	0.63	0.72	79.00	EC20 POPULATION
	3	194	0	50.000	50.000	0.03	0.04	1600.00	LCV FISH
	3	194	0	50.000	50.000	0.03	0.03	1900.00	LTV EC20 DAPHNID
	3	194	0	50.000	50.000	0.02	0.02	2310.00	LTV EC20 FISH
	3	194	0	50.000	50.000	0.01	0.01	5400.00	LCV DAPHNIDS
	4	1	0	50.000	50.000	2.78	2.78	18.00	NAHQ ADVISORY
	4	1	0	50.000	50.000	0.63	0.63	79.00	EC20 POPULATION
	4	1	0	50.000	50.000	0.03	0.03	1600.00	LCV FISH
	4	1	0	50.000	50.000	0.03	0.03	1900.00	LTV EC20 DAPHNID
AROCLOL-1221	4	1	0	50.000	50.000	0.02	0.02	2310.00	LTV EC20 FISH
	4	1	0	50.000	50.000	0.01	0.01	5400.00	LCV DAPHNIDS
	3	210	0	0.500	0.501	172.41	172.91	0.00	NAHQ ADVISORY
	3	210	0	0.500	0.501	0.05	0.05	10.00	EC20 POPULATION
	3	210	0	0.500	0.501	0.01	0.01	60.00	LCV FISH
	3	210	0	0.500	0.501	0.01	0.01	80.00	LTV EC20 FISH
	4	1	0	0.500	0.500	172.41	172.41	0.00	NAHQ ADVISORY
	4	1	0	0.500	0.500	0.05	0.05	10.00	EC20 POPULATION
	4	1	0	0.500	0.500	0.01	0.01	60.00	LCV FISH
	4	1	0	0.500	0.500	0.01	0.01	80.00	LTV EC20 FISH

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
AROCLOR-1232	3	210	0	0.500	0.501	86.21	86.46	0.01	NAHQ ADVISORY
	3	210	0	0.500	0.501	0.03	0.03	16.00	EC20 POPULATION
	3	210	0	0.500	0.501	0.00	0.00	124.00	LCV FISH
	3	210	0	0.500	0.501	0.00	0.00	148.00	LTV EC20 FISH
	4	1	0	0.500	0.500	86.21	86.21	0.01	NAHQ ADVISORY
	4	1	0	0.500	0.500	0.03	0.03	16.00	EC20 POPULATION
	4	1	0	0.500	0.500	0.00	0.00	124.00	LCV FISH
	4	1	0	0.500	0.500	0.00	0.00	148.00	LTV EC20 FISH
	3	210	0	0.500	0.501	50.00	50.14	0.01	NAHQ ADVISORY
	3	210	0	0.500	0.501	0.32	0.32	1.58	EC20 POPULATION
AROCLOR-1242	3	210	0	0.500	0.501	0.17	0.17	2.90	LTV EC20 FISH
	3	210	0	0.500	0.501	0.06	0.06	9.00	LCV FISH
	3	210	0	0.500	0.501	0.06	0.06	0.01	NAHQ ADVISORY
	4	1	0	0.500	0.500	50.00	50.00	0.01	EC20 POPULATION
	4	1	0	0.500	0.500	0.32	0.32	1.58	LTV EC20 FISH
	4	1	0	0.500	0.500	0.17	0.17	2.90	LCV FISH
	4	1	0	0.500	0.500	0.06	0.06	9.00	NAHQ ADVISORY
	3	210	0	0.500	0.501	2500.00	2507.23	0.00	NAHQ ADVISORY
	3	210	0	0.500	0.501	2.50	2.51	0.20	LCV FISH
	3	210	0	0.500	0.501	1.25	1.25	0.40	LTV EC20 FISH
AROCLOR-1248	3	210	0	0.500	0.501	0.40	0.40	1.26	EC20 POPULATION
	3	210	0	0.500	0.501	0.20	0.20	2.50	LTV EC20 DAPHNID
	3	210	0	0.500	0.501	0.12	0.12	4.30	LCV DAPHNIDS
	3	210	0	0.500	0.500	2500.00	2500.00	0.00	NAHQ ADVISORY
	4	1	0	0.500	0.500	2.50	2.50	0.20	LCV FISH
	4	1	0	0.500	0.500	1.25	1.25	0.40	LTV EC20 FISH
	4	1	0	0.500	0.500	0.40	0.40	1.26	EC20 POPULATION
	4	1	0	0.500	0.500	0.20	0.20	2.50	LTV EC20 DAPHNID
	4	1	0	0.500	0.500	0.12	0.12	4.30	LCV DAPHNIDS
	4	1	0	0.500	0.500	50.00	50.00	0.02	NAHQ ADVISORY
AROCLOR-1254	4	1	0	1.000	1.000	1.92	1.92	0.52	LTV EC20 FISH
	4	1	0	1.000	1.000	1.59	1.59	0.63	EC20 POPULATION
	4	1	0	1.000	1.000	1.00	1.00	1.00	LCV FISH
	4	1	0	1.000	1.000	0.83	0.83	1.20	LTV EC20 DAPHNID
	4	1	0	1.000	1.000	0.48	0.48	2.10	LCV DAPHNIDS
	3	210	0	0.500	0.501	2500.00	2507.23	0.00	NAHQ ADVISORY
	3	210	0	0.500	0.501	2.50	2.51	0.20	LCV FISH
	3	210	0	0.500	0.501	1.25	1.25	0.40	LTV EC20 FISH
	3	210	0	0.500	0.501	0.40	0.40	1.26	EC20 POPULATION
	3	210	0	0.500	0.501	0.20	0.20	2.50	LTV EC20 DAPHNID

Table F.4 (continued)

Analyte	Resch	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UCL quotient	Benchmark	Benchmark type
AROCLOP-1260	3	210	0	1.000	1.003	9.09	9.12	0.11	NAHQ ADVISORY
	3	210	0	1.000	1.003	0.48	0.48	2.10	LTV EC20 FISH
	3	210	0	1.000	1.003	0.43	0.44	2.30	LCV FISH
	3	210	0	1.000	1.003	0.00	0.00	316.00	EC20 POPULATION
	4	1	0	1.000	1.000	9.09	9.09	0.11	NAHQ ADVISORY
	4	1	0	1.000	1.000	0.48	0.48	2.10	LTV EC20 FISH
	4	1	0	1.000	1.000	0.43	0.43	2.30	LCV FISH
	4	1	0	1.000	1.000	0.00	0.00	316.00	EC20 POPULATION
	2	24	0	5.000	5.000	0.09	0.09	55.00	SS EC20
	2	24	0	5.000	5.000	0.03	0.03	190.00	NAHQ ACTUAL
ARSENIC	2	24	0	5.000	5.000	0.01	0.01	633.00	LTV EC20 DAPHNID
	2	24	0	5.000	5.000	0.01	0.01	914.10	LCV DAPHNIDS
	2	24	0	5.000	5.000	0.00	0.00	1995.00	EC20 POPULATION
	2	24	0	5.000	5.000	0.00	0.00	2130.00	LTV EC20 FISH
	2	24	0	5.000	5.000	0.00	0.00	2962.00	LCV FISH
	4	32	0	5.000	5.000	0.09	0.09	55.00	SS EC20
	4	32	0	5.000	5.000	0.03	0.03	190.00	NAHQ ACTUAL
	4	32	0	5.000	5.000	0.01	0.01	633.00	LTV EC20 DAPHNID
	4	32	0	5.000	5.000	0.01	0.01	914.10	LCV DAPHNIDS
	4	32	0	5.000	5.000	0.00	0.00	1995.00	EC20 POPULATION
	4	32	0	5.000	5.000	0.00	0.00	2130.00	LTV EC20 FISH
	4	32	0	5.000	5.000	0.00	0.00	2962.00	LCV FISH
	13	8	0	5.000	5.000	0.09	0.09	55.00	SS EC20
	13	8	0	5.000	5.000	0.03	0.03	190.00	NAHQ ACTUAL
	13	8	0	5.000	5.000	0.01	0.01	633.00	LTV EC20 DAPHNID
	13	8	0	5.000	5.000	0.01	0.01	914.10	LCV DAPHNIDS
	13	8	0	5.000	5.000	0.00	0.00	1995.00	EC20 POPULATION
	13	8	0	5.000	5.000	0.00	0.00	2130.00	LTV EC20 FISH
	13	8	0	5.000	5.000	0.00	0.00	2962.00	LCV FISH
BENZENE	2	41	0	4.844	5.053	1.51	1.58	3.20	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.23	0.24	21.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.02	0.02	229.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.02	0.02	299.00	LCV FISH
	2	41	0	4.844	5.053	0.00	0.00	98000.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	1.56	1.56	3.20	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.24	0.24	21.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.02	0.02	229.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.02	0.02	299.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	98000.00	LCV DAPHNIDS

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
BENZENE (continued)	13	2	0	5.000	5.000	1.56	1.56	3.20	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.24	0.24	21.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.02	0.02	229.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.02	0.02	299.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	98000.00	LCV DAPHNIDS
BENZO(A)ANTHRACENE	2	8	0	10.000	10.000	3703.70	3703.70	0.00	NAHQ ADVISORY
	2	8	0	10.000	10.000	15.38	15.38	0.65	LCV DAPHNIDS
	4	10	0	10.000	10.000	3703.70	3703.70	0.00	NAHQ ADVISORY
	4	10	0	10.000	10.000	15.38	15.38	0.65	LCV DAPHNIDS
	13	2	0	10.000	10.000	3703.70	3703.70	0.00	NAHQ ADVISORY
BENZO(A)PYRENE	13	2	0	10.000	10.000	15.38	15.38	0.65	LCV DAPHNIDS
	2	8	0	10.000	10.000	7692.31	7692.31	0.00	NAHQ ADVISORY
	2	8	0	10.000	10.000	33.33	33.33	0.30	LCV DAPHNIDS
	2	8	0	10.000	10.000	3.34	3.34	2.99	LTV EC20 FISH
	3	179	0	10.000	10.148	7692.31	7805.96	0.00	NAHQ ADVISORY
	3	179	0	10.000	10.148	33.33	33.83	0.30	LCV DAPHNIDS
	3	179	0	10.000	10.148	3.34	3.39	2.99	LTV EC20 FISH
	4	10	0	10.000	10.000	7692.31	7692.31	0.00	NAHQ ADVISORY
	4	10	0	10.000	10.000	33.33	33.33	0.30	LCV DAPHNIDS
	4	10	0	10.000	10.000	3.34	3.34	2.99	LTV EC20 FISH
	13	2	0	10.000	10.000	7692.31	7692.31	0.00	NAHQ ADVISORY
	13	2	0	10.000	10.000	33.33	33.33	0.30	LCV DAPHNIDS
BENZOIC ACID	13	2	0	10.000	10.000	3.34	3.34	2.99	LTV EC20 FISH
	3	1	0	50.000	50.000	111.11	111.11	0.45	NAHQ ADVISORY
	3	1	0	50.000	50.000	0.04	0.04	1259.00	EC20 POPULATION
	3	1	0	50.000	50.000	0.01	0.01	7409.00	LTV EC20 FISH
	3	1	0	50.000	50.000	0.00	0.00	12976.00	LCV FISH
BENZYL ALCOHOL	3	1	0	10.000	10.000	68.03	68.03	0.15	NAHQ ADVISORY
	3	1	0	10.000	10.000	0.03	0.03	375.00	EC20 POPULATION
	3	1	0	10.000	10.000	0.02	0.02	550.00	LTV EC20 FISH
	3	1	0	10.000	10.000	0.02	0.02	589.00	LCV FISH

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
BERYLLIUM	2	2	0	0.200	0.885	0.33	1.45	0.61	NAHQ ADVISORY
	2	2	0	0.200	0.885	0.05	0.23	3.80	LTV EC20 DAPHNID
	2	2	0	0.200	0.885	0.04	0.17	5.30	LCV DAPHNIDS
	2	2	0	0.200	0.885	0.01	0.04	21.00	EC20 POPULATION
	2	2	0	0.200	0.885	0.00	0.02	57.00	LCV FISH
	2	2	0	0.200	0.885	0.00	0.01	148.00	LTV EC20 FISH
	4	1	0	0.300	0.300	0.49	0.49	0.61	NAHQ ADVISORY
	4	1	0	0.300	0.300	0.08	0.08	3.80	LTV EC20 DAPHNID
	4	1	0	0.300	0.300	0.06	0.06	5.30	LCV DAPHNIDS
	4	1	0	0.300	0.300	0.01	0.01	21.00	EC20 POPULATION
	4	1	0	0.300	0.300	0.01	0.01	57.00	LCV FISH
	4	1	0	0.300	0.300	0.00	0.00	148.00	LTV EC20 FISH
BIS(2-ETHYLNEXYL)PHTHALATE	2	8	0	10.000	10.000	1000.00	1000.00	0.01	NAHQ ADVISORY
	2	8	0	10.000	10.000	3.33	3.33	3.00	LCV DAPHNIDS
	2	8	0	10.000	10.000	3.33	3.33	3.00	LTV EC20 DAPHNID
	2	8	0	10.000	10.000	1.19	1.19	8.40	LCV FISH
	2	8	0	10.000	10.000	0.20	0.20	50.00	EC20 POPULATION
	2	8	0	10.000	10.000	0.19	0.19	54.00	LTV EC20 FISH
	4	10	0	10.000	10.000	1000.00	1000.00	0.01	NAHQ ADVISORY
	4	10	0	10.000	10.000	3.33	3.33	3.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	3.33	3.33	3.00	LTV EC20 DAPHNID
	4	10	0	10.000	10.000	1.19	1.19	8.40	LCV FISH
	4	10	0	10.000	10.000	0.20	0.20	50.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.19	0.19	54.00	LTV EC20 FISH
	13	2	0	10.000	10.000	1000.00	1000.00	0.01	NAHQ ADVISORY
	13	2	0	10.000	10.000	3.33	3.33	3.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	3.33	3.33	3.00	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	1.19	1.19	8.40	LCV FISH
	13	2	0	10.000	10.000	0.20	0.20	50.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.19	0.19	54.00	LTV EC20 FISH
BORON	1	64	0	80.000	80.000	11.43	11.43	7.00	LTV EC20 DAPHNID
	2	2	0	4.000	524.836	0.57	74.98	7.00	LTV EC20 DAPHNID

Table F.4 (continued)

Analyte	Resch	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UOL quotient	Benchmark	Benchmark type
CADMIUM	2	24	0	2.083	2.083	160.26	160.26	0.01	SS EC20
	2	24	0	2.083	2.083	13.89	13.89	0.15	LCV DAPHNIDS
	2	24	0	2.083	2.083	2.78	2.78	0.75	LTV EC20 DAPHNID
	2	24	0	2.083	2.083	1.89	1.89	1.10	NAHQ ACTUAL
	2	24	0	2.083	2.083	1.23	1.23	1.70	LCV FISH
	2	24	0	2.083	2.083	1.16	1.16	1.80	LTV EC20 FISH
	2	24	0	2.083	2.083	0.48	0.48	4.30	EC20 POPULATION
	2	24	0	2.083	2.083	153.85	153.85	0.01	SS EC20
	4	32	0	2.000	2.000	13.33	13.33	0.15	LCV DAPHNIDS
	4	32	0	2.000	2.000	2.67	2.67	0.75	LTV EC20 DAPHNID
	4	32	0	2.000	2.000	1.82	1.82	1.10	NAHQ ACTUAL
	4	32	0	2.000	2.000	1.18	1.18	1.70	LCV FISH
	4	32	0	2.000	2.000	1.11	1.11	1.80	LTV EC20 FISH
	4	32	0	2.000	2.000	0.47	0.47	4.30	EC20 POPULATION
	4	32	0	2.000	2.000	153.85	153.85	0.01	SS EC20
	13	8	0	2.000	2.000	13.33	13.33	0.15	LCV DAPHNIDS
	13	8	0	2.000	2.000	2.67	2.67	0.75	LTV EC20 DAPHNID
	13	8	0	2.000	2.000	1.82	1.82	1.10	NAHQ ACTUAL
	13	8	0	2.000	2.000	1.18	1.18	1.70	LCV FISH
	13	8	0	2.000	2.000	1.11	1.11	1.80	LTV EC20 FISH
	13	8	0	2.000	2.000	0.47	0.47	4.30	EC20 POPULATION
CARBON TETRACHLORIDE	2	41	0	4.844	5.053	0.08	0.08	63.00	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.07	0.08	65.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.02	0.02	224.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.00	3400.00	LCV FISH
	2	41	0	4.844	5.053	0.00	0.00	5580.00	LCV DAPHNIDS
	3	705	0	5.000	5.000	0.08	0.08	63.00	NAHQ ADVISORY
	3	705	0	5.000	5.000	0.08	0.08	65.00	LTV EC20 FISH
	3	705	0	5.000	5.000	0.02	0.02	224.00	EC20 POPULATION
	3	705	0	5.000	5.000	0.00	0.00	3400.00	LCV FISH
	3	705	0	5.000	5.000	0.00	0.00	5580.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.08	0.08	63.00	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.08	0.08	65.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.02	0.02	224.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	3400.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	5580.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.08	0.08	63.00	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.08	0.08	65.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.02	0.02	224.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	3400.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	5580.00	LCV DAPHNIDS

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
CHLORDANE	3	1	0	0.500	0.500	2.94	2.94	0.17	NAHQ ACTUAL
	3	1	0	0.500	0.500	2.00	2.00	0.25	LTV EC20 FISH
	3	1	0	0.500	0.500	1.00	1.00	0.50	SS EC20
	3	1	0	0.500	0.500	0.70	0.70	0.71	EC20 POPULATION
	3	1	0	0.500	0.500	0.31	0.31	1.60	LCV FISH
	3	1	0	0.500	0.500	0.04	0.04	12.10	LTV EC20 DAPHNID
CHLOROBENZENE	3	1	0	0.500	0.500	0.03	0.03	16.00	LCV DAPHNIDS
	2	41	0	4.844	5.053	0.42	0.44	11.60	NAHQ ADVISORY
	2	41	0	4.844	5.053	0.03	0.03	165.00	EC20 POPULATION
	2	41	0	4.844	5.053	0.00	0.01	1002.00	LTV EC20 FISH
	2	41	0	4.844	5.053	0.00	0.00	1203.00	LCV FISH
	2	41	0	4.844	5.053	0.00	0.00	15042.00	LCV DAPHNIDS
CHLOROFORM	4	10	0	5.000	5.000	0.43	0.43	11.60	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.03	0.03	165.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	1002.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	1203.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	15042.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.43	0.43	11.60	NAHQ ADVISORY
CHROMIUM	13	2	0	5.000	5.000	0.03	0.03	165.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	1002.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	1203.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	15042.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.36	0.36	14.00	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.01	0.01	562.00	EC20 POPULATION
CHLOROFORM	4	10	0	5.000	5.000	0.00	0.00	3578.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	4483.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.00	0.00	8400.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.36	0.36	14.00	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.01	0.01	562.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	3578.00	LCV FISH
CHROMIUM	13	2	0	5.000	5.000	0.00	0.00	4483.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.00	0.00	8400.00	LTV EC20 FISH
	2	24	0	10.000	10.000	37.59	37.59	0.27	SS EC20
	2	24	0	10.000	10.000	20.00	20.00	0.50	LTV EC20 DAPHNID
	2	24	0	10.000	10.000	1.63	1.63	6.13	LCV DAPHNIDS
	2	24	0	10.000	10.000	0.91	0.91	11.00	NAHQ ACTUAL
CHLOROFORM	2	24	0	10.000	10.000	0.20	0.20	51.00	LTV EC20 FISH
	2	24	0	10.000	10.000	0.14	0.14	73.18	LCV FISH
	2	24	0	10.000	10.000	0.03	0.03	316.00	EC20 POPULATION
	2	24	0	10.000	10.000	0.03	0.03	316.00	EC20 POPULATION
	2	24	0	10.000	10.000	0.03	0.03	316.00	EC20 POPULATION
	2	24	0	10.000	10.000	0.03	0.03	316.00	EC20 POPULATION

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
COBALT	1	64	0	4.000	4.000	13.79	13.79	0.29	NAHQ ADVISORY
	1	64	0	4.000	4.000	1.01	1.01	3.98	EC20 POPULATION
	1	64	0	4.000	4.000	0.91	0.91	4.40	LTV EC20 DAPHNID
	1	64	0	4.000	4.000	0.78	0.78	5.10	LCV DAPHNIDS
	1	64	0	4.000	4.000	0.01	0.01	290.00	LCV FISH
	1	64	0	4.000	4.000	0.00	0.00	810.00	LTV EC20 FISH
	2	2	0	4.000	10.853	13.79	37.42	0.29	NAHQ ADVISORY
	2	2	0	4.000	10.853	1.01	2.73	3.98	EC20 POPULATION
	2	2	0	4.000	10.853	0.91	2.47	4.40	LTV EC20 DAPHNID
	2	2	0	4.000	10.853	0.78	2.13	5.10	LCV DAPHNIDS
	2	2	0	4.000	10.853	0.01	0.04	290.00	LCV FISH
	2	2	0	4.000	10.853	0.00	0.01	810.00	LTV EC20 FISH
	3	192	0	5.000	115.367	17.24	397.82	0.29	NAHQ ADVISORY
	3	192	0	5.000	115.367	1.26	28.99	3.98	EC20 POPULATION
	3	192	0	5.000	115.367	1.14	26.22	4.40	LTV EC20 DAPHNID
	3	192	0	5.000	115.367	0.92	22.62	5.10	LCV DAPHNIDS
	3	192	0	5.000	115.367	0.02	0.40	290.00	LCV FISH
	3	192	0	5.000	115.367	0.01	0.14	810.00	LTV EC20 FISH
	4	1	0	5.000	5.000	17.24	17.24	0.29	NAHQ ADVISORY
	4	1	0	5.000	5.000	1.26	1.26	3.98	EC20 POPULATION
	4	1	0	5.000	5.000	1.14	1.14	4.40	LTV EC20 DAPHNID
	4	1	0	5.000	5.000	0.98	0.98	5.10	LCV DAPHNIDS
	4	1	0	5.000	5.000	0.02	0.02	290.00	LCV FISH
	4	1	0	5.000	5.000	0.01	0.01	810.00	LTV EC20 FISH
COPPER	13	8	0	4.000	4.000	19.51	19.51	0.20	LTV EC20 DAPHNID
	13	8	0	4.000	4.000	17.39	17.39	0.23	LCV DAPHNIDS
	13	8	0	4.000	4.000	15.38	15.38	0.26	SS EC20
	13	8	0	4.000	4.000	1.03	1.03	3.87	LCV FISH
	13	8	0	4.000	4.000	0.80	0.80	5.00	LTV EC20 FISH
	13	8	0	4.000	4.000	0.47	0.47	8.60	EC20 POPULATION
	13	8	0	4.000	4.000	0.33	0.33	12.00	NAHQ ACTUAL
	13	8	0	4.000	4.000				

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
DI-N-BUTYLPHTHALATE	2	8	0	10.000	10.000	2.27	2.27	4.40	NAHQ ADVISORY
	2	8	0	10.000	10.000	0.04	0.04	251.00	EC20 POPULATION
	2	8	0	10.000	10.000	0.04	0.04	270.00	LTV EC20 FISH
	2	8	0	10.000	10.000	0.02	0.02	500.00	LTV EC20 DAPHNID
	2	8	0	10.000	10.000	0.01	0.01	717.00	LCV FISH
	2	8	0	10.000	10.000	0.01	0.01	717.00	LCV DAPHNIDS
	2	8	0	10.000	10.000	2.27	2.27	4.40	NAHQ ADVISORY
	4	10	0	10.000	10.000	0.04	0.04	251.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.04	0.04	270.00	LTV EC20 FISH
	4	10	0	10.000	10.000	0.02	0.02	500.00	LTV EC20 DAPHNID
	4	10	0	10.000	10.000	0.01	0.01	717.00	LCV FISH
	4	10	0	10.000	10.000	0.01	0.01	717.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	2.27	2.27	4.40	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.04	0.04	251.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.04	0.04	270.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.02	0.02	500.00	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	0.01	0.01	717.00	LCV FISH
	13	2	0	10.000	10.000	0.01	0.01	717.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	0.10	0.10	100.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.03	0.03	310.00	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	0.02	0.02	570.00	LCV DAPHNIDS
DI-N-OCTYLPHTHALATE	2	8	0	10.000	10.000	0.10	0.10	100.00	LTV EC20 FISH
	2	8	0	10.000	10.000	0.03	0.03	310.00	LTV EC20 DAPHNID
	2	8	0	10.000	10.000	0.02	0.02	570.00	LCV DAPHNIDS
	2	8	0	10.000	10.000	0.01	0.01	1995.00	EC20 POPULATION
	2	8	0	10.000	10.000	0.00	0.00	5700.00	LCV FISH
	2	8	0	10.000	10.000	0.10	0.10	100.00	LTV EC20 FISH
	3	179	0	10.000	10.148	0.03	0.03	310.00	LTV EC20 DAPHNID
	3	179	0	10.000	10.148	0.02	0.02	570.00	LCV DAPHNIDS
	3	179	0	10.000	10.148	0.01	0.01	1995.00	EC20 POPULATION
	3	179	0	10.000	10.148	0.00	0.00	5700.00	LCV FISH
	3	179	0	10.000	10.000	0.10	0.10	100.00	LTV EC20 FISH
	4	10	0	10.000	10.000	0.03	0.03	310.00	LTV EC20 DAPHNID
	4	10	0	10.000	10.000	0.02	0.02	570.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	0.01	0.01	1995.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.00	0.00	5700.00	LCV FISH
	4	10	0	10.000	10.000	0.10	0.10	100.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.10	0.10	100.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.03	0.03	310.00	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	0.02	0.02	570.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	0.01	0.01	1995.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.00	0.00	5706.00	LCV FISH
DIBENZOFURAN	3	1	0	10.000	10.000	5.00	5.00	2.00	NAHQ ADVISORY
	3	1	0	10.000	10.000	0.01	0.01	1003.00	LCV DAPHNIDS

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
DIETHYLPHTHALATE	2	8	0	10.000	10.000	0.37	0.37	27.00	NAHQ ADVISORY
	2	8	0	10.000	10.000	0.01	0.01	1000.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.37	0.37	27.00	NAHQ ADVISORY
	4	10	0	10.000	10.000	0.01	0.01	1000.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.37	0.37	27.00	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.01	0.01	1000.00	EC20 POPULATION
FLUORANTHENE	2	8	0	10.000	10.000	5.88	5.88	1.70	NAHQ ADVISORY
	2	8	0	10.000	10.000	0.31	0.31	32.00	EC20 POPULATION
	2	8	0	10.000	10.000	0.00	0.00	65795.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	5.88	5.88	1.70	NAHQ ADVISORY
	4	10	0	10.000	10.000	0.31	0.31	32.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.00	0.00	65795.00	LCV DAPHNIDS
GAMMA-BHC (LINDANE)	4	10	0	10.000	10.000	5.88	5.88	1.70	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.31	0.31	32.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.00	0.00	65795.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	0.00	0.00	65795.00	LCV DAPHNIDS
	3	1	0	0.050	0.050	0.62	0.62	0.08	NAHQ ADVISORY
	3	1	0	0.050	0.050	0.45	0.45	0.11	SS EC20
HEPTACHLOR	3	1	0	0.050	0.050	0.05	0.05	1.10	LTV EC20 FISH
	3	1	0	0.050	0.050	0.00	0.00	11.00	LTV EC20 DAPHNID
	3	1	0	0.050	0.050	0.00	0.00	14.50	LCV DAPHNIDS
	3	1	0	0.050	0.050	0.00	0.00	14.60	LCV FISH
	3	1	0	0.050	0.050	12.50	12.50	0.00	SS EC20
	3	1	0	0.050	0.050	5.00	5.00	0.01	NAHQ ADVISORY
LEAD	3	1	0	0.050	0.050	0.50	0.50	0.10	EC20 POPULATION
	3	1	0	0.050	0.050	0.06	0.06	0.86	LTV EC20 FISH
	3	1	0	0.050	0.050	0.04	0.04	1.26	LCV FISH
	3	1	0	0.050	0.050	0.02	0.02	3.18	LCV DAPHNIDS
	4	32	0	4.000	4.000	11.43	11.43	0.35	SS EC20
	4	32	0	4.000	4.000	1.25	1.25	3.20	NAHQ ACTUAL
	4	32	0	4.000	4.000	0.33	0.33	12.26	LCV DAPHNIDS
	4	32	0	4.000	4.000	0.21	0.21	18.88	LCV FISH
	4	32	0	4.000	4.000	0.18	0.18	22.00	LTV EC20 FISH
	4	32	0	4.000	4.000	0.06	0.06	71.00	EC20 POPULATION
	4	32	0	4.000	4.000	11.43	11.43	0.35	SS EC20
	13	8	0	4.000	4.000	1.25	1.25	3.20	NAHQ ACTUAL
	13	8	0	4.000	4.000	0.33	0.33	12.26	LCV DAPHNIDS
	13	8	0	4.000	4.000	0.21	0.21	18.88	LCV FISH
	13	8	0	4.000	4.000	0.18	0.18	22.00	LTV EC20 FISH
	13	8	0	4.000	4.000	0.06	0.06	71.00	EC20 POPULATION
	13	8	0	4.000	4.000	11.43	11.43	0.35	SS EC20
	13	8	0	4.000	4.000	1.25	1.25	3.20	NAHQ ACTUAL

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	IDL quotient	Benchmark	Benchmark type
MERCURY	2	24	0	0.200	0.200	666.67	666.67	0.00	MAW ADVISORY
	2	24	0	0.200	0.200	6.67	6.67	0.03	LTV EC20 FISH
	2	24	0	0.200	0.200	5.00	5.00	0.04	LCV DAPHNIDS
	2	24	0	0.200	0.200	0.71	0.71	0.28	EC20 POPULATION
	2	24	0	0.200	0.200	0.38	0.38	0.52	LCV FISH
	2	24	0	0.200	0.200	0.23	0.23	0.87	LTV EC20 DAPHNID
	13	8	0	0.200	0.200	666.67	666.67	0.00	MAW ADVISORY
	13	8	0	0.200	0.200	6.67	6.67	0.03	LTV EC20 FISH
	13	8	0	0.200	0.200	5.00	5.00	0.04	LCV DAPHNIDS
	13	8	0	0.200	0.200	0.71	0.71	0.28	EC20 POPULATION
	13	8	0	0.200	0.200	0.38	0.38	0.52	LCV FISH
	13	8	0	0.200	0.200	0.23	0.23	0.87	LTV EC20 DAPHNID
METHYLENE CHLORIDE	4	10	0	5.000	5.000	0.02	0.02	220.00	MAW ADVISORY
	4	10	0	5.000	5.000	0.01	0.01	410.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.00	0.00	1259.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	42667.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.00	0.00	108000.00	LCV FISH
	13	2	0	5.000	5.000	0.02	0.02	220.00	MAW ADVISORY
	13	2	0	5.000	5.000	0.01	0.01	410.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.00	0.00	1259.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	42667.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.00	0.00	108000.00	LCV FISH
MOLYBDENUM	1	64	0	40.000	40.000	1.33	1.33	30.00	MAW ADVISORY
	1	64	0	40.000	40.000	0.11	0.11	360.00	LTV EC20 DAPHNID
	1	64	0	40.000	40.000	0.05	0.05	880.00	LCV DAPHNIDS
	2	2	0	10.000	215.593	0.33	7.19	30.00	MAW ADVISORY
	2	2	0	10.000	215.593	0.03	0.60	360.00	LTV EC20 DAPHNID
	2	2	0	10.000	215.593	0.01	0.24	880.00	LCV DAPHNIDS
	4	1	0	10.000	10.000	0.33	0.33	30.00	MAW ADVISORY
	4	1	0	10.000	10.000	0.03	0.03	360.00	LTV EC20 DAPHNID
	4	1	0	10.000	10.000	0.01	0.01	880.00	LCV DAPHNIDS
N-NITROSODIPHENYLAMINE	2	8	0	10.000	10.000	4.00	4.00	2.50	MAW ADVISORY
	2	8	0	10.000	10.000	0.25	0.25	40.00	EC20 POPULATION
	2	8	0	10.000	10.000	0.03	0.03	332.00	LCV FISH
	2	8	0	10.000	10.000	0.03	0.03	339.00	LTV EC20 FISH
	2	8	0	10.000	10.000	0.01	0.01	1042.00	LCV DAPHNIDS
	3	179	0	10.000	10.148	4.00	4.06	2.50	MAW ADVISORY
	3	179	0	10.000	10.148	0.25	0.25	40.00	EC20 POPULATION
	3	179	0	10.000	10.148	0.03	0.03	332.00	LCV FISH
	3	179	0	10.000	10.148	0.03	0.03	339.00	LTV EC20 FISH
	3	179	0	10.000	10.148	0.01	0.01	1042.00	LCV DAPHNIDS

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
N-NITROSODIPHENYLAMINE (continued)	4	10	0	10.000	10.000	4.00	4.00	2.50	NAHQ ADVISORY
	4	10	0	10.000	10.000	0.25	0.25	40.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.03	0.03	332.00	LCV FISH
	4	10	0	10.000	10.000	0.03	0.03	339.00	LTV EC20 FISH
	4	10	0	10.000	10.000	0.01	0.01	1042.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	4.00	4.00	2.50	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.25	0.25	40.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.03	0.03	332.00	LCV FISH
	13	2	0	10.000	10.000	0.03	0.03	339.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.01	0.01	1042.00	LCV DAPHNIDS
NAPHTHALENE	2	8	0	10.000	10.000	4.76	4.76	2.10	NAHQ ADVISORY
	2	8	0	10.000	10.000	0.02	0.02	450.00	LTV EC20 FISH
	2	8	0	10.000	10.000	0.02	0.02	600.00	LTV EC20 DAPHNID
	2	8	0	10.000	10.000	0.02	0.02	620.00	LCV FISH
	2	8	0	10.000	10.000	0.01	0.01	1000.00	EC20 POPULATION
	2	8	0	10.000	10.000	0.01	0.01	1163.00	LCV DAPHNIDS
	3	179	0	10.000	10.148	4.76	4.83	2.10	NAHQ ADVISORY
	3	179	0	10.000	10.148	0.02	0.02	450.00	LTV EC20 FISH
	3	179	0	10.000	10.148	0.02	0.02	600.00	LTV EC20 DAPHNID
	3	179	0	10.000	10.148	0.02	0.02	620.00	LCV FISH
	3	179	0	10.000	10.148	0.01	0.01	1000.00	EC20 POPULATION
	3	179	0	10.000	10.148	0.01	0.01	1163.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	4.76	4.76	2.10	NAHQ ADVISORY
	4	10	0	10.000	10.000	0.02	0.02	450.00	LTV EC20 FISH
	4	10	0	10.000	10.000	0.02	0.02	600.00	LTV EC20 DAPHNID
	4	10	0	10.000	10.000	0.02	0.02	620.00	LCV FISH
	4	10	0	10.000	10.000	0.01	0.01	1000.00	EC20 POPULATION
	4	10	0	10.000	10.000	0.01	0.01	1163.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	4.76	4.76	2.10	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.02	0.02	450.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.02	0.02	600.00	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	0.02	0.02	620.00	LCV FISH
	13	2	0	10.000	10.000	0.01	0.01	1000.00	EC20 POPULATION
	13	2	0	10.000	10.000	0.01	0.01	1163.00	LCV DAPHNIDS

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
PHENANTHRENE	2	8	0	10.000	10.000	23.26	23.26	0.43	NAHQ ADVISORY
	2	8	0	10.000	10.000	0.09	0.09	110.00	LTV EC20 DAPHNID
	2	8	0	10.000	10.000	0.05	0.05	200.00	LCV DAPHNIDS
	4	10	0	10.000	10.000	23.26	23.26	0.43	NAHQ ADVISORY
	4	10	0	10.000	10.000	0.09	0.09	110.00	LTV EC20 DAPHNID
	4	10	0	10.000	10.000	0.05	0.05	200.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	23.26	23.26	0.43	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.09	0.09	110.00	LTV EC20 DAPHNID
	13	2	0	10.000	10.000	0.05	0.05	200.00	LCV DAPHNIDS
	13	2	0	10.000	10.000	0.48	0.48	21.00	NAHQ ADVISORY
	13	2	0	10.000	10.000	0.04	0.04	230.00	LTV EC20 FISH
	13	2	0	10.000	10.000	0.01	0.01	1369.00	LCV FISH
	13	2	0	10.000	10.000	0.00	0.00	2005.00	LCV DAPHNIDS
PHENOL	13	2	0	10.000	10.000	0.00	0.00	4467.00	EC20 POPULATION
	4	1	0	5.000	5.000	1.92	1.92	2.60	SS EC20
	4	1	0	5.000	5.000	0.20	0.20	25.00	LTV EC20 DAPHNID
	4	1	0	5.000	5.000	0.14	0.14	35.00	NAHQ ACTUAL
	4	1	0	5.000	5.000	0.13	0.13	40.00	LTV EC20 FISH
	4	1	0	5.000	5.000	0.06	0.06	88.32	LCV FISH
	4	1	0	5.000	5.000	0.05	0.05	91.65	LCV DAPHNIDS
	1	68	0	5.441	5.441	45.34	45.34	0.12	LCV FISH
	1	68	0	5.441	5.441	38.87	38.87	0.14	SS EC20
	1	68	0	5.441	5.441	27.21	27.21	0.20	LTV EC20 FISH
	1	68	0	5.441	5.441	21.76	21.76	0.25	NAHQ ADVISORY
	1	68	0	5.441	5.441	17.00	17.00	0.32	EC20 POPULATION
	1	68	0	5.441	5.441	9.72	9.72	0.56	LTV EC20 DAPHNID
SILVER	1	68	0	5.441	5.441	2.09	2.09	2.60	LCV DAPHNIDS
	1	68	0	5.441	5.441	50.00	50.00	0.12	LCV FISH
	2	1	0	6.000	6.000	42.86	42.86	0.14	SS EC20
	2	1	0	6.000	6.000	30.00	30.00	0.20	LTV EC20 FISH
	2	1	0	6.000	6.000	24.00	24.00	0.25	NAHQ ADVISORY
	2	1	0	6.000	6.000	18.75	18.75	0.32	EC20 POPULATION
	2	1	0	6.000	6.000	10.71	10.71	0.56	LTV EC20 DAPHNID
	2	1	0	6.000	6.000	2.31	2.31	2.60	LCV DAPHNIDS
	2	1	0	6.000	6.000	50.00	50.00	0.12	LCV FISH
	4	1	0	6.000	6.000	42.86	42.86	0.14	SS EC20
	4	1	0	6.000	6.000	30.00	30.00	0.20	LTV EC20 FISH
	4	1	0	6.000	6.000	24.00	24.00	0.25	NAHQ ADVISORY
	4	1	0	6.000	6.000	18.75	18.75	0.32	EC20 POPULATION
	4	1	0	6.000	6.000	10.71	10.71	0.56	LTV EC20 DAPHNID
	4	1	0	6.000	6.000	2.31	2.31	2.60	LCV DAPHNIDS

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
THALLIUM	2	1	0	10.000	10.000	4.00	4.00	2.50	NAHQ ADVISORY
	2	1	0	10.000	10.000	0.18	0.18	57.00	LCV FISH
	2	1	0	10.000	10.000	0.16	0.16	64.00	LTV EC20 DAPHNID
	2	1	0	10.000	10.000	0.15	0.15	67.00	EC20 POPULATION
	2	1	0	10.000	10.000	0.12	0.12	81.00	LTV EC20 FISH
	2	1	0	10.000	10.000	0.07	0.07	135.00	LCV DAPHNIDS
	2	1	0	10.000	10.000	0.07	0.07	135.00	NAHQ ADVISORY
	3	193	0	10.000	11.144	4.00	4.46	2.50	LCV FISH
	3	193	0	10.000	11.144	0.18	0.20	57.00	LTV EC20 DAPHNID
	3	193	0	10.000	11.144	0.16	0.17	64.00	EC20 POPULATION
	3	193	0	10.000	11.144	0.15	0.17	67.00	LTV EC20 FISH
	3	193	0	10.000	11.144	0.12	0.14	81.00	LTV EC20 FISH
	3	193	0	10.000	11.144	0.07	0.08	135.00	LCV DAPHNIDS
	3	193	0	10.000	11.144	0.07	0.08	135.00	NAHQ ADVISORY
	4	1	0	10.000	10.000	4.00	4.00	2.50	LCV FISH
	4	1	0	10.000	10.000	0.18	0.18	57.00	LTV EC20 DAPHNID
	4	1	0	10.000	10.000	0.16	0.16	64.00	EC20 POPULATION
	4	1	0	10.000	10.000	0.15	0.15	67.00	LTV EC20 FISH
	4	1	0	10.000	10.000	0.12	0.12	81.00	LTV EC20 FISH
	4	1	0	10.000	10.000	0.07	0.07	135.00	LCV DAPHNIDS
TOLUENE	4	10	0	5.000	5.000	0.48	0.48	10.40	NAHQ ADVISORY
	4	10	0	5.000	5.000	0.19	0.19	26.00	LTV EC20 FISH
	4	10	0	5.000	5.000	0.02	0.02	200.00	EC20 POPULATION
	4	10	0	5.000	5.000	0.00	0.00	1269.00	LCV FISH
	4	10	0	5.000	5.000	0.00	0.00	25229.00	LCV DAPHNIDS
	4	10	0	5.000	5.000	0.48	0.48	10.40	NAHQ ADVISORY
	13	2	0	5.000	5.000	0.19	0.19	26.00	LTV EC20 FISH
	13	2	0	5.000	5.000	0.02	0.02	200.00	EC20 POPULATION
	13	2	0	5.000	5.000	0.00	0.00	1269.00	LCV FISH
	13	2	0	5.000	5.000	0.00	0.00	25229.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.00	0.00	25229.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.00	0.00	25229.00	LCV DAPHNIDS
	13	2	0	5.000	5.000	0.00	0.00	25229.00	LCV DAPHNIDS
URANIUM	13	12	0	0.667	0.667	95.24	95.24	0.01	NAHQ ADVISORY
	13	12	0	0.667	0.667	0.02	0.02	27.00	EC20 POPULATION
	13	12	0	0.667	0.667	0.00	0.00	142.00	LCV FISH
	13	12	0	0.667	0.667	0.00	0.00	455.00	LTV EC20 FISH

Table F.4 (continued)

Analyte	Reach	Number analyzed in reach (water)	Number detects in reach (water)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UDL quotient	Benchmark	Benchmark type
VANADIUM	2	1	0	2,000	2,000	0.67	0.67	3.00	MAW ADVISORY
	2	1	0	2,000	2,000	0.06	0.06	32.00	EC20 POPULATION
	2	1	0	2,000	2,000	0.05	0.05	41.00	LTV EC20 FISH
	2	1	0	2,000	2,000	0.02	0.02	80.00	LCV FISH
	2	1	0	2,000	2,000	0.00	0.00	430.00	LTV EC20 DAPHNID
	2	1	0	2,000	2,000	0.00	0.00	940.00	LCV DAPHNIDS
	3	1	0	500,000	500,000	166.67	166.67	3.00	MAW ADVISORY
	3	1	0	500,000	500,000	15.63	15.63	32.00	EC20 POPULATION
	3	1	0	500,000	500,000	12.20	12.20	41.00	LTV EC20 FISH
	3	1	0	500,000	500,000	6.25	6.25	80.00	LCV FISH
	3	1	0	500,000	500,000	1.16	1.16	430.00	LTV EC20 DAPHNID
	3	1	0	500,000	500,000	0.53	0.53	940.00	LCV DAPHNIDS
VINYL ACETATE	2	32	0	9,687	9,687	12.11	12.11	0.80	MAW ADVISORY
	2	32	0	9,687	9,687	0.09	0.09	108.00	EC20 POPULATION
	2	32	0	9,687	9,687	0.01	0.01	718.00	LTV EC20 FISH
	2	32	0	9,687	9,687	0.01	0.01	810.00	LCV FISH
	3	2	0	10,000	10,000	12.50	12.50	0.80	MAW ADVISORY
	3	2	0	10,000	10,000	0.09	0.09	108.00	EC20 POPULATION
	3	2	0	10,000	10,000	0.01	0.01	718.00	LTV EC20 FISH
	3	2	0	10,000	10,000	0.01	0.01	810.00	LCV FISH
	2	41	0	9,687	10,106	10.20	10.64	0.95	MAW ADVISORY
	2	41	0	9,687	10,106	0.00	0.00	14520.00	LTV EC20 FISH
	2	41	0	9,687	10,106	0.00	0.00	28879.00	LCV FISH
	4	10	0	10,000	10,000	10.53	10.53	0.95	MAW ADVISORY
VINYL CHLORIDE	4	10	0	10,000	10,000	0.00	0.00	14520.00	LTV EC20 FISH
	4	10	0	10,000	10,000	0.00	0.00	28879.00	LCV FISH
	4	10	0	10,000	10,000	10.53	10.53	0.95	MAW ADVISORY
	13	2	0	10,000	10,000	0.00	0.00	14520.00	LTV EC20 FISH
	13	2	0	10,000	10,000	0.00	0.00	28879.00	LCV FISH
	13	2	0	10,000	10,000	3.42	3.42	1.46	MAW ADVISORY
	3	2	0	5,000	5,000	0.00	0.00	2680.00	LTV EC20 FISH
	3	2	0	5,000	5,000	0.00	0.00	62308.00	LCV FISH
	3	2	0	5,000	5,000	93.90	93.90	0.21	MAW ADVISORY
	2	1	0	20,000	20,000	0.08	0.08	251.00	EC20 POPULATION
	2	1	0	20,000	20,000	0.04	0.04	548.00	LCV FISH
	2	1	0	20,000	20,000	0.01	0.01	2396.00	LTV EC20 FISH
XYLENE	3	2	0	5,000	5,000	3.42	3.42	1.46	MAW ADVISORY
ZIRCONIUM	3	2	0	5,000	5,000	0.00	0.00	2680.00	LTV EC20 FISH
	3	2	0	5,000	5,000	0.00	0.00	62308.00	LCV FISH
	2	1	0	20,000	20,000	93.90	93.90	0.21	MAW ADVISORY
	2	1	0	20,000	20,000	0.08	0.08	251.00	EC20 POPULATION

^aThe maximum detection limit was used in cases where the UCL was missing (i.e., where N = 1).

Table F-5. Mean and upper confidence limit (UCL) concentrations of chemicals detected in fish (ug/L), bioconcentration factors, estimated exposure in water (ug/L), and quotients for exposure to that water concentration

Analyte	Reach (fish)	Number analyzed in reach	Number detects in reach	Mean fish ^a concentration	UCL fish ^a concentration	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag	Benchmark
4,4'-DDD	1	15	5	0.85	5.90	170000.00	0.0000	0.000	0.0119	0.08	0.00	NAMQ ADVISORY EC20 POPULATION
	1	15	5	0.85	5.90	170000.00	0.0000	0.000	0.0000	0.00	0.61	EC20 POPULATION
	1	15	5	0.85	5.90	170000.00	0.0000	0.000	0.0000	0.00	1.69	LCV FISH
	1	15	5	0.85	5.90	170000.00	0.0000	0.000	0.0000	0.00	3.99	LTV EC20 FISH
	3	34	2	0.30	17.08	170000.00	0.0000	0.000	0.0042	0.24	0.00	NAMQ ADVISORY EC20 POPULATION
	3	34	2	0.30	17.08	170000.00	0.0000	0.000	0.0000	0.00	0.61	EC20 POPULATION
	3	34	2	0.30	17.08	170000.00	0.0000	0.000	0.0000	0.00	1.69	LCV FISH
	3	34	2	0.30	17.08	170000.00	0.0000	0.000	0.0000	0.00	3.99	LTV EC20 FISH
	4	18	4	3.67	9.04	170000.00	0.0000	0.000	0.0514	0.13	0.00	NAMQ ADVISORY EC20 POPULATION
	4	18	4	3.67	9.04	170000.00	0.0000	0.000	0.0000	0.00	0.61	EC20 POPULATION
	4	18	4	3.67	9.04	170000.00	0.0000	0.000	0.0000	0.00	1.69	LCV FISH
	5	17	12	1.55	5.83	170000.00	0.0000	0.000	0.0000	0.00	3.99	LTV EC20 FISH
	5	17	12	1.55	5.83	170000.00	0.0000	0.000	0.0218	0.08	0.00	NAMQ ADVISORY EC20 POPULATION
	5	17	12	1.55	5.83	170000.00	0.0000	0.000	0.0000	0.00	0.61	EC20 POPULATION
	5	17	12	1.55	5.83	170000.00	0.0000	0.000	0.0000	0.00	1.69	LCV FISH
	10	8	7	0.69	1.22	170000.00	0.0000	0.000	0.0000	0.00	3.99	LTV EC20 FISH
	10	8	7	0.69	1.22	170000.00	0.0000	0.000	0.0096	0.02	0.00	NAMQ ADVISORY EC20 POPULATION
	10	8	7	0.69	1.22	170000.00	0.0000	0.000	0.0000	0.00	0.61	EC20 POPULATION
	10	8	7	0.69	1.22	170000.00	0.0000	0.000	0.0000	0.00	1.69	LCV FISH
	15	4	1	4.74	200.00	170000.00	0.0000	0.000	0.0664	2.80	0.00	NAMQ ADVISORY EC20 POPULATION
	15	4	1	4.74	200.00	170000.00	0.0000	0.001	0.0000	0.00	0.61	EC20 POPULATION
	15	4	1	4.74	200.00	170000.00	0.0000	0.001	0.0000	0.00	1.69	LCV FISH
	15	4	1	4.74	200.00	170000.00	0.0000	0.001	0.0000	0.00	3.99	LTV EC20 FISH
4,4'-DDE	1	16	8	5.30	64.28	8000000.00	0.0000	0.000	0.0001	0.00	0.01	SS EC20
	1	16	8	5.30	64.28	8000000.00	0.0000	0.000	0.0000	0.00	0.02	LCV DAPHNIDS
	1	16	8	5.30	64.28	8000000.00	0.0000	0.000	0.0000	0.00	0.02	NAMQ ADVISORY
	1	16	8	5.30	64.28	8000000.00	0.0000	0.000	0.0000	0.00	0.35	LTV EC20 FISH
	1	16	8	5.30	64.28	8000000.00	0.0000	0.000	0.0000	0.00	0.73	LCV FISH
	2	13	6	5.29	18.27	8000000.00	0.0000	0.000	0.0001	0.00	0.01	SS EC20
	2	13	6	5.29	18.27	8000000.00	0.0000	0.000	0.0000	0.00	0.02	LCV DAPHNIDS
	2	13	6	5.29	18.27	8000000.00	0.0000	0.000	0.0000	0.00	0.02	NAMQ ADVISORY
	2	13	6	5.29	18.27	8000000.00	0.0000	0.000	0.0000	0.00	0.35	LTV EC20 FISH
	2	13	6	5.29	18.27	8000000.00	0.0000	0.000	0.0000	0.00	0.73	LCV FISH
	3	34	34	1.04	9.10	8000000.00	0.0000	0.000	0.0000	0.00	0.01	SS EC20
	3	34	34	1.04	9.10	8000000.00	0.0000	0.000	0.0000	0.00	0.02	LCV DAPHNIDS
	3	34	34	1.04	9.10	8000000.00	0.0000	0.000	0.0000	0.00	0.02	NAMQ ADVISORY
	3	34	34	1.04	9.10	8000000.00	0.0000	0.000	0.0000	0.00	0.35	LTV EC20 FISH
	3	34	34	1.04	9.10	8000000.00	0.0000	0.000	0.0000	0.00	0.73	LCV FISH
	3	34	34	1.04	9.10	8000000.00	0.0000	0.000	0.0000	0.00	0.00	SS EC20

Table F.5 (continued)

Analyte	Reach	fish)	Number analyzed in reach	Number detects in reach	Mean fish ^a concentration	UCL Fish ^b concentration	BCF	Mean		UCL	Mean	UCL	Benchmark Flag	Benchmark type	
								estimated water conc.	estimated water conc.						
4,4'-DDE (continued)	4	18	12	11.09	24.40	8000000.00	0.0000	0.000	0.0002	0.01	0.00			SS EC20	
	4	18	12	11.09	24.40	8000000.00	0.0000	0.000	0.0001	0.02	0.00		*	LCV DAPHNIDS	
	4	18	12	11.09	24.40	8000000.00	0.0000	0.000	0.0001	0.02	0.00			NAWQ ADVISORY	
	4	18	12	11.09	24.40	8000000.00	0.0000	0.000	0.0000	0.35	0.00			LTV EC20 FISH	
	4	18	12	11.09	24.40	8000000.00	0.0000	0.000	0.0000	0.73	0.00			LCV FISH	
	5	18	9	8.03	13.23	8000000.00	0.0000	0.000	0.0001	0.01	0.00			SS EC20	
	5	18	9	8.03	13.23	8000000.00	0.0000	0.000	0.0001	0.02	0.00		*	LCV DAPHNIDS	
	5	18	9	8.03	13.23	8000000.00	0.0000	0.000	0.0000	0.02	0.00			NAWQ ADVISORY	
	5	18	9	8.03	13.23	8000000.00	0.0000	0.000	0.0000	0.35	0.00			LTV EC20 FISH	
	5	18	9	8.03	13.23	8000000.00	0.0000	0.000	0.0000	0.73	0.00			LCV FISH	
	10	8	2	2.56	2.95	8000000.00	0.0000	0.000	0.0000	0.01	0.00		*	SS EC20	
	10	8	2	2.56	2.95	8000000.00	0.0000	0.000	0.0000	0.02	0.00			LCV DAPHNIDS	
	10	8	2	2.56	2.95	8000000.00	0.0000	0.000	0.0000	0.02	0.00			NAWQ ADVISORY	
	10	8	2	2.56	2.95	8000000.00	0.0000	0.000	0.0000	0.35	0.00			LTV EC20 FISH	
	13	2	2	3.87	6.82	8000000.00	0.0000	0.000	0.0001	0.01	0.00		*	SS EC20	
	13	2	2	3.87	6.82	8000000.00	0.0000	0.000	0.0000	0.02	0.00			LCV DAPHNIDS	
	13	2	2	3.87	6.82	8000000.00	0.0000	0.000	0.0000	0.02	0.00			NAWQ ADVISORY	
	13	2	2	3.87	6.82	8000000.00	0.0000	0.000	0.0000	0.35	0.00			LTV EC20 FISH	
	13	2	2	3.87	6.82	8000000.00	0.0000	0.000	0.0000	0.73	0.00			LCV FISH	
4,4'-DDT	2	13	2	0.47	0.86	340000.00	0.0000	0.000	0.0017	0.01	0.00		*	SS EC20	
	2	13	2	0.47	0.86	340000.00	0.0000	0.000	0.0009	0.02	0.00			LCV DAPHNIDS	
	2	13	2	0.47	0.86	340000.00	0.0000	0.000	0.0007	0.02	0.00			NAWQ ADVISORY	
	2	13	2	0.47	0.86	340000.00	0.0000	0.000	0.0000	0.35	0.00			LTV EC20 FISH	
	2	13	2	0.47	0.86	340000.00	0.0000	0.000	0.0000	0.73	0.00			LCV FISH	
	4	18	1	1.58	200.00	340000.00	0.0000	0.006	0.0058	0.01	0.74		*	SS EC20	
	4	18	1	1.58	200.00	340000.00	0.0000	0.006	0.0029	0.02	0.37			LCV DAPHNIDS	
	4	18	1	1.58	200.00	340000.00	0.0000	0.006	0.0023	0.02	0.29			NAWQ ADVISORY	
	4	18	1	1.58	200.00	340000.00	0.0000	0.006	0.0001	0.35	0.02			LTV EC20 FISH	
	4	18	1	1.58	200.00	340000.00	0.0000	0.006	0.0001	0.73	0.01			LCV FISH	
	5	18	5	0.70	1.29	340000.00	0.0000	0.000	0.0026	0.01	0.00		*	SS EC20	
	5	18	5	0.70	1.29	340000.00	0.0000	0.000	0.0013	0.02	0.00			LCV DAPHNIDS	
	5	18	5	0.70	1.29	340000.00	0.0000	0.000	0.0010	0.02	0.00			NAWQ ADVISORY	
	5	18	5	0.70	1.29	340000.00	0.0000	0.000	0.0001	0.35	0.00			LTV EC20 FISH	
	5	18	5	0.70	1.29	340000.00	0.0000	0.000	0.0000	0.73	0.00			LCV FISH	
	ALDRIN	2	14	2	0.13	0.16	11000.00	0.0000	0.000	0.0002	0.08	0.00			NAWQ ADVISORY
		4	14	2	0.85	1.33	11000.00	0.0001	0.000	0.0010	0.08	0.00			NAWQ ADVISORY
		5	18	2	0.46	5.45	11000.00	0.0000	0.000	0.0006	0.01	0.00			NAWQ ADVISORY
		13	3	2	0.96	1.30	11000.00	0.0001	0.000	0.0012	0.08	0.00			NAWQ ADVISORY

Table F.5 (continued)

Analyte	Reach	Number analyzed in reach	Number detects in reach	Mean fish ^a concentration	UCL Fish ^b concentration	BCF	Mean	UCL	Mean	UCL	Benchmark Flag	Benchmark type	
							estimated water conc.	estimated water conc.		quotient			quotient
AROCLOP-1254	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0002	0.00	0.02	NAHQ ADVISORY	
	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0000	0.00	0.52	LTV EC20 FISH	
	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0000	0.00	0.63	EC20 POPULATION	
	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0000	0.00	1.00	LCV FISH	
	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0000	0.00	1.20	LTV EC20 DAPHNID	
	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0000	0.00	2.10	LCV DAPHNIDS	
	1	16	13	33.28	199.06	10000000.00	0.0000	0.000	0.0000	0.00	0.02	NAHQ ADVISORY	
	2	10	9	121.49	753.09	10000000.00	0.0000	0.000	0.0006	0.00	0.52	LTV EC20 FISH	
	2	10	9	121.49	753.09	10000000.00	0.0000	0.000	0.0000	0.00	0.63	EC20 POPULATION	
	2	10	9	121.49	753.09	10000000.00	0.0000	0.000	0.0000	0.00	1.00	LCV FISH	
	2	10	9	121.49	753.09	10000000.00	0.0000	0.000	0.0000	0.00	1.20	LTV EC20 DAPHNID	
	2	10	9	121.49	753.09	10000000.00	0.0000	0.000	0.0000	0.00	2.10	LCV DAPHNIDS	
	3	31	31	99.03	186.33	10000000.00	0.0000	0.000	0.0000	0.0005	0.00	0.02	NAHQ ADVISORY
	3	31	31	99.03	186.33	10000000.00	0.0000	0.000	0.0000	0.0000	0.00	0.52	LTV EC20 FISH
	3	31	31	99.03	186.33	10000000.00	0.0000	0.000	0.0000	0.0000	0.00	0.63	EC20 POPULATION
	3	31	31	99.03	186.33	10000000.00	0.0000	0.000	0.0000	0.0000	0.00	1.00	LCV FISH
	3	31	31	99.03	186.33	10000000.00	0.0000	0.000	0.0000	0.0000	0.00	1.20	LTV EC20 DAPHNID
	4	18	15	54.39	270.00	10000000.00	0.0000	0.0000	0.000	0.0003	0.00	0.02	NAHQ ADVISORY
	4	18	15	54.39	270.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.52	LTV EC20 FISH
	4	18	15	54.39	270.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.63	EC20 POPULATION
	4	18	15	54.39	270.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.00	LCV FISH
	4	18	15	54.39	270.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.20	LTV EC20 DAPHNID
	5	17	14	58.30	320.00	10000000.00	0.0000	0.0000	0.000	0.0003	0.00	0.02	NAHQ ADVISORY
	5	17	14	58.30	320.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.52	LTV EC20 FISH
	5	17	14	58.30	320.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.63	EC20 POPULATION
	5	17	14	58.30	320.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.00	LCV FISH
	5	17	14	58.30	320.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.20	LTV EC20 DAPHNID
	13	2	2	87.18	190.00	10000000.00	0.0000	0.0000	0.000	0.0004	0.00	0.02	NAHQ ADVISORY
	13	2	2	87.18	190.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.52	LTV EC20 FISH
	13	2	2	87.18	190.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.63	EC20 POPULATION
	13	2	2	87.18	190.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.00	LCV FISH
	13	2	2	87.18	190.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.20	LTV EC20 DAPHNID
	15	4	4	122.34	200.00	10000000.00	0.0000	0.0000	0.000	0.0006	0.00	0.02	NAHQ ADVISORY
	15	4	4	122.34	200.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.52	LTV EC20 FISH
	15	4	4	122.34	200.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	0.63	EC20 POPULATION
	15	4	4	122.34	200.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.00	LCV FISH
	15	4	4	122.34	200.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	1.20	LTV EC20 DAPHNID
	15	4	4	122.34	200.00	10000000.00	0.0000	0.0000	0.000	0.0000	0.00	2.10	LCV DAPHNIDS

Table F.5 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detected in reach (fish)	Mean fish ^a concentration	UCL Fish ^b	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag	Benchmark type
AROCLOL-1260	1	16	16	245.86	896.56	1000000.00	0.0000	0.000	0.0002	0.00	0.11	NAHQ ADVISORY
	1	16	16	245.86	896.56	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	1	16	16	245.86	896.56	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	1	16	16	245.86	896.56	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
	2	14	14	668.53	1406.39	1000000.00	0.0001	0.000	0.0006	0.00	0.11	NAHQ ADVISORY
	2	14	14	668.53	1406.39	1000000.00	0.0001	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	2	14	14	668.53	1406.39	1000000.00	0.0001	0.000	0.0000	0.00	2.30	LCV FISH
	2	14	14	668.53	1406.39	1000000.00	0.0001	0.000	0.0000	0.00	316.00	EC20 POPULATION
	3	36	36	390.85	770.54	1000000.00	0.0000	0.000	0.0004	0.00	0.11	NAHQ ADVISORY
	3	36	36	390.85	770.54	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	3	36	36	390.85	770.54	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	3	36	36	390.85	770.54	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
	4	18	17	433.14	2789.52	1000000.00	0.0000	0.000	0.0004	0.00	0.11	NAHQ ADVISORY
	4	18	17	433.14	2789.52	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	4	18	17	433.14	2789.52	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	4	18	17	433.14	2789.52	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
	5	19	19	347.76	889.60	1000000.00	0.0000	0.000	0.0000	0.00	0.11	NAHQ ADVISORY
	5	19	19	347.76	889.60	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	5	19	19	347.76	889.60	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	5	19	19	347.76	889.60	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
	10	8	7	21.98	73.10	1000000.00	0.0000	0.000	0.0000	0.00	0.11	NAHQ ADVISORY
	10	8	7	21.98	73.10	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	10	8	7	21.98	73.10	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	10	8	7	21.98	73.10	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
	13	4	4	288.12	390.00	1000000.00	0.0000	0.000	0.0003	0.00	0.11	NAHQ ADVISORY
	13	4	4	288.12	390.00	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	13	4	4	288.12	390.00	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	13	4	4	288.12	390.00	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
	15	4	4	310.50	650.00	1000000.00	0.0000	0.000	0.0003	0.00	0.11	NAHQ ADVISORY
	15	4	4	310.50	650.00	1000000.00	0.0000	0.000	0.0000	0.00	2.10	LTV EC20 FISH
	15	4	4	310.50	650.00	1000000.00	0.0000	0.000	0.0000	0.00	2.30	LCV FISH
	15	4	4	310.50	650.00	1000000.00	0.0000	0.000	0.0000	0.00	316.00	EC20 POPULATION
ARSENIC	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.4592	0.73	55.00	SS EC20
	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.1329	0.21	190.00	NAHQ ACTUAL
	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.0399	0.06	633.00	LTV EC20 DAPHNID
	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.0276	0.04	914.10	LCV DAPHNIDS
	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.0127	0.02	1995.00	EC20 POPULATION
	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.0119	0.02	2130.00	LTV EC20 FISH
	1	28	17	61.63	98.23	2.44	25.2574	40.258	0.0085	0.01	2962.00	LCV FISH
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.2828	0.73	55.00	SS EC20
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0819	0.21	190.00	NAHQ ACTUAL
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0819	0.21	190.00	NAHQ ACTUAL

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Mean fish ^a concentration	UCL Fish ^b concentration	BCF	Mean estimated water conc.		UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag	Benchmark type
							estimated	water conc.					
ARSENIC (continued)	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0246	0.06	633.00	LTV EC20 DAPHNID	
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0170	0.04	914.10	LCV DAPHNIDS	
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0078	0.02	1995.00	EC20 POPULATION	
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0073	0.02	2130.00	LTV EC20 FISH	
	2	13	5	37.96	98.02	2.44	15.5556	40.172	0.0053	0.01	2962.00	LCV FISH	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.3955	0.56	55.00	SS EC20	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.1145	0.16	190.00	NAWQ ACTUAL	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.0344	0.05	633.00	LTV EC20 DAPHNID	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.0238	0.03	914.10	LCV DAPHNIDS	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.0109	0.02	1995.00	EC20 POPULATION	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.0102	0.01	2130.00	LTV EC20 FISH	
	3	35	19	53.08	75.04	2.44	21.7539	30.756	0.0073	0.01	2962.00	LCV FISH	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.6552	0.80	55.00	SS EC20	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.1897	0.23	190.00	NAWQ ACTUAL	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.0569	0.07	633.00	LTV EC20 DAPHNID	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.0394	0.05	914.10	LCV DAPHNIDS	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.0181	0.02	1995.00	EC20 POPULATION	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.0169	0.02	2130.00	LTV EC20 FISH	
	4	27	25	87.93	107.22	2.44	36.0373	43.943	0.0122	0.01	2962.00	LCV FISH	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.5559	0.69	55.00	SS EC20	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.1609	0.20	190.00	NAWQ ACTUAL	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.0483	0.06	633.00	LTV EC20 DAPHNID	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.0335	0.04	914.10	LCV DAPHNIDS	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.0153	0.02	1995.00	EC20 POPULATION	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.0144	0.02	2130.00	LTV EC20 FISH	
	5	25	23	74.61	92.28	2.44	30.5769	37.819	0.0103	0.01	2962.00	LCV FISH	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.5323	0.72	55.00	SS EC20	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.1541	0.21	190.00	NAWQ ACTUAL	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.0462	0.06	633.00	LTV EC20 DAPHNID	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.0320	0.04	914.10	LCV DAPHNIDS	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.0147	0.02	1995.00	EC20 POPULATION	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.0137	0.02	2130.00	LTV EC20 FISH	
	10	15	14	71.43	96.82	2.44	29.2742	39.680	0.0099	0.01	2962.00	LCV FISH	
	13	10	10	45.14	59.01	2.44	18.5001	24.185	0.3364	0.44	55.00	SS EC20	
	13	10	4	45.14	59.01	2.44	18.5001	24.185	0.0974	0.13	190.00	NAWQ ACTUAL	
	13	10	4	45.14	59.01	2.44	18.5001	24.185	0.0292	0.04	633.00	LTV EC20 DAPHNID	
	13	10	4	45.14	59.01	2.44	18.5001	24.185	0.0202	0.03	914.10	LCV DAPHNIDS	
	13	10	4	45.14	59.01	2.44	18.5001	24.185	0.0093	0.01	1995.00	EC20 POPULATION	
	13	10	4	45.14	59.01	2.44	18.5001	24.185	0.0087	0.01	2130.00	LTV EC20 FISH	
	13	10	4	45.14	59.01	2.44	18.5001	24.185	0.0062	0.01	2962.00	LCV FISH	
13	10	4	45.14	59.01	2.44	18.5001	24.185	0.4820	0.84	55.00	SS EC20		
15	16	10	64.69	112.68	2.44	26.5123	46.181	0.1395	0.24	190.00	NAWQ ACTUAL		

Table F-5 (continued)

Analyte	Reach (fish)	Number analyzed in reach	Number detects in reach	Mean fish ^a concentration	UCL Fish ^a concentration	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag type
ARSENIC (continued)	15	16	10	64.69	112.68	2.44	26.5123	46.181	0.0419	0.07	633.00
	15	16	10	64.69	112.68	2.44	26.5123	46.181	0.0290	0.05	914.10
	15	16	10	64.69	112.68	2.44	26.5123	46.181	0.0133	0.02	1995.00
	15	16	10	64.69	112.68	2.44	26.5123	46.181	0.0124	0.02	2130.00
	15	16	10	64.69	112.68	2.44	26.5123	46.181	0.0090	0.02	2962.00
BERYLLIUM	1	25	4	1.47	3.26	2.00	0.7338	1.632	1.2029	2.67	0.61
	1	25	4	1.47	3.26	2.00	0.7338	1.632	0.1931	0.43	3.80
	1	25	4	1.47	3.26	2.00	0.7338	1.632	0.1384	0.31	5.30
	1	25	4	1.47	3.26	2.00	0.7338	1.632	0.0349	0.08	21.00
	1	25	4	1.47	3.26	2.00	0.7338	1.632	0.0129	0.03	57.00
	1	25	4	1.47	3.26	2.00	0.7338	1.632	0.0050	0.01	148.00
	3	35	7	0.92	5.45	2.00	0.4608	2.726	0.7554	4.47	0.61
	3	35	7	0.92	5.45	2.00	0.4608	2.726	0.1213	0.72	3.80
	3	35	7	0.92	5.45	2.00	0.4608	2.726	0.0869	0.51	5.30
	3	35	7	0.92	5.45	2.00	0.4608	2.726	0.0219	0.13	21.00
	3	35	7	0.92	5.45	2.00	0.4608	2.726	0.0081	0.05	57.00
	3	35	7	0.92	5.45	2.00	0.4608	2.726	0.0031	0.02	148.00
	5	25	2	0.02	27.00	2.00	0.0080	13.500	0.0132	22.13	0.61
	5	25	2	0.02	27.00	2.00	0.0080	13.500	0.0021	3.55	3.80
	5	25	2	0.02	27.00	2.00	0.0080	13.500	0.0015	2.55	5.30
	5	25	2	0.02	27.00	2.00	0.0080	13.500	0.0004	0.64	21.00
	5	25	2	0.02	27.00	2.00	0.0080	13.500	0.0001	0.24	57.00
	5	25	2	0.02	27.00	2.00	0.0080	13.500	0.0001	0.09	148.00
	10	10	1	0.01	37.00	2.00	0.0048	18.500	0.0079	30.33	0.61
	10	10	1	0.01	37.00	2.00	0.0048	18.500	0.0013	4.87	3.80
	10	10	1	0.01	37.00	2.00	0.0048	18.500	0.0009	3.49	5.30
	10	10	1	0.01	37.00	2.00	0.0048	18.500	0.0002	0.88	21.00
	10	10	1	0.01	37.00	2.00	0.0048	18.500	0.0001	0.32	57.00
	10	10	1	0.01	37.00	2.00	0.0048	18.500	0.0000	0.13	148.00
	15	16	5	1.32	19.00	2.00	0.6620	9.500	1.0853	15.57	0.61
	15	16	5	1.32	19.00	2.00	0.6620	9.500	0.1742	2.50	3.80
	15	16	5	1.32	19.00	2.00	0.6620	9.500	0.1249	1.79	5.30
	15	16	5	1.32	19.00	2.00	0.6620	9.500	0.0315	0.45	21.00
	15	16	5	1.32	19.00	2.00	0.6620	9.500	0.0116	0.17	57.00
	15	16	5	1.32	19.00	2.00	0.6620	9.500	0.0045	0.06	148.00

Table F.5 (continued)

Table 1-3 (continued)													
Analyte	Resch	Number analyzed in reach (fish)	Number detects in reach (fish)	Mean fish ^a concentration	UCL Fish ^b concentration	BCF	Mean estimated		UCL estimated	Mean quotient	UCL quotient	Benchmark Flag	Benchmark type
							water conc.	water conc.					
Bis(2-ETHYLHEXYL)PHTHALATE	3	33	1	47.09	410.00	310.00	0.1519	1.323	15.1912	132.26	0.01		NAHQ ADVISORY
	3	33	1	47.09	410.00	310.00	0.1519	1.323	0.0506	0.44	3.00	<	LCV DAPHNIDS
	3	33	1	47.09	410.00	310.00	0.1519	1.323	0.0506	0.44	3.00	<	LTV EC20 DAPHNID
	3	33	1	47.09	410.00	310.00	0.1519	1.323	0.0181	0.16	8.40		LCV FISH
	3	33	1	47.09	410.00	310.00	0.1519	1.323	0.0030	0.03	50.00		EC20 POPULATION
	3	33	1	47.09	410.00	310.00	0.1519	1.323	0.0028	0.02	54.00	>	LTV EC20 FISH
CHLORDANE	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0206	0.14	0.17		NAHQ ACTUAL
	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0140	0.09	0.25	<	LTV EC20 FISH
	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0070	0.05	0.50		SS EC20
	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0049	0.03	0.71		EC20 POPULATION
	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0022	0.01	1.60		LCV FISH
	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0003	0.00	12.10		LTV EC20 DAPHNID
	1	16	11	16.47	109.45	4702.00	0.0035	0.023	0.0002	0.00	16.00		LCV DAPHNIDS
	1	16	11	16.47	109.45	4702.00	0.0024	0.017	0.0143	0.10	0.17		NAHQ ACTUAL
	2	14	9	11.44	81.60	4702.00	0.0024	0.017	0.0097	0.07	0.25	<	LTV EC20 FISH
	2	14	9	11.44	81.60	4702.00	0.0024	0.017	0.0049	0.03	0.50		SS EC20
	2	14	9	11.44	81.60	4702.00	0.0024	0.017	0.0034	0.02	0.71		EC20 POPULATION
	2	14	9	11.44	81.60	4702.00	0.0024	0.017	0.0015	0.01	1.60		LCV FISH
	2	14	9	11.44	81.60	4702.00	0.0024	0.017	0.0002	0.00	12.10		LTV EC20 DAPHNID
	2	14	9	11.44	81.60	4702.00	0.0024	0.017	0.0002	0.00	16.00		LCV DAPHNIDS
	2	14	9	11.44	81.60	4702.00	0.0085	0.016	0.0499	0.09	0.17		NAHQ ACTUAL
	3	34	34	39.89	74.47	4702.00	0.0085	0.016	0.0339	0.06	0.25	<	LTV EC20 FISH
	3	34	34	39.89	74.47	4702.00	0.0085	0.016	0.0170	0.03	0.50		SS EC20
	3	34	34	39.89	74.47	4702.00	0.0085	0.016	0.0119	0.02	0.71		EC20 POPULATION
	3	34	34	39.89	74.47	4702.00	0.0085	0.016	0.0053	0.01	1.60		LCV FISH
	3	34	34	39.89	74.47	4702.00	0.0085	0.016	0.0007	0.00	12.10		LTV EC20 DAPHNID
	3	34	34	39.89	74.47	4702.00	0.0085	0.016	0.0005	0.00	16.00		LCV DAPHNIDS
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0355	0.12	0.17		NAHQ ACTUAL
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0241	0.08	0.25	<	LTV EC20 FISH
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0121	0.04	0.50		SS EC20
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0085	0.03	0.71		EC20 POPULATION
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0038	0.01	1.60		LCV FISH
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0005	0.00	12.10		LTV EC20 DAPHNID
	4	18	18	28.38	97.19	4702.00	0.0060	0.021	0.0004	0.00	16.00		LCV DAPHNIDS
	5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0133	0.06	0.17		NAHQ ACTUAL
	5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0090	0.04	0.25	<	LTV EC20 FISH
5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0045	0.02	0.50		SS EC20	
5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0032	0.01	0.71		EC20 POPULATION	
5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0014	0.01	1.60		LCV FISH	
5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0002	0.00	12.10		LTV EC20 DAPHNID	
5	18	18	10.62	45.49	4702.00	0.0023	0.010	0.0001	0.00	16.00		LCV DAPHNIDS	

Table F.5 (continued)

Analyte	Reach (fish)	Number analyzed in reach (fish)	Number detects in reach (fish)	Mean fish ^a concentration	UCL Fish ^b	SCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag type	Benchmark
CHLORDANE (continued)	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0104	0.02	0.17	NAHQ ACTUAL
	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0071	0.01	0.25	LTV EC20 FISH
	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0035	0.01	0.50	SS EC20
	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0025	0.00	0.71	EC20 POPULATION
	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0011	0.00	1.60	LCV FISH
	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0001	0.00	12.10	LTV EC20 DAPHNID
	10	8	8	8.32	15.71	4702.00	0.0018	0.003	0.0001	0.00	16.00	LCV DAPHNIDS
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0171	0.09	0.17	NAHQ ACTUAL
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0116	0.06	0.25	LTV EC20 FISH
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0058	0.03	0.50	SS EC20
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0041	0.02	0.71	EC20 POPULATION
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0018	0.01	1.60	LCV FISH
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0002	0.00	12.10	LTV EC20 DAPHNID
	13	4	2	13.69	72.89	4702.00	0.0029	0.016	0.0002	0.00	16.00	LCV DAPHNIDS
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0153	0.03	0.17	NAHQ ACTUAL
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0104	0.02	0.25	LTV EC20 FISH
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0052	0.01	0.50	SS EC20
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0037	0.01	0.71	EC20 POPULATION
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0016	0.00	1.60	LCV FISH
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0002	0.00	12.10	LTV EC20 DAPHNID
	15	4	4	12.25	22.08	4702.00	0.0026	0.005	0.0002	0.00	16.00	LCV DAPHNIDS
COPPER	5	25	6	46.48	170000.00	210.00	0.2213	809.524	1.0797	3948.90	0.20	LTV EC20 DAPHNID
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.9623	3519.67	0.23	LCV DAPHNIDS
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.8513	3113.55	0.26	SS EC20
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.0571	209.02	3.87	LCV FISH
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.0443	161.90	5.00	LTV EC20 FISH
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.0257	94.13	8.60	EC20 POPULATION
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.0184	67.46	12.00	NAHQ ACTUAL
	5	25	6	46.48	170000.00	210.00	0.2213	809.524	0.0000	0.00	100.00	LTV EC20 FISH
DI-N-OCTYLPHTHALATE	3	33	1	24.05	410.00	9300.00	0.0026	0.044	0.0000	0.00	310.00	LTV EC20 DAPHNID
	3	33	1	24.05	410.00	9300.00	0.0026	0.044	0.0000	0.00	570.00	LCV DAPHNIDS
	3	33	1	24.05	410.00	9300.00	0.0026	0.044	0.0000	0.00	1995.00	EC20 POPULATION
	3	33	1	24.05	410.00	9300.00	0.0026	0.044	0.0000	0.00	5700.00	LCV FISH
	3	33	1	24.05	410.00	9300.00	0.0026	0.044	0.0000	0.00	5700.00	LCV FISH

Table F.5 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Mean fish ^a concentration	UCL Fish ^b	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag type	Benchmark
MERCURY	1	28	28	53.67	81.95	63000.00	0.0009	0.001	2.8399	4.34	0.00	NAHQ ADVISORY
	1	28	28	53.67	81.95	63000.00	0.0009	0.001	0.0284	0.04	0.03	LTV EC20 FISH
	1	28	28	53.67	81.95	63000.00	0.0009	0.001	0.0213	0.03	0.04	LCV DAPHNIDS
	1	28	28	53.67	81.95	63000.00	0.0009	0.001	0.0030	0.00	0.28	EC20 POPULATION
	1	28	28	53.67	81.95	63000.00	0.0009	0.001	0.0016	0.00	0.52	LCV FISH
	1	28	28	53.67	81.95	63000.00	0.0009	0.001	0.0010	0.00	0.87	LTV EC20 DAPHNID
	2	13	13	63.84	134.97	63000.00	0.0010	0.002	3.3779	7.14	0.00	NAHQ ADVISORY
	2	13	13	63.84	134.97	63000.00	0.0010	0.002	0.0338	0.07	0.03	LTV EC20 FISH
	2	13	13	63.84	134.97	63000.00	0.0010	0.002	0.0253	0.05	0.04	LCV DAPHNIDS
	2	13	13	63.84	134.97	63000.00	0.0010	0.002	0.0036	0.01	0.28	EC20 POPULATION
	2	13	13	63.84	134.97	63000.00	0.0010	0.002	0.0019	0.00	0.52	LCV FISH
	2	13	13	63.84	134.97	63000.00	0.0010	0.002	0.0012	0.00	0.87	LTV EC20 DAPHNID
	3	35	35	422.59	615.51	63000.00	0.0067	0.010	22.3592	32.57	0.00	NAHQ ADVISORY
	3	35	35	422.59	615.51	63000.00	0.0067	0.010	0.2236	0.33	0.03	LTV EC20 FISH
	3	35	35	422.59	615.51	63000.00	0.0067	0.010	0.1677	0.24	0.04	LCV DAPHNIDS
	3	35	35	422.59	615.51	63000.00	0.0067	0.010	0.0240	0.03	0.28	EC20 POPULATION
	3	35	35	422.59	615.51	63000.00	0.0067	0.010	0.0129	0.02	0.52	LCV FISH
	3	35	35	422.59	615.51	63000.00	0.0067	0.010	0.0077	0.01	0.87	LTV EC20 DAPHNID
	4	27	27	185.34	390.65	63000.00	0.0029	0.006	9.8061	20.67	0.00	NAHQ ADVISORY
	4	27	27	185.34	390.65	63000.00	0.0029	0.006	0.0981	0.21	0.03	LTV EC20 FISH
	4	27	27	185.34	390.65	63000.00	0.0029	0.006	0.0735	0.16	0.04	LCV DAPHNIDS
	4	27	27	185.34	390.65	63000.00	0.0029	0.006	0.0105	0.02	0.28	EC20 POPULATION
	4	27	27	185.34	390.65	63000.00	0.0029	0.006	0.0057	0.01	0.52	LCV FISH
	4	27	27	185.34	390.65	63000.00	0.0029	0.006	0.0034	0.01	0.87	LTV EC20 DAPHNID
	5	25	25	65.98	107.91	63000.00	0.0010	0.002	3.4912	5.71	0.00	NAHQ ADVISORY
	5	25	25	65.98	107.91	63000.00	0.0010	0.002	0.0349	0.06	0.03	LTV EC20 FISH
	5	25	25	65.98	107.91	63000.00	0.0010	0.002	0.0262	0.04	0.04	LCV DAPHNIDS
	5	25	25	65.98	107.91	63000.00	0.0010	0.002	0.0037	0.01	0.28	EC20 POPULATION
	5	25	25	65.98	107.91	63000.00	0.0010	0.002	0.0020	0.00	0.52	LCV FISH
	5	25	25	65.98	107.91	63000.00	0.0010	0.002	0.0012	0.00	0.87	LTV EC20 DAPHNID
	10	15	15	190.08	279.96	63000.00	0.0030	0.004	10.0574	14.81	0.00	NAHQ ADVISORY
	10	15	15	190.08	279.96	63000.00	0.0030	0.004	0.1006	0.15	0.03	LTV EC20 FISH
	10	15	15	190.08	279.96	63000.00	0.0030	0.004	0.0754	0.11	0.04	LCV DAPHNIDS
	10	15	15	190.08	279.96	63000.00	0.0030	0.004	0.0108	0.02	0.28	EC20 POPULATION
	10	15	15	190.08	279.96	63000.00	0.0030	0.004	0.0058	0.01	0.52	LCV FISH
	10	15	15	190.08	279.96	63000.00	0.0030	0.004	0.0035	0.01	0.87	LTV EC20 DAPHNID
	13	10	10	105.51	367.42	63000.00	0.0017	0.006	5.5827	19.44	0.00	NAHQ ADVISORY
	13	10	10	105.51	367.42	63000.00	0.0017	0.006	0.0558	0.19	0.03	LTV EC20 FISH
	13	10	10	105.51	367.42	63000.00	0.0017	0.006	0.0419	0.15	0.04	LCV DAPHNIDS
	13	10	10	105.51	367.42	63000.00	0.0017	0.006	0.0060	0.02	0.28	EC20 POPULATION
	13	10	10	105.51	367.42	63000.00	0.0017	0.006	0.0032	0.01	0.52	LCV FISH
	13	10	10	105.51	367.42	63000.00	0.0017	0.006	0.0019	0.01	0.87	LTV EC20 DAPHNID

Table F.5 (continued)

Analyte	Reach (fish)	Number analyzed in reach (fish)	Number detects in reach (fish)	Mean fish ^a concentration	UCL Fish ^b concentration	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag type	Benchmark
MERCURY (continued)	15	16	16	132.09	218.29	63000.00	0.0021	0.003	6.9888	11.55	0.00	NAHQ ADVISORY
	15	16	16	132.09	218.29	63000.00	0.0021	0.003	0.0699	0.12	0.03	LTV EC20 FISH
	15	16	16	132.09	218.29	63000.00	0.0021	0.003	0.0524	0.09	0.04	LCV DAPHNIDS
	15	16	16	132.09	218.29	63000.00	0.0021	0.003	0.0075	0.01	0.28	EC20 POPULATION
	15	16	16	132.09	218.29	63000.00	0.0021	0.003	0.0040	0.01	0.52	LCV FISH
	15	16	16	132.09	218.29	63000.00	0.0021	0.003	0.0024	0.00	0.87	LTV EC20 DAPHNIDS
SELENIUM	1	28	26	748.33	834.48	8.00	93.5412	104.310	35.9774	40.12	2.60	SS EC20
	1	28	26	748.33	834.48	8.00	93.5412	104.310	3.7416	4.17	25.00	LTV EC20 DAPHNIDS
	1	28	26	748.33	834.48	8.00	93.5412	104.310	2.6726	2.98	35.00	NAHQ ACTUAL
	1	28	26	748.33	834.48	8.00	93.5412	104.310	2.3385	2.61	40.00	LTV EC20 FISH
	1	28	26	748.33	834.48	8.00	93.5412	104.310	1.0591	1.18	88.32	LCV FISH
	1	28	26	748.33	834.48	8.00	93.5412	104.310	1.0206	1.14	91.65	LCV DAPHNIDS
	2	13	11	726.04	850.69	8.00	90.7551	106.337	34.9058	40.90	2.60	SS EC20
	2	13	11	726.04	850.69	8.00	90.7551	106.337	3.6302	4.25	25.00	LTV EC20 DAPHNIDS
	2	13	11	726.04	850.69	8.00	90.7551	106.337	2.5930	3.04	35.00	NAHQ ACTUAL
	2	13	11	726.04	850.69	8.00	90.7551	106.337	2.2689	2.66	40.00	LTV EC20 FISH
	2	13	11	726.04	850.69	8.00	90.7551	106.337	1.0276	1.20	88.32	LCV FISH
	2	13	11	726.04	850.69	8.00	90.7551	106.337	0.9902	1.16	91.65	LCV DAPHNIDS
	3	35	2	231.39	576.88	8.00	28.9237	72.110	11.1245	27.73	2.60	SS EC20
	3	35	2	231.39	576.88	8.00	28.9237	72.110	1.1569	2.88	25.00	LTV EC20 DAPHNIDS
	3	35	2	231.39	576.88	8.00	28.9237	72.110	0.8264	2.06	35.00	NAHQ ACTUAL
	3	35	2	231.39	576.88	8.00	28.9237	72.110	0.7231	1.80	40.00	LTV EC20 FISH
	3	35	2	231.39	576.88	8.00	28.9237	72.110	0.3275	0.82	88.32	LCV FISH
	3	35	2	231.39	576.88	8.00	28.9237	72.110	0.3156	0.79	91.65	LCV DAPHNIDS
	4	27	14	556.53	686.35	8.00	69.5667	85.793	26.7564	33.00	2.60	SS EC20
	4	27	14	556.53	686.35	8.00	69.5667	85.793	2.7827	3.43	25.00	LTV EC20 DAPHNIDS
	4	27	14	556.53	686.35	8.00	69.5667	85.793	1.9876	2.45	35.00	NAHQ ACTUAL
	4	27	14	556.53	686.35	8.00	69.5667	85.793	1.7392	2.14	40.00	LTV EC20 FISH
	4	27	14	556.53	686.35	8.00	69.5667	85.793	0.7877	0.97	88.32	LCV FISH
	4	27	14	556.53	686.35	8.00	69.5667	85.793	0.7590	0.94	91.65	LCV DAPHNIDS
	10	15	8	521.86	582.48	8.00	65.2320	72.810	25.0892	28.00	2.60	SS EC20
	10	15	8	521.86	582.48	8.00	65.2320	72.810	2.6093	2.91	25.00	LTV EC20 DAPHNIDS
	10	15	8	521.86	582.48	8.00	65.2320	72.810	1.8638	2.08	35.00	NAHQ ACTUAL
	10	15	8	521.86	582.48	8.00	65.2320	72.810	1.6308	1.82	40.00	LTV EC20 FISH
	10	15	8	521.86	582.48	8.00	65.2320	72.810	0.7386	0.82	88.32	LCV FISH
	10	15	8	521.86	582.48	8.00	65.2320	72.810	0.7118	0.79	91.65	LCV DAPHNIDS

Table P.5 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number defects in reach (fish)	Mean fish ^a concentration	UCL Fish ^a	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag type	Benchmark
SELENIUM (continued)	15	16	8	524.64	679.48	8.00	65.5796	84.935	25.2229	32.67	2.60	SS EC20
	15	16	8	524.64	679.48	8.00	65.5796	84.935	2.6232	3.40	25.00	LTV EC20 DAPHNID
	15	16	8	524.64	679.48	8.00	65.5796	84.935	1.8737	2.43	35.00	NAHQ ACTUAL
	15	16	8	524.64	679.48	8.00	65.5796	84.935	1.6395	2.12	40.00	LTV EC20 FISH
	15	16	8	524.64	679.48	8.00	65.5796	84.935	0.96	0.96	88.32	LCV FISH
	15	16	8	524.64	679.48	8.00	65.5796	84.935	0.7425	0.93	91.65	LCV DAPHNIDS
	15	16	8	524.64	679.48	8.00	65.5796	84.935	0.7155	0.93	91.65	LCV DAPHNIDS
	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.3138	0.36	21.00	SS EC20
	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.1810	0.21	36.41	LCV FISH
	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.1410	0.16	46.73	LCV DAPHNIDS
ZINC	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.1402	0.16	47.00	LTV EC20 FISH
	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.0824	0.09	80.00	EC20 POPULATION
	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.0599	0.07	110.00	NAHQ ACTUAL
	1	28	28	6589.72	7546.46	1000.00	6.5897	7.546	0.2734	0.31	21.00	SS EC20
	2	13	13	5741.98	6566.14	1000.00	5.7420	6.566	0.1577	0.18	36.41	LCV FISH
	2	13	13	5741.98	6566.14	1000.00	5.7420	6.566	0.1229	0.14	46.73	LCV DAPHNIDS
	2	13	13	5741.98	6566.14	1000.00	5.7420	6.566	0.1222	0.14	47.00	LTV EC20 FISH
	2	13	13	5741.98	6566.14	1000.00	5.7420	6.566	0.0718	0.08	80.00	EC20 POPULATION
	2	13	13	5741.98	6566.14	1000.00	5.7420	6.566	0.0522	0.06	110.00	NAHQ ACTUAL
	2	13	13	5741.98	6566.14	1000.00	5.7420	6.566	0.2745	0.31	21.00	SS EC20
	3	35	35	5763.90	6551.08	1000.00	5.7639	6.551	0.1583	0.18	36.41	LCV FISH
	3	35	35	5763.90	6551.08	1000.00	5.7639	6.551	0.1233	0.14	46.73	LCV DAPHNIDS
	3	35	35	5763.90	6551.08	1000.00	5.7639	6.551	0.1226	0.14	47.00	LTV EC20 FISH
	3	35	35	5763.90	6551.08	1000.00	5.7639	6.551	0.0720	0.08	80.00	EC20 POPULATION
	3	35	35	5763.90	6551.08	1000.00	5.7639	6.551	0.0524	0.06	110.00	NAHQ ACTUAL
	3	35	35	5763.90	6551.08	1000.00	5.7639	6.551	0.2737	0.31	21.00	SS EC20
	4	27	27	5746.66	6545.24	1000.00	5.7467	6.545	0.1578	0.18	36.41	LCV FISH
	4	27	27	5746.66	6545.24	1000.00	5.7467	6.545	0.1230	0.14	46.73	LCV DAPHNIDS
	4	27	27	5746.66	6545.24	1000.00	5.7467	6.545	0.1233	0.14	47.00	LTV EC20 FISH
	4	27	27	5746.66	6545.24	1000.00	5.7467	6.545	0.0718	0.08	80.00	EC20 POPULATION
	4	27	27	5746.66	6545.24	1000.00	5.7467	6.545	0.0522	0.06	110.00	NAHQ ACTUAL
	4	27	27	5746.66	6545.24	1000.00	5.7467	6.545	0.2737	0.31	21.00	SS EC20
	5	24	24	6977.21	10929.98	1000.00	6.9772	10.930	0.3322	0.52	21.00	LCV FISH
	5	24	24	6977.21	10929.98	1000.00	6.9772	10.930	0.1916	0.30	36.41	LCV DAPHNIDS
	5	24	24	6977.21	10929.98	1000.00	6.9772	10.930	0.1493	0.23	46.73	LTV EC20 FISH
	5	24	24	6977.21	10929.98	1000.00	6.9772	10.930	0.1485	0.23	47.00	EC20 POPULATION
	5	24	24	6977.21	10929.98	1000.00	6.9772	10.930	0.0872	0.14	80.00	NAHQ ACTUAL
	5	24	24	6977.21	10929.98	1000.00	6.9772	10.930	0.0634	0.10	110.00	NAHQ ACTUAL

Table F.5 (continued)

Analyte	Reach (fish)	Number analyzed in reach	Number detects in reach	Mean fish ^a concentration	UCL Fish ^b concentration	BCF	Mean estimated water conc.	UCL estimated water conc.	Mean quotient	UCL quotient	Benchmark Flag type	Benchmark
ZINC (continued)	10	15	15	5221.38	5966.77	1000.00	5.2214	5.967	0.2486	0.28	21.00	SS EC20
	10	15	15	5221.38	5966.77	1000.00	5.2214	5.967	0.1434	0.16	36.41	LCV FISH
	10	15	15	5221.38	5966.77	1000.00	5.2214	5.967	0.1117	0.13	46.73	LCV DAPHNIDS
	10	15	15	5221.38	5966.77	1000.00	5.2214	5.967	0.1111	0.13	47.00	LTV EC20 FISH
	10	15	15	5221.38	5966.77	1000.00	5.2214	5.967	0.0653	0.07	80.00	EC20 POPULATION
	10	15	15	5221.38	5966.77	1000.00	5.2214	5.967	0.0475	0.05	110.00	MAWQ ACTUAL
	13	10	10	6313.03	6976.61	1000.00	6.3130	6.977	0.3006	0.33	21.00	SS EC20
	13	10	10	6313.03	6976.61	1000.00	6.3130	6.977	0.1734	0.19	36.41	LCV FISH
	13	10	10	6313.03	6976.61	1000.00	6.3130	6.977	0.1351	0.15	46.73	LCV DAPHNIDS
	13	10	10	6313.03	6976.61	1000.00	6.3130	6.977	0.1343	0.15	47.00	LTV EC20 FISH
	13	10	10	6313.03	6976.61	1000.00	6.3130	6.977	0.0789	0.09	80.00	EC20 POPULATION
	13	10	10	6313.03	6976.61	1000.00	6.3130	6.977	0.0574	0.06	110.00	MAWQ ACTUAL
	15	16	16	5772.05	6898.93	1000.00	5.7720	6.899	0.2749	0.33	21.00	SS EC20
	15	16	16	5772.05	6898.93	1000.00	5.7720	6.899	0.1585	0.19	36.41	LCV FISH
	15	16	16	5772.05	6898.93	1000.00	5.7720	6.899	0.1235	0.15	46.73	LCV DAPHNIDS
	15	16	16	5772.05	6898.93	1000.00	5.7720	6.899	0.1228	0.15	47.00	LTV EC20 FISH
	15	16	16	5772.05	6898.93	1000.00	5.7720	6.899	0.0722	0.09	80.00	EC20 POPULATION
	15	16	16	5772.05	6898.93	1000.00	5.7720	6.899	0.0525	0.06	110.00	MAWQ ACTUAL

^aIn cases of mixed detects and nondetects, statistics are calculated using all values, "as is." The mean concentration is the geometric mean.

^bUCL concentration was compared with the maximum value and the smaller has been used as the "UCL concentration."

Table F.6. Limits of detection of chemicals not detected in fish (ug/L), bioconcentration factors, estimated exposure in water (ug/L), and quotients for exposure to that water concentration

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
2,4-DINITROPHENOL	1	6	0	3400.00	4300.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	2	8	0	3500.00	9300.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	3	34	0	930.00	2100.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	4	13	0	3300.00	4100.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	5	9	0	3300.00	4200.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	13	2	0	3400.00	3500.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	15	4	0	3400.00	3800.00	.	.	.	.	.	150.00	.	NAHQ ACTUAL
	1	6	0	680.00	870.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	1	6	0	680.00	870.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	1	6	0	680.00	870.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
	1	6	0	680.00	870.00	.	.	.	.	.	489.00	*	LCV FISH
	1	6	0	680.00	870.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS
	1	6	0	700.00	1900.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	2	8	0	700.00	1900.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	2	8	0	700.00	1900.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
2-METHYLPHENOL	2	8	0	700.00	1900.00	.	.	.	.	.	489.00	*	LCV FISH
	2	8	0	700.00	1900.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	2	8	0	700.00	1900.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	3	34	0	190.00	410.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
	3	34	0	190.00	410.00	.	.	.	.	.	489.00	*	LCV FISH
	3	34	0	190.00	410.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS
	3	34	0	190.00	410.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	4	13	0	660.00	830.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	4	13	0	660.00	830.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
	4	13	0	660.00	830.00	.	.	.	.	.	489.00	*	LCV FISH
	4	13	0	660.00	830.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS
	4	13	0	660.00	830.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	5	9	0	660.00	830.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	5	9	0	660.00	830.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
	5	9	0	660.00	830.00	.	.	.	.	.	489.00	*	LCV FISH
	5	9	0	660.00	830.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS
	13	2	0	690.00	700.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	13	2	0	690.00	700.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	13	2	0	690.00	700.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
	13	2	0	690.00	700.00	.	.	.	.	.	489.00	*	LCV FISH
	13	2	0	690.00	700.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS
	13	2	0	690.00	700.00	.	.	.	.	.	7.70	.	NAHQ ADVISORY
	15	4	0	670.00	770.00	.	.	.	.	.	74.00	.	EC20 POPULATION
	15	4	0	670.00	770.00	.	.	.	.	.	470.00	*	LTV EC20 FISH
	15	4	0	670.00	770.00	.	.	.	.	.	489.00	*	LCV FISH
	15	4	0	670.00	770.00	.	.	.	.	.	1316.00	*	LCV DAPHNIDS

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
2-NITROPHENOL	1	6	0	680.00	870.00	150	4.533	5.800	0.03	0.04	150.00		NAHQ ACTUAL
	2	8	0	700.00	1900.00	150	4.667	12.667	0.03	0.08	150.00		NAHQ ACTUAL
	3	34	0	190.00	410.00	150	1.267	2.733	0.01	0.02	150.00		NAHQ ACTUAL
	4	13	0	660.00	830.00	150	4.400	5.533	0.03	0.04	150.00		NAHQ ACTUAL
	5	9	0	660.00	830.00	150	4.400	5.533	0.03	0.04	150.00		NAHQ ACTUAL
	13	2	0	690.00	700.00	150	4.600	4.667	0.03	0.03	150.00		NAHQ ACTUAL
4,4'-DDD	15	4	0	670.00	770.00	150	4.467	5.133	0.03	0.03	150.00		NAHQ ACTUAL
	2	14	0	10.00	100.00	170000	0.000	0.001	0.14	1.40	0.00		NAHQ ADVISORY
	2	14	0	10.00	100.00	170000	0.000	0.001	0.00	0.00	0.61		EC20 POPULATION
	2	14	0	10.00	100.00	170000	0.000	0.001	0.00	0.00	0.61	*	LCV FISH
	13	4	0	10.00	100.00	170000	0.000	0.001	0.00	0.00	3.99	*	LTV EC20 FISH
	13	4	0	10.00	100.00	170000	0.000	0.001	0.14	0.14	0.00		NAHQ ADVISORY
4,4'-DDE	13	4	0	10.00	10.00	170000	0.000	0.000	0.00	0.00	0.61	*	EC20 POPULATION
	13	4	0	10.00	10.00	170000	0.000	0.000	0.00	0.00	1.69	*	LCV FISH
	13	4	0	10.00	10.00	170000	0.000	0.000	0.00	0.00	3.99	*	LTV EC20 FISH
	15	4	0	50.00	200.00	8000000	0.000	0.000	0.00	0.00	0.01		SS EC20
	15	4	0	50.00	200.00	8000000	0.000	0.000	0.00	0.00	0.02	*	LCV DAPHNIDS
	15	4	0	50.00	200.00	8000000	0.000	0.000	0.00	0.00	0.02		NAHQ ADVISORY
4,4'-DDT	15	4	0	50.00	200.00	8000000	0.000	0.000	0.00	0.00	0.35		LTV EC20 FISH
	15	4	0	50.00	200.00	8000000	0.000	0.000	0.00	0.00	0.73		LCV FISH
	1	16	0	10.00	200.00	34000	0.000	0.000	0.04	0.74	0.01		SS EC20
	1	16	0	10.00	200.00	34000	0.000	0.000	0.02	0.37	0.02	*	LCV DAPHNIDS
	1	16	0	10.00	200.00	34000	0.000	0.000	0.01	0.29	0.02		NAHQ ADVISORY
	1	16	0	10.00	200.00	34000	0.000	0.000	0.00	0.02	0.35		LTV EC20 FISH
4,4'-DDT	1	16	0	10.00	200.00	34000	0.000	0.000	0.00	0.01	0.73		LCV FISH
	3	34	0	10.00	17.08	34000	0.000	0.001	0.04	0.06	0.01		SS EC20
	3	34	0	10.00	17.08	34000	0.000	0.001	0.02	0.03	0.02	*	LCV DAPHNIDS
	3	34	0	10.00	17.08	34000	0.000	0.001	0.01	0.03	0.02		NAHQ ADVISORY
	3	34	0	10.00	17.08	34000	0.000	0.001	0.00	0.00	0.35		LTV EC20 FISH
	3	34	0	10.00	17.08	34000	0.000	0.001	0.00	0.00	0.73		LCV FISH
	10	8	0	10.00	10.00	34000	0.000	0.000	0.04	0.04	0.01		SS EC20
	10	8	0	10.00	10.00	34000	0.000	0.000	0.02	0.02	0.02	*	LCV DAPHNIDS
	10	8	0	10.00	10.00	34000	0.000	0.000	0.01	0.01	0.02		NAHQ ADVISORY
	10	8	0	10.00	10.00	34000	0.000	0.000	0.00	0.00	0.35		LTV EC20 FISH
	10	8	0	10.00	10.00	34000	0.000	0.000	0.00	0.00	0.73		LCV FISH
	10	8	0	10.00	10.00	34000	0.000	0.000	0.04	0.04	0.01		SS EC20
	13	3	0	10.00	10.00	34000	0.000	0.000	0.02	0.02	0.02	*	LCV DAPHNIDS
	13	3	0	10.00	10.00	34000	0.000	0.000	0.01	0.01	0.02		NAHQ ADVISORY
	13	3	0	10.00	10.00	34000	0.000	0.000	0.00	0.00	0.35		LTV EC20 FISH
	13	3	0	10.00	10.00	34000	0.000	0.000	0.00	0.00	0.73		LCV FISH
	15	4	0	50.00	200.00	34000	0.001	0.006	0.18	0.74	0.01		SS EC20
	15	4	0	50.00	200.00	34000	0.001	0.006	0.09	0.37	0.02	*	LCV DAPHNIDS
	15	4	0	50.00	200.00	34000	0.001	0.006	0.07	0.29	0.02		NAHQ ADVISORY
	15	4	0	50.00	200.00	34000	0.001	0.006	0.00	0.02	0.35		LTV EC20 FISH

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
4,4'-DDT (continued) ACENAPHTHENE	15	4	0	50.00	200.00	34000	0.001	0.006	0.00	0.01	0.73		LCV FISH
	1	6	0	680.00	870.00	390	1.744	2.231	0.79	1.01	2.20	<	NAHQ ADVISORY
	1	6	0	680.00	870.00	390	1.744	2.231	0.01	0.01	197.00		LTV EC20 FISH
	1	6	0	680.00	870.00	390	1.744	2.231	0.00	0.01	413.00		LCV FISH
	1	6	0	680.00	870.00	390	1.744	2.231	0.00	0.00	6646.00	*	LCV DAPHNIDS
	1	6	0	680.00	870.00	390	1.795	4.872	0.82	2.21	2.20		NAHQ ADVISORY
	2	8	0	700.00	1900.00	390	1.795	4.872	0.01	0.02	197.00	<	LTV EC20 FISH
	2	8	0	700.00	1900.00	390	1.795	4.872	0.00	0.01	413.00		LCV FISH
	2	8	0	700.00	1900.00	390	1.795	4.872	0.00	0.00	6646.00	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	390	0.487	1.051	0.22	0.48	2.20		NAHQ ADVISORY
	3	34	0	190.00	410.00	390	0.487	1.051	0.00	0.01	197.00	<	LTV EC20 FISH
	3	34	0	190.00	410.00	390	0.487	1.051	0.00	0.00	413.00		LCV FISH
	3	34	0	190.00	410.00	390	0.487	1.051	0.00	0.00	6646.00	*	LCV DAPHNIDS
	3	34	0	190.00	410.00	390	0.487	1.051	0.00	0.00	197.00	<	NAHQ ADVISORY
	4	6	0	660.00	720.00	390	1.692	1.846	0.01	0.01	197.00	<	LTV EC20 FISH
	4	6	0	660.00	720.00	390	1.692	1.846	0.00	0.00	413.00		LCV FISH
	4	6	0	660.00	720.00	390	1.692	1.846	0.00	0.00	6646.00	*	LCV DAPHNIDS
ALDRIN	15	3	0	670.00	710.00	390	1.718	1.821	0.78	0.83	2.20		NAHQ ADVISORY
	15	3	0	670.00	710.00	390	1.718	1.821	0.01	0.01	197.00	<	LTV EC20 FISH
	15	3	0	670.00	710.00	390	1.718	1.821	0.00	0.00	413.00		LCV FISH
	15	3	0	670.00	710.00	390	1.718	1.821	0.00	0.00	6646.00	*	LCV DAPHNIDS
	1	16	0	10.00	200.00	11000	0.001	0.018	0.01	0.24	0.08		NAHQ ADVISORY
	3	34	0	7.72	8.54	11000	0.001	0.001	0.01	0.01	0.08		NAHQ ADVISORY
	10	8	0	10.00	10.00	11000	0.001	0.001	0.01	0.01	0.08		NAHQ ADVISORY
	15	4	0	50.00	200.00	11000	0.005	0.018	0.06	0.24	0.08		NAHQ ADVISORY
	1	6	0	680.00	870.00	485	1.402	1.794	140206.19	179381.44	0.00		NAHQ ADVISORY
	1	6	0	680.00	870.00	485	1.402	1.794	15.58	19.93	0.09	*	LCV FISH
ANTHRACENE	1	6	0	680.00	870.00	485	1.402	1.794	4.01	5.13	0.35	*	LTV EC20 FISH
	1	6	0	680.00	870.00	485	1.402	1.794	0.67	0.85	2.10	<	LCV DAPHNIDS
	1	6	0	680.00	870.00	485	1.402	1.794	0.17	0.22	8.20	>	LTV EC20 DAPHNID
	1	6	0	700.00	1900.00	485	1.443	3.918	144329.90	391752.58	0.00		NAHQ ADVISORY
	2	8	0	700.00	1900.00	485	1.443	3.918	16.04	43.53	0.09	*	LCV FISH
	2	8	0	700.00	1900.00	485	1.443	3.918	4.12	11.19	0.35	*	LTV EC20 FISH
	2	8	0	700.00	1900.00	485	1.443	3.918	0.69	1.87	2.10	<	LCV DAPHNIDS
	2	8	0	700.00	1900.00	485	1.443	3.918	0.18	0.48	8.20	>	LTV EC20 DAPHNID
	2	8	0	700.00	1900.00	485	0.392	0.845	39175.26	84536.08	0.00		NAHQ ADVISORY
	3	34	0	190.00	410.00	485	0.392	0.845	4.35	9.39	0.09	*	LCV FISH
	3	34	0	190.00	410.00	485	0.392	0.845	1.12	2.42	0.35	*	LTV EC20 FISH
	3	34	0	190.00	410.00	485	0.392	0.845	0.19	0.40	2.10	<	LCV DAPHNIDS
	3	34	0	190.00	410.00	485	0.392	0.845	0.05	0.10	8.20	>	LTV EC20 DAPHNID
	3	34	0	190.00	410.00	485	0.392	0.845	15.12	16.49	0.00		NAHQ ADVISORY
	4	6	0	660.00	720.00	485	1.361	1.485	136082.47	148453.61	0.00	*	LTV EC20 FISH
	4	6	0	660.00	720.00	485	1.361	1.485	3.89	4.24	0.35	*	LCV FISH
	4	6	0	660.00	720.00	485	1.361	1.485	0.65	0.71	2.10	<	LTV EC20 FISH
	4	6	0	660.00	720.00	485	1.361	1.485	0.00	0.00	6646.00	*	LCV DAPHNIDS

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
ANTHRACENE (continued)	4	6	0	660.00	720.00	485	1,361	1,485	0.17	0.18	8.20	>	LTV EC20 DAPHNID
	15	3	0	670.00	710.00	485	1,381	1,464	138144.33	146391.75	0.00	*	NAHQ ADVISORY
	15	3	0	670.00	710.00	485	1,381	1,464	15.35	16.27	0.09	*	LCV FISH
	15	3	0	670.00	710.00	485	1,381	1,464	3.95	4.18	0.35	*	LTV EC20 FISH
	15	3	0	670.00	710.00	485	1,381	1,464	0.66	0.70	2.10	<	LCV DAPHNIDS
ANTIMONY	15	3	0	670.00	710.00	485	1,381	1,464	0.17	0.18	8.20	>	LTV EC20 DAPHNID
	1	28	0	280.00	580.00	1	280.000	580.000	15.56	32.22	18.00		NAHQ ADVISORY
	1	28	0	280.00	580.00	1	280.000	580.000	3.54	7.34	79.00		EC20 POPULATION
	1	28	0	280.00	580.00	1	280.000	580.000	0.17	0.36	1600.00		LCV FISH
	1	28	0	280.00	580.00	1	280.000	580.000	0.15	0.31	1900.00		LTV EC20 DAPHNID
	1	28	0	280.00	580.00	1	280.000	580.000	0.12	0.25	2310.00		LTV EC20 FISH
	1	28	0	280.00	580.00	1	280.000	580.000	0.05	0.11	5400.00		LCV DAPHNIDS
	2	13	0	460.00	600.00	1	460.000	600.000	25.56	33.33	18.00		NAHQ ADVISORY
	2	13	0	460.00	600.00	1	460.000	600.000	5.82	7.59	79.00		EC20 POPULATION
	2	13	0	460.00	600.00	1	460.000	600.000	0.29	0.38	1600.00		LCV FISH
	2	13	0	460.00	600.00	1	460.000	600.000	0.24	0.32	1900.00		LTV EC20 DAPHNID
	2	13	0	460.00	600.00	1	460.000	600.000	0.20	0.26	2310.00		LTV EC20 FISH
	2	13	0	460.00	600.00	1	460.000	600.000	0.09	0.11	5400.00		LCV DAPHNIDS
	3	35	0	300.00	590.00	1	300.000	590.000	16.67	32.78	18.00		NAHQ ADVISORY
	3	35	0	300.00	590.00	1	300.000	590.000	3.80	7.47	79.00		EC20 POPULATION
	3	35	0	300.00	590.00	1	300.000	590.000	0.19	0.37	1600.00		LCV FISH
	3	35	0	300.00	590.00	1	300.000	590.000	0.16	0.31	1900.00		LTV EC20 DAPHNID
	3	35	0	300.00	590.00	1	300.000	590.000	0.13	0.26	2310.00		LTV EC20 FISH
	3	35	0	300.00	590.00	1	300.000	590.000	0.06	0.11	5400.00		LCV DAPHNIDS
	4	27	0	480.00	610.00	1	480.000	610.000	26.67	33.89	18.00		NAHQ ADVISORY
	4	27	0	480.00	610.00	1	480.000	610.000	6.08	7.72	79.00		EC20 POPULATION
	4	27	0	480.00	610.00	1	480.000	610.000	0.30	0.38	1600.00		LCV FISH
	4	27	0	480.00	610.00	1	480.000	610.000	0.25	0.32	1900.00		LTV EC20 DAPHNID
	4	27	0	480.00	610.00	1	480.000	610.000	0.21	0.26	2310.00		LTV EC20 FISH
	4	27	0	480.00	610.00	1	480.000	610.000	0.09	0.11	5400.00		LCV DAPHNIDS
	5	25	0	460.00	630.00	1	460.000	630.000	25.56	35.00	18.00		NAHQ ADVISORY
	5	25	0	460.00	630.00	1	460.000	630.000	5.82	7.97	79.00		EC20 POPULATION
	5	25	0	460.00	630.00	1	460.000	630.000	0.29	0.39	1600.00		LCV FISH
	5	25	0	460.00	630.00	1	460.000	630.000	0.24	0.33	1900.00		LTV EC20 DAPHNID
	5	25	0	460.00	630.00	1	460.000	630.000	0.20	0.27	2310.00		LTV EC20 FISH
	5	25	0	460.00	630.00	1	460.000	630.000	0.09	0.12	5400.00		LCV DAPHNIDS
	10	15	0	350.00	610.00	1	350.000	610.000	19.44	33.89	18.00		NAHQ ADVISORY
	10	15	0	350.00	610.00	1	350.000	610.000	4.43	7.72	79.00		EC20 POPULATION
	10	15	0	350.00	610.00	1	350.000	610.000	0.22	0.38	1600.00		LCV FISH
	10	15	0	350.00	610.00	1	350.000	610.000	0.18	0.32	1900.00		LTV EC20 DAPHNID
	10	15	0	350.00	610.00	1	350.000	610.000	0.15	0.26	2310.00		LTV EC20 FISH
	10	15	0	350.00	610.00	1	350.000	610.000	0.06	0.11	5400.00		LCV DAPHNIDS
	13	10	0	460.00	570.00	1	460.000	570.000	25.56	31.67	18.00		NAHQ ADVISORY
	13	10	0	460.00	570.00	1	460.000	570.000	5.82	7.22	79.00		EC20 POPULATION
	13	10	0	460.00	570.00	1	460.000	570.000	0.29	0.36	1600.00		LCV FISH
	13	10	0	460.00	570.00	1	460.000	570.000	0.24	0.30	1900.00		LTV EC20 DAPHNID

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
ANTIMONY (continued)	13	10	0	440.00	570.00	1	460.000	570.000	0.20	0.25	2310.00		LTV EC20 FISH
	13	10	0	460.00	570.00	1	460.000	570.000	0.09	0.11	5400.00		LCV DAPHNIDS
	15	16	0	350.00	570.00	1	350.000	570.000	19.44	31.67	18.00		NAHQ ADVISORY
	15	16	0	350.00	570.00	1	350.000	570.000	4.43	7.22	79.00		EC20 POPULATION
	15	16	0	350.00	570.00	1	350.000	570.000	0.22	0.36	1600.00		LCV FISH
	15	16	0	350.00	570.00	1	350.000	570.000	0.18	0.30	1900.00		LTV EC20 DAPHNID
	15	16	0	350.00	570.00	1	350.000	570.000	0.15	0.25	2310.00		LTV EC20 FISH
	15	16	0	350.00	570.00	1	350.000	570.000	0.06	0.11	5400.00		LCV DAPHNIDS
	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	0.02		NAHQ ADVISORY
	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	0.52		LTV EC20 FISH
AROCLOX-1254	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	0.63		EC20 POPULATION
	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	1.00		LCV FISH
	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	1.20		LTV EC20 DAPHNID
	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	2.10		LCV DAPHNIDS
	10	8	0	10.00	50.00	10000000	0.000	0.000	0.00	0.00	0.00		NAHQ ADVISORY
	1	6	0	680.00	870.00	10109	0.067	0.086	24.91	31.87	0.00	*	LCV DAPHNIDS
	1	6	0	680.00	870.00	10109	0.067	0.086	0.10	0.13	0.65		NAHQ ADVISORY
	2	8	0	700.00	1900.00	10109	0.069	0.188	25.65	69.61	0.00	*	LCV DAPHNIDS
	3	34	0	190.00	410.00	10109	0.019	0.041	6.96	15.02	0.00	*	NAHQ ADVISORY
	3	34	0	190.00	410.00	10109	0.019	0.041	0.03	0.06	0.00	*	LCV DAPHNIDS
BENZO(A)ANTHRACENE	4	6	0	660.00	720.00	10109	0.065	0.071	24.18	26.38	0.00	*	NAHQ ADVISORY
	4	6	0	660.00	720.00	10109	0.065	0.071	0.10	0.11	0.65	*	LCV DAPHNIDS
	15	3	0	670.00	710.00	10109	0.066	0.070	24.55	26.01	0.00	*	NAHQ ADVISORY
	15	3	0	670.00	710.00	10109	0.066	0.070	0.10	0.11	0.65	*	LCV DAPHNIDS
	1	6	0	680.00	870.00	30	22.667	29.000	17435.90	22307.69	0.00		NAHQ ADVISORY
	1	6	0	680.00	870.00	30	22.667	29.000	75.56	96.67	0.30	*	LCV DAPHNIDS
	1	6	0	680.00	870.00	30	22.667	29.000	7.58	9.70	2.99	>	LTV EC20 FISH
	2	8	0	700.00	1900.00	30	23.333	63.333	17948.72	48717.95	0.00	*	NAHQ ADVISORY
	2	8	0	700.00	1900.00	30	23.333	63.333	77.78	211.11	0.30	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	30	23.333	63.333	7.80	21.18	2.99	>	LTV EC20 FISH
BENZO(A)PYRENE	3	33	0	190.00	410.00	30	6.333	13.667	4871.79	10512.82	0.00	*	NAHQ ADVISORY
	3	33	0	190.00	410.00	30	6.333	13.667	21.11	45.56	0.30	*	LCV DAPHNIDS
	3	33	0	190.00	410.00	30	6.333	13.667	2.12	4.57	2.99	>	LTV EC20 FISH
	4	6	0	660.00	720.00	30	22.000	24.000	16923.08	18461.54	0.00	*	NAHQ ADVISORY
	4	6	0	660.00	720.00	30	22.000	24.000	73.33	80.00	0.30	*	LCV DAPHNIDS
	4	6	0	660.00	720.00	30	22.000	24.000	7.36	8.03	2.99	>	LTV EC20 FISH
	15	3	0	670.00	710.00	30	22.333	23.667	17179.49	18205.13	0.00	*	NAHQ ADVISORY
	15	3	0	670.00	710.00	30	22.333	23.667	74.44	78.89	0.30	*	LCV DAPHNIDS
	15	3	0	670.00	710.00	30	22.333	23.667	7.47	7.92	2.99	>	LTV EC20 FISH
	15	3	0	670.00	710.00	30	22.333	23.667	0.10	0.11	0.65	*	LCV DAPHNIDS

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
BENZOIC ACID	1	6	0	3400.00	4300.00	.	.	.	.	.	0.45	.	NAWQ ADVISORY
	1	6	0	3400.00	4300.00	.	.	.	.	.	1259.00	*	EC20 POPULATION
	1	6	0	3400.00	4300.00	.	.	.	.	.	7409.00	*	LTV EC20 FISH
	1	6	0	3400.00	4300.00	.	.	.	.	.	12976.00	*	LCV FISH
	2	8	0	3500.00	9300.00	.	.	.	.	.	0.45	.	NAWQ ADVISORY
	2	8	0	3500.00	9300.00	.	.	.	.	.	1259.00	*	EC20 POPULATION
	2	8	0	3500.00	9300.00	.	.	.	.	.	7409.00	*	LTV EC20 FISH
	2	8	0	3500.00	9300.00	.	.	.	.	.	12976.00	*	LCV FISH
	3	34	0	930.00	2100.00	.	.	.	.	.	0.45	.	NAWQ ADVISORY
	3	34	0	930.00	2100.00	.	.	.	.	.	1259.00	*	EC20 POPULATION
	3	34	0	930.00	2100.00	.	.	.	.	.	7409.00	*	LTV EC20 FISH
	3	34	0	930.00	2100.00	.	.	.	.	.	12976.00	*	LCV FISH
	4	2	0	3300.00	3600.00	.	.	.	.	.	0.45	.	NAWQ ADVISORY
	4	2	0	3300.00	3600.00	.	.	.	.	.	1259.00	*	EC20 POPULATION
	4	2	0	3300.00	3600.00	.	.	.	.	.	7409.00	*	LTV EC20 FISH
	4	2	0	3300.00	3600.00	.	.	.	.	.	12976.00	*	LCV FISH
BENZYL ALCOHOL	1	6	0	680.00	870.00	.	.	.	.	.	0.15	.	NAWQ ADVISORY
	1	6	0	680.00	870.00	.	.	.	.	.	375.00	*	EC20 POPULATION
	1	6	0	680.00	870.00	.	.	.	.	.	550.00	*	LTV EC20 FISH
	1	6	0	680.00	870.00	.	.	.	.	.	589.00	*	LCV FISH
	2	8	0	700.00	1900.00	.	.	.	.	.	0.15	.	NAWQ ADVISORY
	2	8	0	700.00	1900.00	.	.	.	.	.	375.00	*	EC20 POPULATION
	2	8	0	700.00	1900.00	.	.	.	.	.	550.00	*	LTV EC20 FISH
	2	8	0	700.00	1900.00	.	.	.	.	.	589.00	*	LCV FISH
	3	34	0	190.00	410.00	.	.	.	.	.	0.15	.	NAWQ ADVISORY
	3	34	0	190.00	410.00	.	.	.	.	.	375.00	*	EC20 POPULATION
	3	34	0	190.00	410.00	.	.	.	.	.	550.00	*	LTV EC20 FISH
	3	34	0	190.00	410.00	.	.	.	.	.	589.00	*	LCV FISH
	4	6	0	660.00	720.00	.	.	.	.	.	0.15	.	NAWQ ADVISORY
	4	6	0	660.00	720.00	.	.	.	.	.	375.00	*	EC20 POPULATION
	4	6	0	660.00	720.00	.	.	.	.	.	550.00	*	LTV EC20 FISH
	4	6	0	660.00	720.00	.	.	.	.	.	589.00	*	LCV FISH
	15	3	0	670.00	710.00	.	.	.	.	.	0.15	.	NAWQ ADVISORY
	15	3	0	670.00	710.00	.	.	.	.	.	375.00	*	EC20 POPULATION
	15	3	0	670.00	710.00	.	.	.	.	.	550.00	*	LTV EC20 FISH
	15	3	0	670.00	710.00	.	.	.	.	.	589.00	*	LCV FISH

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
BERYLLIUM	2	13	0	3.00	3.00	2	1.500	1.500	2.46	2.46	0.61		NAHQ ADVISORY
	2	13	0	3.00	3.00	2	1.500	1.500	0.39	0.39	3.80		LTV EC20 DAPHNID
	2	13	0	3.00	3.00	2	1.500	1.500	0.28	0.28	5.30		LTV DAPHNIDS
	2	13	0	3.00	3.00	2	1.500	1.500	0.07	0.07	21.00	*	EC20 POPULATION
	2	13	0	3.00	3.00	2	1.500	1.500	0.03	0.03	57.00	*	LTV FISH
	2	13	0	3.00	3.00	2	1.500	1.500	0.01	0.01	148.00		LTV EC20 FISH
	2	13	0	3.00	3.00	2	1.500	1.500	2.46	2.46	0.61		NAHQ ADVISORY
	4	27	0	3.00	3.00	2	1.500	1.500	0.39	0.39	3.80		LTV EC20 DAPHNID
	4	27	0	3.00	3.00	2	1.500	1.500	0.28	0.28	5.30		LTV DAPHNIDS
	4	27	0	3.00	3.00	2	1.500	1.500	0.07	0.07	21.00	*	EC20 POPULATION
	4	27	0	3.00	3.00	2	1.500	1.500	0.03	0.03	57.00	*	LTV FISH
	4	27	0	3.00	3.00	2	1.500	1.500	0.01	0.01	148.00		LTV EC20 FISH
	13	3	0	3.00	3.00	2	1.500	1.500	2.46	2.46	0.61		NAHQ ADVISORY
	13	3	0	3.00	3.00	2	1.500	1.500	0.39	0.39	3.80		LTV EC20 DAPHNID
	13	3	0	3.00	3.00	2	1.500	1.500	0.28	0.28	5.30		LTV DAPHNIDS
	13	3	0	3.00	3.00	2	1.500	1.500	0.07	0.07	21.00	*	EC20 POPULATION
	13	3	0	3.00	3.00	2	1.500	1.500	0.03	0.03	57.00	*	LTV FISH
	13	3	0	3.00	3.00	2	1.500	1.500	0.01	0.01	148.00		LTV EC20 FISH
BIS(2-ETHYLHEXYL)PHTHALATE	1	6	0	680.00	870.00	310	2.194	2.806	219.35	280.65	0.01		NAHQ ADVISORY
	1	6	0	680.00	870.00	310	2.194	2.806	0.73	0.94	3.00	<	LTV DAPHNIDS
	1	6	0	680.00	870.00	310	2.194	2.806	0.73	0.94	3.00	<	LTV EC20 DAPHNID
	1	6	0	680.00	870.00	310	2.194	2.806	0.26	0.33	8.40		LTV FISH
	1	6	0	680.00	870.00	310	2.194	2.806	0.04	0.06	50.00		EC20 POPULATION
	1	6	0	680.00	870.00	310	2.194	2.806	0.04	0.05	54.00	>	LTV EC20 FISH
	1	6	0	700.00	1900.00	310	2.258	6.129	225.81	612.90	0.01		NAHQ ADVISORY
	2	8	0	700.00	1900.00	310	2.258	6.129	0.75	2.04	3.00	<	LTV DAPHNIDS
	2	8	0	700.00	1900.00	310	2.258	6.129	0.75	2.04	3.00	<	LTV EC20 DAPHNID
	2	8	0	700.00	1900.00	310	2.258	6.129	0.27	0.73	8.40		LTV FISH
	2	8	0	700.00	1900.00	310	2.258	6.129	0.05	0.12	50.00		EC20 POPULATION
	2	8	0	700.00	1900.00	310	2.258	6.129	0.04	0.11	54.00	>	LTV EC20 FISH
	4	6	0	660.00	720.00	310	2.129	2.323	212.90	232.26	0.01		NAHQ ADVISORY
	4	6	0	660.00	720.00	310	2.129	2.323	0.71	0.77	3.00	<	LTV DAPHNIDS
	4	6	0	660.00	720.00	310	2.129	2.323	0.71	0.77	3.00	<	LTV EC20 DAPHNID
	4	6	0	660.00	720.00	310	2.129	2.323	0.25	0.28	8.40		LTV FISH
	4	6	0	660.00	720.00	310	2.129	2.323	0.04	0.05	50.00		EC20 POPULATION
	4	6	0	660.00	720.00	310	2.129	2.323	0.04	0.04	54.00	>	LTV EC20 FISH
	4	6	0	660.00	720.00	310	2.129	2.323	0.04	0.04	54.00		LTV DAPHNIDS
	15	3	0	670.00	710.00	310	2.161	2.290	216.13	229.03	0.01		NAHQ ADVISORY
	15	3	0	670.00	710.00	310	2.161	2.290	0.72	0.76	3.00	<	LTV DAPHNIDS
	15	3	0	670.00	710.00	310	2.161	2.290	0.72	0.76	3.00	<	LTV EC20 DAPHNID
	15	3	0	670.00	710.00	310	2.161	2.290	0.26	0.27	8.40		LTV FISH
	15	3	0	670.00	710.00	310	2.161	2.290	0.04	0.05	50.00		EC20 POPULATION
	15	3	0	670.00	710.00	310	2.161	2.290	0.04	0.04	54.00	>	LTV EC20 FISH

Table F.6 (continued)

Analyte	Reach (fish)	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark Flag	Benchmark type
Cadmium	1	28	0	110.00	230.00	50	2.200	4.600	169.23	353.85	0.01	SS EC20
	1	28	0	110.00	230.00	50	2.200	4.600	14.67	30.67	0.15	LCV DAPHNIDS
	1	28	0	110.00	230.00	50	2.200	4.600	2.93	6.13	0.75	LTV EC20 DAPHNID
	1	28	0	110.00	230.00	50	2.200	4.600	2.00	4.18	1.10	MAWQ ACTUAL
	1	28	0	110.00	230.00	50	2.200	4.600	1.29	2.71	1.70	LCV FISH
	1	28	0	110.00	230.00	50	2.200	4.600	1.22	2.56	1.80	LTV EC20 FISH
	1	28	0	110.00	230.00	50	2.200	4.600	0.51	1.07	4.30	EC20 POPULATION
	2	28	0	110.00	230.00	50	2.200	4.600	276.92	369.23	0.01	SS EC20
	2	13	0	180.00	240.00	50	3.600	4.800	24.00	32.00	0.15	LCV DAPHNIDS
	2	13	0	180.00	240.00	50	3.600	4.800	4.80	6.40	0.75	LTV EC20 DAPHNID
	2	13	0	180.00	240.00	50	3.600	4.800	3.27	4.36	1.10	MAWQ ACTUAL
	2	13	0	180.00	240.00	50	3.600	4.800	2.12	2.82	1.70	LCV FISH
	2	13	0	180.00	240.00	50	3.600	4.800	2.00	2.67	1.80	LTV EC20 FISH
	2	13	0	180.00	240.00	50	3.600	4.800	0.84	1.12	4.30	EC20 POPULATION
	3	35	0	120.00	240.00	50	2.400	4.800	184.62	369.23	0.01	SS EC20
	3	35	0	120.00	240.00	50	2.400	4.800	16.00	32.00	0.15	LCV DAPHNIDS
	3	35	0	120.00	240.00	50	2.400	4.800	3.20	6.40	0.75	LTV EC20 DAPHNID
	3	35	0	120.00	240.00	50	2.400	4.800	2.18	4.36	1.10	MAWQ ACTUAL
	3	35	0	120.00	240.00	50	2.400	4.800	1.41	2.82	1.70	LCV FISH
	3	35	0	120.00	240.00	50	2.400	4.800	1.33	2.67	1.80	LTV EC20 FISH
	3	35	0	120.00	240.00	50	2.400	4.800	0.56	1.12	4.30	EC20 POPULATION
	4	27	0	190.00	240.00	50	3.800	4.800	292.31	369.23	0.01	SS EC20
	4	27	0	190.00	240.00	50	3.800	4.800	25.33	32.00	0.15	LCV DAPHNIDS
	4	27	0	190.00	240.00	50	3.800	4.800	5.07	6.40	0.75	LTV EC20 DAPHNID
	4	27	0	190.00	240.00	50	3.800	4.800	3.45	4.36	1.10	MAWQ ACTUAL
	4	27	0	190.00	240.00	50	3.800	4.800	2.24	2.82	1.70	LCV FISH
	4	27	0	190.00	240.00	50	3.800	4.800	2.11	2.67	1.80	LTV EC20 FISH
	4	27	0	190.00	240.00	50	3.800	4.800	0.88	1.12	4.30	EC20 POPULATION
	5	25	0	190.00	280.00	50	3.800	5.600	292.31	430.77	0.01	SS EC20
	5	25	0	190.00	280.00	50	3.800	5.600	25.33	37.33	0.15	LCV DAPHNIDS
	5	25	0	190.00	280.00	50	3.800	5.600	5.07	7.47	0.75	LTV EC20 DAPHNID
	5	25	0	190.00	280.00	50	3.800	5.600	3.45	5.09	1.10	MAWQ ACTUAL
	5	25	0	190.00	280.00	50	3.800	5.600	2.24	3.29	1.70	LCV FISH
	5	25	0	190.00	280.00	50	3.800	5.600	2.11	3.11	1.80	LTV EC20 FISH
	5	25	0	190.00	280.00	50	3.800	5.600	0.88	1.30	4.30	EC20 POPULATION
	10	15	0	140.00	250.00	50	2.800	5.000	215.38	384.62	0.01	SS EC20
	10	15	0	140.00	250.00	50	2.800	5.000	18.67	33.33	0.15	LCV DAPHNIDS
	10	15	0	140.00	250.00	50	2.800	5.000	3.73	6.67	0.75	LTV EC20 DAPHNID
	10	15	0	140.00	250.00	50	2.800	5.000	2.55	4.55	1.10	MAWQ ACTUAL
	10	15	0	140.00	250.00	50	2.800	5.000	1.65	2.94	1.70	LCV FISH
	10	15	0	140.00	250.00	50	2.800	5.000	1.56	2.78	1.80	LTV EC20 FISH
	10	15	0	140.00	250.00	50	2.800	5.000	0.65	1.16	4.30	EC20 POPULATION
	13	10	0	180.00	240.00	50	3.600	4.800	276.92	369.23	0.01	SS EC20
	13	10	0	180.00	240.00	50	3.600	4.800	24.00	32.00	0.15	LCV DAPHNIDS
	13	10	0	180.00	240.00	50	3.600	4.800	4.80	6.40	0.75	LTV EC20 DAPHNID
	13	10	0	180.00	240.00	50	3.600	4.800	3.27	4.36	1.10	MAWQ ACTUAL
	13	10	0	180.00	240.00	50	3.600	4.800	2.12	2.82	1.70	LCV FISH

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
CADMIUM (continued)	13	10	0	180.00	240.00	50	3.600	4.800	2.00	2.67	1.80		LTV EC20 FISH
	13	10	0	180.00	240.00	50	3.600	4.800	0.84	1.12	4.30		EC20 POPULATION
	15	16	0	140.00	230.00	50	2.800	4.600	215.38	353.85	0.01		SS EC20
	15	16	0	140.00	230.00	50	2.800	4.600	18.67	30.67	0.15		LCV DAPHNIDS
	15	16	0	140.00	230.00	50	2.800	4.600	3.73	6.13	0.75		LTV EC20 DAPHNID
	15	16	0	140.00	230.00	50	2.800	4.600	2.55	4.18	1.10		NAHQ ACTUAL
	15	16	0	140.00	230.00	50	2.800	4.600	1.65	2.71	1.70		LCV FISH
	15	16	0	140.00	230.00	50	2.800	4.600	1.56	2.56	1.80		LTV EC20 FISH
	15	16	0	140.00	230.00	50	2.800	4.600	0.65	1.07	4.30		EC20 POPULATION
	1	28	0	280.00	580.00	200	1.400	2.900	5.26	10.90	0.27		SS EC20
	1	28	0	280.00	580.00	200	1.400	2.900	2.80	5.80	0.50		LTV EC20 DAPHNID
	1	28	0	280.00	580.00	200	1.400	2.900	0.23	0.47	6.13		LCV DAPHNIDS
	1	28	0	280.00	580.00	200	1.400	2.900	0.13	0.26	11.00		NAHQ ACTUAL
	1	28	0	280.00	580.00	200	1.400	2.900	0.03	0.06	51.00		LTV EC20 FISH
	1	28	0	280.00	580.00	200	1.400	2.900	0.02	0.04	73.18		LCV FISH
CHROMIUM	1	28	0	280.00	580.00	200	1.400	2.900	0.00	0.01	316.00		EC20 POPULATION
	2	13	0	460.00	600.00	200	2.300	3.000	8.65	11.28	0.27		SS EC20
	2	13	0	460.00	600.00	200	2.300	3.000	4.60	6.00	0.50		LTV EC20 DAPHNID
	2	13	0	460.00	600.00	200	2.300	3.000	0.38	0.49	6.13		LCV DAPHNIDS
	2	13	0	460.00	600.00	200	2.300	3.000	0.21	0.27	11.00		NAHQ ACTUAL
	2	13	0	460.00	600.00	200	2.300	3.000	0.05	0.06	51.00		LTV EC20 FISH
	2	13	0	460.00	600.00	200	2.300	3.000	0.03	0.04	73.18		LCV FISH
	2	13	0	460.00	600.00	200	2.300	3.000	0.01	0.01	316.00		EC20 POPULATION
	3	35	0	300.00	590.00	200	1.500	2.950	5.64	11.09	0.27		SS EC20
	3	35	0	300.00	590.00	200	1.500	2.950	3.00	5.90	0.50		LTV EC20 DAPHNID
	3	35	0	300.00	590.00	200	1.500	2.950	0.24	0.48	6.13		LCV DAPHNIDS
	3	35	0	300.00	590.00	200	1.500	2.950	0.14	0.27	11.00		NAHQ ACTUAL
	3	35	0	300.00	590.00	200	1.500	2.950	0.03	0.06	51.00		LTV EC20 FISH
	3	35	0	300.00	590.00	200	1.500	2.950	0.02	0.04	73.18		LCV FISH
	3	35	0	300.00	590.00	200	1.500	2.950	0.00	0.01	316.00		EC20 POPULATION
	4	27	0	480.00	610.00	200	2.400	3.050	9.02	11.47	0.27		SS EC20
	4	27	0	480.00	610.00	200	2.400	3.050	4.80	6.10	0.50		LTV EC20 DAPHNID
	4	27	0	480.00	610.00	200	2.400	3.050	0.39	0.50	6.13		LCV DAPHNIDS
	4	27	0	480.00	610.00	200	2.400	3.050	0.22	0.28	11.00		NAHQ ACTUAL
	4	27	0	480.00	610.00	200	2.400	3.050	0.05	0.06	51.00		LTV EC20 FISH
	4	27	0	480.00	610.00	200	2.400	3.050	0.03	0.04	73.18		LCV FISH
	4	27	0	480.00	610.00	200	2.400	3.050	0.01	0.01	316.00		EC20 POPULATION
	5	25	0	460.00	630.00	200	2.300	3.150	8.65	11.84	0.27		SS EC20
	5	25	0	460.00	630.00	200	2.300	3.150	4.60	6.30	0.50		LTV EC20 DAPHNID
	5	25	0	460.00	630.00	200	2.300	3.150	0.38	0.51	6.13		LCV DAPHNIDS
	5	25	0	460.00	630.00	200	2.300	3.150	0.21	0.29	11.00		NAHQ ACTUAL
	5	25	0	460.00	630.00	200	2.300	3.150	0.05	0.06	51.00		LTV EC20 FISH
	5	25	0	460.00	630.00	200	2.300	3.150	0.03	0.04	73.18		LCV FISH
	5	25	0	460.00	630.00	200	2.300	3.150	0.01	0.01	316.00		EC20 POPULATION
	10	15	0	350.00	610.00	200	1.750	3.050	6.58	11.47	0.27		SS EC20
	10	15	0	350.00	610.00	200	1.750	3.050	3.50	6.10	0.50		LTV EC20 DAPHNID

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
CHROMIUM (continued)	10	15	0	350.00	610.00	200	1.750	3.050	0.29	0.50	6.13		LCV DAPHNIDS
	10	15	0	350.00	610.00	200	1.750	3.050	0.16	0.28	11.00		NAHQ ACTUAL
	10	15	0	350.00	610.00	200	1.750	3.050	0.03	0.06	51.00		LTV EC20 FISH
	10	15	0	350.00	610.00	200	1.750	3.050	0.02	0.04	73.18		LCV FISH
	10	15	0	350.00	610.00	200	1.750	3.050	0.01	0.01	316.00		EC20 POPULATION
	13	10	0	450.00	590.00	200	2.250	2.950	8.46	11.09	0.27		SS EC20
	13	10	0	450.00	590.00	200	2.250	2.950	4.50	5.90	0.50		LTV EC20 DAPHNID
	13	10	0	450.00	590.00	200	2.250	2.950	0.37	0.48	6.13		LCV DAPHNIDS
	13	10	0	450.00	590.00	200	2.250	2.950	0.20	0.27	11.00		NAHQ ACTUAL
	13	10	0	450.00	590.00	200	2.250	2.950	0.04	0.06	51.00		LTV EC20 FISH
	13	10	0	450.00	590.00	200	2.250	2.950	0.03	0.04	73.18		LCV FISH
	13	10	0	450.00	590.00	200	2.250	2.950	0.01	0.01	316.00		EC20 POPULATION
	15	16	0	350.00	570.00	200	1.750	2.850	6.58	10.71	0.27		SS EC20
	15	16	0	350.00	570.00	200	1.750	2.850	3.50	5.70	0.50		LTV EC20 DAPHNID
	15	16	0	350.00	570.00	200	1.750	2.850	0.29	0.46	6.13		LCV DAPHNIDS
	15	16	0	350.00	570.00	200	1.750	2.850	0.16	0.26	11.00		NAHQ ACTUAL
	15	16	0	350.00	570.00	200	1.750	2.850	0.03	0.06	51.00		LTV EC20 FISH
	15	16	0	350.00	570.00	200	1.750	2.850	0.02	0.04	73.18		LCV FISH
	15	16	0	350.00	570.00	200	1.750	2.850	0.01	0.01	316.00		EC20 POPULATION
COPPER	1	28	0	280.00	580.00	210	1.333	2.762	6.50	13.47	0.20		LTV EC20 DAPHNID
	1	28	0	280.00	580.00	210	1.333	2.762	5.80	12.01	0.23		LCV DAPHNIDS
	1	28	0	280.00	580.00	210	1.333	2.762	5.13	10.62	0.26		SS EC20
	1	28	0	280.00	580.00	210	1.333	2.762	0.34	0.71	3.87		LCV FISH
	1	28	0	280.00	580.00	210	1.333	2.762	0.27	0.55	5.00		LTV EC20 FISH
	1	28	0	280.00	580.00	210	1.333	2.762	0.16	0.32	8.60		EC20 POPULATION
	1	28	0	280.00	580.00	210	1.333	2.762	0.11	0.23	12.00		NAHQ ACTUAL
	2	13	0	460.00	600.00	210	2.190	2.857	10.69	13.94	0.20		LTV EC20 DAPHNID
	2	13	0	460.00	600.00	210	2.190	2.857	9.52	12.42	0.23		LCV DAPHNIDS
	2	13	0	460.00	600.00	210	2.190	2.857	8.42	10.99	0.26		SS EC20
	2	13	0	460.00	600.00	210	2.190	2.857	0.57	0.74	3.87		LCV FISH
	2	13	0	460.00	600.00	210	2.190	2.857	0.44	0.57	5.00		LTV EC20 FISH
	2	13	0	460.00	600.00	210	2.190	2.857	0.25	0.33	8.60		EC20 POPULATION
	2	13	0	460.00	600.00	210	2.190	2.857	0.18	0.24	12.00		NAHQ ACTUAL
	3	35	0	300.00	590.00	210	1.429	2.810	6.97	13.70	0.20		LTV EC20 DAPHNID
	3	35	0	300.00	590.00	210	1.429	2.810	6.21	12.22	0.23		LCV DAPHNIDS
	3	35	0	300.00	590.00	210	1.429	2.810	5.49	10.81	0.26		SS EC20
	3	35	0	300.00	590.00	210	1.429	2.810	0.37	0.73	3.87		LCV FISH
	3	35	0	300.00	590.00	210	1.429	2.810	0.29	0.56	5.00		LTV EC20 FISH
	3	35	0	300.00	590.00	210	1.429	2.810	0.17	0.33	8.60		EC20 POPULATION
	3	35	0	480.00	610.00	210	2.286	2.905	11.15	14.17	0.20		LTV EC20 DAPHNID
	4	27	0	480.00	610.00	210	2.286	2.905	9.94	12.63	0.23		NAHQ ACTUAL
	4	27	0	480.00	610.00	210	2.286	2.905	8.79	11.17	0.26		LTV EC20 FISH
	4	27	0	480.00	610.00	210	2.286	2.905	0.59	11.17	0.26		SS EC20
	4	27	0	480.00	610.00	210	2.286	2.905	0.46	0.58	3.87		LCV FISH
	4	27	0	480.00	610.00	210	2.286	2.905	0.27	0.34	5.00		LTV EC20 FISH
	4	27	0	480.00	610.00	210	2.286	2.905	0.27	0.34	8.60		EC20 POPULATION

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
COPPER (continued)	4	27	0	480.00	610.00	210	2.286	2.905	0.19	0.24	12.00		NAHQ ACTUAL
	10	15	0	350.00	610.00	210	1.667	2.905	8.13	14.17	0.20		LTV EC20 DAPHNID
	10	15	0	350.00	610.00	210	1.667	2.905	7.25	12.63	0.23		LCV DAPHNIDS
	10	15	0	350.00	610.00	210	1.667	2.905	6.41	11.17	0.26		SS EC20
	10	15	0	350.00	610.00	210	1.667	2.905	0.43	0.75	3.87		LCV FISH
	10	15	0	350.00	610.00	210	1.667	2.905	0.33	0.58	5.00		LTV EC20 FISH
	10	15	0	350.00	610.00	210	1.667	2.905	0.19	0.34	8.60		EC20 POPULATION
	10	15	0	350.00	610.00	210	1.667	2.905	0.14	0.24	12.00		NAHQ ACTUAL
	13	10	0	450.00	590.00	210	2.143	2.810	10.45	13.70	0.20		LTV EC20 DAPHNID
	13	10	0	450.00	590.00	210	2.143	2.810	9.32	12.22	0.23		LCV DAPHNIDS
	13	10	0	450.00	590.00	210	2.143	2.810	8.24	10.81	0.26		SS EC20
	13	10	0	450.00	590.00	210	2.143	2.810	0.55	0.73	3.87		LCV FISH
	13	10	0	450.00	590.00	210	2.143	2.810	0.25	0.33	5.00		LTV EC20 FISH
	13	10	0	450.00	590.00	210	2.143	2.810	0.18	0.23	8.60		EC20 POPULATION
	15	16	0	350.00	570.00	210	1.667	2.714	8.13	13.24	0.20		NAHQ ACTUAL
	15	16	0	350.00	570.00	210	1.667	2.714	7.25	11.80	0.23		LCV DAPHNIDS
	15	16	0	350.00	570.00	210	1.667	2.714	6.41	10.44	0.26		SS EC20
	15	16	0	350.00	570.00	210	1.667	2.714	0.43	0.70	3.87		LCV FISH
	15	16	0	350.00	570.00	210	1.667	2.714	0.33	0.54	5.00		LTV EC20 FISH
	15	16	0	350.00	570.00	210	1.667	2.714	0.19	0.32	8.60		EC20 POPULATION
	15	16	0	350.00	570.00	210	1.667	2.714	0.14	0.23	12.00		NAHQ ACTUAL
DI-N-BUTYLPHTHALATE	1	6	0	680.00	870.00	89	7.640	9.775	1.74	2.22	4.40		NAHQ ADVISORY
	1	6	0	680.00	870.00	89	7.640	9.775	0.03	0.04	251.00		EC20 POPULATION
	1	6	0	680.00	870.00	89	7.640	9.775	0.03	0.04	270.00		LTV EC20 FISH
	1	6	0	680.00	870.00	89	7.640	9.775	0.02	0.02	500.00		LTV EC20 DAPHNID
	1	6	0	680.00	870.00	89	7.640	9.775	0.01	0.01	717.00		LCV FISH
	1	6	0	680.00	870.00	89	7.640	9.775	0.01	0.01	717.00		LCV DAPHNIDS
	2	8	0	700.00	1900.00	89	7.865	21.348	1.79	4.85	4.40		NAHQ ADVISORY
	2	8	0	700.00	1900.00	89	7.865	21.348	0.03	0.09	251.00		EC20 POPULATION
	2	8	0	700.00	1900.00	89	7.865	21.348	0.03	0.08	270.00		LTV EC20 FISH
	2	8	0	700.00	1900.00	89	7.865	21.348	0.02	0.04	500.00		LTV EC20 DAPHNID
	2	8	0	700.00	1900.00	89	7.865	21.348	0.01	0.03	717.00		LCV FISH
	2	8	0	700.00	1900.00	89	7.865	21.348	0.01	0.03	717.00		LCV DAPHNIDS
	3	34	0	190.00	410.00	89	2.135	4.607	0.49	1.05	4.40		NAHQ ADVISORY
	3	34	0	190.00	410.00	89	2.135	4.607	0.01	0.02	251.00		EC20 POPULATION
	3	34	0	190.00	410.00	89	2.135	4.607	0.01	0.02	270.00		LTV EC20 FISH
	3	34	0	190.00	410.00	89	2.135	4.607	0.00	0.01	500.00		LTV EC20 DAPHNID
	3	34	0	190.00	410.00	89	2.135	4.607	0.00	0.01	717.00		LCV FISH
	3	34	0	190.00	410.00	89	2.135	4.607	0.00	0.01	717.00		LCV DAPHNIDS
	4	6	0	660.00	720.00	89	7.416	8.090	1.69	1.84	4.40		NAHQ ADVISORY
	4	6	0	660.00	720.00	89	7.416	8.090	0.03	0.03	251.00		EC20 POPULATION
	4	6	0	660.00	720.00	89	7.416	8.090	0.03	0.03	270.00		LTV EC20 FISH
	4	6	0	660.00	720.00	89	7.416	8.090	0.01	0.02	500.00		LTV EC20 DAPHNID
	4	6	0	660.00	720.00	89	7.416	8.090	0.01	0.01	717.00		LCV FISH
	4	6	0	660.00	720.00	89	7.416	8.090	0.01	0.01	717.00		LCV DAPHNIDS

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	SCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
DI-N-BUTYLPHTHALATE (continued)	15	3	0	670.00	710.00	89	7.528	7.978	1.71	1.81	4.40		NAHQ ADVISORY
	15	3	0	670.00	710.00	89	7.528	7.978	0.03	0.03	251.00		EC20 POPULATION
	15	3	0	670.00	710.00	89	7.528	7.978	0.03	0.03	270.00		LTV EC20 FISH
	15	3	0	670.00	710.00	89	7.528	7.978	0.02	0.02	500.00		LTV EC20 DAPHNID
	15	3	0	670.00	710.00	89	7.528	7.978	0.01	0.01	717.00		LCV FISH
	15	3	0	670.00	710.00	89	7.528	7.978	0.01	0.01	717.00		LCV DAPHNIDS
	1	6	0	680.00	870.00	9300	0.073	0.094	0.00	0.00	100.00	<	LTV EC20 FISH
	1	6	0	680.00	870.00	9300	0.073	0.094	0.00	0.00	310.00		LTV EC20 DAPHNID
	1	6	0	680.00	870.00	9300	0.073	0.094	0.00	0.00	570.00		LCV DAPHNIDS
	1	6	0	680.00	870.00	9300	0.073	0.094	0.00	0.00	1995.00		EC20 POPULATION
DI-N-OCTYLPHTHALATE	1	6	0	82.00	1900.00	9300	0.009	0.204	0.00	0.00	100.00	<	LCV FISH
	2	8	0	82.00	1900.00	9300	0.009	0.204	0.00	0.00	100.00		LTV EC20 FISH
	2	8	0	82.00	1900.00	9300	0.009	0.204	0.00	0.00	310.00		LTV EC20 DAPHNID
	2	8	0	82.00	1900.00	9300	0.009	0.204	0.00	0.00	570.00		LCV DAPHNIDS
	2	8	0	82.00	1900.00	9300	0.009	0.204	0.00	0.00	1995.00		EC20 POPULATION
	2	8	0	82.00	1900.00	9300	0.009	0.204	0.00	0.00	5700.00		LCV FISH
	4	6	0	660.00	720.00	9300	0.071	0.077	0.00	0.00	100.00	<	LTV EC20 FISH
	4	6	0	660.00	720.00	9300	0.071	0.077	0.00	0.00	310.00		LTV EC20 DAPHNID
	4	6	0	660.00	720.00	9300	0.071	0.077	0.00	0.00	570.00		LCV DAPHNIDS
	4	6	0	660.00	720.00	9300	0.071	0.077	0.00	0.00	1995.00		EC20 POPULATION
	4	6	0	660.00	720.00	9300	0.071	0.077	0.00	0.00	5700.00		LCV FISH
	15	3	0	670.00	710.00	9300	0.072	0.076	0.00	0.00	100.00	<	LTV EC20 FISH
	15	3	0	670.00	710.00	9300	0.072	0.076	0.00	0.00	310.00		LTV EC20 DAPHNID
	15	3	0	670.00	710.00	9300	0.072	0.076	0.00	0.00	570.00		LCV DAPHNIDS
	15	3	0	670.00	710.00	9300	0.072	0.076	0.00	0.00	1995.00		EC20 POPULATION
	15	3	0	670.00	710.00	9300	0.072	0.076	0.00	0.00	5700.00		LCV FISH
DIBENZOFURAN	1	6	0	680.00	870.00	.	.	.	.	.	2.00	*	NAHQ ADVISORY
	1	6	0	680.00	870.00	.	.	.	.	.	1003.00	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	.	.	.	.	.	2.00	*	NAHQ ADVISORY
	2	8	0	700.00	1900.00	.	.	.	.	.	1003.00	*	LCV DAPHNIDS
	3	34	0	190.00	410.00	.	.	.	.	.	2.00	*	NAHQ ADVISORY
	3	34	0	190.00	410.00	.	.	.	.	.	1003.00	*	LCV DAPHNIDS
	4	6	0	660.00	720.00	.	.	.	.	.	2.00	*	NAHQ ADVISORY
	4	6	0	660.00	720.00	.	.	.	.	.	1003.00	*	LCV DAPHNIDS
	15	3	0	670.00	710.00	.	.	.	.	.	2.00	*	NAHQ ADVISORY
	15	3	0	670.00	710.00	.	.	.	.	.	1003.00	*	LCV DAPHNIDS
	15	3	0	670.00	710.00	.	.	.	.	.	2.00	*	NAHQ ADVISORY
	15	3	0	670.00	710.00	.	.	.	.	.	1003.00	*	LCV DAPHNIDS

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
DIETHYLPHTHALATE	1	6	0	680.00	870.00	120	5.667	7.250	0.21	0.27	27.00		NAHQ ADVISORY
	1	6	0	680.00	870.00	120	5.667	7.250	0.01	0.01	1000.00		EC20 POPULATION
	2	8	0	80.00	1900.00	120	0.667	15.833	0.02	0.59	27.00		NAHQ ADVISORY
	3	8	0	80.00	1900.00	120	0.667	15.833	0.00	0.02	1000.00		EC20 POPULATION
	3	34	0	190.00	410.00	120	1.583	3.417	0.06	0.13	27.00		NAHQ ADVISORY
	3	34	0	190.00	410.00	120	1.583	3.417	0.00	0.00	1000.00		EC20 POPULATION
	4	6	0	660.00	720.00	120	5.500	6.000	0.20	0.22	27.00		NAHQ ADVISORY
	4	6	0	660.00	720.00	120	5.500	6.000	0.01	0.01	1000.00		EC20 POPULATION
	15	3	0	670.00	710.00	120	5.583	5.917	0.21	0.22	27.00		NAHQ ADVISORY
	15	3	0	670.00	710.00	120	5.583	5.917	0.01	0.01	1000.00		EC20 POPULATION
FLUORANTHENE	1	6	0	680.00	870.00	379	1.794	2.296	1.06	1.35	1.70		NAHQ ADVISORY
	1	6	0	680.00	870.00	379	1.794	2.296	0.06	0.07	32.00	*	EC20 POPULATION
	1	6	0	680.00	870.00	379	1.794	2.296	0.00	0.00	65795.00		LCV DAPHNIDS
	2	8	0	700.00	1900.00	379	1.847	5.013	1.09	2.95	1.70		NAHQ ADVISORY
	2	8	0	700.00	1900.00	379	1.847	5.013	0.06	0.16	32.00	*	EC20 POPULATION
	2	8	0	700.00	1900.00	379	1.847	5.013	0.00	0.00	65795.00		LCV DAPHNIDS
	3	34	0	190.00	410.00	379	0.501	1.082	0.29	0.64	1.70		NAHQ ADVISORY
	3	34	0	190.00	410.00	379	0.501	1.082	0.02	0.03	32.00	*	EC20 POPULATION
	3	34	0	190.00	410.00	379	0.501	1.082	0.00	0.00	65795.00		LCV DAPHNIDS
	4	7	0	660.00	730.00	379	1.741	1.926	1.02	1.13	1.70		NAHQ ADVISORY
	4	7	0	660.00	730.00	379	1.741	1.926	0.05	0.06	32.00	*	EC20 POPULATION
	4	7	0	660.00	730.00	379	1.741	1.926	0.00	0.00	65795.00		LCV DAPHNIDS
	15	3	0	670.00	710.00	379	1.768	1.873	1.04	1.10	1.70		NAHQ ADVISORY
	15	3	0	670.00	710.00	379	1.768	1.873	0.06	0.06	32.00	*	EC20 POPULATION
	15	3	0	670.00	710.00	379	1.768	1.873	0.00	0.00	65795.00		LCV DAPHNIDS
LEAD	1	28	0	280.00	580.00	300	0.933	1.933	2.67	5.52	0.35		SS EC20
	1	28	0	280.00	580.00	300	0.933	1.933	0.29	0.60	3.20		NAHQ ACTUAL
	1	28	0	280.00	580.00	300	0.933	1.933	0.08	0.16	12.26		LCV DAPHNIDS
	1	28	0	280.00	580.00	300	0.933	1.933	0.05	0.10	18.88		LCV FISH
	1	28	0	280.00	580.00	300	0.933	1.933	0.04	0.09	22.00		LTV EC20 FISH
	1	28	0	280.00	580.00	300	0.933	1.933	0.01	0.03	71.00		EC20 POPULATION
	2	13	0	460.00	600.00	300	1.533	2.000	4.38	5.71	0.35		SS EC20
	2	13	0	460.00	600.00	300	1.533	2.000	0.48	0.63	3.20		NAHQ ACTUAL
	2	13	0	460.00	600.00	300	1.533	2.000	0.13	0.16	12.26		LCV DAPHNIDS
	2	13	0	460.00	600.00	300	1.533	2.000	0.08	0.11	18.88		LCV FISH
	2	13	0	460.00	600.00	300	1.533	2.000	0.07	0.09	22.00		LTV EC20 FISH
	2	13	0	460.00	600.00	300	1.533	2.000	0.02	0.03	71.00		EC20 POPULATION
	3	25	0	300.00	590.00	300	1.000	1.967	2.86	5.62	0.35		SS EC20
	3	25	0	300.00	590.00	300	1.000	1.967	0.31	0.61	3.20		NAHQ ACTUAL
	3	25	0	300.00	590.00	300	1.000	1.967	0.08	0.16	12.26		LCV DAPHNIDS
	3	25	0	300.00	590.00	300	1.000	1.967	0.05	0.10	18.88		LCV FISH
	3	25	0	300.00	590.00	300	1.000	1.967	0.01	0.09	22.00		LTV EC20 FISH
	3	25	0	300.00	590.00	300	1.000	1.967	0.05	0.09	22.00		EC20 POPULATION
	4	17	0	480.00	610.00	300	1.600	2.033	4.57	5.81	0.35		SS EC20
	4	17	0	480.00	610.00	300	1.600	2.033	0.50	0.64	3.20		NAHQ ACTUAL

Table F.6 (continued)

Analyte	Resch	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
LEAD (continued)	4	17	0	480.00	610.00	300	1.600	2.033	0.13	0.17	12.26		LCV DAPHNIDS
	4	17	0	480.00	610.00	300	1.600	2.033	0.08	0.11	18.88		LCV FISH
	4	17	0	480.00	610.00	300	1.600	2.033	0.07	0.09	22.00		LTV EC20 FISH
	4	17	0	480.00	610.00	300	1.600	2.033	0.02	0.03	71.00		EC20 POPULATION
	5	5	0	460.00	510.00	300	1.533	1.700	4.38	4.86	0.35		SS EC20
	5	5	0	460.00	510.00	300	1.533	1.700	0.48	0.53	3.20		NAHQ ACTUAL
	5	5	0	460.00	510.00	300	1.533	1.700	0.13	0.14	12.26		LCV DAPHNIDS
	5	5	0	460.00	510.00	300	1.533	1.700	0.08	0.09	18.88		LCV FISH
	5	5	0	460.00	510.00	300	1.533	1.700	0.07	0.08	22.00		LTV EC20 FISH
	5	5	0	460.00	510.00	300	1.533	1.700	0.02	0.02	71.00		EC20 POPULATION
	10	5	0	350.00	440.00	300	1.167	1.467	3.33	4.19	0.35		SS EC20
	10	5	0	350.00	440.00	300	1.167	1.467	0.36	0.46	3.20		NAHQ ACTUAL
	10	5	0	350.00	440.00	300	1.167	1.467	0.10	0.12	12.26		LCV DAPHNIDS
	10	5	0	350.00	440.00	300	1.167	1.467	0.06	0.08	18.88		LCV FISH
	10	5	0	350.00	440.00	300	1.167	1.467	0.05	0.07	22.00		LTV EC20 FISH
	10	5	0	350.00	440.00	300	1.167	1.467	0.02	0.02	71.00		EC20 POPULATION
	13	10	0	450.00	590.00	300	1.500	1.967	4.29	5.62	0.35		SS EC20
	13	10	0	450.00	590.00	300	1.500	1.967	0.47	0.61	3.20		NAHQ ACTUAL
	13	10	0	450.00	590.00	300	1.500	1.967	0.12	0.16	12.26		LCV DAPHNIDS
	13	10	0	450.00	590.00	300	1.500	1.967	0.08	0.10	18.88		LCV FISH
	13	10	0	450.00	590.00	300	1.500	1.967	0.07	0.09	22.00		LTV EC20 FISH
	13	10	0	450.00	590.00	300	1.500	1.967	0.02	0.03	71.00		EC20 POPULATION
	15	16	0	350.00	570.00	300	1.167	1.900	3.33	5.43	0.35		SS EC20
	15	16	0	350.00	570.00	300	1.167	1.900	0.36	0.59	3.20		NAHQ ACTUAL
	15	16	0	350.00	570.00	300	1.167	1.900	0.10	0.15	12.26		LCV DAPHNIDS
	15	16	0	350.00	570.00	300	1.167	1.900	0.06	0.10	18.88		LCV FISH
	15	16	0	350.00	570.00	300	1.167	1.900	0.05	0.09	22.00		LTV EC20 FISH
	15	16	0	350.00	570.00	300	1.167	1.900	0.02	0.03	71.00		EC20 POPULATION
N-NITROSODIPHENYLAMINE	1	6	0	680.00	870.00	91	7.473	9.560	2.99	3.82	2.50		NAHQ ADVISORY
	1	6	0	680.00	870.00	91	7.473	9.560	0.19	0.24	40.00	*	EC20 POPULATION
	1	6	0	680.00	870.00	91	7.473	9.560	0.02	0.03	332.00	*	LCV FISH
	1	6	0	680.00	870.00	91	7.473	9.560	0.02	0.03	339.00	*	LTV EC20 FISH
	1	6	0	680.00	870.00	91	7.473	9.560	0.01	0.01	1042.00	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	91	7.692	20.879	3.08	8.35	2.50		NAHQ ADVISORY
	2	8	0	700.00	1900.00	91	7.692	20.879	0.19	0.52	40.00	*	EC20 POPULATION
	2	8	0	700.00	1900.00	91	7.692	20.879	0.02	0.06	332.00	*	LCV FISH
	2	8	0	700.00	1900.00	91	7.692	20.879	0.02	0.06	339.00	*	LTV EC20 FISH
	2	8	0	700.00	1900.00	91	7.692	20.879	0.01	0.02	1042.00	*	LCV DAPHNIDS
	3	34	0	190.00	410.00	91	2.088	4.505	0.84	1.80	2.50		NAHQ ADVISORY
	3	34	0	190.00	410.00	91	2.088	4.505	0.05	0.11	40.00	*	EC20 POPULATION
	3	34	0	190.00	410.00	91	2.088	4.505	0.01	0.01	332.00	*	LCV FISH
	3	34	0	190.00	410.00	91	2.088	4.505	0.01	0.01	339.00	*	LTV EC20 FISH
	3	34	0	190.00	410.00	91	2.088	4.505	0.00	0.00	1042.00	*	LCV DAPHNIDS
	4	6	0	660.00	720.00	91	7.253	7.912	2.90	3.16	2.50		NAHQ ADVISORY
	4	6	0	660.00	720.00	91	7.253	7.912	0.18	0.20	40.00	*	EC20 POPULATION
	4	6	0	660.00	720.00	91	7.253	7.912	0.02	0.02	332.00	*	LCV FISH

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
N-NITROSDIPHENYLAMINE (continued)	4	6	0	660.00	720.00	91	7.253	7.912	0.02	0.02	339.00	*	LTV EC20 FISH
	4	6	0	660.00	720.00	91	7.253	7.912	0.01	0.01	1042.00	*	LCV DAPHNIDS
	15	3	0	670.00	710.00	91	7.363	7.802	2.95	3.12	2.50		NAHQ ADVISORY
	15	3	0	670.00	710.00	91	7.363	7.802	0.18	0.20	40.00		EC20 POPULATION
	15	3	0	670.00	710.00	91	7.363	7.802	0.02	0.02	332.00	*	LCV FISH
	15	3	0	670.00	710.00	91	7.363	7.802	0.02	0.02	339.00	*	LTV EC20 FISH
	15	3	0	670.00	710.00	91	7.363	7.802	0.01	0.01	1042.00	*	LCV DAPHNIDS
	1	6	0	680.00	870.00	430	1.581	2.023	0.75	0.96	2.10		NAHQ ADVISORY
	1	6	0	680.00	870.00	430	1.581	2.023	0.00	0.00	450.00		LTV EC20 FISH
	1	6	0	680.00	870.00	430	1.581	2.023	0.00	0.00	600.00	>	LTV EC20 DAPHNID
	1	6	0	680.00	870.00	430	1.581	2.023	0.00	0.00	620.00		LCV FISH
	1	6	0	680.00	870.00	430	1.581	2.023	0.00	0.00	1000.00		EC20 POPULATION
	1	6	0	680.00	870.00	430	1.581	2.023	0.00	0.00	1163.00	*	LCV DAPHNIDS
	1	6	0	700.00	1900.00	430	1.628	4.419	0.78	2.10	2.10		NAHQ ADVISORY
	2	8	0	700.00	1900.00	430	1.628	4.419	0.00	0.01	450.00	>	LTV EC20 FISH
NAPHTHALENE	2	8	0	700.00	1900.00	430	1.628	4.419	0.00	0.01	600.00	>	LTV EC20 DAPHNID
	2	8	0	700.00	1900.00	430	1.628	4.419	0.00	0.01	620.00		LCV FISH
	2	8	0	700.00	1900.00	430	1.628	4.419	0.00	0.00	1000.00		EC20 POPULATION
	2	8	0	700.00	1900.00	430	1.628	4.419	0.00	0.00	1163.00	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	430	1.628	4.419	0.00	0.00	2.10		NAHQ ADVISORY
	3	34	0	190.00	410.00	430	0.442	0.953	0.21	0.45	2.10		LTV EC20 FISH
	3	34	0	190.00	410.00	430	0.442	0.953	0.00	0.00	450.00	>	LTV EC20 DAPHNID
	3	34	0	190.00	410.00	430	0.442	0.953	0.00	0.00	600.00	>	LTV EC20 FISH
	3	34	0	190.00	410.00	430	0.442	0.953	0.00	0.00	620.00		LCV FISH
	3	34	0	190.00	410.00	430	0.442	0.953	0.00	0.00	1000.00		EC20 POPULATION
	3	34	0	190.00	410.00	430	0.442	0.953	0.00	0.00	1163.00	*	LCV DAPHNIDS
	4	6	0	660.00	720.00	430	1.535	1.674	0.73	0.80	2.10		NAHQ ADVISORY
	4	6	0	660.00	720.00	430	1.535	1.674	0.00	0.00	450.00	>	LTV EC20 FISH
	4	6	0	660.00	720.00	430	1.535	1.674	0.00	0.00	600.00	>	LTV EC20 DAPHNID
	4	6	0	660.00	720.00	430	1.535	1.674	0.00	0.00	620.00		LCV FISH
	4	6	0	660.00	720.00	430	1.535	1.674	0.00	0.00	1000.00		EC20 POPULATION
	4	6	0	660.00	720.00	430	1.535	1.674	0.00	0.00	1163.00	*	LCV DAPHNIDS
	5	1	0	57.00	57.00	430	0.133	0.133	0.06	0.06	2.10		NAHQ ADVISORY
	5	1	0	57.00	57.00	430	0.133	0.133	0.00	0.00	450.00	>	LTV EC20 FISH
	5	1	0	57.00	57.00	430	0.133	0.133	0.00	0.00	600.00	>	LTV EC20 DAPHNID
	5	1	0	57.00	57.00	430	0.133	0.133	0.00	0.00	620.00		LCV FISH
	5	1	0	57.00	57.00	430	0.133	0.133	0.00	0.00	1000.00		EC20 POPULATION
	5	1	0	57.00	57.00	430	0.133	0.133	0.00	0.00	1163.00	*	LCV DAPHNIDS
	15	3	0	670.00	710.00	430	1.558	1.651	0.74	0.79	2.10		NAHQ ADVISORY
	15	3	0	670.00	710.00	430	1.558	1.651	0.00	0.00	450.00	>	LTV EC20 FISH
	15	3	0	670.00	710.00	430	1.558	1.651	0.00	0.00	600.00	>	LTV EC20 DAPHNID
	15	3	0	670.00	710.00	430	1.558	1.651	0.00	0.00	620.00		LCV FISH
	15	3	0	670.00	710.00	430	1.558	1.651	0.00	0.00	1000.00		EC20 POPULATION
	15	3	0	670.00	710.00	430	1.558	1.651	0.00	0.00	1163.00	*	LCV DAPHNIDS

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark Flag	Benchmark type
NICKEL	1	28	0	280.00	1100.00	100	2.800	11.000	0.56	2.20	<	LCV DAPHNIDS
	1	28	0	280.00	1100.00	100	2.800	11.000	0.25	1.00	<	SS EC20
	1	28	0	280.00	1100.00	100	2.800	11.000	0.08	0.31	<	LCV FISH
	1	28	0	280.00	1100.00	100	2.800	11.000	0.06	0.24	<	LTV EC20 DAPHNID
	1	28	0	280.00	1100.00	100	2.800	11.000	0.05	0.18	<	LTV EC20 FISH
	1	28	0	280.00	1100.00	100	2.800	11.000	0.02	0.07	<	NAHQ ACTUAL
	1	28	0	280.00	1100.00	100	2.800	11.000	0.01	0.05	<	EC20 POPULATION
	1	28	0	280.00	1100.00	100	2.800	11.000	0.92	1.20	<	LCV DAPHNIDS
	2	13	0	460.00	600.00	100	4.600	6.000	0.42	0.55	<	SS EC20
	2	13	0	460.00	600.00	100	4.600	6.000	0.13	0.17	<	LCV FISH
	2	13	0	460.00	600.00	100	4.600	6.000	0.10	0.13	<	LTV EC20 DAPHNID
	2	13	0	460.00	600.00	100	4.600	6.000	0.07	0.10	<	LTV EC20 FISH
	2	13	0	460.00	600.00	100	4.600	6.000	0.03	0.04	<	NAHQ ACTUAL
	2	13	0	460.00	600.00	100	4.600	6.000	0.02	0.03	<	EC20 POPULATION
	2	13	0	460.00	600.00	100	4.600	6.000	0.60	1.18	<	LCV DAPHNIDS
	3	35	0	300.00	590.00	100	3.000	5.900	0.27	0.54	<	SS EC20
	3	35	0	300.00	590.00	100	3.000	5.900	0.09	0.17	<	LCV FISH
	3	35	0	300.00	590.00	100	3.000	5.900	0.07	0.13	<	LTV EC20 DAPHNID
	3	35	0	300.00	590.00	100	3.000	5.900	0.05	0.10	<	LTV EC20 FISH
	3	35	0	300.00	590.00	100	3.000	5.900	0.02	0.04	<	NAHQ ACTUAL
	3	35	0	300.00	590.00	100	3.000	5.900	0.01	0.03	<	EC20 POPULATION
	3	35	0	300.00	590.00	100	3.000	5.900	0.96	1.22	<	LCV DAPHNIDS
	4	27	0	480.00	610.00	100	4.800	6.100	0.44	0.55	<	SS EC20
	4	27	0	480.00	610.00	100	4.800	6.100	0.14	0.17	<	LCV FISH
	4	27	0	480.00	610.00	100	4.800	6.100	0.11	0.14	<	LTV EC20 DAPHNID
	4	27	0	480.00	610.00	100	4.800	6.100	0.08	0.10	<	LTV EC20 FISH
	4	27	0	480.00	610.00	100	4.800	6.100	0.03	0.04	<	NAHQ ACTUAL
	4	27	0	480.00	610.00	100	4.800	6.100	0.02	0.03	<	EC20 POPULATION
	4	27	0	480.00	610.00	100	4.800	6.100	0.92	1.02	<	LCV DAPHNIDS
	5	5	0	460.00	510.00	100	4.600	5.100	0.42	0.46	<	SS EC20
	5	5	0	460.00	510.00	100	4.600	5.100	0.13	0.15	<	LCV FISH
	5	5	0	460.00	510.00	100	4.600	5.100	0.10	0.11	<	LTV EC20 DAPHNID
	5	5	0	460.00	510.00	100	4.600	5.100	0.07	0.08	<	LTV EC20 FISH
	5	5	0	460.00	510.00	100	4.600	5.100	0.03	0.03	<	NAHQ ACTUAL
	5	5	0	460.00	510.00	100	4.600	5.100	0.02	0.02	<	EC20 POPULATION
	5	5	0	350.00	610.00	100	3.500	6.100	0.70	1.22	<	LCV DAPHNIDS
	10	15	0	350.00	610.00	100	3.500	6.100	0.32	0.55	<	SS EC20
	10	15	0	350.00	610.00	100	3.500	6.100	0.10	0.17	<	LCV FISH
	10	15	0	350.00	610.00	100	3.500	6.100	0.08	0.14	<	LTV EC20 DAPHNID
	10	15	0	350.00	610.00	100	3.500	6.100	0.06	0.10	<	LTV EC20 FISH
	10	15	0	350.00	610.00	100	3.500	6.100	0.02	0.04	<	NAHQ ACTUAL
	10	15	0	350.00	610.00	100	3.500	6.100	0.02	0.03	<	EC20 POPULATION
	13	10	0	450.00	590.00	100	4.500	5.900	0.90	1.18	<	LCV DAPHNIDS
	13	10	0	450.00	590.00	100	4.500	5.900	0.41	0.54	<	SS EC20
	13	10	0	450.00	590.00	100	4.500	5.900	0.13	0.17	<	LCV FISH
	13	10	0	450.00	590.00	100	4.500	5.900	0.10	0.13	<	LTV EC20 DAPHNID
	13	10	0	450.00	590.00	100	4.500	5.900	0.07	0.10	<	LTV EC20 FISH

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
NICKEL (continued)	13	10	0	450.00	590.00	100	4.500	5.900	0.03	0.04	160.00		NAWQ ACTUAL
	13	10	0	450.00	590.00	100	4.500	5.900	0.02	0.03	215.00	<	EC20 POPULATION
	15	16	0	350.00	570.00	100	3.500	5.700	0.70	1.14	5.00		LCV DAPHNIDS
	15	16	0	350.00	570.00	100	3.500	5.700	0.32	0.52	11.00		SS EC20
	15	16	0	350.00	570.00	100	3.500	5.700	0.10	0.16	35.00	<	LCV FISH
	15	16	0	350.00	570.00	100	3.500	5.700	0.08	0.13	45.00		LTV EC20 DAPHNID
	15	16	0	350.00	570.00	100	3.500	5.700	0.06	0.09	62.00		LTV EC20 FISH
	15	16	0	350.00	570.00	100	3.500	5.700	0.02	0.04	160.00		NAWQ ACTUAL
	15	16	0	350.00	570.00	100	3.500	5.700	0.02	0.03	215.00		EC20 POPULATION
	1	6	0	680.00	870.00	325	2.092	2.677	4.87	6.23	0.43		NAWQ ADVISORY
	1	6	0	680.00	870.00	325	2.092	2.677	0.02	0.02	110.00		LTV EC20 DAPHNID
	1	6	0	680.00	870.00	325	2.092	2.677	0.01	0.01	200.00		LCV DAPHNIDS
	2	8	0	700.00	1900.00	325	2.154	5.846	5.01	13.60	0.43		NAWQ ADVISORY
	2	8	0	700.00	1900.00	325	2.154	5.846	0.02	0.05	110.00		LTV EC20 DAPHNID
	2	8	0	700.00	1900.00	325	2.154	5.846	0.01	0.03	200.00		LCV DAPHNIDS
	3	34	0	190.00	410.00	325	0.585	1.262	1.36	2.93	0.43		NAWQ ADVISORY
PHENANTHRENE	3	34	0	190.00	410.00	325	0.585	1.262	0.01	0.01	110.00		LTV EC20 DAPHNID
	3	34	0	190.00	410.00	325	0.585	1.262	0.00	0.01	200.00		LCV DAPHNIDS
	4	6	0	660.00	720.00	325	2.031	2.215	4.72	5.15	0.43		NAWQ ADVISORY
	4	6	0	660.00	720.00	325	2.031	2.215	0.02	0.02	110.00		LTV EC20 DAPHNID
	4	6	0	670.00	710.00	325	2.062	2.185	0.01	0.01	200.00		LCV DAPHNIDS
	15	3	0	670.00	710.00	325	2.062	2.185	4.79	5.08	0.43		NAWQ ADVISORY
	15	3	0	670.00	710.00	325	2.062	2.185	0.02	0.02	110.00		LTV EC20 DAPHNID
	15	3	0	670.00	710.00	325	2.062	2.185	0.01	0.01	200.00		LCV DAPHNIDS
	1	6	0	680.00	870.00	780	0.872	1.115	0.04	0.05	21.00		NAWQ ADVISORY
	1	6	0	680.00	870.00	780	0.872	1.115	0.00	0.00	230.00	<	LTV EC20 FISH
	1	6	0	680.00	870.00	780	0.872	1.115	0.00	0.00	1369.00		LCV FISH
	1	6	0	680.00	870.00	780	0.872	1.115	0.00	0.00	2005.00	*	LCV DAPHNIDS
	1	6	0	680.00	870.00	780	0.872	1.115	0.00	0.00	4467.00		EC20 POPULATION
	2	8	0	700.00	1900.00	780	0.897	2.436	0.04	0.12	21.00	<	NAWQ ADVISORY
	2	8	0	700.00	1900.00	780	0.897	2.436	0.00	0.01	230.00		LTV EC20 FISH
	2	8	0	700.00	1900.00	780	0.897	2.436	0.00	0.00	1369.00		LCV FISH
PHENOL	2	8	0	700.00	1900.00	780	0.897	2.436	0.00	0.00	2005.00	*	LCV DAPHNIDS
	2	8	0	700.00	1900.00	780	0.897	2.436	0.00	0.00	4467.00		EC20 POPULATION
	2	8	0	700.00	1900.00	780	0.897	2.436	0.01	0.03	21.00	<	NAWQ ADVISORY
	3	34	0	190.00	410.00	780	0.244	0.526	0.00	0.00	230.00		LTV EC20 FISH
	3	34	0	190.00	410.00	780	0.244	0.526	0.00	0.00	1369.00		LCV FISH
	3	34	0	190.00	410.00	780	0.244	0.526	0.00	0.00	2005.00	*	LCV DAPHNIDS
	3	34	0	190.00	410.00	780	0.244	0.526	0.00	0.00	4467.00		EC20 POPULATION
	4	13	0	660.00	830.00	780	0.846	1.064	0.04	0.05	21.00	<	NAWQ ADVISORY
	4	13	0	660.00	830.00	780	0.846	1.064	0.00	0.00	230.00		LTV EC20 FISH
	4	13	0	660.00	830.00	780	0.846	1.064	0.00	0.00	1369.00		LCV FISH
	4	13	0	660.00	830.00	780	0.846	1.064	0.00	0.00	2005.00	*	LCV DAPHNIDS
	4	13	0	660.00	830.00	780	0.846	1.064	0.00	0.00	4467.00		EC20 POPULATION
	4	13	0	660.00	830.00	780	0.846	1.064	0.00	0.00	21.00		NAWQ ADVISORY
	5	9	0	660.00	830.00	780	0.846	1.064	0.04	0.05	21.00		NAWQ ADVISORY

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
PHENOL (continued)	5	9	0	660.00	830.00	780	0.846	1.064	0.00	0.00	230.00	<	LTV EC20 FISH
	5	9	0	660.00	830.00	780	0.846	1.064	0.00	0.00	1369.00		LCV FISH
	5	9	0	660.00	830.00	780	0.846	1.064	0.00	0.00	2005.00	*	LCV DAPHNIDS
	5	9	0	660.00	830.00	780	0.846	1.064	0.00	0.00	4467.00		EC20 POPULATION
	13	2	0	690.00	700.00	780	0.885	0.897	0.04	0.04	21.00		NAHQ ADVISORY
	13	2	0	690.00	700.00	780	0.885	0.897	0.00	0.00	230.00	<	LTV EC20 FISH
	13	2	0	690.00	700.00	780	0.885	0.897	0.00	0.00	1369.00		LCV FISH
	13	2	0	690.00	700.00	780	0.885	0.897	0.00	0.00	2005.00	*	LCV DAPHNIDS
	13	2	0	690.00	700.00	780	0.885	0.897	0.00	0.00	4467.00		EC20 POPULATION
	15	4	0	670.00	770.00	780	0.859	0.987	0.04	0.05	21.00		NAHQ ADVISORY
	15	4	0	670.00	770.00	780	0.859	0.987	0.00	0.00	230.00	<	LTV EC20 FISH
	15	4	0	670.00	770.00	780	0.859	0.987	0.00	0.00	1369.00		LCV FISH
	15	4	0	670.00	770.00	780	0.859	0.987	0.00	0.00	2005.00	*	LCV DAPHNIDS
	15	4	0	670.00	770.00	780	0.859	0.987	0.00	0.00	4467.00		EC20 POPULATION
SELENIUM	5	25	0	460.00	630.00	8	57.500	78.750	22.12	30.29	2.60		SS EC20
	5	25	0	460.00	630.00	8	57.500	78.750	2.30	3.15	25.00		LTV EC20 DAPHNID
	5	25	0	460.00	630.00	8	57.500	78.750	1.64	2.25	35.00		NAHQ ACTUAL
	5	25	0	460.00	630.00	8	57.500	78.750	1.44	1.97	40.00		LTV EC20 FISH
	5	25	0	460.00	630.00	8	57.500	78.750	0.65	0.89	88.32		LCV FISH
	5	25	0	460.00	630.00	8	57.500	78.750	0.63	0.86	91.65		LCV DAPHNIDS
	5	25	0	460.00	630.00	8	57.500	78.750	21.63	28.37	2.60		SS EC20
	13	10	0	450.00	590.00	8	56.250	73.750	2.25	2.95	25.00		LTV EC20 DAPHNID
	13	10	0	450.00	590.00	8	56.250	73.750	1.61	2.11	35.00		NAHQ ACTUAL
	13	10	0	450.00	590.00	8	56.250	73.750	1.41	1.84	40.00		LTV EC20 FISH
	13	10	0	450.00	590.00	8	56.250	73.750	0.64	0.84	88.32		LCV FISH
	13	10	0	450.00	590.00	8	56.250	73.750	0.61	0.80	91.65		LCV DAPHNIDS
SILVER	1	28	0	110.00	230.00	2	55.000	115.000	458.33	958.33	0.12		LCV FISH
	1	28	0	110.00	230.00	2	55.000	115.000	392.86	821.43	0.14		SS EC20
	1	28	0	110.00	230.00	2	55.000	115.000	275.00	575.00	0.20		LTV EC20 FISH
	1	28	0	110.00	230.00	2	55.000	115.000	220.00	460.00	0.25		NAHQ ADVISORY
	1	28	0	110.00	230.00	2	55.000	115.000	171.87	359.37	0.32		EC20 POPULATION
	1	28	0	110.00	230.00	2	55.000	115.000	98.21	205.36	0.56	<	LTV EC20 DAPHNID
	1	28	0	110.00	230.00	2	55.000	115.000	21.15	44.23	2.60		LCV DAPHNIDS
	1	28	0	110.00	230.00	2	55.000	115.000	750.00	1000.00	0.12		LCV FISH
	2	13	0	180.00	240.00	2	90.000	120.000	642.86	857.14	0.14		SS EC20
	2	13	0	180.00	240.00	2	90.000	120.000	450.00	600.00	0.20		LTV EC20 FISH
	2	13	0	180.00	240.00	2	90.000	120.000	360.00	480.00	0.25		NAHQ ADVISORY
	2	13	0	180.00	240.00	2	90.000	120.000	281.25	375.00	0.32		EC20 POPULATION
	2	13	0	180.00	240.00	2	90.000	120.000	160.71	214.29	0.56	<	LTV EC20 DAPHNID
	2	13	0	180.00	240.00	2	90.000	120.000	34.62	46.15	2.60		LCV DAPHNIDS
	3	35	0	120.00	240.00	2	60.000	120.000	500.00	1000.00	0.12		LCV FISH
	3	35	0	120.00	240.00	2	60.000	120.000	428.57	857.14	0.14		SS EC20
	3	35	0	120.00	240.00	2	60.000	120.000	300.00	600.00	0.20		LTV EC20 FISH
	3	35	0	120.00	240.00	2	60.000	120.000	240.00	480.00	0.25		NAHQ ADVISORY
	3	35	0	120.00	240.00	2	60.000	120.000	187.50	375.00	0.32		EC20 POPULATION

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
SILVER (continued)	3	35	0	120.00	240.00	2	60.000	120.000	107.14	214.29	0.56	<	LTV EC20 DAPHNID
	3	35	0	120.00	240.00	2	60.000	120.000	23.08	46.15	2.60		LCV DAPHNIDS
	4	27	0	190.00	240.00	2	95.000	120.000	791.67	1000.00	0.12		LCV FISH
	4	27	0	190.00	240.00	2	95.000	120.000	678.57	857.14	0.14		SS EC20
	4	27	0	190.00	240.00	2	95.000	120.000	475.00	600.00	0.20		LTV EC20 FISH
	4	27	0	190.00	240.00	2	95.000	120.000	380.00	480.00	0.25		NAMQ ADVISORY
	4	27	0	190.00	240.00	2	95.000	120.000	296.87	375.00	0.32		EC20 POPULATION
	4	27	0	190.00	240.00	2	95.000	120.000	169.64	214.29	0.56	<	LTV EC20 DAPHNID
	4	27	0	190.00	240.00	2	95.000	120.000	36.54	46.15	2.60		LCV DAPHNIDS
	4	27	0	190.00	240.00	2	95.000	120.000	791.67	1041.67	0.12		LCV FISH
	5	25	0	190.00	250.00	2	95.000	125.000	678.57	892.86	0.14		SS EC20
	5	25	0	190.00	250.00	2	95.000	125.000	475.00	625.00	0.20		LTV EC20 FISH
	5	25	0	190.00	250.00	2	95.000	125.000	380.00	500.00	0.25		NAMQ ADVISORY
	5	25	0	190.00	250.00	2	95.000	125.000	296.87	390.62	0.32		EC20 POPULATION
	5	25	0	190.00	250.00	2	95.000	125.000	169.64	223.21	0.56	<	LTV EC20 DAPHNID
	5	25	0	190.00	250.00	2	95.000	125.000	36.54	48.08	2.60		LCV DAPHNIDS
	10	15	0	140.00	250.00	2	70.000	125.000	583.33	1041.67	0.12		LCV FISH
	10	15	0	140.00	250.00	2	70.000	125.000	500.00	892.86	0.14		SS EC20
	10	15	0	140.00	250.00	2	70.000	125.000	350.00	625.00	0.20		LTV EC20 FISH
	10	15	0	140.00	250.00	2	70.000	125.000	280.00	500.00	0.25		NAMQ ADVISORY
	10	15	0	140.00	250.00	2	70.000	125.000	218.75	390.62	0.32		EC20 POPULATION
	10	15	0	140.00	250.00	2	70.000	125.000	125.00	223.21	0.56	<	LTV EC20 DAPHNID
	10	15	0	140.00	250.00	2	70.000	125.000	26.92	48.08	2.60		LCV DAPHNIDS
	10	15	0	140.00	250.00	2	70.000	120.000	750.00	1000.00	0.12		LCV FISH
	13	10	0	180.00	240.00	2	90.000	120.000	857.14	857.14	0.14		SS EC20
	13	10	0	180.00	240.00	2	90.000	120.000	642.86	600.00	0.20		LTV EC20 FISH
	13	10	0	180.00	240.00	2	90.000	120.000	450.00	600.00	0.25		NAMQ ADVISORY
	13	10	0	180.00	240.00	2	90.000	120.000	360.00	480.00	0.32		EC20 POPULATION
	13	10	0	180.00	240.00	2	90.000	120.000	281.25	375.00	0.56	<	LTV EC20 DAPHNID
	13	10	0	180.00	240.00	2	90.000	120.000	160.71	214.29	2.60		LCV DAPHNIDS
	13	10	0	180.00	240.00	2	90.000	120.000	34.62	46.15	2.60		LCV FISH
	15	16	0	140.00	230.00	2	70.000	115.000	583.33	958.33	0.12		SS EC20
	15	16	0	140.00	230.00	2	70.000	115.000	500.00	821.43	0.14		LTV EC20 FISH
	15	16	0	140.00	230.00	2	70.000	115.000	350.00	575.00	0.20		NAMQ ADVISORY
	15	16	0	140.00	230.00	2	70.000	115.000	280.00	460.00	0.25		EC20 POPULATION
	15	16	0	140.00	230.00	2	70.000	115.000	218.75	359.37	0.32		LTV EC20 DAPHNID
	15	16	0	140.00	230.00	2	70.000	115.000	125.00	205.36	0.56	<	LCV DAPHNIDS
	15	16	0	140.00	230.00	2	70.000	115.000	26.92	44.23	2.60		LCV FISH
THALLIUM	1	28	0	20.00	20.00	.	.	.	.	.	2.50		NAMQ ADVISORY
	1	28	0	20.00	20.00	.	.	.	.	.	57.00		LCV FISH
	1	28	0	20.00	20.00	.	.	.	.	.	64.00		LTV EC20 DAPHNID
	1	28	0	20.00	20.00	.	.	.	.	.	67.00		EC20 POPULATION
	1	28	0	20.00	20.00	.	.	.	.	.	81.00		LTV EC20 FISH
	1	28	0	20.00	20.00	.	.	.	.	.	135.00		LCV DAPHNIDS
	1	28	0	20.00	20.00	.	.	.	.	.	2.50		NAMQ ADVISORY
	2	13	0	20.00	20.00	.	.	.	.	.	57.00		LCV FISH
	2	13	0	20.00	20.00	.	.	.	.	.	64.00		LTV EC20 DAPHNID
	2	13	0	20.00	20.00	.	.	.	.	.	.		

Table F.6 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	Maximum detection limit	BCF	Minimum exposure conc.	Maximum exposure conc.	Minimum detection limit quo.	Maximum detection limit quo.	Benchmark	Flag	Benchmark type
THALLIUM (continued)	2	13	0	20.00	20.00	.	.	.	.	.	67.00	.	EC20 POPULATION
	2	13	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	2	13	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS
	3	35	0	20.00	20.00	.	.	.	.	.	2.50	.	NAHQ ADVISORY
	3	35	0	20.00	20.00	.	.	.	.	.	57.00	.	LCV FISH
	3	35	0	20.00	20.00	.	.	.	.	.	64.00	.	LTV EC20 DAPHNID
	3	35	0	20.00	20.00	.	.	.	.	.	67.00	.	EC20 POPULATION
	3	35	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	3	35	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS
	3	35	0	20.00	20.00	.	.	.	.	.	2.50	.	NAHQ ADVISORY
	4	27	0	20.00	20.00	.	.	.	.	.	57.00	.	LCV FISH
	4	27	0	20.00	20.00	.	.	.	.	.	64.00	.	LTV EC20 DAPHNID
	4	27	0	20.00	20.00	.	.	.	.	.	67.00	.	EC20 POPULATION
	4	27	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	4	27	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS
	4	27	0	20.00	20.00	.	.	.	.	.	2.50	.	NAHQ ADVISORY
	5	16	0	20.00	20.00	.	.	.	.	.	57.00	.	LCV FISH
	5	16	0	20.00	20.00	.	.	.	.	.	64.00	.	LTV EC20 DAPHNID
	5	16	0	20.00	20.00	.	.	.	.	.	57.00	.	EC20 POPULATION
	5	16	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	5	16	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS
	5	16	0	20.00	20.00	.	.	.	.	.	2.50	.	NAHQ ADVISORY
	10	15	0	20.00	20.00	.	.	.	.	.	57.00	.	LCV FISH
	10	15	0	20.00	20.00	.	.	.	.	.	64.00	.	LTV EC20 DAPHNID
	10	15	0	20.00	20.00	.	.	.	.	.	67.00	.	EC20 POPULATION
	10	15	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	10	15	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS
	10	15	0	20.00	20.00	.	.	.	.	.	2.50	.	NAHQ ADVISORY
	13	10	0	20.00	20.00	.	.	.	.	.	57.00	.	LCV FISH
	13	10	0	20.00	20.00	.	.	.	.	.	64.00	.	LTV EC20 DAPHNID
	13	10	0	20.00	20.00	.	.	.	.	.	67.00	.	EC20 POPULATION
	13	10	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	13	10	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS
	13	10	0	20.00	20.00	.	.	.	.	.	2.50	.	NAHQ ADVISORY
	15	16	0	20.00	20.00	.	.	.	.	.	57.00	.	LCV FISH
	15	16	0	20.00	20.00	.	.	.	.	.	64.00	.	LTV EC20 DAPHNID
	15	16	0	20.00	20.00	.	.	.	.	.	67.00	.	EC20 POPULATION
	15	16	0	20.00	20.00	.	.	.	.	.	81.00	.	LTV EC20 FISH
	15	16	0	20.00	20.00	.	.	.	.	.	135.00	.	LCV DAPHNIDS

Table F.7. Concentrations of inorganic chemicals detected in sediment (ug/L), toxicological benchmarks (ug/L), and quotients for benthic aquatic life

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Mean ^a sediment conc.	UCL ^b sediment conc.	Mean quotient	UCL quotient	Benchmark	Benchmark type
ANTHONY	5	5	2	676.00	1288.75	0.3380	0.6444	2000	ER-L CONCENTRATION
	5	5	2	676.00	1288.75	0.0270	0.0515	25000	ER-M CONCENTRATION
	5	5	2	676.00	1288.75	0.0270	0.0515	25000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	3	313.12	398.35	0.1566	0.1992	2000	ER-L CONCENTRATION
	7	8	3	313.12	398.35	0.0125	0.0159	25000	ER-M CONCENTRATION
	7	8	3	313.12	398.35	0.0125	0.0159	25000	OVERALL APPARENT EFFECTS THRESHOLD
ARSENIC	1	2	4	6925.00	33925.69	0.2098	1.0281	33000	ER-L CONCENTRATION
	1	2	4	6925.00	33925.69	0.0815	0.3991	85000	ER-M CONCENTRATION
	1	2	4	6925.00	33925.69	0.1385	0.6785	50000	OVERALL APPARENT EFFECTS THRESHOLD
	2	2	4	3475.00	6333.90	0.1053	0.1919	33000	ER-L CONCENTRATION
	2	2	4	3475.00	6333.90	0.0409	0.0745	85000	ER-M CONCENTRATION
	2	2	4	3475.00	6333.90	0.0695	0.1267	50000	OVERALL APPARENT EFFECTS THRESHOLD
	3	2	4	16675.00	179632.08	0.5053	5.4434	33000	ER-L CONCENTRATION
	3	2	4	16675.00	179632.08	0.1962	2.1133	85000	ER-M CONCENTRATION
	3	2	4	16675.00	179632.08	0.3335	3.5926	50000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	5850.00	38250.82	0.1773	1.1591	33000	ER-L CONCENTRATION
	4	2	4	5850.00	38250.82	0.0688	0.4500	85000	ER-M CONCENTRATION
	4	2	4	5850.00	38250.82	0.1170	0.7650	50000	OVERALL APPARENT EFFECTS THRESHOLD
	5	6	10	10233.33	12268.99	0.3101	0.3718	33000	ER-L CONCENTRATION
	5	6	10	10233.33	12268.99	0.1204	0.1443	85000	ER-M CONCENTRATION
	5	6	10	10233.33	12268.99	0.2047	0.2454	50000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	15	11056.25	14128.35	0.3350	0.4281	33000	ER-L CONCENTRATION
	7	8	15	11056.25	14128.35	0.1301	0.1662	85000	ER-M CONCENTRATION
	7	8	15	11056.25	14128.35	0.2211	0.2826	50000	OVERALL APPARENT EFFECTS THRESHOLD
	10	1	2	7400.00	7400.00	0.2242	0.2242	33000	ER-L CONCENTRATION
	10	1	2	7400.00	7400.00	0.0871	0.0871	85000	ER-M CONCENTRATION
	10	1	2	7400.00	7400.00	0.1480	0.1480	50000	OVERALL APPARENT EFFECTS THRESHOLD
CADMIUM	1	2	3	850.00	1485.31	0.1700	0.2971	5000	ER-L CONCENTRATION
	1	2	3	850.00	1485.31	0.0944	0.1650	9000	ER-M CONCENTRATION
	1	2	3	850.00	1485.31	0.1700	0.2971	5000	OVERALL APPARENT EFFECTS THRESHOLD
	2	2	2	375.00	1327.97	0.0750	0.2656	5000	ER-L CONCENTRATION
	2	2	2	375.00	1327.97	0.0417	0.1476	9000	ER-M CONCENTRATION
	2	2	2	375.00	1327.97	0.0750	0.2656	5000	OVERALL APPARENT EFFECTS THRESHOLD
	3	2	4	1075.00	3933.90	0.2150	0.7868	5000	ER-L CONCENTRATION
	3	2	4	1075.00	3933.90	0.1194	0.4371	9000	ER-M CONCENTRATION
	3	2	4	1075.00	3933.90	0.2150	0.7868	5000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	845.00	4720.39	0.1690	0.9441	5000	ER-L CONCENTRATION
	4	2	4	845.00	4720.39	0.0939	0.5245	9000	ER-M CONCENTRATION
	4	2	4	845.00	4720.39	0.1690	0.9441	5000	OVERALL APPARENT EFFECTS THRESHOLD

Table F.7 (continued)

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Mean ^a sediment conc.	UCL ^b sediment conc.	Mean quotient	UCL quotient	Benchmark	Benchmark type
CADMIUM (continued)	5	6	9	2003.33	4906.72	0.4007	0.9813	5000	ER-L CONCENTRATION
	5	6	9	2003.33	4906.72	0.2226	0.5452	9000	ER-M CONCENTRATION
	5	6	9	2003.33	4906.72	0.4007	0.9813	5000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	1	562.50	770.43	0.1125	0.1541	5000	ER-L CONCENTRATION
	7	8	1	562.50	770.43	0.0625	0.0856	9000	ER-M CONCENTRATION
	7	8	1	562.50	770.43	0.1125	0.1541	5000	OVERALL APPARENT EFFECTS THRESHOLD
	1	2	4	11375.00	51081.89	0.1422	0.6385	80000	ER-L CONCENTRATION
	1	2	4	11375.00	51081.89	0.0784	0.3523	145000	ER-M CONCENTRATION
	2	2	4	8500.00	46618.61	0.1062	0.5827	80000	ER-L CONCENTRATION
	2	2	4	8500.00	46618.61	0.0586	0.3215	145000	ER-M CONCENTRATION
CHROMIUM	2	2	4	22750.00	70398.27	0.2844	0.8800	80000	ER-L CONCENTRATION
	3	2	4	22750.00	70398.27	0.1569	0.4855	145000	ER-M CONCENTRATION
	3	2	4	13650.00	35250.55	0.1706	0.4406	80000	ER-L CONCENTRATION
	4	2	4	13650.00	35250.55	0.0941	0.2431	145000	ER-M CONCENTRATION
	4	2	4	24266.67	31106.11	0.3033	0.3888	80000	ER-L CONCENTRATION
	5	6	10	24266.67	31106.11	0.1674	0.2145	145000	ER-M CONCENTRATION
	5	6	10	29775.00	41513.37	0.3722	0.5189	80000	ER-L CONCENTRATION
	7	8	15	29775.00	41513.37	0.2053	0.2863	145000	ER-M CONCENTRATION
	7	8	15	28100.00	28100.00	0.3512	0.3512	80000	ER-L CONCENTRATION
	10	1	2	28100.00	28100.00	0.1938	0.1938	145000	ER-M CONCENTRATION
	10	1	2	37050.00	82157.03	0.5293	1.1737	70000	ER-L CONCENTRATION
	1	2	4	37050.00	82157.03	0.0950	0.2107	390000	ER-M CONCENTRATION
	1	2	4	37050.00	82157.03	0.1235	0.2739	300000	OVERALL APPARENT EFFECTS THRESHOLD
	2	2	4	10500.00	23206.20	0.1500	0.3315	70000	ER-L CONCENTRATION
	2	2	4	10500.00	23206.20	0.0269	0.0595	390000	ER-M CONCENTRATION
	2	2	4	10500.00	23206.20	0.0350	0.0774	300000	OVERALL APPARENT EFFECTS THRESHOLD
	3	2	4	29000.00	105237.23	0.4143	1.5034	70000	ER-L CONCENTRATION
COPPER	3	2	4	29000.00	105237.23	0.0744	0.2698	390000	ER-M CONCENTRATION
	3	2	4	29000.00	105237.23	0.0967	0.3508	300000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	30450.00	181018.53	0.4350	2.5860	70000	ER-L CONCENTRATION
	4	2	4	30450.00	181018.53	0.0781	0.4642	390000	ER-M CONCENTRATION
	4	2	4	30450.00	181018.53	0.1015	0.6034	300000	OVERALL APPARENT EFFECTS THRESHOLD
	5	6	10	32050.00	44295.52	0.4579	0.6328	70000	ER-L CONCENTRATION
	5	6	10	32050.00	44295.52	0.0822	0.1136	390000	ER-M CONCENTRATION
	5	6	10	32050.00	44295.52	0.1068	0.1477	300000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	15	27643.75	37241.37	0.3949	0.5320	70000	ER-L CONCENTRATION
	7	8	15	27643.75	37241.37	0.0709	0.0955	390000	ER-M CONCENTRATION
	7	8	15	27643.75	37241.37	0.0921	0.1241	300000	OVERALL APPARENT EFFECTS THRESHOLD

Table F.7 (continued)

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Mean ^a sediment conc.	UCL ^b sediment conc.	Mean quotient	UCL quotient	Benchmark	Benchmark type
COPPER (continued)	10	1	2	53500.00	53500.00	0.7643	0.7643	70000	ER-L CONCENTRATION
	10	1	2	53500.00	53500.00	0.1372	0.1372	390000	ER-M CONCENTRATION
	10	1	2	53500.00	53500.00	0.1783	0.1783	300000	OVERALL APPARENT EFFECTS THRESHOLD
	1	2	4	29375.00	93858.99	0.8393	2.6817	35000	ER-L CONCENTRATION
	1	2	4	29375.00	93858.99	0.2670	0.8533	110000	ER-M CONCENTRATION
	1	2	4	29375.00	93858.99	0.0979	0.3129	300000	OVERALL APPARENT EFFECTS THRESHOLD
	2	2	4	13500.00	65595.44	0.3857	1.8742	35000	ER-L CONCENTRATION
	2	2	4	13500.00	65595.44	0.1227	0.5963	110000	ER-M CONCENTRATION
	2	2	4	13500.00	65595.44	0.0450	0.2187	300000	OVERALL APPARENT EFFECTS THRESHOLD
LEAD	3	2	4	25975.00	35822.31	0.7421	1.0235	35000	ER-L CONCENTRATION
	3	2	4	25975.00	35822.31	0.2361	0.3257	110000	ER-M CONCENTRATION
	3	2	4	25975.00	35822.31	0.0866	0.1194	300000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	20300.00	200728.11	0.5800	5.7351	35000	ER-L CONCENTRATION
	4	2	4	20300.00	200728.11	0.1845	1.8248	110000	ER-M CONCENTRATION
	4	2	4	20300.00	200728.11	0.0677	0.6691	300000	OVERALL APPARENT EFFECTS THRESHOLD
	5	6	10	47500.00	64534.61	1.3571	1.8438	35000	ER-L CONCENTRATION
	5	6	10	47500.00	64534.61	0.4318	0.5867	110000	ER-M CONCENTRATION
	5	6	10	47500.00	64534.61	0.1583	0.2151	300000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	15	18668.75	22529.67	0.5334	0.6437	35000	ER-L CONCENTRATION
	7	8	15	18668.75	22529.67	0.1697	0.2048	110000	ER-M CONCENTRATION
	7	8	15	18668.75	22529.67	0.0622	0.0751	300000	OVERALL APPARENT EFFECTS THRESHOLD
	10	1	2	39250.00	39250.00	1.1214	1.1214	35000	ER-L CONCENTRATION
	10	1	2	39250.00	39250.00	0.3568	0.3568	110000	ER-M CONCENTRATION
	10	1	2	39250.00	39250.00	0.1308	0.1308	300000	OVERALL APPARENT EFFECTS THRESHOLD
	1	2	4	102.50	769.58	0.6833	5.1305	150	ER-L CONCENTRATION
	1	2	4	102.50	769.58	0.0788	0.5920	1300	ER-M CONCENTRATION
	1	2	4	102.50	769.58	0.1025	0.7696	1000	OVERALL APPARENT EFFECTS THRESHOLD
MERCURY	2	1	2	550.00	550.00	3.6667	3.6667	150	ER-L CONCENTRATION
	2	1	2	550.00	550.00	0.4231	0.4231	1300	ER-M CONCENTRATION
	2	1	2	550.00	550.00	0.5500	0.5500	1000	OVERALL APPARENT EFFECTS THRESHOLD
	3	1	2	3665.00	3665.00	24.4333	24.4333	150	ER-L CONCENTRATION
	3	1	2	3665.00	3665.00	2.8192	2.8192	1300	ER-M CONCENTRATION
	3	1	2	3665.00	3665.00	3.6650	3.6650	1000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	2025.00	4883.90	13.5000	32.5593	150	ER-L CONCENTRATION
	4	2	4	2025.00	4883.90	1.5577	3.7568	1300	ER-M CONCENTRATION
	4	2	4	2025.00	4883.90	2.0250	4.8839	1000	OVERALL APPARENT EFFECTS THRESHOLD
	5	5	8	440.00	867.04	2.9333	5.7803	150	ER-L CONCENTRATION
	5	5	8	440.00	867.04	0.3385	0.6670	1300	ER-M CONCENTRATION
	5	5	8	440.00	867.04	0.4400	0.8670	1000	OVERALL APPARENT EFFECTS THRESHOLD

Table F.7 (continued)

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Mean ^a sediment conc.	UCL ^b sediment conc.	Mean quotient	UCL quotient	Benchmark	Benchmark type
MERCURY (continued)	10	1	2	80.00	80.00	0.5333	0.5333	150	ER-L CONCENTRATION
	10	1	2	80.00	80.00	0.0615	0.0615	1300	ER-M CONCENTRATION
	10	1	2	80.00	80.00	0.0800	0.0800	1000	OVERALL APPARENT EFFECTS THRESHOLD
NICKEL	1	2	4	14800.00	61812.96	0.4933	2.0604	30000	ER-L CONCENTRATION
	1	2	4	14800.00	61812.96	0.2960	1.2363	50000	ER-M CONCENTRATION
	2	2	4	7250.00	48545.17	0.2417	1.6182	30000	ER-L CONCENTRATION
	2	2	4	7250.00	48545.17	0.1450	0.9709	50000	ER-M CONCENTRATION
	3	2	4	36750.00	65338.96	1.2250	2.1780	30000	ER-L CONCENTRATION
	3	2	4	36750.00	65338.96	0.7350	1.3068	50000	ER-M CONCENTRATION
	4	2	4	15725.00	78938.37	0.5242	2.6313	30000	ER-L CONCENTRATION
	4	2	4	15725.00	78938.37	0.3145	1.5788	50000	ER-M CONCENTRATION
	5	6	10	21441.67	27853.92	0.7147	0.9285	30000	ER-L CONCENTRATION
	5	6	10	21441.67	27853.92	0.4288	0.5571	50000	ER-M CONCENTRATION
	7	8	15	25987.50	31632.11	0.8662	1.0544	30000	ER-L CONCENTRATION
	7	8	15	25987.50	31632.11	0.5197	0.6326	50000	ER-M CONCENTRATION
	10	1	2	24250.00	24250.00	0.8083	0.8083	30000	ER-L CONCENTRATION
	10	1	2	24250.00	24250.00	0.4850	0.4850	50000	ER-M CONCENTRATION
SILVER	1	2	4	300.00	1062.37	0.3000	1.0624	1000	ER-L CONCENTRATION
	1	2	4	300.00	1062.37	0.1364	0.4829	2200	ER-M CONCENTRATION
	1	2	4	300.00	1062.37	0.1765	0.6249	1700	OVERALL APPARENT EFFECTS THRESHOLD
	2	2	2	85.00	783.84	0.0850	0.7838	1000	ER-L CONCENTRATION
	2	2	2	85.00	783.84	0.0386	0.3563	2200	ER-M CONCENTRATION
	2	2	2	85.00	783.84	0.0500	0.4611	1700	OVERALL APPARENT EFFECTS THRESHOLD
	3	1	2	230.00	230.00	0.2300	0.2300	1000	ER-L CONCENTRATION
	3	1	2	230.00	230.00	0.1045	0.1045	2200	ER-M CONCENTRATION
	3	1	2	230.00	230.00	0.1353	0.1353	1700	OVERALL APPARENT EFFECTS THRESHOLD
	5	6	8	338.33	640.83	0.3383	0.6408	1000	ER-L CONCENTRATION
	5	6	8	338.33	640.83	0.1538	0.2913	2200	ER-M CONCENTRATION
	5	6	8	338.33	640.83	0.1990	0.3770	1700	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	9	63.12	93.67	0.0631	0.0937	1000	ER-L CONCENTRATION
	7	8	9	63.12	93.67	0.0287	0.0426	2200	ER-M CONCENTRATION
	7	8	9	63.12	93.67	0.0371	0.0551	1700	OVERALL APPARENT EFFECTS THRESHOLD
ZINC	1	2	4	81675.00	341199.23	0.6806	2.8433	120000	ER-L CONCENTRATION
	1	2	4	81675.00	341199.23	0.3025	1.2637	270000	ER-M CONCENTRATION
	1	2	4	81675.00	341199.23	0.3141	1.3123	260000	OVERALL APPARENT EFFECTS THRESHOLD
	2	2	4	39925.00	81537.82	0.3327	0.6795	120000	ER-L CONCENTRATION
	2	2	4	39925.00	81537.82	0.1479	0.3020	270000	ER-M CONCENTRATION
	2	2	4	39925.00	81537.82	0.1536	0.3136	260000	OVERALL APPARENT EFFECTS THRESHOLD

Table F.7 (continued)

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Mean ^a sediment conc.	UCL ^b sediment conc.	Mean quotient	UCL quotient	Benchmark	Benchmark type
ZINC (continued)	3	2	4	76550.00	178199.64	0.6379	1.4850	120000	ER-L CONCENTRATION
	3	2	4	76550.00	178199.64	0.2835	0.6600	270000	ER-M CONCENTRATION
	3	2	4	76550.00	178199.64	0.2944	0.6854	260000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	81425.00	482623.41	0.6785	4.0219	120000	ER-L CONCENTRATION
	4	2	4	81425.00	482623.41	0.3016	1.7875	270000	ER-M CONCENTRATION
	4	2	4	81425.00	482623.41	0.3132	1.8562	260000	OVERALL APPARENT EFFECTS THRESHOLD
	4	2	4	181116.67	248313.51	1.5093	2.0693	120000	ER-L CONCENTRATION
	5	6	10	181116.67	248313.51	0.6708	0.9197	270000	ER-M CONCENTRATION
	5	6	10	181116.67	248313.51	0.6966	0.9551	260000	OVERALL APPARENT EFFECTS THRESHOLD
	5	6	10	73600.00	89511.67	0.6133	0.7459	120000	ER-L CONCENTRATION
	7	8	15	73600.00	89511.67	0.2726	0.3315	270000	ER-M CONCENTRATION
	7	8	15	73600.00	89511.67	0.2831	0.3443	260000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	15	105500.00	105500.00	0.8792	0.8792	120000	ER-L CONCENTRATION
	10	1	2	105500.00	105500.00	0.3907	0.3907	270000	ER-M CONCENTRATION
	10	1	2	105500.00	105500.00	0.4058	0.4058	260000	OVERALL APPARENT EFFECTS THRESHOLD
	10	1	2	105500.00	105500.00	0.4058	0.4058	260000	OVERALL APPARENT EFFECTS THRESHOLD

^aIn cases of mixed detects and nondetects, statistics were calculated using all values, "as is."^bThe maximum concentration was used when the UCL concentration was missing.

Table F.8. Detection limits for inorganic chemicals not detected in sediment (ug/L), toxicological benchmarks (ug/L), and quotients for benthic aquatic life

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Minimum detection limit	UCL ^a detection limit	MDL quotient	UCL quotient	Benchmark	Benchmark type
ANTIMONY	1	1	0	700	700.000	0.35000	0.35000	2000	ER-L CONCENTRATION
	1	1	0	700	700.000	0.02800	0.02800	25000	ER-M CONCENTRATION
	1	1	0	700	700.000	0.02800	0.02800	25000	OVERALL APPARENT EFFECTS THRESHOLD
	3	1	0	300	300.000	0.15000	0.15000	2000	ER-L CONCENTRATION
	3	1	0	300	300.000	0.01200	0.01200	25000	ER-M CONCENTRATION
	3	1	0	300	300.000	0.01200	0.01200	25000	OVERALL APPARENT EFFECTS THRESHOLD
	4	1	0	300	300.000	0.01200	0.01200	2000	ER-L CONCENTRATION
	4	1	0	550	550.000	0.27500	0.27500	25000	ER-M CONCENTRATION
	4	1	0	550	550.000	0.02200	0.02200	25000	OVERALL APPARENT EFFECTS THRESHOLD
	4	1	0	550	550.000	0.02200	0.02200	2000	ER-L CONCENTRATION
	10	1	0	600	600.000	0.30000	0.30000	25000	ER-M CONCENTRATION
	10	1	0	600	600.000	0.02400	0.02400	25000	OVERALL APPARENT EFFECTS THRESHOLD
	10	1	0	600	600.000	0.02400	0.02400	25000	ER-L CONCENTRATION
	10	1	0	600	600.000	0.06667	0.06667	5000	ER-M CONCENTRATION
CADMIUM	10	1	0	600	600.000	0.12000	0.12000	5000	ER-L CONCENTRATION
	10	1	0	600	600.000	0.06667	0.06667	9000	ER-M CONCENTRATION
	10	1	0	600	600.000	0.12000	0.12000	5000	OVERALL APPARENT EFFECTS THRESHOLD
MERCURY	7	8	0	50	237.283	0.33333	1.58189	150	ER-L CONCENTRATION
	7	8	0	50	237.283	0.03846	0.18253	1300	ER-M CONCENTRATION
	7	8	0	50	237.283	0.05000	0.23728	1000	OVERALL APPARENT EFFECTS THRESHOLD
	7	8	0	50	237.283	0.05000	0.23728	1000	OVERALL APPARENT EFFECTS THRESHOLD

^aThe maximum detection limit is used in cases where the UCL detection limit was missing.

Table F-9. Mean and upper confidence limit (UCL) concentrations of organic chemicals detected in sediment (ug/L), toxicological benchmarks (ug/L), and quotients for benthic aquatic life.

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Mean ^a sediment conc.	Mean pore water conc.	UCL conc.	UCL pore water conc.	Mean quotient	UCL quotient	Benchmark type
2-METHYLNAPHTHALENE	3	2	2	.	323.50	.	3455.58	.	.	.	NAHQ ADVISORY
	3	2	2	.	323.50	.	3455.58	.	.	.	NAHQ ACTUAL
	3	2	2	.	323.50	.	3455.58	.	.	.	LCV FISH
	3	2	2	.	323.50	.	3455.58	.	.	.	LCV DAPHNIDS
	3	2	2	.	323.50	.	3455.58	.	.	.	LTV EC20 FISH
	3	2	2	.	323.50	.	3455.58	.	.	.	LTV EC20 DAPHNID
	3	2	2	.	323.50	.	3455.58	.	.	.	SS EC20
	3	2	2	.	323.50	.	3455.58	.	.	.	EC20 POPULATION
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	.	.	NAHQ ACTUAL
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	.	.	LCV DAPHNIDS
4,4'-DDO	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	.	.	LCV EC20 DAPHNID
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	.	.	SS EC20
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	119.943	777.219	0.00 NAHQ ADVISORY
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	0.083	0.535	0.61 EC20 POPULATION
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	0.030	0.193	1.69 LCV FISH
	1	2	1	6.02	527.50	0.0504	3418.16	0.3264	0.013	0.082	3.99 LTV EC20 FISH
	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	.	.	NAHQ ACTUAL
	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	.	.	LTV EC20 DAPHNID
	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	13.463	87.237	0.01 SS EC20
	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	6.731	43.619	0.02 LCV DAPHNIDS
4,4'-DDT	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	5.385	34.895	0.02 NAHQ ADVISORY
	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	0.308	1.994	0.35 LTV EC20 FISH
	1	2	1	5.69	527.50	0.1077	3418.16	0.6979	0.148	0.956	0.73 LCV FISH
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	.	.	NAHQ ACTUAL
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	0.539	4.347	0.02 SS EC20
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	0.021	0.167	0.52 LTV EC20 FISH
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	0.017	0.138	0.63 EC20 POPULATION
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	0.011	0.087	1.00 LCV FISH
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	0.009	0.072	1.20 LTV EC20 DAPHNID
	2	2	1	7.14	1487.50	0.0108	12001.88	0.0869	0.005	0.041	2.10 LCV DAPHNIDS
AROCLOP-1254	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	.	.	NAHQ ACTUAL
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	.	.	SS EC20
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	1.286	14.168	0.02 NAHQ ADVISORY
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	0.049	0.545	0.52 LTV EC20 FISH
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	0.041	0.450	0.63 EC20 POPULATION
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	0.026	0.283	1.00 LCV FISH
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	0.021	0.236	1.20 LTV EC20 DAPHNID
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	0.012	0.135	2.10 LCV DAPHNIDS
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	.	.	NAHQ ACTUAL
	3	2	2	7.14	3551.00	0.0257	39115.67	0.2834	.	.	SS EC20

Table F.9 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H ₂ O and SED	Mean ^a sediment conc.	Mean pore water conc.	UCL sediment conc.	UCL pore water conc.	Mean quotient	UCL quotient	Benchmark type
FLUORANTHENE	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	.	.	NAHQ ACTUAL
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	.	.	LCV FISH
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	.	.	LTV EC20 FISH
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	.	.	LTV EC20 DAPHNID
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	.	.	SS EC20
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	0.350	3.290	1.70 NAHQ ADVISORY
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	0.019	0.175	32.00 EC20 POPULATION
	3	2	4	4.90	472.50	0.5948	4443.19	5.5936	0.000	0.000	65795.00 LCV DAPHNIDS
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	.	.	NAHQ ACTUAL
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	5.529	23.920	SS EC20
NAPHTHALENE	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	0.026	0.112	2.10 NAHQ ADVISORY
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	0.019	0.084	450.00 LTV EC20 FISH
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	0.019	0.084	600.00 LTV EC20 DAPHNID
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	0.019	0.081	620.00 LCV FISH
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	0.012	0.050	1000.00 EC20 POPULATION
	3	2	4	3.59	451.75	11.6118	1954.26	50.2322	0.010	0.043	1163.00 LCV DAPHNIDS
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	.	.	NAHQ ACTUAL
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	.	.	LCV FISH
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	.	.	LTV EC20 FISH
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	.	.	SS EC20
PHENANTHRENE	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	1.909	17.418	EC20 POPULATION
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	0.007	0.068	0.43 NAHQ ADVISORY
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	0.007	0.068	110.00 LTV EC20 DAPHNID
	3	2	4	4.57	305.00	0.8209	2782.71	7.4898	0.004	0.037	200.00 LCV DAPHNIDS
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	NAHQ ADVISORY
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	NAHQ ACTUAL
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	LCV FISH
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	LCV DAPHNIDS
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	LTV EC20 FISH
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	LTV EC20 DAPHNID
PYRENE	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	SS EC20
	3	2	4	4.88	355.00	0.4680	2578.59	3.3992	.	.	EC20 POPULATION

^aIn cases of mixed detects and nondetects, statistics are calculated using all values, "as is."

Table F.10. Detection limits for organic chemicals not detected in sediment (ug/L), estimated pore water concentrations (ug/L), toxicological benchmarks (ug/L), and quotients for benthic aquatic life

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection pore water		Minimum detection pore water		MDL quotient	UCL quotient	Benchmark type
					limit	conc.	limit	conc.			
2,4-DINITROPHENOL	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	NAWQ ADVISORY
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	LCV FISH
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	LCV DAPHNIDS
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	LTV EC20 FISH
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	LTV EC20 DAPHNID
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	SS EC20
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	.	.	EC20 POPULATION
	1	2	0	1.54	3000	8652.09	33838.96	97592.63	57.68	650.62	NAWQ ACTUAL
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	NAWQ ADVISORY
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	LCV FISH
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	LCV DAPHNIDS
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	LTV EC20 FISH
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	LTV EC20 DAPHNID
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	SS EC20
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	.	.	EC20 POPULATION
	2	2	0	1.54	2150	6200.67	2835.31	8177.12	41.34	54.51	NAWQ ACTUAL
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	NAWQ ADVISORY
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	LCV FISH
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	LCV DAPHNIDS
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	LTV EC20 FISH
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	LTV EC20 DAPHNID
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	SS EC20
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	.	.	EC20 POPULATION
	3	2	0	1.54	2750	7931.09	2750.00	7931.09	52.87	52.87	NAWQ ACTUAL
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	NAWQ ADVISORY
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	LCV FISH
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	LCV DAPHNIDS
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	LTV EC20 FISH
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	LTV EC20 DAPHNID
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	SS EC20
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	.	.	EC20 POPULATION
	4	2	0	1.54	2450	7065.88	187826.42	541697.31	47.11	3611.32	NAWQ ACTUAL
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	NAWQ ADVISORY
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	LCV FISH
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	LCV DAPHNIDS
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	LTV EC20 FISH
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	LTV EC20 DAPHNID
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	SS EC20
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	.	.	EC20 POPULATION
	5	4	0	1.54	4150	11968.73	33158.52	95630.22	79.79	637.53	NAWQ ACTUAL

Table F.10 (continued)

Analyte	Resch (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
2,4-DINITROPHENOL (continued)	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	NAHQ ADVISORY
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	LCV FISH
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	LCV DAPHNIDS
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	LTV EC20 FISH
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	LTV EC20 DAPHNID
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	SS EC20
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	EC20 POPULATION
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	596.03	596.03	NAHQ ACTUAL
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	NAHQ ADVISORY
	10	1	0	1.54	31000	89404.98	31000.00	89404.98	.	.	NAHQ ACTUAL
2-METHYLNAPHTHALENE	1	2	0	.	620	.	6993.39	.	.	.	LCV FISH
	1	2	0	.	620	.	6993.39	.	.	.	LCV DAPHNIDS
	1	2	0	.	620	.	6993.39	.	.	.	LTV EC20 FISH
	1	2	0	.	620	.	6993.39	.	.	.	LTV EC20 DAPHNID
	1	2	0	.	620	.	6993.39	.	.	.	SS EC20
	1	2	0	.	620	.	6993.39	.	.	.	EC20 POPULATION
	1	2	0	.	620	.	6993.39	.	.	.	NAHQ ADVISORY
	1	2	0	.	445	.	616.33	.	.	.	NAHQ ACTUAL
	2	2	0	.	445	.	616.33	.	.	.	LCV FISH
	2	2	0	.	445	.	616.33	.	.	.	LCV DAPHNIDS
	2	2	0	.	445	.	616.33	.	.	.	LTV EC20 FISH
	2	2	0	.	445	.	616.33	.	.	.	LTV EC20 DAPHNID
	2	2	0	.	445	.	616.33	.	.	.	SS EC20
	2	2	0	.	445	.	616.33	.	.	.	EC20 POPULATION
	2	2	0	.	445	.	616.33	.	.	.	NAHQ ADVISORY
	2	2	0	.	500	.	38534.72	.	.	.	NAHQ ACTUAL
	4	4	0	.	500	.	38534.72	.	.	.	LCV FISH
	4	4	0	.	500	.	38534.72	.	.	.	LCV DAPHNIDS
	4	4	0	.	500	.	38534.72	.	.	.	LTV EC20 FISH
	4	4	0	.	500	.	38534.72	.	.	.	LTV EC20 DAPHNID
	4	4	0	.	500	.	38534.72	.	.	.	SS EC20
	4	4	0	.	500	.	38534.72	.	.	.	EC20 POPULATION
	4	4	0	.	500	.	38534.72	.	.	.	NAHQ ADVISORY
	5	5	0	.	855	.	6848.50	.	.	.	NAHQ ACTUAL
	5	5	0	.	855	.	6848.50	.	.	.	LCV FISH
	5	5	0	.	855	.	6848.50	.	.	.	LCV DAPHNIDS
	5	5	0	.	855	.	6848.50	.	.	.	LTV EC20 FISH
	5	5	0	.	855	.	6848.50	.	.	.	LTV EC20 DAPHNID
	5	5	0	.	855	.	6848.50	.	.	.	SS EC20
	5	5	0	.	855	.	6848.50	.	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
2-METHYLNAPHTHALENE (continued)												
	10	1	0	.	6500	.	6500.00	.	.	.	NAWQ ADVISORY	.
	10	1	0	.	6500	.	6500.00	.	.	.	NAWQ ACTUAL	.
	10	1	0	.	6500	.	6500.00	.	.	.	LCV FISH	.
	10	1	0	.	6500	.	6500.00	.	.	.	LCV DAPHNIDS	.
	10	1	0	.	6500	.	6500.00	.	.	.	LTV EC20 FISH	.
	10	1	0	.	6500	.	6500.00	.	.	.	LTV EC20 DAPHNID	.
	10	1	0	.	6500	.	6500.00	.	.	.	SS EC20	.
	10	1	0	.	6500	.	6500.00	.	.	.	EC20 POPULATION	.
2-METHYLPHENOL												
	1	2	0	1.95	620	695.65	6993.39	7846.71	.	.	NAWQ ACTUAL	.
	1	2	0	1.95	620	695.65	6993.39	7846.71	.	.	LTV EC20 DAPHNID	.
	1	2	0	1.95	620	695.65	6993.39	7846.71	.	.	SS EC20	.
	1	2	0	1.95	620	695.65	6993.39	7846.71	90.34	1019.05	7.70 NAWQ ADVISORY	74.00 EC20 POPULATION
	1	2	0	1.95	620	695.65	6993.39	7846.71	9.40	106.04	74.00 EC20 POPULATION	74.00 EC20 FISH
	1	2	0	1.95	620	695.65	6993.39	7846.71	1.48	16.70	470.00 LTV EC20 FISH	470.00 LTV EC20 FISH
	1	2	0	1.95	620	695.65	6993.39	7846.71	1.42	16.05	489.00 LCV FISH	489.00 LCV FISH
	1	2	0	1.95	620	695.65	6993.39	7846.71	0.53	5.96	1316.00 LCV DAPHNIDS	1316.00 LCV DAPHNIDS
	1	2	0	1.95	620	695.65	6993.39	7846.71	.	.	NAWQ ACTUAL	.
	2	2	0	1.95	445	499.30	616.33	691.53	.	.	LTV EC20 DAPHNID	.
	2	2	0	1.95	445	499.30	616.33	691.53	.	.	SS EC20	.
	2	2	0	1.95	445	499.30	616.33	691.53	.	.	7.70 NAWQ ADVISORY	74.00 EC20 POPULATION
	2	2	0	1.95	445	499.30	616.33	691.53	64.84	89.81	74.00 EC20 POPULATION	74.00 EC20 FISH
	2	2	0	1.95	445	499.30	616.33	691.53	6.75	9.35	470.00 LTV EC20 FISH	470.00 LTV EC20 FISH
	2	2	0	1.95	445	499.30	616.33	691.53	1.06	1.47	489.00 LCV FISH	489.00 LCV FISH
	2	2	0	1.95	445	499.30	616.33	691.53	1.02	1.41	1316.00 LCV DAPHNIDS	1316.00 LCV DAPHNIDS
	2	2	0	1.95	445	499.30	616.33	691.53	0.38	0.53	NAWQ ACTUAL	.
	3	2	0	1.95	565	633.94	599.27	672.39	.	.	LTV EC20 DAPHNID	.
	3	2	0	1.95	565	633.94	599.27	672.39	.	.	SS EC20	.
	3	2	0	1.95	565	633.94	599.27	672.39	.	.	7.70 NAWQ ADVISORY	74.00 EC20 POPULATION
	3	2	0	1.95	565	633.94	599.27	672.39	82.33	87.32	74.00 EC20 POPULATION	74.00 EC20 FISH
	3	2	0	1.95	565	633.94	599.27	672.39	8.57	9.09	470.00 LTV EC20 FISH	470.00 LTV EC20 FISH
	3	2	0	1.95	565	633.94	599.27	672.39	1.35	1.43	489.00 LCV FISH	489.00 LCV FISH
	3	2	0	1.95	565	633.94	599.27	672.39	1.30	1.38	1316.00 LCV DAPHNIDS	1316.00 LCV DAPHNIDS
	3	2	0	1.95	565	633.94	599.27	672.39	0.48	0.51	NAWQ ACTUAL	.
	4	2	0	1.95	500	561.01	38534.72	43236.66	.	.	LTV EC20 DAPHNID	.
	4	2	0	1.95	500	561.01	38534.72	43236.66	.	.	SS EC20	.
	4	2	0	1.95	500	561.01	38534.72	43236.66	72.86	5615.15	7.70 NAWQ ADVISORY	74.00 EC20 POPULATION
	4	2	0	1.95	500	561.01	38534.72	43236.66	7.58	584.28	74.00 EC20 POPULATION	74.00 EC20 FISH
	4	2	0	1.95	500	561.01	38534.72	43236.66	1.19	91.99	470.00 LTV EC20 FISH	470.00 LTV EC20 FISH
	4	2	0	1.95	500	561.01	38534.72	43236.66	1.15	88.42	489.00 LCV FISH	489.00 LCV FISH
	4	2	0	1.95	500	561.01	38534.72	43236.66	0.43	32.85	1316.00 LCV DAPHNIDS	1316.00 LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection pore water limit	Minimum conc.	UCL ^a limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
2-METHYLPHENOL (continued)	5	3	0	1.95	1070	1200.56	10367.99	11633.07	.	.	NAWQ ACTUAL
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	.	.	LTV EC20 DAPHNID
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	.	.	SS EC20
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	155.92	1510.79	7.70 NAWQ ADVISORY
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	16.22	157.20	74.00 EC20 POPULATION
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	2.55	24.75	470.00 LTV EC20 FISH
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	2.46	23.79	489.00 LCV FISH
	5	3	0	1.95	1070	1200.56	10367.99	11633.07	0.91	8.84	1316.00 LCV DAPHNIDS
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	.	.	NAWQ ACTUAL
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	.	.	LTV EC20 DAPHNID
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	.	.	SS EC20
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	947.16	947.16	7.70 NAWQ ADVISORY
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	98.56	98.56	74.00 EC20 POPULATION
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	15.52	15.52	470.00 LTV EC20 FISH
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	14.91	14.91	489.00 LCV FISH
	10	1	0	1.95	6500	7293.12	6500.00	7293.12	5.54	5.54	1316.00 LCV DAPHNIDS
2-NITROPHENOL	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	NAWQ ADVISORY
	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	LCV FISH
	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	LCV DAPHNIDS
	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	LTV EC20 FISH
	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	LTV EC20 DAPHNID
	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	SS EC20
	1	2	0	1.79	620	1005.52	6993.39	11341.94	6.70	75.61	EC20 POPULATION
	1	2	0	1.79	620	1005.52	6993.39	11341.94	.	.	NAWQ ACTUAL
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	NAWQ ADVISORY
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	LCV FISH
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	LCV DAPHNIDS
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	LTV EC20 FISH
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	LTV EC20 DAPHNID
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	SS EC20
	2	2	0	1.79	445	721.71	616.33	999.57	.	.	EC20 POPULATION
	2	2	0	1.79	445	721.71	616.33	999.57	4.81	6.66	NAWQ ACTUAL
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	NAWQ ADVISORY
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	LCV FISH
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	LCV DAPHNIDS
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	LTV EC20 FISH
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	LTV EC20 DAPHNID
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	SS EC20
	3	2	0	1.79	565	916.32	599.27	971.89	.	.	EC20 POPULATION
	3	2	0	1.79	565	916.32	599.27	971.89	6.11	6.48	NAWQ ACTUAL

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
2-NITROPHENOL (continued)	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	NAWQ ADVISORY
	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	LCV FISH
	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	LCV DAPHNIDS
	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	LTV EC20 FISH
	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	LTV EC20 DAPHNID
	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	SS EC20
	4	2	0	1.79	500	810.91	38534.72	62495.99	5.41	416.64	EC20 POPULATION
	4	2	0	1.79	500	810.91	38534.72	62495.99	.	.	NAWQ ACTUAL
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	NAWQ ADVISORY
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	LCV FISH
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	LCV DAPHNIDS
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	LTV EC20 FISH
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	LTV EC20 DAPHNID
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	SS EC20
	5	4	0	1.79	855	1386.65	6848.50	11106.97	.	.	EC20 POPULATION
	5	4	0	1.79	855	1386.65	6848.50	11106.97	9.24	74.05	NAWQ ACTUAL
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	NAWQ ADVISORY
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	LCV FISH
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	LCV DAPHNIDS
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	LTV EC20 FISH
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	LTV EC20 DAPHNID
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	SS EC20
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	70.28	70.28	EC20 POPULATION
	10	1	0	1.79	6500	10541.77	6500.00	10541.77	.	.	NAWQ ACTUAL
4,4'-DDD	2	2	0	6.02	71	0.01	1170.92	0.11	.	.	NAWQ ACTUAL
	2	2	0	6.02	71	0.01	1170.92	0.11	.	.	LCV DAPHNIDS
	2	2	0	6.02	71	0.01	1170.92	0.11	.	.	LTV EC20 DAPHNID
	2	2	0	6.02	71	0.01	1170.92	0.11	.	.	SS EC20
	2	2	0	6.02	71	0.01	1170.92	0.11	16.14	266.24	0.00 NAWQ ADVISORY
	2	2	0	6.02	71	0.01	1170.92	0.11	0.01	0.18	0.61 EC20 POPULATION
	2	2	0	6.02	71	0.01	1170.92	0.11	0.00	0.07	1.69 LCV FISH
	2	2	0	6.02	71	0.01	1170.92	0.11	0.00	0.03	3.99 LTV EC20 FISH
	3	2	0	6.02	280	0.03	2712.85	0.26	.	.	NAWQ ACTUAL
	3	2	0	6.02	280	0.03	2712.85	0.26	.	.	LCV DAPHNIDS
	3	2	0	6.02	280	0.03	2712.85	0.26	.	.	LTV EC20 DAPHNID
	3	2	0	6.02	280	0.03	2712.85	0.26	.	.	SS EC20
	3	2	0	6.02	280	0.03	2712.85	0.26	63.67	616.85	0.00 NAWQ ADVISORY
	3	2	0	6.02	280	0.03	2712.85	0.26	0.04	0.42	0.61 EC20 POPULATION
	3	2	0	6.02	280	0.03	2712.85	0.26	0.02	0.15	1.69 LCV FISH
	3	2	0	6.02	280	0.03	2712.85	0.26	0.01	0.06	3.99 LTV EC20 FISH
	3	2	0	6.02	280	0.03	2712.85	0.26	.	.	

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
4,4'-DDD (continued)	4	2	0	6.02	240	0.02	6750.45	0.64	.	.	.	NAHQ ACTUAL
	4	2	0	6.02	240	0.02	6750.45	0.64	.	.	.	LCV DAPHNIDS
	4	2	0	6.02	240	0.02	6750.45	0.64	.	.	.	LTV EC20 DAPHNID
	4	2	0	6.02	240	0.02	6750.45	0.64	.	.	.	SS EC20
	4	2	0	6.02	240	0.02	6750.45	0.64	54.57	1534.91	0.00	NAHQ ADVISORY
	4	2	0	6.02	240	0.02	6750.45	0.64	0.04	1.06	0.61	EC20 POPULATION
	4	2	0	6.02	240	0.02	6750.45	0.64	0.01	0.38	1.69	LCV FISH
	4	2	0	6.02	240	0.02	6750.45	0.64	0.01	0.16	3.99	LTV EC20 FISH
	4	2	0	6.02	240	0.02	6750.45	0.64	.	.	.	NAHQ ACTUAL
	5	4	0	6.02	258	0.02	1759.09	0.17	.	.	.	LCV DAPHNIDS
	5	4	0	6.02	258	0.02	1759.09	0.17	.	.	.	LTV EC20 DAPHNID
	5	4	0	6.02	258	0.02	1759.09	0.17	.	.	.	SS EC20
	5	4	0	6.02	258	0.02	1759.09	0.17	58.66	399.98	0.00	NAHQ ADVISORY
	5	4	0	6.02	258	0.02	1759.09	0.17	0.04	0.28	0.61	EC20 POPULATION
	5	4	0	6.02	258	0.02	1759.09	0.17	0.01	0.10	1.69	LCV FISH
	5	4	0	6.02	258	0.02	1759.09	0.17	0.01	0.04	3.99	LTV EC20 FISH
	5	4	0	6.02	258	0.02	1759.09	0.17	.	.	.	NAHQ ACTUAL
	10	1	0	6.02	1130	0.11	1130.00	0.11	.	.	.	LCV DAPHNIDS
	10	1	0	6.02	1130	0.11	1130.00	0.11	.	.	.	LTV EC20 DAPHNID
	10	1	0	6.02	1130	0.11	1130.00	0.11	.	.	.	SS EC20
4,4'-DDE	1	2	0	5.83	300	0.04	3418.16	0.51	256.94	256.94	0.00	NAHQ ADVISORY
	1	2	0	5.83	300	0.04	3418.16	0.51	0.18	0.18	0.61	EC20 POPULATION
	1	2	0	5.83	300	0.04	3418.16	0.51	0.06	0.06	1.69	LCV FISH
	1	2	0	5.83	300	0.04	3418.16	0.51	0.03	0.03	3.99	LTV EC20 FISH
	1	2	0	5.83	300	0.04	3418.16	0.51	.	.	.	NAHQ ACTUAL
	1	2	0	5.83	300	0.04	3418.16	0.51	.	.	.	LTV EC20 DAPHNID
	1	2	0	5.83	300	0.04	3418.16	0.51	.	.	.	EC20 POPULATION
	1	2	0	5.83	300	0.04	3418.16	0.51	5.55	63.20	0.01	SS EC20
	1	2	0	5.83	300	0.04	3418.16	0.51	2.77	31.60	0.02	LCV DAPHNIDS
	1	2	0	5.83	300	0.04	3418.16	0.51	2.22	25.28	0.02	NAHQ ADVISORY
	1	2	0	5.83	300	0.04	3418.16	0.51	0.13	1.44	0.35	LTV EC20 FISH
	1	2	0	5.83	300	0.04	3418.16	0.51	0.06	0.69	0.73	LCV FISH
	2	2	0	5.83	71	0.01	1170.92	0.17	.	.	.	NAHQ ACTUAL
	2	2	0	5.83	71	0.01	1170.92	0.17	.	.	.	LTV EC20 DAPHNID
	2	2	0	5.83	71	0.01	1170.92	0.17	.	.	.	EC20 POPULATION
	2	2	0	5.83	71	0.01	1170.92	0.17	1.31	21.65	0.01	SS EC20
	2	2	0	5.83	71	0.01	1170.92	0.17	0.66	10.82	0.02	LCV DAPHNIDS
	2	2	0	5.83	71	0.01	1170.92	0.17	0.53	8.66	0.02	NAHQ ADVISORY
	2	2	0	5.83	71	0.01	1170.92	0.17	0.03	0.49	0.35	LTV EC20 FISH
	2	2	0	5.83	71	0.01	1170.92	0.17	0.01	0.24	0.73	LCV FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
4,4'-DDE (continued)	3	2	0	5.83	280	0.04	2712.85	0.40	.	.	NAHQ ACTUAL
	3	2	0	5.83	280	0.04	2712.85	0.40	.	.	LTV EC20 DAPHNID
	3	2	0	5.83	280	0.04	2712.85	0.40	.	.	EC20 POPULATION
	3	2	0	5.83	280	0.04	2712.85	0.40	5.18	50.16	0.01 SS EC20
	3	2	0	5.83	280	0.04	2712.85	0.40	2.59	25.08	0.02 LCV DAPHNIDS
	3	2	0	5.83	280	0.04	2712.85	0.40	2.07	20.06	0.02 NAHQ ADVISORY
	3	2	0	5.83	280	0.04	2712.85	0.40	0.12	1.15	0.35 LTV EC20 FISH
	3	2	0	5.83	280	0.04	2712.85	0.40	0.06	0.55	0.73 LCV FISH
	3	2	0	5.83	280	0.04	2712.85	0.40	.	.	NAHQ ACTUAL
	4	2	0	5.83	240	0.04	6750.45	1.00	.	.	LTV EC20 DAPHNID
	4	2	0	5.83	240	0.04	6750.45	1.00	.	.	EC20 POPULATION
	4	2	0	5.83	240	0.04	6750.45	1.00	4.44	124.81	0.01 SS EC20
	4	2	0	5.83	240	0.04	6750.45	1.00	2.22	62.40	0.02 LCV DAPHNIDS
	4	2	0	5.83	240	0.04	6750.45	1.00	1.77	49.92	0.02 NAHQ ADVISORY
	4	2	0	5.83	240	0.04	6750.45	1.00	0.10	2.85	0.35 LTV EC20 FISH
	4	2	0	5.83	240	0.04	6750.45	1.00	0.05	1.37	0.73 LCV FISH
	4	2	0	5.83	258	0.04	1759.09	0.26	.	.	NAHQ ACTUAL
	5	4	0	5.83	258	0.04	1759.09	0.26	.	.	LTV EC20 DAPHNID
	5	4	0	5.83	258	0.04	1759.09	0.26	.	.	EC20 POPULATION
	5	4	0	5.83	258	0.04	1759.09	0.26	4.77	32.52	0.01 SS EC20
	5	4	0	5.83	258	0.04	1759.09	0.26	2.39	16.26	0.02 LCV DAPHNIDS
	5	4	0	5.83	258	0.04	1759.09	0.26	1.91	13.01	0.02 NAHQ ADVISORY
	5	4	0	5.83	258	0.04	1759.09	0.26	0.11	0.74	0.35 LTV EC20 FISH
	5	4	0	5.83	258	0.04	1759.09	0.26	0.05	0.36	0.73 LCV FISH
	5	4	0	5.83	258	0.04	1759.09	0.26	.	.	NAHQ ACTUAL
	10	1	0	5.83	1130	0.17	1130.00	0.17	.	.	LTV EC20 DAPHNID
	10	1	0	5.83	1130	0.17	1130.00	0.17	.	.	EC20 POPULATION
	10	1	0	5.83	1130	0.17	1130.00	0.17	20.89	20.89	0.01 SS EC20
	10	1	0	5.83	1130	0.17	1130.00	0.17	10.45	10.45	0.02 LCV DAPHNIDS
	10	1	0	5.83	1130	0.17	1130.00	0.17	8.36	8.36	0.02 NAHQ ADVISORY
	10	1	0	5.83	1130	0.17	1130.00	0.17	0.48	0.48	0.35 LTV EC20 FISH
	10	1	0	5.83	1130	0.17	1130.00	0.17	0.23	0.23	0.73 LCV FISH
4,4'-DDT	2	2	0	5.69	71	0.01	1170.92	0.24	.	.	NAHQ ACTUAL
	2	2	0	5.69	71	0.01	1170.92	0.24	.	.	LTV EC20 DAPHNID
	2	2	0	5.69	71	0.01	1170.92	0.24	.	.	EC20 POPULATION
	2	2	0	5.69	71	0.01	1170.92	0.24	1.81	29.88	0.01 SS EC20
	2	2	0	5.69	71	0.01	1170.92	0.24	0.91	14.94	0.02 LCV DAPHNIDS
	2	2	0	5.69	71	0.01	1170.92	0.24	0.72	11.95	0.02 NAHQ ADVISORY
	2	2	0	5.69	71	0.01	1170.92	0.24	0.04	0.68	0.35 LTV EC20 FISH
	2	2	0	5.69	71	0.01	1170.92	0.24	0.02	0.33	0.73 LCV FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
4,4'-DDT (continued)	3	2	0	5.69	280	0.06	2712.85	0.55	.	.	NAWQ ACTUAL
	3	2	0	5.69	280	0.06	2712.85	0.55	.	.	LTV EC20 DAPHNID
	3	2	0	5.69	280	0.06	2712.85	0.55	.	.	EC20 POPULATION
	3	2	0	5.69	280	0.06	2712.85	0.55	7.15	69.24	0.01 SS EC20
	3	2	0	5.69	280	0.06	2712.85	0.55	3.57	34.62	0.02 LCV DAPHNIDS
	3	2	0	5.69	280	0.06	2712.85	0.55	2.86	27.69	0.02 NAWQ ADVISORY
	3	2	0	5.69	280	0.06	2712.85	0.55	0.16	0.76	0.35 LTV EC20 FISH
	3	2	0	5.69	280	0.06	2712.85	0.55	0.08	.	0.73 LCV FISH
	3	2	0	5.69	280	0.06	2712.85	0.55	.	.	NAWQ ACTUAL
	3	2	0	5.69	280	0.05	6750.45	1.38	.	.	LTV EC20 DAPHNID
	4	2	0	5.69	240	0.05	6750.45	1.38	.	.	EC20 POPULATION
	4	2	0	5.69	240	0.05	6750.45	1.38	6.13	172.28	0.01 SS EC20
	4	2	0	5.69	240	0.05	6750.45	1.38	3.06	86.14	0.02 LCV DAPHNIDS
	4	2	0	5.69	240	0.05	6750.45	1.38	2.45	68.91	0.02 NAWQ ADVISORY
	4	2	0	5.69	240	0.05	6750.45	1.38	0.14	3.94	0.35 LTV EC20 FISH
	4	2	0	5.69	240	0.05	6750.45	1.38	0.07	1.89	0.73 LCV FISH
	4	2	0	5.69	240	0.05	6750.45	1.38	.	.	NAWQ ACTUAL
	4	2	0	5.69	258	0.05	759.09	0.36	.	.	LTV EC20 DAPHNID
	5	4	0	5.69	258	0.05	759.09	0.36	.	.	EC20 POPULATION
	5	4	0	5.69	258	0.05	759.09	0.36	6.58	44.90	0.01 SS EC20
	5	4	0	5.69	258	0.05	759.09	0.36	3.29	22.45	0.02 LCV DAPHNIDS
	5	4	0	5.69	258	0.05	759.09	0.36	2.63	17.96	0.02 NAWQ ADVISORY
	5	4	0	5.69	258	0.05	759.09	0.36	0.15	1.03	0.35 LTV EC20 FISH
	5	4	0	5.69	258	0.05	759.09	0.36	0.07	0.49	0.73 LCV FISH
	5	4	0	5.69	258	0.05	759.09	0.36	.	.	NAWQ ACTUAL
	5	4	0	5.69	258	0.23	1130.00	0.23	.	.	LTV EC20 DAPHNID
	10	1	0	5.69	1130	0.23	1130.00	0.23	.	.	EC20 POPULATION
	10	1	0	5.69	1130	0.23	1130.00	0.23	28.84	28.84	0.01 SS EC20
	10	1	0	5.69	1130	0.23	1130.00	0.23	14.42	14.42	0.02 LCV DAPHNIDS
	10	1	0	5.69	1130	0.23	1130.00	0.23	11.54	11.54	0.02 NAWQ ADVISORY
	10	1	0	5.69	1130	0.23	1130.00	0.23	0.66	0.66	0.35 LTV EC20 FISH
	10	1	0	5.69	1130	0.23	1130.00	0.23	0.32	0.32	0.73 LCV FISH
	10	1	0	5.69	1130	0.23	1130.00	0.23	.	.	NAWQ ACTUAL
ACENAPHTHENE	1	2	0	3.92	620	7.45	6993.39	84.08	.	.	LTV EC20 DAPHNID
	1	2	0	3.92	620	7.45	6993.39	84.08	.	.	SS EC20
	1	2	0	3.92	620	7.45	6993.39	84.08	.	.	EC20 POPULATION
	1	2	0	3.92	620	7.45	6993.39	84.08	3.39	38.22	2.20 NAWQ ADVISORY
	1	2	0	3.92	620	7.45	6993.39	84.08	0.04	0.43	197.00 LTV EC20 FISH
	1	2	0	3.92	620	7.45	6993.39	84.08	0.02	0.20	413.00 LCV FISH
	1	2	0	3.92	620	7.45	6993.39	84.08	0.00	0.01	6646.00 LCV DAPHNIDS
	1	2	0	3.92	620	7.45	6993.39	84.08	.	.	NAWQ ACTUAL
	1	2	0	3.92	620	7.45	6993.39	84.08	.	.	LTV EC20 DAPHNID
	1	2	0	3.92	620	7.45	6993.39	84.08	.	.	SS EC20

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H ₂ O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
ACENAPHTHENE (continued)	2	2	0	3.92	445	5.35	616.33	7.41	.	.	NAHQ ACTUAL
	2	2	0	3.92	445	5.35	616.33	7.41	.	.	LTV EC20 DAPHNID
	2	2	0	3.92	445	5.35	616.33	7.41	.	.	SS EC20
	2	2	0	3.92	445	5.35	616.33	7.41	.	.	EC20 POPULATION
	2	2	0	3.92	445	5.35	616.33	7.41	2.43	3.37	2.20 NAHQ ADVISORY
	2	2	0	3.92	445	5.35	616.33	7.41	0.03	0.04	197.00 LTV EC20 FISH
	2	2	0	3.92	445	5.35	616.33	7.41	0.01	0.02	413.00 LCV FISH
	2	2	0	3.92	445	5.35	616.33	7.41	0.00	0.00	6646.00 LCV DAPHNIDS
	2	2	0	3.92	445	5.35	616.33	7.41	.	.	NAHQ ACTUAL
	3	2	0	3.92	565	6.79	599.27	7.20	.	.	LTV EC20 DAPHNID
	3	2	0	3.92	565	6.79	599.27	7.20	.	.	SS EC20
	3	2	0	3.92	565	6.79	599.27	7.20	.	.	EC20 POPULATION
	3	2	0	3.92	565	6.79	599.27	7.20	3.09	3.27	2.20 NAHQ ADVISORY
	3	2	0	3.92	565	6.79	599.27	7.20	0.03	0.04	197.00 LTV EC20 FISH
	3	2	0	3.92	565	6.79	599.27	7.20	0.02	0.02	413.00 LCV FISH
	3	2	0	3.92	565	6.79	599.27	7.20	0.00	0.00	6646.00 LCV DAPHNIDS
	3	2	0	3.92	565	6.79	599.27	7.20	.	.	NAHQ ACTUAL
	4	2	0	3.92	500	6.01	38534.72	463.29	.	.	LTV EC20 DAPHNID
	4	2	0	3.92	500	6.01	38534.72	463.29	.	.	SS EC20
	4	2	0	3.92	500	6.01	38534.72	463.29	.	.	EC20 POPULATION
	4	2	0	3.92	500	6.01	38534.72	463.29	2.73	210.59	2.20 NAHQ ADVISORY
	4	2	0	3.92	500	6.01	38534.72	463.29	0.03	2.35	197.00 LTV EC20 FISH
	4	2	0	3.92	500	6.01	38534.72	463.29	0.01	1.12	413.00 LCV FISH
	4	2	0	3.92	500	6.01	38534.72	463.29	0.00	0.07	6646.00 LCV DAPHNIDS
	4	2	0	3.92	500	6.01	38534.72	463.29	.	.	NAHQ ACTUAL
	5	4	0	3.92	855	10.28	6848.50	82.34	.	.	LTV EC20 DAPHNID
	5	4	0	3.92	855	10.28	6848.50	82.34	.	.	SS EC20
	5	4	0	3.92	855	10.28	6848.50	82.34	.	.	EC20 POPULATION
	5	4	0	3.92	855	10.28	6848.50	82.34	4.67	37.43	2.20 NAHQ ADVISORY
	5	4	0	3.92	855	10.28	6848.50	82.34	0.05	0.42	197.00 LTV EC20 FISH
	5	4	0	3.92	855	10.28	6848.50	82.34	0.02	0.20	413.00 LCV FISH
	5	4	0	3.92	855	10.28	6848.50	82.34	0.00	0.01	6646.00 LCV DAPHNIDS
	5	4	0	3.92	855	10.28	6848.50	82.34	.	.	NAHQ ACTUAL
	10	1	0	3.92	6500	78.15	6500.00	78.15	.	.	LTV EC20 DAPHNID
	10	1	0	3.92	6500	78.15	6500.00	78.15	.	.	SS EC20
	10	1	0	3.92	6500	78.15	6500.00	78.15	.	.	EC20 POPULATION
	10	1	0	3.92	6500	78.15	6500.00	78.15	35.52	35.52	2.20 NAHQ ADVISORY
	10	1	0	3.92	6500	78.15	6500.00	78.15	0.40	0.40	197.00 LTV EC20 FISH
	10	1	0	3.92	6500	78.15	6500.00	78.15	0.19	0.19	413.00 LCV FISH
	10	1	0	3.92	6500	78.15	6500.00	78.15	0.01	0.01	6646.00 LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
ALDRIN	1	2	0	3.01	150	14.66	1726.21	168.69	195.45	2249.23	0.08 NAHQ ADVISORY
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	NAHQ ACTUAL
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	LCV FISH
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	LCV DAPHNIDS
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	LTV EC20 FISH
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	LTV EC20 DAPHNID
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	SS EC20
	1	2	0	3.01	150	14.66	1726.21	168.69	.	.	EC20 POPULATION
	2	2	0	3.01	35	3.42	590.10	57.67	45.60	768.89	0.08 NAHQ ADVISORY
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	NAHQ ACTUAL
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	LCV FISH
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	LCV DAPHNIDS
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	LTV EC20 FISH
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	LTV EC20 DAPHNID
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	SS EC20
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	EC20 POPULATION
	2	2	0	3.01	35	3.42	590.10	57.67	.	.	NAHQ ADVISORY
	3	2	0	3.01	135	13.19	1402.82	137.09	175.90	1827.86	0.08 NAHQ ACTUAL
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	LCV FISH
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	LCV DAPHNIDS
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	LTV EC20 FISH
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	LTV EC20 DAPHNID
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	SS EC20
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	EC20 POPULATION
	3	2	0	3.01	135	13.19	1402.82	137.09	.	.	NAHQ ADVISORY
	4	2	0	3.01	120	11.73	4780.11	467.13	156.36	6228.40	0.08 NAHQ ACTUAL
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	NAHQ ACTUAL
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	LCV FISH
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	LCV DAPHNIDS
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	LTV EC20 FISH
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	LTV EC20 DAPHNID
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	SS EC20
	4	2	0	3.01	120	11.73	4780.11	467.13	.	.	EC20 POPULATION
	5	2	0	3.01	129	12.61	1232.35	120.43	168.08	1605.73	0.08 NAHQ ADVISORY
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	NAHQ ACTUAL
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	LCV FISH
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	LCV DAPHNIDS
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	LTV EC20 FISH
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	LTV EC20 DAPHNID
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	SS EC20
	5	2	0	3.01	129	12.61	1232.35	120.43	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between M20 and SED	Minimum detection pore water limit	Minimum water conc.	UCL ^a detection pore water limit	UCL conc.	MDL quotient	UCL quotient	Benchmark type
ANTHRACENE	1	2	0	4.45	620	2.20	6993.39	24.81	.	.	NAHQ ACTUAL
	1	2	0	4.45	620	2.20	6993.39	24.81	.	.	SS EC20
	1	2	0	4.45	620	2.20	6993.39	24.81	.	.	EC20 POPULATION
	1	2	0	4.45	620	2.20	6993.39	24.81	219984.30	2481346.71	0.09 LCV FISH
	1	2	0	4.45	620	2.20	6993.39	24.81	24.44	275.71	0.35 LTV EC20 FISH
	1	2	0	4.45	620	2.20	6993.39	24.81	6.29	70.90	2.10 LCV DAPHNIDS
	1	2	0	4.45	620	2.20	6993.39	24.81	1.05	11.82	8.20 LTV EC20 DAPHNID
	1	2	0	4.45	620	2.20	6993.39	24.81	0.27	3.03	NAHQ ACTUAL
	2	2	0	4.45	445	1.58	616.33	2.19	.	.	SS EC20
	2	2	0	4.45	445	1.58	616.33	2.19	.	.	EC20 POPULATION
	2	2	0	4.45	445	1.58	616.33	2.19	157891.96	218681.27	0.09 LCV FISH
	2	2	0	4.45	445	1.58	616.33	2.19	17.54	24.30	0.35 LTV EC20 FISH
	2	2	0	4.45	445	1.58	616.33	2.19	4.51	6.25	2.10 LCV DAPHNIDS
	2	2	0	4.45	445	1.58	616.33	2.19	0.75	1.04	8.20 LTV EC20 DAPHNID
	2	2	0	4.45	445	1.58	616.33	2.19	0.19	0.27	NAHQ ACTUAL
	3	2	0	4.45	565	2.00	599.27	2.13	.	.	SS EC20
	3	2	0	4.45	565	2.00	599.27	2.13	.	.	EC20 POPULATION
	3	2	0	4.45	565	2.00	599.27	2.13	200469.56	212627.43	0.09 LCV FISH
	3	2	0	4.45	565	2.00	599.27	2.13	22.27	23.63	0.35 LTV EC20 FISH
	3	2	0	4.45	565	2.00	599.27	2.13	5.73	6.08	2.10 LCV DAPHNIDS
	3	2	0	4.45	565	2.00	599.27	2.13	0.95	1.01	8.20 LTV EC20 DAPHNID
	3	2	0	4.45	565	2.00	599.27	2.13	0.24	0.26	NAHQ ACTUAL
	4	2	0	4.45	500	1.77	38534.72	136.73	.	.	SS EC20
	4	2	0	4.45	500	1.77	38534.72	136.73	.	.	EC20 POPULATION
	4	2	0	4.45	500	1.77	38534.72	136.73	177406.69	13672633.95	0.09 LCV FISH
	4	2	0	4.45	500	1.77	38534.72	136.73	19.71	1519.18	0.35 LTV EC20 FISH
	4	2	0	4.45	500	1.77	38534.72	136.73	5.07	390.65	2.10 LCV DAPHNIDS
	4	2	0	4.45	500	1.77	38534.72	136.73	0.84	65.11	8.20 LTV EC20 DAPHNID
	4	2	0	4.45	500	1.77	38534.72	136.73	0.22	16.67	NAHQ ACTUAL
	5	4	0	4.45	855	3.03	6848.50	24.30	.	.	SS EC20
	5	4	0	4.45	855	3.03	6848.50	24.30	.	.	EC20 POPULATION
	5	4	0	4.45	855	3.03	6848.50	24.30	303365.45	2429940.45	0.09 LCV FISH
	5	4	0	4.45	855	3.03	6848.50	24.30	33.71	269.99	0.35 LTV EC20 FISH
	5	4	0	4.45	855	3.03	6848.50	24.30	8.67	69.43	2.10 LCV DAPHNIDS
	5	4	0	4.45	855	3.03	6848.50	24.30	1.44	11.57	8.20 LTV EC20 DAPHNID
	5	4	0	4.45	855	3.03	6848.50	24.30	0.37	2.96	

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between h20 and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
ANTHRACENE (continued)	10	1	0	4.45	6500	23.06	6500.00	23.06	.	.	NAHQ ACTUAL
	10	1	0	4.45	6500	23.06	6500.00	23.06	.	.	SS EC20
	10	1	0	4.45	6500	23.06	6500.00	23.06	.	.	EC20 POPULATION
	10	1	0	4.45	6500	23.06	6500.00	23.06	2306287.03	2306287.03	0.00 NAHQ ADVISORY
	10	1	0	4.45	6500	23.06	6500.00	23.06	256.25	256.25	0.09 LCV FISH
	10	1	0	4.45	6500	23.06	6500.00	23.06	65.89	65.89	0.35 LTV EC20 FISH
	10	1	0	4.45	6500	23.06	6500.00	23.06	10.98	10.98	2.10 LCV DAPHNIDS
	10	1	0	4.45	6500	23.06	6500.00	23.06	2.81	2.81	8.20 LTV EC20 DAPHNID
	10	1	0	4.45	6500	23.06	6500.00	23.06	.	.	NAHQ ACTUAL
	10	1	0	4.45	6500	23.06	6500.00	23.06	.	.	SS EC20
AROCLOL-1221	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	LCV DAPHNIDS
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	LTV EC20 DAPHNID
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	SS EC20
	1	2	0	7.14	1500	0.01	17262.14	0.13	3.75	43.12	0.00 NAHQ ADVISORY
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.01	10.00 EC20 POPULATION
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.00	60.00 LCV FISH
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.00	80.00 LTV EC20 FISH
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	NAHQ ACTUAL
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	LCV DAPHNIDS
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	LTV EC20 DAPHNID
	2	2	0	7.14	325	0.00	6047.34	0.04	0.81	15.11	0.00 NAHQ ADVISORY
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	10.00 EC20 POPULATION
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	60.00 LCV FISH
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	80.00 LTV EC20 FISH
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	NAHQ ACTUAL
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	LCV DAPHNIDS
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	LTV EC20 DAPHNID
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	SS EC20
	3	2	0	7.14	1350	0.01	14028.24	0.10	3.37	35.04	0.00 NAHQ ADVISORY
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.01	10.00 EC20 POPULATION
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.00	60.00 LCV FISH
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.00	80.00 LTV EC20 FISH
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	NAHQ ACTUAL
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	LCV DAPHNIDS
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	LTV EC20 DAPHNID
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	SS EC20
	4	2	0	7.14	1200	0.01	47801.10	0.35	3.00	119.41	0.00 NAHQ ADVISORY
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.03	10.00 EC20 POPULATION
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.01	60.00 LCV FISH
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.00	80.00 LTV EC20 FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
AROCLOL-1221 (continued)	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	NAWQ ACTUAL
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	LCV DAPHNIDS
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	LTV EC20 DAPHNID
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	SS EC20
	5	2	0	7.14	1290	0.01	12323.49	0.09	3.22	30.78	0.00 NAWQ ADVISORY
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.01	10.00 EC20 POPULATION
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00	60.00 LCV FISH
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00	80.00 LTV EC20 FISH
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00	
AROCLOL-1232	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	NAWQ ACTUAL
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	LCV DAPHNIDS
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	LTV EC20 DAPHNID
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	SS EC20
	1	2	0	7.14	1500	0.01	17262.14	0.13	1.87	21.56	0.01 NAWQ ADVISORY
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.01	16.00 EC20 POPULATION
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.00	124.00 LCV FISH
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.00	148.00 LTV EC20 FISH
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.00	NAWQ ACTUAL
	1	2	0	7.14	325	0.00	6047.34	0.04	.	.	LCV DAPHNIDS
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	LTV EC20 DAPHNID
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	SS EC20
	2	2	0	7.14	325	0.00	6047.34	0.04	0.41	7.55	0.01 NAWQ ADVISORY
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	16.00 EC20 POPULATION
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	124.00 LCV FISH
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	148.00 LTV EC20 FISH
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	NAWQ ACTUAL
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	LCV DAPHNIDS
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	LTV EC20 DAPHNID
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	SS EC20
	3	2	0	7.14	1350	0.01	14028.24	0.10	1.69	17.52	0.01 NAWQ ADVISORY
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.01	16.00 EC20 POPULATION
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.00	124.00 LCV FISH
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.00	148.00 LTV EC20 FISH
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.00	NAWQ ACTUAL
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	LCV DAPHNIDS
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	LTV EC20 DAPHNID
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	SS EC20
	4	2	0	7.14	1200	0.01	47801.10	0.35	1.50	59.70	0.01 NAWQ ADVISORY
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.02	16.00 EC20 POPULATION
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.00	124.00 LCV FISH
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.00	148.00 LTV EC20 FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
AROCFLOR-1232 (continued)	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	NAWQ ACTUAL	.
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	LCV DAPHNIDS	.
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	LTV EC20 DAPHNID	.
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	SS EC20	.
	5	2	0	7.14	1290	0.01	12323.49	0.09	1.61	15.39	0.01 NAWQ ADVISORY	0.01
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.01	16.00 EC20 POPULATION	0.01
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00	124.00 LCV FISH	0.00
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00	148.00 LTV EC20 FISH	0.00
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00		
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.00		
AROCFLOR-1242	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	NAWQ ACTUAL	.
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	LCV DAPHNIDS	.
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	LTV EC20 DAPHNID	.
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	SS EC20	.
	1	2	0	7.14	1500	0.01	17262.14	0.13	1.09	12.51	0.01 NAWQ ADVISORY	0.01
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.01	0.08	1.58 EC20 POPULATION	0.01
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.04	2.90 LTV EC20 FISH	0.00
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.01	9.00 LCV FISH	0.00
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	NAWQ ACTUAL	.
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	LCV DAPHNIDS	.
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	LTV EC20 DAPHNID	.
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	SS EC20	.
	2	2	0	7.14	325	0.00	6047.34	0.04	0.24	4.38	0.01 NAWQ ADVISORY	0.01
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.03	1.58 EC20 POPULATION	0.00
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.02	2.90 LTV EC20 FISH	0.00
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.00	9.00 LCV FISH	0.00
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	NAWQ ACTUAL	.
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	LCV DAPHNIDS	.
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	LTV EC20 DAPHNID	.
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	SS EC20	.
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.98	10.16	0.01 NAWQ ADVISORY	0.01
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.01	0.06	1.58 EC20 POPULATION	0.01
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.04	2.90 LTV EC20 FISH	0.00
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.01	9.00 LCV FISH	0.00
	3	2	0	7.14	1200	0.01	47801.10	0.35	.	.	NAWQ ACTUAL	.
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	LCV DAPHNIDS	.
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	LTV EC20 DAPHNID	.
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	SS EC20	.
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.87	34.63	0.01 NAWQ ADVISORY	0.01
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.01	0.22	1.58 EC20 POPULATION	0.01
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.12	2.90 LTV EC20 FISH	0.00
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.04	9.00 LCV FISH	0.00

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between #20 and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
AROCLOL-1242 (continued)	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	NAHQ ACTUAL
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	LCV DAPHNIDS
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	LTV EC20 DAPHNID
	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	SS EC20
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.93	8.93	0.01 NAHQ ADVISORY
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.01	0.06	1.58 EC20 POPULATION
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.03	2.90 LTV EC20 FISH
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.01	9.00 LCV FISH
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.01	
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.01	
AROCLOL-1248	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	NAHQ ACTUAL
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	SS EC20
	1	2	0	7.14	1500	0.01	17262.14	0.13	54.33	625.27	0.00 NAHQ ADVISORY
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.05	0.63	0.20 LCV FISH
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.03	0.31	0.40 LTV EC20 FISH
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.01	0.10	1.26 EC20 POPULATION
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.05	2.50 LTV EC20 DAPHNID
	1	2	0	7.14	1500	0.01	17262.14	0.13	0.00	0.03	4.30 LCV DAPHNIDS
	1	2	0	7.14	1500	0.01	17262.14	0.13	.	.	NAHQ ACTUAL
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	SS EC20
	2	2	0	7.14	325	0.00	6047.34	0.04	11.77	219.05	0.00 NAHQ ADVISORY
	2	2	0	7.14	325	0.00	6047.34	0.04	0.01	0.22	0.20 LCV FISH
	2	2	0	7.14	325	0.00	6047.34	0.04	0.01	0.11	0.40 LTV EC20 FISH
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.03	1.26 EC20 POPULATION
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.02	2.50 LTV EC20 DAPHNID
	2	2	0	7.14	325	0.00	6047.34	0.04	0.00	0.01	4.30 LCV DAPHNIDS
	2	2	0	7.14	325	0.00	6047.34	0.04	.	.	NAHQ ACTUAL
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	SS EC20
	3	2	0	7.14	1350	0.01	14028.24	0.10	48.90	508.13	0.00 NAHQ ADVISORY
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.05	0.51	0.20 LCV FISH
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.02	0.25	0.40 LTV EC20 FISH
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.01	0.08	1.26 EC20 POPULATION
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.04	2.50 LTV EC20 DAPHNID
	3	2	0	7.14	1350	0.01	14028.24	0.10	0.00	0.02	4.30 LCV DAPHNIDS
	3	2	0	7.14	1350	0.01	14028.24	0.10	.	.	NAHQ ACTUAL
	4	2	0	7.14	1200	0.01	47801.10	0.35	.	.	SS EC20
	4	2	0	7.14	1200	0.01	47801.10	0.35	43.47	1731.44	0.00 NAHQ ADVISORY
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.04	1.73	0.20 LCV FISH
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.02	0.87	0.40 LTV EC20 FISH
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.01	0.27	1.26 EC20 POPULATION
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.14	2.50 LTV EC20 DAPHNID
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.08	4.30 LCV DAPHNIDS
	4	2	0	7.14	1200	0.01	47801.10	0.35	0.00	0.08	

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
AROCLOP-1248 (continued)	5	2	0	7.14	1290	0.01	12323.49	0.09	.	.	NAHQ ACTUAL
	5	2	0	7.14	1290	0.01	12323.49	0.09	46.73	446.38	SS EC20
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.05	0.45	0.00 NAHQ ADVISORY
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.02	0.22	0.20 LCV FISH
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.01	0.07	0.40 LTV EC20 FISH
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.04	1.26 EC20 POPULATION
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.02	2.50 LTV EC20 DAPHNID
	5	2	0	7.14	1290	0.01	12323.49	0.09	0.00	0.02	4.30 LCV DAPHNIDS
	1	2	0	7.14	3000	0.02	34181.62	0.25	.	.	NAHQ ACTUAL
	1	2	0	7.14	3000	0.02	34181.62	0.25	1.09	12.38	SS EC20
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.04	0.48	0.02 NAHQ ADVISORY
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.03	0.39	0.52 LTV EC20 FISH
AROCLOP-1254	1	2	0	7.14	3000	0.02	34181.62	0.25	0.02	0.25	0.63 EC20 POPULATION
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.02	0.21	1.00 LCV FISH
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.02	0.12	1.20 LTV EC20 DAPHNID
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.01	0.12	2.10 LCV DAPHNIDS
	5	4	0	7.14	2580	0.02	17590.93	0.13	.	.	NAHQ ACTUAL
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.93	6.37	SS EC20
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.04	0.25	0.02 NAHQ ADVISORY
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.03	0.20	0.52 LTV EC20 FISH
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.03	0.13	0.63 EC20 POPULATION
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.02	0.11	1.00 LCV FISH
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.01	0.06	1.20 LTV EC20 DAPHNID
	10	1	0	7.14	11300	0.08	11300.00	0.08	.	.	NAHQ ACTUAL
	10	1	0	7.14	11300	0.08	11300.00	0.08	4.09	4.09	SS EC20
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.16	0.16	0.02 NAHQ ADVISORY
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.13	0.13	0.52 LTV EC20 FISH
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.08	0.08	0.63 EC20 POPULATION
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.08	0.08	1.00 LCV FISH
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.07	0.07	1.20 LTV EC20 DAPHNID
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.04	0.04	2.10 LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
AROCLOP-1260	1	2	0	7.14	3000	0.02	34181.62	0.25	.	.	NAHQ ACTUAL
	1	2	0	7.14	3000	0.02	34181.62	0.25	.	.	LCV DAPHNIDS
	1	2	0	7.14	3000	0.02	34181.62	0.25	.	.	LTV EC20 DAPHNID
	1	2	0	7.14	3000	0.02	34181.62	0.25	.	.	SS EC20
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.20	2.25	0.11 NAHQ ADVISORY
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.01	0.12	2.10 LTV EC20 FISH
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.01	0.11	2.30 LCV FISH
	1	2	0	7.14	3000	0.02	34181.62	0.25	0.00	0.00	316.00 EC20 POPULATION
	1	2	0	7.14	3000	0.02	34181.62	0.20	.	.	NAHQ ACTUAL
	3	2	0	7.14	2800	0.02	27128.51	0.20	.	.	LCV DAPHNIDS
	3	2	0	7.14	2800	0.02	27128.51	0.20	.	.	LTV EC20 DAPHNID
	3	2	0	7.14	2800	0.02	27128.51	0.20	.	.	SS EC20
	3	2	0	7.14	2800	0.02	27128.51	0.20	0.18	1.79	0.11 NAHQ ADVISORY
	3	2	0	7.14	2800	0.02	27128.51	0.20	0.01	0.09	2.10 LTV EC20 FISH
	3	2	0	7.14	2800	0.02	27128.51	0.20	0.01	0.09	2.30 LCV FISH
	3	2	0	7.14	2800	0.02	27128.51	0.20	0.00	0.00	316.00 EC20 POPULATION
	3	2	0	7.14	2800	0.02	27128.51	0.20	.	.	NAHQ ACTUAL
	4	2	0	7.14	2400	0.02	67504.47	0.49	.	.	LCV DAPHNIDS
	4	2	0	7.14	2400	0.02	67504.47	0.49	.	.	LTV EC20 DAPHNID
	4	2	0	7.14	2400	0.02	67504.47	0.49	.	.	SS EC20
	4	2	0	7.14	2400	0.02	67504.47	0.49	0.16	4.45	0.11 NAHQ ADVISORY
	4	2	0	7.14	2400	0.02	67504.47	0.49	0.01	0.23	2.10 LTV EC20 FISH
	4	2	0	7.14	2400	0.02	67504.47	0.49	0.01	0.21	2.30 LCV FISH
	4	2	0	7.14	2400	0.02	67504.47	0.49	0.00	0.00	316.00 EC20 POPULATION
	4	2	0	7.14	2400	0.02	67504.47	0.49	.	.	NAHQ ACTUAL
	5	4	0	7.14	2580	0.02	17590.93	0.13	.	.	LCV DAPHNIDS
	5	4	0	7.14	2580	0.02	17590.93	0.13	.	.	LTV EC20 DAPHNID
	5	4	0	7.14	2580	0.02	17590.93	0.13	.	.	SS EC20
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.17	1.16	0.11 NAHQ ADVISORY
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.01	0.06	2.10 LTV EC20 FISH
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.01	0.06	2.30 LCV FISH
	5	4	0	7.14	2580	0.02	17590.93	0.13	0.00	0.00	316.00 EC20 POPULATION
	5	4	0	7.14	2580	0.02	17590.93	0.08	.	.	NAHQ ACTUAL
	10	1	0	7.14	11300	0.08	11300.00	0.08	.	.	LCV DAPHNIDS
	10	1	0	7.14	11300	0.08	11300.00	0.08	.	.	LTV EC20 DAPHNID
	10	1	0	7.14	11300	0.08	11300.00	0.08	.	.	SS EC20
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.74	0.74	0.11 NAHQ ADVISORY
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.04	0.04	2.10 LTV EC20 FISH
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.04	0.04	2.30 LCV FISH
	10	1	0	7.14	11300	0.08	11300.00	0.08	0.00	0.00	316.00 EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BENZO(A)ANTHRACENE	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	NAWQ ACTUAL
	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	LCV FISH
	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	LTV EC20 FISH
	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	LTV EC20 DAPHNID
	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	SS EC20
	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	EC20 POPULATION
	1	2	0	5.61	620	0.15	6993.39	1.72	56.37	635.80	0.00 NAWQ ADVISORY
	1	2	0	5.61	620	0.15	6993.39	1.72	0.23	2.64	0.65 LCV DAPHNIDS
	1	2	0	5.61	620	0.15	6993.39	1.72	-	-	NAWQ ACTUAL
	2	2	0	5.61	445	0.11	616.33	0.15	-	-	LCV FISH
	2	2	0	5.61	445	0.11	616.33	0.15	-	-	LTV EC20 FISH
	2	2	0	5.61	445	0.11	616.33	0.15	-	-	LTV EC20 DAPHNID
	2	2	0	5.61	445	0.11	616.33	0.15	-	-	SS EC20
	2	2	0	5.61	445	0.11	616.33	0.15	-	-	EC20 POPULATION
	2	2	0	5.61	445	0.11	616.33	0.15	40.46	56.03	0.00 NAWQ ADVISORY
	2	2	0	5.61	445	0.11	616.33	0.15	0.17	0.23	0.65 LCV DAPHNIDS
	2	2	0	5.61	565	0.14	599.27	0.15	-	-	NAWQ ACTUAL
	3	2	0	5.61	565	0.14	599.27	0.15	-	-	LCV FISH
	3	2	0	5.61	565	0.14	599.27	0.15	-	-	LTV EC20 FISH
	3	2	0	5.61	565	0.14	599.27	0.15	-	-	LTV EC20 DAPHNID
	3	2	0	5.61	565	0.14	599.27	0.15	-	-	SS EC20
	3	2	0	5.61	565	0.14	599.27	0.15	-	-	EC20 POPULATION
	3	2	0	5.61	565	0.14	599.27	0.15	51.37	54.48	0.00 NAWQ ADVISORY
	3	2	0	5.61	565	0.14	599.27	0.15	0.21	0.23	0.65 LCV DAPHNIDS
	3	2	0	5.61	565	0.14	599.27	0.15	-	-	NAWQ ACTUAL
	4	2	0	5.61	500	0.12	38534.72	9.46	-	-	LCV FISH
	4	2	0	5.61	500	0.12	38534.72	9.46	-	-	LTV EC20 FISH
	4	2	0	5.61	500	0.12	38534.72	9.46	-	-	LTV EC20 DAPHNID
	4	2	0	5.61	500	0.12	38534.72	9.46	-	-	SS EC20
	4	2	0	5.61	500	0.12	38534.72	9.46	-	-	EC20 POPULATION
	4	2	0	5.61	500	0.12	38534.72	9.46	45.46	3503.39	0.00 NAWQ ADVISORY
	4	2	0	5.61	500	0.12	38534.72	9.46	0.19	14.55	0.65 LCV DAPHNIDS
	4	2	0	5.61	855	0.21	6848.50	1.68	-	-	NAWQ ACTUAL
	5	4	0	5.61	855	0.21	6848.50	1.68	-	-	LCV FISH
	5	4	0	5.61	855	0.21	6848.50	1.68	-	-	LTV EC20 FISH
	5	4	0	5.61	855	0.21	6848.50	1.68	-	-	LTV EC20 DAPHNID
	5	4	0	5.61	855	0.21	6848.50	1.68	-	-	SS EC20
	5	4	0	5.61	855	0.21	6848.50	1.68	-	-	EC20 POPULATION
	5	4	0	5.61	855	0.21	6848.50	1.68	77.73	622.63	0.00 NAWQ ADVISORY
	5	4	0	5.61	855	0.21	6848.50	1.68	0.32	2.59	0.65 LCV DAPHNIDS
	5	4	0	5.61	855	0.21	6848.50	1.68	-	-	NAWQ ACTUAL

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL* detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BENZO(A)ANTHRACENE (continued)	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	NAWQ ACTUAL
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	LCV FISH
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	LTV EC20 FISH
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	LTV EC20 DAPHNID
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	SS EC20
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	EC20 POPULATION
	10	1	0	5.61	6500	1.60	6500.00	1.60	590.95	590.95	0.00 NAWQ ADVISORY
	10	1	0	5.61	6500	1.60	6500.00	1.60	2.45	2.45	0.65 LCV DAPHNIDS
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	NAWQ ACTUAL
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	LCV FISH
	10	1	0	5.61	6500	1.60	6500.00	1.60	.	.	LTV EC20 DAPHNID
BENZO(A)PYRENE	1	2	0	6.04	620	0.06	6993.39	0.64	.	.	NAWQ ACTUAL
	1	2	0	6.04	620	0.06	6993.39	0.64	.	.	LCV FISH
	1	2	0	6.04	620	0.06	6993.39	0.64	.	.	LTV EC20 DAPHNID
	1	2	0	6.04	620	0.06	6993.39	0.64	.	.	SS EC20
	1	2	0	6.04	620	0.06	6993.39	0.64	.	.	EC20 POPULATION
	1	2	0	6.04	620	0.06	6993.39	0.64	43.50	490.62	0.00 NAWQ ADVISORY
	1	2	0	6.04	620	0.06	6993.39	0.64	0.19	2.13	0.30 LCV DAPHNIDS
	1	2	0	6.04	620	0.06	6993.39	0.64	0.02	0.21	2.99 LTV EC20 FISH
	1	2	0	6.04	445	0.04	616.33	0.06	.	.	NAWQ ACTUAL
	2	2	0	6.04	445	0.04	616.33	0.06	.	.	LCV FISH
	2	2	0	6.04	445	0.04	616.33	0.06	.	.	LTV EC20 DAPHNID
	2	2	0	6.04	445	0.04	616.33	0.06	.	.	SS EC20
	2	2	0	6.04	445	0.04	616.33	0.06	.	.	EC20 POPULATION
	2	2	0	6.04	445	0.04	616.33	0.06	31.22	43.24	0.00 NAWQ ADVISORY
	2	2	0	6.04	445	0.04	616.33	0.06	0.14	0.19	0.30 LCV DAPHNIDS
	2	2	0	6.04	445	0.04	616.33	0.06	0.01	0.02	2.99 LTV EC20 FISH
	3	2	0	6.04	565	0.05	599.27	0.05	.	.	NAWQ ACTUAL
	3	2	0	6.04	565	0.05	599.27	0.05	.	.	LCV FISH
	3	2	0	6.04	565	0.05	599.27	0.05	.	.	LTV EC20 DAPHNID
	3	2	0	6.04	565	0.05	599.27	0.05	.	.	SS EC20
	3	2	0	6.04	565	0.05	599.27	0.05	.	.	EC20 POPULATION
	3	2	0	6.04	565	0.05	599.27	0.05	39.64	42.04	0.00 NAWQ ADVISORY
	3	2	0	6.04	565	0.05	599.27	0.05	0.17	0.18	0.30 LCV DAPHNIDS
	3	2	0	6.04	565	0.05	599.27	0.05	0.02	0.02	2.99 LTV EC20 FISH
	3	2	0	6.04	565	0.05	599.27	0.05	.	.	NAWQ ACTUAL
	4	2	0	6.04	500	0.05	38534.72	3.51	.	.	LCV FISH
	4	2	0	6.04	500	0.05	38534.72	3.51	.	.	LTV EC20 DAPHNID
	4	2	0	6.04	500	0.05	38534.72	3.51	.	.	SS EC20
	4	2	0	6.04	500	0.05	38534.72	3.51	.	.	EC20 POPULATION
	4	2	0	6.04	500	0.05	38534.72	3.51	35.08	2703.39	0.00 NAWQ ADVISORY
	4	2	0	6.04	500	0.05	38534.72	3.51	0.15	11.71	0.30 LCV DAPHNIDS
	4	2	0	6.04	500	0.05	38534.72	3.51	0.02	1.18	2.99 LTV EC20 FISH

Table F.10 (continued)

Analyte	Reach	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection pore water limit	Minimum conc.	UCL ^a detection pore water limit	UCL conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
BENZO(A)PYRENE (continued)	5	4	0	6.04	855	0.08	6848.50	0.62	.	.	NAHQ ACTUAL	.
	5	4	0	6.04	855	0.08	6848.50	0.62	.	.	LCV FISH	.
	5	4	0	6.04	855	0.08	6848.50	0.62	.	.	LTV EC20 DAPHNID	.
	5	4	0	6.04	855	0.08	6848.50	0.62	.	.	SS EC20	.
	5	4	0	6.04	855	0.08	6848.50	0.62	.	.	EC20 POPULATION	.
	5	4	0	6.04	855	0.08	6848.50	0.62	59.98	480.45	0.00 NAHQ ADVISORY	0.00
	5	4	0	6.04	855	0.08	6848.50	0.62	0.26	2.08	0.30 LCV DAPHNIDS	0.30
	5	4	0	6.04	855	0.08	6848.50	0.62	0.03	0.21	2.99 LTV EC20 FISH	2.99
	5	4	0	6.04	855	0.08	6848.50	0.62	.	.	NAHQ ACTUAL	.
	10	1	0	6.04	6500	0.59	6500.00	0.59	.	.	LCV FISH	.
	10	1	0	6.04	6500	0.59	6500.00	0.59	.	.	LTV EC20 DAPHNID	.
	10	1	0	6.04	6500	0.59	6500.00	0.59	.	.	SS EC20	.
	10	1	0	6.04	6500	0.59	6500.00	0.59	.	.	EC20 POPULATION	.
	10	1	0	6.04	6500	0.59	6500.00	0.59	456.01	456.01	0.00 NAHQ ADVISORY	0.00
	10	1	0	6.04	6500	0.59	6500.00	0.59	1.98	1.98	0.30 LCV DAPHNIDS	0.30
	10	1	0	6.04	6500	0.59	6500.00	0.59	0.20	0.20	2.99 LTV EC20 FISH	2.99
	10	1	0	6.04	6500	0.59	6500.00	0.59	.	.	NAHQ ACTUAL	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	NAHQ ADVISORY	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	NAHQ ACTUAL	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LCV FISH	.
BENZO(B)FLUORANTHENE	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LCV DAPHNIDS	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LTV EC20 FISH	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LTV EC20 DAPHNID	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	SS EC20	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	EC20 POPULATION	.
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	NAHQ ADVISORY	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	NAHQ ACTUAL	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LCV FISH	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LCV DAPHNIDS	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LTV EC20 FISH	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LTV EC20 DAPHNID	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	SS EC20	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	EC20 POPULATION	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	NAHQ ADVISORY	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	NAHQ ACTUAL	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LCV FISH	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LCV DAPHNIDS	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LTV EC20 FISH	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LTV EC20 DAPHNID	.
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	SS EC20	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	EC20 POPULATION	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	NAHQ ADVISORY	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	NAHQ ACTUAL	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	LCV FISH	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	LCV DAPHNIDS	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	LTV EC20 FISH	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	LTV EC20 DAPHNID	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	SS EC20	.
	3	3	0	6.84	565	0.01	599.27	0.01	.	.	EC20 POPULATION	.

Table F.10 (continued)

Analyte	Resch (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between W20 and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BENZO(B)FLUORANTHENE (continued)	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	NAWQ ADVISORY
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	NAWQ ACTUAL
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LCV FISH
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LCV DAPHNIDS
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LTV EC20 FISH
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LTV EC20 DAPHNID
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	SS EC20
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	EC20 POPULATION
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	NAWQ ADVISORY
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	NAWQ ACTUAL
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LCV FISH
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LCV DAPHNIDS
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LTV EC20 FISH
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LTV EC20 DAPHNID
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	SS EC20
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	EC20 POPULATION
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	NAWQ ADVISORY
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	NAWQ ACTUAL
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LCV FISH
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LCV DAPHNIDS
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LTV EC20 FISH
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LTV EC20 DAPHNID
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	SS EC20
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	EC20 POPULATION
BENZO(G,H,I)PERYLENE	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	NAWQ ADVISORY
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	NAWQ ACTUAL
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LCV FISH
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LCV DAPHNIDS
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LTV EC20 FISH
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LTV EC20 DAPHNID
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	SS EC20
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	EC20 POPULATION
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	NAWQ ADVISORY
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	NAWQ ACTUAL
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LCV FISH
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LCV DAPHNIDS
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LTV EC20 FISH
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LTV EC20 DAPHNID
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	SS EC20
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BENZO(G,H,I)PERYLENE (continued)	3	2	0	6.58	565	0.01	599.27	0.02	.	.	NAHQ ADVISORY
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	NAHQ ACTUAL
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LCV FISH
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LCV DAPHNIDS
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LTV EC20 FISH
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LTV EC20 DAPHNID
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	SS EC20
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	EC20 POPULATION
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	NAHQ ADVISORY
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	NAHQ ACTUAL
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LCV FISH
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LCV DAPHNIDS
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LTV EC20 FISH
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LTV EC20 DAPHNID
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	SS EC20
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	EC20 POPULATION
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	NAHQ ADVISORY
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	NAHQ ACTUAL
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LCV FISH
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LCV DAPHNIDS
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LTV EC20 FISH
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LTV EC20 DAPHNID
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	SS EC20
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	EC20 POPULATION
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	NAHQ ADVISORY
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	NAHQ ACTUAL
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LCV FISH
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LCV DAPHNIDS
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LTV EC20 FISH
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LTV EC20 DAPHNID
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	SS EC20
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BENZO(K)FLUORANTHENE	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	NAHQ ADVISORY
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	NAHQ ACTUAL
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LCV FISH
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LCV DAPHNIDS
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LTV EC20 FISH
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	LTV EC20 DAPHNID
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	SS EC20
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	EC20 POPULATION
	1	2	0	6.84	620	0.01	6993.39	0.10	.	.	NAHQ ADVISORY
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	NAHQ ACTUAL
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LCV FISH
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LCV DAPHNIDS
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LTV EC20 FISH
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	LTV EC20 DAPHNID
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	SS EC20
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	EC20 POPULATION
	2	2	0	6.84	445	0.01	616.33	0.01	.	.	NAHQ ADVISORY
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	NAHQ ACTUAL
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	LCV FISH
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	LCV DAPHNIDS
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	LTV EC20 FISH
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	LTV EC20 DAPHNID
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	SS EC20
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	EC20 POPULATION
	3	2	0	6.84	565	0.01	599.27	0.01	.	.	NAHQ ADVISORY
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	NAHQ ACTUAL
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LCV FISH
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LCV DAPHNIDS
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LTV EC20 FISH
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	LTV EC20 DAPHNID
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	SS EC20
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	EC20 POPULATION
	4	2	0	6.84	500	0.01	38534.72	0.56	.	.	NAHQ ADVISORY
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	NAHQ ACTUAL
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LCV FISH
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LCV DAPHNIDS
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LTV EC20 FISH
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	LTV EC20 DAPHNID
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	SS EC20
	5	4	0	6.84	855	0.01	6848.50	0.10	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a limit	UCL conc.	MOL quotient	UCL quotient	Benchmark type
BENZO(K) FLUORANTHENE (continued)	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	NAHQ ADVISORY
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	NAHQ ACTUAL
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LCV FISH
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LCV DAPHNIDS
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LTV EC20 FISH
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	LTV EC20 DAPHNID
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	SS EC20
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	EC20 POPULATION
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	
	10	1	0	6.84	6500	0.09	6500.00	0.09	.	.	
BENZOIC ACID	1	2	0	1.87	3000	4046.89	33838.96	45647.50	.	.	NAHQ ACTUAL
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	.	.	LCV DAPHNIDS
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	.	.	LTV EC20 DAPHNID
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	.	.	SS EC20
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	8993.09	101438.89	0.45 NAHQ ADVISORY
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	3.21	36.26	1259.00 EC20 POPULATION
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	0.55	6.16	7409.00 LTV EC20 FISH
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	0.31	3.52	12976.00 LCV FISH
	1	2	0	1.87	3000	4046.89	33838.96	45647.50	.	.	NAHQ ACTUAL
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	.	.	LCV DAPHNIDS
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	.	.	LTV EC20 DAPHNID
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	.	.	SS EC20
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	6445.04	8499.40	0.45 NAHQ ADVISORY
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	2.30	3.04	1259.00 EC20 POPULATION
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	0.39	0.52	7409.00 LTV EC20 FISH
	2	2	0	1.87	2150	2900.27	2835.31	3824.73	0.22	0.29	12976.00 LCV FISH
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	.	.	NAHQ ACTUAL
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	.	.	LCV DAPHNIDS
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	.	.	LTV EC20 DAPHNID
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	.	.	SS EC20
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	8243.66	8243.66	0.45 NAHQ ADVISORY
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	2.95	2.95	1259.00 EC20 POPULATION
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	0.50	0.50	7409.00 LTV EC20 FISH
	3	2	0	1.87	2750	3709.65	2750.00	3709.65	0.29	0.29	12976.00 LCV FISH
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	.	.	NAHQ ACTUAL
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	.	.	LCV DAPHNIDS
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	.	.	LTV EC20 DAPHNID
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	.	.	SS EC20
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	7344.35	563046.37	0.45 NAHQ ADVISORY
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	2.63	201.25	1259.00 EC20 POPULATION
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	0.45	34.20	7409.00 LTV EC20 FISH
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	0.25	19.53	12976.00 LCV FISH
	4	2	0	1.87	2450	3304.96	187826.42	253370.87	.	.	

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a limit	UCL conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
BENZOIC ACID (continued)	5	4	0	1.87	4150	5598.20	33158.52	44729.62	.	.	.	NAHQ ACTUAL
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	.	.	.	LCV DAPHNIDS
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	.	.	.	LTV EC20 DAPHNID
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	.	.	.	SS EC20
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	12440.44	99399.15	0.45	NAHQ ADVISORY
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	4.45	35.53	1259.00	EC20 POPULATION
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	0.76	6.04	7409.00	LTV EC20 FISH
	5	4	0	1.87	4150	5598.20	33158.52	44729.62	0.43	3.45	12976.00	LCV FISH
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	.	.	.	NAHQ ACTUAL
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	.	.	.	LCV DAPHNIDS
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	.	.	.	LTV EC20 DAPHNID
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	92928.55	92928.55	0.45	NAHQ ADVISORY
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	33.22	33.22	1259.00	EC20 POPULATION
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	5.64	5.64	7409.00	LTV EC20 FISH
	10	1	0	1.87	31000	41817.85	31000.00	41817.85	3.22	3.22	12976.00	LCV FISH
BENZYL ALCOHOL	1	2	0	1.10	620	4924.84	6993.39	55550.43	.	.	.	NAHQ ACTUAL
	1	2	0	1.10	620	4924.84	6993.39	55550.43	.	.	.	LCV DAPHNIDS
	1	2	0	1.10	620	4924.84	6993.39	55550.43	.	.	.	LTV EC20 DAPHNID
	1	2	0	1.10	620	4924.84	6993.39	55550.43	.	.	.	SS EC20
	1	2	0	1.10	620	4924.84	6993.39	55550.43	33502.28	377894.10	0.15	NAHQ ADVISORY
	1	2	0	1.10	620	4924.84	6993.39	55550.43	13.13	148.13	375.00	EC20 POPULATION
	1	2	0	1.10	620	4924.84	6993.39	55550.43	8.95	101.00	550.00	LTV EC20 FISH
	1	2	0	1.10	620	4924.84	6993.39	55550.43	8.36	94.31	589.00	LCV FISH
	2	2	0	1.10	445	3534.76	616.33	4895.66	.	.	.	NAHQ ACTUAL
	2	2	0	1.10	445	3534.76	616.33	4895.66	.	.	.	LCV DAPHNIDS
	2	2	0	1.10	445	3534.76	616.33	4895.66	.	.	.	LTV EC20 DAPHNID
	2	2	0	1.10	445	3534.76	616.33	4895.66	.	.	.	SS EC20
	2	2	0	1.10	445	3534.76	616.33	4895.66	24045.99	33303.84	0.15	NAHQ ADVISORY
	2	2	0	1.10	445	3534.76	616.33	4895.66	9.43	13.06	375.00	EC20 POPULATION
	2	2	0	1.10	445	3534.76	616.33	4895.66	6.43	8.90	550.00	LTV EC20 FISH
	2	2	0	1.10	445	3534.76	616.33	4895.66	6.00	8.31	589.00	LCV FISH
	3	2	0	1.10	565	4487.95	599.27	4760.14	.	.	.	NAHQ ACTUAL
	3	2	0	1.10	565	4487.95	599.27	4760.14	.	.	.	LCV DAPHNIDS
	3	2	0	1.10	565	4487.95	599.27	4760.14	.	.	.	LTV EC20 DAPHNID
	3	2	0	1.10	565	4487.95	599.27	4760.14	30530.30	32381.87	0.15	NAHQ ADVISORY
	3	2	0	1.10	565	4487.95	599.27	4760.14	11.97	12.69	375.00	EC20 POPULATION
	3	2	0	1.10	565	4487.95	599.27	4760.14	8.16	8.65	550.00	LTV EC20 FISH
	3	2	0	1.10	565	4487.95	599.27	4760.14	7.62	8.08	589.00	LCV FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BENZYL ALCOHOL (continued)	4	2	0	1.10	500	3971.64	38534.72	306092.15	.	.	NAWQ ACTUAL
	4	2	0	1.10	500	3971.64	38534.72	306092.15	.	.	LCV DAPHNIDS
	4	2	0	1.10	500	3971.64	38534.72	306092.15	.	.	LTV EC20 DAPHNID
	4	2	0	1.10	500	3971.64	38534.72	306092.15	.	.	SS EC20
	4	2	0	1.10	500	3971.64	38534.72	306092.15	27017.97	2082259.50	0.15 NAWQ ADVISORY
	4	2	0	1.10	500	3971.64	38534.72	306092.15	10.59	816.25	375.00 EC20 POPULATION
	4	2	0	1.10	500	3971.64	38534.72	306092.15	7.22	556.53	550.00 LTV EC20 FISH
	4	2	0	1.10	500	3971.64	38534.72	306092.15	6.74	519.68	589.00 LCV FISH
	4	2	0	1.10	500	3971.64	38534.72	306092.15	.	.	NAWQ ACTUAL
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	.	.	LCV DAPHNIDS
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	.	.	LTV EC20 DAPHNID
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	.	.	SS EC20
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	57818.45	560243.81	0.15 NAWQ ADVISORY
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	22.66	219.62	375.00 EC20 POPULATION
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	15.45	149.74	550.00 LTV EC20 FISH
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	14.43	139.82	589.00 LCV FISH
	5	3	0	1.10	1070	8499.31	10367.99	82355.84	.	.	NAWQ ACTUAL
	10	1	0	1.10	6500	51631.34	6500.00	51631.34	.	.	LCV DAPHNIDS
	10	1	0	1.10	6500	51631.34	6500.00	51631.34	.	.	LTV EC20 DAPHNID
	10	1	0	1.10	6500	51631.34	6500.00	51631.34	.	.	SS EC20
BIS(2-ETHYLHEXYL)PHTHALATE	1	2	0	4.89	1180	1.52	5428.92	6.99	.	.	NAWQ ACTUAL
	1	2	0	4.89	1180	1.52	5428.92	6.99	.	.	SS EC20
	1	2	0	4.89	1180	1.52	5428.92	6.99	152.01	699.38	0.01 NAWQ ADVISORY
	1	2	0	4.89	1180	1.52	5428.92	6.99	0.51	2.33	3.00 LCV DAPHNIDS
	1	2	0	4.89	1180	1.52	5428.92	6.99	0.51	2.33	3.00 LTV EC20 DAPHNID
	1	2	0	4.89	1180	1.52	5428.92	6.99	0.18	0.83	8.40 LCV FISH
	1	2	0	4.89	1180	1.52	5428.92	6.99	0.03	0.14	50.00 EC20 POPULATION
	1	2	0	4.89	1180	1.52	5428.92	6.99	0.03	0.13	54.00 LTV EC20 FISH
	1	2	0	4.89	1180	1.52	5428.92	6.99	.	.	NAWQ ACTUAL
	4	2	0	4.89	500	0.64	27638.29	35.61	.	.	SS EC20
	4	2	0	4.89	500	0.64	27638.29	35.61	64.41	3560.50	0.01 NAWQ ADVISORY
	4	2	0	4.89	500	0.64	27638.29	35.61	0.21	11.87	3.00 LCV DAPHNIDS
	4	2	0	4.89	500	0.64	27638.29	35.61	0.21	11.87	3.00 LTV EC20 DAPHNID
	4	2	0	4.89	500	0.64	27638.29	35.61	0.08	4.24	8.40 LCV FISH
	4	2	0	4.89	500	0.64	27638.29	35.61	0.01	0.71	50.00 EC20 POPULATION
	4	2	0	4.89	500	0.64	27638.29	35.61	0.01	0.66	54.00 LTV EC20 FISH
	4	2	0	4.89	500	0.64	27638.29	35.61	.	.	NAWQ ACTUAL
	4	2	0	4.89	500	0.64	27638.29	35.61	.	.	SS EC20
	4	2	0	4.89	500	0.64	27638.29	35.61	64.41	3560.50	0.01 NAWQ ADVISORY
	4	2	0	4.89	500	0.64	27638.29	35.61	0.21	11.87	3.00 LCV DAPHNIDS
	4	2	0	4.89	500	0.64	27638.29	35.61	0.08	4.24	8.40 LCV FISH
	4	2	0	4.89	500	0.64	27638.29	35.61	0.01	0.71	50.00 EC20 POPULATION
	4	2	0	4.89	500	0.64	27638.29	35.61	0.01	0.66	54.00 LTV EC20 FISH

Table F.10 (continued)

Analyte	Resch (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL* detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
BIS(2-ETHYLHEXYL)PHTHALATE (cont.)	10	1	0	4.89	6500	8.37	6500.00	8.37	.	.	NAHQ ACTUAL
	10	1	0	4.89	6500	8.37	6500.00	8.37	.	.	SS EC20
	10	1	0	4.89	6500	8.37	6500.00	8.37	837.36	837.36	0.01 NAHQ ADVISORY
	10	1	0	4.89	6500	8.37	6500.00	8.37	2.79	2.79	3.00 LCV DAPHNIDS
	10	1	0	4.89	6500	8.37	6500.00	8.37	2.79	2.79	3.00 LTV EC20 DAPHNID
	10	1	0	4.89	6500	8.37	6500.00	8.37	1.00	1.00	8.40 LCV FISH
	10	1	0	4.89	6500	8.37	6500.00	8.37	0.17	0.17	50.00 EC20 POPULATION
	10	1	0	4.89	6500	8.37	6500.00	8.37	0.16	0.16	54.00 LTV EC20 FISH
	10	1	0	4.89	6500	8.37	6500.00	8.37	.	.	NAHQ ACTUAL
	10	1	0	4.89	6500	8.37	6500.00	8.37	837.36	837.36	0.01 NAHQ ADVISORY
CHLORDANE	1	2	0	2.78	1500	248.94	17262.14	2864.80	1464.34	16851.77	0.17 NAHQ ACTUAL
	1	2	0	2.78	1500	248.94	17262.14	2864.80	995.75	11459.21	0.25 LTV EC20 FISH
	1	2	0	2.78	1500	248.94	17262.14	2864.80	497.88	5729.60	0.50 SS EC20
	1	2	0	2.78	1500	248.94	17262.14	2864.80	350.62	4034.93	0.71 EC20 POPULATION
	1	2	0	2.78	1500	248.94	17262.14	2864.80	155.59	1790.50	1.60 LCV FISH
	1	2	0	2.78	1500	248.94	17262.14	2864.80	20.57	236.76	12.10 LTV EC20 DAPHNID
	1	2	0	2.78	1500	248.94	17262.14	2864.80	15.56	179.05	16.00 LCV DAPHNIDS
	2	2	0	2.78	325	53.94	6047.34	1003.61	.	.	NAHQ ADVISORY
	2	2	0	2.78	325	53.94	6047.34	1003.61	317.27	5903.58	0.17 NAHQ ACTUAL
	2	2	0	2.78	325	53.94	6047.34	1003.61	215.75	4014.43	0.25 LTV EC20 FISH
	2	2	0	2.78	325	53.94	6047.34	1003.61	107.87	2007.22	0.50 SS EC20
	2	2	0	2.78	325	53.94	6047.34	1003.61	75.97	1413.53	0.71 EC20 POPULATION
	2	2	0	2.78	325	53.94	6047.34	1003.61	33.71	627.26	1.60 LCV FISH
	2	2	0	2.78	325	53.94	6047.34	1003.61	4.46	82.94	12.10 LTV EC20 DAPHNID
	2	2	0	2.78	325	53.94	6047.34	1003.61	3.37	62.73	16.00 LCV DAPHNIDS
	3	2	0	2.78	1350	224.04	14028.24	2328.11	.	.	NAHQ ADVISORY
	3	2	0	2.78	1350	224.04	14028.24	2328.11	1317.91	13694.75	0.17 NAHQ ACTUAL
	3	2	0	2.78	1350	224.04	14028.24	2328.11	896.18	9312.43	0.25 LTV EC20 FISH
	3	2	0	2.78	1350	224.04	14028.24	2328.11	448.09	4656.22	0.50 SS EC20
	3	2	0	2.78	1350	224.04	14028.24	2328.11	315.56	3279.03	0.71 EC20 POPULATION
	3	2	0	2.78	1350	224.04	14028.24	2328.11	140.03	1455.07	1.60 LCV FISH
	3	2	0	2.78	1350	224.04	14028.24	2328.11	18.52	192.41	12.10 LTV EC20 DAPHNID
	3	2	0	2.78	1350	224.04	14028.24	2328.11	14.00	145.51	16.00 LCV DAPHNIDS
	4	2	0	2.78	1200	199.15	47801.10	7933.01	.	.	NAHQ ADVISORY
	4	2	0	2.78	1200	199.15	47801.10	7933.01	1171.47	46664.75	0.17 NAHQ ACTUAL
	4	2	0	2.78	1200	199.15	47801.10	7933.01	796.60	31732.03	0.25 LTV EC20 FISH
	4	2	0	2.78	1200	199.15	47801.10	7933.01	398.30	15866.01	0.50 SS EC20
	4	2	0	2.78	1200	199.15	47801.10	7933.01	280.49	11173.25	0.71 EC20 POPULATION
	4	2	0	2.78	1200	199.15	47801.10	7933.01	124.47	4958.13	1.60 LCV FISH
	4	2	0	2.78	1200	199.15	47801.10	7933.01	16.46	655.62	12.10 LTV EC20 DAPHNID
	4	2	0	2.78	1200	199.15	47801.10	7933.01	12.45	495.81	16.00 LCV DAPHNIDS

Table F.10 (continued)

Analyte	Resch (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
CHLORDANE (continued)	5	2	0	2.78	1290	214.09	12323.49	2045.19	1259.33	12030.54	NAHQ ADVISORY	NAHQ ADVISORY
	5	2	0	2.78	1290	214.09	12323.49	2045.19	856.35	8180.76	0.17 NAHQ ACTUAL	0.17 NAHQ ACTUAL
	5	2	0	2.78	1290	214.09	12323.49	2045.19	428.17	4090.38	0.25 LTV EC20 FISH	0.25 LTV EC20 FISH
	5	2	0	2.78	1290	214.09	12323.49	2045.19	301.53	2880.55	0.50 SS EC20	0.50 SS EC20
	5	2	0	2.78	1290	214.09	12323.49	2045.19	133.80	1278.24	0.71 EC20 POPULATION	0.71 EC20 POPULATION
	5	2	0	2.78	1290	214.09	12323.49	2045.19	17.69	169.02	1.60 LCV FISH	1.60 LCV FISH
	5	2	0	2.78	1290	214.09	12323.49	2045.19	13.38	127.82	12.10 LTV EC20 DAPHNID	12.10 LTV EC20 DAPHNID
	5	2	0	2.78	1290	214.09	12323.49	2045.19			16.00 LCV DAPHNIDS	16.00 LCV DAPHNIDS
	1	2	0	5.91	620	0.08	6993.39	0.86			NAHQ ADVISORY	NAHQ ADVISORY
	1	2	0	5.91	620	0.08	6993.39	0.86			NAHQ ACTUAL	NAHQ ACTUAL
CHRYSENE	1	2	0	5.91	620	0.08	6993.39	0.86			LCV FISH	LCV FISH
	1	2	0	5.91	620	0.08	6993.39	0.86			LCV DAPHNIDS	LCV DAPHNIDS
	1	2	0	5.91	620	0.08	6993.39	0.86			LTV EC20 FISH	LTV EC20 FISH
	1	2	0	5.91	620	0.08	6993.39	0.86			LTV EC20 DAPHNID	LTV EC20 DAPHNID
	1	2	0	5.91	620	0.08	6993.39	0.86			SS EC20	SS EC20
	1	2	0	5.91	620	0.08	6993.39	0.86			EC20 POPULATION	EC20 POPULATION
	1	2	0	5.91	445	0.05	616.33	0.08			NAHQ ADVISORY	NAHQ ADVISORY
	2	2	0	5.91	445	0.05	616.33	0.08			NAHQ ACTUAL	NAHQ ACTUAL
	2	2	0	5.91	445	0.05	616.33	0.08			LCV FISH	LCV FISH
	2	2	0	5.91	445	0.05	616.33	0.08			LCV DAPHNIDS	LCV DAPHNIDS
	2	2	0	5.91	445	0.05	616.33	0.08			LTV EC20 FISH	LTV EC20 FISH
	2	2	0	5.91	445	0.05	616.33	0.08			LTV EC20 DAPHNID	LTV EC20 DAPHNID
	2	2	0	5.91	445	0.05	616.33	0.08			SS EC20	SS EC20
	2	2	0	5.91	445	0.05	616.33	0.08			EC20 POPULATION	EC20 POPULATION
	2	2	0	5.91	445	0.05	616.33	0.08			NAHQ ADVISORY	NAHQ ADVISORY
	3	2	0	5.91	565	0.07	599.27	0.07			NAHQ ACTUAL	NAHQ ACTUAL
	3	2	0	5.91	565	0.07	599.27	0.07			LCV FISH	LCV FISH
	3	2	0	5.91	565	0.07	599.27	0.07			LCV DAPHNIDS	LCV DAPHNIDS
	3	2	0	5.91	565	0.07	599.27	0.07			LTV EC20 FISH	LTV EC20 FISH
	3	2	0	5.91	565	0.07	599.27	0.07			LTV EC20 DAPHNID	LTV EC20 DAPHNID
	3	2	0	5.91	565	0.07	599.27	0.07			SS EC20	SS EC20
	3	2	0	5.91	565	0.07	599.27	0.07			EC20 POPULATION	EC20 POPULATION
	3	2	0	5.91	565	0.07	599.27	0.07			NAHQ ADVISORY	NAHQ ADVISORY
	4	2	0	5.91	500	0.06	38534.72	4.74			NAHQ ACTUAL	NAHQ ACTUAL
	4	2	0	5.91	500	0.06	38534.72	4.74			LCV FISH	LCV FISH
	4	2	0	5.91	500	0.06	38534.72	4.74			LCV DAPHNIDS	LCV DAPHNIDS
	4	2	0	5.91	500	0.06	38534.72	4.74			LTV EC20 FISH	LTV EC20 FISH
	4	2	0	5.91	500	0.06	38534.72	4.74			LTV EC20 DAPHNID	LTV EC20 DAPHNID
	4	2	0	5.91	500	0.06	38534.72	4.74			SS EC20	SS EC20
	4	2	0	5.91	500	0.06	38534.72	4.74			EC20 POPULATION	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
CHRYSENE (continued)	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	NAHQ ADVISORY
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	NAHQ ACTUAL
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	LCV FISH
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	LCV DAPHNIDS
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	LTV EC20 FISH
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	LTV EC20 DAPHNID
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	SS EC20
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	EC20 POPULATION
	5	4	0	5.91	855	0.11	6791.58	0.84	.	.	NAHQ ADVISORY
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	NAHQ ACTUAL
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	LCV FISH
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	LCV DAPHNIDS
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	LTV EC20 FISH
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	LTV EC20 DAPHNID
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	SS EC20
	10	1	0	5.91	6500	0.80	6500.00	0.80	.	.	EC20 POPULATION
	1	2	0	5.20	620	0.39	6993.39	4.41	.	.	NAHQ ACTUAL
	1	2	0	5.20	620	0.39	6993.39	4.41	0.09	1.00	SS EC20
	1	2	0	5.20	620	0.39	6993.39	4.41	0.00	0.02	4.40 NAHQ ADVISORY
	1	2	0	5.20	620	0.39	6993.39	4.41	0.00	0.02	251.00 EC20 POPULATION
DI-N-BUTYLPHTHALATE	1	2	0	5.20	620	0.39	6993.39	4.41	0.00	0.01	270.00 LTV EC20 FISH
	1	2	0	5.20	620	0.39	6993.39	4.41	0.00	0.01	500.00 LTV EC20 DAPHNID
	1	2	0	5.20	620	0.39	6993.39	4.41	0.00	0.01	717.00 LCV FISH
	1	2	0	5.20	620	0.39	6993.39	4.41	0.00	0.01	717.00 LCV DAPHNIDS
	2	2	0	5.20	445	0.28	616.33	0.39	.	.	NAHQ ACTUAL
	2	2	0	5.20	445	0.28	616.33	0.39	.	.	SS EC20
	2	2	0	5.20	445	0.28	616.33	0.39	0.06	0.09	4.40 NAHQ ADVISORY
	2	2	0	5.20	445	0.28	616.33	0.39	0.00	0.00	251.00 EC20 POPULATION
	2	2	0	5.20	445	0.28	616.33	0.39	0.00	0.00	270.00 LTV EC20 FISH
	2	2	0	5.20	445	0.28	616.33	0.39	0.00	0.00	500.00 LTV EC20 DAPHNID
	2	2	0	5.20	445	0.28	616.33	0.39	0.00	0.00	717.00 LCV FISH
	2	2	0	5.20	445	0.28	616.33	0.39	0.00	0.00	717.00 LCV DAPHNIDS
	3	2	0	5.20	565	0.36	599.27	0.38	.	.	NAHQ ACTUAL
	3	2	0	5.20	565	0.36	599.27	0.38	.	.	SS EC20
	3	2	0	5.20	565	0.36	599.27	0.38	0.08	0.09	4.40 NAHQ ADVISORY
	3	2	0	5.20	565	0.36	599.27	0.38	0.00	0.00	251.00 EC20 POPULATION
	3	2	0	5.20	565	0.36	599.27	0.38	0.00	0.00	270.00 LTV EC20 FISH
	3	2	0	5.20	565	0.36	599.27	0.38	0.00	0.00	500.00 LTV EC20 DAPHNID
	3	2	0	5.20	565	0.36	599.27	0.38	0.00	0.00	717.00 LCV FISH
	3	2	0	5.20	565	0.36	599.27	0.38	0.00	0.00	717.00 LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore conc.	UCL ^a detection limit	UCL conc.	HDI quotient	UCL quotient	Benchmark type
DI-N-BUTYLPHTHALATE (continued)											
	4	2	0	5.20	500	0.32	38534.72	24.31	.	.	NAWQ ACTUAL
	4	2	0	5.20	500	0.32	38534.72	24.31	.	.	SS EC20
	4	2	0	5.20	500	0.32	38534.72	24.31	0.07	5.53	4.40 NAWQ ADVISORY
	4	2	0	5.20	500	0.32	38534.72	24.31	0.00	0.10	251.00 EC20 POPULATION
	4	2	0	5.20	500	0.32	38534.72	24.31	0.00	0.09	270.00 LTV EC20 FISH
	4	2	0	5.20	500	0.32	38534.72	24.31	0.00	0.05	500.00 LTV EC20 DAPHNID
	4	2	0	5.20	500	0.32	38534.72	24.31	0.00	0.03	717.00 LCV FISH
	4	2	0	5.20	500	0.32	38534.72	24.31	0.00	0.03	717.00 LCV DAPHNIDS
	5	4	0	5.20	855	0.54	6848.50	4.32	.	.	NAWQ ACTUAL
	5	4	0	5.20	855	0.54	6848.50	4.32	.	.	SS EC20
	5	4	0	5.20	855	0.54	6848.50	4.32	0.12	0.98	4.40 NAWQ ADVISORY
	5	4	0	5.20	855	0.54	6848.50	4.32	0.00	0.02	251.00 EC20 POPULATION
	5	4	0	5.20	855	0.54	6848.50	4.32	0.00	0.02	270.00 LTV EC20 FISH
	5	4	0	5.20	855	0.54	6848.50	4.32	0.00	0.01	500.00 LTV EC20 DAPHNID
	5	4	0	5.20	855	0.54	6848.50	4.32	0.00	0.01	717.00 LCV FISH
	5	4	0	5.20	855	0.54	6848.50	4.32	0.00	0.01	717.00 LCV DAPHNIDS
	10	1	0	5.20	6500	4.10	6500.00	4.10	.	.	NAWQ ACTUAL
	10	1	0	5.20	6500	4.10	6500.00	4.10	.	.	SS EC20
	10	1	0	5.20	6500	4.10	6500.00	4.10	0.93	0.93	4.40 NAWQ ADVISORY
	10	1	0	5.20	6500	4.10	6500.00	4.10	0.02	0.02	251.00 EC20 POPULATION
	10	1	0	5.20	6500	4.10	6500.00	4.10	0.02	0.02	270.00 LTV EC20 FISH
	10	1	0	5.20	6500	4.10	6500.00	4.10	0.01	0.01	500.00 LTV EC20 DAPHNID
	10	1	0	5.20	6500	4.10	6500.00	4.10	0.01	0.01	717.00 LCV FISH
	10	1	0	5.20	6500	4.10	6500.00	4.10	0.01	0.01	717.00 LCV DAPHNIDS
DI-N-OCTYLPHTHALATE											
	1	2	0	5.55	620	0.17	6993.39	1.97	.	.	NAWQ ADVISORY
	1	2	0	5.55	620	0.17	6993.39	1.97	.	.	NAWQ ACTUAL
	1	2	0	5.55	620	0.17	6993.39	1.97	.	.	SS EC20
	1	2	0	5.55	620	0.17	6993.39	1.97	0.00	0.02	100.00 LTV EC20 FISH
	1	2	0	5.55	620	0.17	6993.39	1.97	0.00	0.01	310.00 LTV EC20 DAPHNID
	1	2	0	5.55	620	0.17	6993.39	1.97	0.00	0.00	570.00 LCV DAPHNIDS
	1	2	0	5.55	620	0.17	6993.39	1.97	0.00	0.00	1995.00 EC20 POPULATION
	1	2	0	5.55	620	0.17	6993.39	1.97	0.00	0.00	5700.00 LCV FISH
	2	2	0	5.55	445	0.13	616.33	0.17	.	.	NAWQ ADVISORY
	2	2	0	5.55	445	0.13	616.33	0.17	.	.	NAWQ ACTUAL
	2	2	0	5.55	445	0.13	616.33	0.17	.	.	SS EC20
	2	2	0	5.55	445	0.13	616.33	0.17	0.00	0.00	100.00 LTV EC20 FISH
	2	2	0	5.55	445	0.13	616.33	0.17	0.00	0.00	310.00 LTV EC20 DAPHNID
	2	2	0	5.55	445	0.13	616.33	0.17	0.00	0.00	570.00 LCV DAPHNIDS
	2	2	0	5.55	445	0.13	616.33	0.17	0.00	0.00	1995.00 EC20 POPULATION
	2	2	0	5.55	445	0.13	616.33	0.17	0.00	0.00	5700.00 LCV FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark	
											Benchmark type	Benchmark
DI-N-OCTYLPHTHALATE (continued)	3	2	0	5.55	565	0.16	599.27	0.17	-	-	-	NAHQ ADVISORY
	3	2	0	5.55	565	0.16	599.27	0.17	-	-	-	NAHQ ACTUAL
	3	2	0	5.55	565	0.16	599.27	0.17	-	-	-	SS EC20
	3	2	0	5.55	565	0.16	599.27	0.17	0.00	0.00	100.00	LTV EC20 FISH
	3	2	0	5.55	565	0.16	599.27	0.17	0.00	0.00	310.00	LTV EC20 DAPHNID
	3	2	0	5.55	565	0.16	599.27	0.17	0.00	0.00	570.00	LCV DAPHNIDS
	3	2	0	5.55	565	0.16	599.27	0.17	0.00	0.00	1995.00	EC20 POPULATION
	3	2	0	5.55	565	0.16	599.27	0.17	0.00	0.00	5700.00	LCV FISH
	3	2	0	5.55	565	0.16	599.27	0.17	-	-	-	NAHQ ADVISORY
	4	2	0	5.55	500	0.14	38534.72	10.86	-	-	-	NAHQ ACTUAL
	4	2	0	5.55	500	0.14	38534.72	10.86	-	-	-	SS EC20
	4	2	0	5.55	500	0.14	38534.72	10.86	0.00	0.00	100.00	LTV EC20 FISH
	4	2	0	5.55	500	0.14	38534.72	10.86	0.00	0.04	310.00	LTV EC20 DAPHNID
	4	2	0	5.55	500	0.14	38534.72	10.86	0.00	0.02	570.00	LCV DAPHNIDS
	4	2	0	5.55	500	0.14	38534.72	10.86	0.00	0.01	1995.00	EC20 POPULATION
	4	2	0	5.55	500	0.14	38534.72	10.86	0.00	0.00	5700.00	LCV FISH
	4	2	0	5.55	500	0.14	38534.72	10.86	-	-	-	NAHQ ADVISORY
	5	4	0	5.55	855	0.24	6848.50	1.93	-	-	-	NAHQ ACTUAL
	5	4	0	5.55	855	0.24	6848.50	1.93	-	-	-	SS EC20
	5	4	0	5.55	855	0.24	6848.50	1.93	0.00	0.02	100.00	LTV EC20 FISH
	5	4	0	5.55	855	0.24	6848.50	1.93	0.00	0.01	310.00	LTV EC20 DAPHNID
	5	4	0	5.55	855	0.24	6848.50	1.93	0.00	0.00	570.00	LCV DAPHNIDS
	5	4	0	5.55	855	0.24	6848.50	1.93	0.00	0.00	1995.00	EC20 POPULATION
	5	4	0	5.55	855	0.24	6848.50	1.93	0.00	0.00	5700.00	LCV FISH
10	1	0	5.55	6500	1.83	6500.00	1.83	-	-	-	NAHQ ADVISORY	
10	1	0	5.55	6500	1.83	6500.00	1.83	-	-	-	NAHQ ACTUAL	
10	1	0	5.55	6500	1.83	6500.00	1.83	-	-	-	SS EC20	
10	1	0	5.55	6500	1.83	6500.00	1.83	0.02	0.02	100.00	LTV EC20 FISH	
10	1	0	5.55	6500	1.83	6500.00	1.83	0.01	0.01	310.00	LTV EC20 DAPHNID	
10	1	0	5.55	6500	1.83	6500.00	1.83	0.00	0.00	570.00	LCV DAPHNIDS	
10	1	0	5.55	6500	1.83	6500.00	1.83	0.00	0.00	1995.00	EC20 POPULATION	
10	1	0	5.55	6500	1.83	6500.00	1.83	0.00	0.00	5700.00	LCV FISH	
DIBENZOFURAN	1	2	0	4.12	620	4.70	6993.39	53.05	-	-	-	NAHQ ACTUAL
	1	2	0	4.12	620	4.70	6993.39	53.05	-	-	-	LCV FISH
	1	2	0	4.12	620	4.70	6993.39	53.05	-	-	-	LTV EC20 FISH
	1	2	0	4.12	620	4.70	6993.39	53.05	-	-	-	LTV EC20 DAPHNID
	1	2	0	4.12	620	4.70	6993.39	53.05	-	-	-	SS EC20
	1	2	0	4.12	620	4.70	6993.39	53.05	-	-	-	EC20 POPULATION
	1	2	0	4.12	620	4.70	6993.39	53.05	26.53	26.53	2.00	NAHQ ADVISORY
	1	2	0	4.12	620	4.70	6993.39	53.05	0.00	0.05	1003.00	LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
DIBENZOFURAN (continued)	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	NAHQ ACTUAL
	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	LCV FISH
	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	LTV EC20 FISH
	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	LTV EC20 DAPHNID
	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	SS EC20
	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	EC20 POPULATION
	2	2	0	4.12	445	3.38	616.33	4.68	1.69	2.34	2.00	NAHQ ADVISORY
	2	2	0	4.12	445	3.38	616.33	4.68	0.00	0.00	1003.00	LCV DAPHNIDS
	2	2	0	4.12	445	3.38	616.33	4.68	.	.	.	NAHQ ACTUAL
	3	2	0	4.12	565	4.29	599.27	4.55	.	.	.	LCV FISH
	3	2	0	4.12	565	4.29	599.27	4.55	.	.	.	LTV EC20 FISH
	3	2	0	4.12	565	4.29	599.27	4.55	.	.	.	LTV EC20 DAPHNID
	3	2	0	4.12	565	4.29	599.27	4.55	.	.	.	SS EC20
	3	2	0	4.12	565	4.29	599.27	4.55	.	.	.	EC20 POPULATION
	3	2	0	4.12	565	4.29	599.27	4.55	2.14	2.27	2.00	NAHQ ADVISORY
	3	2	0	4.12	565	4.29	599.27	4.55	0.00	0.00	1003.00	LCV DAPHNIDS
	3	2	0	4.12	565	4.29	599.27	4.55	.	.	.	NAHQ ACTUAL
	4	2	0	4.12	500	3.79	38534.72	292.32	.	.	.	LCV FISH
	4	2	0	4.12	500	3.79	38534.72	292.32	.	.	.	LTV EC20 FISH
	4	2	0	4.12	500	3.79	38534.72	292.32	.	.	.	LTV EC20 DAPHNID
	4	2	0	4.12	500	3.79	38534.72	292.32	.	.	.	SS EC20
	4	2	0	4.12	500	3.79	38534.72	292.32	.	.	.	EC20 POPULATION
	4	2	0	4.12	500	3.79	38534.72	292.32	1.90	146.16	2.00	NAHQ ADVISORY
	4	2	0	4.12	500	3.79	38534.72	292.32	0.00	0.29	1003.00	LCV DAPHNIDS
	4	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	NAHQ ACTUAL
	5	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	LCV FISH
	5	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	LTV EC20 FISH
	5	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	LTV EC20 DAPHNID
	5	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	SS EC20
	5	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	EC20 POPULATION
	5	4	0	4.12	855	6.49	6848.50	51.95	3.24	25.98	2.00	NAHQ ADVISORY
	5	4	0	4.12	855	6.49	6848.50	51.95	0.01	0.05	1003.00	LCV DAPHNIDS
	5	4	0	4.12	855	6.49	6848.50	51.95	.	.	.	NAHQ ACTUAL
	10	1	0	4.12	6500	49.31	6500.00	49.31	.	.	.	LCV FISH
	10	1	0	4.12	6500	49.31	6500.00	49.31	.	.	.	LTV EC20 FISH
	10	1	0	4.12	6500	49.31	6500.00	49.31	.	.	.	LTV EC20 DAPHNID
	10	1	0	4.12	6500	49.31	6500.00	49.31	.	.	.	SS EC20
	10	1	0	4.12	6500	49.31	6500.00	49.31	.	.	.	EC20 POPULATION
	10	1	0	4.12	6500	49.31	6500.00	49.31	24.65	24.65	2.00	NAHQ ADVISORY
	10	1	0	4.12	6500	49.31	6500.00	49.31	0.05	0.05	1003.00	LCV DAPHNIDS
	10	1	0	4.12	6500	49.31	6500.00	49.31	.	.	.	NAHQ ACTUAL

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number defects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a limit	UCL conc.	MDL quotient	UCL quotient	Benchmark type
DIETHYLPHTHALATE	1	2	0	2.47	620	210.08	6993.39	2369.67	.	.	NAWQ ACTUAL
	1	2	0	2.47	620	210.08	6993.39	2369.67	.	.	LCV FISH
	1	2	0	2.47	620	210.08	6993.39	2369.67	.	.	LCV DAPHNIDS
	1	2	0	2.47	620	210.08	6993.39	2369.67	.	.	LTV EC20 FISH
	1	2	0	2.47	620	210.08	6993.39	2369.67	.	.	LTV EC20 DAPHNID
	1	2	0	2.47	620	210.08	6993.39	2369.67	.	.	SS EC20
	1	2	0	2.47	620	210.08	6993.39	2369.67	7.78	87.77	27.00 NAWQ ADVISORY
	1	2	0	2.47	620	210.08	6993.39	2369.67	0.21	2.37	1000.00 EC20 POPULATION
	1	2	0	2.47	445	150.79	616.33	208.84	.	.	NAWQ ACTUAL
	2	2	0	2.47	445	150.79	616.33	208.84	.	.	LCV FISH
	2	2	0	2.47	445	150.79	616.33	208.84	.	.	LCV DAPHNIDS
	2	2	0	2.47	445	150.79	616.33	208.84	.	.	LTV EC20 FISH
	2	2	0	2.47	445	150.79	616.33	208.84	.	.	LTV EC20 DAPHNID
	2	2	0	2.47	445	150.79	616.33	208.84	.	.	SS EC20
	2	2	0	2.47	445	150.79	616.33	208.84	5.58	7.73	27.00 NAWQ ADVISORY
	2	2	0	2.47	445	150.79	616.33	208.84	0.15	0.21	1000.00 EC20 POPULATION
	3	2	0	2.47	565	191.45	599.27	203.06	.	.	NAWQ ACTUAL
	3	2	0	2.47	565	191.45	599.27	203.06	.	.	LCV FISH
	3	2	0	2.47	565	191.45	599.27	203.06	.	.	LCV DAPHNIDS
	3	2	0	2.47	565	191.45	599.27	203.06	.	.	LTV EC20 FISH
	3	2	0	2.47	565	191.45	599.27	203.06	.	.	LTV EC20 DAPHNID
	3	2	0	2.47	565	191.45	599.27	203.06	.	.	SS EC20
	3	2	0	2.47	565	191.45	599.27	203.06	7.09	7.52	27.00 NAWQ ADVISORY
	3	2	0	2.47	565	191.45	599.27	203.06	0.19	0.20	1000.00 EC20 POPULATION
	4	2	0	2.47	500	169.42	38534.72	13057.26	.	.	NAWQ ACTUAL
	4	2	0	2.47	500	169.42	38534.72	13057.26	.	.	LCV FISH
	4	2	0	2.47	500	169.42	38534.72	13057.26	.	.	LCV DAPHNIDS
	4	2	0	2.47	500	169.42	38534.72	13057.26	.	.	LTV EC20 FISH
	4	2	0	2.47	500	169.42	38534.72	13057.26	.	.	LTV EC20 DAPHNID
	4	2	0	2.47	500	169.42	38534.72	13057.26	.	.	SS EC20
	4	2	0	2.47	500	169.42	38534.72	13057.26	6.27	483.60	27.00 NAWQ ADVISORY
	4	2	0	2.47	500	169.42	38534.72	13057.26	0.17	13.06	1000.00 EC20 POPULATION
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	NAWQ ACTUAL
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	LCV FISH
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	LCV DAPHNIDS
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	LTV EC20 FISH
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	LTV EC20 DAPHNID
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	SS EC20
	5	4	0	2.47	855	289.71	6848.50	2320.58	10.73	85.95	27.00 NAWQ ADVISORY
	5	4	0	2.47	855	289.71	6848.50	2320.58	0.29	2.32	1000.00 EC20 POPULATION
	5	4	0	2.47	855	289.71	6848.50	2320.58	.	.	

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	HDL quotient	UCL quotient	Benchmark type	Benchmark
DIETHYLPHTHALATE (continued)	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	NAHQ ACTUAL	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	LCV FISH	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	LCV DAPHNIDS	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	LTV EC20 FISH	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	LTV EC20 DAPHNID	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	SS EC20	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	81.57	81.57	NAHQ ADVISORY	27.00
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	2.20	2.20	EC20 POPULATION	1000.00
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	NAHQ ACTUAL	.
	10	1	0	2.47	6500	2202.49	6500.00	2202.49	.	.	LCV FISH	.
FLUORANTHENE	1	2	0	4.90	620	0.78	6993.39	8.80	.	.	NAHQ ACTUAL	.
	1	2	0	4.90	620	0.78	6993.39	8.80	.	.	LCV FISH	.
	1	2	0	4.90	620	0.78	6993.39	8.80	.	.	LTV EC20 FISH	.
	1	2	0	4.90	620	0.78	6993.39	8.80	.	.	LTV EC20 DAPHNID	.
	1	2	0	4.90	620	0.78	6993.39	8.80	.	.	SS EC20	.
	1	2	0	4.90	620	0.78	6993.39	8.80	0.46	5.18	1.70 NAHQ ADVISORY	1.70
	1	2	0	4.90	620	0.78	6993.39	8.80	0.02	0.28	EC20 POPULATION	32.00
	1	2	0	4.90	620	0.78	6993.39	8.80	0.00	0.00	LCV DAPHNIDS	65795.00
	2	2	0	4.90	445	0.56	616.33	0.78	.	.	NAHQ ACTUAL	.
	2	2	0	4.90	445	0.56	616.33	0.78	.	.	LCV FISH	.
	2	2	0	4.90	445	0.56	616.33	0.78	.	.	LTV EC20 FISH	.
	2	2	0	4.90	445	0.56	616.33	0.78	.	.	LTV EC20 DAPHNID	.
	2	2	0	4.90	445	0.56	616.33	0.78	.	.	SS EC20	.
	2	2	0	4.90	445	0.56	616.33	0.78	0.33	0.46	1.70 NAHQ ADVISORY	1.70
	2	2	0	4.90	445	0.56	616.33	0.78	0.02	0.02	EC20 POPULATION	32.00
	2	2	0	4.90	445	0.56	616.33	0.78	0.00	0.00	LCV DAPHNIDS	65795.00
	2	2	0	4.90	445	0.56	616.33	0.78	.	.	NAHQ ACTUAL	.
	4	2	0	4.90	500	0.63	38534.72	48.51	.	.	LCV FISH	.
	4	2	0	4.90	500	0.63	38534.72	48.51	.	.	LTV EC20 FISH	.
	4	2	0	4.90	500	0.63	38534.72	48.51	.	.	LTV EC20 DAPHNID	.
	4	2	0	4.90	500	0.63	38534.72	48.51	.	.	SS EC20	.
	4	2	0	4.90	500	0.63	38534.72	48.51	0.37	28.54	1.70 NAHQ ADVISORY	1.70
	4	2	0	4.90	500	0.63	38534.72	48.51	0.02	1.52	EC20 POPULATION	32.00
	4	2	0	4.90	500	0.63	38534.72	48.51	0.00	0.00	LCV DAPHNIDS	65795.00
	4	2	0	4.90	500	0.63	38534.72	48.51	.	.	NAHQ ACTUAL	.
	5	4	0	4.90	855	1.08	6848.50	8.62	.	.	LCV FISH	.
	5	4	0	4.90	855	1.08	6848.50	8.62	.	.	LTV EC20 FISH	.
	5	4	0	4.90	855	1.08	6848.50	8.62	.	.	LTV EC20 DAPHNID	.
	5	4	0	4.90	855	1.08	6848.50	8.62	.	.	SS EC20	.
	5	4	0	4.90	855	1.08	6848.50	8.62	0.63	5.07	1.70 NAHQ ADVISORY	1.70
	5	4	0	4.90	855	1.08	6848.50	8.62	0.03	0.27	EC20 POPULATION	32.00
	5	4	0	4.90	855	1.08	6848.50	8.62	0.00	0.00	LCV DAPHNIDS	65795.00
	5	4	0	4.90	855	1.08	6848.50	8.62	.	.	NAHQ ACTUAL	.

Table F.10 (continued)

Analyte	Resch (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
FLUORANTHENE (continued)	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	NAWQ ACTUAL
	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	LCV FISH
	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	LTV EC20 FISH
	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	LTV EC20 DAPHNID
	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	SS EC20
	10	1	0	4.90	6500	8.18	6500.00	8.18	4.81	4.81	1.70 NAWQ ADVISORY
	10	1	0	4.90	6500	8.18	6500.00	8.18	0.26	0.26	32.00 EC20 POPULATION
	10	1	0	4.90	6500	8.18	6500.00	8.18	0.00	0.00	65795.00 LCV DAPHNIDS
	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	NAWQ ACTUAL
	10	1	0	4.90	6500	8.18	6500.00	8.18	.	.	EC20 POPULATION
GAMMA-BHC (LINDANE)	1	2	0	3.72	150	2.86	1726.21	32.89	.	.	NAWQ ACTUAL
	1	2	0	3.72	150	2.86	1726.21	32.89	35.73	411.15	0.08 NAWQ ADVISORY
	1	2	0	3.72	150	2.86	1726.21	32.89	25.98	299.02	0.11 SS EC20
	1	2	0	3.72	150	2.86	1726.21	32.89	2.60	29.90	1.10 LTV EC20 FISH
	1	2	0	3.72	150	2.86	1726.21	32.89	0.26	2.99	11.00 LTV EC20 DAPHNID
	1	2	0	3.72	150	2.86	1726.21	32.89	0.20	2.27	14.50 LCV DAPHNIDS
	1	2	0	3.72	150	2.86	1726.21	32.89	0.20	2.25	14.60 LCV FISH
	2	2	0	3.72	35	0.67	590.10	11.24	.	.	NAWQ ACTUAL
	2	2	0	3.72	35	0.67	590.10	11.24	.	.	EC20 POPULATION
	2	2	0	3.72	35	0.67	590.10	11.24	8.34	140.55	0.08 NAWQ ADVISORY
	2	2	0	3.72	35	0.67	590.10	11.24	6.06	102.22	0.11 SS EC20
	2	2	0	3.72	35	0.67	590.10	11.24	0.61	10.22	1.10 LTV EC20 FISH
	2	2	0	3.72	35	0.67	590.10	11.24	0.06	1.02	11.00 LTV EC20 DAPHNID
	2	2	0	3.72	35	0.67	590.10	11.24	0.05	0.78	14.50 LCV DAPHNIDS
	2	2	0	3.72	35	0.67	590.10	11.24	0.05	0.77	14.60 LCV FISH
	3	2	0	3.72	135	2.57	1402.82	26.73	.	.	NAWQ ACTUAL
	3	2	0	3.72	135	2.57	1402.82	26.73	.	.	EC20 POPULATION
	3	2	0	3.72	135	2.57	1402.82	26.73	32.15	334.13	0.08 NAWQ ADVISORY
	3	2	0	3.72	135	2.57	1402.82	26.73	23.39	243.00	0.11 SS EC20
	3	2	0	3.72	135	2.57	1402.82	26.73	2.34	24.30	1.10 LTV EC20 FISH
	3	2	0	3.72	135	2.57	1402.82	26.73	0.23	2.43	11.00 LTV EC20 DAPHNID
	3	2	0	3.72	135	2.57	1402.82	26.73	0.18	1.84	14.50 LCV DAPHNIDS
	3	2	0	3.72	135	2.57	1402.82	26.73	0.18	1.83	14.60 LCV FISH
	4	2	0	3.72	120	2.29	4780.11	91.08	.	.	NAWQ ACTUAL
	4	2	0	3.72	120	2.29	4780.11	91.08	.	.	EC20 POPULATION
	4	2	0	3.72	120	2.29	4780.11	91.08	28.58	1138.54	0.08 NAWQ ADVISORY
	4	2	0	3.72	120	2.29	4780.11	91.08	20.79	828.03	0.11 SS EC20
	4	2	0	3.72	120	2.29	4780.11	91.08	2.08	82.80	1.10 LTV EC20 FISH
	4	2	0	3.72	120	2.29	4780.11	91.08	0.21	8.28	11.00 LTV EC20 DAPHNID
	4	2	0	3.72	120	2.29	4780.11	91.08	0.16	6.28	14.50 LCV DAPHNIDS
	4	2	0	3.72	120	2.29	4780.11	91.08	0.16	6.24	14.60 LCV FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
GAMMA-BHC (LINDANE) (continued)	5	2	0	3.72	129	2.46	1232.35	23.48	.	.	NAHQ ACTUAL
	5	2	0	3.72	129	2.46	1232.35	23.48	30.73	293.52	EC20 POPULATION
	5	2	0	3.72	129	2.46	1232.35	23.48	22.35	213.47	0.08 NAHQ ADVISORY
	5	2	0	3.72	129	2.46	1232.35	23.48	2.23	21.35	0.11 SS EC20
	5	2	0	3.72	129	2.46	1232.35	23.48	0.22	2.13	1.10 LTV EC20 FISH
	5	2	0	3.72	129	2.46	1232.35	23.48	0.17	1.62	11.00 LTV EC20 DAPHNID
	5	2	0	3.72	129	2.46	1232.35	23.48	0.17	1.61	14.50 LCV DAPHNIDS
	5	2	0	3.72	129	2.46	1232.35	23.48	0.17	1.61	14.60 LCV FISH
	1	2	0	5.44	150	0.05	1726.21	0.63	.	.	NAHQ ACTUAL
	1	2	0	5.44	150	0.05	1726.21	0.63	13.62	156.69	LTV EC20 DAPHNID
	1	2	0	5.44	150	0.05	1726.21	0.63	5.45	62.68	0.00 SS EC20
	1	2	0	5.44	150	0.05	1726.21	0.63	0.54	6.27	0.01 NAHQ ADVISORY
	1	2	0	5.44	150	0.05	1726.21	0.63	0.06	0.73	0.10 EC20 POPULATION
	1	2	0	5.44	150	0.05	1726.21	0.63	0.04	0.50	0.86 LTV EC20 FISH
	1	2	0	5.44	150	0.05	1726.21	0.63	0.02	0.20	1.26 LCV FISH
HEPTACHLOR	1	2	0	5.44	150	0.05	1726.21	0.63	.	.	NAHQ ACTUAL
	2	2	0	5.44	35	0.01	590.10	0.21	3.18	53.56	LTV EC20 DAPHNID
	2	2	0	5.44	35	0.01	590.10	0.21	1.27	21.43	0.00 SS EC20
	2	2	0	5.44	35	0.01	590.10	0.21	0.13	2.14	0.01 NAHQ ADVISORY
	2	2	0	5.44	35	0.01	590.10	0.21	0.01	0.25	0.10 EC20 POPULATION
	2	2	0	5.44	35	0.01	590.10	0.21	0.01	0.17	0.86 LTV EC20 FISH
	2	2	0	5.44	35	0.01	590.10	0.21	0.00	0.07	1.26 LCV FISH
	2	2	0	5.44	35	0.01	590.10	0.21	.	.	NAHQ ACTUAL
	3	2	0	5.44	135	0.05	1402.82	0.51	12.25	127.33	LTV EC20 DAPHNID
	3	2	0	5.44	135	0.05	1402.82	0.51	4.90	50.93	0.00 SS EC20
	3	2	0	5.44	135	0.05	1402.82	0.51	0.49	5.09	0.01 NAHQ ADVISORY
	3	2	0	5.44	135	0.05	1402.82	0.51	0.06	0.59	0.10 EC20 POPULATION
	3	2	0	5.44	135	0.05	1402.82	0.51	0.04	0.40	0.86 LTV EC20 FISH
	3	2	0	5.44	135	0.05	1402.82	0.51	0.02	0.16	1.26 LCV FISH
	3	2	0	5.44	135	0.05	1402.82	0.51	.	.	NAHQ ACTUAL
	4	2	0	5.44	120	0.04	4780.11	1.74	10.89	433.89	LTV EC20 DAPHNID
	4	2	0	5.44	120	0.04	4780.11	1.74	4.36	173.56	0.01 SS EC20
	4	2	0	5.44	120	0.04	4780.11	1.74	0.44	17.36	0.01 NAHQ ADVISORY
	4	2	0	5.44	120	0.04	4780.11	1.74	0.05	2.02	0.10 EC20 POPULATION
	4	2	0	5.44	120	0.04	4780.11	1.74	0.03	1.38	0.86 LTV EC20 FISH
	4	2	0	5.44	120	0.04	4780.11	1.74	0.01	0.55	1.26 LCV FISH
	4	2	0	5.44	120	0.04	4780.11	1.74	.	.	NAHQ ACTUAL
	4	2	0	5.44	120	0.04	4780.11	1.74	10.89	433.89	LTV EC20 DAPHNID
	4	2	0	5.44	120	0.04	4780.11	1.74	4.36	173.56	0.01 SS EC20
	4	2	0	5.44	120	0.04	4780.11	1.74	0.44	17.36	0.01 NAHQ ADVISORY
	4	2	0	5.44	120	0.04	4780.11	1.74	0.05	2.02	0.10 EC20 POPULATION
	4	2	0	5.44	120	0.04	4780.11	1.74	0.03	1.38	0.86 LTV EC20 FISH
	4	2	0	5.44	120	0.04	4780.11	1.74	0.01	0.55	1.26 LCV FISH
	4	2	0	5.44	120	0.04	4780.11	1.74	.	.	NAHQ ACTUAL
	4	2	0	5.44	120	0.04	4780.11	1.74	10.89	433.89	LTV EC20 DAPHNID
	4	2	0	5.44	120	0.04	4780.11	1.74	4.36	173.56	0.01 SS EC20
	4	2	0	5.44	120	0.04	4780.11	1.74	0.44	17.36	0.01 NAHQ ADVISORY
	4	2	0	5.44	120	0.04	4780.11	1.74	0.05	2.02	0.10 EC20 POPULATION
	4	2	0	5.44	120	0.04	4780.11	1.74	0.03	1.38	0.86 LTV EC20 FISH
	4	2	0	5.44	120	0.04	4780.11	1.74	0.01	0.55	1.26 LCV FISH

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection pore water limit	Minimum conc.	UCL ^a detection pore water limit	UCL conc.	MDL quotient	UCL quotient	Benchmark type
HEPTACHLOR (continued)	5	2	0	5.44	129	0.05	1232.35	0.45	.	.	NAHQ ACTUAL
	5	2	0	5.44	129	0.05	1232.35	0.45	.	.	LTV EC20 DAPHNID
	5	2	0	5.44	129	0.05	1232.35	0.45	11.71	111.86	0.00 SS EC20
	5	2	0	5.44	129	0.05	1232.35	0.45	4.68	44.74	0.01 NAHQ ADVISORY
	5	2	0	5.44	129	0.05	1232.35	0.45	0.47	4.47	0.10 EC20 POPULATION
	5	2	0	5.44	129	0.05	1232.35	0.45	0.05	0.52	0.86 LTV EC20 FISH
	5	2	0	5.44	129	0.05	1232.35	0.45	0.04	0.36	1.26 LCV FISH
	5	2	0	5.44	129	0.05	1232.35	0.45	0.01	0.14	3.18 LCV DAPHNIDS
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	NAHQ ADVISORY
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	NAHQ ACTUAL
INDENO(1,2,3-CD)PYRENE	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LCV FISH
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LCV DAPHNIDS
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LTV EC20 FISH
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	LTV EC20 DAPHNID
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	SS EC20
	1	2	0	6.58	620	0.02	6993.39	0.18	.	.	EC20 POPULATION
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	NAHQ ADVISORY
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	NAHQ ACTUAL
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LCV FISH
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LCV DAPHNIDS
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LTV EC20 FISH
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	LTV EC20 DAPHNID
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	SS EC20
	2	2	0	6.58	445	0.01	616.33	0.02	.	.	EC20 POPULATION
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	NAHQ ADVISORY
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	NAHQ ACTUAL
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LCV FISH
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LCV DAPHNIDS
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LTV EC20 FISH
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	LTV EC20 DAPHNID
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	SS EC20
	3	2	0	6.58	565	0.01	599.27	0.02	.	.	EC20 POPULATION
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	NAHQ ADVISORY
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	NAHQ ACTUAL
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LCV FISH
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LCV DAPHNIDS
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LTV EC20 FISH
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	LTV EC20 DAPHNID
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	SS EC20
	4	2	0	6.58	500	0.01	38534.72	1.01	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	HDL quotient	UCL quotient	Benchmark type
INDENO(1,2,3-CD)PYRENE (continued)	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	NAHQ ADVISORY
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	NAHQ ACTUAL
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LCV FISH
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LCV DAPHNIDS
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LTV EC20 FISH
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	LTV EC20 DAPHNID
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	SS EC20
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	EC20 POPULATION
	5	4	0	6.58	855	0.02	6848.50	0.18	.	.	NAHQ ADVISORY
	10	4	0	6.58	855	0.02	6848.50	0.18	.	.	NAHQ ACTUAL
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LCV FISH
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LCV DAPHNIDS
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LTV EC20 FISH
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	LTV EC20 DAPHNID
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	SS EC20
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	EC20 POPULATION
	10	1	0	6.58	6500	0.17	6500.00	0.17	.	.	NAHQ ADVISORY
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	NAHQ ACTUAL
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	LCV FISH
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	LCV DAPHNIDS
N-NITROSO-DI-N-PROPYLAMINE	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	LTV EC20 FISH
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	LTV EC20 DAPHNID
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	SS EC20
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	EC20 POPULATION
	1	2	0	1.36	620	2706.40	6993.39	30527.23	.	.	NAHQ ADVISORY
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	NAHQ ACTUAL
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	LCV FISH
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	LCV DAPHNIDS
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	LTV EC20 FISH
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	LTV EC20 DAPHNID
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	SS EC20
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	EC20 POPULATION
	2	2	0	1.36	445	1942.50	616.33	2690.37	.	.	NAHQ ADVISORY
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	NAHQ ACTUAL
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	LCV FISH
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	LCV DAPHNIDS
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	LTV EC20 FISH
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	LTV EC20 DAPHNID
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	SS EC20
	3	2	0	1.36	565	2466.31	599.27	2615.89	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
N-NITROSO-DI-N-PROPYLAMINE (continued)											
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	NAWQ ADVISORY
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	NAWQ ACTUAL
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	LCV FISH
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	LCV DAPHNIDS
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	LTV EC20 FISH
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	LTV EC20 DAPHNID
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	SS EC20
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	EC20 POPULATION
	4	2	0	1.36	500	2182.58	38534.72	168210.15	.	.	NAWQ ADVISORY
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	NAWQ ACTUAL
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	LCV FISH
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	LCV DAPHNIDS
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	LTV EC20 FISH
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	LTV EC20 DAPHNID
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	SS EC20
	5	4	0	1.36	855	3732.21	6848.50	29894.80	.	.	EC20 POPULATION
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	NAWQ ADVISORY
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	NAWQ ACTUAL
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	LCV FISH
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	LCV DAPHNIDS
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	LTV EC20 FISH
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	LTV EC20 DAPHNID
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	SS EC20
	10	1	0	1.36	6500	28373.53	6500.00	28373.53	.	.	EC20 POPULATION
N-NITROSO-DI-N-PROPYLAMINE											
	1	2	0	3.13	620	45.96	6993.39	518.43	.	.	NAWQ ACTUAL
	1	2	0	3.13	620	45.96	6993.39	518.43	.	.	LTV EC20 DAPHNID
	1	2	0	3.13	620	45.96	6993.39	518.43	.	.	SS EC20
	1	2	0	3.13	620	45.96	6993.39	518.43	207.37	2.50	NAWQ ADVISORY
	1	2	0	3.13	620	45.96	6993.39	518.43	12.96	40.00	EC20 POPULATION
	1	2	0	3.13	620	45.96	6993.39	518.43	1.56	332.00	LCV FISH
	1	2	0	3.13	620	45.96	6993.39	518.43	1.53	339.00	LTV EC20 FISH
	1	2	0	3.13	620	45.96	6993.39	518.43	0.50	1042.00	LCV DAPHNIDS
	1	2	0	3.13	445	32.99	616.33	45.69	.	.	NAWQ ACTUAL
	2	2	0	3.13	445	32.99	616.33	45.69	.	.	LTV EC20 DAPHNID
	2	2	0	3.13	445	32.99	616.33	45.69	18.28	2.50	NAWQ ADVISORY
	2	2	0	3.13	445	32.99	616.33	45.69	1.14	40.00	EC20 POPULATION
	2	2	0	3.13	445	32.99	616.33	45.69	0.82	332.00	LCV FISH
	2	2	0	3.13	445	32.99	616.33	45.69	0.10	339.00	LTV EC20 FISH
	2	2	0	3.13	445	32.99	616.33	45.69	0.13	1042.00	LCV DAPHNIDS
	2	2	0	3.13	445	32.99	616.33	45.69	0.04	0.04	LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
N-NITROSDIPHENYLAMINE (continued)	3	2	0	3.13	565	41.88	599.27	44.42	.	.	NAWQ ACTUAL
	3	2	0	3.13	565	41.88	599.27	44.42	.	.	LTV EC20 DAPHNID
	3	2	0	3.13	565	41.88	599.27	44.42	.	.	SS EC20
	3	2	0	3.13	565	41.88	599.27	44.42	16.75	17.77	2.50 NAWQ ADVISORY
	3	2	0	3.13	565	41.88	599.27	44.42	1.05	1.11	40.00 EC20 POPULATION
	3	2	0	3.13	565	41.88	599.27	44.42	0.13	0.13	332.00 LCV FISH
	3	2	0	3.13	565	41.88	599.27	44.42	0.12	0.13	339.00 LTV EC20 FISH
	3	2	0	3.13	565	41.88	599.27	44.42	0.04	0.04	1042.00 LCV DAPHNIDS
	3	2	0	3.13	565	41.88	599.27	44.42	.	.	NAWQ ACTUAL
	4	2	0	3.13	500	37.07	38534.72	2856.62	.	.	LTV EC20 DAPHNID
	4	2	0	3.13	500	37.07	38534.72	2856.62	.	.	SS EC20
	4	2	0	3.13	500	37.07	38534.72	2856.62	14.83	1142.65	2.50 NAWQ ADVISORY
	4	2	0	3.13	500	37.07	38534.72	2856.62	0.93	71.42	40.00 EC20 POPULATION
	4	2	0	3.13	500	37.07	38534.72	2856.62	0.11	8.60	332.00 LCV FISH
	4	2	0	3.13	500	37.07	38534.72	2856.62	0.11	8.43	339.00 LTV EC20 FISH
	4	2	0	3.13	500	37.07	38534.72	2856.62	0.04	2.74	1042.00 LCV DAPHNIDS
	4	2	0	3.13	500	37.07	38534.72	2856.62	.	.	NAWQ ACTUAL
	5	4	0	3.13	855	63.38	6848.50	507.69	.	.	LTV EC20 DAPHNID
	5	4	0	3.13	855	63.38	6848.50	507.69	.	.	SS EC20
	5	4	0	3.13	855	63.38	6848.50	507.69	25.35	203.07	2.50 NAWQ ADVISORY
	5	4	0	3.13	855	63.38	6848.50	507.69	1.58	12.69	40.00 EC20 POPULATION
	5	4	0	3.13	855	63.38	6848.50	507.69	0.19	1.53	332.00 LCV FISH
	5	4	0	3.13	855	63.38	6848.50	507.69	0.19	1.50	339.00 LTV EC20 FISH
	5	4	0	3.13	855	63.38	6848.50	507.69	0.06	0.49	1042.00 LCV DAPHNIDS
	5	4	0	3.13	855	63.38	6848.50	507.69	.	.	NAWQ ACTUAL
	10	1	0	3.13	6500	481.85	6500.00	481.85	.	.	LTV EC20 DAPHNID
	10	1	0	3.13	6500	481.85	6500.00	481.85	.	.	SS EC20
	10	1	0	3.13	6500	481.85	6500.00	481.85	192.74	192.74	2.50 NAWQ ADVISORY
	10	1	0	3.13	6500	481.85	6500.00	481.85	12.05	12.05	40.00 EC20 POPULATION
	10	1	0	3.13	6500	481.85	6500.00	481.85	1.45	1.45	332.00 LCV FISH
	10	1	0	3.13	6500	481.85	6500.00	481.85	1.42	1.42	339.00 LTV EC20 FISH
	10	1	0	3.13	6500	481.85	6500.00	481.85	0.46	0.46	1042.00 LCV DAPHNIDS
	10	1	0	3.13	6500	481.85	6500.00	481.85	.	.	NAWQ ACTUAL
NAPHTHALENE	1	2	0	3.59	620	15.94	6993.39	179.76	.	.	SS EC20
	1	2	0	3.59	620	15.94	6993.39	179.76	7.59	85.60	2.10 NAWQ ADVISORY
	1	2	0	3.59	620	15.94	6993.39	179.76	0.04	0.40	450.00 LTV EC20 FISH
	1	2	0	3.59	620	15.94	6993.39	179.76	0.03	0.30	600.00 LTV EC20 DAPHNID
	1	2	0	3.59	620	15.94	6993.39	179.76	0.03	0.29	620.00 LCV FISH
	1	2	0	3.59	620	15.94	6993.39	179.76	0.02	0.18	1000.00 EC20 POPULATION
	1	2	0	3.59	620	15.94	6993.39	179.76	0.01	0.15	1163.00 LCV DAPHNIDS
	1	2	0	3.59	620	15.94	6993.39	179.76	.	.	NAWQ ACTUAL
	1	2	0	3.59	620	15.94	6993.39	179.76	.	.	SS EC20
	1	2	0	3.59	620	15.94	6993.39	179.76	7.59	85.60	2.10 NAWQ ADVISORY

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL conc.	MDL quotient	UCL quotient	Benchmark type
NAPHTHALENE (continued)	2	2	0	3.59	445	11.44	616.33	15.84	.	.	NAHQ ACTUAL
	2	2	0	3.59	445	11.44	616.33	15.84	.	.	SS EC20
	2	2	0	3.59	445	11.44	616.33	15.84	5.45	7.54	2.10 NAHQ ADVISORY
	2	2	0	3.59	445	11.44	616.33	15.84	0.03	0.04	450.00 LTV EC20 FISH
	2	2	0	3.59	445	11.44	616.33	15.84	0.02	0.03	600.00 LTV EC20 DAPHNID
	2	2	0	3.59	445	11.44	616.33	15.84	0.02	0.03	620.00 LCV FISH
	2	2	0	3.59	445	11.44	616.33	15.84	0.01	0.02	1000.00 EC20 POPULATION
	2	2	0	3.59	445	11.44	616.33	15.84	0.01	0.01	1163.00 LCV DAPHNIDS
	2	2	0	3.59	445	11.44	616.33	15.84	.	.	NAHQ ACTUAL
	4	4	0	3.59	500	12.85	38534.72	990.49	.	.	SS EC20
	4	4	0	3.59	500	12.85	38534.72	990.49	6.12	471.66	2.10 NAHQ ADVISORY
	4	4	0	3.59	500	12.85	38534.72	990.49	0.03	2.20	450.00 LTV EC20 FISH
	4	4	0	3.59	500	12.85	38534.72	990.49	0.02	1.65	600.00 LTV EC20 DAPHNID
	4	4	0	3.59	500	12.85	38534.72	990.49	0.02	1.60	620.00 LCV FISH
	4	4	0	3.59	500	12.85	38534.72	990.49	0.01	0.99	1000.00 EC20 POPULATION
	4	4	0	3.59	500	12.85	38534.72	990.49	0.01	0.85	1163.00 LCV DAPHNIDS
	4	4	0	3.59	500	12.85	38534.72	990.49	.	.	NAHQ ACTUAL
	5	5	0	3.59	855	21.98	6848.50	176.03	.	.	SS EC20
	5	5	0	3.59	855	21.98	6848.50	176.03	10.47	83.83	2.10 NAHQ ADVISORY
	5	5	0	3.59	855	21.98	6848.50	176.03	0.05	0.39	450.00 LTV EC20 FISH
	5	5	0	3.59	855	21.98	6848.50	176.03	0.04	0.29	600.00 LTV EC20 DAPHNID
	5	5	0	3.59	855	21.98	6848.50	176.03	0.04	0.28	620.00 LCV FISH
	5	5	0	3.59	855	21.98	6848.50	176.03	0.02	0.18	1000.00 EC20 POPULATION
	5	5	0	3.59	855	21.98	6848.50	176.03	0.02	0.15	1163.00 LCV DAPHNIDS
	10	10	0	3.59	6500	167.08	6500.00	167.08	.	.	NAHQ ACTUAL
	10	10	0	3.59	6500	167.08	6500.00	167.08	79.56	79.56	2.10 NAHQ ADVISORY
	10	10	0	3.59	6500	167.08	6500.00	167.08	0.37	0.37	450.00 LTV EC20 FISH
	10	10	0	3.59	6500	167.08	6500.00	167.08	0.28	0.28	600.00 LTV EC20 DAPHNID
	10	10	0	3.59	6500	167.08	6500.00	167.08	0.27	0.27	620.00 LCV FISH
	10	10	0	3.59	6500	167.08	6500.00	167.08	0.17	0.17	1000.00 EC20 POPULATION
	10	10	0	3.59	6500	167.08	6500.00	167.08	0.14	0.14	1163.00 LCV DAPHNIDS

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
PHENANTHRENE	1	2	0	4.57	620	1.67	6993.39	18.82	.	.	NAHQ ACTUAL
	1	2	0	4.57	620	1.67	6993.39	18.82	.	.	LCV FISH
	1	2	0	4.57	620	1.67	6993.39	18.82	.	.	LTV EC20 FISH
	1	2	0	4.57	620	1.67	6993.39	18.82	.	.	SS EC20
	1	2	0	4.57	620	1.67	6993.39	18.82	.	.	EC20 POPULATION
	1	2	0	4.57	620	1.67	6993.39	18.82	3.88	43.77	0.43 NAHQ ADVISORY
	1	2	0	4.57	620	1.67	6993.39	18.82	0.02	0.17	110.00 LTV EC20 DAPHNID
	1	2	0	4.57	620	1.67	6993.39	18.82	0.01	0.09	200.00 LCV DAPHNIDS
	1	2	0	4.57	620	1.67	6993.39	18.82	.	.	NAHQ ACTUAL
	2	2	0	4.57	445	1.20	616.33	1.66	.	.	LCV FISH
	2	2	0	4.57	445	1.20	616.33	1.66	.	.	LTV EC20 FISH
	2	2	0	4.57	445	1.20	616.33	1.66	.	.	SS EC20
	2	2	0	4.57	445	1.20	616.33	1.66	.	.	EC20 POPULATION
	2	2	0	4.57	445	1.20	616.33	1.66	2.79	3.86	0.43 NAHQ ADVISORY
	2	2	0	4.57	445	1.20	616.33	1.66	0.01	0.02	110.00 LTV EC20 DAPHNID
	2	2	0	4.57	445	1.20	616.33	1.66	0.01	0.01	200.00 LCV DAPHNIDS
	2	2	0	4.57	445	1.20	616.33	1.66	.	.	NAHQ ACTUAL
	4	2	0	4.57	500	1.35	38534.72	103.72	.	.	LCV FISH
	4	2	0	4.57	500	1.35	38534.72	103.72	.	.	LTV EC20 FISH
	4	2	0	4.57	500	1.35	38534.72	103.72	.	.	SS EC20
	4	2	0	4.57	500	1.35	38534.72	103.72	.	.	EC20 POPULATION
	4	2	0	4.57	500	1.35	38534.72	103.72	3.13	241.20	0.43 NAHQ ADVISORY
	4	2	0	4.57	500	1.35	38534.72	103.72	0.01	0.94	110.00 LTV EC20 DAPHNID
	4	2	0	4.57	500	1.35	38534.72	103.72	0.01	0.52	200.00 LCV DAPHNIDS
	4	2	0	4.57	500	1.35	38534.72	103.72	.	.	NAHQ ACTUAL
	5	4	0	4.57	855	2.30	6848.50	18.43	.	.	LCV FISH
	5	4	0	4.57	855	2.30	6848.50	18.43	.	.	LTV EC20 FISH
	5	4	0	4.57	855	2.30	6848.50	18.43	.	.	SS EC20
	5	4	0	4.57	855	2.30	6848.50	18.43	.	.	EC20 POPULATION
	5	4	0	4.57	855	2.30	6848.50	18.43	5.35	42.87	0.43 NAHQ ADVISORY
	5	4	0	4.57	855	2.30	6848.50	18.43	0.02	0.17	110.00 LTV EC20 DAPHNID
	5	4	0	4.57	855	2.30	6848.50	18.43	0.01	0.09	200.00 LCV DAPHNIDS
	5	4	0	4.57	855	2.30	6848.50	18.43	.	.	NAHQ ACTUAL
	10	1	0	4.57	6500	17.49	6500.00	17.49	.	.	LCV FISH
	10	1	0	4.57	6500	17.49	6500.00	17.49	.	.	LTV EC20 FISH
	10	1	0	4.57	6500	17.49	6500.00	17.49	.	.	SS EC20
	10	1	0	4.57	6500	17.49	6500.00	17.49	.	.	EC20 POPULATION
	10	1	0	4.57	6500	17.49	6500.00	17.49	40.69	40.69	0.43 NAHQ ADVISORY
	10	1	0	4.57	6500	17.49	6500.00	17.49	0.16	0.16	110.00 LTV EC20 DAPHNID
	10	1	0	4.57	6500	17.49	6500.00	17.49	0.09	0.09	200.00 LCV DAPHNIDS
	10	1	0	4.57	6500	17.49	6500.00	17.49	.	.	NAHQ ACTUAL

Table F.10 (continued)

Analyte	Reach (sediment)	Number		Log coefficient between H ₂ O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type	Benchmark
		analyzed in reach	Number detects in reach (sediment)									
PHENOL	1	2	0	1.46	620	2149.77	6993.39	24248.64	.	.	.	NAWQ ACTUAL
	1	2	0	1.46	620	2149.77	6993.39	24248.64	.	.	.	LTV EC20 DAPHNID
	1	2	0	1.46	620	2149.77	6993.39	24248.64	.	.	.	SS EC20
	1	2	0	1.46	620	2149.77	6993.39	24248.64	102.37	1154.70	21.00	NAWQ ADVISORY
	1	2	0	1.46	620	2149.77	6993.39	24248.64	9.35	105.43	230.00	LTV EC20 FISH
	1	2	0	1.46	620	2149.77	6993.39	24248.64	1.57	17.71	1369.00	LCV FISH
	1	2	0	1.46	620	2149.77	6993.39	24248.64	1.07	12.09	2005.00	LCV DAPHNIDS
	1	2	0	1.46	620	2149.77	6993.39	24248.64	0.48	5.43	4467.00	EC20 POPULATION
	1	2	0	1.46	620	2149.77	6993.39	24248.64	.	.	.	NAWQ ACTUAL
	2	2	0	1.46	445	1542.98	616.33	2137.03	.	.	.	LTV EC20 DAPHNID
	2	2	0	1.46	445	1542.98	616.33	2137.03	.	.	.	SS EC20
	2	2	0	1.46	445	1542.98	616.33	2137.03	73.48	101.76	21.00	NAWQ ADVISORY
	2	2	0	1.46	445	1542.98	616.33	2137.03	6.71	9.29	230.00	LTV EC20 FISH
	2	2	0	1.46	445	1542.98	616.33	2137.03	1.13	1.56	1369.00	LCV FISH
	2	2	0	1.46	445	1542.98	616.33	2137.03	0.77	1.07	2005.00	LCV DAPHNIDS
	2	2	0	1.46	445	1542.98	616.33	2137.03	0.35	0.48	4467.00	EC20 POPULATION
	2	2	0	1.46	445	1542.98	616.33	2137.03	.	.	.	NAWQ ACTUAL
	3	2	0	1.46	565	1959.06	599.27	2077.87	.	.	.	LTV EC20 DAPHNID
	3	2	0	1.46	565	1959.06	599.27	2077.87	.	.	.	SS EC20
	3	2	0	1.46	565	1959.06	599.27	2077.87	93.29	98.95	21.00	NAWQ ADVISORY
	3	2	0	1.46	565	1959.06	599.27	2077.87	8.52	9.03	230.00	LTV EC20 FISH
	3	2	0	1.46	565	1959.06	599.27	2077.87	1.43	1.52	1369.00	LCV FISH
	3	2	0	1.46	565	1959.06	599.27	2077.87	0.98	1.04	2005.00	LCV DAPHNIDS
	3	2	0	1.46	565	1959.06	599.27	2077.87	0.44	0.47	4467.00	EC20 POPULATION
	3	2	0	1.46	565	1959.06	599.27	2077.87	.	.	.	NAWQ ACTUAL
	4	2	0	1.46	500	1733.68	38534.72	133614.07	.	.	.	LTV EC20 DAPHNID
	4	2	0	1.46	500	1733.68	38534.72	133614.07	.	.	.	SS EC20
	4	2	0	1.46	500	1733.68	38534.72	133614.07	82.56	6362.57	21.00	NAWQ ADVISORY
	4	2	0	1.46	500	1733.68	38534.72	133614.07	7.54	580.93	230.00	LTV EC20 FISH
	4	2	0	1.46	500	1733.68	38534.72	133614.07	1.27	97.60	1369.00	LCV FISH
	4	2	0	1.46	500	1733.68	38534.72	133614.07	0.86	66.64	2005.00	LCV DAPHNIDS
	4	2	0	1.46	500	1733.68	38534.72	133614.07	0.39	29.91	4467.00	EC20 POPULATION
	4	2	0	1.46	500	1733.68	38534.72	133614.07	.	.	.	NAWQ ACTUAL
	5	4	0	1.46	855	2964.60	6848.50	23746.28	.	.	.	LTV EC20 DAPHNID
	5	4	0	1.46	855	2964.60	6848.50	23746.28	.	.	.	SS EC20
	5	4	0	1.46	855	2964.60	6848.50	23746.28	141.17	1130.78	21.00	NAWQ ADVISORY
	5	4	0	1.46	855	2964.60	6848.50	23746.28	12.89	103.24	230.00	LTV EC20 FISH
	5	4	0	1.46	855	2964.60	6848.50	23746.28	2.17	17.35	1369.00	LCV FISH
	5	4	0	1.46	855	2964.60	6848.50	23746.28	1.48	11.84	2005.00	LCV DAPHNIDS
	5	4	0	1.46	855	2964.60	6848.50	23746.28	0.66	5.32	4467.00	EC20 POPULATION
	5	4	0	1.46	855	2964.60	6848.50	23746.28	.	.	.	NAWQ ACTUAL

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach (sediment)	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
PHENOL (continued)	10	1	0	1.46	6500	22537.90	6500.00	22537.90	.	.	NAHQ ACTUAL
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	.	.	LTV EC20 DAPHNID
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	.	.	SS EC20
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	1073.23	1073.23	21.00 NAHQ ADVISORY
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	97.99	97.99	230.00 LTV EC20 FISH
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	16.46	16.46	1369.00 LCV FISH
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	11.24	11.24	2005.00 LCV DAPHNIDS
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	5.05	5.05	4467.00 EC20 POPULATION
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	.	.	NAHQ ADVISORY
	10	1	0	1.46	6500	22537.90	6500.00	22537.90	.	.	NAHQ ACTUAL
PYRENE	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	NAHQ ACTUAL
	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	NAHQ FISH
	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	LCV DAPHNIDS
	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	LTV EC20 FISH
	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	LTV EC20 DAPHNID
	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	SS EC20
	1	2	0	4.88	620	0.82	6993.39	9.22	.	.	EC20 POPULATION
	1	2	0	4.88	445	0.59	616.33	0.81	.	.	NAHQ ADVISORY
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	NAHQ ACTUAL
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	LCV FISH
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	LCV DAPHNIDS
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	LTV EC20 FISH
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	LTV EC20 DAPHNID
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	SS EC20
	2	2	0	4.88	445	0.59	616.33	0.81	.	.	EC20 POPULATION
	2	2	0	4.88	500	0.66	38534.72	50.80	.	.	NAHQ ADVISORY
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	NAHQ ACTUAL
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	LCV FISH
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	LCV DAPHNIDS
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	LTV EC20 FISH
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	LTV EC20 DAPHNID
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	SS EC20
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	EC20 POPULATION
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	NAHQ ADVISORY
	4	2	0	4.88	500	0.66	38534.72	50.80	.	.	NAHQ ACTUAL
	5	4	0	4.88	855	1.13	6848.50	9.03	.	.	LCV FISH
	5	4	0	4.88	855	1.13	6848.50	9.03	.	.	LCV DAPHNIDS
	5	4	0	4.88	855	1.13	6848.50	9.03	.	.	LTV EC20 FISH
	5	4	0	4.88	855	1.13	6848.50	9.03	.	.	LTV EC20 DAPHNID
	5	4	0	4.88	855	1.13	6848.50	9.03	.	.	SS EC20
	5	4	0	4.88	855	1.13	6848.50	9.03	.	.	EC20 POPULATION

Table F.10 (continued)

Analyte	Reach (sediment)	Number analyzed in reach	Number detects in reach (sediment)	Log coefficient between H2O and SED	Minimum detection limit	Minimum pore water conc.	UCL ^a detection limit	UCL pore water conc.	MDL quotient	UCL quotient	Benchmark type
PYRENE (continued)	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	NAHQ ADVISORY
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	NAHQ ACTUAL
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	LCV FISH
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	LCV DAPHNIDS
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	LTV EC20 FISH
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	LTV EC20 DAPHNID
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	SS EC20
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	EC20 POPULATION
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	
	10	1	0	4.88	6500	8.57	6500.00	8.57	.	.	

^aThe maximum detection limit is used in cases where the UCL detection limit was missing.

Table F.11. Mean and upper confidence limit (UCL) concentrations of chemicals in fish filets (mg/kg), toxicological benchmarks for wildlife (mg/kg), and quotients for hazards to piscivorous wildlife

Analyte	Reach	Number analyzed in reach (fish)	Number detected in reach (fish)	Mean ^a fish concentration	UCL ^b fish concentration	Lower piscivore criterion	Mean piscivore quotient	UCL piscivore quotient	Reference dose	RFD dietary concentration	Mean RFD piscivore quotient	UCL RFD piscivore quotient
4,4'-DDO	1	15	5	0.00085	0.006	0.01	0.0849	0.5900	.	.	.	.
	3	34	2	0.00030	0.017	0.01	0.0300	1.7080	.	.	.	.
	4	18	4	0.00367	0.009	0.01	0.3671	0.9039	.	.	.	.
	5	17	12	0.00155	0.006	0.01	0.1555	0.5826	.	.	.	.
	10	8	7	0.00069	0.001	0.01	0.0687	0.1219	.	.	.	.
	15	4	1	0.00474	0.200	0.01	0.4738	20.0000	.	.	.	.
4,4'-DDE	1	16	8	0.00530	0.064	0.01	0.5298	6.4285	.	.	.	.
	2	13	6	0.00529	0.018	0.01	0.5292	1.8272	.	.	.	.
	3	34	34	0.00104	0.009	0.01	0.1044	0.9104	.	.	.	.
	4	18	12	0.01109	0.024	0.01	1.1089	2.4405	.	.	.	.
	5	18	9	0.00803	0.013	0.01	0.8030	1.3229	.	.	.	.
	10	8	2	0.00256	0.003	0.01	0.2561	0.2948	.	.	.	.
4,4'-DDT	13	2	2	0.00387	0.007	0.01	0.3873	0.6820	.	.	.	.
	2	13	2	0.00047	0.001	0.01	0.0467	0.0859	0.00050	0.0063	0.075	0.137
	4	18	1	0.00158	0.200	0.01	0.1577	20.0000	0.00050	0.0063	0.252	32.000
	5	18	5	0.00070	0.001	0.01	0.0703	0.1291	0.00050	0.0063	0.113	0.207
	2	14	2	0.00013	0.000	.	.	.	0.00003	0.0004	0.343	0.440
	4	14	2	0.00085	0.001	.	.	.	0.00003	0.0004	2.265	3.559
ALDRIN	5	18	2	0.00046	0.005	.	.	.	0.00003	0.0004	1.238	14.537
	13	3	2	0.00096	0.001	.	.	.	0.00003	0.0004	2.561	3.478
AROCFLOR-1254	1	16	13	0.03328	0.199	0.04	0.8320	4.9765	.	.	.	.
	2	10	9	0.12149	0.753	0.04	3.0373	18.8272	.	.	.	.
	3	31	31	0.09903	0.186	0.04	2.4757	4.6583	.	.	.	.
	4	18	15	0.05439	0.270	0.04	1.3596	6.7500	.	.	.	.
	5	17	14	0.05830	0.320	0.04	1.4576	8.0000	.	.	.	.
	13	2	2	0.08718	0.190	0.04	2.1794	4.7500	.	.	.	.
AROCFLOR-1260	15	4	4	0.12234	0.200	0.04	3.0585	5.0000	.	.	.	.
	1	16	16	0.24586	0.897	0.04	6.1464	22.4140	.	.	.	.
	2	14	14	0.66853	1.406	0.04	16.7132	35.1598	.	.	.	.
	3	36	36	0.39085	0.771	0.04	9.7714	19.2634	.	.	.	.
	4	18	17	0.43314	2.790	0.04	10.8284	69.7380	.	.	.	.
	5	19	19	0.34776	0.890	0.04	8.6939	22.2399	.	.	.	.
AROCFLOR-1260	10	8	7	0.02198	0.073	0.04	0.5494	1.8275	.	.	.	.
	13	4	4	0.28812	0.390	0.04	7.2030	9.7500	.	.	.	.
	15	4	4	0.31050	0.650	0.04	7.7625	16.2500	.	.	.	.

Table F.11 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number defects in reach (fish)	Mean ^f fish concentration	UCL ^b fish concentration	Lower piscivore criterion	Mean piscivore quotient	UCL piscivore quotient	Reference dose	RFD dietary concentration	Mean RFD piscivore quotient	UCL RFD piscivore quotient
ARSENIC	1	28	17	0.06163	0.098	3.00	0.0205	0.0327	0.00100	0.0125	4.930	7.858
	2	13	5	0.03796	0.098	3.00	0.0127	0.0327	0.00100	0.0125	3.036	7.842
	3	35	19	0.05308	0.075	3.00	0.0177	0.0250	0.00100	0.0125	4.246	6.004
	4	27	25	0.08793	0.107	3.00	0.0293	0.0357	0.00100	0.0125	7.034	8.578
	5	25	23	0.07461	0.092	3.00	0.0249	0.0308	0.00100	0.0125	5.969	7.382
	10	15	14	0.07143	0.097	3.00	0.0238	0.0323	0.00100	0.0125	5.714	7.745
	13	10	4	0.04514	0.059	3.00	0.0150	0.0197	0.00100	0.0125	3.611	4.721
	15	16	10	0.06469	0.113	3.00	0.0216	0.0376	0.00100	0.0125	5.175	9.014
	1	25	4	0.00147	0.003	25.00	0.0001	0.0001	0.00500	0.0625	0.023	0.052
	3	35	7	0.00092	0.005	25.00	0.0000	0.0002	0.00500	0.0625	0.015	0.087
	5	25	2	0.00002	0.027	25.00	0.0000	0.0011	0.00500	0.0625	0.000	0.432
	10	10	1	0.00001	0.037	25.00	0.0000	0.0015	0.00500	0.0625	0.000	0.592
	15	16	5	0.00132	0.019	25.00	0.0001	0.0008	0.00500	0.0625	0.021	0.304
	3	33	1	0.04709	0.410	.	.	.	.	.	.	.
	1	16	11	0.01647	0.109	.	.	.	.	.	.	.
BIS(2-ETHYLHEXYL)PHTHALATE	2	14	9	0.01144	0.082	.	.	.	.	.	.	.
	3	34	34	0.03989	0.074	.	.	.	.	.	.	.
	4	18	18	0.02838	0.097	.	.	.	.	.	.	.
	5	18	18	0.01062	0.045	.	.	.	.	.	.	.
	10	8	8	0.00832	0.016	.	.	.	.	.	.	.
	13	4	2	0.01369	0.073	.	.	.	.	.	.	.
	15	4	4	0.01225	0.022	.	.	.	.	.	.	.
	5	25	6	0.04648	170.000	50.00	0.0009	3.4000	1.30000	16.2500	0.003	10.462
	3	33	1	0.02405	0.410	.	.	.	.	.	.	.
	1	28	28	0.05367	0.082	0.01	5.3673	8.1945	0.00030	0.0037	14.313	21.852
	2	13	13	0.06384	0.135	0.01	6.3843	13.4967	0.00030	0.0037	17.025	35.991
	3	35	35	0.42259	0.616	0.01	42.2590	61.5506	0.00030	0.0037	112.691	164.135
	4	27	27	0.18534	0.391	0.01	18.5336	39.0647	0.00030	0.0037	49.423	104.173
	5	25	25	0.06598	0.108	0.01	6.5984	10.7910	0.00030	0.0037	17.596	28.776
	10	15	15	0.19008	0.280	0.01	19.0085	27.9963	0.00030	0.0037	50.689	74.657
	13	10	10	0.10551	0.367	0.01	10.5512	36.7416	0.00030	0.0037	28.137	97.978
	15	16	16	0.13209	0.218	0.01	13.2087	21.8292	0.00030	0.0037	35.223	58.211
COPPER	5	25	6	0.04648	170.000	50.00	0.0009	3.4000	1.30000	16.2500	0.003	10.462
	3	33	1	0.02405	0.410	.	.	.	.	.	.	.
DI-N-OCTYLPHTHALATE	1	28	28	0.05367	0.082	0.01	5.3673	8.1945	0.00030	0.0037	14.313	21.852
	2	13	13	0.06384	0.135	0.01	6.3843	13.4967	0.00030	0.0037	17.025	35.991
	3	35	35	0.42259	0.616	0.01	42.2590	61.5506	0.00030	0.0037	112.691	164.135
	4	27	27	0.18534	0.391	0.01	18.5336	39.0647	0.00030	0.0037	49.423	104.173
	5	25	25	0.06598	0.108	0.01	6.5984	10.7910	0.00030	0.0037	17.596	28.776
	10	15	15	0.19008	0.280	0.01	19.0085	27.9963	0.00030	0.0037	50.689	74.657
MERCURY	13	10	10	0.10551	0.367	0.01	10.5512	36.7416	0.00030	0.0037	28.137	97.978
	15	16	16	0.13209	0.218	0.01	13.2087	21.8292	0.00030	0.0037	35.223	58.211

Table F.11 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Mean ^a fish concentration	UCL ^b fish concentration	Lower piscivore criterion	Mean piscivore quotient	UCL piscivore quotient	Reference dose	RFD dietary concentration	Mean RFD piscivore quotient	UCL RFD piscivore quotient
SELENIUM	1	28	26	0.74833	0.834	0.50	1.4967	1.6690	0.00300	0.0375	19.955	22.253
	2	13	11	0.72604	0.851	0.50	1.4521	1.7014	0.00300	0.0375	19.361	22.685
	3	35	2	0.23139	0.577	0.50	0.4628	1.1538	0.00300	0.0375	6.170	15.383
	4	27	14	0.55653	0.686	0.50	1.1131	1.3727	0.00300	0.0375	14.841	18.303
	10	15	8	0.52186	0.582	0.50	1.0437	1.1650	0.00300	0.0375	13.916	15.533
	15	16	8	0.52464	0.679	0.50	1.0493	1.3590	0.00300	0.0375	13.990	18.119
ZINC	1	28	28	6.58972	7.546	60.00	0.1098	0.1258	0.20000	2.5000	2.636	3.019
	2	13	13	5.74198	6.566	60.00	0.0957	0.1094	0.20000	2.5000	2.297	2.626
	3	35	35	5.76390	6.551	60.00	0.0961	0.1092	0.20000	2.5000	2.306	2.620
	4	27	27	5.74666	6.545	60.00	0.0958	0.1091	0.20000	2.5000	2.299	2.618
	5	24	24	6.97721	10.930	60.00	0.1163	0.1822	0.20000	2.5000	2.791	4.372
	10	15	15	5.22138	5.967	60.00	0.0870	0.0994	0.20000	2.5000	2.089	2.387
	13	10	10	6.31303	6.977	60.00	0.1052	0.1163	0.20000	2.5000	2.525	2.791
	15	16	16	5.77205	6.899	60.00	0.0962	0.1150	0.20000	2.5000	2.309	2.760

^aThe mean concentration is the geometric mean.^bUCL concentration was compared with the maximum value and the smaller has been used as the "UCL concentration."

Table F-12. Minimum and upper confidence limit (UCL) detection limits for chemicals not detected in fish filets (mg/kg), toxicological benchmarks for wildlife (mg/kg), and quotients for hazards to piscivorous wildlife

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ^a detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
1,2,4-TRICHLOROBENZENE	1	6	0	0.68000	0.87000	0.00131	0.0164	.	0.0164	41.527	53.130
	2	8	0	0.70000	1.90000	0.00131	0.0164	.	0.0164	42.748	116.031
	3	34	0	0.19000	0.41000	0.00131	0.0164	.	0.0164	11.603	25.038
	4	6	0	0.66000	0.72000	0.00131	0.0164	.	0.0164	40.305	43.969
	15	3	0	0.67000	0.71000	0.00131	0.0164	.	0.0164	40.916	43.359
1,2-DICHLOROBENZENE	1	6	0	0.68000	0.87000	0.09000	1.1250	.	1.1250	0.604	0.773
	2	8	0	0.70000	1.90000	0.09000	1.1250	.	1.1250	0.622	1.689
	3	34	0	0.19000	0.41000	0.09000	1.1250	.	1.1250	0.169	0.364
	4	6	0	0.66000	0.72000	0.09000	1.1250	.	1.1250	0.587	0.640
	15	3	0	0.67000	0.71000	0.09000	1.1250	.	1.1250	0.596	0.631
1,3-DICHLOROBENZENE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
1,4-DICHLOROBENZENE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
2,4,5-TRICHLOROPHENOL	1	6	0	3.40000	4.30000	0.10000	1.2500	.	1.2500	2.720	3.440
	2	8	0	3.50000	9.30000	0.10000	1.2500	.	1.2500	2.800	7.440
	3	34	0	0.93000	2.10000	0.10000	1.2500	.	1.2500	0.744	1.680
	4	13	0	3.30000	4.10000	0.10000	1.2500	.	1.2500	2.640	3.280
	5	9	0	3.30000	4.20000	0.10000	1.2500	.	1.2500	2.640	3.360
	13	2	0	3.40000	3.50000	0.10000	1.2500	.	1.2500	2.720	2.800
	15	4	0	3.40000	3.80000	0.10000	1.2500	.	1.2500	2.720	3.040
2,4,6-TRICHLOROPHENOL	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	13	0	0.66000	0.83000	.	.	.	.	.	.
	5	9	0	0.66000	0.83000	.	.	.	.	.	.
	13	2	0	0.69000	0.70000	.	.	.	.	.	.
	15	4	0	0.67000	0.77000	.	.	.	.	.	.

Table F.12 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
2,4-DICHLOROPHENOL	1	6	0	0.68000	0.87000	0.00300	0.0375	.	0.0375	18.133	23.200
	2	8	0	0.70000	1.90000	0.00300	0.0375	.	0.0375	18.667	50.667
	3	34	0	0.19000	0.41000	0.00300	0.0375	.	0.0375	5.067	10.933
	4	14	0	0.66000	0.83000	0.00300	0.0375	.	0.0375	17.600	22.133
	5	9	0	0.66000	0.83000	0.00300	0.0375	.	0.0375	17.600	22.133
	13	2	0	0.69000	0.70000	0.00300	0.0375	.	0.0375	18.400	18.667
	15	4	0	0.67000	0.77000	0.00300	0.0375	.	0.0375	17.867	20.533
	1	6	0	0.68000	0.87000	0.02000	0.2500	.	0.2500	2.720	3.480
	2	8	0	0.70000	1.90000	0.02000	0.2500	.	0.2500	2.800	7.600
	3	34	0	0.19000	0.41000	0.02000	0.2500	.	0.2500	1.640	1.640
2,4-DIMETHYLPHENOL	1	6	0	0.68000	0.87000	0.02000	0.2500	.	0.2500	2.720	3.480
	2	8	0	0.70000	1.90000	0.02000	0.2500	.	0.2500	2.800	7.600
	3	34	0	0.19000	0.41000	0.02000	0.2500	.	0.2500	1.640	1.640
	4	14	0	0.66000	0.83000	0.02000	0.2500	.	0.2500	2.640	3.320
	5	9	0	0.66000	0.83000	0.02000	0.2500	.	0.2500	2.640	3.320
	13	2	0	0.69000	0.70000	0.02000	0.2500	.	0.2500	2.760	2.800
	15	4	0	0.67000	0.77000	0.02000	0.2500	.	0.2500	2.680	3.080
	1	6	0	3.40000	4.30000	.	.	.	.	.	.
	2	8	0	3.50000	9.30000	.	.	.	.	.	.
	3	34	0	0.93000	2.10000	.	.	.	.	.	.
2,4-DINITROTOLUENE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	6	3	0	0.67000	0.71000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
2,6-DINITROTOLUENE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	6	3	0	0.67000	0.71000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	0.08000	1.0000	.	1.0000	0.680	0.870
	2	8	0	0.70000	1.90000	0.08000	1.0000	.	1.0000	0.700	1.900
	3	34	0	0.19000	0.41000	0.08000	1.0000	.	1.0000	0.190	0.410
	4	6	0	0.66000	0.72000	0.08000	1.0000	.	1.0000	0.660	0.720
2-CHLORONAPHTHALENE	1	6	0	0.68000	0.87000	0.08000	1.0000	.	1.0000	0.680	0.870
	2	8	0	0.70000	1.90000	0.08000	1.0000	.	1.0000	0.700	1.900
	3	34	0	0.19000	0.41000	0.08000	1.0000	.	1.0000	0.190	0.410
	4	6	0	0.66000	0.72000	0.08000	1.0000	.	1.0000	0.660	0.720
	6	3	0	0.67000	0.71000	0.08000	1.0000	.	1.0000	0.670	0.710
	15	3	0	0.67000	0.71000	0.08000	1.0000	.	1.0000	0.670	0.710
	1	6	0	0.68000	0.87000	0.08000	1.0000	.	1.0000	0.680	0.870
	2	8	0	0.70000	1.90000	0.08000	1.0000	.	1.0000	0.700	1.900
	3	34	0	0.19000	0.41000	0.08000	1.0000	.	1.0000	0.190	0.410
	4	6	0	0.66000	0.72000	0.08000	1.0000	.	1.0000	0.660	0.720

Table F.12 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
3-NITROANILINE	1	6	0	3.40000	4.30000	.	.	.	.	.	.
	3	27	0	0.93000	2.10000	.	.	.	.	.	.
	4	2	0	3.30000	3.60000	.	.	.	.	.	.
4,4'-DDD	2	14	0	0.01000	0.10000	.	.	0.01	0.0100	1.000	10.000
	13	4	0	0.01000	0.01000	.	.	0.01	0.0100	1.000	1.000
	15	4	0	0.05000	0.20000	.	.	0.01	0.0100	5.000	20.000
4,4'-DDE	1	16	0	0.01000	0.20000	0.00050	0.0063	0.01	0.0063	1.600	32.000
	3	34	0	0.01000	0.01708	0.00050	0.0063	0.01	0.0063	1.600	2.733
	10	8	0	0.01000	0.01000	0.00050	0.0063	0.01	0.0063	1.600	1.600
4,4'-DDT	13	3	0	0.01000	0.01000	0.00050	0.0063	0.01	0.0063	1.600	1.600
	15	4	0	0.05000	0.20000	0.00050	0.0063	0.01	0.0063	8.000	32.000
4,6-DINITRO-2-METHYLPHENOL	1	6	0	3.40000	4.30000	.	.	.	.	.	.
	2	8	0	3.50000	9.30000	.	.	.	.	.	.
	3	34	0	0.93000	2.10000	.	.	.	.	.	.
4-BROMOPHENYL-PHENYLETHER	4	13	0	3.30000	4.10000	.	.	.	.	.	.
	5	8	0	3.30000	4.20000	.	.	.	.	.	.
	13	2	0	3.40000	3.50000	.	.	.	.	.	.
4-CHLORO-3-METHYLPHENOL	15	4	0	3.40000	3.80000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
4-CHLORO-3-METHYLPHENOL	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	15	0	0.66000	0.83000	.	.	.	.	.	.
	5	9	0	0.66000	0.83000	.	.	.	.	.	.
4-CHLOROANILINE	13	2	0	0.69000	0.70000	.	.	.	.	.	.
	15	4	0	0.67000	0.77000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	0.00400	0.0500	.	0.0500	13.600	17.400
4-CHLOROANILINE	2	2	0	0.70000	1.80000	0.00400	0.0500	.	0.0500	14.000	36.000
	3	34	0	0.19000	0.41000	0.00400	0.0500	.	0.0500	3.800	8.200
	4	6	0	0.66000	0.72000	0.00400	0.0500	.	0.0500	13.200	14.400
4-CHLOROANILINE	15	3	0	0.67000	0.71000	0.00400	0.0500	.	0.0500	13.400	14.200

Analyte	Reach	Number analyzed in reach (fish)	Number defects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
4-CHLOROPHENYL-PHENYLETHER	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
4-METHYLPHENOL	1	6	0	0.68000	0.87000	0.05000	0.6250	.	0.6250	1.088	1.392
	2	8	0	0.70000	1.90000	0.05000	0.6250	.	0.6250	1.120	3.040
	3	34	0	0.19000	0.41000	0.05000	0.6250	.	0.6250	0.304	0.656
	4	14	0	0.66000	0.83000	0.05000	0.6250	.	0.6250	1.056	1.328
	5	9	0	0.66000	0.83000	0.05000	0.6250	.	0.6250	1.056	1.328
	13	2	0	0.69000	0.70000	0.05000	0.6250	.	0.6250	1.104	1.120
	15	4	0	0.67000	0.77000	0.05000	0.6250	.	0.6250	1.072	1.232
4-NITROANILINE	1	6	0	3.40000	4.30000	0.00131	0.0164	.	0.0164	207.634	262.595
	2	8	0	3.50000	9.30000	0.00131	0.0164	.	0.0164	213.740	567.939
	3	34	0	0.93000	2.10000	0.00131	0.0164	.	0.0164	56.794	128.244
	4	2	0	3.30000	3.60000	0.00131	0.0164	.	0.0164	201.527	219.847
4-NITROPHENOL	1	6	0	3.40000	4.30000	.	.	.	.	.	.
	2	8	0	3.50000	9.30000	.	.	.	.	.	.
	3	34	0	0.93000	2.10000	.	.	.	.	.	.
	4	13	0	3.30000	4.10000	.	.	.	.	.	.
	5	9	0	3.30000	4.20000	.	.	.	.	.	.
ACENAPHTHENE	13	2	0	3.40000	3.50000	.	.	.	.	.	.
	15	4	0	3.40000	3.80000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	0.06000	0.7500	.	0.7500	0.907	1.160
	2	8	0	0.70000	1.90000	0.06000	0.7500	.	0.7500	0.933	2.533
	3	34	0	0.19000	0.41000	0.06000	0.7500	.	0.7500	0.253	0.547
ACENAPHTHYLENE	4	6	0	0.66000	0.72000	0.06000	0.7500	.	0.7500	0.880	0.960
	15	3	0	0.67000	0.71000	0.06000	0.7500	.	0.7500	0.893	0.947
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
ALDRIN	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	16	0	0.01000	0.20000	0.00003	0.0004	.	0.0004	26.667	533.333
	3	34	0	0.00772	0.00854	0.00003	0.0004	.	0.0004	20.587	22.773
	10	8	0	0.01000	0.01000	0.00003	0.0004	.	0.0004	26.667	26.667
15	4	0	0.05000	0.20000	0.00003	0.0004	.	0.0004	133.333	533.333	

Table F.12 (continued)

Analyte	Reech	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
ANTHRACENE	1	6	0	0.68000	0.87000	0.30000	3.7500	.	3.7500	0.181	0.232
	2	8	0	0.70000	1.90000	0.30000	3.7500	.	3.7500	0.187	0.507
	3	34	0	0.19000	0.41000	0.30000	3.7500	.	3.7500	0.051	0.109
	4	6	0	0.66000	0.72000	0.30000	3.7500	.	3.7500	0.176	0.192
	15	3	0	0.67000	0.71000	0.30000	3.7500	.	3.7500	0.179	0.189
ANTHRONY	1	28	0	0.28000	0.58000	0.00040	0.0050	0.30	0.0050	56.000	116.000
	2	13	0	0.46000	0.60000	0.00040	0.0050	0.30	0.0050	92.000	120.000
	3	35	0	0.30000	0.59000	0.00040	0.0050	0.30	0.0050	60.000	118.000
	4	27	0	0.48000	0.61000	0.00040	0.0050	0.30	0.0050	96.000	122.000
	5	25	0	0.46000	0.63000	0.00040	0.0050	0.30	0.0050	92.000	126.000
	10	15	0	0.35000	0.61000	0.00040	0.0050	0.30	0.0050	70.000	122.000
	13	10	0	0.46000	0.57000	0.00040	0.0050	0.30	0.0050	92.000	114.000
	15	16	0	0.35000	0.57000	0.00040	0.0050	0.30	0.0050	70.000	114.000
	10	8	0	0.01000	0.05000	.	.	0.04	0.0400	0.250	1.250
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
BENZO(A)ANTHRACENE	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
BENZO(A)PYRENE	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	33	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
BENZO(B)FLUORANTHENE	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	33	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
BENZO(G,H,I)PERYLENE	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	32	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.

Table F-12 (continued)

Analyte	Reech	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
BENZ(a)FLUORANTHENE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	33	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
BENZOIC ACID	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	3.40000	4.30000	4.00000	50.0000	.	50.0000	0.068	0.086
	2	8	0	3.50000	9.30000	4.00000	50.0000	.	50.0000	0.070	0.186
	3	34	0	0.93000	2.10000	4.00000	50.0000	.	50.0000	0.019	0.042
BENZYL ALCOHOL	4	2	0	3.30000	3.60000	4.00000	50.0000	.	50.0000	0.066	0.072
	1	6	0	0.68000	0.87000	0.09000	1.1250	.	1.1250	0.604	0.773
	2	8	0	0.70000	1.90000	0.09000	1.1250	.	1.1250	0.622	1.689
	3	34	0	0.19000	0.41000	0.09000	1.1250	.	1.1250	0.169	0.364
BERYLLIUM	4	6	0	0.66000	0.72000	0.09000	1.1250	.	1.1250	0.587	0.640
	15	3	0	0.67000	0.71000	0.09000	1.1250	.	1.1250	0.596	0.631
	2	13	0	0.00300	0.00300	0.00500	0.0625	25.00	0.0625	0.048	0.048
	4	27	0	0.00300	0.00300	0.00500	0.0625	25.00	0.0625	0.048	0.048
BIS(2-CHLOROETHOXY)METHANE	13	3	0	0.00300	0.00300	0.00500	0.0625	25.00	0.0625	0.048	0.048
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
BIS(2-CHLOROETHYL)ETHER	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
BIS(2-CHLOROISOPROPYL)ETHER	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
BIS(2-ETHYLHEXYL)PHTHALATE	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
BIS(2-ETHYLHEXYL)PHTHALATE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.

Table F.12 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
BUTYLBENZYLPHTHALATE	1	6	0	0.68000	0.87000	0.20000	2.5000	.	2.5000	0.272	0.348
	2	8	0	0.70000	1.90000	0.20000	2.5000	.	2.5000	0.280	0.760
	3	34	0	0.19000	0.41000	0.20000	2.5000	.	2.5000	0.076	0.164
	4	6	0	0.66000	0.72000	0.20000	2.5000	.	2.5000	0.264	0.288
	15	3	0	0.67000	0.71000	0.20000	2.5000	.	2.5000	0.268	0.284
CALCIUM	1	28	0	0.11000	0.23000	0.00050	0.0063	0.25	0.0063	17.600	36.800
	2	13	0	0.18000	0.24000	0.00050	0.0063	0.25	0.0063	28.800	38.400
	3	35	0	0.12000	0.24000	0.00050	0.0063	0.25	0.0063	19.200	38.400
	4	27	0	0.19000	0.24000	0.00050	0.0063	0.25	0.0063	30.400	38.400
	5	25	0	0.19000	0.28000	0.00050	0.0063	0.25	0.0063	30.400	44.800
	10	15	0	0.14000	0.25000	0.00050	0.0063	0.25	0.0063	22.400	40.000
	13	10	0	0.18000	0.24000	0.00050	0.0063	0.25	0.0063	28.800	38.400
	15	16	0	0.14000	0.23000	0.00050	0.0063	0.25	0.0063	22.400	36.800
	1	28	0	0.28000	0.58000	.	.	1.00	1.0000	0.280	0.580
	2	13	0	0.46000	0.60000	.	.	1.00	1.0000	0.460	0.600
CHROMIUM	3	35	0	0.30000	0.59000	.	.	1.00	1.0000	0.300	0.590
	4	27	0	0.48000	0.61000	.	.	1.00	1.0000	0.480	0.610
	5	25	0	0.46000	0.63000	.	.	1.00	1.0000	0.460	0.630
	10	15	0	0.35000	0.61000	.	.	1.00	1.0000	0.350	0.610
	13	10	0	0.45000	0.59000	.	.	1.00	1.0000	0.450	0.590
	15	16	0	0.35000	0.57000	.	.	1.00	1.0000	0.350	0.570
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
CHRYSENE	15	3	0	0.67000	0.71000	.	.	.	.	.	.
	1	28	0	0.28000	0.58000	1.30000	16.2500	50.00	16.2500	0.017	0.036
	2	13	0	0.46000	0.60000	1.30000	16.2500	50.00	16.2500	0.028	0.037
	3	35	0	0.30000	0.59000	1.30000	16.2500	50.00	16.2500	0.018	0.036
	4	27	0	0.48000	0.61000	1.30000	16.2500	50.00	16.2500	0.030	0.038
	5	25	0	0.46000	0.63000	1.30000	16.2500	50.00	16.2500	0.022	0.038
	10	15	0	0.35000	0.61000	1.30000	16.2500	50.00	16.2500	0.028	0.036
	13	10	0	0.45000	0.59000	1.30000	16.2500	50.00	16.2500	0.022	0.035
	15	16	0	0.35000	0.57000	1.30000	16.2500	50.00	16.2500	0.022	0.035
	1	6	0	0.68000	0.87000	0.10000	1.2500	.	1.2500	0.544	0.696
COPPER	2	8	0	0.70000	1.90000	0.10000	1.2500	.	1.2500	0.560	1.520
	3	34	0	0.19000	0.41000	0.10000	1.2500	.	1.2500	0.152	0.328
	4	6	0	0.66000	0.72000	0.10000	1.2500	.	1.2500	0.528	0.576
	6	3	0	0.66000	0.72000	0.10000	1.2500	.	1.2500	0.536	0.568
	15	3	0	0.67000	0.71000	0.10000	1.2500	.	1.2500	0.536	0.568
D1-N-BUTYLPHTHALATE	1	6	0	0.68000	0.87000	0.10000	1.2500	.	1.2500	0.544	0.696
	2	8	0	0.70000	1.90000	0.10000	1.2500	.	1.2500	0.560	1.520
	3	34	0	0.19000	0.41000	0.10000	1.2500	.	1.2500	0.152	0.328
	4	6	0	0.66000	0.72000	0.10000	1.2500	.	1.2500	0.528	0.576
	15	3	0	0.67000	0.71000	0.10000	1.2500	.	1.2500	0.536	0.568

Table F.12 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
HEXACHLOROBENZENE											
	1	6	0	0.68000	0.87000	0.00080	0.0100	.	0.0100	68.000	87.000
	2	8	0	0.70000	1.90000	0.00080	0.0100	.	0.0100	70.000	190.000
	3	34	0	0.19000	0.41000	0.00080	0.0100	.	0.0100	19.000	41.000
	4	6	0	0.66000	0.72000	0.00080	0.0100	.	0.0100	66.000	72.000
	15	3	0	0.67000	0.71000	0.00080	0.0100	.	0.0100	67.000	71.000
HEXACHLOROBUTADIENE											
	1	6	0	0.68000	0.87000	0.00200	0.0250	.	0.0250	27.200	34.800
	2	8	0	0.70000	1.90000	0.00200	0.0250	.	0.0250	28.000	76.000
	3	34	0	0.19000	0.41000	0.00200	0.0250	.	0.0250	7.600	16.400
	4	6	0	0.66000	0.72000	0.00200	0.0250	.	0.0250	26.400	28.800
	15	3	0	0.67000	0.71000	0.00200	0.0250	.	0.0250	26.800	28.400
HEXACHLOROCYCLOPENTADIENE											
	1	6	0	0.68000	0.87000	0.00700	0.0875	.	0.0875	7.771	9.943
	2	8	0	0.70000	1.90000	0.00700	0.0875	.	0.0875	8.000	21.714
	3	34	0	0.19000	0.41000	0.00700	0.0875	.	0.0875	2.171	4.686
	4	6	0	0.66000	0.72000	0.00700	0.0875	.	0.0875	7.543	8.229
	15	3	0	0.67000	0.71000	0.00700	0.0875	.	0.0875	7.657	8.114
HEXACHLOROETHANE											
	1	6	0	0.68000	0.87000	0.00100	0.0125	.	0.0125	54.400	69.600
	2	8	0	0.70000	1.90000	0.00100	0.0125	.	0.0125	56.000	152.000
	3	34	0	0.19000	0.41000	0.00100	0.0125	.	0.0125	15.200	32.800
	4	6	0	0.66000	0.72000	0.00100	0.0125	.	0.0125	52.800	57.600
	15	3	0	0.67000	0.71000	0.00100	0.0125	.	0.0125	53.600	56.800
INDENO(1,2,3-CD)PYRENE											
	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	33	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
ISOPHORONE											
	1	6	0	0.68000	0.87000	0.20000	2.5000	.	2.5000	0.272	0.348
	2	8	0	0.70000	1.90000	0.20000	2.5000	.	2.5000	0.280	0.760
	3	34	0	0.19000	0.41000	0.20000	2.5000	.	2.5000	0.076	0.164
	4	6	0	0.66000	0.72000	0.20000	2.5000	.	2.5000	0.264	0.288
	15	3	0	0.67000	0.71000	0.20000	2.5000	.	2.5000	0.268	0.284
LEAD											
	1	28	0	0.28000	0.58000	.	.	15.00	15.0000	0.019	0.039
	2	13	0	0.46000	0.60000	.	.	15.00	15.0000	0.031	0.040
	3	25	0	0.30000	0.59000	.	.	15.00	15.0000	0.020	0.039
	4	17	0	0.48000	0.61000	.	.	15.00	15.0000	0.032	0.041
	5	5	0	0.46000	0.51000	.	.	15.00	15.0000	0.031	0.034
	10	5	0	0.35000	0.44000	.	.	15.00	15.0000	0.023	0.029
	13	10	0	0.45000	0.59000	.	.	15.00	15.0000	0.030	0.039
	15	16	0	0.35000	0.57000	.	.	15.00	15.0000	0.023	0.038

Table F.12 (continued)

Analyte	Resch	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ¹ detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
N-NITROSO-DI-N-PROPYLAMINE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
N-NITROSDIPHENYLAMINE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
NAPHTHALENE	1	6	0	0.68000	0.87000	0.00400	0.0500	.	0.0500	13.600	17.400
	2	8	0	0.70000	1.90000	0.00400	0.0500	.	0.0500	14.000	38.000
	3	34	0	0.19000	0.41000	0.00400	0.0500	.	0.0500	3.800	8.200
	4	6	0	0.66000	0.72000	0.00400	0.0500	.	0.0500	13.200	14.400
	5	1	0	0.05700	0.05700	0.00400	0.0500	.	0.0500	1.140	1.140
NICKEL	15	3	0	0.67000	0.71000	0.00400	0.0500	.	0.0500	13.400	14.200
	1	28	0	0.28000	1.10000	0.02000	0.2500	50.00	0.2500	1.120	4.400
	2	13	0	0.46000	0.60000	0.02000	0.2500	50.00	0.2500	1.840	2.400
	3	35	0	0.30000	0.59000	0.02000	0.2500	50.00	0.2500	1.200	2.360
	4	27	0	0.48000	0.61000	0.02000	0.2500	50.00	0.2500	1.920	2.440
NITROBENZENE	5	5	0	0.46000	0.51000	0.02000	0.2500	50.00	0.2500	1.840	2.040
	10	15	0	0.35000	0.61000	0.02000	0.2500	50.00	0.2500	1.400	2.440
	13	10	0	0.45000	0.59000	0.02000	0.2500	50.00	0.2500	1.800	2.360
	15	16	0	0.35000	0.57000	0.02000	0.2500	50.00	0.2500	1.400	2.280
	1	6	0	0.68000	0.87000	0.00050	0.0063	.	0.0063	108.800	139.200
PENTACHLOROPHENOL	2	8	0	0.70000	1.90000	0.00050	0.0063	.	0.0063	112.000	304.000
	3	34	0	0.19000	0.41000	0.00050	0.0063	.	0.0063	30.400	65.600
	4	6	0	0.66000	0.72000	0.00050	0.0063	.	0.0063	105.600	115.200
	15	3	0	0.67000	0.71000	0.00050	0.0063	.	0.0063	107.200	113.600
	1	6	0	3.40000	4.30000	0.03000	0.3750	.	0.3750	9.067	11.467
	2	8	0	3.50000	9.30000	0.03000	0.3750	.	0.3750	9.333	24.800
	3	34	0	0.93000	2.10000	0.03000	0.3750	.	0.3750	2.480	5.600
	4	13	0	3.30000	4.10000	0.03000	0.3750	.	0.3750	8.800	10.933
	5	9	0	3.30000	4.20000	0.03000	0.3750	.	0.3750	8.800	11.200
	13	2	0	3.40000	3.50000	0.03000	0.3750	.	0.3750	9.067	9.333
	15	4	0	3.40000	3.80000	0.03000	0.3750	.	0.3750	9.067	10.133

Table F.12 (continued)

Analyte	Reach	Number analyzed in reach (fish)	Number detects in reach (fish)	Minimum detection limit	UCL ^a detection limit	Reference dose	RFD dietary concentration	Lower piscivore criterion	Minimum criterion	Minimum detection limit quo.	UCL detection limit quo.
PHENANTHRENE	1	6	0	0.68000	0.87000	.	.	.	.	.	.
	2	8	0	0.70000	1.90000	.	.	.	.	.	.
	3	34	0	0.19000	0.41000	.	.	.	.	.	.
	4	6	0	0.66000	0.72000	.	.	.	.	.	.
	15	3	0	0.67000	0.71000	.	.	.	.	.	.
PHENOL	1	6	0	0.68000	0.87000	0.60000	7.5000	.	7.5000	0.091	0.116
	2	8	0	0.70000	1.90000	0.60000	7.5000	.	7.5000	0.093	0.253
	3	34	0	0.19000	0.41000	0.60000	7.5000	.	7.5000	0.025	0.055
	4	13	0	0.66000	0.83000	0.60000	7.5000	.	7.5000	0.088	0.111
	5	9	0	0.66000	0.83000	0.60000	7.5000	.	7.5000	0.088	0.111
	13	2	0	0.69000	0.70000	0.60000	7.5000	.	7.5000	0.092	0.093
	15	4	0	0.67000	0.77000	0.60000	7.5000	.	7.5000	0.089	0.103
PYRENE	1	6	0	0.68000	0.87000	0.03000	0.3750	.	0.3750	1.813	2.320
	2	8	0	0.70000	1.90000	0.03000	0.3750	.	0.3750	1.867	5.067
	3	34	0	0.19000	0.41000	0.03000	0.3750	.	0.3750	0.507	1.093
	4	6	0	0.66000	0.72000	0.03000	0.3750	.	0.3750	1.760	1.920
	15	3	0	0.67000	0.71000	0.03000	0.3750	.	0.3750	1.787	1.893
SELENIUM	5	25	0	0.46000	0.63000	0.00300	0.0375	0.50	0.0375	12.267	16.800
	13	10	0	0.45000	0.59000	0.00300	0.0375	0.50	0.0375	12.000	15.733
SILVER	1	28	0	0.11000	0.23000	0.00300	0.0375	10.00	0.0375	2.933	6.133
	2	13	0	0.18000	0.24000	0.00300	0.0375	10.00	0.0375	4.800	6.400
	3	35	0	0.12000	0.24000	0.00300	0.0375	10.00	0.0375	3.200	6.400
	4	27	0	0.19000	0.24000	0.00300	0.0375	10.00	0.0375	5.067	6.400
	5	25	0	0.19000	0.25000	0.00300	0.0375	10.00	0.0375	5.067	6.667
	10	15	0	0.14000	0.25000	0.00300	0.0375	10.00	0.0375	3.733	6.667
	13	10	0	0.18000	0.24000	0.00300	0.0375	10.00	0.0375	4.800	6.400
	15	16	0	0.14000	0.23000	0.00300	0.0375	10.00	0.0375	3.733	6.133
THALLIUM	1	28	0	0.02000	0.02000	.	.	.	.	.	.
	2	13	0	0.02000	0.02000	.	.	.	.	.	.
	3	35	0	0.02000	0.02000	.	.	.	.	.	.
	4	27	0	0.02000	0.02000	.	.	.	.	.	.
	5	16	0	0.02000	0.02000	.	.	.	.	.	.
	10	15	0	0.02000	0.02000	.	.	.	.	.	.
	13	10	0	0.02000	0.02000	.	.	.	.	.	.
	15	16	0	0.02000	0.02000	.	.	.	.	.	.

^aThe maximum detection limit is used in cases where the UCL detection limit was missing.

Appendix G
BRADEN BRANCH

CONTENTS OF APPENDIX G

G.1	STUDY AREA	G-3
G.2	HISTORICAL BACKGROUND	G-3
G.3	RECENT DATA	G-5
G.4	REFERENCE CONCENTRATIONS	G-5
G.5	DATA SUMMARY	G-5
G.6	SUMMARY AND INTERPRETATION	G-12
G.7	HUMAN HEALTH RISK	G-12
G.8	REFERENCES FOR APPENDIX G	G-16

TABLES

G.1	Data listing for CR-ERP Braden Branch sediment grab sample radiological analyses	G-6
G.2	Summary statistics for Braden Branch sediment core data	G-7
G.3	Data listing for CR-ERP Braden Branch sediment core samples radiological analyses	G-8
G.4	Summary statistics for Braden Branch sediment grab data	G-9
G.5	Results of adult nonconservative screening for carcinogens in surface sediment of Braden Branch where at least one measurement was above detection limits	G-13
G.6	Results of adult conservative screening for carcinogens in surface sediment of Braden Branch where at least one measurement was above detection limits	G-14

FIGURES

G.1	Braden Branch near-shore sample sites	G-4
G.2	Cobalt-60 concentrations at Braden Branch	G-10
G.3	Cobalt concentrations in Braden Branch core (no. 18700G)	G-11

BRADEN BRANCH SCREENING-LEVEL RISK ASSESSMENT

Braden Branch is a known source of ^{60}Co contamination to Melton Hill Reservoir at Clinch River Mile (CRM) 50.5. The source of this contamination was the site of the former American Nuclear Corporation (ANC), which released ^{60}Co into the watershed of Braden Branch. Samples were collected in Braden Branch during Phase 1 of the Clinch River Remedial Investigation to characterize this site. The purpose of this appendix is to summarize the data from sampling of Braden Branch and to present a screening-level assessment of the risk to human health associated with exposure to the contaminated areas.

G.1 STUDY AREA

Braden Branch Creek is a small stream that enters Melton Hill Reservoir at approximately CRM 50.5 (Fig. G.1). This is several hundred meters upstream and on the opposite shoreline from the Oak Ridge Marina. The creek is separated from Melton Hill Reservoir by a railroad berm. Surface water is exchanged between the two water bodies via a culvert that passes through the berm. Damming caused by the berm creates an embayment on the upstream side, which acts as a settling pond for particles washed from the Braden Branch watershed.

G.2 HISTORICAL BACKGROUND

In 1968, personnel from ORNL noted abnormally high radioactivity in a water sample from the Braden Branch area of the Clinch River. The source of this radioactivity was traced to the former ANC located on Braden Branch Creek. Under the jurisdiction of the U.S. Atomic Energy Commission (until 1965) and the State of Tennessee (between 1965 and 1970 when ANC closed), ANC used ^{60}Co as a radiation source for medical instruments between 1962 and 1970.

After the contamination was detected in the Clinch River, radiation surveys near the site revealed exposure rates in the mR/hr to R/hr range. The ANC facility was closed in 1970, and clean-up efforts commenced. The clean up was restricted to the immediate area around the plant. No effort was made to clean up contamination in the Braden Branch embayment or areas downstream. The Division of Radiological Health of the Tennessee Department of Environment and Conservation (TDEC) obtains results of monthly radiation surveys in the area (approximately 50 acres) and from annual measurements near the building site. Public access to the ANC building and surrounding land is restricted.

Sediment samples were collected from the Braden Branch embayment in 1977 and 1987. In 1977, ^{60}Co concentrations in surface sediment within the Braden Branch embayment were as high as 1287 pCi/g dry wt. The highest concentration found in surface sediment in the embayment during a 1987 survey was 152.0 pCi/g. Explanations for this drop in concentration are (1) radioactive decay; the half-life of ^{60}Co is approximately five years, which means that 1287 pCi/g dry wt in 1977 would have been reduced to 320 pCi/g by 1987, and (2) the redistribution and dilution of contaminated sediment could be responsible for spreading and mixing contaminated sediment within the embayment and into Melton Hill Reservoir. The observation that the highest ^{60}Co concentrations in 1977 were in the upper end of the

Fig. G.1. Braden Branch near-shore sample sites.

embayment but in 1987 were in the lower end suggests that there is some redistribution of ^{60}Co . Finally, along with redistribution, contaminated sediments may be buried by "cleaner" sediments. There is some historic evidence of this as well. For example, in the 1987 survey of the embayment, a ^{60}Co concentration in the top 6 in. of sediment at one site was 39.9 pCi/g dry wt, while the concentration in the 6–12 in. segment was 383.0 pCi/g dry wt. However, this pattern is by no means consistent throughout the embayment.

G.3 RECENT DATA

Sediment sampling was conducted in the immediate vicinity of Braden Branch as Phase 1 of the CRRI. A total of 20 sediment grab samples were taken between July 25 and August 10, 1990 (Table G.1). These samples were taken with a Ponar grab sampler for the purpose of characterizing the radionuclide concentrations in the surface sediment only. A single sediment core was also taken within 300 m of the confluence of Braden Branch and Melton Hill Reservoir (Table G.2). This core was divided into separate samples by depth intervals (2 cm to a depth of 48 cm) and analyzed to provide a vertical profile of sediment contamination (Table G.3). Sample locations for grab samples (ID nos. 109–125, and 188–190) and for the core sample (ID no. 187) are shown on Fig. G.1.

G.4 REFERENCE CONCENTRATIONS

Radionuclides are present throughout the world due to natural occurrence and to man-made sources. Sediment samples taken from Norris Reservoir provide reference or "background" concentrations for radionuclides and other contaminants being examined in the CRRI. Norris Reservoir background levels of ^{60}Co , which originated from atmospheric testing of nuclear weapons and not from DOE facilities in Oak Ridge, are below minimum detectable activity.

G.5 DATA SUMMARY

Table G.4 summarizes ^{60}Co surface sediment concentrations from the 20 grab samples and the top 2 cm of the core sample collected in 1990. Fig. G.2 shows the ^{60}Co concentration at each of the sample locations. Locations with the highest concentrations (>1.01 pCi/g) are, with the exception of sample no. 190 (^{60}Co concentration = 4.53 pCi/g), within 200 m of the culvert under the railroad bridge where Braden Branch enters Melton Hill Reservoir.

Figure G.3 shows ^{60}Co concentration by depth in core no. 18700G and compares these concentrations with those from a core taken from Melton Hill Reservoir at CRM 23.9 (core no. 8200G). The profile for the Braden Branch core indicates that there are two concentration peaks. The first is at a depth of 22 cm with a concentration of 20.07 pCi/g. The second peak is at 33 cm with an associated concentration of 39.33 pCi/g. The concentrations from the two downstream cores (near Solway Bridge and Melton Hill Dam) are all less than 1.0 pCi/g dry wt, and most of the concentrations in these cores are below the detectable limit. The concentrations of ^{60}Co in the Braden Branch core are considerably higher than the minimum detectable levels (0–0.3 pCi/g) to a depth of 45 cm.

Table G.1. Data Listing For CR-ERP Braden Branch Sediment Grab Sample Radiological Analyses

RIVER SYSTEM POSITION	RIVER MILE/KM	SAMPLE MATRIX	CR-ERP SAMPLE ID	ANALYSIS	RESULT	RAD UNCERTAINTY (1 S.D.)	UNITS	DATA QUALIFIER
1	CRM 50.1	GRAB	124008	CO-60	0.0434	0.0167	PCI/G (DRY)	J
1	CRM 50.1	GRAB	124008	CS-137	0.1512	0.0114	PCI/G (DRY)	V/V
1	CRM 50.1	GRAB	125008	CO-60	0.0242	0.0092	PCI/G (DRY)	J
1	CRM 50.1	GRAB	125008	CS-137	0.0798	0.0117	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	109008	CO-60	7.4079	0.0596	PCI/G (DRY)	J
1	CRM 50.5	GRAB	109008	CS-137	1.4640	0.0292	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	110008	CO-60	9.0599	0.0564	PCI/G (DRY)	J
1	CRM 50.5	GRAB	110008	CS-137	0.9334	0.0249	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	111008	CO-60	6.8871	0.0592	PCI/G (DRY)	J
1	CRM 50.5	GRAB	111008	CS-137	1.0347	0.0254	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	112008	CO-60	2.3061	0.0146	PCI/G (DRY)	J
1	CRM 50.5	GRAB	112008	CS-137	0.3763	0.0063	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	113008	CO-60	1.0990	0.0202	PCI/G (DRY)	J
1	CRM 50.5	GRAB	113008	CS-137	0.2218	0.0094	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	114008	CO-60	6.8564	0.0367	PCI/G (DRY)	J
1	CRM 50.5	GRAB	114008	CS-137	1.9501	0.0188	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	115008	CO-60	8.2378	0.0336	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	115008	CS-137	1.2485	0.0145	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	116008	CO-60	3.5694	0.0380	PCI/G (DRY)	J
1	CRM 50.5	GRAB	116008	CS-137	0.4006	0.0147	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	117008	CO-60	0.1940	0.0094	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	117008	CS-137	1.4336	0.0349	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	118008	CO-60	0.5020	0.0165	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	118008	CS-137	1.0119	0.0221	PCI/G (DRY)	J
1	CRM 50.5	GRAB	119008	CO-60	0.2024	0.0112	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	119008	CS-137	0.0493	0.0087	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	120008	CO-60	0.3352	0.0176	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	120008	CS-137	0.1221	0.0099	PCI/G (DRY)	J
1	CRM 50.5	GRAB	122008	CO-60	0.0554	0.0056	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	122008	CS-137	0.2999	0.0187	PCI/G (DRY)	J
1	CRM 50.5	GRAB	123008	CO-60	0.0294	0.0128	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	123008	CS-137	0.1727	0.0061	PCI/G (DRY)	J
1	CRM 50.5	GRAB	188008	CO-60	0.0383	0.0051	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	189008	CO-60	0.1455	0.0252	PCI/G (DRY)	J
1	CRM 50.5	GRAB	189008	CS-137	4.5291	0.0131	PCI/G (DRY)	V/V
1	CRM 50.5	GRAB	190008	CO-60	0.7173	0.0653	PCI/G (DRY)	J
1	CRM 50.5	GRAB	190008	CS-137		0.0287	PCI/G (DRY)	V/V

Includes only routine samples and detected values

Table 6.2. Summary Statistics For Braden Branch Sediment Core Data
Summarized for All Reaches

TYPE OF ANALYSIS	ANALYSIS	OBS.	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	SAMPLE UNITS
RAD	CO-60	24	0.088889	UJ	39.3257	V/V	10.3075	PCI/G (DRY)
	CS-137	24	0.037668	V/V	1.6667	V/V	0.9500	PCI/G (DRY)

Table G.3. Data Listing For CR-ERP Braden Branch Sediment Core Sample Radiological Analyses

RIVER SYSTEM POSITION	RIVER MILE/KM	CR-ERP SAMPLE ID	DEPTH INCREMENT (CM)	ANALYSIS	RESULT	RAD UNCERTAINTY (1 S.D.)	UNITS	DATA QUALIFIER
1	CRM 50.5	187016	000-002	CO-60	6.2766	0.1383	PCI/G (DRY)	V/V
1	CRM 50.5	187016	000-002	CS-137	1.6667	0.0638	PCI/G (DRY)	V/V
1	CRM 50.5	187026	002-004	CO-60	5.1992	0.0922	PCI/G (DRY)	V/V
1	CRM 50.5	187026	002-004	CS-137	1.3208	0.0503	PCI/G (DRY)	V/V
1	CRM 50.5	187036	004-006	CO-60	4.9715	0.2657	PCI/G (DRY)	V/V
1	CRM 50.5	187036	004-006	CS-137	1.4611	0.1632	PCI/G (DRY)	V/V
1	CRM 50.5	187046	006-008	CO-60	5.3959	0.2762	PCI/G (DRY)	V/V
1	CRM 50.5	187046	006-008	CS-137	1.4549	0.1492	PCI/G (DRY)	V/V
1	CRM 50.5	187056	008-010	CO-60	5.8007	0.2669	PCI/G (DRY)	V/V
1	CRM 50.5	187056	008-010	CS-137	1.4235	0.1317	PCI/G (DRY)	V/V
1	CRM 50.5	187066	010-012	CO-60	5.9770	0.2627	PCI/G (DRY)	V/V
1	CRM 50.5	187066	010-012	CS-137	1.5107	0.1412	PCI/G (DRY)	V/V
1	CRM 50.5	187076	012-014	CO-60	7.0478	0.2605	PCI/G (DRY)	V/V
1	CRM 50.5	187076	012-014	CS-137	1.3025	0.1143	PCI/G (DRY)	V/V
1	CRM 50.5	187086	014-016	CO-60	9.0371	0.2889	PCI/G (DRY)	V/V
1	CRM 50.5	187086	014-016	CS-137	1.1554	0.1210	PCI/G (DRY)	V/V
1	CRM 50.5	187096	016-018	CO-60	13.9271	0.3647	PCI/G (DRY)	V/V
1	CRM 50.5	187096	016-018	CS-137	1.0659	0.1403	PCI/G (DRY)	V/V
1	CRM 50.5	187106	018-020	CO-60	19.3943	0.3995	PCI/G (DRY)	V/V
1	CRM 50.5	187106	018-020	CS-137	0.8505	0.1418	PCI/G (DRY)	V/V
1	CRM 50.5	187116	020-022	CO-60	20.0659	0.4084	PCI/G (DRY)	V/V
1	CRM 50.5	187116	020-022	CS-137	0.8959	0.1581	PCI/G (DRY)	V/V
1	CRM 50.5	187126	022-024	CO-60	9.7592	0.3371	PCI/G (DRY)	V/V
1	CRM 50.5	187126	022-024	CS-137	1.4125	0.1493	PCI/G (DRY)	V/V
1	CRM 50.5	187136	024-026	CO-60	6.8679	0.2773	PCI/G (DRY)	V/V
1	CRM 50.5	187136	024-026	CS-137	1.2887	0.1387	PCI/G (DRY)	V/V
1	CRM 50.5	187146	026-028	CO-60	8.6687	0.2941	PCI/G (DRY)	V/V
1	CRM 50.5	187146	026-028	CS-137	1.5325	0.1502	PCI/G (DRY)	V/V
1	CRM 50.5	187156	028-030	CO-60	13.1120	0.3320	PCI/G (DRY)	V/V
1	CRM 50.5	187156	028-030	CS-137	0.9544	0.1383	PCI/G (DRY)	V/V
1	CRM 50.5	187166	030-032	CO-60	21.1804	0.4244	PCI/G (DRY)	V/V
1	CRM 50.5	187166	030-032	CS-137	1.2865	0.1326	PCI/G (DRY)	V/V
1	CRM 50.5	187176	032-034	CO-60	39.3257	0.5852	PCI/G (DRY)	V/V
1	CRM 50.5	187176	032-034	CS-137	1.0687	0.1781	PCI/G (DRY)	V/V
1	CRM 50.5	187186	034-036	CO-60	27.2194	0.3912	PCI/G (DRY)	V/V
1	CRM 50.5	187186	034-036	CS-137	0.5357	0.1020	PCI/G (DRY)	V/V
1	CRM 50.5	187196	036-038	CO-60	8.4264	0.2171	PCI/G (DRY)	V/V
1	CRM 50.5	187196	036-038	CS-137	0.1783	0.0620	PCI/G (DRY)	V/V
1	CRM 50.5	187206	038-040	CO-60	6.2693	0.1973	PCI/G (DRY)	V/V
1	CRM 50.5	187216	040-042	CO-60	2.4843	0.1525	PCI/G (DRY)	V/V
1	CRM 50.5	187216	040-042	CS-137	0.0377	0.0422	PCI/G (DRY)	V/V
1	CRM 50.5	187226	042-044	CO-60	0.6897	0.0821	PCI/G (DRY)	V/V
1	CRM 50.5	187236	044-046	CO-60	0.1954	0.0570	PCI/G (DRY)	V/V

Includes only routine samples and detected values

Table G.4. Summary Statistics For Braden Branch Sediment Grab Data
Summarized for All Reaches

TYPE OF ANALYSIS	ANALYSIS	OBS.	MINIMUM VALUE	MIN QUAL.	MAXIMUM VALUE	MAX QUAL.	AVERAGE VALUE	SAMPLE UNITS
RAD	CO-60	20	0.024233	J	9.05986	J	2.73849	PCI/G (DRY)
	CS-137	20	0.049297	V/V	1.95006	V/V	0.52404	PCI/G (DRY)

Fig. G.2. Cobalt-60 concentrations at Braden Branch.

Fig. G.3. Cobalt concentrations in Braden Branch core (no. 18700G).

G.6 SUMMARY AND INTERPRETATION

The ^{60}Co sediment concentrations in the Melton Hill Reservoir in the immediate vicinity of the Braden Branch culvert are at least an order of magnitude above the minimum detection levels (0.3 pCi/g). The highest concentrations are within approximately 200 m of the culvert under the railroad bridge where Braden Branch flows into Melton Hill Reservoir. The concentration at the sample 1900G location suggests that contaminant transport has occurred in a southeasterly direction within the small channel. Low concentrations at the channel outlets suggest that downstream transport of ^{60}Co into the main channel of Melton Hill Reservoir may have been minimal in the recent past.

The ^{60}Co concentration profile from the sediment core indicates that there are several peaks in ^{60}Co concentrations that are two orders of magnitude above background levels. While concentrations in downstream cores suggest that downstream movement of contamination has been minimal, the results are inconclusive. The near-background concentrations in these cores could indicate that contaminant dilution has occurred or simply that the cores were not taken where contaminants had deposited.

These results and the results from previous studies indicate that Braden Branch has been a source of ^{60}Co contamination to Melton Hill Reservoir. Furthermore, the elevated concentrations in the surface sediments suggest that Braden Branch may still be contributing measurable amounts of ^{60}Co to the reservoir.

G.7 SCREENING HEALTH RISK ASSESSMENT

Quantitative assessment of human health risk from exposure to contaminants in the environment is extremely difficult because the data necessary to determine exposure and doses are difficult to obtain. These data include the speed and efficiency of contamination transport along various exposure routes, the magnitude and duration of the exposure, and the relationship between exposure and health risk. Because of the difficulty of acquiring such data, a set of assumptions are typically used to provide screening estimates of lifetime risk of an incremental cancer incidence. The Environmental Protection Agency has developed risk factors for radionuclides and provides guidance in their use to allow the conversion of contaminant concentrations in the environment into estimates of human health risk resulting from various exposure routes (EPA 1991a, b).

Results from screening level risk analysis for 1990 samples, which were collected in Braden Branch embayment, are presented in Table G.5 for nonconservative screening and in Table G.6 for conservative screening. The pathway of most concern is the external exposure pathway. Conservative screening risk due to direct exposure to sediments is 1×10^{-4} , which indicates it may be of concern. Nonconservative screening risk due to external exposure is 5×10^{-7} . All screening risk values from ingestion and inhalation pathways are $<1 \times 10^{-7}$ and, therefore, are of little concern.

It should be noted that the sediments in the Braden Branch area are exposed for some part of the year but probably not enough to allow for the 2000 h/year exposure included in the calculations; therefore, our calculations are conservative and probably are relevant only to the case in which sediments are dredged and used as a source of topsoil.

Table G.5 Results of adult nonconservative screening for carcinogens in surface sediment of Braden Branch where at least one measurement in a reach was above detection limits.

Media	Contaminant type	Contaminant	ratio of detects to total samples	Concentration (type)	Cancer slope factor	Intake rate	Carcinogen screening index
				Bq/kg	1/Bq	Bq/yr	
Sed. Exposure	Radionuclides	Co-60	20/20	2.92E+01 (G M)	2.3E-07	2.3E+00	5.3E-07
		Cs-137	20/20	1.18E+01 (G M)	5.4E-08	2.6E+00	1.4E-07
		TOTAL					6.7E-07

Sed. Exposure							6.7E-07

Sed. Ingestion	Radionuclides	Co-60	20/20	2.92E+01 (G M)	4.1E-10	3.2E+00	1.3E-09
		Cs-137	20/20	1.18E+01 (G M)	7.6E-10	1.3E+00	9.8E-10
		TOTAL					2.3E-09

Sed. Ingestion							2.3E-09

							6.7E-07
=====							

There is virtually no risk from ^{60}Co in deep, undisturbed sediment. Exchange with the water column is minimal, and water acts as an effective shield to gamma radiation. If the sediment is exposed due to draw-down of the water, the potential risk is similar to that of ^{60}Co in exposed sediment. The external exposure is then dominated by the size of the area containing ^{60}Co and the length of time an individual remains in the area of contamination as well as any physical features that act to partially or completely shield an individual from the gamma radiation.

G.8 REFERENCES FOR APPENDIX G

Miller, L.F., E.S. Stetar, J.H. Coleman, A. Payne, J. Hoffelt, and D.J. Auslander. 1987. Monitoring of Radionuclide Migration and Exposure Rates From the American Nuclear Corporation Plant Site. In: *Proceedings of the 1987 Oak Ridge Model Conference Volume III, Health and Safety*. CONF-871075—Vol. 3).

Environmental Protection Agency, 1991a. *Health Effects Assessment Summary Tables. Annual FY-1991*. OERR 9200.6-303 (91-1).

Environmental Protection Agency, 1991b. *OSWER Directive 9285.6-303*. March 25, 1991.

DISTRIBUTION

1. S. M. Adams
2. M. L. Ambrose
3. L. D. Bates
4. J. J. Beauchamp
5. D. T. Bell
6. M. S. Bevelhimer
7. B. G. Blaylock
8. C. C. Brandt
9. A. L. Brenkert
10. J. B. Cannon
- 11-15. R. B. Cook
16. W. D. Crumby
17. J. H. Cushman
18. M. F. P. DeLozier
19. C. J. Ford
20. D. E. Fowler
21. M. L. Frank
22. R. H. Gardner
23. C. W. Gehrs
24. M. J. Gentry
25. C. D. Goins
26. J. M. Grebmeier
27. R. S. Halbrook
28. P. J. Halsey
29. R. A. Harris
30. S. G. Hildebrand
31. S. K. Holladay
32. L. A. Hook
33. R. N. Hull
34. P. Kanciruk
35. B. L. Kimmel
36. L. A. Kszos
37. I. L. Larsen
38. D. A. Levine
39. J. M. Loar
40. R. C. Longman
41. S. E. Madix
- 42-44. D. M. Matteo
45. C. W. McGinn
46. M. E. Mitchell
47. S. L. Niemela
- 48-49. P. T. Owen
50. D. E. Reichle
51. K. A. Rose
52. G. E. Rymer
53. F. E. Sharples
54. D. S. Shriner
55. J. L. Skiles
56. S. H. Stow
57. G. W. Suter
58. R. I. Van Hook
59. D. R. Watkins
60. R. K. White
61. L. F. Williams
62. D. C. Wood
- 63-65. ESD Library
- 66-70. ER Document Management Center
71. Clinch River ER DMC
- 72-73. ORNL Laboratory Records
74. ORNL Patent Section
75. Office of Assistant Manager for Energy Research and Development, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8600
76. M. Ferre, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8541
77. J. F. Franklin, Bloedel Professor of Ecosystem Analysis, College of Forest Resources, University of Washington, Anderson Hall AR-10, Seattle, WA 98195
- 78-83. C. S. Gist, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8541
84. R. C. Harriss, Institute for the Study of Earth, Oceans, and Space, Science and Engineering Research Building, University of New Hampshire, Durham, NH 03824
85. F. O. Hoffman, SENES Oak Ridge, Inc., 677 Emory Valley Road, Suite C, Oak Ridge, TN 37830
86. G. Y. Jordy, Director, Office of Program Analysis, Office of Energy Research, ER-30, G-226, U.S. Department of Energy, Washington, DC 20545
87. K. Murray, Radian Corporation, 120 S. Jefferson Circle, Oak Ridge, TN 37831

- 88-89. R. L. Nace, Branch Chief, Nonenrichment Facilities, Oak Ridge Program Division, Office of Eastern Area Programs, Office of Environmental Restoration, EM-423, Trevion 2, U.S. Department of Energy, Washington, DC 20585
- 90. C. R. Olsen, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
- 91. R. H. Olsen, Professor, Microbiology and Immunology Department, University of Michigan, Medical Sciences II, #5605, 1301 East Catherine Street, Ann Arbor, MI 48109-0620
- 92. A. Patrinos, Director, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
- 93-94. R. C. Sleeman, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8541
- 95. D. W. Swindle, Radian Corporation, 120 S. Jefferson Circle, Oak Ridge, TN 37831
- 96-97. H. M. Thron, Chief, Enrichment Facilities, Oak Ridge Program Division, Office of Eastern Area Programs, Office of Environmental Restoration, EM-423, Trevion 2, U.S. Department of Energy, Washington, DC 20585
- 98. F. J. Wobber, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
- 99-100. Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831

**DATE
FILMED**

5 / 13 / 93