89
17.78

14.62

LA-7125-MS

Informal Report

UC-20i

Issued: April 1978

A Summary of Scyllac Results

R. E. Siemon

los alamos
scientific laboratory

of the University of California

LOS ALAMOS, NEW MEXICO 87545

An Affirmative Action/Equal Opportunity Employer

UNITED STATES
DEPARTMENT OF ENERGY
CONTRACT W-7405-ENG. 36

MASTER

DISTRIBUTION STATEMENT 1

Printed in the United States of America. Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

		Microfiche		\$ 3.00							
001-025	4.00	126-150	7.25	251-275	10.75	376-400	13.00	501-525	15.25		
026-050	4.50	151-175	8.00	276-300	11.00	401-425	13.25	526-550	15.50		
051-075	5.25	176-200	9.00	301-325	11.75	426-450	14.00	551-575	16.25		
076-100	6.00	201-225	9.25	326-350	12.00	451-475	14.50	576-600	16.50		
101-125	6.50	226-250	9.50	351-375	12.50	476-500	15.00	601-up	--1		

1. Add \$2.50 for each additional 100-page increment from 601 pages up.

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

A SUMMARY OF SCYLLAC RESULTS

by

R. E. Siemon

ABSTRACT

A series of theta-pinch high-beta stellarator experiments has been recently completed. They demonstrated the existence of a toroidal equilibrium at high beta (about 70%) and reveal the expected instability of gross plasma motion. Feedback stabilization, by means of optical sensors for plasma position coupled to current amplifiers for magnetic field control, works effectively to keep the plasma column near the center of the discharge vessel. Confinement of plasma in the recent feedback experiments is limited by classical end losses associated with the finite length toroidal sectors. This document gives a simplified history of Scyllac much as it was presented in an invited paper at the 1977 Plasma Physics Division meeting of the American Physical Society held in Atlanta, Georgia.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

INTRODUCTION TO THE TOROIDAL THETA PINCH

There have been many important contributions to the Scyllac work but F. L. Ribe is the person who deserves the most credit for the original idea and the experiments that have tested the concept. Figure 1 lists alphabetically only those experimenters who worked directly on Scyllac for a significant period of time (one year or more) and even so there are over 30 names. Not listed are the theorists who have made extremely valuable contributions in the planning and analysis of experiments and not listed are many scientists who have worked on experiments related to Scyllac such as the staged theta-pinch or the implosion heating experiment. The principal investigators during the most recent feedback stabilization experiments are listed at the bottom

SCYLLAC EXPERIMENTERS

F. L. RIBE

BARTSCH	HARRIS	McCONNELL
CALL	JAHODA	MILLER
CANTRELL	JOHNSON	QUINN
ELLIS	KAUFMANN	RAWCLIFFE
FREEMAN	KEMP	SAWYER
FREESE	KEWISH	SEIBEL
GRIBBLE	KLARE	SIEMON
GUTSCHER	KRISTAL	THOMAS
HAMMER	KUTAC	WELDON
HANDY	LILLBERG	ZIMMERMANN
HARDER	MACHALEK	

PRINCIPAL INVESTIGATORS FEEDBACK EXPERIMENTS

BARTSCH, CANTRELL, GRIBBLE, KLARE, KUTAC,
MILLER, AND QUINN

Fig. 1.

The main contributors to the Scyllac experiment.

of Fig. 1, and a complete description of results is available in a recent article by these authors submitted to Nuclear Fusion.

The original Scyllac concept grew out of a series of linear theta-pinch experiments (Scylla devices) in which hot (keV), dense ($\sim 10^{16}\text{-cm}^{-3}$) plasmas were created but lost rapidly because of open ends. The philosophy chosen was to bend a theta pinch gently into a closed torus (Scylla - closed or Scyllac) with as little perturbation as possible, consistent with the requirements of toroidal equilibrium and stability. Figure 2, a photograph of the Scyllac experiment during its operation as a full torus, gives the overall scale of a theta pinch when operated as a large aspect ratio torus. Not apparent in the

Fig. 2.
Scyllac, a toroidal theta pinch.

picture are the thousands of capacitors and switching circuitry required to produce the theta-pinch discharge.

THEORY OF THE HIGH-BETA STELLARATOR

Bending a linear stable theta pinch into a circle leads immediately to well-known problems. Figure 3 shows schematically a plasma of radius a in a torus with major radius R . The gradient in magnetic field strength due to toroidal curvature causes unfavorable particle drifts or equivalently a body force per unit length $F_R = \beta B_0^2 a^2 / 4R$. Written this way it is clear that plasma beta is troublesome and reducing beta would aid the equilibrium problem. On the other hand, plasma beta is essential to the economy of fusion reactors and a theta pinch readily creates a high-beta plasma. It was thus decided to seek a high-beta confinement scheme.

The method chosen for equilibrium was the high-beta stellarator. Figure 4 is a sketch of the plasma perturbations required by the theory. The main additional magnetic field requirement is termed helical $L=1$ and leads to a helical plasma shape. The shape repeats

Fig. 3.
Fundamental toroidal force proportional to beta.

Fig. 4.
High-beta stellarator plasma shape parameters δ_1 and δ_0 .

along the axis with a wavelength λ that has ranged in various experiments from 0.3 to 1.0 m. The actual helical distortion in experiments resembles that in Fig. 4 but is more elongated and thus more gently curved. The shape characterization δ_1 tells how far the plasma column center has moved away from the original unperturbed position in units of plasma radius.

According to the theory of a high-beta stellarator, either $L=0$ or $L=2$ fields must be added to the $L=1$ field for equilibrium. Most Scyllac experiments have used $L=0$, although recently $L=2$ has been favored. The plasma shape caused by $L=0$ fields as shown in Fig. 4 is a humpiness or variation in plasma cross-sectional area and is characterized by δ_0 . Not shown is the $L=2$ perturbation that leads to an elliptical plasma shape. An important distinction to be discussed

below is between the fields applied by control of external conductors ($L=1$, $L=0$, etc.) and the plasma response (δ_1 , δ_0 , etc.). Clearly the fields can be established precisely, but the plasma response is estimated by theory and measured by experiment.

OVERVIEW OF SCYLLAC EXPERIMENTS

Before turning to data and experimental details we start with a brief and oversimplified summary of the entire sequence of Scyllac experiments (Fig. 5) as they were carried out between 1970 and 1977. The initial task was to bend the linear theta into a circle with the good result that end losses by plasma flow and thermal conduction were eliminated. The immediate negative consequence is that plasma is not in equilibrium and moves rapidly to the wall of the quartz discharge tube.

As a matter of fact that was actually checked experimentally in the first Scyllac experiment. No helical equilibrium fields were applied. The plasma was observed to implode to the center of the quartz discharge tube. Then while being heated by compression from the rising field, the plasma at the same time accelerated within μs away from the axis until it impacted the walls of the tube.

The next step was to apply the high-beta stellarator fields, and the good result obtained was that the plasma no longer went immediately to the wall. Unfortunately, however, the theoretically anticipated result emerged experimentally, that the equilibrium was both dynamic and unstable. Given the many degrees of freedom for plasma instability, there was one encouraging aspect to the results: the only observed plasma instability was a long wavelength, whole-body or gross plasma motion. In the presence of helical fields the plasma was observed to oscillate in the horizontal plane (plane of the torus) until finally it moves as a whole--up, down, in, or out--until reaching the wall.

This type of $m=1$, $k_z=0$ instability suggests the use of feedback stabilization. Feedback in this context means the use of optical sensors to determine the plasma displacement, followed by adjustments in helical fields to force the plasma back to the axis of the vacuum vessel.

SUMMARY OF SCYLLAC

- BEND THETA PINCH INTO A CIRCLE (GOOD).
- SEE IT RUSH TO THE WALL (BAD).
- ADD EXTRA FIELDS - HELICAL $l=1$, $l=0$, $l=2$, ETC. (GOOD).
- FIND A DYNAMIC UNSTABLE EQUILIBRIUM (BAD).
 - BASIC PROBLEM LONG WAVELENGTH MODES.
 - SUGGESTS FEEDBACK STABILIZATION.
- IMPROVE EXPERIMENT (GOOD).
 - MACHINED FLUX SURFACES.
 - HELICAL QUARTZ DISCHARGE TUBE.
 - FAST "JONES" CIRCUIT TO TRIM EQUILIBRIUM.
- DISCOVER OUR FEEDBACK IS NOT POWERFUL ENOUGH FOR keV PLASMA (BAD).
- SUCCESSFULLY MODIFY EXPERIMENT TO DEMONSTRATE FEEDBACK STABILIZATION (VERY GOOD).

Fig. 5.
Simplified history of Scyllac.

To insure that the complications of feedback stabilization were necessary, a variety of improvements were incorporated into the experiments and further tests of high-beta stellarator theory were

performed. The preciseness of the helical fields was greatly improved by machining flux surfaces into the metal theta-pinch coil instead of applying fields with wire coils. Guided by work at Garching, a machine was built which forms a helical quartz discharge tube. The plasma which implodes to the center of the discharge tube is then formed in a shape which initially resembles the helical shape required by the high-beta stellarator. Oscillations in equilibrium forces were overcome by fast rising and accurately controlled "Jones" circuits used to power coils for trimming the equilibrium. Meanwhile, feedback power amplifiers with impressive parameters were constructed and feedback position detectors were developed. These necessary, painstaking, and time-consuming developments are the essence of experimental work although they tend to be underestimated in the planning of a research program.

By 1974 experiments on the full torus were completed and despite the continuing improvements in technique as just described, there emerged a somber conclusion in the work. Given the size of torus that Los Alamos Scientific Laboratory (LASL) could afford and the practical level of power that could be developed for the feedback system, there was no way to control the instability of the keV plasma produced in the theta pinch. It was back to the drawing boards to find a modification of the apparatus that would give feedback stabilization a reasonable test.

Fortunately the modifications were worked out successfully, and the bottom line of this research is that feedback stabilization works very much as originally conceived. This naturally is judged as "very good" by the researchers involved, and the remainder of the discussion presents evidence to support this conclusion.

CHARACTERISTICS OF EQUILIBRIUM

All of the early Scyllac experiments and some of the recent ones have been tested successfully in toroidal sector experiments. The sectors are smaller and thus permit more variations and results per unit time. This approach is limited to the well-established time scale associated with end effects (typically tens of μ s) but provides a reasonable basis for observing MHD equilibrium and stability behavior

Fig. 6.
Toroidal sector experiment showing machined flux surfaces.

with typical time constants of a few μ s. Figure 6 is a photograph of a recent Scyllac sector. The top half of the compression coil has been removed to show the shape of an $L=1,0$ flux surface used to create the equilibrium fields.

Some confusion in early experiments resulted from the dynamic nature of the toroidal equilibrium. Figure 7 summarizes the few equations needed to describe the problem. The plasma acceleration in the horizontal plane, $\ddot{x}$, is determined by the difference between the basic outward force, F_R , and the inward high-beta stellarator force, $F_{1,0}$, given by the helical $L=1$ and $L=0$ fields. Note that the $F_{1,0}$ force results from three terms. The first, $b_0 b_1$ represents the product of applied magnetic fields. The values of b_0 and b_1 represent the fraction of main toroidal field used in the helical $L=0$ and $L=1$ perturbations. If the fields are created by a machined flux surface, the product $b_0 b_1$ is independent of time. The other two terms in the $F_{1,0}$ force expression involve first a product of field, b_0 , and plasma shape, δ_1 , and last a product of field b_1 and shape δ_0 . The important

DYNAMICAL ASPECTS OF TOROIDAL EQUILIBRIUM

$$\ddot{X} = F_R - F_{I,0}$$

$$F_R = \beta V_A^2 / R$$

$$F_{I,0} = h V_A^2 \left(b_0 b_1 - \frac{h a}{2} b_0 \delta_1 + \delta_0 b_1 \right)$$

- b_0, b_1 PARAMETERIZE MAGNETIC FIELD

- δ_0, δ_1 PARAMETERIZE PLASMA SHAPES

Fig. 7.
Equations for force transients.

feature of these last two terms is that the plasma shapes δ_0 and δ_1 are involved. The initial transients associated with the plasma implosion in a theta pinch cause δ_0 and δ_1 to vary in time. There results a time variation in the $F_{I,0}$ force and, therefore, some accelerations of the plasma column as the plasma shape settles to its equilibrium value. The plasma tends eventually to steady-state values of δ_1 and δ_0 in response to fixed values of b_1 and b_0 .

Streak photographs as shown in Fig. 8 are an ideal diagnostic for these effects. The white stripe, created by the plasma luminosity, represents a direct measure of plasma displacement vs time. The horizontal displacement with time running left to right is shown in the left-hand side (plasma viewed from above) and the vertical displacement is shown on the right. The double stripe results from beam splitters that show two plasma views on each camera. The important characteristics seen on the upper pair of photographs are rapid oscillations in the first 10 μ s that upon analysis are found to be excursions in the helical plasma shape. The discharge tube in the upper photograph was not shaped as a helix. The lower streaks show the

Fig. 8.
Streak photographs showing plasma transients.

nearly complete elimination of helical oscillations by the use of a helical quartz discharge tube. There remains in the lower streaks a reproducible tendency for the plasma to move in the horizontal plane first one way and then the other before eventually moving unstably to the wall.

In Fig. 9 the acceleration of the plasma in the horizontal plane is plotted as a function of time. The net force clearly oscillates in time with a half-period of about $5 \mu s$. The oscillations of force and corresponding complex trajectories observed in streak pictures result from the transient plasma bumpiness or δ_0 response. The transients would not be so harmful except that a net force (outward) remains acting at late times (after about $8 \mu s$). The net force prevents effective feedback operation because large feedback currents are required just to maintain equilibrium. Therefore it was found necessary to develop circuits which fine tune the equilibrium at late times.

Detailed measurements support the interpretation of the time varying force as being a result of δ_0 transients. In Fig. 10 direct measurement of δ_0 is shown. The excluded flux is given as a function

Fig. 9.
Plasma acceleration in response to δ_0 transient.

of time at two different axial plasma positions where the plasma should be its maximum and minimum diameter due to bumpiness or δ_0 . Excluded flux depends on both plasma beta and density profile in a somewhat complicated way, but for the present purpose it is sufficient to think of it as directly measuring the plasma size or cross-sectional area. The curve labeled $hz=0$ is where magnetic field is locally maximum and plasma should be smallest. The data (solid lines) clearly indicate that the plasma has a smaller diameter than where $hz=\pi$. However, the ratio of diameters is not constant in time and shows directly the δ_0 transient. Notice that the half-period is about $5 \mu s$ in agreement with the force transients. One could guess that this time is about a

Fig. 10.

Plasma δ_0 transient observed directly by excluded flux.

helical half-wavelength divided by the Alfvén velocity, and, in fact, detailed MHD calculation as shown by the dotted curves gives that result in agreement with the experiment.

DEFINITIVE FEEDBACK RESULTS

Given that the equilibrium is adequately understood and measures are available to tune out transient forces at late times, the experiment is ready for the application of feedback stabilization. Figure 11 lists the characteristics of the $L=0,1,2$ sector on which the most definitive feedback results were obtained in about February of 1977. Equilibrium $L=0,1$ fields were machined into the compression coil. A helical quartz discharge tube suppressed δ_1 transients. The δ_0 transients were controlled by application of programmed $L=2$ currents. Feedback was applied with $L=2$ windings using about 50 MW/m

SCYLLAC $l=0,1,2$ SECTOR
CONFIGURATION

- HELICAL FIELDS MACHINED INTO THETA-PINCH COMPRESSION COIL.
- HELICAL QUARTZ DISCHARGE TUBE.
- $l=2$ EQUILIBRIUM TRIMMING WINDINGS.
- $l=2$ FEEDBACK ABOUT 50 MW/M.

PARAMETERS

R	4m
λ	63cm
B_0	14 kG
n_e	$2 \times 10^{16} \text{ cm}^{-3}$
a	1cm
T_e, T_i	100 eV
β	0.7

Fig. 11.
Features of $L=0,1,2$ sector experiment.

of feedback power. The other parameters of importance are listed at the bottom of Fig. 11.

Figure 12 shows how the equilibrium obtained using the "Jones circuits" is defined phenomenologically. The dotted lines represent plasma position in the horizontal plane on sequential shots with

Fig. 12.

Data showing that equilibrium is properly adjusted.

slightly varied trimming current. The solid curves show the current as it is applied after $8 \mu s$ to counter the remaining δ_0 transients. One level of current results in unstable plasma acceleration, in one direction, and a slightly different level of current leads to unstable acceleration in the opposite direction. With this type of fine-tuned equilibrium one can expect feedback to control the dynamics and prevent the unstable excursions.

The photograph in Fig. 13 shows the experiment in operation. Five cylindrical tubes containing position detectors can be discerned as they are mounted 45 degrees from vertical. The position detectors are

Fig. 13.
Feedback apparatus installed on L=0,1,2 sector.

arrays of silicon photodiodes which generate microamps per millimeter of plasma displacement. Cables in copper conduit relay the signal to the feedback modules--large rectangular boxes at the top of the photo showing round cooling fans on their sides. These modules amplify the position detector signals and produce hundreds of amps in the L=2 windings that are located as closely as possible to the plasma. Figure 14 shows how the L=2 feedback windings are wound on the helical quartz discharge tube. The oval shaped perturbations are required to permit plasma diagnostic access for Thomson scattering, interferometry, streak photos, and so forth.

In Fig. 15 a schematic layout of the sector is shown. The point here is to explain what is meant by "local feedback." The first position detector station, PD_1 , actually consisting of two position detectors viewing the plasma in orthogonal directions, is used to control the feedback currents in wavelengths one and two of the sector. The next station, PD_2 , controls current in wavelengths three and four, and so forth around the sector. The feedback is "local" in the sense

Fig. 14.
helical discharge tube wrapped with $L=2$ coils.

that information about behavior in one part of the sector is used to control currents in only that part of the sector. We shall see that this approach leads to feedback generated oscillations where adjacent position detectors struggle against one another.

In Fig. 16 streak photographs show the dramatic improvement that results with the application of feedback. The upper pair of streaks without feedback show a typical unstable excursion leading to a wall encounter in the horizontal plane at about 20 μ s. In the lower pair of streaks, at the same observation stations, and with no other change except the application of feedback power, we see plasma confinement extended to 50 or so μ s. Classical end effects dominate the plasma after about 40 μ s, so no better confinement than that shown can be expected in the sector experiment.

For the same shots just used to compare feedback on and off, we have position detector data and magnitude of $L=2$ field shown in Fig. 17. The b_2 field shown in the lower half of Fig. 17 results from the equilibrium trimming in the absence of feedback. The modified b_2

Fig. 15.
Arrangement of position detectors for local feedback.

Fig. 16.
Comparison of streak photos with and without feedback.

Fig. 17.
Plasma position response to adjustments in b_2 field by feedback.

waveform in the upper half of the figure is a result of feedback responding to plasma motion. Again it is clear that plasma behavior, as seen by position detectors in this case, is improved by feedback in that the unstable excursion after 20 μs without feedback is eliminated by the application of feedback control.

Various tests of the feedback systems were made to insure that we were interpreting the results correctly. For example, the polarity of position detectors was reversed so that, say, an upward excursion results in an upward force. The results were unambiguous--the plasma was driven hard against the walls of the discharge tube. A more subtle variation is to increase the feedback gain, i.e., proportionality between position and $L=2$ current. According to feedback theory, too much gain leads to oscillations in the system. In the lowest streak photo of Fig. 18, we see the onset of these oscillations when the gain has been increased by a factor of 4.

VERTICAL PLANE

Fig. 18.
Streak photos showing response to increased feedback gain.

In Fig. 19 we have data which indicate some limitations in local feedback control. Plasma displacement in the vertical direction (shown with solid lines) is plotted for three adjacent position detector stations. Without feedback there are no force transients in the vertical direction. If the column starts upward, it continues upward with a simple form of unstable motion. In Fig. 19 with feedback applied, we see oscillations in vertical position. They are clearly associated with the feedback current oscillations shown with dashed curves. Notice that position detectors one and two are out of phase. This corresponds to an oscillation with an axial wavelength of about

Fig. 19.
 Position detector data showing short wavelength modes excited by feedback system.

four helical wavelengths. Such oscillations would be stable without feedback but are driven into oscillation by the feedback system primarily because the module response time is finite.

RECENT SCYLLAC IMPROVEMENTS

Based on the growing appreciation of subtle problems in the $L=0,1$ equilibrium and the nature of local feedback control, a new configuration was designed. The main features are indicated in Fig. 20. The equilibrium has only $L=1$ and $L=2$ fields, and feedback is applied with $L=2$ windings. By not using $L=0$ fields one eliminates the troublesome δ_0 transient, and according to theory, δ_2 transients are inconsequential to the force balance.

A fairly complex modal feedback system was developed for use on the $L=1,2$ sector. Position detector data from many axial stations are analyzed by a fast processor--essentially an analog computer that does fast fourier transforms on the plasma axial mode structure. In response to each measured axial mode, a current pattern in the $L=2$ windings is generated to suppress that particular plasma motion. In this case short wavelength oscillations like those generated by local feedback control are ignored and thus not excited by the analog processor as it analyzes position information.

MAIN FEATURES $l=1,2$ EXPERIMENT

- NO $l=0$ FIELD.
 - ELIMINATES δ_0 TRANSIENT.
- MODAL FEEDBACK.
 - STOPS FEEDBACK OSCILLATIONS.

Fig. 20.
Improvements in $L=1,2$ experiment.

Unfortunately, results from the L=1,2 sector are limited because not enough time was available to completely diagnose the experiment. It was agreed to extend the Scyllac operation through the end of August for the measurements, and in fact, most data were obtained between August 20th and 40th.

Figure 21 shows streak photos of plasma motion without feedback. The qualitative improvement that resulted from eliminating δ_0 transients was striking. Even without feedback it was possible to obtain significant confinement. However, analyses of position detector data on shots like the one in Fig. 21 generally indicate some wall contact somewhere in the sector at typically 30 or so μs . The motion is qualitatively more "gentle" in the L=1,2 sector than in L=0,1 sector and the growth rate is less than one expects from theory.

Fig. 21.
Streak photos of equilibrium behavior in L=1,2 sector.

Improvement with feedback is shown in Fig. 22. The upper streaks indicate a general level of motion that is suppressed by feedback, as shown in the lower set of streaks. Confinement is extended in these experiments to about 70 μ s--a limit that is again consistent with classical end loss.

Data in Fig. 23 show the drop in electron temperature due to end effects as measured by Thomson scattering. Considerable scatter in the data is unavoidable because the laser scattering source is fixed in space while the plasma moves around a few centimeters in the discharge tube. Because the plasma column is cooler on the edge than in the center, measurements at late times tend to underestimate the central plasma temperature. In any case, data in Fig. 23 show a cooling trend for 60 μ s, which is consistent with classical radial confinement and cooling from the ends of the finite length sector.

Other data from the L=1,2 sector are still being analyzed and will be published in the near future. One feature of interest is a surprising level of undamped oscillations in the helical δ_1 distortion. The oscillations are independent of feedback and are not clearly detrimental to the plasma confinement.

Fig. 22.
Streak photos of feedback effects on L=1,2 sector.

Fig. 23.
Electron temperature vs time shows cooling caused by sector length.

CONCLUSIONS

Figure 24 states the main conclusions to be drawn from work of the last year on the Scyllac experiment. The clear evidence for a toroidal high-beta stellarator equilibrium and the efficacy of feedback stabilization are important advances in our quest for magnetic confinement schemes that can lead to practical fusion power. A reasonable case for a fusion reactor based on these experiments has been made elsewhere.

CONCLUSIONS

- SCYLLAC $l=0,1,2$ SECTOR EXPERIMENT.
 - DEMONSTRATES EQUILIBRIUM.
 - FEEDBACK IMPROVES CONFINEMENT.
 - CONFINEMENT LIMITED BY SECTOR LENGTH.
 - SUGGESTS NEED FOR MODAL FEEDBACK.

- SCYLLAC $l=1,2$ EXPERIMENT (LIMITED STUDY).
 - SHOWS EQUILIBRIUM LESS DYNAMIC.
 - SUGGESTS STABILITY BETTER THEN THEORY PREDICTS.
 - CONFINEMENT LIMITED BY SECTOR LENGTH.
 - SUBJECT TO UNEXPECTED OSCILLATIONS INDEPENDENT OF FEEDBACK.

- UNRESOLVED QUESTIONS.
 - CONFINEMENT WITH FEEDBACK IN FULL TORUS.
 - WALL STABILIZATION (EXPERIMENT PROPOSED AT GARCHING).

Fig. 24.

Main conclusions of the recent Scyllac sector experiments.

Important unresolved questions at the present level of understanding are listed. Because feedback was only tested in sector experiments it is not certain what confinement limits would prevail in a full-torus experiment.

A convincing result of the high-beta stellarator theory is the existence of wall stabilization with no requirement for feedback when plasma parameters are properly adjusted. An experiment to test this

option has been proposed at Garching and we thoroughly endorse that proposal.

Acknowledgments

Particular credit is due G. Miller for translation of the high-beta stellarator theory into expressions suitable for comparison with experiment. This talk was prepared with the assistance of R. Bartsch, E. Cantrell, R. Gribble, G. Miller, and W. Quinn.