2
12/2/78

CONF-780946--5

MASTER

MASTER

PREPRINT UCRL-81586

Lawrence Livermore Laboratory

The Physics of Field Reversed Mirrors

R. F. Post

September 8, 1978

This paper was prepared for submission to the Fourth International Conference on Driven Magnetic Fusion Reactors, Erice, Italy, September 18-26, 1978.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

Commission of the European Communities
Fourth International Conference on
Driven Magnetic Fusion Reactors
Lisbon, Italy September 15-26, 1976

THE THEORY OF FIELD-REVERSED MIRRORS*

R. L. BOYD
University of California
Lawrence Livermore Laboratory
Livermore, California 94550

NOTICE
This report was prepared for the U.S. Department of Energy by the Lawrence Livermore Laboratory under contract number W-7405-ENG-48. The U.S. Government is authorized to reproduce and distribute reprints for government purposes not withstanding any copyright notation that may appear hereon. This report is available from the U.S. Department of Energy, Office of Scientific and Technical Information, Springfield, Virginia 22161.

ABSTRACT

Since the earliest days of fusion research it has been hoped that diamagnetic currents flowing in a plasma could be used to help confine the plasma. Recently this hope has been strengthened both by theoretical advances and by experimental results made possible by technological developments. On the theoretical front analytical treatments and computer simulation studies have demonstrated equilibrium solutions existing both in the finite orbit and in the large-orbit limit. Progress has also been made in identifying the conditions required for the stability of field-reversed states. It appears that configurations of the general form of fig. 1(b) might, possibly elongated to napkin-ring form, represent stable states. Building on previous experimental work, several investigators have been able to create field-reversed states. One method, based on the FIRM idea of Christofilos, traps an intense relativistic electron beams (REB) to create a field-reversing current ring. Other approaches use either the reversed field theta pinch technique or REB pulses to create field-reversing diamagnetic currents in a long cylindrical plasma. In the former method, millisecond-long field-reversing electron rings have been achieved; in the latter method field-reversed plasma states lasting 30 to 50 microseconds have been achieved. Another approach under investigation is the Field Reversed Mirror (FRM) created by the tangential injection of high current neutral beams. Plasma states that approach field reversal have been achieved by this technique. The motivations for achieving field reversed states are strong: The possibility of achieving compact high density confined plasma states for fusion power generation that make maximal effective use of the confining field.

*Work performed under the auspices of the U. S. Department of Energy by the Lawrence Livermore Laboratory under contract number W-7405-ENG-48.

1. INTRODUCTION

Since the earliest days of magnetic fusion research the idea of using magnetic fields arising from within a plasma to help to confine that plasma has intrigued fusion researchers. Perhaps the earliest example of this concept was the so-called toroidal pinch, originally conceived as a thin bicycle-tire-like plasma entity self-confined by currents flowing within the plasma. In the toroidal pinch the plasma was to exist in pressure equilibrium with its self magnetic field, an equilibrium calculated by an extension of the so-called Bennett pinch equilibrium.

As was soon evident experimentally, the simple toroidal pinch was doomed to fail, owing to the rapid growth of hydromagnetic instabilities in the pinched column. As is evident from first principles, such a column, if sustained in pressure equilibrium solely by its self magnetic field, is subject to two types of MHD instability modes, the kink mode and the pinch-off mode, illustrated schematically in Fig. 1. Driven by magnetic forces, these modes are fast-growing and catastrophic in their effect on the pinched column - either causing it to writhe like a snake (the kink mode) until the column dashes itself against the chamber wall, or to neck down (the pinch-off mode) with the generation of intense electric fields through violent current interruptions.

Although research into field-reversed plasma states has gone far beyond the primitive example of the toroidal pinch, the primary physics issue involved in researching these states is the generic descendant of the critical issue in the toroidal pinch that we have discussed. The issue: defining equilibrium states that are at the same time stable against magnetically driven MHD modes.

From a practical point-of-view an issue equally as important as that of equilibrium and stability is the question of how to create a field-reversed state. Technical means must be found to initiate and subsequently to sustain a pattern of electrical currents within the field-reversed entity that will, in combination with externally generated magnetic fields, create the desired magnetic field configuration. Among the means being used to accomplish this end are the following:

- Trapping high energy electrons (or ions) injected into the con-

finement region, thereby creating a diamagnetic current shell (Christofilos' ASTRON and its descendants).

- Utilizing the flux-conserving property of a conducting plasma to capture oppositely-directed magnetic flux in an externally reversed transient magnetic field (reversed-field theta pinch).
- Employing tangentially-injected high energy neutral beams to maintain steady-state diamagnetic currents within a field-reversed entity (The Field Reversed Mirror as studied at the Lawrence Livermore Laboratory).

From a fusion power standpoint the motivations for achieving field reversal are economic in nature: Field reversed entities, being intrinsically high beta states, make very efficient use of the magnetic field. For the same reason they are compact; a field reversed entity generating tens of megawatts of fusion power might be about the size of a soccer ball. Finally, plasma confinement times in field reversed entities could be substantially enhanced as compared to confinement times in the non-reversed mirror-confined states from which they may be derived (as in the Field Reversed Mirror).

Despite the esthetically and economically attractive features of field reversal it must be recognized that the physics issues raised in attempting to achieve such states are profound ones. To achieve a stable field-reversed state is clearly a more difficult task than to achieve a stable low beta plasma state where the plasma diamagnetic currents only slightly perturb the confining field. But the potential rewards, both in simplicity and in economic terms, are so great that the pursuit of field reversal is well worth the effort. In the section to follow we will review selected examples of theoretical work on equilibrium and stability of field reversed states and cite examples of experiments of various kinds in which transient field reversed states have been achieved, or in which a state of field reversal has been approached.

II. FIELD REVERSED EQUILIBRIA

The most elementary field reversal equilibrium to visualize would be that of a bicycle tire toroidal pinch immersed in a constant B field directed along the major axis of the toroid. But since such a general

configuration, in this limit, is evidently MHD unstable, attention has been turned toward configurations that should be less subject to the kink and pinching modes that spelled the death knell of the simple pinch. On intuitive grounds one might expect flat fattened rings, more resembling a rotund doughnut than a torus, with an elongated field reversing current layers, as visualized for the A120N, would be less subject to kinking and pinching off. The general kind of field configuration we are here describing is illustrated schematically in Fig. 2, as it might be created in a Field-Reversed Mirror (FRM) as visualized at Livermore. Here the field-reversed plasma entity, that individual region bounded by closed magnetic lines, is itself bounded by the field lines of the mirror field within which it is formed.

Beyond qualitative visualization, however, the problem of calculating specific equilibria in quantitative terms is a relatively difficult one, one where analytical results have been obtained only in special cases. Two general classes of solutions have been investigated, representing two limits of the class of field-reversed entities.

- The Fluid Limit. Here the plasma is treated as a structureless resistanceless electrically conducting fluid. Field reversal results from the presence of electrical currents that exist in the body of the plasma in order to satisfy the requirements of pressure balance.
- The Maxwell-Vlasov or particle kinetic limit. Here the reversed field configuration is determined through self-consistent evaluation of the trajectories of energetic particles (electrons or ions) executing orbits in a magnetic field configuration which is a superposition of the externally applied field and the field associated with the particles' orbital motions.

In actuality, real reversed field plasma states lie in a continuum which must encompass both limits, since real plasmas will contain particles of both signs of charge spanning a wide range of energies. Nevertheless the various special cases that have been worked out, though not wholly realistic, provide valuable insight and represent starting points for more realistic evaluations of the problems of field reversed equilibria and stability. What all of the solutions have in

common is the fact that they must exploit the diamagnetic properties of gyrating particles to the point that they not only depress the field toward zero in the interior of the plasma ($r > 1$), but that they have sufficient capability to actually reverse the field direction in regions near the outer edge of the cavity. In what follows we will not attempt to cover more than a few illustrative examples.

A. The Fluid Limit

An informative relation can be derived in the fluid limit, using equations (1) and (2), the determination of general, axially-symmetric, toroidal equilibria. Grad (1) and Shafranov (2) have derived a partial differential equation for the flux function ψ , to be derived below, from which the equilibria may be determined. Our starting point is the pressure balance relationship for a static plasma pressure:

$$\nabla p = -\frac{1}{c} \nabla \psi \times \nabla \psi \quad (1)$$

From this relationship we immediately find that the lines of B_z , lying on constant ψ surfaces, are also surfaces of constant p . Considering now one of the flux (ψ) or constant p surfaces we note that any closed curve on this surface is linked both by the same flux and the same total current. We may therefore use the variables ψ and I to specify the relation itself, that is, the self-consistent distribution of magnetic field. In addition, and if we may use Maxwell's equations and Stokes' Theorem, obtaining

$$\oint \vec{B} \cdot d\vec{l} = (4\pi/c) I \quad (2)$$

$$\text{and} \quad \oint \vec{A} \cdot d\vec{l} = \psi, \quad (3)$$

where the line integral is taken around any axis-encircling closed contour lying on the ψ surface in question. From (2) and (3) we find immediately:

$$B_\theta = 2I/cr, \quad A_z = c/2\pi r. \quad (4)$$

Then, since $\vec{B} = \nabla \psi \times \vec{A}$ and $\nabla \times \vec{B} = (4\pi/c) \vec{j}$ we have

$$B_z = \frac{1}{2\pi} \left(\frac{1}{r} \frac{\partial \psi}{\partial r} \right) \quad B_r = -\frac{1}{2\pi} \left(\frac{1}{r} \frac{\partial \psi}{\partial z} \right) \quad (5)$$

$$j_z = \frac{1}{2\pi r} \frac{\partial I}{\partial r} \quad j_r = -\frac{1}{2\pi r} \frac{\partial I}{\partial z}$$

Knowing I and ψ all field quantities and current densities can be found. The remaining current density, j_θ , is related to p and ψ through the equation

$$j_\theta = C(\psi) + D(\psi) \quad (6)$$

$$\text{where } C(\psi) = 2\pi c \frac{dp(\psi)}{d\psi}$$

$$D(\psi) = \frac{1}{c} \frac{dI^2(\psi)}{d\psi}$$

Note that we are here permitted to write $p = p(\psi)$ and $I = I(\psi)$ since these are all surface quantities, that is p and I are constant on surfaces of constant ψ .

The scalar partial differential equation that results from the above considerations is

$$\begin{aligned} \frac{\partial^2 \psi}{\partial z^2} + \frac{\partial^2 \psi}{\partial r^2} - \frac{1}{r} \frac{\partial \psi}{\partial r} &= -\frac{8\pi^2}{c} r j_\theta \\ &= -\frac{8\pi^2}{c} [C(\psi)r^2 + D(\psi)] \end{aligned} \quad (7)$$

Outside the plasma $j_\theta = 0$ and $p = 0$ and the equation is of homogeneous form.

General toroidal equilibrium can be found by choosing various forms for $C(\psi)$ and $D(\psi)$ as appropriate to the situation at hand. For our purposes a special case, obtained by setting $C(\psi) = \text{constant}$, $D(\psi) = 0$, illustrates a field reversed equilibrium. From the definition of $C(\psi)$ above we see that this case corresponds to the case $j_\theta \sim r$ within the plasma, a not unreasonable physical assumption.. The resulting equilibrium is sometimes called the "Hill's Vortex Solution" because of its mathematical duality with Hill's [3] solution of an hydrodynamic equilibrium involving stable vortex motion in a fluid. Within the plasma ($j_\theta \neq 0$) the solution obtained is

$$\psi = -\frac{1}{2} \left(\frac{r^2}{R^2} \right) (2R^2 - r^2 - 4z^2) \quad (8)$$

Here R and z are parameters determining the physical size and the ellipticity (elongation), respectively, of the field reversed entity.

Outside the plasma $p = 0$ and $j_z = 0$ and the above solution must be replaced by the solution to the homogeneous equation, matching solutions at the plasma boundary.

In the special case of a spherical plasma boundary, of radius a , where the field reversed entity is embedded in a uniform vacuum magnetic field $B_z = B_0$, the solutions take on a relatively simple form

$$\psi = -\frac{3B_0}{2a^2} r^2 \left[a^2 - (r^2 + z^2) \right], \quad (r^2 + z^2) \leq a^2 \quad (9)$$

$$\psi = -\frac{3B_0}{2} r^2 \left[1 - \frac{a^2}{(r^2 + z^2)^{3/2}} \right], \quad (r^2 + z^2) \geq a^2 \quad (10)$$

Field lines of this configuration, corresponding to $\psi = \text{constant}$ surfaces, are shown plotted in Fig. 3. From the solution above we may also determine the variation of B within the plasma, thus, the degree of field reversal achieved. At the midplane ($z = 0$), for example, we find

$$\begin{aligned} B = B_z &= \frac{1}{2} \cdot \left(\frac{1}{r} \frac{\partial \psi}{\partial r} \right) \bigg|_{z=0} \\ &= -\frac{3B_0}{2a^2} \left[a^2 - 2r^2 \right], \quad r \leq a \end{aligned} \quad (11)$$

Evaluated at $r = 0$, $B(0) = -(3/2)B_0$, i.e. the field reversal factor $B(0)/B_0 = -3/2$. Thus at $r = 0$, $z = 0$, the field is reversed and has an intensity $3/2$ times that of the initial (vacuum) field.

The plasma pressure profile may also be evaluated, since

$$\frac{dp}{d\psi} = \frac{1}{2\pi c} C(\psi).$$

Thus we find $p \sim \psi + \text{constant}$

$$p = \left(\frac{B_0^2}{8\pi} \right) \left\{ -\frac{3 \cdot 15}{4} \left(\frac{r^2}{a^2} \right) \left[1 - \left(\frac{r^2}{a^2} + \frac{z^2}{a^2} \right) \right] \right\} \quad (12)$$

The maximum in p occurs at $z = 0$, $r = a/\sqrt{2}$ where

$$p_{\max} = \left(\frac{3 \cdot 15}{16} \right) \left(\frac{B_0^2}{8\pi} \right) \approx 2.8 \left(\frac{B_0^2}{8\pi} \right) \quad (13)$$

We see that owing to strong self-confining forces that arise as a result of the field-reversing diamagnetic currents, the value of $p_{\max} = p_{\max}/(B_0^2/8\pi) \approx 2.8$. This illustrates a property of field-reversed states of considerable potential economic significance. Such states can in principle confine plasmas with peak pressures that exceed the magnetic pressure of the vacuum fields within which they are embedded.

In Fig. 4 B_z/B_0 , B_z^2/B_0^2 and $p_0 = p_0/(B_0^2/8\pi)$ evaluated at $z = 0$ are plotted as a function of r/a . Note that the null point in B_z and B_z^2 occurring at $r/a = 1/\sqrt{2}$ coincides with the maximum value of p , as might have been expected intuitively. That is, $p_{\max}$ occurs at the low point in the self-dug magnetic well created by the field-reversed entity.

The solution just discussed is but a special case of a very wide class of axially symmetric field-reversed equilibria derivable from Eq. (7). This equation also describes equilibrium states appropriate to the tokamak. In fact, the tokamak plasma is a kind of "field reversed" entity, one in which the poloidal field reverses direction between the inner and outer surface of the confined plasma.

Thus we note particularly that Eq. (7) is not restricted to solutions of the type we have discussed, where the toroidal B field, B_θ , has been assumed equal to zero. In fact, field reversed entities of the same general configuration as we have been discussing exist for $B_\theta \neq 0$. An example is the "Spheromak" configuration discussed by Bussac, Furth, et al., [4] and others. As mentioned later, the incorporation of a toroidal field within a field-reversed entity may enhance its MHD stability and may therefore be an important element in achieving a practical configuration.

8. The Particle Orbit Limit

Analytic solutions for field-reversed equilibria may also be obtained in what might be called the "particle orbit" limit. That is to say, it is possible to find sets of self-consistent particle orbits that generate field reversal in steady state through their own motion

in the net magnetic field created by that motion and by external coils. This problem represents an extension into the domain of field reversal of the general problem of calculating high beta equilibria. As such it represents a difficult task, one which has thus far only been accomplished for special models.

One such model, which the author has considered, Post [5], is the case of a long (z-independent) cylindrical entity. The current carriers (ions) are characterized by a distribution function that is the product of delta functions in energy and canonical angular momentum, two of the three invariants of the problem. The third invariant is the longitudinal momentum, $p_z = Mv_z$. Since any distribution function expressed in terms of the invariants of the particle motion is a solution to the Vlasov equation, it is only necessary to simultaneously satisfy Maxwell's equation to generate equilibrium solutions.

The present problem has been earlier discussed by Marx, [6] using a different approach. The approach taken here starts with the invariants P_ϕ and E , canonical angular momentum and particle energy, and solves for the self-consistent particle orbits. We have

$$P_\phi = \left[p_\phi + (e/c) A_z \right] = \left[Mv_\phi r + \frac{e}{2c} \cdot \right] = \text{constant} \quad (14)$$

$$E = (1/2M) (p_r^2 + p_z^2 + p_\phi^2) = \frac{1}{2} Mv^2 = \text{constant} \quad (15)$$

and, for the magnetic field,

$$\begin{aligned} B(r) &= B_0 + 4\pi j(r) & r < a \\ &= B_0 & r > a \end{aligned}$$

$$I_\phi(r) = \int_r^a j_\phi(r') dr' \quad (16)$$

Then,

$$\begin{aligned} \psi(r) &= \int_0^r 2\pi r' B(r') dr' \\ &= \pi r^2 B_0 + 8\pi^2 \int_0^r r' dr' \int_{r'}^a j_\phi(r'') dr'' \quad (17) \end{aligned}$$

Thus we have

$$p_\theta = Mv_\theta r + \frac{e}{2\pi c} \left\{ r^2 B_0 + 8\pi^2 \int_0^r r' dr' \int_{r'}^a j_z(r'') dr'' \right\} \quad (18)$$

To proceed further we must establish a relationship between j_θ and v_θ , thereby allowing us to pose the problem as an integral equation for v_θ . This can be accomplished, for the chosen distribution function, as follows:

We have, in general, that

$$\vec{j} = \frac{e}{M} \int \vec{p} f(\vec{p}) d\vec{p}, \quad (19)$$

so that

$$j_\theta = \frac{e}{M} \iiint p_\theta f(p_r, p_\theta, p_z) dp_r dp_\theta dp_z \quad (20)$$

We may write this in terms of the three constants of the motion, E , p_θ and p_z as

$$j_\theta = \int_{p_z} \int_{p_\theta} \int_E p_\theta(E, p_\theta) f(E, p_\theta, p_z) J \left[\frac{p_r, p_\theta}{E, p_\theta} \right] dE dp_\theta dp_z, \quad (21)$$

where the Jacobian

$$J \left(\begin{matrix} p_r, p_\theta \\ E, p_\theta \end{matrix} \right) = \frac{M}{rp_r},$$

and

$$p_\theta = \left(\frac{p_\theta}{r} - \frac{e}{c} A_\theta \right) = Mv_\theta \quad (21)$$

$$p_r^2 = 2ME - \left(\frac{p_\theta}{r} - \frac{e}{c} A_\theta \right)^2 - p_z^2 \quad (22)$$

$$j_\theta = \frac{e}{r} \int_{p_z} \int_{p_\theta} \int_E \frac{\left(\frac{p_\theta}{r} - \frac{e}{c} A_\theta \right) f(E, p_\theta, p_z)}{\left\{ 2ME - \frac{p_\theta^2}{r^2} - \frac{e^2}{c^2} A_\theta^2 - p_z^2 \right\}^{1/2}} dE dp_\theta dp_z \quad (23)$$

we now take

$$f(E, p_z, p_z) = A(E - E_0) \delta(p_z - p_0) g(p_z) \quad (24)$$

There results

$$j_z = \frac{e}{r} \int_{-p_{z0}}^{p_{z0}} \frac{\left(\frac{p_0}{r} - \frac{e}{c} A_z\right) g(p_z) dp_z}{\left\{ 2ME_0 - \left(\frac{p_0}{r} - \frac{e}{c} A_z\right)^2 - p_z^2 \right\}^{1/2}} \quad (24)$$

where

$$p_{z0}^2 = 2ME_0 - \left(\frac{p_0}{r} - \frac{e}{c} A_z\right)^2$$

Take $g(p_z) = f_0 = \text{constant}$. There results

$$j_z = \frac{ef_0}{r} \left(\frac{p_0}{r} - \frac{e}{c} A_z \right) \quad (25)$$

$$\text{But } \left(\frac{p_0}{r} - \frac{e}{c} A_z \right) = p_z = mv \quad (26)$$

Thus we find

$$j_z = Mef_0 (v_z/r), \quad -v_0 \leq v \leq v_0 \quad (27)$$

Similarly, we find

$$\begin{aligned} n(r) &= -Mf_0/r & s \leq r \leq a, \\ &= 0 & \text{otherwise,} \end{aligned} \quad (28)$$

where s and a are the inner and outer boundaries of the layer. We thus find that $j_z \sim (v_z/r)$ and that $n(r) \sim 1/r$ between the inner and outer boundaries. Now knowing the relationship between j_z and v_z , Eq. (18) may be converted to an integral equation from which $v_z(r)$ and all other relevant quantities may be determined. Defining

$$y = v_z/v, \quad -1 \leq y \leq 1,$$

Eq. (18) becomes an integral equation for y :

$$y = \frac{K_0}{r} - \left\{ K_1 r + \frac{K_2}{r} \int_0^r r' dr' + \int_{r'}^a \frac{y dr'}{r'^{1/2}} \right\} \quad (29)$$

Here $K_0 = P_0/Mv$, $K_1 = v_{c0}/2v = 1/(2a_0)$, $v_{c0} = eB_0/Mc$; $a_0 = v/v_{c0}$, and $K_2 = 4\pi K_0 v_{c0}/3v$, with K_1 being the proportionality constant between j_θ and v_θ that will determine the strength of the layer;

$$j_\theta = K_0 v_\theta / r.$$

This integral equation may be converted by differentiation to a differential equation of Bessel form. Let $x = K_2 r$, a dimensionless radius, then (29) becomes

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} - \left(\frac{1}{x} + \frac{1}{x^2} \right) y = 0 \quad (30)$$

The general solution to this equation is

$$y = AI_2(2x^{1/2}) + BK_2(2x^{1/2}); \quad |y| \leq 1. \quad (31)$$

The constants A and B may be determined by inserting the boundary conditions. Considering the extremum orbits, that is those particles which traverse between the innermost and outermost boundaries of the layers, these are: at $r = a$ $y = -1$, i.e. $v_\theta = -v$, corresponding to purely θ motion, in the diamagnetic direction. At $r = s$, if we are looking for field-reversed solutions, we must also require $y = -1$ (diamagnetic current direction). The boundary condition $y = +1$ at $r = s$ corresponds to a field-enhancing rather than a field-reversing condition.

One more condition must be imposed to obtain a solution in closed form. We must establish a relationship between the strength of the current layer (proportional to K_0) and the dimensions of the layer. This relationship may be obtained by noting that at the outer turnaround point, $r = a$, the outermost boundary of the layer, the radius of curvature of the particles at their turnaround must be a_0 , their gyroradius in the vacuum field. It may readily be shown that the desired relationship may be expressed in the form

$$\frac{a}{a_0} = \left[1 - a \frac{dy}{dr} \right]_{r=a} \quad (31)$$

Defining now the dimensionless layer strength parameter $\lambda = K_2 a_0 (a_0/a)$, and performing the indicated differentiation of the solution, Eq. (31), we find a transcendental equation for λ in terms of the dimensionless

radius parameters of the layer, (a/a_0) and (s/a) . We have

$$\lambda^{1/2} = AK_1 \left[2 \left(\frac{a_-}{a_0} \right)^{1/2} \right] - A_1 \left[2 \left(\frac{a_-}{a_0} \right)^{1/2} \right], \quad (32)$$

$$\text{where } A = - \left[K_2'(s) - K_2'(a) \right] / D$$

$$B = - \left[I_2'(s) - I_2'(a) \right] / D$$

$$D = K_2'(s) I_2'(a) - I_2'(s) K_2'(a)$$

$$A = 2 \left(\frac{a_-}{a_0} \right) \left(\frac{s}{a} \right)^{1/2} \lambda^{1/2}$$

$$B = 2 \left(\frac{a_-}{a_0} \right) \lambda^{1/2}$$

Knowing the extremum solution for y the value of the self-consistent magnetic field may readily be found through the easily-derived relationship:

$$\frac{B(r)}{B_0} = 1 + \left(\frac{a_-}{a_0} \right) \int_r^a \frac{y \, dr'}{r'^2}, \quad (33)$$

where $y(r)$ is the solution for the extremum orbits, Eq. (31).

Using the above formulation we may now specify any possible self-consistent equilibrium solution with the chosen distribution function in terms of the two dimensionless parameters, a/a_0 and s/a , and the additional specification whether $y = -1$ (diamagnetic turnaround) or $y = +1$ (paramagnetic turnaround) at $r = s$. Given these, λ may be determined through Eq. (32) and knowing λ , $y(r)$ for the extremum orbits and $B(r)/B_0$ may be evaluated, through Eqs. (31) and (33).

Equations (32), (31), and (33) above for λ , $y(r)$, and $B(r)/B_0$, have been solved by computer programs. Figures 5, 6 and 7 show plots of $B(r)/B_0$ for various values of a/a_0 , with s/a as a parameter, for diamagnetic inner turnaround (DIT). Fig. 8 is a similar plot for paramagnetic inner turnaround (PIT).

Given the self-consistent magnetic field, the self-consistent particle orbits in this field may be determined, either by integration of the solution $y(r) = v_{\perp}(r)/v$, or by computer routines such as the Livermore TIBRO program which integrates the equations of motion of

particles in a specified magnetic field. Figs. 9 and 10 are examples of such orbits, for the extremum orbits for DIT; Fig. 11 illustrates a case for PIT.

Before discussing the general conclusions from this case-study of field reversal, there are some additional results, derived from the above equations, that should be given.

From Eq. (28), the total number of particles/cm associated with a layer of given (a/a_0) and (s/a) may be determined. Since $j_z = K/(v_\theta/r) = m\hbar e f_0 (v_\theta/r)$, it follows that $f_0 = K_0/(\pi e M)$. Thus $n(r) = K_0/(\pi e r)$. Thus we find

$$N = \int_s^a 2\pi n(r) r dr = 2\pi (K_0/e) (a - s). \quad (34)$$

From the relationship between K_0 and λ we then find:

$$N = \frac{1}{2} (Mc^2/e^2) \left[\lambda \left(\frac{a}{a_0} \right)^2 \left(1 - \frac{s}{a} \right) \right] \text{ cm}^{-1} \quad (35)$$

for deuterons this becomes

$$N = 0.65 \times 10^{16} \left[\lambda \left(\frac{a}{a_0} \right)^2 \left(1 - \frac{s}{a} \right) \right] \text{ cm}^{-1} \quad (36)$$

Similarly, for $n(r)$ we have

$$n(r) = \frac{1}{4\pi} (Mc^2/e^2) \lambda \left(\frac{a}{a_0} \right)^2 (1/ra) \text{ cm}^{-3} \quad (37)$$

Solving this relationship for λ and inserting the definition of a_0 we then find

$$\lambda = \frac{\frac{1}{2} n(a) M v^2}{(B_0^2/8\pi)}, \quad (38)$$

That is λ measures the "beta" value of the layer. Since in the last analysis, for this long-layer configuration where self-pinching forces are not operative, the external field must sustain the outward kinetic pressure, we find $\lambda \sim 1.0$ for $(a/a_0) \gg 1$. On the other hand, because of the cylindrical convergence that occurs ($n(r) \sim 1/r$), on the inner surface of the layer the diamagnetic current, having a sheath-like character as can be seen from Fig. 5, is amplified. Thus in principle any desired level of field reversal (or field enhancement)

may be achieved, dependent upon the value of s/a . In the limit $(a/ag) \rightarrow 1$ we find for the field-reversed case:

$$\frac{B(a)}{B_0} = \sqrt{\frac{a}{s}} \quad (39)$$

Note that this implies that

$$\frac{B_z^2(s)}{B_0^2} = \left(\frac{a}{s}\right), \quad (40)$$

i.e. it simply reflects the necessity to satisfy kinetic pressure balance considerations at the inner surface of the layer as well as at the outer surface.

Finally, completely to specify all aspects of the orbital behavior of the particles, we must establish the parameters for particles that have their inner and outer turnaround radii within the layer, rather than at the inner and outer surfaces, as is the case with the extremum orbits. These orbits (those with turnaround radii inside the layer) have non-zero values of v_z at turnaround, that is they move in roughly helical paths within the layer. The relevant parameters may be readily determined, as follows:

We have, in general,

$$p_r^2 + p_z^2 + p_\theta^2 = 2ME_0 \quad (41)$$

At turnaround, $p_r = 0$.

$$p_\theta^2 + p_z^2 = 2ME_0 \quad (42)$$

Also, at $r = a$, $p_z = 0$, $v_\theta = -v$

$$p_{\theta \max}^2 = 2ME_0 \quad (43)$$

since

$$p_\theta = rp_z + (e/2\pi c) \psi(r)$$

$$p_\theta = -a(2ME_0)^{1/2} + \frac{e}{2\pi c} \psi(a) = \text{constant}$$

$$= -r(2ME_0 - p_z^2)^{1/2} + \frac{e}{2-c} \psi(r) ,$$

for turnaround at $r = a$. From this relationship, and calculating $\psi(r)$ by integration of $B(r)$ we find a transcendental equation for $x = (r/a)$, the turnaround radii.

For diamagnetic turnaround at both inner and outer radii this equation is

$$\left(\frac{a}{a_0}\right) \int_x^1 u \left[\frac{B(u)}{B_0} \right] du + x \sin \phi = 1.0 \quad (44)$$

where the pitch angle

$$\phi = \sin^{-1} \left[1 - \frac{v_z^2}{v^2} \right]^{1/2}$$

for the particles at their turnaround points. Thus $\sin \phi = 1.0$ for the extremum orbits. For field-reversing cases the particles have only a limited range of ϕ accessible to them, i.e. there exists a minimum value of $\phi = \phi_{\min}$ for which the inner and outer turnaround radii are coincident. This limiting group of particles thus moves on purely helical paths with pitch angle $\phi_{\min}$. We may determine $\phi_{\min}$, thus determine the allowable range of ϕ for the particle orbits by noting that at $\phi_{\min}$, $d \sin \phi / dr = 0$. Thus from Eq. (44) above we find, by differentiation, a transcendental equation for x at the boundary surface on which $\phi = \phi_{\min}$. This equation is

$$x_0^2 = \left(\frac{a_0}{a}\right) \left[\frac{B_0}{B(x_0)} \right] \left\{ 1 - \left(\frac{a}{a_0}\right) \int_{x_0}^1 u \left[\frac{B(u)}{B_0} \right] du \right\} \quad (45)$$

The above problem may also be solved for PIT orbits also. But in this case the equation from which the inner turnaround radius x_1 is determined is modified to

$$\left(\frac{a}{a_0}\right) \int_{x_1}^1 u \left[\frac{B(u)}{B_0} \right] du - x_1 \sin \phi = 1.0 \quad (46)$$

Also for this case all values of $\sin \phi$ are allowed, and the particle motion on the bounding surface (where $\sin \phi = 0$) is purely in the z direction.

From the above discussion we see that it is possible to generate field reversed equilibria analytically for which all details of the particle motions can be determined. Although the model is idealized, the angular distributions found are not greatly different from ones that might be generated in the laboratory, say by neutral beam injection. There is some reason to believe therefore that insight into more realistic problems may be gained by examining characteristics of the solutions here obtained. Some general results emerge:

- Field Reversal is not possible for entities with $a/a_0 > 2.0$.
- Strong field reversal requires $a/a_0 \gg 2.0$ and $s/a \gg 1.0$ (thick layers of large diameter as measured in vacuum orbit radii).
- The minimum number of ions/cm required to achieve field reversal is of order $3 \times 10^{15} \text{ cm}^{-1}$ (deuterons).
- Depending on the details of how the particles are launched in the layer, either field reversal or field enhancement can be achieved.

The above analytical approach can be extended to the case of an axially-symmetric vacuum field with a radial gradient. This case could model the creation of a long field-reversed equilibrium in a high-order multipole field, a problem of interest in connection with the stability of the layer.

C. The Rigid Rotor

Another class of field reversing plasma configuration that has been studied analytically is the so-called Rigid Rotor. It is another example of an equilibrium state that can be analyzed by the Maxwell-Vlasov equation formulation. The term "Rigid Rotor" is clearly a misnomer when applied to a plasma state. In our context here it means that we are discussing a plasma state where the average angular rotation frequency, ω , of the current carriers of the plasma is independent of

radius, i.e. there is no rotational shear so that $v_\theta = r\omega$ at all radii within the plasma. Some of the rationale behind looking at such states is the fact that, unless maintained by external sources, collisions within a rotating plasma would tend to damp out shear in the rotational velocity, while constancy of angular momentum would tend to preserve any initial rotation of the entity as a whole. As with the previous treatment, the theoretical treatment of rigid rotor equilibria is usually limited to cylindrical symmetry, i.e. to long cylindrical layers where end effects are ignored. Rigid rotor equilibria have been calculated by a number of investigators; in the discussion to follow we will present a simplified treatment, referring the reader to the results of Kapetanakis, Golden and Chu [7] for a more detailed treatment.

As with the previous treatment the starting point for the analysis is the recognition that any function of the constants of the motion, E , P_θ and p_z , is a solution of the Vlasov equation. For the rigid rotor it turns out that functions of the form $F(E - \omega P_\theta) g(p_z)$, where ω is a constant, will generate rigid rotor solutions. We will here choose a distribution function of the form

$$f(E - P_\theta) g(p_z) \quad (47)$$

to illustrate the rigid rotor effect. As before (see Eq. (23)) we have for j_θ

$$j_\theta = \frac{e}{M} \int_{p_z} g(p_z) dp_z \int_{P_\theta} \int_E \left\{ \frac{\left(\frac{P_\theta}{r} - \frac{e}{c} A_\theta \right) (E - P_\theta)}{2ME - \left(\frac{P_\theta}{r} - \frac{e}{c} A_\theta \right)^2 - p_z^2} \right\}^{1/2} dP_\theta dE \quad (49)$$

Upon integration over E this becomes

$$j_\theta = \frac{e}{M} \int_{p_z} g(p_z) dp_z \int_{P_\theta} \left\{ \frac{\left(\frac{P_\theta}{r} - \frac{e}{c} A_\theta \right) dP_\theta}{2MP_\theta - \left(\frac{P_\theta}{r} - \frac{e}{c} A_\theta \right)^2 - p_z^2} \right\}^{1/2} dP_\theta \quad (50)$$

subject to the requirement

$$2MP_\theta - \left(\frac{P_\theta}{r} - \frac{e}{c} A_\theta \right)^2 - p_z^2 \geq 0$$

Letting $\chi = \left(\frac{p}{r} - \frac{e}{c} A_z \right)$ the integral over P_z becomes

$$\int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} x dx \left\{ -\chi^2 + 4\pi q_z - \left[\frac{2\pi}{c} \frac{e}{r} A_z - \frac{p^2}{r^2} \right] \right\}^{1/2} \quad (51)$$

which may be integrated to yield the value $M(r)$. Following Kapetanakis, et al., if we now take $q(p_z) = 0$ (no z velocity) we find, upon normalization,

$$j = n_0 e r, \quad (52)$$

i.e. the azimuthal current associated with the rotation of charge density n_0 at an angular frequency ω .

In the same way the density turns out to be

$$n(r) = n_0 = \text{constant}$$

Thus distributions of the general type chosen lead to rigid rotor behavior. From the above expression for j , and from the imposition of boundary conditions, it is possible to define the inner and outer boundaries of field-reversing layers and to calculate the variation of magnetic field within them. As can be seen by integration of Eq. (52), the variation of $B(r)$ in the present case will be of the form

$$\frac{B(r)}{B_0} = 1 - \gamma \left(\frac{a}{a_0} \right)^2 \left(1 - \frac{r^2}{a^2} \right), \quad r \leq a, \quad (53)$$

where γ measures the strength of the layer, i.e. is proportional to its density and its rotation velocity. Several authors, including Kapetanakis et al. have discussed more general cases, ones in which the layer is surrounded by a flux-conserving shell, as well as layers of limited radial extent, i.e. current shells. Fig. 12, taken from their work illustrates some of the equilibria they have found.

Rigid rotor equilibria thus represent another separate class of field-reversing entity that may model physical situations of interest.

In this connection it should be noted that the first equilibrium discussed, the Hill's Vortex model, is in some sense also a rigid rotor case in that the solution was derived subject to the assumption that $j_z = 0$ within the plasma.

D. Computer Simulation Models

Another successful approach to the generation of field-reversed equilibria has been via the use of computers to simulate the plasma through numerical integration of the motions of a large number of "particles" moving in the self-consistent magnetic fields generated by their motion and by external conductors. We will not discuss this technique in detail but mention only an example, the "Superlayer" code at Livermore, Byers [8]. Although axial symmetry is assumed, and only ionic currents are included, this code otherwise simulates most of the important physical processes going on in a real plasma, including the introduction of energetic ions through neutral injection and the subsequent slowing of these ions by dynamical friction with the plasma electrons.

The advantage of codes such as the Superlayer code over analytical approaches is that they can generally more realistically model the actual plasma situation, thus in some sense represent "experiments" paralleling laboratory experimentation. Their disadvantage is that it is not as readily possible to reach general conclusions as to the physics involved as it is with analytical results.

Studies with Superlayer have been made simulating the buildup of field-reversed states by neutral beam injection in a magnetic field modeling that is the Livermore 2XIIB experiment. It was found possible to create field-reversed states in this case provided the neutral beams were tangentially injected (see Fig. 2) at intensities of order 600 amperes equivalent or higher. We will later show a comparison between 2XIIB experimental results (where field-reversal was approached, but not achieved) with Superlayer.

Fig. 13 shows an example of a field reversed state found by Superlayer.

D. Electron Currents

In the last three examples, where field reversal has been assumed

to be created solely by ionic diamagnetic currents, a tacit assumption has been made that may not correspond to what actually will occur. The assumption is that any electron currents carried by the electrons of the plasma are small compared to the ionic terms. This assumption might not be valid, even when the electron pressure is small compared to the ion current. The issue here is the following: While it is probably true that in regions of the field reversed entity where $B \neq 0$ the electrons, being trapped on the magnetic lines, cannot significantly modify the ion diamagnetic current, this assumption may not be valid in regions where $B = 0$. In any finite- ℓ high- ℓ reversed field configuration without an entrapped toroidal field there will be a closed curve in space where $B = 0$. For example in the Hill's Vortex solution this is a circle centered at $z = 0$ with relative radius, $r/a = 1/\sqrt{2}$. But at such a null the electrons are not inhibited from "catching up" with the ions, owing to collisional drag on the ions, thereby locally cancelling the ion current. Some have conjectured that this effect could propagate beyond the initial null line, eventually destroying the ion diamagnetism. Baldwin and Fowler (9) have considered the effect, including the ionization effect of higher ℓ impurities (the so-called "Okawa current"), concluding that partial current cancellation by the electrons could increase the reversed β/β_0 of a field-reversed entity by of order a factor of 10. At this juncture, however, there is neither sufficient theoretical work nor corroborative experimental data to confirm or deny the electron current cancellation phenomenon in this context.

III. STABILITY OF FIELD-REVERSED STATES

There is no doubt that field-reversed states in pressure equilibrium with a confining magnetic field exist. The central physics issue then becomes the question of the stability of such states. Unfortunately, owing to the difficulty of analyzing such questions in a rigorous manner, there is as yet no sufficiently comprehensive treatment of the stability issue. Treatments addressing specific instability modes have, however, been given.

Of primary concern are the MHD modes, those which lead to catastrophically rapid gross motions of the field-reversed entity. We may

make a rough division of the MHD modes into two categories, is appropriate to the limiting case of equilibrium which they include. The first category, mentioned earlier, is the kinking or pinching instability of a thin bicycle-tire equilibrium, one in which the pinching forces are dominant and instability occurs in the form of a magnetically-driven kinking or pinching off of the column. At the other extreme is the long layer, where in equilibrium axial pinching forces are weak or absent. Here the modes are either flute-like in nature, i.e. gross transverse displacements, or "tearing" modes, in which the column tends to break up into rings. Marx [6] has analyzed the stability of an infinitely long layer against breaking up into rings, finding that this will not occur if the layer is surrounded by a close-fitting conducting shell. On the other hand, Fowler [10] has given a proof that for realistic, i.e. finite length, equilibria the balance between axial pinching forces and longitudinal kinetic pressure is such as to prevent the tearing mode. It seems likely therefore that this kind of mode - i.e. breakup of a field-reversed layer into rings - can be avoided, and that the main concern is with link-like MHD modes.

One general approach to the stabilization of a field-reversed entity against kinking is to fabricate it so that its major circumference is large compared to its minor circumference. This excellent places strong geometrical constraints on the wavelength of the kinking mode and certainly promotes stability. No definitive treatment of this question has as yet appeared, but Sudan and Rosenbluth [11] have developed a theoretical treatment based on an energy-principle approach that gives a sufficient condition for stability. Their work strongly suggests that flat field-reversed states stable against MHD kinking modes can be found.

Another factor which should tend to promote the stability of field-reversed equilibria derives from finite-orbit effects. As we have seen, field reversed entities should exist in which energetic ions may execute complex orbits traversing the entire field-reversed region. Analogous to the case of the finite-orbit stabilization of mirror systems, F-O effects should help stabilize field-reversed systems which are a few gyroradii in dimension. Because of the potentially high beta values of such states useful fusion power could be derived from field-reversed entities only 5 or so orbit radii in dimensions ($a/a_0 = 5.0$

in the previous notation). At this value E-G effects might be quite important.

1.1. EXPERIMENTAL APPROACHES

Field-reversed plasma configurations have now been demonstrated by a number of techniques. These experiments have shown clearly that the concept of high-beta field-reversing plasmas is not merely a theoretical construct. It would appear that, in the context of fusion power research, the task from here on will be either that of developing configurations exceeding conventional magnetic confinement systems, or that it will be a search for configurations that are both stable and feasible to create and maintain by practical techniques of plasma generation and heating. In the discussion to follow we will not attempt to cover the experimental work in field reversal in detail but will instead mention only highlights.

A. Relativistic Beams

One of the earliest approaches to field reversal was the ASIPPON of Christofilos, in which a laser or electron-beam relativistic energies was to produce field reversal. Although the original ASIPPON device at Livermore turned out to be not as effective, the basic idea embodied in it has been employed at a number of places elsewhere. In the work of Davis, Maguire and Smith (see [12]) for example, a high-intensity burst of electrons is captured in a magnetic field produced by millisecond-long field reversal. Both axially symmetric and quadrupolar magnetic well geometries have been tested. Stabilization of the rings is accomplished by creating them in a plasma background and using either a close-fitting conducting sleeve or the quadrupolar field configuration to suppress low frequency instabilities. In recent experiments the Connell Group [13] has demonstrated the possibility of translating and compressing field-reversed electron rings, magnetically moving them into regions of higher vacuum and higher magnetic field. This results in longer lifetime against scattering losses arising from collisions with the background plasma. In these latter experiments field reversal persisted for over 1 millisecond, a time far longer than the characteristic growth time of MHD-type instabilities.

Another kind of field-reversed entity has also been created using intense relativistic electron beams (REB). In the work of Cetinian et al. at NRL [14] a 1 MeV 50 kA beam of 60 ns duration is fired axially through a cusp field to induce rotation. This rotating electron burst, while passing through a chamber filled with gas, induces field-reversing currents in the plasma created by the passage of the beam. These currents persist for several microseconds after the electron burst has left through the far end of the tube. The observed decay time, 5 microseconds, is consistent with classical diffusion rates at the density ($5 \times 10^{15} \text{ cm}^{-3}$) and temperature (12 eV) of the field-reversed plasma. The investigators believe that the REB technique can be extended to the creation of hotter, longer lived, field-reversed plasma states.

B. Reversed Field Theta Pinch

Another approach, one which seems to have been discovered by accident in the early days of fusion research, is the so-called Reversed Field Theta Pinch (RFTP). In the conventional theta pinch a plasma embedded in an initial magnetic field is created by shock ionization and heating of a gas by a rapidly rising solenoidal magnetic field. If this magnetic field is then rapidly reversed in direction the trapped flux inside the plasma is now oppositely directed to the applied field. After reconnection of the magnetic lines at the ends there results a field-reversed configuration, as shown schematically in Fig. 14 taken from the work of Linford [15]. Depending on the temperature and the purity of the plasma this entity may last for 30 microseconds or more, at which point it is terminated by the rapid growth of an $m = 2$ rotational instability. While it exists the field reversed configuration reduces the rate of end loss of the plasma by more than a factor of 10, relative to the loss rate for a conventional theta pinch plasma configuration. There is evidence that the onset of the rotational instability is influenced by impurities and that it may therefore be amenable to being further delayed or even suppressed by technical improvements. Whereas the REB reversed field entities of the ASTRON type consist of large-orbit particles, the RFTP configuration seems to be created solely by diamagnetic currents in a plasma where $a/a_0 \gg 1$. Thus the two techniques appear to model the two extremes

we have previously discussed.

C. Neutral Beam Injection

While field reversal has probably not yet been achieved through this technique, a different and important approach to the problem is the Field Reversed Mirror (FRM) achieved by the tangential injection of neutral beams (see Fig. 2). In experiments of this type carried out in the 2XII-B facility at Livermore, Simonen et al. [25] states in which $B/B_0 = 1.0$, i.e., where the diamagnetic field nearly canceled the confining field were achieved, a reversal of the neutral beam current (see Fig. 3) shows measured B/B_0 values as a function of neutral beam current, taken with a vacuum magnetic field of 1.35 kG gauss, as compared with predictions of the Sub-layer model. On the plot $B/B_0 = 1.0$ represents the threshold for field reversal. At the present time it is not known whether the $B/B_0 = 1.0$ value seen in Fig. 3 is a result of technical limitations or whether it represents effects associated with the onset of field reversal. Thus, the new ideas taken to ameliorate the technical limitations through increasing the neutral beam current and through the introduction of neutral beam startup penning lamps (plasma guns) that may permit the achievement of field reversal, using the neutral beam technique, are still in the stage of laboratory applications in fusion, the existing FRM being a 100 m long, low power, compact plasma confined experiment, being modified to a source of fusion power could be the result.

This document contains information of a preliminary nature and is not to be distributed outside the Laboratory without the approval of the U.S. Department of Energy to the extent that it covers material to be classified.

NOTES

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, method, or process disclosed, or represents that its use would not infringe privately owned rights.

References

- [1] GRAD, H., RUBIN, H., "Hydromagnetic Equilibria and Force-free Fields," Proc. U.N. Intern. Conf. Peaceful Uses At. Energy, 2nd, Geneva, 1958 (United Nations, Geneva, 1958), vol. 31, p. 190.
- [2] SHAFRANOV, V.V., Reviews of Plasma Physics, vol. 2, pp. 112-117, Consultants Bureau, N.Y. (1966), edited by M.A. Leontovich.
- [3] HILL, M.J., Phil. Trans. Roy. Soc. A, Pt. 1, 213 (1994).
- [4] BUSSAC, M.N., FURTH, H.P., et al., paper CN-37-X-1, "Low-Aspect-Ratio Limit of the Toroidal Reactor: The Spheromak," presented at the Seventh IAEA Conf. on Plasma Physics and Controlled Fusion Research, Innsbruck, Austria, Aug. 23-30, 1978.
- [5] POST, R.F., "Field-Reversed Equilibria Created by Neutral Beam Injection - A Theoretical Model," A.P.S. Bull. 22, 1192 (1977).
- [6] MARX, K.O., Phys. Fluids 11, 357 (1968).
- [7] KAPETANAKOS, C.A., GOLDEN, J., CHU, K.R., Plasma Physics 19, 387 (1976).
- [8] BYERS, J.A., Phys. Rev. Lett. 39, 1476 (1977).
- [9] BALDWIN, D.E., FOWLER, T.K. (unpublished).
- [10] FOWLER, T.K., Phys. Fluids 15, 2075 (1972).
- [11] SUDAN, R.N., ROSENBLUTH, M.N., Cornell Univ. Rpt. LPS 230, March 1978 (unpublished).
- [12] DAVIS, H.A., MEGER, R.A., FLEISCHMANN, H.H., Phys. Rev. Lett. 37, 542 (1976).
- [13] TUSZEWSKI, M., et al., paper IAEA-CN-37-5-4 at Seventh IAEA Conf. on Plasma Physics and Controlled Fusion, Innsbruck, Austria, Aug. 23-30, 1978.
- [14] SETHIAN, J.D., et al., Phys. Fluids 21, 1227 (1978).
- [15] LINFORD, R.K., et al., paper CN-S-1-1 at Seventh IAEA Conf. on Plasma Physics and Controlled Fusion, Innsbruck, Austria, Aug. 23-30, 1978.
- [16] SIMONEN, T.C., et al., paper CN-37-J-1, "2XIIB Plasma Confinement Experiments," presented at the Seventh IAEA Conf. on Plasma Physics and Controlled Fusion Research, Innsbruck, Austria, Aug. 23-30, 1978.

Figure Captions

- Fig. 1 Schematic illustration of kinking and pinch off instability modes.
- Fig. 2 Schematic of field-reversed mirror configuration embedded in magnetic well mirror field.
- Fig. 3 Plot of field lines in spherical Hill's Vortex field-reversed configuration.
- Fig. 4 Plots of B_z/B_0 , B_z^2/B_0^2 and β_0 (evaluated at $z = 0$) as a function of r/a in spherical Hill's vortex field-reversed configuration. (Figure courtesy G. Miley, Univ. of Ill.)
- Fig. 5 Plot of B_z/B_0 for long-layer field-reversed (DIT) configuration; $a/a_0 = 10.0$, s/a as parameter; vacuum orbit (circle) shown for scale.
- Fig. 6 Plot of B_z/B_0 for long-layer field-reversed (DIT) configuration; $a/a_0 = 3.0$, s/a as parameter; vacuum orbit (circle) shown for scale.
- Fig. 7 Plot of B_z/B_0 for long-layer high beta (DIT) configuration; $a/a_0 = 2.0$, s/a as parameter; vacuum orbit (circle) shown for scale.
- Fig. 8 Plot of B_z/B_0 for long-layer field-enhanced (PIT) configuration; $a/a_0 = 10.0$, s/a as parameter; vacuum orbit (circle) shown for scale.
- Fig. 9 Plot of extremum orbit in long-layer field-reversed equilibrium; $a/a_0 = 10.0$, $s/a = 0.2$; vacuum orbit (circle) shown for scale.
- Fig. 10 Plot of extremum orbit in long-layer field-reversed equilibrium; $a/a_0 = 5.0$, $s/a = 0.4$; vacuum orbit (circle) shown for scale.
- Fig. 11 Plot of extremum orbit in long-layer field-enhanced equilibrium; $a/a_0 = 10.0$, $s/a = 0.2$; vacuum orbit (circle) shown for scale.
- Fig. 12 Examples of B_z/B_0 for rigid-rotor field-reversed layers, (Figure taken from Kapatzenakos, et al. (Ref. 7)).
- Fig. 13 Example of equilibrium field configuration generated by Superlayer code.

Fig. 14 Schematic of reversed-field theta pinch configuration.

Fig. 15 Comparison between 2X1B data and Superlayer code predictions; $\Delta B/B_0$ at $B_0 = 4.35$ kilogauss.

Figure 1: Schematic illustration of kinking and pinch off instability modes.

FIELD REVERSED MIRROR

Figure 1: Schematic of field-reversed mirror configuration embedded in magnetic well mirror field.

Figure 3: Plot of field lines in spherical Hill's Vortex field-reversed configuration.

Figure 4: Plots of B_z/B_0 , B_z^2/B_0^2 and β_0 (evaluated at $z = 0$) as a function of r/a in spherical Hill's vortex field-reversed configuration. (Figure courtesy G. Miley, Univ. of Ill.)

Figure 5: Plot of B_z/B_0 for long-layer field-reversed (DIT) configuration; $a/a_0 = 10.0$, s/a as parameter; vacuum orbit (circle) shown for scale.

Figure 6: Plot of B_z/B_0 for long-layer field-reversed (DIT) configuration; $a/a_0 = 3.0$, s/a as parameter; vacuum orbit (circle) shown for scale.

Figure 7: Plot of B_z/B_0 for long-layer high beta (DIT) configuration; $a/a_0 = 2.0$, s/a as parameter; vacuum orbit (circle) shown for scale.

Figure 8: Plot of B_z/B_0 for long-layer field-enhanced (PIT) configuration; $a/a_0 = 10.0$, s/a as parameter; vacuum orbit (circle) shown for scale.

Figure 9: Plot of extremum orbit in long-layer field-reversed equilibrium; $a/a_0 = 10.0$, $s/a = 0.2$; vacuum orbit (circle) shown for scale.

Figure 10: Plot of extremum orbit in long-layer field-reversed equilibrium; $a/a_0 = 5.0$, $s/a = 0.4$; vacuum orbit (circle) shown for scale.

$$\begin{aligned} a/a_0 &= 10.0 \\ s/a &= 0.2 \end{aligned}$$

Figure 11: Plot of extremum orbit in long-layer field-reversed equilibrium; $a/a_0 = 10.0$, $s/a = 0.2$; vacuum orbit (circle) shown for scale.

Figure 12: Examples of B_z/B_0 for rigid-rotor field-reversed layers, (Figure taken from Kapatnakos, et al. (Ref. 7)).

Figure 13: Example of equilibrium field configuration generated by Superlayer code.

Figure 14: Schematic of reversed-field theta pinch configuration.

Figure 15: Comparison between 2XIIB data and Superlayer code predictions; $\Delta B/B_0$ at $B_0 = 4.35$ kilogauss.