05
9/1/84
PPPL-2116

UC20-F

PPPL-2116

DR# 0258-5

①
I-16207

NOTICE
PORTIONS OF THIS REPORT ARE ILLEGIBLE. It
has been reproduced from the best available
copy to permit the broadest possible avail-
ability.

HARMONIC GENERATION AND PARAMETRIC DECAY
IN THE ION CYCLOTRON FREQUENCY RANGE

By

F.N. Skiff, K.L. Wong, and M. Ono

JUNE 1984

PLASMA
PHYSICS
LABORATORY

PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America.

Available from:

National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151

Price: Printed Copy \$ ____; Microfiche \$3.50

<u>*PAGES</u>	<u>NTIS</u> <u>Selling Price</u>
1-25	\$5.00
26-50	\$6.50
51-75	\$8.00
76-100	\$9.50
101-125	\$11.00
126-150	\$12.50
151-175	\$14.00
176-200	\$15.50
201-225	\$17.00
226-250	\$18.50
251-275	\$20.00
276-300	\$21.50
301-325	\$23.00
326-350	\$24.50
351-375	\$26.00
376-400	\$27.50
401-425	\$29.00
426-450	\$30.50
451-475	\$32.00
476-500	\$33.50
500-525	\$35.00
526-550	\$36.50
551-575	\$38.00
576-600	\$39.50

For documents over 600
pages, add \$1.50 for each
additional 25 page increment.

HARMONIC GENERATION AND PARAMETRIC DECAY
IN THE ION CYCLOTRON FREQUENCY RANGE

PPPL--2116
DEG4 015147

F. N. Skiff, K. L. Wong, and M. Ono
Plasma Physics Laboratory, Princeton University
Princeton, New Jersey 08544

ABSTRACT

Harmonic generation and parametric decay are examined in a toroidal ACT-I plasma using electrostatic plate antennas. The harmonic generation, which is consistent with sheath rectification, is sufficiently strong that the nonlinearly generated harmonic modes themselves decay parametrically. Resonant and nonresonant parametric decay of the second harmonic are observed and compared with uniform pump theory. Resonant decay of lower hybrid waves into lower hybrid waves and slow ion cyclotron waves is seen for the first time. Surprisingly, the decay processes are nonlinearly saturated, indicating absolute instability.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

The successful application of high power rf heating schemes generally requires an understanding of nonlinear mechanisms which come into play in plasmas subjected to strong electric fields. Fortunately many nonlinear processes can be studied on small-scale devices where measurements are more easily performed. In this letter we report observations of harmonic generation and subsequent parametric decay of the second harmonic at low power levels in the low density toroidal ACT-1 plasma using electrostatic plate antennas. Resonant decay of lower hybrid waves into lower hybrid waves and slow ion cyclotron (electron-ion Buchsbaum) waves is seen for the first time.

The harmonic generation is consistent with plasma sheath rectification at the antenna, and uniform pump theory provides accurate predictions of the unstable frequencies for parametric decay. The saturated amplitude of the decay, however, does not scale exponentially with the pump power, but linearly, even at pump levels close to threshold. This scaling indicates a nonlinear saturation mechanism rather than convective saturation of the decay.

The plasma used in this experiment was produced in the ACT-1 device using a biased hot filament source in the toroidal magnetic field which is adjustable up to 4.2 kG. Plasma temperatures of $T_e = T_i = 1$ eV, density $n = 0.5 - 1 \times 10^{10}$, and low collisionality $\nu_i/\Omega_{ci} = 10^{-3}$ were typical. Two curved metal plates, separated toroidally by 1 port (16.7 cm) and driven in antiphase through a matching network, served as an antenna. The driver frequency ω_D was chosen between $1 - 2 \times \Omega_{ci}$ which, for the slow wave and $\omega_{pi} \approx 2 \times \Omega_{ci}$, corresponds to a second harmonic ion Bernstein wave. For the harmonics, however, $n\omega_D > \omega_{LH} = (\omega_{pi}^2 + \Omega_{ci}^2)^{1/2}$ so that these frequencies correspond to lower hybrid waves. Edge regions of laboratory plasmas are in this regime. A schematic of the experimental setup is shown in Fig. 1(a).

To model the coupling of an electrostatic antenna one may start with the equation for the potential Φ

$$\nabla \cdot \hat{\epsilon} \cdot \nabla \Phi = 4\pi \rho_{\text{free}},$$

where $\hat{\epsilon}$ is the hot uniform-plasma dielectric tensor.² ρ_{free} refers to electric space charge not included in the uniform plasma model, in particular, to the charge on the antenna conductors and on the surface of the plasma sheath. Because of the behavior of the sheath, both the displacement and conduction currents from the antenna have a nonlinear component, the former due to nonlinear sheath capacitance³ and the latter to sheath rectification.⁴ Because sheath rectification is dominant, the current-voltage characteristic of the antenna can be estimated by

$$i = \frac{n_0 v_t A}{4} \times e^{\frac{eV_a}{kT_e}},$$

where i is the antenna current, n_0 is the density in the sheath, v_t is the electron thermal velocity corresponding to the electron temperature T_e , and A is the effective plate area. (An accurate determination of A requires consideration of the effect of the magnetic field.) With antenna voltage $V_a = V_0 \times e^{i\omega t}$ the amplitude of the n^{th} harmonic in the current is

$$\frac{I_n}{I_0} = I_{n-1} \left(\frac{eV_0}{kT_e} \right) / I_0 \left(\frac{eV_0}{kT_e} \right) .$$

I_m denotes the modified Bessel function of order m . Figure 1(b) shows the observed harmonic amplitude ratios A_2/A_1 , A_3/A_1 , and A_4/A_1 (A_1 being the driver amplitude, etc.) as a function of eV_a/kT_e , the ratio of antenna voltage to plasma electron temperature. Shown for comparison are theoretical values of the above ratios, based on sheath rectification and evaluating the single parameter, eV_a/kT_e , for 1 W of antenna power. Observation of the resonance cones of the harmonics using radial scanning rf probes confirm that the harmonics are being generated near the antenna. It is also evident from the presence of resonance cones and the magnitude of the cone angles, given by

$$\tan^2 \theta = \frac{\omega_{pe}^2}{\omega^2} \times \left(\frac{\omega^2 - \omega_{ci}^2}{\omega^2 - \omega_{LH}^2} \right)$$

that the harmonics are propagating as lower hybrid waves.

As the density is increased, the amount of harmonic generation decreases. Simultaneous changes in the linear loading make the density effect hard to quantify, but it is consistent with our simple model which predicts that at constant antenna power the sheath will exhibit less nonlinearity at higher densities. In principal, the nonlinearity can also be reduced by increasing T_e and/or by reducing A .

While harmonic generation is confined to the sheath region, the spectrum is additionally enriched by nonlinear wave-wave and wave-particle interactions in the plasma bulk. In particular, in this experiment we observed both resonant and nonresonant parametric decay of the second harmonic. The growth rate for these decays is given by

$$\gamma = \frac{-\text{Im}(\epsilon) + \Gamma_{\sigma, \eta} \frac{1}{B} |\mu_{\sigma}^* - \mu_{\eta}^*| \text{Im}(\frac{\chi_{\sigma} \chi_{\eta}}{\epsilon})_{\omega}}{|\partial \epsilon / \partial \omega_-|}, \quad (1)$$

$$\text{with } \mu_{\sigma} = \left(\frac{q}{m}\right)_{\sigma} \left[\left(\frac{E_{0\parallel} k_{\parallel}}{\omega_0^2} + \frac{E_{0\perp} \cdot \hat{k}_{\perp}}{\omega_0^2 - \Omega_{\sigma}^2} \right)^2 + \frac{|E_{0\perp} \times \hat{k}_{\perp}|^2 \Omega_{\sigma}^2}{(\omega_0^2 - \Omega_{\sigma}^2)^2 \omega_0^2} \right]^{1/2}.$$

$\epsilon = 1 + \chi_e + \chi_i = \hat{k} \cdot \hat{\epsilon} \cdot \hat{k}$ is the scalar projection of the dielectric tensor, ω_0 , ω_- , ω are the angular frequencies of the pump, lower sideband and low frequency wave or quasimode, respectively, and Ω_{σ} is the cyclotron frequency of species σ .

Two distinct decay processes were observed involving the second harmonic as pump. Figure 2(a) shows the complete spectrum including driver (F_0), second harmonic ($2F_0$), upper and lower sidebands (USB_1 , USB_2 , LSB_1 , LSB_2), and corresponding low frequency excitations (QM_1 , LF_2). One process (subscript 1) is the well-known decay of a lower hybrid wave into a lower hybrid wave and an ion cyclotron quasimode.⁷ The other process (2) (not seen previously) is the resonant decay of a lower hybrid wave into a lower hybrid wave and an ion cyclotron wave (sometimes called a Buchsbaum mode)⁸ with $\omega < \Omega_{ci}$ and

$$\frac{k_{\perp}}{k_{\parallel}} = \frac{\omega_{pe}}{\omega} \times \left(\frac{\omega^2 - \Omega_{ci}^2}{\omega^2 - \omega_{LH}^2} \right)^{1/2}.$$

The two decay processes can be readily distinguished by the frequencies of the decay waves. Data plotted in Fig. 2(b) pertain to both decays as observed over a range of toroidal field strengths. Observed decay wave frequencies are compared with the frequencies determined by Eq. (1) to be the most unstable. It is found that the quasimode frequency is always close to f_{c1} and that the frequency of the ion cyclotron wave is strongly constrained by the matching condition $k_0 = k_- + k$.

Observation of the decay wave amplitudes also provides information concerning the nature of the decay process. Resonant decays are constrained by the Manley-Rowe relations $N_0 + N_- = \text{const.}$ and $N_0 + N = \text{const.}$ where N_0 refers to the wave "action" of the pump (etc.).⁹ Wave amplitude is proportional to wave action. In Fig. 2(c) the ratios $A_{\text{LSB2}}/A_{\text{LF2}}$, $A_{\text{LSB1}}/A_{\text{QM1}}$, $A_{\text{LSB2}}/A_{2F\phi}$, and $A_{\text{LSB1}}/A_{2F\phi}$ are plotted as functions of the squared amplitude of the second harmonic. For the resonant decay (subscript 2), the ratio of the decay wave amplitudes is a constant, in agreement with Manley-Rowe, since $N_- = N = 0$ and $N_0 = N_0^i$ at $t=0$ gives for $t>0$

$$A_{\text{LF}}/A_{\text{LSB}} \propto \frac{N}{N_-} = \frac{(N_0^i - N_0)}{(N_0^i - N_0)} = 1$$

For the nonresonant decay this scaling does not hold and the decay wave amplitudes are determined by the decay saturation mechanism.¹⁰ Above threshold the observed sideband amplitudes are proportional to the pump amplitude $A_{2F\phi}$ [Fig. 2(c)] indicating nonlinear saturation for both decay types. This behavior is surprising in view of the fact that parametric decay of lower hybrid waves is expected to be convective.¹¹ The same amplitude scaling has, however, been seen on an earlier experiment.¹¹ One possible

explanation is that the decay is made absolute by reflective trapping of the decay waves. (In contrast to the earlier experiments, ACT-1 has no ends.) The finite spatial extent of the pump can accomplish reflective trapping through the effect of the pump on the decay wave wave numbers.^{12,13} Treating the pump as a square potential "well," a rough estimate of the threshold for absolute instability is:

$$|r_1 r_2 e^{rL}| > 1 ,$$

where r_1 and r_2 are the (unequal) well wall reflectivities, r is the spatial growth rate $r = \gamma/v_{\text{group}}$, and L is the pump cone width as transited by the decay wave. For absolute instability the group velocities of the decay waves must be oppositely directed. The threshold condition reduces for our experiment to:

$$r_1 r_2 e^{rL} \sim \frac{\gamma^2}{8\omega\omega_-} e^{2\pi \gamma/\omega} > 1.$$

This threshold is very close to the convective threshold and very similar in scaling. For lower hybrid waves at higher frequency it is interesting to note that one of the reflections (nearly tangent) becomes strong and the other weak thereby reducing the convective threshold by a factor of two. Experimentally it is often the case (when there are resonance cones) that the observed

thresholds are lower than convective estimates.¹¹ Further work is required to resolve this issue.

In summary, we have observed substantial harmonic generation associated with the use of an electrostatic antenna in the ion cyclotron frequency range. These harmonics have sufficient intensity to decay parametrically through both resonant and nonresonant channels. The decay wave amplitude scaling indicates absolute rather than convective instability.

ACKNOWLEDGMENTS

We thank G. A. Wurden for first noticing the spectrum, Liu Chen for helpful discussions, and J. Taylor and W. Kineyko for their technical assistance.

This work has been supported by the U.S. Department of Energy Contract No. DE-AC02-76-CH0-3073.

F. Skiff is supported by a Fannie and John Hertz Foundation Fellowship.

REFERENCES

- 1 M. Ono, K. L. Wong, and G. A. Wurden, Phys. Fluids 26, 298 (1983).
- 2 T. H. Stix, Theory of Plasma Waves (McGraw Hill, New York, 1962).
- 3 A. J. Cohen and G. Bekefi, Phys. Fluids 14, 1512 (1971).
- 4 S. J. Crawford and R. F. Meadnosky, J. Geophys. Res. 69, 2765 (1964).
- 5 J. L. Hirshfield and I. B. Bernstein, Phys. Fluids 18, 1787 (1975).
- 6 Khorasanov et al., Sov. Tech. Phys. Lett. 3(11), 470 (1977).
- 7 K. L. Wong and M. Ono, Phys. Rev. Lett. 47, 842 (1981).
- 8 S. J. Buchsbaum, Phys. Fluids 3, 418 (1960).
- 9 A. A. Galeev and R. S. Sagdeev, "Nonlinear Plasma Theory" in Reviews of Plasma Physics, 7 (Plenum, New York, 1979) 16.
- 10 K. Nishikawa and C. S. Liu, in Advances in Plasma Physics, Vol. 6, (John Wiley, New York, 1976) 67.
- 11 K. L. Wong, J. R. Wilson, and M. Porkolab, Phys. Fluids 23, 96 (1980).
- 12 D. F. Dubois, D. W. Forslund, and E. A. Williams, Phys. Rev. Lett. 33, 1013 (1974).
- 13 G. Picard and T. W. Johnston, Phys. Rev. Lett. 51, 574 (1983).

FIGURE CAPTIONS

1. Observations of Harmonic Generation

- (a) Experimental setup on ACT-I with electrostatic launchers.
- (b) Solid line harmonic ratios from sheath rectification. Points are amplitudes from the spectrum analyzer.

2. Observations of Parametric Decay

- (a) Parametric decay spectrum showing driver pump (second harmonic), and resonant(2) and nonresonant(1) decay products.
- (b) Decay product frequencies as a function of magnetic field intensity. Square points are calculated from uniform pump theory. Solid curves are based on data (x marks).
- (c) Decay product amplitude ratios vs antenna power.

Fig. 1

Fig. 2

EXTERNAL DISTRIBUTION IN ADDITION TO TIC UC-20

Plasma Res Lab, Austr Nat'l Univ, AUSTRALIA
 Dr. Frank J. Paoloni, Univ of Wollongong, AUSTRALIA
 Prof. I.R. Jones, Flinders Univ., AUSTRALIA
 Prof. M.H. Brennan, Univ Sydney, AUSTRALIA
 Prof. F. Cap, Inst Theo Phys, AUSTRIA
 Prof. Frank Verheest, Inst theoretische, BELGIUM
 Dr. D. Palumbo, Dg XII Fusion Prog, BELGIUM
 Ecole Royale Militaire, Lab de Phys Plasmas, BELGIUM
 Dr. P.H. Sekanaka, Univ Estadual, BRAZIL
 Dr. C.R. James, Univ of Alberta, CANADA
 Prof. J. Teichmann, Univ of Montreal, CANADA
 Dr. H.M. Skarsgard, Univ of Saskatchewan, CANADA
 Prof. S.R. Greenivasan, University of Calgary, CANADA
 Prof. Tudor W. Johnston, INRS-Energie, CANADA
 Dr. Hannes Bernard, Univ British Columbia, CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Zhengwu Li, SN Inst Physics, CHINA
 Library, Tsing Hua University, CHINA
 Librarian, Institute of Physics, CHINA
 Inst Plasma Phys, Academia Sinica, CHINA
 Dr. Peter Lukac, Komenského Univ, CZECHOSLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Prof. Schatzman, Observatoire de Nice, FRANCE
 J. Raetz, CEH-BP6, FRANCE
 AM Dupes Library, AM Dupes Library, FRANCE
 Dr. Tom Musil, Academy Bibliographic, HONG KONG
 Preprint Library, Cent Res Inst Phys, HUNGARY
 Dr. S.K. Trehan, Panjab University, INDIA
 Dr. Indra, Mohan Lal Das, Banaras Hindu Univ, INDIA
 Dr. L.K. Chavda, South Gujarat Univ, INDIA
 Dr. R.K. Chhajlani, Var Ruchi Marg, INDIA
 P. Kow, Physical Research Lab, INDIA
 Dr. Phillip Rosenau, Israel Inst Tech, ISRAEL
 Prof. S. Cuperman, Tel Aviv University, ISRAEL
 Prof. G. Rostagni, Univ DI Padova, ITALY
 Librarian, Int'l Ctr Theo Phys, ITALY
 Miss Clella De Palo, Assoc EURATOM-CNEN, ITALY
 Biblioteca, del CNR EURATOM, ITALY
 Dr. H. Yameto, Toshiba Res & Dev, JAPAN
 Prof. M. Yoshikawa, JAERI, Tokai Res Est, JAPAN
 Prof. T. Uchida, University of Tokyo, JAPAN
 Research Info Center, Nagoya University, JAPAN
 Prof. Kyoji Nishikawa, Univ of Hiroshima, JAPAN
 Prof. Sigoru Mori, JAERI, JAPAN
 Library, Kyoto University, JAPAN
 Prof. Ichiro Kawakami, Nihon Univ, JAPAN
 Prof. Satoshi Itoh, Kyushu University, JAPAN
 Tech Info Division, Korea Atomic Energy, KOREA
 Dr. R. England, Ciudad Universitaria, MEXICO
 Bibliotheek, Fom-Inst Voor Plasma, NETHERLANDS
 Prof. B.S. Lilley, University of Waikato, NEW ZEALAND
 Dr. Suresh C. Sharma, Univ of Calabar, NIGERIA
 Prof. J.A.C. Cabral, Inst Superior Tech, PORTUGAL
 Dr. Octavian Petrus, ALI Cluza University, ROMANIA
 Prof. M.A. Hellberg, University of Natal, SO AFRICA
 Dr. Johan de Villiers, Atomic Energy Bd, SO AFRICA
 Fusion Div, Library, JEN, SPAIN
 Prof. Hans Wilhelmson, Chalmers Univ Tech, SWEDEN
 Dr. Lennart Stenflo, University of UMEA, SWEDEN
 Library, Royal Inst Tech, SWEDEN
 Dr. Erik T. Karlson, Uppsala Universitet, SWEDEN
 Centre de Recherches, Ecole Polytech Fed, SWITZERLAND
 Dr. W.L. Weise, Nat'l Bur Stand, USA
 Dr. W.M. Stacey, Georg Inst Tech, USA
 Dr. S.T. Wu, Univ Alabama, USA
 Prof. Norman L. Oleson, Univ S Florida, USA
 Dr. Benjamin Ma, Iowa State Univ, USA
 Prof. Magne Kristiansen, Texas Tech Univ, USA
 Dr. Raymond Askew, Auburn Univ, USA
 Dr. V.T. Tolok, Kharkov Phys Tech Ins, USSR
 Dr. D.D. Ryutov, Siberian Acad Sci, USSR
 Dr. G.A. Eliseev, Kurchatov Institute, USSR
 Dr. V.A. Glukhikh, Inst Electro-Physical, USSR
 Institute Gen. Physics, USSR
 Prof. T.J. Boyd, Univ College N Wales, WALES
 Dr. K. Schindler, Ruhr Universitat, W. GERMANY
 Nuclear Res Estab, Julich Ltd, W. GERMANY
 Librarian, Max-Planck Institut, W. GERMANY
 Dr. H.J. Kaeppeler, University Stuttgart, W. GERMANY
 Bibliothek, Inst Plasmaforschung, W. GERMANY