006006

1 OF 1

Technical Report

DE93 006006

June 1, 1992 through August 31, 1992

Contract No.: DE FG-22-91PC-91289
Project Title: Characterization of the Organic-Sulfur-Degrading Enzymes
Principal Investigator: Nancy W.Y. Ho
LORRE
Purdue University
1295 Potter Building
West Lafayette, IN 47907-1295

OBJECTIVES

Our immediate objective of this project is to characterize and purify the enzymes involved in degrading organic sulfur in coal from two well characterized organic sulfur-degrading strains, IGTS8 and K3B. The purification of these enzymes will facilitate the analysis of the structural organization of the genes encoding these enzymes. The characterization of these enzymes will be able to determine the true value of these enzymes towards the development of a practical process for coal desulfurization. Our long-term goal for this project is to overexpress the genes encoding these enzymes in their original host as well as in other microorganisms that can grow very well under the conditions suitable for coal desulfurization.

RESEARCH PERFORMED

We have obtained a lot of exciting results this quarter. First of all, we have carried out electron microscope ultrastructural analysis of K3B cells. Our results showed that K3B cells cultured in DBT sulfone did not contain the similar protuberant structures that

MASTER

appeared on the cell wall of IGTS8 cells when the latter cells were cultured in medium with DBT as their sulfur source.

We have also carried out more ultrastructural analysis on IGTS8 cells which have been cultured for many generations in medium containing sulfate or DBT sulfone as the sole sulfur source; to study whether the use of these compounds as the sole sulfur source will change the pattern of the protuberant structures. These new results further support our previous observations. IGTS8 cells, originally cultured in DBT but subsequently cultured in sulfate medium for numerous generations, still retain some, but less, of the protuberant structures on their cell wall, and some contain no such protuberant structures at all. We are planning to carry out more electron microscope analysis of IGTS8 cells cultured in different media as well as to carry out similar analysis on other species of *Rhodococcus rhodochorous* to determine whether the cell wall protuberant structures of IGTS8 are related to the enzymes for organic sulfur degradation.

In preparation for the forthcoming electron microscope studies, we ordered two *R. rhodochorous* strains from ATCC. They are ATCC 184 and ATCC 13808. The selection of these two strains are totally random. In order to be able to interpret the electron microscope results, we finished some preliminary characterization of these two strains. To our surprise, we found that both these strains can grow on DBT, although very poorly. However, they also cannot grow very well in medium with sulfate as the sole sulfur source (poorer than in DBT medium). Our results might imply that most *R. rhodochorous* species contain the enzymes required for the conversion of the sulfur moiety of DBT to sulfate but can not use sulfate as their sulfur source.

We also found that the major product of IGTS8 degradation of DBT seemed to be benzoic acid but not 2-hydroxy-biphenyl as reported previously by Kilbane and others. We have already repeated the same analysis (by HPLC) quite a few times and the same results were obtained. We are continuing to carry out more analysis with sophisticated controls in order to solve this dilemma.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

IGT58 w
DBT as the
sulfur source

IGT58 w
 $SO_4^{=}$ as the
sulphur source

K3B in
DBTS

K3B in
 SO_4^{2-}

Mobile : 45% CH₃CN
UV : 254

CH. 1 C.S 2.50 ATT 8 OFF5 10 09/02/92 15:08

- Standards : Mixture in 45% CH₃CN
- ①. Benzoic Acid
 - ②. DBT sulfoxide
 - ③. DBT Sulfone
 - ④. D,O'-Biphenol (Lower layer)
 - ⑤. DBT

D-2500

09/02/92 15:08

METHOD: JING LU TAG: 27 CH: 1

FILE: 1 CALC-METHOD: EXT-STD TABLE: 8 CONC: AREA

NO.	RT	AREA	HEIGHT	G/DL	BC	NAME
2	1.95	11638	368	*****	BU	
3	2.88	10830684	761227	*****	UU	
4	3.46	3403	790	3403.000	TBB	
5	3.76	2998	358	2998.000	TBB	
6	4.33	4396	445	4396.000	TBB	
7	4.82	2889520	215770	*****	UU	
8	6.13	1134694	81048	*****	UB	
9	8.32	10106496	524694	*****	BB	
14	30.87	1269591	21355	*****	BB	
TOTAL		26253420	1606055	*****		
PEAK REJ :		2000				
SF :		1.000				
SAMP-AMT :		1.000				

RAW DATA STORAGE NO. 15

Flow rate : 1.0 ml/min.
Mobile : 45% CH₃CN
UV : 254

CH. 1 C.S 2.50 ATT 8 OFF5 10 09/02/92 11:38

DBT sulfo

CH. 1 C.S 2.50 ATT 8 OFFS 10 09/26/92 12:55

*Leads will be used
The supernatant analyzed*

up layer

(9/26)

0-1900

09/26/92 12:55

METHOD: NANCY

TAG: 3 CH: 1

FILE: 1 CALC-METHOD: EXT-STD TABLE: 8 CONC: AREA

NO.	RT	AREA	HEIGHT	G/DL	BC	NAME
1	2.66	6530749	763560	0.000	BU	B-ACID
2	3.00	159384	24146	0.159	TBB	
3	3.69	46552	3525	0.046	TBB	
4	4.16	1164434	73437	0.000	UU	S-OXID
5	5.92	778715	31208	0.778	UU	
6	7.02	6678	357	0.006	TBB	
7	13.09	3203	32	0.003	BB	
8	17.13	161725	4221	0.161	BB	

TOTAL 8851440 900486 1.156

PEAK REJ : 2000

SF : 1.000

SAMP-AMT : 1.000

RAW DATA STORAGE NO. 15

CH. 1 C.S 2.50 ATT 8 OFFS 10 09/26/92 13:33

DBT with 15T

CH. 1 C.5 2.50 ATT 8 OFFS 10 09/26/92 12:55

BT with 16758 (11.35)
up layer

0-2500

09/26/92 12:55

METHOD: NANCY

TAG: 3 CH: 1

FILE: 1 CALC-METHOD: EXT-STD TABLE: 8 CONC: AREA

NO.	RT	AREA	HEIGHT	G/DL	BC	NAME
1	2.66	6530749	763560	0.000	BU	B-ACID
2	3.00	159384	24146	0.159	TBB	
3	3.69	46552	3525	0.046	TBB	
4	4.16	1164434	73437	0.000	UU	S-OXID
5	5.92	778715	31208	0.778	UU	
6	7.02	6678	357	0.006	TBB	
7	13.09	3203	32	0.003	BB	
8	17.13	161725	4221	0.161	BB	

TOTAL 8851440 900486 1.156

PEAK REJ : 2000

SF : 1.000

SAMP-AMT : 1.000

RAW DATA STORAGE NO. 15

CH. 1 C.5 2.50 ATT 8 OFFS 10 09/26/92 13:33

DBT with 167

SAMP-AMT : 1.000

RAW DATA STORAGE NO. 15

CH. 1 C.S 2.50 ATT 8 OFFS 10 09/26/92 13:33

D-2500

09/26/92 13:33

METHOD: NANCY TAG: 4 CH: 1

FILE: 1 CALC-METHOD: EXT-STD TABLE: 8 CONC: AREA

NO.	RT	AREA	HEIGHT	WID	BC	NAME
1	1.58	19140	1185	0.019	BB	
2	2.91	3988332	189087	3.988	BU	
3	4.22	17017	1099	0.000	TBC	S-OXID
TOTAL		4024489	191379	4.007		
PEAK REJ :		2000				
SF :		1.000				
SAMP-AMT :		1.000				

RAW DATA STORAGE NO. 16

**DATE
FILMED**

21/7/93