D.B. Moore-Shedrow, Manager
Authorized Derivative Classifier

**THE CONCENTRATION OF RADIONUCLIDES AND METALS IN VEGETATION
ADJACENT TO AND IN THE SRL SEEPAGE BASINS (U)**

Charles E. Murphy Jr.

Publication Date: December 14, 1992

Approved By: D.B. Moore, Manager
Environmental Sciences Section
Savannah River Laboratory

Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808

SAVANNAH RIVER SITE

**THIS DOCUMENT WAS PREPARED IN CONNECTION WITH WORK UNDER THE
U.S. DEPARTMENT OF ENERGY (CONTRACT DE-AC09-89SR18035).**

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

CONTENTS

ABSTRACT	1
INTRODUCTION.....	1
Description.....	2
History of Operation	3
Soil Concentrations	3
<i>Description of Sampling Sites</i>	3
<i>Differences between Basins</i>	4
<i>Variation with Depth</i>	5
METHODS.....	6
Sampling	6
Analysis.....	6
<i>Preparation</i>	6
<i>Radionuclides</i>	6
<i>Metals</i>	7
RESULTS.....	7
Comparison of Leaves and Wood	7
Comparison of Pines and Hardwoods	8
Relationship of Location to Concentration	9
Vegetation /Soil Concentration Ratios	10
<i>Vegetation on the Bottom of Basin Four</i>	10
<i>Vegetation on the Edge of the Basins</i>	12
Estimation of Total Radionuclides and Metallic Elements in the Vegetation	13
REFERENCES	14
APPENDIX	15

LIST OF FIGURES

1. Layout and Sampling Scheme for Collection of Vegetation from the SRL Seepage Basins.	2
2. Distribution of Radionuclides among Basins (Waste-Water entered at Basin 1 and only rarely reached Basin 4).	4
3. The Decrease in ¹³⁷ Cs with Distance from the Surface.	5
4. The Slope (Logarithmic) of Contaminant Concentration versus Distance from the Surface in Each Basin.	5
5. Relationship of Leaf to Wood Concentration for Strontium-90.	7
6. The Effect of Location on Average Radionuclide Contaminant Concentrations in Basin 4.	9
7. The Effect of Location on Average Metallic Contaminant Concentrations in Basin 4.	9
8. Vegetation/Soil Concentration Ratios for Trees growing in the Sediments of Basin 4.	11
9. Geometric Mean Vegetation/Soil Concentration Ratios for Trees growing on the Edge of All the Basins.	12

LIST OF TABLES

1. The Dimensions and Approximate Capacity of the SRL Seepage Basins.	2
2. Radioactive Releases to the SRL Seepage Basins from 1958 to 1980 (DPSTD-84-110).	3
3. Estimated Releases of Selected Metals (DPSTD-84-110).	4
4. Ratio of Concentration in Leaves to Concentration in Wood.	8
5. Ratio of Concentrations in Hardwood/Pine in Basin 4.	8
6. Concentration Ratios Measured at SRL Seepage Basins and a Literature Survey.	11
7. Total Contaminants in SRL Seepage Basin Trees.	13
8. Radionuclide Concentrations (pCi/gm-dry) in Trees from the SRL Seepage Basins.	16
9. Metal Concentrations (mg/kgm-dry) in Trees from the SRL Seepage Basins.	18

THE CONCENTRATION OF RADIONUCLIDES AND METALS IN VEGETATION ADJACENT TO AND IN THE SRL SEEPAGE BASINS

by
Charles E. Murphy Jr.
Savannah River Technology Center
Aiken, S.C.

ABSTRACT

In 1991 the trees on the dikes surrounding the SRL Seepage Basins were sampled and analyzed to inventory the contaminants transported from the basins into the vegetation. Tree leaves and wood were collected and analyzed for ^{90}Sr , ^{60}Co , ^{137}Cs , ^{238}Pu , $^{239,240}\text{Pu}$, $^{242,244}\text{Cm}$, ^{241}Am , Ba, Cr, Hg, Mg, Mn, Ni, and Pb. The concentrations of contaminants were influenced by sample type (leaves versus wood), species type (pines versus hardwoods), and location relative to distance from the basin.

The concentration in the vegetation and the sediments were used to calculate concentration ratios for trees growing on the sediments in the bottom of Basin 4, the only basin that was not flooded with water. With the exception of Mg, Cr, and Ba, which are significantly higher, the concentration ratios all fall within the range of previously determined ratios reported in the literature; however, in some cases the mean concentration ratio measured at the SRL Basins is higher than the mean literature values.

Concentration ratios were also calculated for the trees growing on the edge of all four basins. The mean concentration ratios for the contaminants in trees growing at the edge of the basin reflected both the effect of type of contaminant and location, as seen in Basin 4. Ba, ^{90}Sr , Mg, and Mn have high concentration ratios similar to those found in the trees growing directly on the sediments. This suggests a relatively high mobility and transport of these materials through roots extending under the basin. Other contaminants, such as ^{137}Cs , Pu isotopes, and ^{241}Am have lower concentration ratios for the trees growing on the edge of the Basins than for trees growing in the bottom of the basin, suggesting that these contaminants are less mobile.

The total inventory of each contaminant in the trees was estimated. The relationships between leaf and wood, pines and hardwood, location, and mass of the material in each of these classes were used to weight the total inventory estimate. The radionuclide with the largest inventory was 0.7 mCi for ^{90}Sr . The metallic contaminant with the largest inventory was Mn at 200 gm.

INTRODUCTION

Seepage basins are designed to contain aqueous waste so that the wastewater will flow into the soil at the bottom of the basin. Soils can be very effective filters for particulates suspended in water and generally are highly absorptive of cations in solution. Most soils have relatively low absorptive capacity for anionic solutes. Even though seepage basins have been very effective in removing the majority of radionuclides, such as ^{137}Cs and the transuranic nuclides, from wastewater, the use of seepage basins for treatment has been discontinued

at Savannah River Site (SRS). This is, in part, because of the inability to remove some other radionuclides, such as ^{90}Sr and ^3H . The measurements of radionuclide and metal contamination of vegetation growing adjacent to and in the Savannah River Laboratory (SRL, now the Savannah River Technology Center) Seepage Basins is part of the plan for closing this basin system. The total radionuclide and metallic element content of the vegetation will be used to choose disposal options for the vegetation when the basins are closed.

Description

The SRL Basins and the sediments in the basins are described (Bransford et al. 1988). The following description of the SRL Basins is largely taken from this document.

The four Seepage Basins are located south of Road A-1 and west of Road D-1 in the northwestern section of SRS. The approximate coordinates of the southeast corner of Basin 1 are 33.345128°N and 81.733505°W.

The basins are rectangular in shape and were constructed by removing the earth from within the basins to form the surrounding dikes. The construction of Basins 1, 2, and 3 principally involved excavation of natural soils and the construction of limited perimeter dikes. The construction of Basin 4 required substantial filling at the north end (adjacent to Tims Branch) to achieve both the basin bottom and the dike crest elevation.

Table 1. The Dimensions and Approximate Capacity of the SRL Seepage Basins.

Basin	Dimensions (LxWxD, m)	Volume (m³)
1	40 x 19 x 2.0	1,520
2	40 x 40 x 2.0	3,200
3	53 x 38 x 2.7	5,440
4	94 x 46 x 3.4	14,700

Waste-water entered Basin 1 by a pipeline from SRL. The water overflowing Basin 1 entered Basin 2 and then sequentially into Basins 3 and 4 when the water flow was greater than the infiltration capacity of the preceding basins (see figure 1). Waste-water seldomly entered Basin 4 because the infiltration capacity of the first three basins equaled or exceeded the input volume. Water entering the sediment at the base of the water column appears to

Figure 1. Layout and Sampling Scheme for Collection of Vegetation from the SRL Seepage Basins.

have infiltrated downward, toward the ground water. There is no evidence that water from the basins reached Tims Branch, a small stream that is near down the hydrologic gradient from Basin 4.

Basins 1 and 2 have no vegetation growing on the basin bottom. The bottom of these basins has generally remained covered with water, even after discharges to the basins stopped. The slopes of the basin dikes are densely covered with young trees. Most of the trees were less than ten years in age at the time of this sampling (June 1991). The majority of the trees are deciduous angiosperms (hardwoods), with a few scattered conifers (pines).

The bottom of Basin 3 also lacks any vegetation. At the time of this study, the basin bottom was intermittently covered with water. The vegetative cover on the slopes of the dikes surrounding Basin 3 contains less woody vegetation with a majority of the surface covered by grasses and herbaceous vegetation. The woody vegetation is largely hardwood trees with some scattered pines.

Basin 4 is the only basin that has trees growing on the bottom of the basin. Most of the basin is covered by a dense stand of young pines. A few hardwoods are also growing in the basin. The age of the trees was estimated to be less than ten years at the time of this study. The edge of the basin was characterized by a less dense stand of the same species. There is a small stand of older pine trees, estimated age 25 to 35 years, growing on the high dike separating the north side of Basin 4 from Tims Branch.

History of Operation

The first two basins were placed into operation in 1954. Basins 3 and 4 were added in 1958 and 1960, respectively. SRL used the seepage basins to dispose of low-level radioactive liquid wastes. Pipes that tied into the low-level drains in the laboratories transferred the waste into one of four underground tanks. When one of the tanks accumulated approximately 20 m³ of waste, the contents were sampled. If the analysis of the wastewater indicated concentrations not exceeding 100 d/m/mL alpha or 50 d/m/mL beta-gamma, the waste-

water was discharged to the seepage basins. If concentrations exceeded these limits, the wastewater was sent to another facility.

Releases to the SRL Seepage Basins stopped after 1982. During the 28-year loading history, approximately 130,000 m³ of water were discharged to the basins. Table 2 lists the radioactive releases between 1958 and 1980. An estimate of the total discharge of other contaminants was made on the basis of the concentrations in the low-level waste stream sampled during October, 1982. The loadings of selected metals are listed in Table 3.

Soil Concentrations

Description of Sampling Sites

Shelby tube samplers bored five 20 ft. sample cores from the bottom of each basin, a total of twenty cores. The top 2 ft of each Shelby sampler was cut into 3-inch segments which were removed from the tube segments in the field. Below the top 2 ft, 3-ft. intervals were used for analysis. A total of 280 sediment samples were collected.

Following collection, the samples were shipped to a contract laboratory (Envirodyne) for analysis. A description of the analytical techniques

Table 2. Radioactive Releases to the SRL Seepage Basins from 1958 to 1980 (Bransford et al. 1988).

<u>Radionuclide</u>	<u>Released Activity (Ci)</u>
³ H	105
^{89,90} Sr	0.4
¹³⁷ Cs	4.7
Natural U	0.022
²³⁸ Pu	0.009
²³⁹ Pu	0.003
²⁴¹ Am	0.001
^{242,244} Cm	0.001
^{103,106} Ru	1.4
⁶⁰ Co	0.1
^{141,144} Ce	2.7
Alpha (unidentified)	4.2
Beta-gamma	10.6

Table 3. Estimated Releases of Selected Metals (Bransford et al. 1988).

Element	Release (kg)
Barium	64
Chromium	1,220
Mercury	26
Magnesium	810
Manganese	438
Nickel	670
Lead	515
Zinc	219

can be found in Bransford et al. (1988)

Differences between Basins

Figure 2 illustrates the average concentration of the seven radionuclides measured in the top two feet of the sediments, relative to the maximum

value measured. The concentrations fall into three patterns. The concentration of ^{137}Cs , ^{238}Pu , and $^{239,240}\text{Pu}$ increase from Basin 1 to a maximum in Basin 2 and decrease in Basin 3 and further to a minimum in Basin 4. ^{90}Sr concentrations peak in Basin 3.

^{241}Am and $^{242,244}\text{Cm}$ show the pattern of a peak in concentration in Basins 2 but also show an increase in concentration between Basins 3 and 4. The variability in cores from the same basins is large enough that the increase between Basins 3 and 4 may be an artifact of the sampling procedure.

Similar analysis of the six elemental contaminants, Ba, Cr, Hg, Mg, Mn, and Ni show similar patterns. All have peak concentrations in Basin 2. Ba, and Cr, show an increase in concentration between Basins 3 and 4, while the other elements show a continued decrease. The levels of two contaminants that show up in vegetation, Hg and Pb, are too low in the sediments to be able to

Figure 2. Distribution of Radionuclides among Basins. Waste-Water entered at Basin 1 and only rarely reached Basin 4.

determine their spatial distribution among the Basins.

Variation with Depth

With a few exceptions, the concentration of all the contaminants measured in both the vegetation and the sediments show a marked decrease in concentration with distance from the surface. This is expected if these radionuclides and metallic elements are strongly absorbed by the sediments. In many cases, a straight line relationship is found between concentration and depth when the concentration of a radionuclide or a metal collected at a particular depth is plotted against depth in logarithmic coordinates. Figure 3 illustrates this for the average concentration of ^{137}Cs in the cores from Basin 4.

Figure 3. The Decrease in ^{137}Cs with Distance from the Surface.

Figure 4. The Slope (Logarithmic) of Contaminant Concentration versus Distance from the Surface in Each Basin.

concentration with depth, are very similar for most of the radionuclides. The slopes of this relationship for the metallic elements shows a greater variability between elements and between the same element in different basins. Only ^{90}Sr and Mn in Basin 4 show a positive slope (increasing concentration with depth). No explanation is available to explain the reason for the positive slopes. The fact that Basin 4 only rarely received wastewater and it is the basin having vegetation growing on the bottom of the basin may partially explain this behavior.

METHODS

Sample Collection

One kilogram of leaves and one kilogram of branch wood were collected for analysis. Each sample consisted of material from a single tree placed in double polyethylene bags which were sealed before moving to the next sampling location.

Vegetation samples were taken from the edge of the basins at each of the four corners. The corners were selected because they could be easily located if more sampling was needed later and because the trees closest to the edge of the basins were likely to show the highest concentrations. In addition to the 16 locations at the edge of each basin, 12 trees growing on the bottom of Basin 4 were sampled. Four trees were selected closest to the three soil sampling locations identified as 4-1, 4-3, and 4-5 in Bransford et al. (1988), identified as sample location 7 through 9, 10 through 14, and 15 through 18 respectively in Figure 1. Two additional trees were sampled near the top of the dike on the north side of Basin 3. A total of 60 samples were collected, 30 leaf samples and 30 branch wood samples (see figure 1).

The samples were sealed in a DOT type 7A drum. The drum was shipped to Thermo Analytical, Inc., Richmond, CA where the vegetation was analyzed for selected radionuclides and metallic elements.

Analysis

Sample Preparation

The total weight of the vegetation sample was taken. The sample was homogenized. A measured fraction was taken and dried. A dried weight was obtained to determine percent moisture. From this fraction an aliquot was taken and ashed for dissolution.

Radionuclides

The required radiological analyses were performed on aliquots of undried vegetation, dried vegetation, or dissolved vegetation depending on the volatility of the elements. A 500 mL sample of dried vegetation was prepared for gamma counting (^{137}Cs , ^{60}Co) and placed in a Marinelli beaker and weighed. The sample was counted for at least 400 minutes on a Ge(Li) high-resolution gamma spectrometer. The gamma spectrum was evaluated and quantified by a computer program. Positive peaks above background were calculated to desired units. The instruments were calibrated using multiple gamma standard solutions.

Ashed samples were leached with HNO_3 and H_2O_2 solutions and filtered. The leached sample was then ashed and sequentially treated with HF / HNO_3 , H_3BO_3 , HCl , and HNO_3 . The solution remaining after these treatments was added to the leachate. A sequential chemical separation of the dissolved vegetation was used to separate fractions for Sr, Pu, and Am-Cm analysis. The tracers ^{85}Sr , ^{243}Am , and ^{242}Pu were used in the analyses. Plutonium is separated on an ion exchange column. The eluted Pu was electroplated onto a stainless steel disc for counting by alpha spectrometry.

The fraction passing through the Pu ion exchange column was back extracted and concentrated. The Am-Cm fraction was separated from the Y carrier on an anion exchange column. After further purification, the Am-Cm fraction was electroplated on a stainless steel disc for counting by alpha spectrometry.

Each of the alpha-emitting actinides, which has been deposited on stainless steel discs, was

counted for at least 1000 minutes on an alpha spectrometer. The spectrum was analyzed by a computer program that used the peak intensities and the yields calculated from the tracers to calculate the concentration of these nuclides.

Problems were encountered with the ^{241}Am and $^{242,244}\text{Cm}$ analysis because of interference by other radioisotopes in the samples. The comparatively high levels of ^{244}Cm in some of the samples interfered with the ^{241}Am spectra. This raised the minimum detectable level in three of the samples. The laboratory also found unusually high recoveries of the ^{243}Am tracer. It is suspected that the presence of ^{245}Cm interfered with the ^{243}Am spectra. The samples showing the high recovery of the tracer were reanalyzed, without tracer, and the peak was corrected by comparison with the ^{244}Cm peak.

Further decontamination of the ^{90}Y from the Am-Cm separation is done by coprecipitation of Y. The precipitate is ashed to Y_2O_3 and mounted in a copper backed Mylar film sandwich on a Nylon planchet for beta counting. Each sample is counted up to 5 times to check the decay of ^{90}Y . This is used to calculate the amount of ^{90}Sr at the time of collection.

Metals

The metal concentration in the vegetation samples, except Hg, were determined by atomic absorption analysis using EPA Method's 7080, 7190, 7450, 7460, 7520, 7420, and 7950, respectively for Ba, Cr, Mg, Mn, Ni, Pb, and Zn. Mercury analysis was done by the cold vapor technique described in EPA method 7470 (EPA 1986).

RESULTS

The primary goal of this project was determining the total radionuclide and metallic element content of the vegetation. The best estimate of the total quantity of these materials requires a knowledge of the distribution of the materials. For instance, if there is a consistent difference in the concentration of a contaminant between the tree

leaves and the wood, an estimate based only on the wood might be biased. Three factors were investigated to improve the accuracy of the final estimate of the total inventory of each contaminant:

- 1) the relationship between concentration in leaves and wood
- 2) the difference in concentration between pines and hardwoods, and
- 3) the location of the trees relative to the basins.

Comparison of Leaves and Wood

The concentration of each contaminant in the wood was plotted against the concentration in the leaves collected from the same branch. Figures 5 is an example of the relationship found for ^{90}Sr . A straight line was fit to this relationship by least-squares regression. The slopes of the regression lines are an estimate of the concentration expected in leaf tissue given the concentration of wood tissue from the same branch. If the slope is 1.0 the concentrations would be expected to be identical.

Figure 5. Relationship of Leaf to Wood Concentration for Strontium-90.

Table 4 lists the slope for each of the analyzed contaminants. All of the slopes except ^{238}Pu and Pb are statistically significantly different from unity, suggesting that leaf concentrations of contaminants are different from wood concentrations. In the case of these two contaminants there is no choice but to pool the leaf and branch data when estimating the total contaminant concentration in the vegetation. However, even though the slopes calculated for $^{239,240}\text{Pu}$, $^{242,244}\text{Cm}$, ^{241}Am , Cr, and Ni are significantly different from unity, the low regression coefficients, shown in column 3, indicate that the slopes do not provide reliable predictions of leaf concentrations from wood concentrations. The relationships for ^{90}Sr , ^{60}Co , ^{137}Cs , Ba, Mg, and Mn are stronger and can be used to estimate different concentrations for the leaves and the wood. Even in the case of these contaminants it should be recognized that the slopes for ^{90}Sr and Ba are so close to unity that there is no practical difference between leaf and wood concentrations.

Table 4. Ratio of Concentration in Leaves to Concentration in Wood.

<u>Contaminant</u>	<u>Slope</u>	<u>R-square</u>
^{90}Sr	1.05*	0.81
^{60}Co	3.21*	0.91
^{137}Cs	2.72*	0.87
^{238}Pu	0.39	0.05
$^{239,240}\text{Pu}$	1.06*	0.29
$^{242,244}\text{Cm}$	0.63*	0.49
^{241}Am	1.63*	0.62
Ba	1.09*	0.84
Cr	1.03*	0.35
Hg	1.00*	0.77
Mg	4.11*	0.77
Mn	6.18*	0.87
Ni	2.55*	0.57
Pb	0.13	0.19

* Slope significant at the 0.99 confidence level.

Comparison of Pines and Hardwoods

Preliminary inspection of the contaminant concentrations suggested that the concentrations of some contaminants were higher in the hardwood trees than in the pines (Table 5). Most of the trees on the edges of Basins 1, 2, and 3 are hardwoods. Only in and around Basin 4 were there enough trees of each type to test the hypothesis that the mean concentrations were different.

Ratios of the mean concentration in hardwoods to the mean concentration in pines were calculated for each contaminant. Then the hypothesis of different mean values was tested by applying a Student t-test. Table 5 shows the ratios and the probability that the mean values are the same. The t-test suggests that the means are different for ^{137}Cs , Ba, Mg, and Mn. The differences for ^{90}Sr , Hg, Pb and perhaps ^{60}Co are marginal, while all other differences are within the range expected by random variation.

There are actually two sources of variation in the data from Basin 4. Most of the pines sampled are growing on the bottom of the basin while most of the hardwoods are on the edge of the basin. One would expect the pines to have higher concentra-

Table 5. Ratio of Concentrations in Hardwood/Pine in Basin 4.

<u>Contaminant</u>	<u>Ratio</u>	<u>P(T<=t) one-tail</u>
^{90}Sr	2.77	0.071
^{60}Co	3.39	0.103
^{137}Cs	0.36	0.038
^{238}Pu	3.79	0.411
$^{239,240}\text{Pu}$	3.41	0.416
$^{242,244}\text{Cm}$	2.24	0.365
^{241}Am	3.64	0.341
Ba	13.89	0.002
Hg	0.70	0.055
Mg	3.06	0.005
Mn	3.44	0.012
Ni	1.46	0.690
Pb	0.59	0.064

tions because they are rooted in sediments with higher concentrations of contaminants. Therefore, if the pine trees have higher concentrations than the hardwoods of a particular contaminant, this could be caused by either the pine trees being rooted in soil with a higher concentration of the contaminant or by the pine trees being more efficient in removing the contaminant from the soil or both. However, if hardwoods have higher concentrations of the contaminant and, on the average, are growing in sediment with a lower concentration of the contaminant, then the increased uptake is likely to have been caused by an actual increase in the efficiency of the hardwoods in removing this contaminant from the sediment. Therefore, the lower concentration of ^{137}Cs in pines (ratio less than 1) may have been caused by either sediment or vegetation conditions while the higher concentration of ^{90}Sr , Ba, Mn, and Mg are probably caused by vegetative processes.

Relationship of Location to Concentration

The effect of location is important only at Basin 4 which is the only basin with trees growing on its bottom. At Basins 1 and 2, most of the trees are growing on the slope of the dike surrounding the basin and are not very far from the basin edge. Basin 3 has comparatively few trees, and most are on the edge of the basin. Therefore the effect of location was evaluate for only Basin 4. Because of the difference in concentration between different species types, noted in the previous section, only hardwood trees were used in this analysis. The concentration of contaminants were not measured in pines at all the other locations. Figures 6 and 7 illustrate the effect of location for the radionuclides and metallic elements, respectively.

With the exception of ^{90}Sr , there is a large decrease in concentration of the radionuclides as the location of the tree is further from the basin. The relatively high concentrations of ^{90}Sr in the trees growing on the basin edge and even the trees higher up on the dike, may be related to the relatively high mobility of ^{90}Sr in soils.

Figure 7. The Effect of Location on Average Metallic Contaminant Concentrations in Basin 4.

Figure 6. The Effect of Location on Average Radionuclide Contaminant Concentrations in Basin 4.

The relationship of concentration in the vegetation and the location of the trees is very different for the four metallic elements (Ba, Mg, Mn, and Ni). The concentration in the vegetation was actually higher for trees growing on the edge of the basin than on the bottom. This is difficult to explain by any known process. Since Basin 4 received only limited amounts of wastewater, the concentrations of these naturally occurring elements may be higher in the soil used for the dikes than the soil in the bottoms of the basin.

Vegetation/Soil Concentration Ratios

Concentration ratios are used to estimate the concentration of contaminant in one component of the environment given the concentration in another component that is the primary source of the contaminant to the former component. Vegetation-soil concentration ratios are used when the soil is the primary source of the contaminant of interest. Vegetation-soil uptake ratios vary greatly because of the complex relationship between environmental factors that effect the uptake of materials from soils. One important factor under field conditions is the depth of vegetation rooting. The ratios presented here were calculated on the basis of an average concentration over the top 1 ft of sediment. The same calculations for an average over 2 ft of sediment will give ratios that are approximate twice the ratio for contaminants retained in the upper few inches of the soil, such as ^{137}Cs , but give little difference for contaminants that are more uniformly distributed with depth, such as ^{90}Sr . For this reason concentration ratios developed under similar condition of soil type, water regime, soil contaminant distribution, and vegetation are used whenever possible.

Normally, this approach is used when vegetation is grown in soil with a well-characterized radionuclide concentration. Concentration ratios will also be calculated for vegetation growing on the edge of the SRL Basins. As described in the previous section, the effects of location, species type and soil will influence the vegetation concen-

tration relative to the concentration in the sediment on the bottom of the basin. While not an ideal situation, the concentration ratios derived may still be the best way of estimating radionuclide and, perhaps, metallic element concentrations around similar seepage basins.

The results of the concentration ratios presented in Table 6 and Figures 8 and 9, are give as geometric means with logarithmic-transformed one standard deviation error bars. This was done because the frequency distribution of the data was limited at the lower end of the range by minimum values that are near zero. This leads to skewed frequency distributions that resembles a log-normal distribution.

Vegetation on the Bottom of Basin Four

Table 6 and Figure 8 illustrate the concentration ratios calculated for the pine trees growing on the bottom of Basin 4. The uptake ratios are calculated for the leaves. Ratios for wood can be calculated by dividing the leaf ratios by the factors developed in the previous section. The concentration ratios are compared to ratios and ranges of ratios from several other sources; few results are available for trees. The quoted means and ranges are, for the most part, from the vegetative parts of food crops or forage. The occasional tree values fall in this range.

With the exception of Mg, Cr, and Ba, which are significantly higher, the concentration ratios fall within the range of previously determined ratios. In the case of Mg, Cr, and Ba the range of concentration ratios were only available in Baes et al. (1984), which generally indicated a lower range of ratios than Coughtrey et al. (1985) or Till and Meyer (1983).

However, even for contaminants where the concentration ratio falls within the range of previous measurements, the mean concentration ratio at the SRL Basins is higher than the mean literature values for a number of the contaminants. The most likely explanation is that many of the soils used in previous studies are fertilized for crop production.

Figure 8. Geometric Mean Vegetation/Soil Concentration Ratios for Trees growing in the Sediment of Basin 4.

Table 6. Concentration Ratios Measured at SRL Seepage Basins and a Literature Survey.

Nuclide/Element	SRL Basins		Literature	
	Basin 4	Edge	Mean	Range
⁹⁰ Sr	1.69	0.88	2.93(2)	(0.1-20)(2)
⁶⁰ Co	0.61	0.018	0.06(1)	(0.01-1)(3)
¹³⁷ Cs	2.37	0.0052	0.26(2)	(0.1-10)(3)
²³⁸ Pu	0.0047	0.00015	0.00045(1)	(0.006-0.1)(2)
^{239,240} Pu	0.0073	0.00012	0.00045(1)	(0.006-0.1)(2)
^{242,244} Cm	0.021	0.0052	0.00085(1)	(10 ⁻⁶ -1.7)(2)
²⁴¹ Am	0.00014	0.00025	0.0055(1)	(10 ⁻⁴ -10 ⁻¹)(2)
Ba	0.43	5.0	0.005(1)	(0.056-0.16)(1)
Cr	ND	0.049	0.02(3)	(10 ⁻³ -0.0145)(1)
Mg	67.	60.	1.0(1)	(0.2-4.0)(1)
Mn	16	17	1.48(2)	(1-100)(2)
Ni	1.12	0.43	0.06(1)	(0.1-10)(2)
Pb	0.58	0.15	0.10(1)	(0.013-0.9)(1)

Sources: (1) Baes et al. 1984, (2) Coughtrey et al. 1985, and (3) Till and Meyer 1983.
 ND, no data above detection limit for either soil or vegetation.

Figure 9. Geometric Mean Vegetation/Soil Concentration Ratios for Trees growing on the Edge of All the Basins.

Fertilization should increase the concentration of Ca and K, which are known to compete with Sr and Cs uptake respectively. The increased uptake of Cs in plants grown in low K soil is particularly pronounced on sandy soils, such as those found at SRS. Under these conditions the upper end of the uptake range has been found in the literature reviewed.

Vegetation on the Edge of the Basins

Table 6 and Figure 9 show the geometric mean concentration ratios calculated for the radionuclides and metallic elements measured in both trees and soil. The concentration factors are for leaves. The concentration ratios for wood can be found from the ratios developed in the section dealing with leaf-wood differences. The variability in concentration ratios, as illustrated by the one standard deviation (logarithmic transform) error bars in Figure 9, as expected, is large. However, it

is possible to use the results to provide an upper bound to the expected concentrations under similar conditions.

The average concentration ratio of the trees located at the edge the basins, show the same relationship to the concentration ratios for the trees in the bottom of basin 4 as the relationship to concentration with location at Basin 4. ⁹⁰Sr, Ba, Mg, and Mn have high concentration ratios consistent with the relatively high concentrations of these contaminants in vegetation at the edge of the basin. These relatively high concentrations could be explained by either high mobility or high transport of these materials through roots extending under the basin. Conversely, contaminants such as ¹³⁷Cs, Pu isotopes, and ²⁴¹Am which appear to be less mobile have lower concentration ratios for the trees growing on the edge of the basins than for trees growing in the bottom of the basin.

Estimation of Total Radionuclides and Metallic Elements in the Vegetation

Table 7 shows the quantity of each radionuclide and element in each basin. The volume of each component was estimated on the basis of a vegetation survey of the seepage basins. All trees were tallied by height and diameter (at the soil line for small trees and at breast height for larger trees). Stem wood volume was estimate from Monk (1966) for smaller trees and from Wahlenberg (1960) for larger trees. Wood density was taken from Koch (1972). Based on data from these same sources, 20% was added for branch wood and 10% of the

total was used to estimate the leaf mass. The 10% used for the leaf mass will under estimate the mass for small trees and over estimate it in the larger trees; however, the value is a reasonable mean for trees of the size growing in and adjacent to the basins.

The concentrations used in estimating the total inventory in any one category is the average of trees in that category, rather than the regression slope estimates extrapolated to each category. The greatest influence on the total inventories of each contaminant is the amount of tree mass growing in or adjacent to the basin.

Table 7. Total Contaminants in SRL Seepage Basin Trees.

	<u>Basin Number</u>				
<u>mCi</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>Total</u>
Sr90	8.64E-02	5.31E-02	6.61E-02	4.91E-01	6.96E-01
Co60	2.94E-05	3.16E-05	4.99E-04	4.29E-04	9.89E-04
Cs137	1.51E-03	1.86E-03	1.40E-03	5.65E-02	6.13E-02
Pu238	2.44E-07	1.60E-07	3.87E-08	1.12E-06	1.57E-06
Pu239	7.56E-07	3.10E-07	7.44E-08	3.17E-06	4.31E-06
Cm244	1.14E-04	5.04E-04	1.35E-04	4.35E-03	5.10E-03
Am241	1.10E-06	6.34E-07	3.99E-07	2.20E-05	2.41E-05
 <u>gm</u>					
Ba	4.86E+01	5.09E+01	2.66E+01	5.50E+02	6.76E+02
Cr	2.05E-01	3.50E-01	1.09E-01	4.33E+00	5.00E+00
Hg	1.51E-02	1.96E-02	3.91E-03	3.24E-01	3.63E-01
Mg	7.09E+02	5.04E+02	3.16E+02	5.59E+03	7.12E+03
Mn	1.79E+02	2.34E+02	4.63E+01	1.59E+03	2.05E+03
Ni	2.99E+00	2.48E+00	8.83E-01	2.80E+01	3.44E+01
Pb	6.60E-01	5.00E-01	3.84E-01	6.59E+01	6.75E+01

REFERENCES

- Baes, C.G., R.D. Sharp, A.L. Sjoreen, and R.W. Shor. 1984. A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides through Agriculture. ORNL-5786, Oak Ridge National Laboratory, Oak Ridge, Tenn. 150 pp.
- Bransford, J.L., H.W. Bledsoe, R.V. Simmons. 1988. Technical Data Summary, Chemical Characterization of sediments and Groundwater at the Savannah River Laboratory Seepage Basins. DPSTD-84-110, E.I. du Pont de Nemours, Inc., Savannah River Laboratory, Aiken, S.C.
- Coughtrey, P.J., D. Jackson, M.C. Thorne. 1985. Radionuclide Distribution and Transport in Terrestrial and Aquatic Ecosystems. A.A. Balkema Publ., Boston, 6 Volumes.
- EPA. 1986. Test Methods for Evaluating Solid Wastes, Vol. 1A, USEPA, Superintendent of Documents, Washington, D.C.
- Koch, P. 1972. Utilization of Southern Pines. Agriculture Handbook No. 420, Vol I, U.S.D.A., Superintendent of Documents, Washington, D.C., pp 238, 582.
- Monk, C.D. 1966. Root-shoot dry weights in loblolly pine. Bot. Gaz. 127: 246-248.
- Till, J.E. and H.R. Meyer 1983. Radiological Assessment, NUREG/CR-3332, Superintendent of Documents, U.S. Government Printing Office, Washington, D.C.
- Wahlenberg, W.G. 1960. Loblolly Pine. The School of Forestry, Duke University, Durham, N.C., p 270.
- U.S. Nuclear Regulatory Commission. 1977. Regulatory Guide 1.109, Calculation of Annual Doses to Man from Routine Releases of Reactor Effluents for the Purpose of Evaluating Compliance with 10 CFR Part 50 Superintendent of Documents, U.S. Government Printing Office, Washington, D.C.

APPENDIX
Concentration of Radionuclides and Metal in Trees

Table 8. Radionuclide Concentrations (pCi/gm-dry) and percentage error in Trees from the SRL Seepage Basin.

Basin	Tree Species	Type	Sr90	Co60	Cs137	Pu238	Pu239	Cm244	Am241								
1	27	elm	leaf	354.0	1.2	<0.113	14.40	0.8	0.00284	5.9	0.01080	3.8	0.23	3.7	0.00550	5.5	
1	28	maple	leaf	12.3	2.1	<0.038	0.01	11.0	0.00093	12	0.00090	13.0	0.01	4.2	0.00028	17.0	
1	29	sweetgum	leaf	53.7	1.5	0.038	40.0	3.40	1.4	0.00091	11	0.00115	8.9	0.24	5.8	0.00422	13.0
1	30	maple	leaf	13.5	1.3	<0.028	0.13	13.0	0.00027	23	0.00071	14.0	0.05	4.1	0.00057	17.0	
2	23	cherry	leaf	182.0	1.3	6.064	32.0	15.20	0.6	0.00042	13	0.00050	13.0	0.49	6.6	0.00150	18.0
2	24	pine	leaf	3.5	1.3	<0.066	1.92	2.4	0.00045	16	0.00031	20.0	0.03	3.4	0.00016	28.0	
2	25	sweetgum	leaf	26.0	1.7	<0.030	0.71	3.4	0.00018	27	0.00012	30.0	0.04	3.6	0.00039	19.0	
2	26	maple	leaf	84.0	1.3	<0.047	1.38	2.8	0.00042	17	0.00101	11.0	1.36	11.0	0.00091	60.0	
3	19	sweetgum	leaf	170.0	1.3	0.056	38.0	0.94	3.5	0.00008	43	0.00008	43.0	0.59	11.0	0.00111	72.0
3	20	chinaberry	leaf	320.0	1.2	<0.034	1.57	2.1	0.00014	21	0.00021	21.0	0.64	5.0	0.00115	11.0	
3	21	pine	leaf	22.9	0.8	0.032	40.0	7.45	0.7	0.00007	56	0.00008	36.0	0.12	3.4	0.00040	20.0
3	22	sweetgum	leaf	102.0	1.3	<0.031	19.50	0.6	0.00073	14	0.00077	14.0	1.59	7.1	0.00094	59.0	
4	1	sweetgum	leaf	18.7	1.0	<0.047	0.14	15.0	0.00010	43	0.00014	30.0	0.22	5.0	<0.00161	<0.00357	
4	2	cherry	leaf	125.0	1.0	<0.051	1.89	2.0	0.00015	24	0.00044	14.0	0.78	8.0	<0.00357	<0.00357	
4	3	elm	leaf	64.4	1.4	0.058	32.0	0.90	3.5	0.00005	60	0.00047	14.0	0.67	7.3	-0.00100	80.0
4	4	sweetgum	leaf	34.6	2.0	<0.029	0.50	3.7	0.00017	30	0.00027	25.0	0.92	8.5	0.00140	24.0	
4	5	sweetgum	leaf	101.0	1.3	<0.044	0.24	11.0	0.00004	50	0.00012	27.0	0.13	3.4	-0.00014	155.0	
4	6	pine	leaf	61.1	0.8	0.118	14.0	7.73	1.0	0.00020	25	0.00074	11.0	5.48	60.0	-0.00195	382.0
4	7	pine	leaf	66.9	0.9	<0.017	134.00	0.2	0.00073	10	0.00112	8.4	0.63	3.7	0.00017	106.0	
4	8	pine	leaf	23.6	2.0	0.285	8.0	87.80	0.2	0.00085	14	0.00128	11.0	0.18	3.0	<0.00046	<0.00046
4	9	pine	leaf	42.8	1.5	0.319	6.6	46.50	0.3	0.00083	10	0.00157	7.2	0.74	4.8	0.00075	42.0
4	10	willow	leaf	165.0	1.3	1.380	3.4	157.00	0.2	0.00702	5	0.01010	4.4	4.73	12.0	0.00021	184.0
4	11	pine	leaf	63.6	2.0	0.363	4.0	104.00	0.2	0.00068	15	0.00205	9.0	0.35	4.0	<0.00147	<0.00147
4	12	pine	leaf	28.3	0.8	0.319	5.7	150.00	0.2	0.00079	10	0.00123	8.6	0.56	3.9	0.00014	154.0
4	13	pine	leaf	34.4	1.5	0.282	9.9	108.00	0.3	0.00060	11	0.00057	11.0	0.53	4.4	0.00084	14.0
4	14	willow	leaf	184.0	1.3	0.639	5.6	49.30	0.4	0.00527	4.8	0.00809	4.1	6.12	24.0	<0.04260	<0.04260
4	15	pine	leaf	32.0	1.4	0.437	7.8	194.00	0.2	0.00068	10	0.00073	10.0	1.32	7.9	0.00024	148.0
4	16	pine	leaf	19.5	1.1	0.406	9.9	191.00	0.1	0.00079	11	0.00120	8.5	0.70	4.1	0.00012	191.0
4	17	pine	leaf	7.3	3.0	0.133	10.0	82.90	0.2	0.00054	14	0.00068	12.0	0.38	5.0	<0.00202	<0.00202
4	18	pine	leaf	20.4	2.0	0.312	5.2	165.00	0.1	0.00064	10	0.00092	8.7	1.03	7.2	0.00078	24.0
1	27	elm	wood	200.0	1.4	0.027	44.0	3.28	1.3	0.00014	21	0.00071	11.0	0.17	6.3	0.00150	20.0
1	28	maple	wood	6.8	2.9	0.028	38.0	< 0.03	0.00007	33	0.00005	33.0	0.00	9.2	0.00003	133.0	133.0
1	29	sweetgum	wood	115.0	0.8	0.029	32.0	1.37	1.7	0.00021	19	0.00044	14.0	0.28	3.3	0.00209	20.0
1	30	maple	wood	14.2	1.3	<0.022	< 0.03	< 0.03	0.00012	28	0.00066	10.0	0.01	6.7	0.00011	56.0	56.0
2	23	cherry	wood	59.5	1.6	0.025	35.0	4.13	1.0	0.00002	100	0.00008	33.0	0.04	3.7	0.00021	25.0
2	24	pine	wood	3.5	2.2	<0.028	<0.028	0.42	3.5	0.00001	300	0.00002	100.0	0.13	4.6	0.00024	35.0
2	25	sweetgum	wood	27.3	0.9	<0.027	<0.027	0.29	5.4	0.00006	50	0.00041	15.0	1.32	22.0	0.00032	111.0

Table 8. Radionuclide Concentrations (pCi/gm-dry) and percentage error in Trees from the SRL Seepage Basin (continued).

Basin	Tree Species	Type	Sr90	Co60	Cs137	Pu238	Pu239	Cm244	Am241							
2	26	maple	wood	113.0	0.9	<0.038	1.28	2.2	0.00046	13	0.00065	10.0	0.54	3.7	0.00172	15.0
3	19	sweetgum	wood	143.0	0.9	<0.041	0.36	9.8	0.00008	30	0.00002	67.0	0.13	4.4	<0.00096	
3	20	chinaberry	wood	277.0	0.8	<4.310	0.40	9.1	0.00008	33	0.00028	16.1	0.32	9.2	0.00084	25.0
3	21	pine	wood	16.4	1.1	<0.032	2.78	1.2	0.00001	400	0.00004	33.0	0.02	3.8	0.00001	400.0
3	22	sweetgum	wood	83.2	0.8	0.036	26.0	0.8	0.00007	40	0.00020	19.0	0.40	3.6	0.00134	11.0
4	1	sweetgum	wood	34.4	1.0	<0.034	0.07	21.0	0.00005	40	0.00013	31.0	0.08	4.0	<0.00193	
4	2	cherry	wood	42.6	2.0	<0.026	0.36	7.0	-0.00001	200	0.00016	37.0	0.20	3.0	<0.00040	
4	3	elm	wood	41.9	2.0	<0.028	0.39	4.9	0.00001	100	0.00010	25.0	0.42	19.0	<0.00477	
4	4	sweetgum	wood	75.7	1.5	<0.029	0.32	5.9	0.00005	43	0.00052	12.0	0.62	7.0	<0.00075	
4	5	sweetgum	wood	245.0	1.4	0.051	35.0	11.0	0.00052	11	0.00041	12.0	0.35	5.7	0.00088	20.0
4	6	pine	wood	52.2	0.8	0.041	34.0	1.0	0.00009	31	0.00028	15.0	0.45	6.9	-0.00024	213.0
4	7	pine	wood	60.4	0.7	0.099	7.2	0.2	0.00040	14	0.00085	9.0	0.23	3.6	0.00039	21.0
4	8	pine	wood	27.6	2.0	0.143	7.0	0.3	0.00289	6	0.00425	5.0	0.56	4.0	<0.00017	0.0
4	9	pine	wood	28.0	1.8	0.118	10.0	0.3	0.00027	18	0.00056	13.0	0.34	4.0	0.00068	16.0
4	10	willow	wood	112.0	1.7	0.324	5.5	0.2	0.00034	15	0.00176	6.6	10.40	46.0	<0.00337	
4	11	pine	wood	45.5	2.0	0.166	9.0	0.3	0.00428	5	0.00571	5.0	0.44	5.0	<0.00085	
4	12	pine	wood	14.8	1.6	0.177	11.0	0.3	0.00017	19	0.00050	12.0	0.45	12.0	0.00029	191.0
4	13	pine	wood	48.6	1.3	0.133	9.0	0.3	0.00156	6.9	0.00175	6.6	0.42	3.6	0.00018	137.0
4	14	willow	wood	193.0	0.8	0.186	6.6	0.4	0.00018	24	0.00418	4.4	3.78	9.0	<0.00244	
4	15	pine	wood	24.8	1.5	0.100	11.0	0.3	0.00010	37	0.00026	21.0	0.22	3.4	0.00025	24.0
4	16	pine	wood	22.0	1.5	0.115	12.0	0.3	0.00006	40	0.00018	22.0	0.33	3.7	0.00008	250.0
4	17	pine	wood	6.7	3.0	0.055	20.0	0.4	0.00005	33	0.00011	25.0	0.59	3.0	<0.00036	
4	18	pine	wood	23.1	1.4	0.109	10.0	0.4	0.00031	15	0.00020	20.0	0.26	3.3	0.00042	17.0

Table 9. Metal Concentrations (mg/kg-dry) in Trees from the SRL Seepage Basin.

Basin	Tree	Species	Type	Ba	Cr	Hg	Mg	Mn	Ni	Pb
1	27	elm	leaf	43.1	0.4	0.018	1180	401.0	6.0	0.6
1	28	maple	leaf	60.6	< 0.2	0.022	3280	630.0	3.6	1.3
1	29	sweetgum	leaf	82.2	< 0.2	0.026	3330	1500.0	16.0	< 0.5
1	30	maple	leaf	89.9	< 0.2	0.014	2930	901.0	5.0	0.6
2	23	cherry	leaf	79.0	2.5	0.019	2350	465.0	7.2	< 0.5
2	24	pine	leaf	5.7	< 0.2	0.052	789	725.0	1.1	< 0.5
2	25	sweetgum	leaf	95.9	< 0.2	<0.010	2330	1040.0	6.2	< 0.5
2	26	maple	leaf	108.0	2.1	0.020	2200	1420.0	11.0	< 0.5
3	19	sweetgum	leaf	73.3	< 0.2	<0.010	1840	373.0	3.9	1.3
3	20	chinaberry	leaf	41.9	0.8	<0.010	1730	60.5	< 0.8	3.6
3	21	pine	leaf	7.8	0.3	0.036	726	270.0	< 0.8	< 0.5
3	22	sweetgum	leaf	46.2	< 0.2	0.014	1650	617.0	6.6	1.9
4	1	sweetgum	leaf	74.3	0.9	0.036	2870	1550.0	< 0.3	< 0.5
4	2	cherry	leaf	70.2	1.3	0.017	2530	626.0	< 0.3	1.5
4	3	elm	leaf	131.0	2.0	0.026	2620	1340.0	1.0	< 0.5
4	4	sweetgum	leaf	23.2	18.5	0.028	4120	856.0	18.2	0.8
4	5	sweetgum	leaf	41.6	< 0.2	0.016	2840	412.0	4.2	0.8
4	6	pine	leaf	4.0	< 0.2	0.050	795	315.0	< 0.8	< 0.5
4	7	pine	leaf	5.5	< 0.2	0.052	714	201.0	5.3	< 0.5
4	8	pine	leaf	1.9	< 0.2	0.052	1060	238.0	< 0.3	1.4
4	9	pine	leaf	3.2	< 0.2	0.050	591	249.0	2.9	0.9
4	10	willow	leaf	6.0	< 0.2	0.048	1220	256.0	4.6	< 0.5
4	11	pine	leaf	2.8	< 0.2	0.051	1310	166.0	< 0.3	1.2
4	12	pine	leaf	1.8	< 0.2	0.040	872	197.0	4.1	< 0.5
4	13	pine	leaf	1.5	< 0.2	0.042	955	156.0	4.6	< 0.5
4	14	willow	leaf	16.5	0.3	0.024	1750	457.0	4.6	< 0.5
4	15	pine	leaf	1.8	< 0.2	0.034	750	239.0	3.0	< 0.5
4	16	pine	leaf	1.3	< 0.2	0.046	710	235.0	5.5	2.6
4	17	pine	leaf	0.9	0.4	0.042	733	59.0	< 0.3	< 0.5
4	18	pine	leaf	0.9	< 0.2	0.020	699	200.0	12.6	0.7
1	27	elm	wood	48.1	< 0.2	0.043	645	113.0	5.1	1.1
1	28	maple	wood	24.2	< 0.2	<0.005	395	52.1	1.0	< 0.5
1	29	sweetgum	wood	72.8	< 0.2	<0.005	546	173.0	2.0	< 0.5
1	30	maple	wood	40.4	< 0.2	<0.005	375	77.3	1.8	< 0.5
2	23	cherry	wood	35.9	0.4	<0.006	244	83.4	2.4	< 0.5
2	24	pine	wood	7.6	< 0.2	0.043	438	178.0	1.0	< 0.5

Table 9. Metal Concentrations (mg/kg-dry) in Trees from the SRL Seepage Basin (continued).

Basin	Tree	Species	Type	Ba	Cr	Hg	Mg	Mn	Ni	Pb
2	25	sweetgum	wood	52.0	< 0.2	<0.008	361	110.0	2.4	< 0.5
2	26	maple	wood	98.7	< 0.2	0.019	347	264.0	2.4	< 0.5
3	19	sweetgum	wood	116.0	< 0.2	<0.010	681	78.0	1.8	0.9
3	20	chinaberry	wood	61.8	< 0.2	<0.007	727	30.6	1.2	0.7
3	21	pine	wood	1.9	< 0.2	<0.005	259	53.0	< 0.8	< 0.5
3	22	sweetgum	wood	37.6	< 0.2	<0.005	478	103.0	2.7	< 0.5
4	1	sweetgum	wood	78.2	1.6	0.031	465	243.0	4.2	4.3
4	2	cherry	wood	29.0	0.4	0.059	215	62.6	4.3	15.2
4	3	elm	wood	87.2	< 0.2	<0.010	498	96.4	1.9	1.3
4	4	sweetgum	wood	27.3	< 0.2	0.014	509	105.0	1.8	0.8
4	5	sweetgum	wood	60.9	< 0.2	0.018	592	112.0	4.0	1.8
4	6	pine	wood	1.9	< 0.2	0.021	227	36.9	< 0.8	< 0.5
4	7	pine	wood	2.5	< 0.2	0.039	265	30.8	1.9	< 0.5
4	8	pine	wood	2.2	0.3	0.055	457	40.7	7.5	0.6
4	9	pine	wood	0.9	< 0.2	0.019	281	31.8	1.7	< 0.5
4	10	willow	wood	6.6	< 0.2	0.022	284	55.7	1.9	< 0.5
4	11	pine	wood	2.0	< 0.2	0.042	271	40.3	3.8	< 0.5
4	12	pine	wood	1.1	< 0.2	<0.010	241	20.7	1.5	< 0.5
4	13	pine	wood	1.4	< 0.2	0.033	267	29.8	2.2	< 0.5
4	14	willow	wood	25.5	< 0.2	0.016	564	124.0	1.4	0.7
4	15	pine	wood	3.3	0.3	<0.007	416	69.5	2.1	9.0
4	16	pine	wood	1.2	< 0.2	0.025	281	53.0	1.6	< 0.5
4	17	pine	wood	1.7	< 0.2	0.038	340	41.4	0.9	< 0.5
4	18	pine	wood	1.3	< 0.2	0.038	273	47.9	1.0	< 0.5

**DATE
FILMED**

6 / 17 / 93