AUGUST 1978

PPPL-1471

UC-20g

MASTER

**ALFVÉN RESONANCE HEATING VIA
MAGNETOSONIC MODES IN
LARGE TOKAMAKS**

BY

C. F. F. KARNEY AND F. W. PERKINS

MASTER

**PLASMA PHYSICS
LABORATORY**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

**PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY**

**This work was supported by the U. S. Department of Energy
Contract No. AT-(40-1)-3073. Reproduction, translation,
distribution, use and disposal, in whole or in part, by or
for the United States Government is unlimited.**

Alfvén Resonance Heating Via Magnetosonic Modes in
Large Tokamaks

Charles F. F. Karney and Francis W. Perkins
Plasma Physics Laboratory, Princeton University
Princeton, New Jersey 08540

NOTICE
This report was prepared in an account of work
sponsored by the United States Government. Neither the
United States nor the United States Department of
Energy, nor any of their employees, nor any of their
contractors, subcontractors, or their employees, makes
any warranty, express or implied, or assumes any legal
liability or responsibility for the accuracy, completeness
or usefulness of any information, apparatus, product or
process disclosed, or represents that its use would not
infringe privately owned rights.

ABSTRACT

The theory of Alfvén resonance heating of tokamaks is extended beyond the incompressible MHD model to include finite ω/Ω_1 effects, which lead to off-diagonal terms in the conductivity tensor, and compressibility, which permits the fast Alfvén mode. The finite frequency effects can greatly change the dissipation resulting from the shear Alfvén resonance. With an appropriate choice of parameters, the dissipation can vanish allowing high-Q toroidal eigenmodes in large tokamaks such as PLT.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *y b*

Tokamaks of the size of PLT are the first to nominally permit propagation of magnetosonic (i.e., compressional Alfvén) modes at frequencies that are a modest fraction of the ion-cyclotron frequency. (Typically $\omega/\Omega_i \approx 0.5$.)

In this frequency range, the transition from the high density propagation region for magnetosonic waves to the low density evanescent region does not occur via a simple cutoff, but rather by a closely-spaced cutoff-resonance-cutoff triplet. The resonance, which occurs at the point in the plasma profile where $\omega \approx k_{\parallel} v_A$, is called the shear Alfvén resonance and generally leads to energy absorption. The goal of this work is to evaluate how much absorption the shear Alfvén resonance causes. We accomplish this by calculating the reflection coefficient for magnetosonic waves incident on the cutoff-resonance-cutoff triplet.

Previous analyses of the problem^{1,2} treated the MHD limit in which $\omega/\Omega_i \rightarrow 0$, and, by and large, ignored compressibility thus suppressing magnetosonic modes. The recent work of Ott, et al.,² does recognize that the shear Alfvén resonance can damp magnetosonic modes but does not include finite frequency effects, which we find to be important, and specializes its results to the $m = 0$ ($k_y = 0$) case.

Since the dissipation depends on the fields close to the region of the cutoffs and resonance, which typically occupy only a small fraction of the minor radius of a tokamak, we may accurately treat the problem by a slab geometry model in which x is parallel to the density gradient and z is parallel to the constant magnetic field, B_0 . A linear

variation of density with x is employed. The neglect of the shear of the magnetic field is justifiable for a tokamak as long as $n \gg m/q$ (n and m are the toroidal and poloidal mode numbers and q is the safety factor). The electrons and ions are treated as separate cold fluids. All field quantities vary as $f(x) \exp(ik_y y + ik_z z - i\omega t)$. The parallel electrical field is neglected (it is shorted out by the high parallel electron conductivity). The remaining components of the electric field are described by

$$(A - k_y^2) E_x + i(D - k_y d/dx) E_y = 0, \quad (1)$$

$$-i(D + k_y d/dx) E_x + (A + d^2/dx^2) E_y = 0, \quad (2)$$

where $A = (\omega^2/v_A^2) [\Omega_i^2/(\Omega_i^2 - \omega^2)] - k_z^2$, $D = (A + k_z^2)\omega/\Omega_i$ and $v_A = c\Omega_i/\omega_{pi}$. Making the substitutions $E = E_y$ and

$$\psi = (A - k_y^2)^{-1} (A d/dx - k_y D) E_y, \quad (3)$$

we may recast (1) and (2) as a coupled set of first order differential equations,

$$\frac{d}{dx} E - \frac{k_y D}{A} E = \frac{A - k_y^2}{A} \psi \quad (4)$$

$$\frac{d}{dx} \psi + \frac{k_y D}{A} \psi = - \left(\frac{A^2 - D^2}{A} \right) E \quad (5)$$

The right-hand side of Eq. (5) is the cutoff-resonance-cutoff triplet.

We put Eqs. (4) and (5) into dimensionless form by defining a scalelength $\ell = [A'(x_0)]^{-1/3}$ and a new coordinate $\xi = (x - x_0)/\ell$, where $A' = dA/dx$ and x_0 is the point at which $A = 0$ (the finite frequency generalization of the shear Alfvén resonance). For small ξ we retain only the leading order terms in the series expansions, for A and D . Thus $A \approx \xi/\ell^2$ and $D \approx S/\ell^2$, where $S = k_{\parallel}^2 \ell^2 (\omega/\omega_i)$. Equations (4) and (5) then become

$$\frac{d}{d\xi} E - \frac{MS}{\xi} E - \frac{\xi - M^2}{\xi} \Psi = 0 \quad (6)$$

$$\frac{d}{d\xi} \Psi + \frac{MS}{\xi} \Psi + \frac{\xi^2 - S^2}{\xi} E = 0. \quad (7)$$

where $M = k_y \ell$ and $\Psi = \ell \psi$. The equation for E reads

$$\frac{d^2}{d\xi^2} E - \frac{M^2}{\xi(\xi - M^2)} \frac{d}{d\xi} E + \left(\frac{\xi^2 - S^2}{\xi} - M^2 + \frac{MS}{\xi(\xi - M^2)} \right) E = 0. \quad (8)$$

We note the following points about Eq. (8). The equation is singular at $\xi = 0$, the shear Alfvén resonance. The point $\xi = M^2$ is only an apparent singularity as may be seen from Eqs. (6) and (7). This point is the cutoff for the magnetosonic wave. In the limit $|\xi| \rightarrow \infty$ the equation becomes the Airy equation, $d^2 E/d\xi^2 + (\xi - M^2)E = 0$, which physically describes a propagating magnetosonic wave for $\xi > M^2$ and an evanescent mode for $\xi < M^2$.

We wish to determine the reflection coefficient, for this will determine Q of magnetosonic modes. To do this we solve Eq. (8) with the boundary condition that as $\xi \rightarrow -\infty$, $E \propto \text{Ai}(M^2 - \xi)$, the evanescent Airy function. We ignore the small portion of the Bi function which should be added since the boundary of the plasma where $E = 0$ is at the finite distance $\xi = -x_0/\ell$. By integrating Eq. (8) through the Alfvén resonance layer we find what combination of the Ai and Bi functions we have at $\xi \rightarrow \infty$. (Treating the frequency as a Laplace transform variable, we find that causality requires us to go above the singularity at $\xi = 0$ in the complex ξ plane.) We decompose the Ai and Bi functions into a wave incident on the layer $\xi = M^2$ and a reflected wave. We define an amplitude reflection coefficient, R , and from it the fraction, q , of incident power dissipated upon reflection $q = 1 - |R|^2$.

In cases of interest, S and M are small. An analytical estimate of q is then possible. We obtain this by finding series solutions to Eq. (8) about the singular point and by matching these onto the Airy functions at large ξ . In the neighborhood of the singularities, the solutions are (to lowest orders in M and S)

$$E_1 = 1 + \left(\frac{S}{M} + S^2\right) \xi + O(S^4, \xi^3, S^2 \xi^2), \quad (9)$$

$$E_2 = 1 + MS(1 - MS) E_1 \log(\xi) + O(S^4, \xi^3, S^2 \xi^2). \quad (10)$$

[We assume $M = O(S)$.] The appropriate combination of E_1 and E_2 that matches onto $Ai(-\xi)$ in the region $M^2 \ll -\xi \sim \xi_0 \ll 1$ is

$$E = [c_1 - (M/S)c_2]E_2 + [(M/S)c_2 + \log(-\xi_0)(M^2c_2 - MS c_1)]E_1, \quad (11)$$

where $c_1 = Ai(0) = 0.355$ and $c_2 = -Ai'(0) = 0.259$. We write $\log(-\xi_0) = \log|\xi_0| + i\pi$, and then ignore the $\log|\xi_0|$ term since it is smaller than the other real terms. For $M^2 \ll \xi \ll 1$ we then have

$$E = Ai(-\xi) + i(\pi/2\sqrt{3})[M(c_2/c_1)^{1/2} - S(c_1/c_2)^{1/2}]^2 Bi(-\xi), \quad (12)$$

from which we determine the fractional power absorbed,

$$q = (2\pi/\sqrt{3})[M(c_2/c_1)^{1/2} - S(c_1/c_2)^{1/2}]^2. \quad (13)$$

The MHD limit is given by $S \rightarrow 0$. We see that the effect of the finite frequency is either to increase or to decrease q depending on the sign of M . If $S = Mc_2/c_1$ we have $q = 0$.

In Fig. 1 we compare Eq. (13) with the value of q obtained by numerically integrating Eqs. (6) and (7). We see that there is good agreement for $|S|, |M| \leq 0.5$. In particular the line along which $q=0$ is accurately given by Eq. (13).

To illustrate our result let us compute q for PLT taking $k_y = -m/a$, $k_{||} = n/R$, $a = 40\text{cm}$, $R = 1.3\text{m}$ and assuming a parabolic density profile. We find

$$q = 0.1 m^2 \left(\frac{B_{40}}{a_{40} f_{25}} \right)^{4/3} \left(\frac{1}{n_{14}} \right)^{2/3} \left\{ 1 + \left(\frac{a}{R} \right)^2 \frac{n^2}{10m} \left[\frac{f_{25}}{B_{40} n_{14} a_{40}^2} \right]^{1/3} \right\}^2 \quad (14)$$

where B_{40} is the toroidal field in units of 40kG, f_{25} is the frequency of the wave generator in units of 25 MHz, n_{14} is the central density in units of 10^{14} cm^{-3} and a_{40} the minor radius in units of 40cm. Thus if $m = -1$, q is small for $n \approx \pm 10$. With this value of n the wave energy propagates away from the antenna reasonably quickly (since $k_{\perp} \sim k_{\parallel}$) so that the parallel damping length is long. We expect, therefore, that a high-Q eigenmode will be excited. Practically, this allows us to combine the high antenna loading associated with toroidal eigenmodes³ with the Alfvén wave dissipation process.

The physics of the dissipation mechanism has not been discussed here. Inclusion of finite temperature and parallel electric field effects⁴ shows that the dissipated energy is linearly mode-converted⁵ into a thermal wave which propagates into the plasma from the resonance. This wave damps via linear electron Landau damping. Hence the energy absorbed by the shear Alfvén resonance shows up as electron heating near the resonance.

This work was supported by the U. S. Department of Energy Contract No. EY-76-C-02-3073.

REFERENCES

- ¹L. Chen and A. Hasegawa, Phys. Fluids 17, 1399 (1974);
J. A. Tataronis and W. Grossman, Nucl. Fusion 16, 667 (1976);
J. A. Tataronis, J. Plasma Phys. 13, 87 (1975);
J. M. Kappraff and J. A. Tataronis, J. Plasma Phys. 18,
209 (1977).
- ²E. Ott, J.-M. Wersinger, and P. T. Bonoli, Laboratory of
Plasma Studies Report LPS 236, Cornell University (March, 1978).
- ³J. Adam et al., in Plasma Physics and Controlled
Nuclear Fusion Research 1974 (IAEA, Vienna, 1975) p. 65.
- ⁴A. Hasegawa and L. Chen, Phys. Fluids 19, 1924 (1976).
- ⁵T. H. Stix, Phys. Rev Lett. 15, 878 (1965).

Fig. 1. Contours of constant $\sqrt{q}$. (a) Results obtained by numerically integrating (6) and (7). (b) Analytic result, (13).

782194