TECHNICAL REPORT**MONOCHROMOSOMAL HYBRIDS FOR THE ANALYSIS OF THE HUMAN GENOME****RAGHBIR S. ATHWAL, PH.D.****Department of Microbiology and Molecular Genetics
New Jersey Medical School****PROJECT SUMMARY**

The objective of this research project is to transfer individual human chromosomes to mouse and/or Chinese hamster cells to produce hybrid cell lines each harboring a single different human chromosome. The human chromosome present in rodent cells will be marked with a dominant selectable marker and maintained by selection.

The production of monochromosomal hybrids is achieved by microcell mediated transfer (MMCT) of single human chromosomes to rodent cells. In these experiments chromosomes are first "tagged" with a selectable marker in human cells and marked chromosomes are subsequently transferred to rodent cells by microcell fusion. Several different experimental schemes have been developed to 'tag' human chromosomes with a selectable marker. Amphotropic retroviral vectors provide a highly efficient system to introduce selectable markers into normal diploid human cells. The integration of a retroviral vector into the cell genome occurs by recombination at a defined nucleotide sequence in the LTRs and only a single copy of the vector integrates in a cell. This property of retroviral vectors allows one to isolate a segment of the chromosomal DNA flanking the vector integration site by PCR amplification.

In the studies described in this project, the amphotropic retroviral vector pZIPgpt, carrying the dominant selectable marker gpt, is used to tag the human chromosomes in normal diploid cells. Human DNA flanking the integrated vector is rescued by PCR amplification and cloned into a plasmid vector. Cloned human DNA is then used to probe Southern blots of DNAs from a panel of hybrid cell lines to identify the chromosome of its origin. This allows us to identify clonal human cell lines, each carrying the marker integrated into a different chromosome. Marked chromosomes are then transferred to rodent cells by MMCT. Identification of the marked human chromosome in human cells prior to its transfer into rodent cells will eliminate the redundancies in the recovery of the chromosomes in hybrid cells and expedite the completion of our hybrid cell panel. Once a chromosome is rescued in mouse cells, it can be transferred to Chinese hamster and/or other cell lines of interest.

Using the proposed experimental approach, we have already produced monochromosomal hybrids for 16 different human chromosomes which include 1,2,5,6,7,8,9,12,13,14,15,17,19,21 and X. Each hybrid cell line is analyzed by cytogenetic and molecular methods to identify and determine the integrity of the human chromosome.

A panel of monochromosomal hybrid cell lines will provide an invaluable biological resource for complementation analysis, cloning of chromosome specific DNA fragments, physical mapping of human chromosomes and fractionation of the human genome. The DNAs from hybrid cell lines will be made available freely for studies on the analysis of the human genome and gene mapping.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

A. OBJECTIVE

The overall objective of our project is to transfer individual human chromosomes to rodent cells to generate a panel of hybrid cell lines each harboring a single different human chromosome (monochromosomal hybrid cell lines). The human chromosome present in rodent cells will carry a dominant selectable marker, *Ecogpt*, and will be retained by selection. We have already produced such hybrid cell lines for 16 different human chromosomes (Table 1). The future goals for the continuation of this program are:

1. Complete two panels of monochromosomal hybrid cell lines (i.e. mouse/human and Chinese hamster/human) to represent all human chromosomes.
2. Analysis of all hybrid cell lines for chromosome identification and structural integrity. Alu-PCR - Southern blot method will be used to detect the presence of non-selectable chromosomes or chromosomal segments. In situ hybridization of Alu-PCR products from hybrid cell lines to metaphase spread of normal human cells will be used to determine the structural integrity of the transferred chromosome.
3. Transfer human chromosomes to Marsupial and/or Muntjac cells to generate Marsupial/human and/or Muntjac/human monochromosomal hybrid cell lines.

Marsupial and Muntjac cells have a diploid chromosome complement of $2n=14$ and $2n=6$ respectively. The chromosomes of these species are much larger as compared to those of human. A human chromosome present among Marsupial or Muntjac chromosome can be easily separated by unit gravity (lg) sedimentation method or flow sorting. These monochromosomal hybrids will be extremely useful to fractionate the human genome to construct chromosome specific large insert DNA libraries in a YAC vector.

4. An extension of our program will be to generate a set of radiation hybrids for human chromosome 2 each containing different segment of this chromosome. These hybrids will be used to construct a physical map of chromosome 2.

Monochromosomal hybrids are used to map genes on the human chromosomes and to construct a physical maps. The material produced in this project will serve as a national resource for various laboratories in the human genome project and are freely available to the research community. Hybrid cell lines for several chromosomes have already been made available to other laboratories for gene mapping studies.

B. PROGRESS REPORT

Since the inception of this project, we have produced monochromosomal hybrids for 16 different human chromosomes. We have also improved our methods and develop new techniques which will enhance our ability to fulfill our commitment to construct panels of hybrid cell lines. During this budget period (i.e. 1992-1993), we have produced 6 new hybrids which include chromosomes 8, 12, 14, 16, 19 and X. At the rate the project is now proceeding, we anticipate that the hybrid panels will be completed during this grant period. In addition, we have developed new techniques for the analysis of hybrid cell lines. Following is a summary of our progress to date:

1. **Monochromosomal Hybrids Produced to Date:**

The experimental approach used to produce monochromosomal hybrid cell lines is given in figure 1 and hybrids produced to date are listed in Table 1.

A retroviral vector carrying the selectable marker Ecogpt was introduced into the normal diploid human fibroblast cell line GM03468 and a total of 210 independent gene transfer clones were isolated by selection in MX medium. Normal human cells have a limited life span. In order to perpetuate the marked human chromosomes, gene transfer clones were fused with an immortal human tumor cell line HT1080TG (Fig. 1). In these studies, a total of 125 human x human hybrid cell lines have been produced.

As an intermediary step, clones of marked diploid human cells were also simultaneously hybridized with mouse A9 cells to generate mouse/human somatic cell hybrids (Fig.1). Mouse/human hybrids are stored frozen while human/human hybrid clones are analyzed to identify the marked chromosomes. Once a marked chromosome is identified in human cells, the corresponding mouse/human hybrid cell line is used to transfer the marked chromosome to another recipient cell line by microcell fusion.

Table 1 : Monochromosomal hybrids containing marked human chromosomes.

Mono. Hybrids in Mouse cells	Human Chromosome	No. of Independent Hybrid Cell Lines
RA-1	1	2
RA-2	2	5
RA-5	5	3
RA-6	6	4
RA-6q	6q	1
RA-Iso 6q	Iso 6q	1
RA-7	7	1
RA-8	8	2
RA-9	9	1
RA-12	12	2
RA-12q	12q	1
RA-13	13	2
RA-14	14	2
RA-15	15	2
RA-16	16	3
RA-17	17	3
RA-19	19	2
RA-21	21	1
RA-X	X	2
Hamster/Human Mono. Hybrids	Human Chromosome	
CHA-2	2	
CHA-5	5	
CHA-6q	6q	
CHA-6	6	
CHA-13	13	
CHA-15	15	

Mouse cell lines in general are easy to induce micronuclei formation. In addition, segregation of unmarked human chromosomes in whole cell hybrids facilitate the transfer of single chromosomes. Thus, mouse/human whole cell hybrid generation is an essential intermediary step for the production of monochromosomal hybrids. Since our mission is to generate a panel of monochromosomal hybrid cell

lines with marked chromosomes, hybrid cell lines containing unmarked single chromosomes and hybrids with two human chromosomes are not included in Table 1.

Once a chromosome is rescued into mouse cells, it can be easily transferred to any other cell type of interest. We have already transferred several chromosomes to Chinese hamster cells to generate hamster/human monochromosomal hybrid cell lines (Table 1).

2. New Techniques Developed

a. Identification of the Marked Chromosomes In Human Cells: (Fig.2)

In the production of monochromosomal hybrid cell lines, we have been dependent upon fortuitous recovery of different chromosomes in microcell hybrids. Thus, hybrid cell lines for some chromosomes were obtained in duplicate or even triplicate while others were never recovered. The redundancy in the production of microcell hybrids can be reduced or even eliminated if the marked chromosomes can be identified in human cells prior to their transfer to mouse cells.

We have developed a new method using PCR technology to determine the identity of the marked chromosome in human cells. This will allow us to generate a bank of 24 human cell lines each carrying the selectable marker integrated into a different chromosome. The experimental strategy outlined in Fig.2 involves PCR mediated amplification and cloning of the human DNA flanking the integrated vector. The cloned DNA is then mapped to a chromosome by Southern blot analysis of a panel of rodent/human hybrid cell lines.

The rationale for the cloning of the junction sequence is based upon the recombination dependent integration of the retroviral vectors into cellular DNA involving a precise nucleotide sequence in LTRs (Fig.2). Cellular DNA, isolated from the marked cells is digested with a frequent cutting restriction endonuclease which has unique sites in vector LTRs. As depicted in Fig.2, SAU3A1 cuts each LTR into two segments resulting in four DNA fragments. In two of these fragments, part of the LTR sequence is linked to an internal vector segment while in the other two fragments, part of the LTR is linked to the human cellular DNA flanking the integration site. Self ligation of the digested DNA, under conditions that promote circularization, yield unique DNA circles each containing a part of the LTR sequence. PCR amplification of ligated DNA with a set of oligonucleotide primers complementary to the end of 3' LTR yield DNA products comprised of 79bp vector DNA fragment and a human DNA fragment linked to a part of the LTR sequence. The human DNA product of PCR amplification is separated by gel electrophoresis and cloned into a plasmid. Cloned human DNA is used to probe the Southern blots of a panel of hybrid cell lines to determine the chromosome of its origin. Marked chromosomes identified in this manner are subsequently transferred to mouse cells by microcell fusion. The method will significantly expedite the production of monochromosomal hybrids. A manuscript describing the results of these experiments is currently being prepared for publication.

b. Analysis of Hybrid Cell Lines:

The usefulness of monochromosomal hybrids depends upon the fact that only a single intact "tagged" chromosome is present in each cell line. Presence of chromosomal fragments other than the marked chromosome or rearranged chromosomes would lead to erroneous interpretation of the data obtained with these hybrids. Thus, appropriate evaluation of each hybrid cell line is essential to ascertain the purity of the hybrid cell lines. Therefore, all hybrid cell lines are analysed by several different methods.

We have developed a highly sensitive method to determine human chromosomal content of hybrid cells. In this method a ^{32}P labeled probe, prepared for the Alu-PCR amplified DNA products of a new hybrid, is hybridized to the Southern blots of Alu-PCR amplified DNA from a hybrid cell panel (Fig. 3). Hybridization in this case occurs only with the DNA of the hybrid containing a corresponding human chromosome. As such, the method allows one to detect the presence of multiple chromosomes or chromosomal fragments in putative monochromosomal hybrid cell lines. All of the hybrids produced in our lab are subjected to this type of analysis. It helps to identify hybrids containing non-selectable chromosomal fragments or rearranged chromosomes. An Alu-PCR Southern blot that shows the presence of human chromosome 6 in a monochromosomal hybrid is shown in Fig. 3.

All of the hybrid cell lines are also analyzed by in situ hybridization using a total human DNA probe for chromosome painting to detect the presence of chromosomal fragments other than the marked chromosomes. Our results show that each of these hybrids contained only one human chromosome and no additional chromosomal fragments were observed.

c. Fingerprinting Human Chromosomes:

Chromosome identification in hybrid cells is generally achieved by cytogenetic or molecular analysis for the presence of a limited number of markers. Both of these methods are inadequate when subchromosomal fragments or rearranged chromosomes are present in the hybrid cells. We have developed an approach for fingerprinting individual human chromosomes which allows the assignment of PCR amplified DNA fragments to the regions of a chromosome. A fingerprint map developed for a specific chromosome can also be used to identify chromosomal segments present in radiation-hybrids originating from a monochromosomal hybrid. A description of the method and its utility in gene mapping has been published in Genomics and a reprint is appended with this report.

d. Chromosome "Tagging" by Homologous Insertion

We have developed a plasmid vector that carries human repeat sequences and the selectable markers NEO and HSV-TK. The vector when introduced into mouse/human hybrid cell lines, preferentially integrates into the human chromosome by homologous recombination. However, in addition to homologous insertion, random integration of the vector into mouse chromosomes also occurs. The vector is designed so that homologous insertion leads to the exclusion of HSV-TK gene while retaining the neo marker. On the other hand non-homologous insertion of the vector retains both NEO and TK genes. The cells expressing the neo marker can be selected by growth in medium containing G418. Addition of Gancyclovir (GANC) into the medium selects against HSV-TK. Thus, hybrid cells in which the neo marker is integrated into a human chromosome by homologous insertion can be selectively separated by growth in medium containing G418 and GANC. We have used this method to insert neo marker into human chromosome 9 in hybrid RA-9. Homologous insertion into human chromosome 9 occurred in 5% of the clones. We are using this method to insert a second marker into the human chromosomes already marked with gpt or to 'tag' unmarked human chromosomes present individually in hybrid cells. The presence of two or multiple markers in a human chromosome will provide a method to generate subchromosomal hybrids by double selection (i.e. for the retention of a marker and loss of the second marker) following X-irradiation. This approach will also expedite completion of hybrid cell panel.

3. PUBLICATIONS

The results of our research towards the production and use of the monochromosomal hybrids were presented in several conferences and published in a number of papers. A list of publications and meeting abstracts during this budget period is given here and a copy of each is enclosed with this report.

a. Papers Published or Accepted

1. Jeggo, P., M. Hafezparast, A.F. Thompson G.P. Kaur, A.K. Sandhu and R.S. Athwal (1992). Hamster-human hybrid cell lines bearing different fragments of human chromosome 2. *Somatic Cell & Mol. Genetics* (In Press).
2. Kaur, G.P. and R.S. Athwal (1992). Complementation of DNA repair defect in XP-C by transfer of human chromosome 5. *Somatic Cell and Molecular Genetics* (In Press).
3. Jeggo, P.A., M. Hafezparast, A.F. Thompson, B.C. Broughton, G.P. Kaur, M. Zdzienicka, and R.S. Athwal (1992). Localization of a DNA repair gene (XRCC5) involved in double strand break rejoining to human chromosome 2. *Proc. Natl. Acad. Sci. U.S.A.* 89:64-23-6427.
4. Sidhu, M.S., B. Helen, and R.S. Athwal (1992). Fingerprinting human chromosomes by PCR mediated DNA amplification. *Genomics* 14-:728-732.
5. Kaur, G.P., A.Rinaldy, R.S.Lloyd, and R.S. Athwal (1992) A gene that partially complements defects in Xeroderma pigmentosum group A cells maps to human chromosome 8. *Somatic Cell and Mol. Genet.* 18:4:371-379.
6. Mori, T., T.L. Rinaldy, R.S. Athwal, G.P. Kaur, O. Nikaido, R.S. Lloyd and A. Rinaldy (1992). An XP complementation group A related gene: Confirmation using monoclonal antibodies against the Cyclobutane dimer and (6-4) photoproduct. *Mutat. Res.* 293:2:143-150.

b. Manuscripts in Preparation:

1. Athwal, R.S., A.K. Sandhu and G.P.Kaur. Monochromosomal hybrids for the genetic analysis of the human genome: Integration of a selectable marker and transfer of individual human chromosomes to rodent cells.
2. Kaur, G.P., A.K. Sandhu, K.D. Dobler, R. Day III and R.S. Athwal. Transfer of individual human chromosomes into tumor cells: Identification of a suppressor gene on chromosome 9 for malignant glioma.
3. Sandhu, A.K., G.P. Kaur, K. Hubbard-Smith, K. Jha, H.L. Ozer and R.S. Athwal. Suppression of immortalization in SV40 transformed human fibroblasts by the transfer of human chromosome 6.
4. Sidhu, M.S., A.K. Sandhu, and R.S. Athwal. Mapping of Vector integration sites in human genome and transfer of the marked chromosomes to rodent cells by microcell fusion.

c. Abstracts of Presentations In Scientific Meetings:

1. Kaur, G.P. and R.S. Athwal (1992) Identification of a DNA repair gene on human chromosome 5 that complements Xeroderma Pigmentosum cells of group C. Amer. J. Hum. Genet. Suppl.51(4):192.
2. Bagga, P.S. and R.S. Athwal (1992) Complementation of DNA repair defect in Xeroderma Pigmentosum and Cockayne Syndrome Cells by bacteriophage T4 gene denV. Amer. J. Hum. Genet. Suppl.51(4):212.
3. Sandhu, A.K., G.P. Kaur, M.S. Kansara and R.S. Athwal (1991) Transformation of Normal Human Fibroblasts: Vector mediated disruption of suppressor gene Proc. Inter. Cong. Hum. Genet. Washington, D.C. Amer. J. Hum. Genet. 49 (Suppl): 4:455 (Abs#2587).
4. Sidhu, M.S. Helen, B.K. and Athwal, R.S. (1991). Fingerprinting human chromosomes by PCR mediated DNA amplification. Int. Cong. Human Genet. Washington D.C., Amer. J. Hum. Genet. 49 (Suppl.): 4:293 (Abs#1627).
5. Sandhu, A.K., G.P. Kaur, K. Hubbard-Smith, K.K. Jha, H.L. Ozer and R.S. Athwal (1991). Suppression of immortalization in human fibroblasts by the transfer of human chromosome 6. Int. Cong. Human Genet., Washington, D.C.. Amer. J. Hum. Genet. 49 (Suppl): 4:457 (Abs #2601).
6. Kaur, G.P., Sandu, A.K., Dobler, K.D., Day III, R. and Athwal, R.S. (1991). Suppression of transformed phenotype in human malignant glioma by microcell mediated transfer to chromosome 9. Proc. Int. Cong. Human Genet., Washington, D.C.. Amer. J. Hum. Genet. 49 (Suppl): 4:452 (Abs#2568).
7. Kaur, G.P., Bagga, P.S. Rinaldy, A., Lloyd, R.S. and Athwal, R.S. (1991). Two gene hypothesis for DNA repair defect in XP-A: Identification of second gene on human chromosomes 8. Int. Cong. Human Genet. Washington D.C., Amer. J. Hum. Genet. 49 (Supp.): 4:409 (Abs. #2314).

4. Conclusion:

In conclusion, we have already produced monochromosomal hybrids for 2/3 of the human genome. In addition, we have generated sufficient biological materials to complete the proposed panels of hybrid cell lines.

We have developed experimental procedures to identify marked chromosomes in human cell lines prior to their transfer to rodent cells. This would eliminate redundancy in the production of monochromosomal hybrids and therefore help expedite completion of the hybrid cell panels. We have also developed a highly sensitive method to identify human chromosomes in hybrid cells.

Monochromosomal hybrids produced in our lab are used in a number of laboratories for experiments on gene mapping, gene isolation, chromosome fractionation and genetic analysis for complementation of cellular phenotypes such as DNA repair and regulation of cell growth. Monochromosomal hybrid cell lines are freely available to scientific community for experiments on gene mapping and analysis of the human genome. We are preparing large quantities of DNA from each hybrid cell line which will be available to the research community for various experiments.

FIGURE 1

EXPERIMENTAL APPROACH TO GENERATE MONOCHROMOSOMAL HYBRID CELL LINES

FIGURE 1

FIGURE 2

schematic representation of LTR-PCR amplification of chromosomal DNA sequences flanking integrated vector. Amplified Human DNA is used to probe the Southern blots of hybrid cell panel.

Fig. 3

Autoradiograph of a Southern blot showing hybridization of Alu-primed PCR product of a microcell hybrid RA6 to a panel of somatic cell hybrids. Probe hybridizes only to hybrid containing chromosome 6. Therefore RA6 does not contain fragments from other chromosomes.

END

DATE
FILMED

2 / 22 / 93