

MASTER

**A Model for the Movement and
Distribution of Fish in a
Body of Water**

D. L. DeAngelis

ENVIRONMENTAL SCIENCES DIVISION
Publication No. 1173

OAK RIDGE NATIONAL LABORATORY
OPERATED BY UNION CARBIDE CORPORATION · FOR THE DEPARTMENT OF ENERGY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

27
6-15-78
25 D. DeAngelis

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Printed in the United States of America. Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
Price: Printed Copy \$6.00; Microfiche \$3.00

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, contractors, subcontractors, or their employees, makes any warranty, express or implied, nor assumes any legal liability or responsibility for any third party's use or the results of such use of any information, apparatus, product or process disclosed in this report, nor represents that its use by such third party would not infringe privately owned rights.

Contract No. W-7405-eng-26

A MODEL FOR THE MOVEMENT AND DISTRIBUTION
OF FISH IN A BODY OF WATER

D. L. DeAngelis

ENVIRONMENTAL SCIENCES DIVISION
Publication No. 1173

DATE PUBLISHED - JUNE, 1978

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37830
operated by
UNION CARBIDE CORPORATION
for the
DEPARTMENT OF ENERGY

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

ACKNOWLEDGEMENTS

I wish to thank Dr. C. C. Coutant for his encouragement and help in this work. I also thank Drs. J. S. Suffern and R. H. Gardner, who have offered many useful criticisms. This work was sponsored by the Division of Biological and Environmental Research, U.S. Department of Energy, under contract W-7405-eng-26 with Union Carbide Corporation.



THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

ABSTRACT

DEANGELIS, D. L. 1978. A model for the movement and distribution of fish in a body of water. ORNL/TM-6310. Oak Ridge National Laboratory, Oak Ridge, Tennessee. 78 pp.

A Monte Carlo mathematical model tracks the movement of fish in a body of water (e.g., a pond or reservoir) which is represented by a two-dimensional grid. For the case of a long, narrow reservoir, depth and length along the reservoir are the logical choices for coordinate axes. In the model, it is assumed that the movement of fish is influenced by gradients of temperature and dissolved oxygen, as well as food availability and habitat preference. The fish takes one spatial "step" at a time, the direction being randomly selected, but also biased by the above factors.

In trial simulations, a large number of simulated fish were allowed to distribute themselves in a hypothetical body of water. Assuming only temperature was influencing the movements of the fish, the resultant distributions are compared with experimental data on temperature preferences.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGMENTS	iii
ABSTRACT	v
LIST OF TABLES	ix
LIST OF FIGURES	xi
GENERAL DESCRIPTION OF THE MODEL	4
MATHEMATICAL DESCRIPTION OF THE MODEL	9
COMPUTER PROGRAM	16
Part A. Input Cards	19
Part B. Output	24
Part C. Computer program listing	24
TRIAL SIMULATIONS	28
Fish movements	28
Fish distribution patterns	28
SUMMARY	37
REFERENCES	43
APPENDIX	45

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	The Program Variables	29
2	Parameter Values Used in the Example shown in Figure 7	34
3	Changes in parameter values from Table 2 relevant to the example shown in Fig. 11	41

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1	A hypothetical reservoir. The vertical dimension is depth in meters (disproportionately scaled), and the horizontal dimension is length along the reservoir in kilometers, with the downstream dam at the left. Isotherms in degrees Centigrade (solid lines) and dissolved oxygen isobars in parts per thousand (dotted lines) are sketched in. The shaded region denotes high food availability. A power plant is assumed located at the upstream end of the reservoir 5
2	The point (i,j) in a grid of points, with the adjacent points to which the fish can move in one step 7
3	A grid of points representing a portion of the reservoir. The shaded circles are water, while the open circles are above the water surface. The shaded circles surrounded by larger circles represent points along the preferred temperature isotherm 8
4	A plot of food availability to fish in a hypothetical reservoir. The peaks and plateaus represent the regions of high food availability 18
5	Input data for a sample trial simulation as it appears on the data cards 20
6	Input data for a sample trial simulation as it is printed out by the computer program 25
7	Plot of simulated motions of two fish initially placed at points A and B. The assumed preferred temperature is $TEMP_p = 29.0^{\circ}C$ and the force of temperature attraction, P_T , is 1.0 for case A and 50.0 for case B 26
8	The distribution of 500 fish influenced only by temperature in the reservoir after 200 steps. The assumed preferred temperature is $29.0^{\circ}C$. Other parameters of the model are given in Table 2 27

<u>Figure</u>	<u>Page</u>
9 Histogram of percent distribution of fish in Fig. 8 about the preferred temperature of 29.0°C	38
10 Histogram of relative frequency of largemouth bass in ambient water temperatures during daytime (from Reynolds and Casterlin 1977)	39
11 The distribution of 500 fish influenced by temperature, dissolved oxygen, food availability and habitat favorability in the reservoir after 200 steps. Parameter values are given in Table 3	40

INTRODUCTION

The distribution of fish populations in bodies of water is interesting to sportsmen, commercial fishermen, and ecologists alike. Several factors that may influence fish movements and spatial population distribution have been proposed, including temperature, dissolved oxygen in the water, pH values, the availability of food, the presence of cover for protection from predators, and the occurrence of competitors. These are not all independent. Dissolved oxygen is to some extent related to water temperature, as is the availability of certain types of prey. If the locomotor responses of fish to each of the factors were known in detail, then one could feasibly predict the average motions of a fish in a given body of water. The task of identifying and quantifying all the influences on fish locomotor behavior will not be easy, but significant progress has been made, thanks to ingenious laboratory experiments and telemetry methods useful for the field.

As the factors involved in the spatial behavior of fish begin to be understood, it can be applied to a host of practical matters. For example, one would like to know where in a body of water fish population densities will be highest at a given time of year. Also, how will the population distribution in space respond to slow or rapid changes in the condition of the water, either through natural processes such as seasonal variations, or artificial changes such as those induced by power plant operations?

Both basic research and practical applications in the area of fish movements will rely on techniques of mathematical modeling. Models incorporating specific hypotheses will form a framework for experimental research, from which the data can be used to test the hypotheses. When the fundamental parameters of the models have been quantified, the model can be used predictively. This report describes a mathematical model capable of being used in conjunction with laboratory experiments and field studies, and later, for predictive purposes.

Much experimental research has gone into the study of the effects of temperature on locomotor behavior in fishes. Temperature has been called the most important influence on the behavior of many freshwater fish (e.g., Coutant 1975). It has long been noticed that fish move to different areas of a body of water as water temperature changes. For example, largemouth bass overwinter in deep water, where the temperature is warmest. Using underwater telemetry, Warden and Lorio (1973) found that largemouth bass tend to move great distances to new home ranges in spring and fall, when water temperature is changing most rapidly. In winter, the population of largemouth bass congregates around the thermal discharges of power plants (Gibbons, Hook and Forney 1972). In a Texas cooling reservoir, it was noticed that largemouth bass sought out the cooler shoreline zones in summer mornings when the remainder of the reservoir had temperatures exceeding 37.8°C (Smith 1972).

Laboratory studies have been performed to refine the data on temperature selection of several centrarchid species (Reynolds and Casterlin 1976, Stuntz and Magnuson 1976). Researchers have also sought to relate temperature preference with thermoregulation and the optimization of physiological processes (e.g., McCauley and Huggins 1976, Reynolds and Casterline 1976). Growth rates of largemouth bass usually seem to be optimal near their temperature preference (Coutant and Cox 1976), although this does not seem to be the case for bluegills in thermal discharge areas during the summer months (Kitchell et al. 1974). Bluegills were shown to actively avoid lethal temperatures (Peterson and Schutsky 1976), and to vary their temperature preference according to their daily rations (Stuntz and Magnuson 1976).

A question that has bearing on attempts to model fish movements is what is the precise mechanism by which fish tend to center around their preferred temperatures? Neill (1976) discusses different mechanisms in detail and describes one-dimensional computer models based on some of these mechanisms. Thermoregulatory movements can be broadly categorized as predictive or reactive. In the former case, the fish is assumed to have some knowledge, by prior experience or instinct, of the

temperature distribution in the body of water, and will use this knowledge to move toward the desired temperature range. For example, since lower water strata are normally cooler than upper layers, the fish should automatically move downwards when it feels too warm. Reactive behavior presupposes no prior knowledge of the temperature distribution, but only that the fish responds to different temperature regimes by altering its locomotory behavior. Several models of reactive movements have been developed. One type of model has been termed orthokinetic by Fraenkel and Gunn (1961). According to this model, fish slow their movements when in the preferred temperature range, increasing their chances of staying there. Both Fraenkel and Gunn (1961) and Neill (1976) have pointed out the inefficiency of this model for producing aggregation about the preferred temperature. Fish whose direction of motion was originally oriented away from the preferred temperature would continue to move away from it. Neill was able to obtain realistic aggregation only when his model specified a high probability of changing directions when the fish was moving away from the preferred temperature range. This form of behavior is called klinokinesis.

Dissolved oxygen and pH in the water are important to the health of the fish and, therefore, presumably influence its movements. While fish have not been shown to exhibit dissolved oxygen and pH preferences, they might be expected to avoid unfavorable conditions. For example, at 25°C the minimum oxygen requirement of small largemouth bass is almost 0.92 ppm (Moss and Scott 1961); it would be advantageous for such fish to preferentially move away from areas with dissolved oxygen levels below this minimum.

The movement of fish in response to food availability and habitat preference probably involve learning where favorable conditions exist in a body of water. It is harder to develop models for response to these factors than it is for motion in temperature gradients, since it is difficult to know the extent of learning in the fish.

The model described in this report assumes that the fish acts as if it can sense temperature gradients and will move along a temperature

gradient in the direction of its preferred temperature. We do not specify whether the fish acts this way because it actually can perceive temperature gradients or because its klinokinetic activity increases as it moves into less preferable temperature ranges. On the scale length we are dealing with (meters vertically and kilometers horizontally), the precise mechanisms of motion on the small scale may be unimportant. We also assume that the fish will move away from dissolved oxygen levels below that which is the minimum tolerable, and that they will have a general tendency to move toward areas of greater available food and more favorable habitat. These several influences can either reinforce each other or, to some extent, cancel each other under particular circumstances. Aside from these basic assumptions, the model is very general and can be parameterized to suit a variety of situations.

The present model is offered not as a description of the way fish behave, but as a device by which a variety of hypothetical descriptions of locomotor behavior can be tested. A few examples are given to illustrate the way in which the model is used. More thorough exploration of the model will be undertaken later, in combination with field studies.

GENERAL DESCRIPTION OF THE MODEL

The intent of this model is to predict the average spatial distribution of a fish population in a closed body of water. To do this we simulate the movements of individual fish, allowing a large number of fish to start from random positions in the body of water, and to move for a certain period of time. We assume that a small number of factors influence the movements of the fish; temperature, dissolved oxygen, food availability and habitat preference.

The model is designed to apply to a two-dimensional representation of a hypothetical reservoir (Fig. 1). The two dimensions are depth and either length along the reservoir or width across a cross section. A three-dimensional representation would be preferable, but would pose




Fig. 1. A hypothetical reservoir. The vertical dimension is depth in meters (disproportionately scaled), and the horizontal dimension is length along the reservoir in kilometers, with the downstream dam at the left. Isotherms in degrees Centigrade (solid lines) and dissolved oxygen isobars in parts per thousand (dotted lines) are sketched in. The shaded region denotes high food availability. A power plant is assumed located at the upstream end of the reservoir.

problems both computationally and graphically. It is hoped that this model will eventually be extended to three dimensions, but the present two-dimensional model is useful. Note that the scaling in the vertical (depth) dimension is greatly exaggerated relative to the horizontal coordinate. Typical temperature and dissolved oxygen isoclines are sketched in, and the area in which food availability is greatest (usually the shallow water along shore lines) is shaded. We assume that the position of those factors are stable over the time scale in which a fish can move considerable distances. A typical fish will have a preferred range of temperatures, will tend to avoid very low levels of dissolved oxygen, will be attracted by high food availability, and will prefer habitats that give it sufficient cover from predators. On this basis, the average distribution of a model fish population may be reliably predicted, though the path of a given fish is unique.

For modeling purposes, it is necessary to represent the two-dimensional space by a grid of points. Consider a fish located at some point (i,j) in the grid points (Fig. 2). The fish can move to one of eight adjacent points $(i+\delta, j+\epsilon)$, where δ and ϵ take on the values $-1, 0$ and $+1$ (but both cannot be 0 simultaneously). It is assumed that the following factors influence the next location of the fish:

1. The tendency of the fish to continue moving in the general direction in which it is already moving. This can be termed the "forward inertia" of motion.
2. The preferred temperature of the fish and the temperature at the present location of the fish, (i,j) , and the eight surrounding points.
3. The location of food supplies and cover.
4. The boundary of the water body, which sets limits on the motion of the fish.

These factors can be elucidated to some extent by examination of Fig. 3. Assume the fish is located at point (i,j) and has just moved from the point $(i,j-1)$. The black points in this figure are those in the body of water while the white dots are above its surface. The isotherm of the preferred temperature is represented by black dots

ORNL-DWG 77-2861




Fig. 2. The point (i, j) in a grid of points, with the adjacent points to which the fish can move in one step.

ORNL-DWG 78-1682




Fig. 3. A grid of points representing a portion of the reservoir. The shaded circles are water, while the open circles are above the water surface. The shaded circles surrounded by larger circles represent points along the preferred temperature isotherms.

surrounded by a circle. The most likely next "step" of the fish is to the point $(i,j+1)$, since this is in its direction of preferred temperature as well as its direction of inertia. The fish also has a high probability of moving to point $(i-1,j+1)$. Of course, the fish cannot move to points $(i+1,j-1)$, $(i+1,j)$, or $(i+1,j+1)$ because these lie above the surface of the water.

It is conceptually and mathematically advantageous to discuss fish movements in terms of the four factors listed above, but these factors have not been quantified in detail (except for factor 4; the fish we are dealing with cannot normally leave the water). Data are available on the response of some fish species to temperature and dissolved oxygen variations, but other factors, such as food availability and habitat preferences, complicate the situation in natural bodies of water, making predictions based on mathematical models less reliable.

MATHEMATICAL DESCRIPTION OF THE MODEL

It is convenient to represent the probability of a fish moving one step from a point (i,j) to another (k,m) in a two-dimensional grid as an element of a transition matrix, $P_{ij,km}$. Since the fish can move from one grid point only to an adjacent one in a single step, k and m are constrained as follows:

$$k = i + \delta \quad (\delta = -1,0,+1) \quad (1a)$$

$$m = j + \epsilon \quad (\epsilon = -1,0,+1), \quad (1b)$$

(see Fig. 1). In all future discussion, k and m will be implicitly subject to the limitations (1a,1b).

The sum over all probabilities for direction of motion must equal unity:

$$\sum_{k=i-1}^{i+1} \sum_{m=j-1}^{j+1} P_{ij,km} = 1.0 \quad (2)$$

The model is event-oriented, where an event is a step in space. This means that, given a fish initially at point (i,j) , the next moment of interest occurs only when the fish has moved to an adjacent grid point. Therefore, the probability of the fish being in its same position at the next locomotory event in the model is identically zero, or

$$P_{ij,ij} = 0.0. \quad (3)$$

All of the transition elements together define a transition matrix, \underline{P} . Let $\underline{X}(1)$ be the probability vector for the position of the fish at a given moment. The elements of $\underline{X}(1)$, which are $x_{ij}(1)$, represent the probabilities of the fish being located at any given point (i,j) . The condition

$$\sum_{i=-\infty}^{+\infty} \sum_{j=-\infty}^{+\infty} x_{ij}(1) = 1.0 \quad (4)$$

must hold since the fish must be somewhere in the water body. Then

$$\underline{X}(2) = \sum_{j=m-1}^{m+1} \sum_{i=k-1}^{k+1} x_{ij} P_{ij,km} = \underline{P} \cdot \underline{X}(1) \quad (5)$$

is the probability matrix for the position of the fish after its next movement to a new grid point.

If the movement of the fish from one grid point to the next is purely random (i.e., "random walk"), then

$$P_{ij,km} = 1.0/8.0 = 0.125 ; \quad (6)$$

that is, there is an equal probability of 0.125 of the fish going to any of the eight adjacent points. However, the motion of the fish is biased by its forward inertia, temperature and dissolved oxygen gradients, the location of food and favored habitat, and boundaries of the body of water.

Consider first only the influence of forward inertia. It introduces a directional bias on top of random motion. The transition probability can be written

$$P_{ij,km} = \{1.0 + I(k,m)\} / \xi , \quad (7)$$

where ξ is the normalization factor,

$$\xi = \sum_{k=i-1}^{i+1} \sum_{m=j-1}^{j+1} P'_{ij,km} , \quad (8)$$

and

$$P'_{ij,ij} = 0.0 \quad (9a)$$

$$P_{ij,km} = 1.0 + I(k,m) \quad (m \neq j, \text{ if } k = i) . \quad (9b)$$

The term $I(k,m)$ is a measure of the strength of forward inertia relative to random effects in determining the next grid point in the fish's course of movement. If $I(k,m) \ll 1.0$, then the random effects dominate the movement. On the other hand, if, say, $I(i+1,j+1) \gg 1.0$ and $I(i+1,j+1) \gg I(k,m)$ for all seven other pertinent values of k and m , then the fish is likely to move upward and to the right on its next step. The magnitude of $I(k,m)$ for particular values of k and m depends on the past motion of the fish. For this reason, \underline{P} is not a Markov process matrix.

In a similar manner, the effects of temperature and dissolved oxygen can be incorporated into this mathematical scheme. If $T(k,m)$, $DO(k,m)$, $F(k,m)$ and $H(k,m)$ represent the strengths with which temperature gradients, dissolved oxygen gradients and gradients in distribution of food availability and habitat desirability, respectively, then one can write

$$P_{ij,km} = \{1.0 + I(k,m) + T(k,m) + DO(k,m) + F(k,m) + H(k,m)\} / \xi \quad (10)$$

where ξ is defined by Eq. (8) and now

$$P'_{ij,km} = 1.0 + I(k,m) + T(k,m) + DO(k,m) + F(k,m) + H(k,m) \quad (11)$$

The effects of the boundary of the body of water on fish movement is incorporated as follows. Define $B(k,m)$ as the boundary factor, and now write $P_{ij,km}$ as

$$P_{ij,km} = \{1.0 + I(k,m) + T(k,m) + DO(k,m) + F(k,m) + H(k,m)\} B(k,m) / \xi, \quad (12)$$

where ξ is defined by Eq. (8) and now

$$P'_{ij,km} = \{1.0 + I(k,m) + T(k,m) + DO(k,m) + F(k,m) + H(k,m)\} B(k,m), \quad (13)$$

where

$$B(k,m) = \begin{cases} 1, & (k,m) \text{ in the body of water} \\ 0, & (k,m) \text{ outside the body of water} \end{cases} \quad (14)$$

It is now appropriate to discuss the detailed formulations of $I(k,m)$, $T(k,m)$, $D(k,m)$, $F(k,m)$ and $H(k,m)$. These are developed in as simple and practical a manner as possible in the absence of definitive field measurements. Subsequent studies may require alterations of these formulations.

Inertia of forward movement, $I(k,m)$

Assume the fish is at point (i,j) and its preceding location was (i',j') , where

$$i = i' + \delta' \quad (15a)$$

$$j = j' + \epsilon' \quad (15b)$$

and where δ' and ϵ' have the same ranges of values as δ and ϵ [see Eqs. (1a,1b)]. Then $I(k,m)$, where k and m are given by Eqs. (1a,1b), is a conditional probability,

$$I(k,m) = \text{Probability } (\delta, \epsilon \text{ given } \delta', \epsilon'), \quad (16)$$

where this probability is higher the more positive the correlation between (δ, ϵ) and (δ', ϵ') . In the model, a quantity, C , is defined, where,

$$C = |\delta - \delta'| + |\epsilon - \epsilon'|. \quad (17)$$

The bars represent absolute values of the enclosed differences. The quantity C can take on one of five different integer values, for each of which $I(k,m)$ is assigned a different value, e_i , as represented in Eq.(18),

$$I(k,m) = \begin{cases} e_1 & (C = 0) \\ e_2 & (C = 1) \\ e_3 & (C = 2) \\ e_4 & (C = 3) \\ e_5 & (C = 4), \end{cases} \quad (18)$$

where the constants e_i are chosen so that $e_1 > e_2 > e_3 > e_4 > e_5$. The model fish is likely to continue in the same general direction because $I(k,m)$ is greatest when $\delta = \delta'$ and $\epsilon = \epsilon'$.

Temperature term, $T(k,m)$

Assume the fish has a preferred temperature, $TEMP_p$. The temperature at point (i,j) is defined as $TEMP(i,j)$. Define the absolute difference between the temperature at (i,j) and the optimal temperature by $d_T(i,j) = |TEMP(i,j) - TEMP_p|$. Then, if (k,m) is a neighboring point of (i,j) , we define the temperature effect, $T(k,m)$, by

$$T(k,m) = \begin{cases} s_T > 0.0 & d_T(k,m) < d_T(i,j) \\ 0.0 & d_T(k,m) \geq d_T(i,j). \end{cases} \quad (19)$$

The quantitative value of the constant s_T is assigned to reflect the strength of the effect of the temperature gradient on the fish. Estimates of values might be obtained from experiments in which only temperature effects are present.

Dissolved oxygen term, $DO(k,m)$

We have no information on the existence of a "preferred" DO level, but there is evidence on minimum tolerable levels. Define by $DISOX_{\min}$ the minimum tolerable level and by $DISOX(i,j)$ the dissolved oxygen at point (i,j) . Then if the fish is in a spatial region in which the dissolved oxygen is below the minimum tolerable limit, (i.e., $DISOX(i,j) < DISOX_{\min}$), then define the dissolved oxygen effect, $DO(k,m)$, by

$$DO(k,m) = \begin{cases} s_{DO} > 0.0 & DISOX(k,m) > DISOX(i,j) \\ 0.0 & DISOX(k,m) < DISOX(i,j). \end{cases} \quad (20)$$

If the fish is in a region in which the amount of dissolved oxygen in the water is above the minimum tolerable limit, then $DO(k,m) = 0$ for all values of k and m . The constant s_{DO} is a measure of the strength of avoidance by fish of low dissolved oxygen levels.

Food availability terms, $F_q(k,m)$

Assume that there are q regions in the body of water that are attractive to fish because of high food availability. We assume that the closest of these to the current position of the fish will exert some attraction on the fish. Define by $d_{F,q}(i,j)$ the level of food availability at point (i,j) . Then if (k,m) is a point neighboring (i,j) , the force of attraction of the food is

$$F_q(k,m) = \begin{cases} s_{F,q} > 0.0 & d_{F,q}(k,m) < d_{F,q}(i,j) \\ 0.0 & d_{F,q}(k,m) > d_{F,q}(i,j). \end{cases} \quad (21)$$

Habitat preference terms, $H_p(k,m)$

Assume that there are \bar{P} regions in the body of water that are attractive to fish because of their favorability as habitat. We assume that the closest of these to the current position of the fish will exert some attraction on the fish. Define by $d_{H,p}(i,j)$ the level of habitat favorability at point (i,j) . Then if (k,m) is a point neighboring (i,j) , the force of attraction of habitat is

$$H_p(k,m) = \begin{cases} s_{H,p} > 0.0 & d_{H,p}(k,m) < d_{H,p}(i,j) \\ 0.0 & d_{H,p}(k,m) > d_{H,p}(i,j). \end{cases} \quad (22)$$

COMPUTER PROGRAM

The computer program consists of a MAIN PROGRAM and three subroutines, SUBROUTINE RANSET, FUNCTION URAND, SUBROUTINE PLOTT and SUBROUTINE HIST.

The MAIN PROGRAM first reads in the input data, which is described in Part A below, and then prints it out (see Part B, below). There are two ways in which data on temperature and dissolved oxygen can be entered; either by specifying each grid point values, or by using mathematical functions to express their spatial variation. As an example of the latter, temperature might be given by the function

$$\text{TEMP} = 40000. / \{10000. + (i-85.)^2 + 5.0(j-55.)^2\}, \quad (23)$$

which leads to the isotherms shown in Fig. 1. Similar functions are used for dissolved oxygen. Food distribution might be modeled by functions of the form

$$\text{FOOD}_q = F_D / (1.0 + 1.0 \exp\{-\alpha \left(\frac{i-I_q}{F_q}\right)^2 - \beta_{Fq} \left(\frac{j-J_q}{F_q}\right)^2\}), \quad (24)$$

which are plotted in Fig. 4. The peaks and plateaus in this figure represent regions of high food availability. Similar functions are used to describe habitat preferences.

In the input data, the user specifies how many fish are released at random locations in the body of water and how many spatial steps they are allowed to take. The user also chooses whether or not the paths of the fish are to be plotted. If they are not, only the final positions of the fish will be shown by a dot. The user can also have the computer print out the isotherms, if desired.

The program first randomly selects, using a pseudo-random number generator, the position and direction of motion of the fish. Thereafter, the movement of the fish from point to point on the grid is determined by the pseudo-random number generator, in combination with the transition probabilities, $p_{ij,km}$, which are computed at each step.

Information on the paths and final positions of the fish is stored for later printing.

The only purpose of SUBROUTINE RANSET and FUNCTION URAND is to generate pseudo-random numbers on the interval (0,1). These subroutines have been described elsewhere (McGarth and Irving, 1975) and so will not be discussed here. The type of simulation that uses a pseudo-random number generator is commonly referred to as a Monte Carlo simulation. SUBROUTINE PLOTT handles the plotting of the outline of the body of water, while SUBROUTINE HIST plots a histogram of the final temperature distribution of the fish.

The computer program is meant to be very general. If changes in the program are necessary, however, the documentation of the program below should be complete enough to enable the user to make these changes.

The remainder of this section consists of a description of the data input cards (Part A), the printed output of the program (Part B), and a listing of the computer program (Part C). In the next section, the use of the program is demonstrated by means of some trial simulations.

ORNL-DWG 78-2644




Fig. 4. A plot of food availability to fish in a hypothetical reservoir. The peaks and plateaus represent the regions of high food availability.

Part A. Input Cards

Figure 5 is a listing of the input cards relevant to an example given in the next section. These input cards are described below:

Card A

Input parameters: NHOR, NVER, NREG

Format: 4I5

NHOR = number of horizontal grid points

NVER = number of vertical grid points

NREG = number of environmental regions (usually there will be only two;
(1) the body of water, and (2) the surrounding air and land

Card B

Input parameters: NREGP

Format: I5

NREGP = the number of points on the line to be drawn to define the
boundary of the body of water

Card Set C

Input parameters: (ARRAYX(I), I=1,NREGP)

Format: 7E10.0

ARRAYX(I) = the horizontal coordinates of points on the line defining
the boundary of the body of water

Card Set D

Input parameters: (ARRAYY(I), I=1,NREGP)

Format: 7E10.0

ARRAYY(I) = the vertical coordinates of points on the line defining
the boundary of the body of water

Card Set E

Input parameters: NVER cards containing the information IREG,
(IBEG(I), IEND(I), TYPE(I), I=1,IREG)

Format: I2, 8X, 6(2I2,F5.1,1X)

90	60	2								A
17										B
2.0		5.0	10.0	15.0	20.0	25.0	30.0			C
35.0		43.0	55.0	70.0	77.0	82.0	88.0			C
88.0		2.0	2.0							C
3.0		3.0	4.0	5.0	6.0	7.0	9.0			D
10.0		16.0	22.0	30.0	36.0	38.0	39.0			D
55.0		55.0	3.0							D
1		0190	1.0							E
1		0190	1.0							E
3		0102	1.0	0305	3.0	0690	1.0			E
3		0102	1.0	0307	3.0	0890	1.0			E
3		0102	1.0	0312	3.0	1390	1.0			E
3		0102	1.0	0314	3.0	1590	1.0			E
3		0102	1.0	0322	3.0	2390	1.0			E
3		0102	1.0	0326	3.0	2790	1.0			E
3		0102	1.0	0329	3.0	3090	1.0			E
3		0102	1.0	0331	3.0	3290	1.0			E
3		0102	1.0	0334	3.0	3590	1.0			E
3		0102	1.0	0336	3.0	3790	1.0			E
3		0102	1.0	0338	3.0	3990	1.0			E
3		0102	1.0	0339	3.0	4090	1.0			E
3		0102	1.0	0340	3.0	4190	1.0			E
3		0102	1.0	0342	3.0	4390	1.0			E
3		0102	1.0	0344	3.0	4590	1.0			E
3		0102	1.0	0345	3.0	4690	1.0			E
3		0102	1.0	0347	3.0	4890	1.0			E
3		0102	1.0	0349	3.0	5090	1.0			E
3		0102	1.0	0354	3.0	5290	1.0			E
3		0102	1.0	0353	3.0	5490	1.0			E
3		0102	1.0	0355	3.0	5690	1.0			E
3		0102	1.0	0357	3.0	5890	1.0			E
3		0102	1.0	0359	3.0	6090	1.0			E
3		0102	1.0	0361	3.0	6290	1.0			E
3		0102	1.0	0363	3.0	6490	1.0			E
3		0102	1.0	0365	3.0	6690	1.0			E
3		0102	1.0	0366	3.0	6790	1.0			E
3		0102	1.0	0368	3.0	6990	1.0			E
3		0102	1.0	0369	3.0	7090	1.0			E
3		0102	1.0	0371	3.0	7290	1.0			E
3		0102	1.0	0372	3.0	7390	1.0			E
3		0102	1.0	0373	3.0	7490	1.0			E
3		0102	1.0	0374	3.0	7590	1.0			E
3		0102	1.0	0375	3.0	7690	1.0			E
3		0102	1.0	0376	3.0	7790	1.0			E
3		0102	1.0	0380	3.0	8190	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
3		0102	1.0	0387	3.0	8890	1.0			E
1		0190	1.0							F
1		0190	1.0							F
1		0190	1.0							F
1		0190	1.0							F
1		0190	1.0							F
		0	0							F
		29.0	1.0							P
		2.0	0.0							I
		0								J
		0.0								K
		0								L
		0.0								N
		0.001	0.001	10.						O
		0.1	0.1	0.1						O
		98765								R
		500	200							S
		0	1							T
		16.	44.	4.0						U
		2	200							V
		1	1							T

Fig. 5. Input data for a sample trial simulation as it appears on the data cards.

IREG = number of different environmental types along a given line of grid points

IBEG(I) = the horizontal coordinate of the first grid point of a particular environmental type along a given horizontal line

IEND(I) = The horizontal coordinate of the last grid point of a particular environmental type along a given horizontal line

TYPE(I) = a numerical label attached to each environmental type to distinguish it from others

Card F

Input parameters: ITEM, IDISOX

Format: 2I5

ITEM = 0 if spatial temperature data is given by an equation in the program
 = 1 if spatial temperature data is read in point by point

IDISOX = 0 if spatial dissolved oxygen is given by an equation in the program
 = 1 if spatial dissolved oxygen data is read in point by point

Card Set G (included only if ITEM = 1)

Input parameters: (TEMPA(I,J), I=1,NHOR), J=1,NVER

Format: 7E10.0

TEMPA(I,J) = temperature at grid point (I,J)

Card Set H (included only if IDISOX = 1)

Input parameters: (DISOX(I,J), I=1,NHOR), J=1,NVER

Format: 7E10.0

DISOX(I,J) = dissolved oxygen level at grid point (I,J)

Card I

Input parameters: TEMPRF, TEMFOR

Format: 2E10.0

TEMPFR = preferred temperature of fish

TEMFOR = force of attraction of preferred temperature of fish

Card J

Input parameters: DOXMIN, DOXFOR

Format: 2E10.0

DOXMIN = minimum tolerable dissolved oxygen level for fish

DOXFOR = attractive force of higher dissolved oxygen levels on fish

Card K

Input parameter: NFOOD

Format: I5

NFOOD = number of centers of high food availability

Card L

Input parameter: FDATCT

Format: E10.0

FDATCT = force of attraction of food availability on fish movements

Card Set M

Input parameters: FDNUM(I), FDALP(I), FDBET(I), FDIQ(I), FDJQ(I)

Format: 5E10.0

FDNUM(I)

FDALP(I) parameters describing spatial distributions of

FDBET(I) = food about each of the centers of food

FDIQ(I) availability (see Eq. 24 and Table 1)

FDJQ(I)

Card N

Input parameter: NHAB

Format: I5

NHAB = number of centers of high habitat favorability

Card O

Input parameter: HBATCT

Format: E10.0

HBATCT = force of attraction of habitat favorability on fish movements

Card Set P

Input parameters: HBNUM(I), HBALP(I), HBBET(I), HBIQ(I), HBJQ(I)

Format: 5E10.0

HBNUM(I)

HBALP(I) parameters describing the spatial distribution

HBBET(I) = of habitat favorability about the high habitat

HBIQ(I) favorability centers (analogous to Eq. (24); also

HBJQ(I) see Table 1 for definitions)

Card Set Q

Input parameters: RES(I), I=1,NREG

Format: 7E10.0

RES(I) = boundary crossing factors (causing fish to remain in the body of water)

Card R

Input parameters: ERTIA(I), I=1,5

Format: 5E10.0

ERTIA(I) = Inertia of forward motion, e_j (see Eq. 18)

Card S

Input parameter: IX

Format: I5

IX = pseudo-random number generator initialization or "seed". It must be an odd integer. A different value of IX should be used each time the program is run

Card T

Input parameters: NFISH, NSTEP

Format: 2I5

NFISH = number of fish considered in the body of water

NSTEP = number of steps in space each fish is allowed to take

Card U

Input parameters: I PLOT, ISOTH

Format: 2I5

I PLOT = 1 if the fish paths are to be plotted, 0 otherwise

ISOTH = 1 if the isotherms are to be plotted, 0 otherwise

Card V

Input parameters: T EML, T EMH, T EMINT

Format: 3E10.0

T EML = minimum isotherm to be plotted

T EMH = maximum isotherm to be plotted

T EMINT = width of intervals between isotherms

Part B. Output

The printed output consists of two parts. First, the input data is printed out (Fig. 6). Second, a schemata of the body of water is plotted, into which fish paths or spatial population distribution are plotted (Figs. 7 and 8). The plotting is done using the DISSPLA graphics package (Integrated Software Systems Corporation 1970) which is available at many computer installations. Programming changes would be necessary to adapt the program to other graphics packages.

Part C. Computer program details

The complete computer program listing is printed in the Appendix. The comment cards interspersed through the program should enable the user to understand its general design. However, some additional comments may be useful.

1. The arrays are dimensioned to permit a maximum of 90x60 grid points at present. This can be changed if desired.
2. A typical run dispersing 500 fish takes about 3 minutes of CPU time in the IBM 360/91 computer, although this changes to some extent as some of the model parameters are varied. The GO step uses less than 230K of computer core.

FISH MOVEMENT IN A BODY OF WATER

NUMBER OF HORIZONTAL GRID POINTS, NHOR = 90
 NUMBER OF VERTICAL GRID POINTS, NVER = 60
 NUMBER OF ENVIRONMENTAL REGIONS, NREG = 2

 TEMPERATURE IS DESCRIBED BY A MATHEMATICAL FUNCTION

 DISSOLVED OXYGEN AMOUNTS DESCRIBED BY A MATHEMATICAL FUNCTION

 PREFERRED TEMPERATURE, TEMPRF = 29.0000
 FORCE OF ATTRACTION OF PREFERRED TEMPERATURE, TEMPOR = 1.0000

 MINIMUM TOLERABLE DISSOLVED OXYGEN LEVEL, DOXMIN = 2.0000
 FORCE OF ATTRACTION OF HIGHER DISSOLVED OXYGEN LEVELS = 0.0

 FORCE OF ATTRACTION OF GREATER FOOD AVAILABILITY, FDATCT = 0.0
 NFOOD = 0

 FORCE OF ATTRACTION OF HABITAT PREFERENCES, HBATCT = 0.0
 NHAB = 0

 BOUNDARY CROSSING FACTORS, RES(I) = 0.001 0.001

 VALUES OF FORWARD INERTIA, ERTIA = 0.10 0.10 0.10 0.0 0.0

 RANDOM NUMBER INITIATOR, IX = 98765

 NUMBER OF FISH IN BODY OF WATER, NFISH = 500
 NUMBER OF STEPS EACH FISH IS ALLOWED TO TAKE, NSTEPS = 200

 ISOTHERMS ARE PLOTTED

 MINIMUM ISOTHERM PLOTTED, TEML = 16.0000
 MAXIMUM ISOTHERM PLOTTED, TEMH = 44.0000
 DISTANCE BETWEEN ISOTHERMS, TEMINT = 4.0000

Fig. 6. Input data for a sample trial simulation as it is printed out by the computer program.

FISH DISTRIBUTION




Fig. 7. Plot of simulated motions of two fish initially placed at points A and B. The assumed preferred temperature is $TEMP_p = 29.0^{\circ}C$ and the force of temperature attraction, p_T , is 1.0 for case A and 50.0 for case B.




Fig. 8. The distribution of 500 fish influenced only by temperature in the reservoir after 200 steps. The assumed preferred temperature is 29.0°C. Other parameters of the model are given in Table 2.

Table 1 is a compilation of the principal FORTRAN variables in the computer program. The equivalent mathematical symbols, of any, and definitions are given as well.

TRIAL SIMULATIONS

The fundamental question that must be asked of this model is how accurately it can simulate the movements of individual fish and the spatial distribution patterns of populations of fish. There is not enough data on either of these phenomena in natural environments to allow parameters for a model to be thoroughly tested. However, laboratory experiments provide some data on fish distributions in environments in which only thermal effects are important. We shall focus on the thermal influences on the fish in our model and only briefly note how the other factors influence fish distributions in space.

Fish Movements

Consider the reservoir pictured in Fig. 7, with only the temperature gradient assumed to have an effect on the fish. The temperature isoclines are given by Eq. (23) and the remaining parameters of the model are given in Table 2. A simulated fish is placed in the reservoir at the position A; it moves, with a fair amount of meandering, toward the preferred temperature, $TEMP_p = 29.0^\circ C$. The amount of meandering can be decreased by increasing the force of the temperature gradient on the fish movement; that is, by increasing p_T . When p_T is increased from $p_T = 1.0$ to $p_T = 50.$, and a fish is released at point B, it moves more directly toward the preferred temperature.

Fish Distribution Patterns

Allow 500 fish to be released at randomly selected initial positions in the body of water, and to move in response to temperature gradients only. After 200 steps, they have all had a chance to respond

Table 1. Principal program variables

Fortran variable name	Dimension (if array)	Mathematical symbol	Definition
ARAX	(50)		Storage array for horizontal coordinates of isotherm curves for later plotting
ARAY	(50)		Storage array for vertical coordinates of isotherm curves for later plotting
ARRAYX	(50)		Storage array for horizontal coordinates of outline of body of water
ARRAYY	(50)		Storage array for vertical coordinates of outline of body of water
D			Random number chosen from uniform distribution on the interval (0,1)
DDIFF			Difference between the dissolved oxygen level at the current position of the fish and its minimum tolerable dissolved oxygen level
DDIFFA			Difference between dissolved oxygen level of any of the next eight possible positions of the fish and its minimum tolerable dissolved oxygen level
DDR	(3,3)	$I(k,m)$	Measure of the strength of the inertia of forward movement of the fish
DIR	(3,3)	$DO(k,m)$	Attraction of point (k,m) on fish because of the difference in the dissolved oxygen level from that of the current location of the fish
DISOX	(100)	$DISOX(k,m)$	Storage array for dissolved oxygen levels along some given horizontal line, k
DOX		$DISOX(i,j)$	Level of dissolved oxygen at the current position of the fish

Table 1. (continued)

Fortran variable name	Dimension (if array)	Mathematical symbol	Definition
DOXFOR		S_{DO}	Attractive force of higher dissolved oxygen level on fish movements
DOXMIN		$DISOX_{min}$	Minimum tolerable dissolved oxygen level for fish
ERTIA	(5)	e_i	Strength of forward inertia of fish
FDALP	(20)	α_{Fq}	Parameter describing the spatial distribution of food about each of the centers of food availability (see Eq. 24)
FCATCT		$\beta_{F,q}$	Force of attraction of food availability on fish movements
FDBET	(20)	β_{Fq}	Parameter describing the spatial distribution of food about each of the centers of food availability (see Eq. 24)
FDIQ	(20)	I_{Fq}	Parameter (horizontal coordinate) describing the spatial distribution of food about each of the centers of food availability (see Eq. 24)
FDJQ	(20)	J_{Fq}	Same as above definition (vertical coordinate)
FDNUM	(20)	F_D	Parameter describing the spatial distribution of favorable habitat about each of the centers of food availability (see Eq. 24)
FDR	(3,3)	$F_Q(k,m)$	Attraction of point (k,m) or fish because of the difference in food availability from the current location (i,j)
FOOD			Measure of the amount of food available to the fish at its current location
FOODA			Measure of the amount of food available to fish in its possible next location

Table 1. (continued)

Fortran variable name	Dimension (if array)	Mathematical symbol	Definition
GRID	(90,60)		Array that stores information on the type of region each grid point is in, as well as its temperature and dissolved oxygen level
HAB			Measure of the favorability of habitat at the current location of the fish
HABA			Measure of the favorability of habitat at the possible next location of the fish
HABALP	(20)	$\alpha_{H,q}$	Parameter describing the spatial distribution of favorable habitat about each of the centers of favorable habitat (in equation analogous to Eq. 24)
HBATCT		$s_{H,q}$	Force of attraction of habit favorability on fish movements
HBBET	(20)	$\beta_{H,q}$	Parameters describing the spatial distribution of favorable habitat about each of the centers of favorable habitat (in equation analogous to Eq. 24)
HBIQ	(20)	$I_{H,q}$	Center of a region of favorable habitat (horizontal coordinate)
HBJQ	(20)	$J_{H,q}$	Center of a region of favorable habitat (vertical coordinate)
HBNUM	(20)	H_D	Parameters describing the spatial distribution of favorable habitat about each of the centers of favorable habitat (in equation analogous to Eq. 24)
IDISOX			Logical variable specifying whether dissolved oxygen levels are described by a mathematical function (IDISOX=0) or point by point (IDISOX=1)

Table 1. (continued)

Fortran variable name	Dimension (if array)	Mathematical symbol	Definition
IPLOT			Logical variable specifying whether or not fish paths are to be plotted
IPRES			Current position of the fish (horizontal coordinate)
ISOTH			Logical variable specifying whether or not the isotherms are to be plotted
ISTR			Horizontal coordinate of the starting position of a given fish
ITEM			Logical variable specifying whether temperature is described by a mathematical function (ITEM=0) or by point-by-point data (ITEM=1)
IX			Pseudo-random number generator initiator
JPRES			Current position of the fish (vertical coordinate)
JSTR			Starting position of the fish (vertical coordinate)
NDIST			Integer variable that increases by 1 for each step a particular fish takes. When NDIST=NSTEPS, no further steps are taken
NFISH			Number of fish simulated in the body of water
NFOOD			Number of centers of food availability
NHAB			Number of centers of high habitat favorability
NHOR			Number of horizontal grid lines
NREG			Number of environmental regions [usually there will be only two; (1) the body of water, and (2) the surrounding air and land]
NSV			Integer variable that increases by 1 for each fish that is "inserted" into the body of water. When NSV = NFISH, no further fish are inserted.

Table 1. (continued)

Fortran variable name	Dimension (if array)	Mathematical symbol	Definition
NSTEPS			Number of steps in space that each fish is allowed to take
NVER			Number of vertical grid lines
RES	(50)	$B(k,m)$	Boundary crossing factors (causing fish to remain in the body of water)
SAVI	(500)		Array that stores horizontal coordinates of fish movement for later plotting
SAVJ	(500)		Array that stores vertical coordinates of fish movement for later plotting
TDIFF		$d_T(i,j)$	Difference between temperature of current position of fish and its preferred temperature
TDIFFA		$d_T(k,m)$	Difference between temperature of possible next position of the fish and its preferred temperature
TDR	(3,3)	$T_q(k,m)$	Attraction of point (k,m) on the fish because of the difference on temperature from its current position
TEMH			Temperature of maximum isotherm to be plotted
TEMINT			Width of intervals between isotherms
TEML			Temperature of minimum isotherm to be plotted
TEMPOR		s_T	Force of attraction of preferred temperature of fish
TEMP			Temperature at current location of fish
TEMPA	(100)	$TEMP(i,j)$	Storage array for temperature data along a given horizontal line, i
TEMPRF		$TEMP_p$	Preferred temperature of the fish
V	(3,3)	$P_{ij,km}$	Transition probability from grid point (i,j) to grid point (k,m)

Table 2. Parameter values for the example in Fig. 5

NHOR = 90 NVER = 60 NREG = 2
NREGP = 17
ARRAYX(I) (I=1,17) = 2.0, 5.0, 10.0, 15.0, 20.0, 25.0, 30.0, 35.0, 43.0, 55.0,
70.0, 77.0, 82.0, 88.0, 2.0, 2.0
ARRAYY(I) (I=1,17) = 3.0, 3.0, 4.0, 5.0, 6.0, 7.0, 9.0, 10.0, 16.0, 22.0, 30.0,
36.0, 38.0, 39.0, 55.0, 55.0, 3.0

IREG	IBEG(1)	IEND(1)	TYPE(1)	IBEG(2)	IEND(2)	TYPE(2)	IBEG(3)	IEND(3)	TYPE(3)
1	01	90	1.0						
1	01	90	1.0						
3	01	02	1.0	03	05	3.0	06	90	1.0
3	01	02	1.0	03	07	3.0	08	90	1.0
3	01	02	1.0	03	12	3.0	13	90	1.0
3	01	02	1.0	03	14	3.0	15	90	1.0
3	01	02	1.0	03	22	3.0	23	90	1.0
3	01	02	1.0	03	26	3.0	27	90	1.0
3	01	02	1.0	03	29	3.0	30	90	1.0
3	01	02	1.0	03	31	3.0	32	90	1.0
3	01	02	1.0	03	34	3.0	35	90	1.0
3	01	02	1.0	03	36	3.0	37	90	1.0
3	01	02	1.0	03	38	3.0	39	90	1.0
3	01	02	1.0	03	39	3.0	40	90	1.0
3	01	02	1.0	03	40	3.0	41	90	1.0
3	01	02	1.0	03	42	3.0	43	90	1.0
3	01	02	1.0	03	44	3.0	45	90	1.0
3	01	02	1.0	03	45	3.0	46	90	1.0
3	01	02	1.0	03	47	3.0	48	90	1.0
3	01	02	1.0	03	49	3.0	50	90	1.0
3	01	02	1.0	03	51	3.0	52	90	1.0
3	01	02	1.0	03	53	3.0	54	90	1.0
3	01	02	1.0	03	55	3.0	56	90	1.0
3	01	02	1.0	03	57	3.0	58	90	1.0

Table 2. (continued)

IREG	IBEG(1)	IEND(1)	TYPE(1)	IBEG(2)	IEND(2)	TYPE(2)	IBEG(3)	IEND(3)	TYPE(3)
3	01	02	1.0	03	59	3.0	60	90	1.0
3	01	02	1.0	03	61	3.0	62	90	1.0
3	01	02	1.0	03	63	3.0	64	90	1.0
3	01	02	1.0	03	65	3.0	66	90	1.0
3	01	02	1.0	03	66	3.0	67	90	1.0
3	01	02	1.0	03	68	3.0	69	90	1.0
3	01	02	1.0	03	69	3.0	70	90	1.0
3	01	02	1.0	03	71	3.0	72	90	1.0
3	01	02	1.0	03	72	3.0	73	90	1.0
3	01	02	1.0	03	73	3.0	74	90	1.0
3	01	02	1.0	03	74	3.0	75	90	1.0
3	01	02	1.0	03	75	3.0	76	90	1.0
3	01	02	1.0	03	76	3.0	77	90	1.0
3	01	02	1.0	03	80	3.0	81	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	02	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
3	01	02	1.0	03	87	3.0	88	90	1.0
1	01	90	1.0						
1	01	90	1.0						
1	01	90	1.0						
1	01	90	1.0						
1	01	90	1.0						

Table 2. (continued)

ITEM = 0 IDISOX = 0
TEMPA(I,J) not entered
DISOX(I,J) not entered
TEMPRF = 29.0 TEMFOR = 1.0
DOXMIN = 2.0 DOXFOR = 0.0
NFOOD = 0 FDATCT = 0.0
FDNUM(I), FDALP(I), FDBET(I), FDIQ(I), FDJQ(I) not entered
NHAB = 0 HBATCT = 0.0
HBNUM(I), HBALP(I), HBBET(I), HBIQ(I), HBJQ(I) not entered
RES(I) (I=1,3) = 0.001, 0.001, 10.0
ERTIA(I) (I=1,5) = 0.1, 0.1, 0.1, 0.0, 0.0
IX = 98765
NFISH = 2 NSTEP = 200
IPLOT = 1 ISOTH = 1
TEML = 16.0 TEMH = 44.0 TEMINT = 4.0

to the preferred temperature. The distribution of fish after 200 steps is shown in Fig. 8, for parameter values given in Table 2, except that now NFISH = 500 and IPLIT = 0. It is interesting to look at the histogram describing the percent distribution of fish about the preferred temperature of 29.0°C (Fig. 9), since this can be compared with laboratory data, such as that shown in Fig. 10 for largemouth bass (Reynolds and Casterlin 1975). The agreement is not bad (although the model results are more peaked and lack the skewing seen in the experiment), which is some indication that we have chosen a reasonable set of parameters for our model; however, other choices of parameter values may give better results.

Next we add in the effects of dissolved oxygen (Fig. 1), food availability (Fig. 4), and habitat preferenda, with the appropriate changes in parameter values from Table 2 shown in Table 3. The ultimate average distribution of fish is now greatly altered (Fig. 11).

DISCUSSION AND SUMMARY

The model described in this report is designed to simulate the movements of individual fish in a body of water and to predict the spatial patterns of a population of fish under the influence of temperature, dissolved oxygen levels, food availability and habitat preferences. The body of water is represented by a two-dimensional grid of points, with water depth and longitudinal axis being the coordinates. The simulated fish takes one spatial step at a time, the direction of travel being chosen by a pseudo-random number generator, but biased by the initial direction of motion of the fish, as well as its response to temperature gradients and the other factors mentioned above. Model output is plotted in graphs.

This model is designed for use in planning and evaluating the results of experimental laboratory and field studies of fish movement and spatial distribution. The application of the model to experimental data is still in a preliminary stage, and the development of the model into an effective predictive tool will take continued work. The model




Fig. 9. Histogram of percent distribution of fish in Fig. 8 about the preferred temperature of 29.0°C.




Fig. 10. Histogram of relative frequency of largemouth bass in ambient water temperatures during daytime (from Reynolds and Casterlin 1977).

FISH DISTRIBUTION




Fig. 11. The distribution of 500 fish influenced by temperature, dissolved oxygen, food availability and habitat favorability in the reservoir after 200 steps. Parameter values are given in Table 3.

Table 3. Changes in parameter values from Table 2. relevant to the case shown in Fig. 11

NFOOD = 1

FDNUM(1) = 10.0

FDALP(1) = 0.02

FDBET(1) = 0.2

FDIQ(1) = 60.0

FDJQ(1) = 45.0

FDNUM(2) = 10.0

FDALP(2) = 0.10

FDBET(2) = 0.20

FDIQ(2) = 70.0

FDJQ(2) = 50.0

NHAB = 1

HBNUM(1) = 10.0

HBALP(1) = 0.05

HBET(1) = 0.20

HBIQ(1) = 45.0

HBJQ(1) = 53.0

HBNUM(2) = 10.0

HBALP(2) = 0.05

HBBET(2) = 0.2

HBIQ(2) = 45.0

HBJQ(2) = 53.0

NFISH = 500

IPLLOT = 0

is flexible enough to take into account most of the important factors influencing fish movement, but considerable effort needs to be expended in quantifying these factors.

REFERENCES

- Coutant, C. C. 1975. Responses of bass to natural and artificial temperature regimes. pp. 272-285. IN: Stroud, R. H., and H. Clepper (eds.), Black Bass Biology and Management. Sports Fishing Institute, Washington, DC. 534 pp.
- Coutant, C. C., and D. K. Cox. 1976. Growth rates of subadult largemouth bass at 24 to 35.5 C. pp. 188-120. IN: Esch, G. W., and R. W. McFarlane (eds.), Thermal Ecology II. ERDA Symposium Series CONF-750425. National Technical Information System, Springfield, VA.
- Fraenkel, G. S., and D. L. Gunn. 1961. The Orientation of Animals (revised edition). Dover Pub., Inc., New York. 376 pp.
- Gibbons, J. W., J. T. Hook, and D. L. Forney. 1972. Winter responses of largemouth bass to heated effluent from a nuclear reactor. Prog. Fish Cult. 34(2):88-90.
- Kitchell, J. F., J. F. Koonce, R. V. O'Neill, H. H. Shugart, Jr., J. J. Magnuson, and R. S. Booth. 1974. Model of fish biomass energetics. Trans. Am. Fish. Soc. 103(4):786-798.
- McCauley, R., and N. Huggins. 1976. Behavioral thermoregulation by rainbow trout in a temperature gradient. pp. 171-175. IN: Esch, G. W., and R. W. McFarlane (eds.), Thermal Ecology II. ERDA Symposium Series CONF-750425. National Technical Information System, Springfield, VA.
- McGarth, E. J., and D. C. Irving. 1975. Techniques for efficient Monte Carlo simulation. Vol. II. Random number generation for selected probability distributions. ORNL/RSIC-38 (Vol. 2). Oak Ridge National Laboratory, Oak Ridge, TN.
- Moss, D. D., and D. C. Scott. 1961. Dissolved oxygen requirements of three species of fish. Trans. Am. Fish. Soc. 90(4):377-393.
- Neill, W. H. 1976. Mechanisms of behavioral thermoregulation in fishes. pp. 156-169. IN: Sigma Research Inc. (ed), Report of a Workshop on the Impact of Thermal Power Plant Cooling Systems on Aquatic Environments. Electric Power Research Institute, Palo Alto, CA.

- Peterson, S. E., and R. M. Schutsky. 1976. Some relationships of upper thermal tolerances to preference and avoidance responses of the bluegill. pp. 148-153. IN: Esch, G. W., and R. W. McFarlane (eds.), Thermal Ecology II. ERDA Symposium Series CONF-750425. National Technical Information System, Springfield, VA.
- Reynolds, W. W., and M. E. Casterlin. 1976. Thermal preferenda and behavioral thermoregulation in three centrarchid fishes. IN: Esch, G. W., and R. W. McFarlane (eds.), Thermal Ecology II. ERDA Symposium Series CONF-750425. National Technical Information System, Springfield, VA.
- Smith, S. F. 1972. Effects of a thermal effluent on aquatic life in an East Tennessee reservoir. Proc. 25th Annu. Conf. S. E. Game and Fish Comm., 374-384.
- Stuntz, W. E., and J. J. Magnuson. 1976. Daily ration, temperature selection, and activity of bluegill. pp. 180-184. IN: Esch, G. W., and R. W. McFarlane (eds.), Thermal Ecology II. ERDA Symposium Series CONF-750425. National Technical Information System, Springfield, VA.
- Warden, R. L., Jr., and W. J. Lorio. 1975. Movements of largemouth bass (Micropterus salmoides) in impounded waters as determined by underwater telemetry. Trans. Am. Fish. Soc. 104(4):696-702.

APPENDIX: THE COMPUTER PROGRAM

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

```

0001      IMPLICIT REAL*4(A-H,O-Z)
          C
          C *****
          C
          C THIS PROGRAM COMPUTES FISH DISTRIBUTIONS IN A BODY OF WATER.
          C
          C WRITTEN BY D. L. DEANGELIS, 1977
          C
          C *****
0002      DIMENSION GRID(90,60),IREG(40),IEND(40),TYPE(40)
0003      DIMENSION ARAX(50), ARAY(50)
0004      DIMENSION DHOR(100),DVER(100),RES(40)
0005      DIMENSION IA(40),JA(40),ATINDX(40)
0006      DIMENSION ADIR(40),DIR(3,3),RDIR(3,3),BDY(3,3),ATDIR(3,3),V(3,3),
          C 1SAVI(500),SAVJ(500),SPRESI(1),SPRESJ(1)
0007      DIMENSION IGR(90),SDF(40),AGR(100),AG(40),ERTIA(20)
0008      DIMENSION ITBEG(50),ITEND(50),TDR(3,3)
0009      DIMENSION TEMPA(100),DISOX(100),FDR(3,3),HDR(3,3),DDR(3,3)
0010      DIMENSION FDALP(20),FDBET(20),FDNUM(20),FDIQ(20),FDJQ(20),
          C 1HBALP(20),HBBET(20),HBNUM(20),HBIQ(20),HBJQ(20)
0011      DIMENSION TEMSAV(100)
0012      COMMON/FINBLK/NFIN,NREGP
0013      COMMON/NFBLOK/NFISH
0014      COMMON/DRAW/ARRAYX(50,50),ARRAYY(50,50)
0015      COMMON/THERM/KA,KB,NPTYP
          C
0016      NSV = 0
0017      DO 3 I=1,100
0018      TEMSAV(I) = 0.0
0019      3 CONTINUE
          C
          C
          C
          C.....READ IN THE NUMBER OF HORIZONTAL AND VERTICAL GRID POINTS AND THE
          C NUMBER OF DISTINCT AREA TYPES
          C
0020      READ(5,1000) NHOR,NVER,NREG
0021      1000 FORMAT(14I5)
0022      WRITE(6,2000)
0023      2000 FORMAT(1H1,10X,'FISH MOVEMENT IN A BODY OF WATER',///)
0024      WRITE(6,2001) NHOR,NVER,NREG
0025      2001 FORMAT(1H ,5X,'NUMBER OF HORIZONTAL GRID POINTS, NHOR = ',I5,/,
          C 1 6X,'NUMBER OF VERTICAL GRID POINTS, NVER = ',I5,/,
          C 2 6X,'NUMBER OF ENVIRONMENTAL REGIONS, NREG = ',I5,/)
          C
          C.....READ IN CARDS CONTAINING THE OUTLINE OF THE BODY OF WATER (LONGITUDE
          C VERSUS DEPTH)
          C
0026      READ(5,1000) NREGP
0027      I = 1
0028      READ(5,1004) (ARRAYX(I,J),J=1,NREGP)
0029      READ(5,1004) (ARRAYY(I,J),J=1,NREGP)

```

```

0030      1004 FORMAT (7E10.0)
C
C
C.....READ IN CARDS SPECIFYING WHICH HABITAT EACH GRID POINT BELONGS TO
C
0031      DO 10 J=1,NVER
0032      READ (5,1001) IREG, (IBEG(I), IEND(I), TYPE(I), I=1, IREG)
0033      1001 FORMAT (I2, 8X, 6(2I2, F5.1, 1X))
0034      DO 5 I=1, IREG
0035      IE = IREG(I)
0036      IJ = IEND(I)
0037      DO 5 K=IB, IE
0038      GRID(K, J) = TYPE(I)
0039      5 CONTINUE.
0040      10 CONTINUE

C
C
C.....READ IN ITEM=0 AND IDISOX=0 IF TEMPERATURE AND DISSOLVED OXYGEN ARE
C DESCRIBED BY MATHEMATICAL FUNCTIONS AND ITEM=1 AND IDISOX=1 IF THEY
C ARE SPECIFIED POINT-BY-POINT IN SPACE
C
0041      READ (5,1000) ITEM, IDISOX
0042      IF (ITEM .EQ. 1) GO TO 12
0043      WRITE (6, 2002)
0044      2002 FORMAT (1H ,5X, 'TEMPERATURE IS DESCRIBED BY A MATHEMATICAL FUNCTION
0045      1', //)
0046      GO TO 14
0047      12 CONTINUE
0048      WRITE (6, 2003)
0049      2003 FORMAT (1H ,5X, 'TEMPERATURE IS READ IN GRID POINT-BY-GRID POINT', //
0050      1)
0051      14 CONTINUE
0052      IF (IDISOX .EQ. 1) GO TO 16
0053      WRITE (6, 2004)
0054      2004 FORMAT (1H ,5X, 'DISSOLVED OXYGEN AMOUNTS DESCRIBED BY A MATHEMATICA
0055      1L FUNCTION', //)
0056      GO TO 18
0057      16 CONTINUE
0058      WRITE (6, 2005)
0059      2005 FORMAT (1H ,5X, 'DISSOLVED OXYGEN AMOUNTS READ IN GRID POINT-BY-GRID
0060      1 POINT', //)
0061      18 CONTINUE

C
C
C.....READ IN THE TEMPERATURE VALUES AT EACH GRID POINT
C EITHER FROM INPUT DATA OR AN EQUATION
C
0062      DO 25 J=1, NVER
0063      SJ = J
0064      IF (ITEM .EQ. 0) GO TO 22
0065      READ (5, 1004) (TEMPA(I), I=1, NHOR)
0066      22 CONTINUE
0067      DO 25 I=1, NHOR
0068      SI = I
0069      IF (ITEM .EQ. 0) GO TO 23
0070      TEMP = TEMPA(I)
0071      GO TO 24
0072      23 CONTINUE

```



```

0069      TEMP = 400000. / (10000. + 0.8*(SI-85.)**2 + 6.*(SJ-55.)**2)
0070      24 CONTINUE
0071      LTEM = TEMP*10.
0072      TEMP = LTEM
0073      TEMP = 0.1*TEMP
0074      GRID(I,J) = GRID(I,J) + 0.01*TEMP
0075      25 CONTINUE
C
C.....READ IN DISSOLVED OXYGEN VALUES AT EACH POINT
C      EITHER FROM INPUT DATA OR AN EQUATION
C
0076      DO 30 J=1,NVER
0077      IF (IDISOX .EQ. 0) GO TO 27
0078      READ(5,1004) (DISOX(I),I=1,NHOR)
0079      27 CONTINUE
0080      DO 30 I=1,NHOR
0081      IF (ITEM .EQ. 0) GO TO 28
0082      DOX = DISOX(I)
0083      GO TO 29
0084      28 CONTINUE
0085      SJ = J
0086      DOX = 1.0 + 0.1*SJ
0087      29 CONTINUE
0088      LDOX = DOX*10.
0089      DOX = LDOX
0090      DOX = 0.1*DOX
0091      GRID(I,J) = GRID(I,J) + 0.000001*DOX
0092      30 CONTINUE
C
C.....READ IN OPTIMAL TEMPERATURE AND ITS ATTRACTIVE EFFECT ON FISH
C
0093      READ(5,1002) TEMPRF, TEMPOR
0094      WRITE(6,2006) TEMPRF, TEMPOR
0095      2006 FORMAT(1H ,5X,'PREFERRED TEMPERATURE, TEMPRF = ',F10.4,/,/,6X,
1'FORCE OF ATTRACTION OF PREFERRED TEMPERATURE, TEMPOR = ',F10.4,/,/
2)
C
C.....READ IN THE VALUE OF THE MINIMUM TOLERABLE DISSOLVED OXYGEN LEVEL
C      FOR THE FISH IN QUESTION AND THE FORCE OF ATTRACTION OF HIGHER
C      DISSOLVED OXYGEN LEVELS
C
0096      READ(5,1004) DOXMIN,DOXPOR
0097      WRITE(6,2007) DOXMIN,DOXPOR
0098      2007 FORMAT(1H ,5X,'MINIMUM TOLERABLE DISSOLVED OXYGEN LEVEL, DOXMIN =
1',F10.4,/,/,6X,'FORCE OF ATTRACTION OF HIGHER DISSOLVED OXYGEN LEVE
2LS = ',F10.4,/)
C
C.....READ IN CARDS SPECIFYING THE EFFECTS OF THE ATTRACTION OF FOOD
C      DISTRIBUTED THROUGH THE BODY OF WATER ON THE MOVEMENTS OF FISH
C
0099      READ(5,1000) NFOOD
0100      READ(5,1002) FDATCT
0101      WRITE(6,2008) FDATCT,NFOOD
0102      2008 FORMAT(1H ,5X,'FORCE OF ATTRACTION OF GREATER FOOD AVAILABILITY, F
1DATCT = ',F10.4,/,/,6X,'NFOOD = ',I5,/)
0103      IF (NFOOD .EQ. 0) GO TO 36
0104      WRITE(6,2009)

```

```

0105      2009 FORMAT (1H ,12X,'FOOD DISTRIBUTION COEFFICIENTS',/,6X,'PNUM',12X,
          1'FDALP',12X,'FDBET',12X,'FDIQ',13X,'FDJQ',/)
0106      DO 35 I=1,NFOOD
0107      READ (5,1002) PNUM(I),FDALP(I),FDBET(I),FDIQ(I),FDJQ(I)
0108      WRITE (6,2010) PNUM(I),FDALP(I),FDBET(I),FDIQ(I),FDJQ(I)
0109      2010 FORMAT (1H ,5X,6(E15.8,2X))
0110      35 CONTINUE
0111      36 CONTINUE

C
C.....READ IN CARDS SPECIFYING THE EFFECTS OF THE ATTRACTION OF HABITATS
C      IN THE BODY OF WATER ON THE MOVEMENTS OF FISH
C
0112      WRITE (6,2020)
0113      2020 FORMAT (/)
0114      READ (5,1000) NHAB
0115      READ (5,1002) HBATCT
0116      WRITE (6,2012) HBATCT,NHAB
0117      2012 FORMAT (1H ,5X,'FORCE OF ATTRACTION OF HABITAT PREFERENCES, HBATCT
          1= ',F10.4,/,6X,'NHAB = ',I5,/)
0118      IF (NHAB.EQ. 0) GO TO 41
0119      WRITE (6,2011)
0120      2011 FORMAT (1H ,12X,'HABITAT DISTRIBUTION COEFFICIENTS',/,6X,'HNUM',
          112X,'HBALP',12X,'HBBET',12X,'HBIQ',13X,'HBJQ',/)
0121      DO 40 I=1,NHAB
0122      READ (5,1002) HNUM(I),HBALP(I),HBBET(I),HBIQ(I),HBJQ(I)
0123      WRITE (6,2010) HNUM(I),HBALP(I),HBBET(I),HBIQ(I),HBJQ(I)
0124      40 CONTINUE
0125      41 CONTINUE

C
C.....READ IN BOUNDARY CROSSING FACTORS FOR KEEPING FISH IN THE BODY OF WATER
C
0126      READ (5,1002) (RES(I),I=1,NREG)
0127      1002 FORMAT (7E10.0)
0128      WRITE (6,2020)
0129      WRITE (6,2013) (RES(I),I=1,NREG)
0130      2013 FORMAT (1H ,5X,'BOUNDARY CROSSING FACTORS, RES(I) = ',4(P9.3,2X),/
          1)
0131      WRITE (6,2020)

C
C
C
C
C.....READ IN THE 'INERTIA' VALUE, OR TENDENCY TO CONTINUE MOVING IN SAME
C      DIRECTION
C
0132      READ (5,1002) (ERTIA(I),I=1,5)
0133      WRITE (6,2014) (ERTIA(I),I=1,5)
0134      2014 FORMAT (1H ,5X,'VALUES OF FORWARD INERTIA, ERTIA = ',5(F6.2,1X),/)

C
C
C
C.....READ IN RANDOM NUMBER GENERATOR INITIALIZATION
C      MMM INITIATES SUBROUTINE RANSET AND SHOULD BE LEFT AT THE VALUE BELOW
C
0135      READ (5,1000) IX
0136      WRITE (6,2015) IX
0137      2015 FORMAT (1H ,5X,'RANDOM NUMBER INITIATOR, IX = ',I5,/)
0138      MMM = 2147483647

```

```

0139          CALL RANSET(MMM,IX)
C
C
C.....READ IN THE NUMBER OF OF FISH IN THE BODY OF WATER AND THE NUMBER
C          OF STEPS EACH TAKES
C
0140          READ(5,1000) NFISH,NSTEPS
0141          WRITE(6,2016) NFISH, NSTEPS
0142          2016 FORMAT(1H ,5X,'NUMBER OF FISH IN BODY OF WATER, NFISH = ',I5, '//,
          16X,'NUMBER OF STEPS EACH FISH IS ALLOWED TO TAKE, NSTEPS = ',I5,
          2//)
C
C.....READ IN IPLOT=1 IF FISH PATHS ARE TO BE PLOTTED, IPLOT=0 OTHERWISE
C
C.....READ IN ISOTH=1 IF ISOTHERMS ARE TO BE PLOTTED, ISOTH=0 OTHERWISE
C
0143          READ(5,1000) IPLOT, ISOTH
0144          IF(IPLOT .NE. 1) GO TO 45
0145          WRITE(6,2017)
0146          2017 FORMAT(1H ,5X,'FISH PATHS ARE PLOTTED', '//)
0147          45 CONTINUE
0148          IF(ISOTH .NE. 1) GO TO 47
0149          WRITE(6,2018)
0150          2018 FORMAT(1H ,5X,'ISOTHERMS ARE PLOTTED', '//)
0151          47 CONTINUE
C
C.....READ IN THE MINIMUM AND MAXIMUM ISOTHERMS TO BE PLOTTED, AS WELL AS
C          THE TEMPERATURE INTERVALS BETWEEN THEM
C
0152          IF(ISOTH .EQ. 0) GO TO 59
0153          READ(5,1004) TEML,TEMH,TEMINT
0154          WRITE(6,2019) TEML,TEMH,TEMINT
0155          2019 FORMAT(1H ,5X,'MINIMUM ISOTHERM PLOTTED, TEML = ',F10.4, '//,6X,
          1          'MAXIMUM ISOTHERM PLOTTED, TEMH = ',F10.4, '//,6X,
          2          'DISTANCE BETWEEN ISOTHERMS, TEMINT = ',F10.4, '//)
0156          59 CONTINUE
C
C
C
0157          CALL PLOTT
C
C.....CHOOSE AN INITIAL POSITION AND SENSE OF DIRECTION OF THE FISH RANDOMLY
C
0158          60 CONTINUE
0159          ISV = 1
0160          NDIST = 0
0161          SHOR = NHOR
0162          SVER = NVER
0163          65 CONTINUE
0164          D = URAND(DUMY)
0165          DH = SHOR*D
0166          D = URAND(DUMY)
0167          DV = SVER*D
0168          ISTRT = DH

```

```

0169      JSTRT = DV
0170      IPAST = ISTRT
0171      JPAST = JSTRT
0172      SAVI (ISV) = ISTRT
0173      SAVJ (ISV) = JSTRT
0174      D = URAND(DUMY)
0175      FRAC = .125
0176      DO 80 I=1,3
0177      DO 80 J=1,3
0178      IF (I .EQ. 2 .AND. J .EQ. 2) GO TO 80
0179      IF (D .LT. FRAC) GO TO 85
0180      FRAC = FRAC + .125
0181      80 CONTINUE
0182      85 CONTINUE
0183      IDR = I
0184      JDR = J
0185      IPRES = ISTRT - 2 + IDR
0186      JPRES = JSTRT - 2 + JDR
0187      IF (GRID(ISTRT,JSTRT) .LT. 2.0) GO TO 65
C
C
C.....BEGINNING OF TIME ITERATION
C
0100      100 CONTINUE
C
C.....DETERMINE CURRENT TEMPERATURE AND DISSOLVED OXYGEN
C
0189      IF (GRID(IPRES,JPRES) .GT. 3.0) GO TO 105
0190      TEMP = (GRID(IPRES,JPRES) - 2.0)*1000.
0191      LTEM = TEMP
0192      STEM = LTEM
0193      DOX = (TEMP-STEM) * 100.
0194      GO TO 107
0195      105 CONTINUE
0196      TEMP = (GRID(IPRES,JPRES) - 3.0)*1000.
0197      LTEM = TEMP
0198      STEM = LTEM
0199      DOX = (TEMP-STEM) * 100.
0200      107 CONTINUE
0201      TEMP = 0.1*TEMP
0202      TDIFF = ABS(TEMP - TEMPRF)
0203      DDIFF = DOX - DOXMIN
0204      SJPAST = JPAST
0205      SIPAST = IPAST
0206      SISTRT = ISTRT
0207      SJSTRT = JSTRT
0208      SIPRES = IPRES
0209      SJPRES = JPRES
C
C.....DETERMINE CURRENT FOOD AVAILABILITY
C
0210      IF (NFOOD .EQ. 0) GO TO 111
0211      FOOD = 0.0
0212      DO 110 I=1,NFOOD
0213      DISTI = ABS(SIPRES-PDIO(I))
0214      DISTJ = ABS(SJPRES - FDJO(I))
0215      EXPARG = EXP(-FDALP(I) *DISTI - FDBET(I) *DISTJ)
0216      FOOD = FOOD + FDNUM(I) *EXPARG

```

```

0217       110 CONTINUE
0218       111 CONTINUE
C
C.....DETERMINE CURRENT HABITAT FAVORABILITY
C
0219       IF(NHAB .EQ. 0) GO TO 116
0220       HAB = 0.0
0221       DO 115 I=1,NHAB
0222         DISTI = ABS(SIPRES - HBIQ(I))
0223         DISTJ = ABS(SJPRES - HBJQ(I))
0224         EXPARG = EXP(-HBALP(I)*DISTI - HBBET(I)*DISTJ)
0225         HAB = HAB + HENUM(I)*EXPARG
0226       115 CONTINUE
0227       116 CONTINUE
C
C
C.....CALCULATION OF THE EFFECTS OF INERTIA OF FORWARD MOTION, THE EFFECTS
C OF TEMPERATURE AND DISSOLVED OXYGEN GRADIENTS, AND THE INFLUENCE OF
C SPATIAL FOOD DISTRIBUTION AND HABITAT PREFERENCES
C
0228       DO 200 I=1,3
0229       DO 200 J=1,3
0230       IPI = IPRES + I - 2
0231       JPI = JPRES + J - 2
0232       SIPI = IPI
0233       SJPI = JPI
C
C.....INERTIA EFFECTS
C
0234       SI = I
0235       SJ = J
0236       SIDR = IDR
0237       SJDR = JDR
0238       C = ABS(SI-SIDR) + ABS(SJ-SJDR)
0239       IF(C .GT. 0.0) GO TO 123
0240       IERT = 1
0241       GO TO 129
0242       123 IF(C .GT. 1.0) GO TO 124
0243       IERT = 2
0244       GO TO 129
0245       124 IF(C .GT. 2.0) GO TO 125
0246       IERT = 3
0247       GO TO 129
0248       125 IF(C .GT. 3.0) GO TO 126
0249       IERT = 4
0250       GO TO 129
0251       126 IF(C .GT. 4.0) GO TO 127
0252       IERT = 5
0253       GO TO 129
0254       127 IERT = 5
0255       129 CONTINUE
0256       DIR(I,J) = ERTIA(IERT)
C
C.....BOUNDARY EFFECTS
C
0257       IF(IPI .GE. NHOR .OR. JPI .GE. NVER) GO TO 60
0258       IF(IPI .LE. 0 .OR. JPI .LE. 0) GO TO 60
0259       IGRIDF = GRID(IPI,JPI)

```

```

0260      BDY(I,J) = RES(IGRIDP)
C
C
C.....EFFECTS OF FOOD ATTRACTION
C
0261      IF(NFOOD .EQ. 0) GO TO 141
0262      FOODA = 0.0
0263      DO 140 K=1,NFOOD
0264          DISTI = ABS(SIPI - FDIO(K))
0265          DISTJ = ABS(SJPI - FDJO(K))
0266          EXPARG = EXP(-FDALP(K)*DISTI - FDBET(K)*DISTJ)
0267          FOODA= FOODA+ FDNUM(K)*EXPARG
0268      140 CONTINUE
0269          IF(FOODA .LT. FOOD) GO TO 141
0270          FDR(I,J) = FEATCT
0271          GO TO 142
0272      141 CONTINUE
0273          FDR(I,J) = 0.0
0274      142 CONTINUE
C
C.....EFFECTS OF HABITAT PREFERENCE
C
0275      IF(NHAE .EQ. 0) GO TO 151
0276      HABA = 0.0
0277      DO 150 K=1,NHAB
0278          DISTI = ABS(SIPI - HBIQ(K))
0279          DISTJ = ABS(SJPI - HBJQ(K))
0280          EXPARG = EXP(-HBALP(K)*DISTI - HBBET(K)*DISTJ)
0281          HABA= HABA+ HBNUM(K)*EXPARG
0282      150 CONTINUE
0283          IF(HABA .LT. HAB) GO TO 151
0284          HDR(I,J) = HBATCT
0285          GO TO 152
0286      151 CONTINUE
0287          HDR(I,J) = 0.0
0288      152 CONTINUE
C
C.....TEMPERATURE AND DISSOLVED OXYGEN EFFECTS
C
0289      TDR(I,J) = 0.0
0290      IF(GRID(IPI,JPI) .GT. 3.0) GO TO 180
0291      TEMP = (GRID(IPI,JPI) - 2.0)*1000.
0292      LTEM = TEMP
0293      STEM = LTEM
0294      DOX = (TEMP-STEM)*100.
0295      GO TO 185
0296      180 CONTINUE
0297      TEMP = (GRID(IPI,JPI) - 3.0)*1000.
0298      LTEM = TEMP
0299      STEM = LTEM
0300      DOX = (TEMP-STEM)*100.
0301      185 CONTINUE
0302      TEMP = 0.1*TEMP
0303      TDIFFA = ABS(TEMP- TEMPRF)
0304      DDIFFA = DOX - DOXMIN
0305      IF(TDIFFA .GT. TDIFFF) GO TO 190
0306      TDR(I,J) = TEMPOR
0307      GO TO 191

```

```

0308      190 CONTINUE
0309          TDR(I,J) = 0.0
0310      191 CONTINUE
0311          IF(DDIFF .GT. 0.0) GO TO 199
0312          DDR(I,J) = 0.0
0313          IF(DDIPFA .LT. 0.0) GO TO 195
0314          DDR(I,J) = DOXFOR
0315          GO TO 199
0316      195 CONTINUE
0317          DDR(I,J) = 0.0
0318      199 CONTINUE
0319      200 CONTINUE
C
C
C.....USING THE ABOVE CALCULATIONS TO OBTAIN THE PROBABILITIES FOR THE
C      DIRECTION OF THE NEXT STEP IN EACH OF THE EIGHT POSSIBLE DIRECTIONS.
C
0320          VSUM = 0.0
0321          DO 210 I=1,3
0322          DO 210 J=1,3
0323          V(I,J) = (1.0 + TDR(I,J) + DIR(I,J) + DDR(I,J) + FDR(I,J)
1 + HDR(I,J))*BDY(I,J)
0324          IF(I .EQ. 2 .AND. J .EQ. 2) GO TO 210
0325          VSUM = VSUM + V(I,J)
0326      210 CONTINUE
0327          Y = URAND(DUMY)
0328          PER = 0.0
0329          DO 250 I=1,3
0330          DO 250 J=1,3
0331          IF(I .EQ. 2 .AND. J .EQ. 2) GO TO 245
0332          PER = (V(I,J)/VSUM) + PER
0333          IF(Y .GT. PER) GO TO 245
0334          GO TO 251
0335      245 CONTINUE
0336      250 CONTINUE
0337      251 CONTINUE
0338          IP = IPRES
0339          JP = JPRES
0340          IPRES = IPRES - 2 + I
0341          JPRES = JPRES - 2 + J
0342          IPAST = IP
0343          JPAST = JP
0344          IDR = I
0345          JDR = J
C
C
C.....COMPUTING THE DISTRIBUTION OF FISH
C
0346          NDIST = NDIST + 1
0347          IF(NDIST .GT. NSTEPS) GO TO 280
0348          IF(IPRES .LE. 0 .OR. JPRES .LE. 0) GO TO 280
0349          IF(IPRES .GT. NHOR .OR. JPRES .GT. NVER) GO TO 280
0350          ISV = ISV + 1
0351          SAVI(ISV) = IPRES
0352          SAVJ(ISV) = JPRES
0353          GO TO 100
0354      280 CONTINUE
0355          P1 = URAND(DUMY)

```

```

0356      F2 = URAND(DUMY)
0357      SIP = IPRES
0358      SJP = JPRES
0359      SPRESI(1) = SIP - 0.5 + P1
0360      SPRESJ(1) = SJP - 0.5 + P2
0361      CALL MARKER(1)
0362      CALL SCLPIC(.125)
0363      CALL CURVE(SPRESI,SPRESJ,1,-1)
0364      NSV = NSV + 1
0365      TEM = 0.0
0366      DO 285 I=1,80
0367      TEMPLU = TEM + 0.5
0368      IF(TEMP .LT. IEM .OR. TEMP .GT. TEMPLU) GO TO 284
0369      TEMSAV(I) = TEMSAV(I) + 1.0
0370      284 CONTINUE
0371      TEM = TEM + 0.5
0372      285 CONTINUE
0373      IF(IPLOT .EQ. 0) GO TO 290
0374      CALL CURVE(SAVI,SAVJ,ISV,0)
0375      290 CONTINUE
0376      IF(NSV .EQ. NFISH) GO TO 300
0377      GO TO 60
0378      300 CONTINUE

C
C.....PLOTTING OF ISOTHERMS
C
0379      TEM = IEM1
0380      NK = (TEMH - TEM1)/TEMINT
0381      KB = 1
0382      DO 400 K=1,NK
0383      SJ = 0.0
0384      KA = 0
0385      DO 350 J=1,NVER
0386      SJ = J
0387      ARGT = (40000.- (6*(SJ-55.))**2+10000.)*TEM)/(TEM*0.8)
0388      IF(ARGT .LE. 0.0) GO TO 320
0389      SI = 85. - SQRT(ARGT)
0390      IF(SI .LT. 1.0 .OR. SI .GT. 90.0) GO TO 320
0391      I = SI
0392      IF(GRID(I,J) .LT. 2.0) GO TO 320
0393      KA = KA + 1
0394      ARAX(KA) = SI
0395      ARAY(KA) = SJ
0396      GO TO 345
0397      320 CONTINUE
0398      IF(KA .EQ. 0) GO TO 340
0399      CALL CURVE(ARAX,ARAY,KA,0)
0400      KA = 0
0401      KB = KB + 1
0402      340 CONTINUE
0403      345 CONTINUE
0404      SJ = SJ + 1.0
0405      350 CONTINUE
0406      TEM = TEM + TEMINT
0407      400 CONTINUE
0408      401 CONTINUE
0409      CALL ENDPL(1)
0410      CALL DCNEPL

```



```
C
C.....PLOTING OF TEMPERATURE HISTOGRAM
C
0411      DO 500 I=1,80
0412      500 CONTINUE
0413      CALL HIST (TEMSAV)
0414      CALL DONEPL
0415      STOP
0416      END
```

```

0001      FUNCTION URAND(FRAN)
C
C
C.....E.J.MCGARTH AND D.C.IRVING. 1975. TECHNIQUES FOR EFFICIENT MONTE CARLO
C      SIMULATION. VOL. 2. RANDOM NUMBER GENERATION FOR SELECTED PROBABILITY
C      DISTRIBUTIONS. ORNL-RSIC-38
C
0002      COMMON/NIRNG/RAN(10),GEN(10),NWRD,BASE,MOD,FBASE,FMOD
0003      DIMENSION SUM(10)
0004      INTEGER RAN,GEN,BASE,CARRY,SUM,PROD,HPROD
0005      DO 30 IS=1,NWRD
0006      30    SUM(IS)=0.
0007          DO 1 IG=1,NWRD
0008          N2=NWRD-IG+1
0009          DO 1 IE=1,N2
0010          IS=IR+IG-1
0011          PROD=RAN(IR)*GEN(IG)
0012          HPROD=PROD/BASE
0013          LPROD=PROD-HPROD*BASE
0014          SUM(IS)=SUM(IS)+LPROD
0015          IF (IS.LT.NWRD) SUM(IS+1)=SUM(IS+1)+HPROD
0016      1    CONTINUE
0017          N2=NWRD-1
0018          DO 5 IS=1,N2
0019          CARRY=SUM(IS)/BASE
0020          SUM(IS)=SUM(IS)-CARRY*BASE
0021          SUM(IS+1)=SUM(IS+1)+CARRY
0022      5    CONTINUE
0023          SUM(NWRD)=SUM(NWRD)-MOD*(SUM(NWRD)/MOD)
0024          DO 20 IS=1,NWRD
0025          20    RAN(IS)=SUM(IS)
0026          FRAN=SUM(1)
0027          DO 10 IS=2,NWRD
0028          10    FRAN=FRAN/FBASE+SUM(IS)
0029          FRAN=FRAN/FMOD
0030          URAND=FRAN
0031          RETURN
0032          END

```

```

0001          SUBROUTINE RANSET (MAXINT,NSTRT)
C
C
C.....E.J.MCGARTH AND D.C.IRVING. 1975. TECHNIQUES FOR EFFICIENT MONTE CARLO
C SIMULATION. VOL. 2. RANDOM NUMBER GENERATION FOR SELECTED PROBABILITY
C DISTRIBUTIONS. ORNL-RSIC-38
C
C
0002          COMMON/MIRNG/ RAN(10),GEN(10),NWRD,BASE,MOD,FBASE,FMOD
0003          INTEGER RAN,GEN,BASE,CARRY,REM
0004          MAXI=MAXINT/4
0005          IB=0
0006          BASE=1
0007          99  IF (BASE.GT.MAXI) GO TO 100
0008          BASE=BASE*4
0009          IB=IB+1
0010          GO TO 99
0011          100 BASE=2**IB
0012          FBASE=BASE
0013          NWRD=47/IB+1
0014          REM=47-IB*(NWRD-1)
0015          MOD=2**REM
0016          FMOD=MCD
0017          DO 101 N=1,10
0018          RAN(N)=0
0019          101  GEN(N)=0
0020          GEN(1)=5
0021          DO 200 I=1,14
0022          CARRY=0
0023          DO 190 N=1,NWRD
0024          GEN(N)=GEN(N)*5+CARRY
0025          CARRY=0
0026          IF (GEN(N).LT.BASE) GO TO 190
0027          CARRY=GEN(N)/BASE
0028          GEN(N)=GEN(N)-BASE*CARRY
0029          190  CONTINUE
0030          200  CONTINUE
0031          NSTART=NSTRT
0032          IF (NSTART.LE.0) NSTART=2001
0033          NSTART=2*(NSTART/2)+1
0034          DO 300 N=1,NWRD
0035          NTEMP=NSTART/BASE
0036          RAN(N)=NSTART-NTEMP*BASE
0037          300  NSTART=NTEMP
0038          RETURN
0039          END

```

```
0001      SUBROUTINE PLCTT
0002      COMMON/FINBLK/NFIN,NREGP
0003      COMMON/DRAW/ARRAYX(50,50),ARRAYY(50,50)
0004      COMMON/THERM/KA,KB,NPTYP
0005      DIMENSION XX(500),YY(500)
0006      CALL CALCF
0007      CALL BGNPL(-1)
0008      CALL PAGE(14.,11.)
0009      CALL TITLE(' FISH DISTRIBUTIONS$',100,'LENGTH',6,'DEPTH',5,10.,6.)
0010      CALL GRAP(0.,'SCALE',100.,0.,'SCALE',65.)
0011      XMAX = 100.
0012      XMIN = 0.0
0013      YMAX = 60.
0014      YMIN = 0.0
0015      I = 1
0016      DO 50 J=1,NREGP
0017      XX(J) = ARRAYX(I,J)
0018      YY(J) = ARRAYY(I,J)
0019      50 CONTINUE
0020      CALL CURVE(XX,YY,NREGP,0)
0021      RETURN
0022      END
```

```
0001      SUBROUTINE HIST(TEMSAV)
0002      DIMENSION TEMSAV(100)
0003      DIMENSION CLASS(100),FREQ(100)
0004      COMMON/NFBLOK/NFISH
0005      SNFISH = NFISH
0006      NCLASS = 40
0007      NDAY = 1
0008      FRQMX = 1.0
0009      XSTEP = 2.0
0010      DO 10 I=1,40
0011      SI = I
0012      CLASS(I) = 20. + 0.5*SI
0013      FREQ(I) = TEMSAV(I+40)/SNFISH
0014      10 CONTINUE
0015      CALL BGNPL(-1)
0016      CALL PAGE(8.,11.)
0017      CALL TITLE('TEMPERATURE SELECTIONS',100,
0018      1'TEMPERATURES',100,'NUMBER OF FISHS',100,7.,7.)
0019      CALL XAXANG(45.)
0020      CALL GRAP(20.,XSTEP,40.,0.,'SCALE',FRQMX)
0021      CALL INTNO(NDAY,9.50,9.6)
0022      BWIDTH = 7./(NCLASS-1)
0023      CALL BARS(BWIDTH)
0024      CALL CURVE(CLASS,FREQ,NCLASS,0)
0025      CALL ENDPL(0)
0026      RETURN
0027      END
```

INTERNAL DISTRIBUTION

- | | | | |
|--------|-------------------|--------|-----------------------------|
| 1. | S. M. Adams | 48. | H. Postma |
| 2-11. | S. I. Auerbach | 49. | D. E. Reichle |
| 12. | L. W. Barnthouse | 50. | C. R. Richmond |
| 13. | R. W. Brocksen | 51. | B. Schaich |
| 14. | R. L. Burgess | 52. | H. H. Shugart |
| 15. | D. S. Carroll | 53. | R. H. Strand |
| 16. | S. W. Christensen | 54. | E. G. Struxness |
| 17-26. | C. C. Coutant | 55. | J. S. Suffern |
| 27. | D. K. Cox | 56. | R. I. Van Hook, Jr. |
| 28-38. | D. L. DeAngelis | 57. | W. Van Winkle, Jr. |
| 39. | J. W. Elwood | 58. | D. S. Vaughan |
| 40. | W. R. Emanuel | 59. | H. Waddle |
| 41. | R. H. Gardner | 60. | J. A. Watts |
| 42. | W. F. Harris | 61. | Biology Library |
| 43. | K. D. Kumar | 62-63. | Central Research Library |
| 44. | J. B. Mankin | 64-65. | Laboratory Records Dept. |
| 45. | J. S. Mattice | 66. | Laboratory Records, ORNL-RC |
| 46. | R. B. McLean | 67. | ORNL Y-12 Technical Library |
| 47. | R. V. O'Neill | 68. | ORNL Patent Office |

EXTERNAL DISTRIBUTION

69. Richard O. Anderson, Cooperative Fishery Research Unit, University of Missouri, Columbia, MO 65201
70. C. A. Barans, South Carolina Marine Resources Research Institute, P. O. Box 12559, Charleston, SC 29412
71. Roger A. Barnhart, Cooperative Fishery Research Unit, Humboldt State University, Arcata, CA 95521
72. T. L. Beitinger, Dept. of Fisheries and Wildlife, Texas A & M University, College Station, TX 77840
73. D. H. Bennett, Virginia Polytechnic Institute and State University, Blacksburg, VA 24060
74. Theodore C. Bjornn, Cooperative Fishery Research Unit, University of Idaho, Moscow, ID 83843
75. Charles F. Bryan, Cooperative Fishery Research Unit, Louisiana State University, Baton Rouge, LA 70803
76. Robert L. Butler, Cooperative Fishery Research Unit, Pennsylvania State University, University Park, PA 16802
77. D. S. Cherry, Center for Environmental Studies, Virginia Polytechnic Institute and State University, Blacksburg, VA 24060
78. J. P. Clugston, Southeast Reservoir Investigations, Clemson University, Clemson, SC 29631
79. Daniel W. Coble, Cooperative Fishery Research Unit, University of Wisconsin, Stevens Point, WI 54481
80. A. E. Dizon, National Marine Fisheries Service, Honolulu, HI 96800

81. L. Farges, IAEA, Karnter Ring 11, P. O. Box 590 A-1011, Vienna, Austria
82. R. Don Estes, Cooperative Fishery Research Unit, Tennessee Technological University, Cookeville, TN 38501
83. F. E. J. Fry, University of Toronto, Toronto, Ontario, Canada
84. J. R. Gammon, Biology Department, DePauw University, Greencastle, IN 46135
85. J. W. Gibbons, Savannah River Ecological Laboratory, Aiken, SC 29801
86. J. Gift, Ichthyological Associates, Inc., New Jersey Marine Ecological Study, 3201 Bayshore Ave., Brigantine, NJ 08203
87. C. P. Goodyear, Power Plant Team, U. S. Department of the Interior, Ann Arbor, MI 48105
88. Richard W. Gregory, Cooperative Fishery Research Unit, Montana State University, Bozeman, MT 59715
89. Bernard L. Griswold, Cooperative Fishery Research Unit, Ohio State University, Columbus, OH 43210
90. Donald C. Hales, Cooperative Fishery Research Unit, South Dakota State University, Brookings, SD 57006
91. Richard W. Hatch, Cooperative Fishery Research Unit, University of Maine at Orono, Orono, ME 04473
92. Melvin T. Huish, Cooperative Fishery Research Unit, North Carolina State University, Raleigh, NC 27606
93. P. R. Kamath, Bhabha Atomic Research Center, Trombay, Bombay, India
94. J. R. M. Kelso, Canada Centre for Inland Waters, Burlington, Ontario, L7R 4A6, Canada
95. W. C. Leggett, Biology Department, McGill University, Montreal, Canada
96. John A. Maciolek, Cooperative Fishery Research Unit, University of Hawaii, Honolulu, HI 96822
97. J. J. Magnuson, Laboratory of Limnology, Univ. of Wisconsin, Madison, WI 53703
98. O. Eugene Maughan, Cooperative Fishery Research Unit, Oklahoma State University, Stillwater, OK 74074
99. James A. McCann, Chief, Division of Fishery Research, U. S. Department of the Interior, Fish and Wildlife Service, Washington, DC 20240
100. R. W. McCauley, Biology Department, Wilfred Laurier University, Waterloo, Ontario, N2L 3C5, Canada
101. William J. McConnell, Cooperative Fishery Research Unit, Colorado State University, Fort Collins, CO 80521
102. J. H. McCormick, USEPA, Environmental Research Laboratory, Duluth, MN 55800
103. J. McMaho, Atomic Energy of Canada, Ltd., Chalk River, Canada
104. J. W. Meldrin, Ichthyological Associates, 100 S. Cass St., Middletown, DE 19709
105. D. Miller, USEPA, Environmental Research Laboratory, Narragansett, RI 02882
106. Robert J. Muncy, Cooperative Fishery Research Unit, Iowa State University, Ames, IA 50010

107. W. Neill, Dept. of Fisheries and Wildlife, Texas A & M University, College Station, TX 77840
108. John G. Nickum, Cooperative Fishery Research Unit, Cornell University, Ithaca, NY 14850
109. Garland B. Pardue, Cooperative Fishery Research Unit, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061
110. John S. Ramsey, Cooperative Fishery Research Unit, Auburn University, Auburn, AL 36830
111. Roger J. Reed, Cooperative Fishery Research Unit, University of Massachusetts, Amherst, MA 01002
112. Robert E. Reinert, Cooperative Fishery Research Unit, University of Georgia, Athens, GA 30601
113. J. M. Reutter, F. T. Stone Laboratory, Put-In-Bay, OH 43456
114. W. Reynolds, Biology Dept., Pennsylvania State University, Wilkesbarre, PA 18708
115. F. P. Richards, Ecological Analysts, Melville, NY 11746
116. W. Schikarski, Nuclear Research Center, Karlsruhe, Federal Republic of Germany
117. M. Schneider, Ecosystems Dept., Battelle Northwest, Richland, WA 99352
118. Carl B. Schreck, Cooperative Fishery Research Unit, Oregon State University, Corvallis, OR 97331
119. S. A. Spigarelli, Argonne National Laboratory, Argonne, IL 60439
120. J. R. Stauffer, Gunter Hall, Frostburg State College, Frostburg, MD 21532
121. Jerry C. Tash, Cooperative Fishery Research Unit, University of Arizona, Tucson, AZ 85721
122. M. Van deu Avyle, Cooperative Fisheries Unit, Tennessee Tech. University, Cookeville, TN 38501
123. L. Wilson, Dept. of Forestry, Wildlife, and Fisheries, University of Tennessee, Knoxville, TN 37916
124. Richard R. Whitney, Cooperative Fishery Research Unit, University of Washington, Seattle, WA 98105
125. Richard S. Wydoski, Cooperative Fishery Research Unit, Utah State University, Logan, UT 84321
126. Research and Technical Support Division, DOE-ORO.
- 127-153. Technical Information Center, Oak Ridge, TN 37830