DOE Technical Report No. DOE/PC/88921-15

Fifteenth Quarterly Report on Research Grant No. DE-FG22-88PC88921

Title: SPIN-MAPPING OF COAL STRUCTURES WITH ESE AND ENDOR

Principal Investigators: R. L. Belford and R. B. Clarkson

Institution: University of Illinois at Urbana-Champaign

Date: June 1, 1992

US/DOE Patent Clearance is not required prior to publication of this document.

RECEIVED

MAY 03 1993

ABSTRACT

OSTI

Advanced electron paramagnetic resonance (EPR) techniques — ENDOR, ESE, and VHF-EPR — are used to probe the molecular structure and surface properties of coals and of model systems. Typically, the model compounds under investigation and their analogues are found in coals as stable free radicals which give rise to an EPR signal. In some cases, as in the work reported this quarter, the model compounds are selected because they have some characteristic, such as a particular functional group or heteroatom which may be found in coals, which fits them to serve as test materials for methods development.

While this research group continues to catalogue and analyze EPR and ESE spectra of coal and coal maceral samples, it has very recently also made progress in technique development and application. The work reported for this quarter is such a development - an examination of the applicability of W-band EPR and of low-frequency ESE (electron spin echo) spectroscopy to characterization of heteroatoms in a heterogeneous environment. Nitroaromatic radicals catalytically produced in the presence of oxygen as asymmetric surface species on a particulate catalyst surface was probed by both techniques, and it is shown that the electronic structure in the vicinity of the nitrogen and oxygen atoms can be elucidated in this way. By use of isotopically labeled molecular oxygen, the possible role of oxygen in the catalysis and in the binding of the nitroaromatic species to the surface are explored. The following report summarizes the highlights of this study.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

REPORT

NITROGEN COUPLINGS OF NITROAROMATIC RADICALS ON POWDERED CATALYST SURFACES. - Low-frequency (S-band) ESEEM (electron spin echo envelope modulation) from EPR- active *m*-dinitrobenzene on gamma-alumina.

Meta-dinitrobenzene (*m*-DNB) radical anions (*m*-DNB $\cdot^-$) were generated in disordered state by catalytic action of oxygen with *m*-DNB (both N-14 and N-15 versions) on activated surface sites of activated γ -alumina powder. Companion studies of nitrobenzene on activated γ -alumina powder and *m*-DNB on activated MgO powder are under way. Electron Spin Echo Envelope Modulation (ESEEM) spectra of these species at low magnetic fields and frequencies (S-band, 2-4 GHz) were acquired at near-liquid-helium temperatures. This study concerns our DOE-PETC project in two ways. Firstly, we have been using the low-field ESE technique to probe properties of specimens of native and treated coal and coal macerals, and particularly the effects of heteroatoms such as nitrogen, oxygen, and sulfur in the coal structures. Secondly, one of our most successful and widely-used method for the production of aromatic radical species as EPR-active models for similar species in coals has been the oxygen-catalyzed production from parent aromatic neutral species on activated metal-oxide particulates. The focus in all these studies is the extraction of information about the heteroatom, be it sulfur, nitrogen, or oxygen. Thus the study summarized here is a benchmark investigation and methods study.

The three-pulse ESEEM experiments were performed at four different microwave frequencies from 2 to 4 GHz: 3.85, 3.24, 3.01, and 2.81 GHz. These ESEEM patterns were processed, Fourier-transformed, and mapped against the S-band spectrometer frequency to produce radiofrequency-domain nitrogen nuclear modulation line positions vs. microwave frequency. Spectral simulations were performed with NANGSEL, a computer program written by Dr. Jeffrey B. Cornelius, and modified in this study to accommodate more general spin systems. The calculations, employing a powder-average algorithm, permit anisotropic *g*-tensors and small anisotropic super-hyperfine coupling tensors, which are suitable in our case. The patterns clearly were from only one nitrogen atom in the two-nitrogen-atom *m*-DNB $\cdot^-$ species. Very-high-frequency (95 GHz) EPR spectroscopy previously done in this laboratory on the same species, prepared with N-14 or N-15, had shown much larger resolved splittings from the other nitrogen atom, showing that this supposedly symmetrical species cannot be at all symmetric on this catalytic surface. Clearly, one nitrogen has most of the electron spin density and presumably is attached to and greatly affected by the oxide surface, while the other nitrogen, the one being observed in the ESEEM experiments, is only weakly coupled to the electron spin, is remote from the surface, and is presumably very like the nitrogen atoms in neutral aromatic nitro groups. Using the N-15 ESEEM patterns, we were able to analyze for the hyperfine and Zeeman parameters characterizing the remote N atom. Then, from the ESEEM patterns of the species containing the quadrupolar N-14 nuclei, we were able to extract the nitrogen quadrupole coupling parameters. There is clearly some sensitivity to catalyst surface effects, since aging of the surfaces produced some modifications in the patterns.

Derived coupling parameters for the remote nitrogen nucleus in the dinitrobenzene species on alumina are summarized in Table I. Generally, the quadrupole coupling parameters revealed local electron density around the nitrogen atom much like neutral aromatic nitro species in other environments, giving us some confidence that the catalytic surface method of producing these model hetero compounds need not produce pathologically different species than those which might be found in other states (eg., in coal).

One matter of scientific interest is the use of the so-called exact cancellation condition by which one may tune an ESEEM spectrometer to provide nuclear modulations which are essentially pure nuclear quadrupole frequencies. This phenomenon arises as follows. The magnitudes of the two electron spin manifolds will go in opposite directions as the external magnetic field changes. This means that for quadrupolar nuclei ($I \geq 1$), an external magnetic field of magnitude B_0 can be chosen such that the nuclear Zeeman interaction cancels the super-hyperfine interaction in one of the two electron spin states on resonance. The ensuing nuclear energy levels, in the electron spin state in which the cancellation occurs, correspond to the pure nuclear quadrupole states, $\hat{H}_n^{\text{eff}} = \mathbf{I} \cdot \mathbf{P} \cdot \mathbf{I}$. This permits a direct observation of the NQR transitions, the frequencies of which can be used to calculate the quadrupole parameters (e^2qQ and η). In the S-band range, we have been able to achieve this condition. Moreover, in the process of doing so, we have developed a diagnostic procedure for locating this exact cancellation condition. The procedure involves making plots of certain modulation frequencies and intensities against spectrometer frequency. There are stationary points (extrema in some modulations) which signal the exact cancellation and at which, the radiofrequencies are the same as would be obtained in a pure quadrupole resonance experiment. By tracking the S-band ESEEM modulations with the microwave frequencies, we find that the "exact cancellation" condition can be achieved in the presence of small g-anisotropy.

An important, hitherto unanswered question in the catalytic oxygen-aided production of aromatic radical surface species has been this: "What is the role and fate of the oxygen?" To our knowledge, it has never been identified in surface-bound radical systems. To address this, we prepared the same surface species with molecular oxygen prepared with the O-17 isotope. In the S-band ESEEM patterns, we found modulations corresponding to the O-17 nuclear g-value and were able use these patterns to define an effective distance from the surface-bound nitrogen atom to the O-17 in molecular oxygen as well as to the aluminum atom in the surface.

Table I. ESEEM parameters for the weakly coupled nitrogen-14 atom of *m*-DNB radical formed on activated gamma-alumina surface sites.

Surface Radical Conditioning	Experimental or Calculated from N-15 Data	Hyperfine Coupling A_{iso} (MHz)	Quadrupole Coupling e^2qQ (MHz)	Asymmetry Parameter η
Freshly-made	exp.		1.243	0.36
	cal.	0.65	1.245	0.36
Aged for ca. 1 year	exp.		1.219	0.37
	cal.	0.65	1.220	0.37

Results of this study will be published in a series of journal articles (W. Shi, R. L. Belford, R. B. Clarkson, et al.; manuscripts in preparation) and in the doctoral dissertation of Wenjun Shi (W. Shi, PhD Thesis, Department of Chemistry, University of Illinois, Urbana, IL, 1993).

END

DATE
FILMED

5 / 20 / 93