LAMONT-DOHERTY GEOLOGICAL OBSERVATORY
OF
COLUMBIA UNIVERSITY
PALISADES, NEW YORK 10964

MASTER

REPORT SUBMITTED TO

ACQUISITION AND ASSISTANCE-OPERATIONS DIVISION

U. S. DEPARTMENT OF ENERGY

NATURAL CATALYSIS OF THE $\text{CO}_2 - \text{HCO}_3^-$

REACTION IN SURFACE SEA WATER

Final Report
February 1981

Principal Investigator:

W.S. Broecker

Co-Investigator:

J. Shepherd

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

DISCLAIMER

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MSW

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

REPORT

Contract No. 79 EV10242

Natural Catalysis of the $\text{CO}_2 - \text{HCO}_3^-$

Reaction in Surface Sea Water

Berger and Libby (1969) proposed that the exchange rate of CO_2 between ocean and atmosphere might be enhanced by the same catalyst (carbonic anhydrase) that operates in our lungs. They envisioned the loss of this catalyst from marine animals as the sea water source and proposed that the effect would be larger in highly productive waters than in low productivity waters. Although their idea received little comment or interest from the oceanographic community it has never been disproven. New interest in it was engendered by the following observations.

1) The mean CO_2 invasion rate calculated from $\text{Rn}^{222}/\text{Ra}^{226}$ measurements in the surface sea yields a somewhat lower value (~30%) than that obtained from the distribution of bomb radiocarbon between air and sea.

2) The large difference between the bomb ^{14}C inventory in the equatorial and temperate water columns and in the excess surface water $^{14}\text{C}/\text{C}$ ratio (i.e., bomb ^{14}C content) of these two areas has focused interest in modeling them separately. To do this the difference in their CO_2 invasion rates must be known.

In order to obtain experimental evidence in this regard we undertook to measure simultaneously the O_2 , CO_2 , $^{14}\text{CO}_2$ and Rn-222 in large recently collected sea water samples during the April 1979 FGGE expedition. The object of this exercise was to see whether the transfer rates for CO_2

or $C^{14}O_2$ were higher than those expected from the O_2 and Rn^{222} measurements. The evasion rate of radon was measured by spiking sea water samples with Ra^{226} and then following the Rn^{222}/Ra^{226} ratio in the water with time. The O_2 invasion rate was measured by following the O_2 content of sea water (stripped with N_2) with time. The CO_2 invasion rate was measured by adding acid to the sea water and following the pH, P_{CO_2} , ΣCO_2 , and alkalinity with time. The $C^{14}O_2$ uptake was estimated by observing the change in ΔC^{14} in the water samples from beginning to end of the experiment (this was done for us by Paul Quay and Minze Stuiver).

The ^{14}C results are now on hand from one eight day run of the experiment. During this run two of the four stainless steel barrels (~160% volume) were run with submersible circulating pumps to increase the gas exchange rate to near average ocean conditions (i.e., a "piston velocity" of ~2 m/day). The two remaining barrels did not have pumps and were manually stirred every two hours in order to keep the temperature and salinity as uniform as possible (piston velocity ~0.4 m/day). All four barrels were filled with surface water from 0° , $158^\circ W$ at the same time. Water from the two barrels with pumps were combined for one ^{14}C sample, likewise for the barrels without pumps.

The ^{14}C results showed surprisingly that the barrels with pumps increased in ^{14}C content from 91‰ to 105‰ while the barrels without pumps increased from 91‰ to 106‰ in spite of the lower exchange rate. Under the conditions of the experiment we would expect to see an increase of ^{14}C in the barrels with pumps of about 30‰ with no enhancement and in the barrels w/o pumps an increase of about 6‰ . The result

is not as totally perplexing to us as it may seem. A preliminary attempt to perform this experiment in 1978 using plastic barrels revealed that a source of "dead" CO_2 (no ^{14}C) must be contaminating the water. We suspected the plastic of the containers. Thus we redid the experiment in 1979 with stainless steel containers. The pumps, unfortunately have a plastic housing. Our examination of the ^{14}C and CO_2 chemistry data suggests that the odd results are caused by the pump housing.

Nevertheless, if one assumes the input of dead CO_2 into the water is constant with time, the blank can be determined and the actual exchange rates estimated. The data can be summarized as follows:

<u>Barrels</u>	<u>Average ^{222}Rn Film Thickness</u>	<u>Blank</u>	<u>Estimated ^{14}C Film Thickness</u>
1 & 4 w/pumps	65±15	~0.13 $\mu\text{moles CO}_2/\text{kg}\cdot\text{hr}$	60-80
2 & 3 no pumps	330±20	0	110-220

¹ Units are in microns of film thickness. One standard theory of gas exchange parameterizes the rate in terms of the thickness of a stagnant film of water at the air/water interface. The thicker the film, the slower the exchange.

In a nutshell, we find no evidence of a CO_2 enhancement at conditions approximating the real ocean. At slower exchange rates, however, we observe a possible increase of the CO_2 exchange rate by a factor of 2 to 3. This is in accord with present theory of inorganic enhancement (Bolin, 1961) in the sense that an enhancement, if observed, should be larger at slower exchange rates. Under the most quiescent ocean conditions exchange rates parameterized by 300 μ film thicknesses are observed. However, this result would not lead to significant changes in our estimation of CO_2 input into the ocean.