ORNL/CDIAC--49

DE93 003112

Trends '91: A Compendium of Data on Global Change Highlights

Editors

Thomas A. Boden
Robert J. Sepanski
Frederick W. Stoss

M A R C H 1 9 9 2

Carbon Dioxide
Information Analysis Center
Environmental Sciences Division
Oak Ridge National Laboratory
Oak Ridge, Tennessee 37831-6335

Publication No. ORNL/CDIAC-49
ESD Publication No. 3797

MASTER

Sponsored by the
U.S. Department of Energy
Environmental Sciences Division

REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

Contents

Preface	vii
Abstract	ix
Contributing Investigators— <i>Trends '91</i>	1
Abstract— <i>Trends '91</i>	3
Foreword— <i>Trends '91</i>	5
Acknowledgments— <i>Trends '91</i>	7
ATMOSPHERIC CO ₂	9
ATMOSPHERIC CH ₄	19
OTHER TRACE GASES	29
CO ₂ EMISSIONS	39
TEMPERATURE	49
Appendix A: Carbon Dioxide Information Analysis Center	A-1
Appendix B: Contents— <i>Trends '91</i>	B-1
Appendix C: Order Form	C-1

Preface

The Carbon Dioxide Information Analysis Center (CDIAC) at Oak Ridge National Laboratory (ORNL) has been prompted to produce the series *Trends*, a concise inventory of data in response to heightened concern about global environmental issues, in particular climate changes induced by the greenhouse effect. In 1990 CDIAC presented its data in the inaugural edition of the *Trends* series entitled, *Trends '90: A Compendium of Data on Global Change*. The data were subsequently revised and updated in the 1991 issue and the coverage of topics related to global environmental data was expanded.

Highlights contains extracts from *Trends '91* to illustrate the content, style, and presentation of data contained in the full 700-page report. *Highlights* includes a listing of the investigators contributing data for *Trends '91*. In addition, it contains the abstract, foreword, and acknowledgments, as well as the introduction and a sample data record from each of the report's five chapters. The chapters are "Atmospheric CO₂," "Atmospheric CH₄," "Other Trace Gases," "CO₂ Emissions," and "Temperature." Appendix A provides information about CDIAC and its activities related to global environmental issues. Appendix B lists the contents of the full report. An order form for obtaining a free copy of *Trends '91* is found in Appendix C.

It is hoped that this overview of *Trends '91* will stimulate your interest in the full report. Please route this summary to your colleagues and bring it to the attention of your institution's library. We would be most happy to see that they, too, receive a copy of *Trends '91*.

— Thomas A. Boden
— Robert J. Sepanski
— Frederick W. Stoss

Editors

Carbon Dioxide Information Analysis Center

March 1992

Abstract

Boden, T.A., R.J. Sepanski,^a and F.W. Stoss^a (eds.) 1992. *Trends '91: A Compendium of Data on Global Change-Highlights*. ORNL/CDIAC-049. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory,^b Oak Ridge, Tennessee, 80 pp.

This document is an excerpt of the full report and contains extracts from *Trends '91: A Compendium of Data on Global Change* to illustrate the content, style, and presentation of the data contained in the full 700-page report. *Highlights* includes a listing of the investigators contributing data for *Trends '91* and also contains the abstract, foreword, and acknowledgments, as well as the introduction and a sample data record from each of the report's five chapters: "Atmospheric CO₂," "Atmospheric CH₄," "Other Trace Gases," "CO₂ Emissions," and "Temperature." Appendix A provides information about CDIAC, Appendix B lists the contents of the full report, and Appendix C is an order form for ordering a free copy of *Trends '91*.

Keywords: Air pollution, ambient temperature, atmospheric chemistry, carbon cycle, carbon dioxide, climate change, climates, chlorofluorocarbons, Earth atmosphere, emissions, flaring, fossil fuels, geophysical surveys, global aspects, global warming, greenhouse effect, meteorology, methane, monitoring, natural gas, regional analysis, temperature monitoring, temperature surveys, trace gases.

^aUniversity of Tennessee, Energy, Environment, and Resources Center

^bThe Carbon Dioxide Information Analysis Center (CDIAC) is supported by the U.S. Department of Energy, Office of Energy Research, Office of Health and Environmental Research, Environmental Sciences Division, Carbon Dioxide Research Program. CDIAC is housed in the Environmental Sciences Division of Oak Ridge National Laboratory (ORNL) which is managed by Martin Marietta Energy Systems, Inc. for the U.S. Department of Energy under contract DE-AC05-84OR21400. *Trends '91: A Compendium of Data on Global Change-Highlights* is ORNL Publication No. ORNL/CDIAC-049 and Environmental Sciences Division Publication No. 3797.

Contributing Investigators

We extend our gratitude to the 78 individuals from 30 institutions in 11 countries who contributed data for inclusion in *Trends '91*. We also thank the laboratories, agencies, institutions, and organizations that have supported the efforts of this dedicated research community. Were it not for the cooperation and dedication of this group of researchers, this document, as well as other derived information products that support *Trends '91*, could not have been produced. Complete addresses of each investigator are provided with the data sets they have contributed.

Y. S. Korotkevich

Arctic and Antarctic Research Institute
St. Petersburg, Russia

Neil B. A. Trivett

Atmospheric Environment Service
Downsview, Ontario, Canada

R. Rosen

Atmospheric and Environmental Research, Inc.
Cambridge, Massachusetts, U.S.A.

Beatriz Navascués

Centro Meteorológico Zonal de La Coruña
(*Meteorological Center of La Coruña*)
Ciudad Jardin, Spain

Ramón Juega**Carmen Rus**

Centro Meteorológico Zonal de Sta. Cruz de Tenerife
(*Meteorological Center of Tenerife*)
Santa Cruz de Tenerife, Canary Islands, Spain

Chemical Manufacturers Association

Fluorocarbon Program
Washington, D.C., U.S.A.

André Gaudry**P. Monfray****G. Polian****Gerard Lambert**

Centre des Faibles Radioactivites
(*Center for Studies in Weak Radioactivity*)
Gif-sur-Yvette, France

David J. Beardsmore**Paul Fraser****Graeme I. Pearman****Paul Steele**

Commonwealth Scientific and Industrial Research Organization
Division of Atmospheric Research
Aspendale, Victoria, Australia

H. J. Kanter**K. Munzert****Reinhold Reiter****Rudolf Sladkovic**

Fraunhofer-Institut für Atmosphärische Umweltforschung
(*Fraunhofer Institute for Atmospheric Research*)
Garmisch-Partenkirchen, Germany

F. N. Alyea**Derek M. Cunnold**

Georgia Institute of Technology
Atlanta, Georgia, U.S.A.

A. M. Brounshtein**E. V. Faber****A. A. Shashkov**

Hydrometeorological Service of the Russia
St. Petersburg, Russia

C. S. Wong

Institute of Ocean Sciences
Sidney, British Columbia, Canada

Vincenzo Cundari**Tiziano Colombo**

Italian Meteorological Service
Sestola, Italy

László Haszpra

Központi Légtörfizikai Intézet
(*Institute for Atmospheric Physics*)
Budapest, Hungary

J. M. Barnola**J. Chappellaz****C. Lorius****D. Raynaud**

Laboratoire de Glaciologie et de Géophysique de l'Environnement
(*Laboratory of Glaciology and Environmental Geophysics*)
Saint Martin d'Heres, France

Contributing Investigators

Roy W. Spencer

*Marshall Space Flight Center
Huntsville, Alabama, U.S.A.*

R. G. Prinn

*Massachusetts Institute of Technology
Cambridge, Massachusetts, U.S.A.*

Rainer Schmitt

*Meteorological Consultant
Glashütten, Germany*

James Hansen

Reto Ruedy

Helene Wilson

*National Aeronautic and Space Administration,
Goddard Space Flight Center
New York, New York, U.S.A.*

James K. Angell

*National Oceanic and Atmospheric
Administration,
Air Resources Laboratory
Silver Spring, Maryland, U.S.A.*

Thomas J. Conway

Ed Dlugokencky

Dale A. Gillette

Thomas B. Harris

Walter D. Komhyr

Patricia M. Lang

Russell C. Martin

Ken A. Masarie

Donald W. Nelson

James T. Peterson

Pieter P. Tans

Kirk W. Thoning

Lee S. Waterman

*National Oceanic and Atmospheric
Administration,
Environmental Research Laboratories
Boulder, Colorado, U.S.A.*

Ronald G. Baldwin

Michael G. Burgin

Thomas R. Karl

*National Oceanic and Atmospheric
Administration,
National Climatic Data Center
Asheville, North Carolina, U.S.A.*

Thomas A. Boden

Gregg Marland

*Oak Ridge National Laboratory
Oak Ridge, Tennessee, U.S.A.*

S. Crawford

M. Aslam K. Khalil

Rei A. Rasmussen

*Oregon Graduate Institute of Science and
Technology
Beaverton, Oregon, U.S.A.*

Charles D. Keeling

Timothy P. Whorf

*Scripps Institution of Oceanography
La Jolla, California, U.S.A.*

Pavel Ya. Groisman

K. M. Lugina

Konstantin Ya. Vinnikov

*State Hydrological Institute
St. Petersburg, Russia*

John R. Christy

*University of Alabama
Huntsville, Alabama, U.S.A.*

G. Fischer

H. Friedli

E. Lochbrunner

Hans Lötscher

E. Moor

A. Neftel

H. Oeschger

J. Schwander

Ulrich Siegenthaler

Berhard Stauffer

*University of Bern
Bern, Switzerland*

P. Simmons

*University of Bristol
Bristol, U.K.*

Philip D. Jones

Tom M. L. Wigley

*University of East Anglia
Norwich, U.K.*

Abstract—Trends '91

Boden, T.A., R.J. Sepanski,^a and F.W. Stoss^a (eds.). 1991. *Trends '91: A Compendium of Data on Global Change*. ORNL/CDIAC-46. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 700 pp.

This document is a source of frequently used global-change data. This second issue of the *Trends* series expands the coverage of sites recording atmospheric concentrations of carbon dioxide (CO₂) and methane (CH₄), and it updates records reported in the first issue. New data for other trace atmospheric gases have been included in this issue: historical data on nitrous oxide (N₂O) from ice cores, modern records of atmospheric concentrations of chlorofluorocarbons (CFC-11 and CFC-12) and N₂O, and estimates of global emissions of CFC-11 and CFC-12. The estimates for global and national CO₂ emissions from the burning of fossil fuels, the production of cement, and gas flaring have been revised and updated. Regional CO₂ emission estimates have been added, and long-term temperature records have been updated and expanded.

Data records are presented in four- to six-page formats, each dealing with a specific site, region, or emissions species. The data records include tables and graphs; discussion of methods for collecting, measuring, and reporting the data; trends in the data; and references to literature that provides further information.

All data appearing in the document are available on digital media from the Carbon Dioxide Information Analysis Center.

Keywords: Air pollution, ambient temperature, atmospheric chemistry, carbon cycle, carbon dioxide, climate change, climates, chlorofluorocarbons, Earth atmosphere, geophysical surveys, global aspects, global warming, greenhouse effect, meteorology, methane, monitoring, regional analysis, temperature monitoring, temperature surveys, trace gases.

^aUniversity of Tennessee, Energy, Environment, and Resources Center.

Foreword—*Trends '91*

Trends '91: A Compendium of Data on Global Change is part of the Carbon Dioxide Information Analysis Center's (CDIAC's) continued effort to distribute, in an accessible format, scientific data critical to global-change issues. *Trends '91* is intended for researchers, policy makers, educators, and others interested in the observational data underlying the issues related to our changing global environment.

Response to last year's inaugural issue of *Trends* has been gratifying. About 8,000 copies were distributed to users in 150 countries. *Trends '90* has been used by researchers worldwide, but perhaps most satisfying is its use by policy makers and educators. *Trends '90* has been cited in *America's Climate Change Strategy: An Action Agenda*, distributed by the Office of the President, and cited in *Environmental Quality, 1990*, the Annual Report of the President's Council on Environmental Quality. *Trends '90* has also been used as a textbook for Earth Systems Education, a graduate education course at The Ohio State University. It is with this multifaceted user community in mind that *Trends* was originally created.

Trends '91 expands the number of sites for which atmospheric concentrations of carbon dioxide (CO₂) and methane (CH₄) are recorded, and it updates the records reported in *Trends '90*. The estimates for global and national CO₂ emissions from the burning of fossil fuels, the production of cement, and gas flaring have been revised and updated. In addition, long-term temperature records have been revised and expanded. For the first time, this report also includes estimates of regional CO₂ emissions. *Trends '91* presents new data for other trace atmospheric gases as well. Historical data from ice cores are reported for nitrous oxide (N₂O), as are modern records of atmospheric concentrations of chlorofluorocarbons (CFC-11 and CFC-12) and N₂O. Global emissions of CFC-11 and CFC-12 are also presented.

Those of you familiar with *Trends '90* will note a significant change in style and format. Because the content and format of future issues of *Trends* will continue to be largely determined by user response, we seek your comments about *Trends '91*.

One of the goals of CDIAC is to provide ready access to timely data and information pertinent to global environmental changes. CDIAC endeavors to provide comprehensive reports, numeric data packages (NDPs), computer model packages (CMPs), and other information center products and services to the worldwide user community. CDIAC's publications, including *Trends*, are frequently and extensively put to use by a diverse international audience.

CDIAC acknowledges the support of the international science community in supplying data for its NDPs and for *Trends '91*. The data in *Trends '91* are most properly referenced by citing the contributing investigators as well as the section and subsequent subsection in which the data are found. The following citation for atmospheric CO₂ concentrations from the Scripps Institution of Oceanography's continuous monitoring program at Mauna Loa is presented as an example:

Keeling, C.D., and T.P. Whorf. 1991. Atmospheric CO₂—Modern Record, Mauna Loa. pp. 12–15. In T.A. Boden, R.J. Sepanski, and F.W. Stoss (eds.), *Trends '91: A Compendium of Data on Global Change*, ORNL/CDIAC-46. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A.

Foreword—*Trends '91*

Some of the CO₂ and CH₄ data in *Trends '91* are considered preliminary. Users should pay close attention to the table footnotes to avoid misinterpretation or misrepresentation of the accuracy of some of these data. More-detailed discussions on the applications and limitations for most data sets in *Trends '91* are documented in Numerical Data Packages (NDPs), which are available on digital media from CDIAC. If you have any questions concerning data contained in this report or would like to discuss a particular aspect of the data's application, please contact us.

CDIAC produces a periodic newsletter reporting on our activities, data packages we produce, and other CO₂ and climate topics. The newsletter is free; if you would like to be on our mailing list, just contact CDIAC.

Data requests and inquiries concerning *Trends '91*, our newsletter, or any other CDIAC information products may be sent to:

Carbon Dioxide Information Analysis Center
Oak Ridge National Laboratory
Building 1000, MS-6335
P.O. Box 2008
Oak Ridge, Tennessee 37831-6335, U.S.A.

(615) 574-0390
624-0390 (FTS)
(615) 574-2232 (FAX)
624-2232 (FTS FAX)

BITNET: CDP@ORNLSTC
OMNET: CDIAC
INTERNET: CDP@STC10.CTD.ORNL.GOV

To understand and resolve complex environmental problems, the international community needs reliable access to the underlying data upon which conclusions are based. We hope you find *Trends '91* a useful document.

— Paul Kanciruk, Director
— Robert M. Cushman, Deputy Director
Carbon Dioxide Information Analysis Center

Acknowledgments—*Trends '91*

As with any collective effort, this report has been prepared with the help of many people over many months. Our greatest debt is to the contributing investigators whose data compose the core of this report.

We acknowledge another group of important people, the readers for whom *Trends* is intended. We are delighted with the response received from the publication of the first issue of *Trends*. Many of the readers' recommendations on content, format, and style have been incorporated in *Trends '91*. We will continue to consider suggestions and comments from the readers of this and future issues.

We express our gratitude to the following personnel at Oak Ridge National Laboratory and Martin Marietta Energy Systems, whose collective efforts have brought this document into being. Our deepest thanks go to Cheryl Buford from the Publications Division's Advanced Publishing Technology group. Were it not for Cheryl's specialized electronic publishing design, composition, and production skills, this document could not have been completed. We also recognize Frances Littleton, Robert Eldridge, and Larry Davis's skills in electronic publishing and appreciate their assistance in preparing this report. We are grateful for the creative talents of a team of graphic artists for their input in the style and design of *Trends '91*, with special thanks to Kim Sneed for the design of the cover and new CDIAC logo and to Mitchell Williams, Dami Rich, and Jamie Payne for their assistance in the preparation of maps and other illustrations. Our profound gratitude is extended to the staff of the Graphics Division, especially to Chuck Strayer, Paul Baker, and Ronnie Hooks, all of whom have maintained the utmost in cordiality while accommodating us and meeting difficult schedules in the printing of this document. In addition, we are indebted to Gay Marie Logsdon, from the Environmental Sciences Division Publications Office, for her editorial and design suggestions. Through her efforts, the technical aspect of the subject matter has been transformed into a text suited for a diverse audience with multidisciplinary interests in the topic of global change. We also appreciate the efforts of Nathan Schulman in editing this report. We acknowledge the original concept for *Trends* was conceived by our former CDIAC director, Michael P. Farrell.

We again thank Tom Gross, program manager with the U.S. Department of Energy's Office of Health and Environmental Research, (Environmental Sciences Division) Carbon Dioxide Research Program, for his continued support of CDIAC activities, including the production of *Trends*. Tom has long supported international exchange of scientific information among an audience with multidisciplinary interests. His visionary promotion of the value-added process of providing information has been a mainstay in developing CDIAC's services and derived-information products.

CDIAC is supported by the U.S. Department of Energy, Office of Health and Environmental Research, Environmental Sciences Division, Carbon Dioxide Research Program.

CDIAC is housed in the Environmental Sciences Division (ESD) at Oak Ridge National Laboratory (ORNL), which is managed by Martin Marietta Energy Systems, Inc., for the U.S. Department of Energy under Contract DE-AC05-84OR21400. *Trends '91* is ORNL Publication Number ORNL/CDIAC-046 and is ESD Publication No. 3746.

Atmospheric CO₂

This section of the highlights contains the introduction to the *Trends '91* chapter on atmospheric CO₂ concentrations and a sample record of data for concentrations measured at the Mauna Loa Observatory (Hawaii, U.S.A.) as provided by Charles D. Keeling and Timothy P. Whorf of the Scripps Institution of Oceanography. The sample illustrates the type of data and its presentation as found in the full report. The complete list of data sets and site locations for the chapter can be found in Appendix B of this document. An order form to obtain a free copy of *Trends '91* is found in Appendix C.

Introduction

Atmospheric CO₂

Precise records of past and present atmospheric CO₂ concentrations are critical to studies attempting to model and understand the global carbon cycle and possible CO₂-induced climate change. Attempts to determine past levels of atmospheric CO₂ concentrations have been made by using a variety of techniques, including direct measurements of trapped air in polar ice cores, indirect determinations from carbon isotopes in tree rings, spectroscopic data, and carbon and oxygen isotopic changes in carbonate sediments in deep-ocean cores. The modern period of precise atmospheric CO₂ measurements began during the International Geophysical Year (1958) with Keeling's (Scripps Institution of Oceanography) pioneering determinations at Mauna Loa, Hawaii, and at the South Pole. The Mauna Loa record remains the single most valuable CO₂ time series. Since Keeling's initial efforts, other agencies and organizations have implemented programs to monitor background levels of atmospheric CO₂ concentrations. Two of the larger programs are the World Meteorological Organization's (WMO's) Background Air Pollution Monitoring Network (BAPMoN) and the National Oceanic and Atmospheric Administration's Climate Monitoring and Diagnostics Laboratory (CMDL—formerly Geophysical Monitoring for Climate Change), which operates a flask monitoring network and a continuous monitoring network.

In the following section, two historical atmospheric CO₂ records derived from ice cores and 48 modern records of monthly and annual atmospheric CO₂ concentrations from 36 globally distributed sites are provided (see below). Collectively, these records document the

Location of the sites for which atmospheric CO₂ records are presented in this section.

Atmospheric CO₂

atmospheric CO₂ record during the past 160,000 years (see figure below). For the modern records it seemed appropriate to present Keeling's Mauna Loa record first, because it represents the longest continuous modern record available. Most records are indicative of background air conditions in a particular region or location. Most sites are situated in remote locations and are subject to large, uncontaminated air masses (see figure on facing page). The data presented on the pages that follow represent a significant portion, although not all, of the modern and historical atmospheric CO₂ records and provide irrefutable evidence that atmospheric levels of CO₂ concentrations have risen over the past three decades.

Annual atmospheric CO₂ concentrations during the past 160,000 years (as derived from the Vostok and Siple ice cores and Keeling's Mauna Loa record).

Atmospheric CO₂

The data presented in this section were made available to the Carbon Dioxide Information Analysis Center (CDIAC) by the principal investigator(s) listed at the beginning of each data record, and CDIAC acknowledges their kind cooperation. We urge readers to credit the principal investigators and their organizations when using these data. Users are encouraged to contact CDIAC before using data for specific model exercises or research exercises. Some of the data are preliminary and subject to adjustment (values could change by several parts per million). All the data presented here are available in digitized form from CDIAC.

Mt. Cimone Observatory, Italy.

Mauna Loa

Charles D. Keeling and Timothy P. Whorf

Scripps Institution of Oceanography, University of California, La Jolla, California 92093, U.S.A.

Period of Record

1958–1990

Methods

Air samples are collected continuously from air intakes at the top of four 7-m towers and one 27-m tower. Four air samples are collected each hour for the purpose of determining the CO₂ concentration. Determinations of CO₂ are made by using an Applied Physics Corporation nondispersive infrared gas analyzer with a water vapor freeze trap. This analyzer registers the concentration of CO₂ in a stream of air flowing at approximately 0.5 L/min.⁻¹ Every 20 minutes the flow is replaced by a stream of calibrating gas or "working reference gas." In December 1983, CO₂-in-N₂ calibration gases were replaced with the presently used CO₂-in-air calibration gases. These calibration gases and other reference gases are compared periodically to determine the instrument sensitivity and to check for possible contamination in the air-handling system. These reference gases are themselves calibrated against specific standard gases whose CO₂ concentrations are determined manometrically. Greater details about the sampling methods are given in Keeling et al. (1982).

Hourly averages of atmospheric CO₂, wind speed, and direction are plotted as a basis for selecting data for further processing. Data are selected for periods of steady hourly data to within approximately 0.5 parts per million by volume (ppmv), with at least six consecutive hours of steady data required to form a daily average. Greater details about the data selection criteria are given in Bacastow et al. (1985).

The Mauna Loa data presented on the following pages are reported in the 1987 Scripps mole fraction scale.

**Mauna Loa
Observatory**
Hawaii, U.S.A.

*Barren lava field of an
active volcano*

19°32' N, 155°35' W
3397 m above MSL

Mauna
Loa

Mauna Loa

Trends

The Mauna Loa atmospheric CO₂ measurements constitute the longest continuous record of atmospheric CO₂ concentrations available in the world. The Mauna Loa site is considered one of the most favorable locations for measuring undisturbed air because possible local influences of vegetation or human activities on atmospheric CO₂ concentrations are minimal and any influences from volcanic vents may be excluded from the records. The methods and equipment used to obtain these measurements have been essentially unchanged over the 32-year record.

Because of the favorable site location, continuous monitoring, and careful selection and scrutiny of the data, the Mauna Loa record is considered to be a precise record and a reliable indicator of the regional trend in the concentrations of atmospheric CO₂ in the middle layers of the troposphere. The Mauna Loa record shows a 12% increase in the mean annual concentration in 32 years, from 315.83 parts per million by volume of dry air (ppmv) in 1959 to 353.95 ppmv in 1990.

Mauna Loa

References

- Bacastow, R.B., and C.D. Keeling. 1981. Atmospheric carbon dioxide concentration and the observed airborne fraction. pp. 103–12. In B. Bolin (ed.), *Carbon Cycle Modelling*, SCOPE 16. John Wiley and Sons, New York.
- Bacastow, R.B., C.D. Keeling, and T.P. Whorf. 1985. Seasonal amplitude increase in atmospheric CO₂ concentration at Mauna Loa, Hawaii, 1959–1982. *Journal of Geophysical Research* 90(D6):10529–40.
- Keeling, C.D. 1986. Atmospheric CO₂ concentrations—Mauna Loa Observatory, Hawaii 1958–1986. NDP-001/R1. Carbon Dioxide Information Analysis Center. Oak Ridge National Laboratory. Oak Ridge, Tennessee.
- Keeling, C.D., R.B. Bacastow, A.E. Bainbridge, C.A. Ekdahl, Jr., P.R. Guenther, L.S. Waterman, and J.F.S. Chin. 1976. Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii. *Tellus* 28(6):538–51.
- Keeling, C.D., R.B. Bacastow, and T.P. Whorf. 1982. Measurements of the concentration of carbon dioxide at Mauna Loa Observatory, Hawaii. pp. 377–85. In W.C. Clark (ed.), *Carbon Dioxide Review: 1982*. Oxford University Press, New York.
- Keeling, C.D., R.B. Bacastow, A.F. Carter, S.C. Piper, T.P. Whorf, M. Heimann, W.G. Mook, and H. Roeloffzen. 1989. A three-dimensional model of atmospheric CO₂ transport based on observed winds: 1. Analysis of observational data. In D.H. Peterson (ed.), *Aspects of Climate Variability in the Pacific and the Western Americas*. Geophysical Monograph 55:165–235.
- Pales, J.C., and C.D. Keeling. 1965. The concentration of atmospheric carbon dioxide in Hawaii. *Journal of Geophysical Research* 70:6053–76.

Mauna Loa

Atmospheric Concentrations of Carbon Dioxide^a

Year	CO ₂ Concentrations (ppmv)												
	Jan	Feb	Mar	April	May	June	July	Aug	Sept	Oct	Nov	Dec	Ann
1958			315.55	317.29	317.34		315.69	314.78	313.03		313.18	314.50	
1959	315.42	316.32	316.49	317.56	318.13	318.00	316.39	314.66	313.68	313.18	314.66	315.43	315.83
1960	316.27	316.81	317.42	318.87	319.87	319.43	318.01	315.75	314.00	313.68	314.84	316.03	316.75
1961	316.73	317.54	318.38	319.31	320.42	319.61	318.42	316.64	314.83	315.15	315.95	316.85	317.49
1962	317.78	318.40	319.53	320.41	320.85	320.45	319.44	317.25	316.12	315.27	316.53	317.53	318.30
1963	318.58	318.92	319.70	321.22	322.08	321.31	319.58	317.61	316.05	315.83	316.91	318.20	318.83
1964	319.41				322.06	321.73	320.27	318.54	316.54	316.71	317.53	318.55	
1965	319.27	320.28	320.73	321.97	322.00	321.71	321.05	318.71	317.65	317.14	318.71	319.25	319.87
1966	320.46	321.43	322.22	323.54	323.91	323.59	322.26	320.21	318.48	317.94	319.63	320.87	321.21
1967	322.17	322.34	322.88	324.25	324.83	323.93	322.39	320.76	319.10	319.23	320.56	321.80	322.02
1968	322.40	322.99	323.73	324.86	325.41	325.19	323.97	321.92	320.10	319.96	320.97	322.48	322.83
1969	323.52	323.89	325.04	326.01	326.67	325.96	325.13	322.90	321.61	321.01	322.08	323.37	323.93
1970	324.34	325.30	326.29	327.54	327.54	327.21	325.98	324.42	322.91	322.90	323.85	324.96	325.27
1971	326.01	326.51	327.01	327.62	328.76	328.40	327.20	325.28	323.20	323.40	324.64	325.85	326.16
1972	326.60	327.47	327.58	329.56	329.90	328.92	327.89	326.17	324.68	325.04	326.34	327.39	327.29
1973	328.37	329.40	330.14	331.33	332.31	331.90	330.70	329.15	327.34	327.02	327.99	328.48	329.51
1974	329.18	330.55	331.32	332.48	332.92	332.08	331.02	329.24	327.28	327.21	328.29	329.41	330.08
1975	330.23	331.24	331.87	333.14	333.80	333.42	331.73	329.90	328.40	328.17	329.32	330.59	330.99
1976	331.58	332.39	333.33	334.41	334.71	334.17	332.88	330.77	329.14	328.77	330.14	331.52	331.98
1977	332.75	333.25	334.53	335.90	336.57	336.10	334.76	332.59	331.41	330.98	332.24	333.68	333.73
1978	334.80	335.22	336.47	337.59	337.84	337.72	336.37	334.51	332.60	332.37	333.75	334.79	335.34
1979	336.05	336.59	337.79	338.71	339.30	339.12	337.56	335.92	333.74	333.70	335.13	336.56	336.68
1980	337.84	338.19	339.90	340.60	341.29	341.00	339.39	337.43	335.72	335.84	336.93	338.04	338.52
1981	339.06	340.30	341.21	342.33	342.74	342.07	340.32	338.27	336.52	336.68	338.19	339.44	339.76
1982	340.57	341.44	342.53	343.39	343.96	343.18	341.88	339.65	337.80	337.69	339.09	340.32	340.96
1983	341.20	342.35	342.93	344.77	345.58	345.14	343.81	342.22	339.69	339.82	340.98	342.82	342.61
1984	343.52	344.33	345.11	346.88	347.25	346.61	345.22	343.11	340.90	341.17	342.80	344.04	344.25
1985	344.79	345.82	347.25	348.17	348.75	348.07	346.38	344.52	342.92	342.63	344.06	345.38	345.73
1986	346.12	346.79	347.69	349.38	350.04	349.38	347.78	345.75	344.70	344.01	345.50	346.75	346.99
1987	347.86	348.32	349.26	350.84	351.70	351.11	349.37	347.97	346.31	346.22	347.68	348.82	348.79
1988	350.29	351.51	352.08	353.45	354.08	353.66	352.25	350.30	348.58	348.74	349.93	351.21	351.35
1989	352.62	352.93	353.54	355.27	355.52	354.97	353.74	351.51	349.63	349.82	351.12	352.35	352.75
1990	353.47	354.51	355.18	355.98	356.94	355.99	354.58	352.68	350.72	350.92	352.55	353.91	353.95

^aAverages of data adjusted to the 15th of each month expressed in parts per million by volume. Annual averages are based on monthly means. Numeric data supporting *Trends '91* are available in electronic form from CDIAC. These monthly and annual values, as well as daily concentrations, may be obtained from CDIAC (Keeling 1986) or from the principal investigators.

Note: The SIO Mauna Loa record presented in *Trends '90* did not contain corrections made to the Mauna Loa record at the time of the Mauna Loa Report. The corrections included updates of the calibration factor and of the working tank indices back to the beginning of the record as well as adjustments made for a drifting detector in the 1960s.

Atmospheric CH₄

This section of the highlights contains the introduction to the *Trends '91* chapter on atmospheric (CH₄) concentrations. It also provides a sample record of data used in constructing a historical record of atmospheric CH₄ concentrations from ice cores. The data for this sample record was provided by D. Raynaud, J. Chappellaz, J.M. Barnola, and C. Lorius from the Glaciology and Environmental Geophysics Laboratory (Saint Martin d'Hères, France) and Y.S. Korotkevich from the Arctic and Antarctic Research Institute (St. Petersburg, Russia). The sample illustrates the type of data and its presentation as found in the full report. The complete list of data sets and site locations for the chapter can be found in Appendix B of this document. An order form to obtain a free copy of *Trends '91* is found in Appendix C.

Introduction

Atmospheric CH₄

Methane (CH₄) is recognized as one of the most important radiatively active trace gases that might significantly affect climatic change within the next century. There is compelling evidence that the concentration of CH₄ in the atmosphere is rising rapidly and that present levels are approximately double those of a few hundred years ago (see figure on page 206). The most important methane sources are enteric fermentation in ruminants; anaerobic decay of organic matter in rice paddies, natural wetlands, and landfills; coal mining; natural gas production and distribution; oil exploration and production; and biomass burning. Precise records of past and present CH₄ concentrations are critical to studies attempting to determine the potential effects that greenhouse gases would have on climate. Past concentrations of atmospheric CH₄ have been determined by analyzing air bubbles trapped in ice from Greenland and Antarctica (see figure on page 207). Since 1978, when the modern CH₄ record begins, systematic investigations have found on average a 1–2% increase in CH₄ concentrations per year. Most of these measurements have been made in the troposphere by using the method of gas chromatography with a flame ionization detector. Other measurements have relied upon an optical absorption technique that uses the sun as a source of infrared radiation and yields an average atmospheric abundance along the entire path length traversed through the atmosphere.

The following pages provide historical and modern CH₄ records, including CH₄ concentrations derived from ice cores drilled at 6 sites, global monthly averages derived from 7 monitoring sites, and monthly atmospheric CH₄ concentration records from 25 globally distributed sites (see figure on bottom of page 207). Data from these 25 sites were collected as part of the monitoring program of the National Oceanic and Atmospheric Administration's Climate Monitoring and Diagnostics Laboratory (NOAA/CMDL).

We urge readers to credit the principal investigators and their organizations listed at the beginning of the data records when using these data. In addition, users are encouraged to contact CDIAC before applying the data in specific model exercises or research exercises. Some of the data are preliminary and are subject to change by several parts per billion. All of the data presented here are available in digitized form from CDIAC.

Other Trace Gases—Chlorofluorocarbon Release Estimates

CFC-11

Atmospheric CFC-11 Releases^a

Releases (million kilograms)					Releases (million kilograms)				
Year	Total	Refrigeration	Blowing agents (closed-cell foam)	Open-cell foam, aerosols, and all other	Year	Total	Refrigeration	Blowing agents (closed-cell foam)	Open-cell foam, aerosols, and all other
1931	0.0	0.0	0.0	0.0	1961	52.1	3.2	0.6	48.2
1932	0.0	0.0	0.0	0.0	1962	65.4	3.9	1.3	60.3
1933	0.0	0.0	0.0	0.0	1963	80.0	4.5	1.8	73.6
1934	0.0	0.0	0.0	0.0	1964	95.0	5.2	2.7	87.2
1935	0.0	0.0	0.0	0.0	1965	108.3	5.8	3.5	98.9
1936	0.0	0.0	0.0	0.0	1966	121.3	6.5	4.6	110.2
1937	0.0	0.0	0.0	0.0	1967	137.6	7.1	5.8	124.7
1938	0.1	0.1	0.0	0.0	1968	156.8	7.9	7.6	141.3
1939	0.1	0.1	0.0	0.0	1969	181.9	8.7	9.5	163.6
1940	0.1	0.1	0.0	0.0	1970	206.6	9.7	11.4	185.6
1941	0.1	0.1	0.0	0.0	1971	226.9	10.5	14.2	202.2
1942	0.1	0.1	0.0	0.0	1972	255.8	11.8	17.4	226.6
1943	0.2	0.2	0.0	0.0	1973	292.4	13.0	21.2	258.1
1944	0.2	0.2	0.0	0.0	1974	321.4	14.5	24.0	283.0
1945	0.3	0.3	0.0	0.0	1975	310.9	15.8	25.8	269.3
1946	0.6	0.3	0.0	0.4	1976	316.7	17.6	27.8	271.4
1947	1.3	0.3	0.0	1.0	1977	303.9	19.0	32.0	252.8
1948	2.3	0.3	0.0	2.0	1978	283.6	20.7	35.0	228.0
1949	3.8	0.2	0.0	3.6	1979	263.7	22.4	40.1	201.1
1950	5.5	0.2	0.0	5.2	1980	250.8	24.3	44.0	182.6
1951	7.6	0.3	0.0	7.4	1981	248.2	25.9	49.9	172.4
1952	11.0	0.3	0.0	10.6	1982	239.5	26.8	53.3	159.4
1953	15.0	0.4	0.1	14.5	1983	252.8	27.4	57.9	167.4
1954	18.6	0.6	0.1	17.9	1984	271.1	27.2	63.5	180.5
1955	23.0	0.8	0.1	22.2	1985	280.8	26.7	69.0	185.2
1956	28.7	1.0	0.1	27.5	1986	295.1	26.0	75.1	193.9
1957	32.2	1.3	0.2	30.7	1987	310.6	25.9	84.7	200.1
1958	30.2	1.7	0.2	28.3	1988	314.5	26.2	91.4	196.9
1959	30.9	2.1	0.3	28.5	1989	265.2	26.5	97.5	141.1
1960	40.5	2.7	0.5	37.4					

^aRelease figures for refrigeration represent the sum of releases for hermetically sealed and nonhermetically sealed refrigeration. These data are available upon request from the Chemical Manufacturers Association. The data as presented in Trends '91 are also available in electronic form from CRIAC.

CO₂ Emissions

This section of the highlights contains the introduction to the *Trends '91* chapter on CO₂ emissions and a sample record of global CO₂ emissions data provided by Gregg Marland and Tom Boden from Oak Ridge National Laboratory (Oak Ridge, Tennessee, U.S.A.). This sample illustrates the type of data and its presentation as found in *Trends '91*. The complete list of data sets and site regions for the chapter can be found in Appendix B. An order form to obtain a free copy of *Trends '91* is found in Appendix C.

Introduction

CO₂ Emissions

Atmospheric CO₂ is produced both from natural sources and from human activities. Of human activities, the most important source of CO₂ is the release of carbon during the burning of fossil fuels. Since the Industrial Revolution, global CO₂ emissions from fossil fuel consumption have risen dramatically (see figure below). In 1989, 5.97 gigatons (1 gigaton = 1000 million metric tons) of carbon was emitted to the atmosphere as a result of fossil fuel burning, cement manufacturing, and gas flaring. These emissions, although small in comparison with the amounts of carbon stored in the oceans and in the terrestrial biosphere, represent a significant component of global carbon cycling and the greenhouse effect and, more important, represent the largest human contribution to the CO₂ present in the atmosphere.

Keeling (1973) was the first to establish a systematic method for estimating the amount of CO₂ emitted from fossil fuel consumption. He used energy data from the U.N. Department of International Economic and Social Affairs. Since 1973, both the energy data collection and the procedures for estimating CO₂ emissions have been refined and improved (Marland and Rotty 1984; Marland et al. 1989; United Nations 1990).

The following section presents Keeling's 1860–1953 global emission estimates along with more-recent global, regional, and national annual emission estimates calculated by Marland and Boden. These more recent estimates cover the period 1950–1989 and were calculated by using the methods of Marland and Rotty (1984). The data used for the calculations were energy and population statistics from the U.N., cement manufacturing data from the U.S. Department of Interior's Bureau of Mines, and gas flaring data from the U.N. and from the U.S. Department of Energy's Energy Information Administration.

CO₂ Emissions

Emission estimates for 1950–1989 are provided for nine regions. The regional definitions are consistent with those previously used by Rotty et al. (1984), Rotty and Marland (1986), and by the United Nations. The data are presented in descending order with the highest emitting region in 1989, North America, which is presented first. These estimates illustrate the changing patterns in regional CO₂ emissions over the last four decades, particularly in developing regions.

For emphasis and brevity, national estimates are provided only for the 20 highest emitting countries in 1989 (see figure below). The data are presented in descending order, with the highest emitting country, the United States, presented first. These 20 countries contributed slightly over 80% of all the 1989 world emissions from fossil fuel consumption. The top three countries—the United States, the U.S.S.R., and the People's Republic of China—were responsible for over half (50.6%) of the world emissions from fossil fuel burning in 1989. Spain, the 20th highest emitting nation, contributed slightly less than 1% to this total.

Ranking of the twenty highest emitting countries in 1989 and their rank in 1950, contributions to global emissions, and per capita emission rates.

We urge readers to credit the principal investigators and their organizations (listed at the beginning of each data record) when using these data. Users are cautioned to pay close attention to table footnotes concerning the nature of each data record and are encouraged to contact CDIAC before applying the data in specific model exercises or research exercises.

Global

Gregg Marland and Tom Boden

Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee
37831-6335, U.S.A.

Sponsored by the U.S. Department of Energy, Carbon Dioxide Research Program

Period of Record

1950–1989

Methods

These CO₂ emission estimates, derived primarily from energy statistics published by the United Nations (1991), were calculated by using the methods of Marland and Rotty (1984). The energy statistics were compiled primarily from annual questionnaires distributed by the U.N. Statistical Office and supplemented by official national statistical publications. As stated in the introduction of the *1989 Energy Statistics Yearbook*, “where official data are not available or are inconsistent, estimates are made by the Statistical Office based on governmental, professional or commercial materials. Estimates include, but are not limited to, extrapolated data based on partial year information, use of annual trends, trade data based on partner country reports, breakdowns of aggregated data as well as analysis of current energy events and activities.” Data from the U.S. Department of Interior’s Bureau of Mines (Johnson 1990) were used to estimate CO₂ emitted during cement production. Emissions from gas flaring were derived primarily from U.N. data but were supplemented with data from the U.S. Department of Energy’s Energy Information Administration (1991), Rotty (1974), and with a few national estimates provided by G. Marland.

Global

Trends

Global carbon emissions from fossil fuel consumption, cement manufacturing, and gas flaring have risen significantly since 1950. The 1989 estimate for global emissions of carbon, ~5.97 gigatons, marks the sixth consecutive year that global emissions have increased. The 1989 estimate is the highest annual emission estimate since the data record began in 1950. The previous highest estimate was for 1988, in which 5.90 gigatons of carbon was emitted to the atmosphere from fossil fuel burning, cement manufacturing, and gas flaring. The 1989 estimate represents a 1.2% growth over 1988, a notable drop from growth rates of the last several years.

Globally, liquid and solid fuels accounted for approximately 81% of the emissions from fossil fuel burning in 1989. Combustion of gas fuels (e.g., natural gas) accounted for 15.9% (946 million metric tons) of the total emissions from fossil fuels in 1989. This estimate represents an increase of 8.0% over emissions from gas fuels in 1987. Emissions from cement manufacturing increased slightly, while global emissions from gas flaring seem to be rising during the last two years. This last trend is the result of very uncertain data on the U.S.S.R. and should be treated lightly. Collectively, emissions from cement production and gas flaring contributed less than 4% to the total emissions for 1989.

Global CO₂
emissions.

Global

References

- Johnson, W. 1990. *Minerals Yearbook*, Cement 1989. U.S. Department of the Interior Bureau of Mines, Washington, D.C.
- Keeling, C.D. 1973. Industrial production of carbon dioxide from fossil fuels and limestone. *Tellus* 25:174–98.
- Marland, G., and R.M. Rotty. 1984. Carbon dioxide emissions from fossil fuels: A procedure for estimation and results for 1950–1982. *Tellus* 36(B):232–61.
- Marland, G., T.A. Boden, R.C. Griffin, S.F. Huang, P. Kanciruk, and T.R. Nelson. 1989. Estimates of CO₂ emissions from fossil fuel burning and cement manufacturing, based on the United Nations energy statistics and the U.S. Bureau of Mines cement manufacturing data. ORNL/CDIAC-25, NDP-030. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Rotty, R.M. 1974. First estimates of global flaring of natural gas. *Atmospheric Environment* 8:681–86.
- Rotty, R.M., and G. Marland. 1986. Fossil fuel combustion: Recent amounts, patterns, and trends of CO₂. pp. 484–500. In J.R. Trabalka and D.E. Reichle (eds.), *The Changing Carbon Cycle: A Global Analysis*, Springer-Verlag, New York.
- United Nations. 1991. *1989 Energy Statistics Yearbook*. United Nations Department of International Economic and Social Affairs, Statistical Office, New York.
- U.S. Department of Energy. 1991. *International Energy Annual 1989*. DOE/EIA-0219(89). Energy Information Administration, Office of Energy Markets and End Use, Washington, D.C.

Global Carbon Dioxide Emission Estimates^a

Year	Emissions (million metric tons of carbon)						Per capita emissions (metric tons of carbon)
	Total	Solid	Liquid	Gas	Cement	Gas flaring	
1950	1638	1077	423	97	18	23	0.65
1951	1775	1137	479	115	20	24	0.69
1952	1803	1127	504	124	22	26	0.69
1953	1848	1132	533	131	24	27	0.70
1954	1871	1123	557	138	27	27	0.69
1955	2050	1215	625	150	30	31	0.74
1956	2185	1281	679	161	32	32	0.78
1957	2278	1317	714	178	34	35	0.80
1958	2338	1344	732	192	36	35	0.80
1959	2471	1390	790	214	40	36	0.83
1960	2586	1419	850	235	43	39	0.86
1961	2602	1356	905	254	45	42	0.85
1962	2708	1358	981	277	49	44	0.86
1963	2855	1404	1053	300	51	47	0.89
1964	3016	1442	1138	328	57	51	0.92
1965	3154	1468	1221	351	59	55	0.95
1966	3314	1485	1325	380	63	60	0.97
1967	3420	1455	1424	410	65	66	0.98
1968	3596	1456	1552	445	70	73	1.01
1969	3809	1494	1674	487	74	80	1.05
1970	4091	1571	1838	516	78	87	1.11
1971	4242	1571	1946	554	84	88	1.12
1972	4409	1587	2056	583	89	94	1.15
1973	4648	1594	2240	608	95	110	1.18
1974	4656	1591	2244	618	96	107	1.16
1975	4629	1686	2131	623	95	93	1.13
1976	4895	1723	2313	647	103	109	1.18
1977	5034	1786	2390	646	108	104	1.19
1978	5082	1802	2383	674	116	107	1.18
1979	5366	1899	2535	714	119	100	1.23
1980	5264	1921	2409	725	120	89	1.18
1981	5129	1930	2272	735	121	72	1.13
1982	5094	1993	2178	734	121	69	1.11
1983	5085	1998	2163	736	125	63	1.09
1984	5243	2088	2186	785	128	57	1.10
1985	5369	2196	2169	819	131	55	1.11
1986	5551	2253	2274	836	136	52	1.12
1987	5661	2310	2285	876	142	48	1.13
1988	5897	2388	2388	916	150	55	1.15
1989	5967	2392	2419	946	152	56	1.15

^aEmission estimates rounded and expressed in million metric tons of carbon; per capita estimates rounded and expressed in metric tons of carbon. Data appearing in *Trends '91* are available in electronic form from CDIAC.

Temperature

This section of the highlights contains the introduction to the *Trends '91* chapter on temperature and a sample record of global and hemispheric temperature anomalies provided by Philip D. Jones and Tom M.L. Wigley from the Climatic Research Unit of the School of Environmental Sciences at the University of East Anglia (Norwich, U.K.). The sample illustrates the type of data and its presentation as found in *Trends '91*. The complete list of data sets and regions for this chapter can be found in Appendix B. An order form to obtain a free copy *Trends '91* is found in Appendix C.

Introduction

Temperature

During the past 20 years, much attention has been given to the potential climatic effects of increasing concentrations of atmospheric greenhouse gases. This is not surprising because these potential climatic effects could have far-reaching environmental, economic, and social consequences. Much of the work that has been performed during this period has involved (1) climate modeling or empirical analyses attempting to attribute climatic changes to various astronomical and atmospheric factors and (2) assembly and examination of new geological, historical, and instrumental data. From these studies, researchers have gained important new insights into the possible climatic response to the increasing concentrations of CO₂ and other trace gases, together with a better understanding of the sensitivity of the overall climate system to both human and natural perturbations.

In order to isolate the “greenhouse signal,” or that part of the climate change that is indicative of increased concentrations of greenhouse gases, it is imperative that we have long-term observational records to identify the climate changes that have already occurred. It is also imperative that these observational records have sufficient spatial coverage. Although local and regional variations in temperature are important in assessing the impacts of climate and climate change on society, large-scale averages are most likely to indicate detectable changes in climate that result from changes in atmospheric trace gas concentrations. Fortunately, for the past century many climatic variables (e.g., surface air temperature) have been measured at a large number of meteorological stations, mostly at land locations in the Northern Hemisphere. Researchers have compiled and analyzed several time series of global average surface air temperatures largely on the basis of land-based data and data from a few fixed-position weather ships. Previous attempts to determine large-scale tropospheric and stratospheric temperatures have typically used radiosonde and rocketsonde data. Recently, global temperature series have been compiled on the basis of satellite data. Despite the possible shortcomings of each of these individual temperature records, collectively they provide a good indication of global warming trends. Indeed, many of the series show warming trends in the troposphere, particularly during the 1980s, yet it remains unclear whether these trends result from a buildup of greenhouse gases or are due to other factors [e.g., greater frequency of El Niño/Southern Oscillation (ENSO) events in the Pacific Ocean] (see figure on following page).

The following pages provide six records of surface air temperatures and surface air temperature anomalies. Their spatial coverage ranges from the globe to U.S. regions and from the Earth’s surface to the low stratosphere. The first three records are global and hemispheric surface air temperature anomalies. They are followed by a record containing global and hemispheric temperature anomalies derived from satellite data and a record containing annual and seasonal temperature anomalies for the Earth’s surface, the troposphere, the tropopause, and the low stratosphere. For these vertical profiles, global anomaly estimates are presented first, followed by northern zonal averages, southern zonal averages, and hemispheric averages. Finally, national estimates of seasonal and annual temperatures for the United States are presented, followed by estimates for the subregions that comprise the aggregated West, Central, and Eastern regions within the United States. All of these temperature records have been frequently referenced in studies addressing potential greenhouse gas-induced climate changes.

Temperature

Global annual temperature anomalies from Spencer and Christy (A), Angell (B), Vinnikov et al. (C), Hansen et al. (D), and Jones et al. (E).

Temperature

All of the data presented in this section were made available to CDIAC by the principal investigator(s) listed at the beginning of the data records. We urge readers to credit the principal investigators and their organizations when using these data. Users are cautioned to pay close attention to table footnotes concerning the nature of each data record and are encouraged to contact CDIAC before applying the data in specific model exercises or research exercises.

Global and Hemispheric Anomalies

Philip D. Jones and Tom M. L. Wigley

Climatic Research Unit, School of Environmental Sciences, University of East Anglia,
Norwich, United Kingdom NR4 7TJ

Sponsored by the U.S. Department of Energy, Carbon Dioxide Research Program

Period of Record

1854–1990 (relative to a 1950–1979 reference period)

Methods

These global and hemispheric temperature anomaly estimates were compiled by Jones et al. (1986a,b,c), Jones (1988), Jones and Wigley (1990), and Jones et al. (1991) and are based on corrected land and marine data. Land data were derived from meteorological data and fixed-position weather ship data that were corrected for nonclimatic errors, such as station shifts and/or instrument changes. The marine data were from the Comprehensive Ocean-Atmosphere Data Set (COADS) compilation, which extends (with updates) to 1986 (Woodruff et al. 1987). Updates to 1988 were based on hemispheric sea surface temperature (SST) estimates produced by the United Kingdom Meteorological Office. Both SSTs and marine air temperatures were corrected, but only SST data were used in the combined land-marine data set. Two series of global and hemispheric temperature anomaly estimates are provided, one adjusted for the influence of the El Niño/Southern Oscillation (ENSO) events and the other uncorrected for ENSO events.

The ENSO influence can be extracted by using regression techniques (Jones 1988, Angell 1990). According to Jones (1988), the Southern Oscillation Index (SOI) explains 20–30% of the high-frequency temperature variance.

Global and Hemispheric Anomalies

Trends

These variations in global and hemispheric annual temperatures compiled by Jones and Wigley show little trend during the nineteenth century, marked warming to 1940, relatively steady conditions to the mid-1970s, followed by a rapid warming during the 1980s. The 1980s was a decade of unprecedented warmth, with 1990 the warmest year since comparable records began in the middle of the 19th century. Both the long-term and the recent warming trends have been common to the two hemispheres. The annual value for 1990 is clearly the warmest for the Northern Hemisphere and global records. In the global record, the provisional value 0.39°C for 1990 is 0.05°C warmer than that for the previous warmest year (1988). All six of the warmest years in the global record have now occurred since 1980. In descending order they are 1990, 1988, 1987, 1983, 1989, and 1981.

After correction for ENSO events, the four most recent years become the four warmest in the global record, and every year since 1985 shows an increase in temperature.

Temperature

Global and Hemispheric Anomalies

Northern Hemisphere annual temperature anomalies, 1854–1990.

Southern Hemisphere annual temperature anomalies, 1854–1990.

Temperature

Global and Hemispheric Anomalies

References

- Angell, J.K. 1990. Variation in global tropospheric temperature after adjustment for El Niño influence. *Geophysical Research Letters* 17:1093–96.
- Jones, P.D. 1988. The influence of ENSO on global temperatures. *Climate Monitor* 17(3):80–89.
- Jones, P.D. and T.M.L. Wigley. 1990. Global warming trends. *Scientific American* 263:84–91.
- Jones, P.D., S.C.B. Raper, R.S. Bradley, H.F. Diaz, P.M. Kelly, and T.M.L. Wigley. 1986a. Northern Hemisphere surface air temperature variations: 1851–1984. *Journal of Climate and Applied Meteorology* 25(2):161–79.
- Jones, P.D., S.C.B. Raper, and T.M.L. Wigley. 1986b. Southern Hemisphere surface air temperature variations: 1851–1984. *Journal of Climate and Applied Meteorology* 25(9):1213–30.
- Jones, P.D., T.M.L. Wigley, and P.B. Wright. 1986c. Global temperature variations between 1861 and 1984. *Nature* 322:430–34.
- Jones, P.D., P.Ya. Groisman, M. Coughlan, N. Plummer, W.-C. Wang, and T.R. Karl. 1990a. Assessment of urbanization effects in time series of surface air temperatures over land. *Nature* 347:169–72.
- Jones, P.D., T.M.L. Wigley, and P.B. Wright. 1990b. Global and hemispheric annual temperature variations between 1861 and 1988. NDP-022/R1, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Jones, P.D., T.M.L. Wigley, and G. Farmer. 1991. Marine and land temperature data sets: A comparison and a look at recent trends. pp. 153–72 In M.E. Schlesinger (ed.), *Greenhouse-Gas-Induced Climatic Change: A Critical Appraisal of Simulations and Observations*. Elsevier Science Publishers, Amsterdam, Netherlands.
- Woodruff, S.D., R.J. Slutz, R.J. Jenne, and P.M. Steurer. 1987. A comprehensive ocean-atmosphere data set. *Bulletin of the American Meteorological Society* 80:1239–47.

Global and Hemispheric Anomalies

Annual Temperature Anomalies^a

Year	North- ern Hem.	South- ern Hem.	Global	ENSO-subtracted ^b			Year	North- ern Hem.	South- ern Hem.	Global	ENSO-subtracted ^b		
				North- ern Hem.	South- ern Hem.	Global					North- ern Hem.	South- ern Hem.	Global
1854	-0.15	-0.08	-0.12				1892	-0.42	-0.32	-0.37	-0.41	-0.31	-0.36
1855	-0.51	-0.49	-0.50				1893	-0.48	-0.37	-0.43	-0.46	-0.35	-0.41
1856	-0.33	-0.21	-0.27				1894	-0.35	-0.32	-0.34	-0.34	-0.31	-0.33
1857	-0.32	-0.60	-0.46				1895	-0.40	-0.20	-0.30	-0.41	-0.21	-0.31
1858	-0.37	-0.46	-0.42				1896	-0.17	-0.09	-0.13	-0.21	-0.13	-0.17
1859	-0.02	-0.26	-0.14				1897	-0.14	-0.10	-0.12	-0.22	-0.18	-0.20
1860	-0.43	-0.31	-0.37				1898	-0.23	-0.40	-0.31	-0.21	-0.38	-0.29
1861	-0.25	-0.58	-0.41				1899	-0.15	-0.22	-0.19	-0.14	-0.21	-0.18
1862	-0.67	-0.41	-0.54				1900	-0.03	-0.09	-0.06	-0.05	-0.11	-0.08
1863	-0.09	-0.30	-0.20				1901	-0.09	-0.24	-0.17	-0.08	-0.23	-0.16
1864	-0.47	-0.50	-0.49				1902	-0.31	-0.25	-0.28	-0.30	-0.24	-0.27
1865	-0.20	-0.25	-0.23				1903	-0.35	-0.47	-0.41	-0.36	-0.47	-0.42
1866	-0.17	-0.21	-0.19				1904	-0.46	-0.52	-0.49	-0.40	-0.46	-0.43
1867	-0.30	-0.20	-0.25	-0.31	-0.21	-0.26	1905	-0.29	-0.40	-0.34	-0.36	-0.47	-0.41
1868	-0.11	-0.39	-0.25	-0.14	-0.42	-0.28	1906	-0.18	-0.29	-0.23	-0.20	-0.30	-0.25
1869	-0.10	-0.34	-0.22	-0.13	-0.37	-0.25	1907	-0.47	-0.37	-0.42	-0.43	-0.33	-0.38
1870	-0.27	-0.41	-0.34	-0.26	-0.40	-0.33	1908	-0.44	-0.48	-0.46	-0.43	-0.47	-0.45
1871	-0.44	-0.39	-0.41	-0.44	-0.39	-0.41	1909	-0.36	-0.31	-0.34	-0.34	-0.29	-0.32
1872	-0.17	-0.38	-0.28	-0.16	-0.37	-0.27	1910	-0.35	-0.33	-0.34	-0.30	-0.28	-0.29
1873	-0.24	-0.35	-0.29	-0.19	-0.30	-0.24	1911	-0.38	-0.42	-0.40	-0.34	-0.38	-0.36
1874	-0.30	-0.51	-0.41	-0.31	-0.52	-0.42	1912	-0.42	-0.18	-0.30	-0.50	-0.26	-0.38
1875	-0.61	-0.34	-0.48	-0.59	-0.32	-0.46	1913	-0.43	-0.21	-0.32	-0.46	-0.24	-0.35
1876	-0.39	-0.47	-0.43	-0.38	-0.46	-0.42	1914	-0.22	-0.17	-0.20	-0.26	-0.21	-0.24
1877	-0.10	-0.25	-0.17	-0.14	-0.29	-0.21	1915	-0.04	-0.12	-0.08	-0.11	-0.18	-0.15
1878	0.19	-0.17	0.01	0.10	-0.25	-0.08	1916	-0.32	-0.27	-0.30	-0.26	-0.21	-0.24
1879	-0.32	-0.42	-0.37	-0.24	-0.34	-0.29	1917	-0.59	-0.40	-0.50	-0.46	-0.27	-0.37
1880	-0.36	-0.30	-0.33	-0.30	-0.24	-0.27	1918	-0.40	-0.31	-0.36	-0.29	-0.20	-0.25
1881	-0.33	-0.25	-0.29	-0.31	-0.23	-0.27	1919	-0.33	-0.15	-0.24	-0.39	-0.21	-0.30
1882	-0.30	-0.28	-0.29	-0.32	-0.30	-0.31	1920	-0.29	-0.19	-0.24	-0.33	-0.22	-0.28
1883	-0.44	-0.31	-0.38	-0.46	-0.33	-0.40	1921	-0.16	-0.26	-0.21	-0.12	-0.22	-0.17
1884	-0.48	-0.39	-0.44	-0.50	-0.41	-0.46	1922	-0.30	-0.31	-0.31	-0.28	-0.29	-0.29
1885	-0.41	-0.34	-0.38	-0.44	-0.37	-0.41	1923	-0.25	-0.30	-0.28	-0.22	-0.27	-0.25
1886	-0.33	-0.16	-0.25	-0.35	-0.18	-0.27	1924	-0.28	-0.35	-0.31	-0.32	-0.39	-0.35
1887	-0.31	-0.48	-0.40	-0.25	-0.42	-0.34	1925	-0.12	-0.25	-0.19	-0.07	-0.20	-0.14
1888	-0.29	-0.38	-0.34	-0.32	-0.41	-0.37	1926	0.02	-0.07	-0.03	-0.09	-0.18	-0.14
1889	-0.20	-0.12	-0.16	-0.26	-0.18	-0.22	1927	-0.09	-0.21	-0.15	-0.10	-0.22	-0.16
1890	-0.38	-0.40	-0.39	-0.31	-0.33	-0.32	1928	-0.10	-0.24	-0.17	-0.09	-0.23	-0.16
1891	-0.31	-0.36	-0.34	-0.30	-0.35	-0.33	1929	-0.32	-0.34	-0.33	-0.28	-0.30	-0.29

Temperature

Global and Hemispheric Anomalies

Annual Temperature Anomalies (continued)^a

Year	North- ern Hem.	South- ern Hem.	Global	ENSO-subtracted ^b			Year	North- ern Hem.	South- ern Hem.	Global	ENSO-subtracted ^b		
				North- ern Hem.	South- ern Hem.	Global					North- ern Hem.	South- ern Hem.	Global
1930	-0.02	-0.28	-0.15	0.00	-0.26	-0.13	1961	0.12	0.11	0.12	0.13	0.12	0.13
1931	0.06	-0.14	-0.04	0.08	-0.12	-0.02	1962	0.17	0.10	0.14	0.20	0.13	0.17
1932	-0.04	-0.17	-0.11	-0.04	-0.17	-0.11	1963	0.19	0.08	0.14	0.21	0.10	0.16
1933	-0.22	-0.18	-0.20	-0.24	-0.20	-0.22	1964	-0.14	-0.14	-0.14	-0.15	-0.15	-0.15
1934	-0.10	-0.14	-0.12	-0.08	-0.12	-0.10	1965	-0.14	-0.08	-0.11	-0.14	-0.07	-0.11
1935	-0.14	-0.20	-0.17	-0.14	-0.20	-0.17	1966	0.03	-0.09	-0.03	-0.04	-0.16	-0.10
1936	-0.07	-0.19	-0.13	-0.05	-0.17	-0.11	1967	0.02	-0.10	-0.04	0.03	-0.09	-0.03
1937	0.08	-0.08	0.00	0.08	-0.08	0.00	1968	-0.02	-0.12	-0.07	0.00	-0.10	-0.05
1938	0.13	-0.02	0.05	0.15	0.00	0.07	1969	0.02	0.15	0.09	0.00	0.13	0.07
1939	0.01	-0.10	-0.05	0.07	-0.04	0.01	1970	0.00	0.08	0.04	-0.03	0.05	0.01
1940	-0.09	0.05	-0.02	-0.15	0.00	-0.08	1971	-0.14	-0.08	-0.11	-0.07	-0.01	-0.04
1941	0.01	0.09	0.05	-0.09	-0.01	-0.05	1972	-0.16	0.19	0.02	-0.14	0.21	0.04
1942	0.05	0.02	0.04	-0.01	-0.04	-0.02	1973	0.09	0.22	0.16	0.05	0.18	0.12
1943	0.07	-0.04	0.02	0.10	-0.01	0.05	1974	-0.20	-0.07	-0.14	-0.11	0.02	-0.04
1944	0.19	0.16	0.17	0.19	0.16	0.17	1975	-0.11	-0.10	-0.11	-0.06	-0.05	-0.06
1945	0.02	0.11	0.06	0.02	0.11	0.06	1976	-0.29	-0.14	-0.22	-0.20	-0.05	-0.13
1946	0.07	-0.17	-0.05	0.07	-0.17	-0.05	1977	0.08	0.11	0.09	0.03	0.06	0.04
1947	0.02	-0.11	-0.05	-0.02	-0.15	-0.09	1978	0.00	0.06	0.03	-0.06	0.00	-0.03
1948	0.02	-0.14	-0.06	0.03	-0.13	-0.05	1979	0.09	0.16	0.13	0.09	0.16	0.13
1949	0.03	-0.14	-0.05	0.01	-0.16	-0.07	1980	0.12	0.20	0.16	0.10	0.18	0.14
1950	-0.12	-0.14	-0.13	-0.07	-0.09	-0.08	1981	0.28	0.14	0.21	0.27	0.13	0.20
1951	0.06	-0.12	-0.03	0.12	-0.06	0.03	1982	0.06	0.14	0.10	0.06	0.14	0.10
1952	0.12	0.05	0.09	0.09	0.02	0.06	1983	0.23	0.35	0.29	0.09	0.21	0.15
1953	0.22	0.01	0.12	0.18	-0.03	0.08	1984	0.00	0.23	0.12	0.00	0.23	0.12
1954	-0.05	-0.21	-0.13	-0.07	-0.23	-0.15	1985	-0.04	0.23	0.10	-0.04	0.23	0.10
1955	-0.02	-0.26	-0.14	0.02	-0.22	-0.10	1986	0.10	0.23	0.17	0.10	0.23	0.17
1956	-0.23	-0.22	-0.23	-0.14	-0.13	-0.14	1987	0.27	0.39	0.33	0.19	0.31	0.25
1957	0.06	0.09	0.08	0.08	0.11	0.10	1988	0.32	0.36	0.34	0.29	0.33	0.31
1958	0.19	0.11	0.15	0.14	0.07	0.10	1989	0.25	0.24	0.25	0.35	0.32	0.33
1959	0.08	0.06	0.07	0.06	0.04	0.05	1990	0.47	0.31	0.39	0.46	0.30	0.38
1960	0.08	-0.02	0.03	0.09	-0.01	0.04							

^aExpressed in degrees Celsius.

^bData corrected for the influences of El Niño/Southern Oscillation (ENSO).

Appendices

Appendix A

Carbon Dioxide Information Analysis Center

The Carbon Dioxide Information Analysis Center (CDIAC), located within the Environmental Sciences Division of Oak Ridge National Laboratory, has been in operation since 1982. CDIAC provides information support to the international research, policy, and education communities for the evaluation of complex environmental issues associated with elevated levels of atmospheric CO₂, including potential climate change. CDIAC activities include obtaining and evaluating data, articles, and reports; producing digital numeric data and computer model packages (NDPs and CMPs); distributing CO₂-related reports; and producing the newsletter, *CDIAC Communications*, which has a worldwide distribution of over 5500 in 151 countries. To date, 38 NDPs and CMPs have been compiled, 12 issues of *CDIAC Communications* have been distributed, and CDIAC has responded to more than 16,000 information requests.

CDIAC is funded by the U.S. Department of Energy (DOE) to support its Carbon Dioxide Research Program (CDRP). The goal of CDRP is to develop sound scientific information for policy formation and governmental action in response to changes of atmospheric CO₂. CDRP's thrust during the past decade has been to (1) elucidate the processes that control the global carbon cycle and provide predictions of future atmospheric CO₂ change; (2) develop data and models of the processes by which changes in the Earth's radiative balance may change climate at global and regional scales and predict rates of potential climate change; and (3) develop the data and models required to define and predict the combined effect of climate and CO₂ on plants, crops, and ecosystems.

In 1989, CDRP moved from the DOE Office of Basic Energy Sciences into the DOE Office of Health and Environmental Research. At the same time, it was combined with other atmospheric research activities under the Environmental Sciences Division, which is directed by Ari Patrinos. The CDRP managers (and their areas of responsibility) are Roger C. Dahlman (global carbon research, vegetation research, and education), Michael R. Riches (climate research, quantitative links, oceans research, and resource analysis), and Thomas J. Gross (information and integration, including responsibility for CDIAC).

The mailing address for the program is:

Carbon Dioxide Research Program
Environmental Sciences
Division (ER-74)
U.S. Department of Energy
Washington, DC 20585, U.S.A.

Contributing Investigators

Abstract

Foreword

Acknowledgments

ATMOSPHERIC CO₂

Historical Record from Ice Cores

Vostok (*Antarctica*)

Siple Station (*Antarctica*)

Modern Record

Scripps Institution of Oceanography (U.S.A.)

Mauna Loa (*Hawaii, U.S.A.*)

American Samoa (*U.S. Territory, South Pacific Ocean*)

Point Barrow (*Alaska, U.S.A.*)

South Pole (*Antarctica*)

National Oceanic and Atmospheric Administration, Climate Monitoring and Diagnostics Laboratory, Continuous Monitoring Program (U.S.A.)

Amundsen Scott (*South Pole, Antarctica*)

Cape Matatula (*American Samoa, U.S. Territory, South Pacific Ocean*)

Mauna Loa (*Hawaii, U.S.A.*)

Point Barrow (*Alaska, U.S.A.*)

National Oceanic and Atmospheric Administration, Climate Monitoring and Diagnostics Laboratory, Flask Sampling Program (U.S.A.)

Alert (*Northwest Territories, Canada*)

Amsterdam Island (*Indian Ocean*)

Amundsen Scott (*South Pole, Antarctica*)

Ascension Island (*South Atlantic Ocean*)

Cape Grim (*Tasmania, Australia*)

Cape Kumukahi (*Hawaii, U.S.A.*)

Cape Matatula (*American Samoa, U.S. Territory, South Pacific Ocean*)

Cape Meares (*Oregon, U.S.A.*)

Christmas Island (*Kiribati, South Pacific Ocean*)

Cold Bay (*Alaska, U.S.A.*)

Halley Bay (*Antarctica*)

Key Biscayne (*Florida, U.S.A.*)

Mahé Island (*Seychelles, Indian Ocean*)

Appendix B

National Oceanic and Atmospheric Administration, Climate Monitoring and Diagnostics Laboratory, Flask Sampling Program (U.S.A.), cont'd.

Mariana Islands (*Guam, Pacific Ocean*)
Mauna Loa (*Hawaii, U.S.A.*)
Mould Bay (*Northwest Territories, Canada*)
Niwt Ridge (*Colorado, U.S.A.*)
Ocean Station "M" (*North Atlantic Ocean*)
Palmer Station (*Anvers Island, Antarctica*)
Point Barrow (*Alaska, U.S.A.*)
Ragged Point (*Barbados, Atlantic Ocean*)
Sand Island (*Midway Islands, South Pacific Ocean*)
Shemya Island (*Alaska, U.S.A.*)
St. Croix (*U.S. Virgin Islands, Atlantic Ocean*)
Terceira Island (*Azores, North Atlantic Ocean*)

Other Monitoring Programs

Cape Grim (*Tasmania, Australia*)
Alert (*Northwest Territories, Canada*)
Cape St. James (*British Columbia, Canada*)
Sable Island (*Nova Scotia, Canada*)
Amsterdam Island (*Indian Ocean, France*)
Garmisch-Partenkirchen (*Germany*)
Wank Peak (*Germany*)
Zugspitze (*Germany*)
K-Pusztá (*Hungary*)
Mt. Cimone (*Italy*)
Izaña (*Canary Islands, Spain*)
Bering Island (*U.S.S.R.*)
Kotelny Island (*U.S.S.R.*)
Ocean Station "C" (*North Atlantic Ocean, U.S.S.R.*)
Teriberka Station (*U.S.S.R.*)

ATMOSPHERIC CH₄

Historical Record from Ice Cores

Vostok (*Antarctic*)
Byrd Station (*Antarctica*)
Dye 3 (*Greenland*)
Siple Station (*Antarctica*)
Crête (*Greenland*), Camp Century (*Greenland*), and Byrd Station (*Antarctica*)

Contents—Trends '91

Appendix B

Modern Record

Oregon Graduate Institute of Science and Technology

Global

*National Oceanic and Atmospheric Administration, Climate Monitoring
and Diagnostics Laboratory, Flask Sampling Program*

Alert (Northwest Territories, Canada)

Amsterdam Island (Indian Ocean)

Amundsen Scott (South Pole, Antarctica)

Ascension Island (South Atlantic Ocean)

Cape Grim (Tasmania, Australia)

Cape Kumukahi (Hawaii, U.S.A.)

Cape Meares (Oregon, U.S.A.)

Christmas Island (Kiribati, South Pacific Ocean)

Cold Bay (Alaska, U.S.A.)

Halley Bay (Antarctica)

Key Biscayne (Florida, U.S.A.)

Kitt Peak (Arizona, U.S.A.)

Mahé Island (Seychelles, Indian Ocean)

Mariana Islands (Guam, North Pacific Ocean)

Matatula Point (American Samoa, U.S. Territory, South Pacific Ocean)

Mauna Loa (Hawaii, U.S.A.)

Mould Bay (Northwest Territories, Canada)

Niwot Ridge (Colorado, U.S.A.)

Ocean Station "M" (North Atlantic Ocean)

Olympic Peninsula (Washington, U.S.A.)

Palmer Station (Antarctica)

Point Barrow (Alaska U.S.A.)

St. Croix (U.S. Virgin Islands, Atlantic Ocean)

Syowa (Antarctica)

Terceira Island (Azores, North Atlantic Ocean)

OTHER TRACE GASES

Atmospheric Concentrations—Historical Record: N₂O from Ice Cores

Crête (Greenland), Camp Century (Greenland), and Byrd Station (Antarctica)

Atmospheric Concentrations—Modern Record

Adrigole (Ireland)

Cape Grim (Tasmania, Australia)

Cape Meares (Oregon, U.S.A.)

Appendix B

Atmospheric Concentrations—Modern Record, cont'd.

Point Matatula (*American Samoa, U.S. Territory, South Pacific Ocean*)

Ragged Point (*Barbados, Atlantic Ocean*)

Chlorofluorocarbon Release Estimates

CFC-11 (CFCl₃)

CFC-12 (CF₂Cl₂)

CO₂ EMISSIONS

Historical Record

Global

Modern Record

Global

Regional (ranked in 1989 descending order)

North America

Eastern Europe

Western Europe

Centrally Planned Asia

Far East

Oceania

Developing America

Middle East

Africa

National (the highest 20, ranked in 1989 descending order)

United States

Union of Soviet Socialist Republics

People's Republic of China

Japan

India

Federal Republic of Germany

United Kingdom

Canada

Poland

Italy

France

German Democratic Republic

Mexico

South Africa

Contents—Trends '91

Appendix B

National (the highest 20, ranked in 1989 descending order), cont'd.

Australia
Czechoslovakia
Republic of Korea
Romania
Brazil
Spain

TEMPERATURE

Global and Hemispheric Anomalies

Global, Hemispheric, and Zonal Anomalies

Global (90° N–90° S)
North Polar (60° N–90° N)
North Temperate (30° N–60° N)
North Subtropics (10° N–30° N)
Equatorial (10° N–10° S)
South Subtropics (30° S–10° S)
South Temperate (60° S–30° S)
South Polar (90° S–60° S)
Tropical (30° N–30° S)
Northern Hemisphere (0°–90° N)
Southern Hemisphere (0°–90° S)

U.S. National

National
North Pacific Coast
South Pacific Coast
North Cascades
California Interior Valleys
East Slope North Cascades
Great Basin
Southern Desert
Northern Rockies
Southern Rockies
Northern Steppes
Southern Steppes
Northern Plains
Southern Plains
South Coastal Plain
Gulf Coast

Appendix B

U.S. National, cont'd.

- Great Lakes
- Eastern Prairies
- Northern Appalachians
- Southern Appalachians
- Northern Piedmont
- Southern Piedmont
- Coastal Northeast
- Coastal Southeast

APPENDICES

Appendix A: Carbon Dioxide Information Analysis Center

Appendix B: Numeric Data Packages

Appendix C: Acronyms, Abbreviations, and Chemical Formulas

Order Form

Appendix C

To order additional copies of the highlights or the full copy of *Trends '91: A Compendium of Data on Global Change*, simply complete this order form and return to CDIAC. Inquire about bulk orders of either the summary or full document for classroom instruction.

Request for *Trends '91*:

- ☐ Please send a copy of the full document
- ☐ Please send a copy of the numeric data in *Trends '91* in PC format (specify floppy diskettes):
 - ☐ 5¼ in. ☐ high density
 - ☐ 3½ in. ☐ low density

Other requests:

- ☐ *Catalog of Data Bases and Reports*
- ☐ Sample issue of DOE Research Summary series
- ☐ *Glossary: Carbon Dioxide and Climate*
- ☐ Sample issue of newsletter, *CDIAC Communications*

Name: _____

Organization: _____

Mailing Address: _____

City: _____ State: _____ ZIP: _____ Country: _____

Telephone: _____ e-mail: _____

Send completed form to:

Carbon Dioxide Information Analysis Center
Oak Ridge National Laboratory
P.O. Box 2008, MS-6335
Oak Ridge, Tennessee 37831-6335, U.S.A.
(615) 574-0390
(615) 574-2232 FAX

BITNET: CDP@ORNLSTC
OMNET: CDIAC
INTERNET: CDP@STC10.CTD.ORN.LGOV.

END

DATE
FILMED

4 / 1 / 93