NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

DOE/ER/40281--T?

DE92 016285

Progress Report

"Relativistic Heavy Ion Research"

DE FG02 86ER40281

Prof. Shoji Nagamiya
Principal Investigator

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

1 Overview of Current Activities

The main focus of our group at BNL during the past five years has been participation in the E802/859/866 experiment, which is one of the major heavy ion experiments at the BNL-AGS. Data-taking for E802/859 is now completed, and extensive data analysis is in progress. The Nevis group is actively involved in this effort. During CY92, the construction of various detector elements for E866 will be finished and in CY93 it is expected that the data taking of E866 will start. In E866 the Nevis group will construct wire chambers for the existing spectrometer arm. In addition, we plan to contribute to the data acquisition system upgrade and to maintain our investment in the Level II triggering system developed for E859.

At BNL the Nevis group has played a dominant role. For example, S. Nagamiya was the co-spokesman of E802 (with O. Hansen at the initial stage and C. Chasman at a later stage) and W. Zajc was the co-spokesman of E859 (with L. Remsberg). The Nevis group constructed key detectors, such as the segmented Cerenkov counter and the streamer-tube-based back counter. The Nevis group made major contributions to the design and implementation of the Level II Trigger in E859. The major portion of the E802/E859 Monte Carlo was written at Nevis, and the group has, in cooperation with MIT and BNL, participated extensively in the offline data analysis.

The most exciting event at the BNL-AGS this year was the successful acceleration of Au beams. We took preliminary data in April '92 using beams of 11.7 A-GeV/c Au ions, as part of a survey run for E866. E866 is an extension of the E802-style experiment to Au beams. We plan to maintain our active involvement in this experiment as we did for E802/859. With the advent of truly heavy ion beams, we feel that the next few years will be a most exciting time for the BNL-AGS heavy ion program. This excitement, along with the chance for direct involvement with the experiment, make this program very attractive to graduate students. It is fortunate indeed that this opportunity to study nuclear matter at extreme baryon density has a timescale optimized for the training of those graduate students who will go on to post-doctoral careers at RHIC. The recent appointment of B. Cole as a post-doc significantly strengthens the Nevis contribution to the forthcoming E866 experiment.

Since 1989 the group also performed serious design work on a RHIC experiment. After various events, we are now playing a key role in the experiment called PHENIX; one of the two major experiments at RHIC. S. Nagamiya is the spokesman of this experiment. In addition, W. Zajc and C.Y. Chi are playing an important role in the design of the triggering system in PHENIX, while T. Nayak has spent over a year developing new TOF systems for RHIC. An overview of the detector concept as of January 1992 may be found in Attachment #1.

The Columbia group has also participated in a small-scale experiment, E858/878 at the BNL-AGS. Participation in this experiment has been, however, on an individual basis rather than based on the involvement of the entire group. One of our students (P. Stankus) is writing a thesis based on data from E858, and one of our Chinese visitors (W. L. Zhan) is now participating in E878.

At CERN, the ongoing program in heavy ion physics consists of experiment NA44. The goal of this experiment is to measure two-particle correlations with high statistics and high momentum resolution for pairs of positive and negative K-mesons, as well as pions. The experiment was designed to provide optimum results with the heavy nuclear beams, up to lead, which are expected to be available at the SPS in 1994. It utilizes large aperture superconducting quadrupoles to achieve relatively large angular coverage in the center-of-mass system, at high momentum resolution of 0.25%, but for relatively small coverage of the particle momentum ($\Delta p < 0.15p$). This work at CERN is also one of the major activities of the Nevis heavy ion group.

The current members of the Nevis heavy ion group are summarized in the table on the following page.

Members of the Nevis Heavy Ion Group				
Name	Status	DOE no. (FY92)	Main Research Activity	Comments
S. Nagamiya	Prof.	2	RHIC, AGS	PHENIX Spokesman
W. Willis	Prof.	1	CERN + SSC	
W. Zajc	Assoc. Prof.	2	AGS, RHIC	E859 Spokesman
J. Dodd	Postdoc	12	CERN	
C. Y. Chi	Postdoc	6	RHIC, AGS	50% System Eng. for Nevis Computer
B. Cole	Postdoc	12	AGS, RHIC	
T. Nayak	Postdoc	0	RHIC, AGS	Paid by RHIC R&D
P. Stankus	GRA	8	AGS-E858	ORNL postdoc soon
O. Vossnack	GRA	12	AGS-E859	
F. Q. Wang	GRA	12	AGS-E866	
Y. F. Wang	GRA	12	AGS-E859/866	
Y. D. Wu	GRA	7	AGS-E866	
B. Budick	Visiting	0	AGS	Professor, NYU
X. H. Yang	Visiting	10	RHIC	Peking U.
Q. M. Li	Visiting	12	RHIC	IHEP, Beijing
W. L. Zhan	Visiting	12	AGS, RHIC	Lanzhou

(In addition, W. Sippach, the senior Nevis electronics engineer is actively involved in the RHIC design effort.)

2 Review of the Past Three-Year Activity

2.1 Physics from E802

The highlights for the first three years in E802 (1986-9) were

- Nuclear transparency and stopping power from energy flow data: It was discovered that the nucleus is "black" in terms of pion production at the AGS energy, suggesting that large target nuclei such as Au completely stop the Si projectile nucleus. This, coupled with the significant particle production observed at AGS energies, indicates that these energies are optimal for the production of high baryon density nuclear matter.
- Anomalous K^+ enhancement: The observed K^+/π^+ ratio is larger than 20%, which is an enhancement of more than a factor of 3 as compared to the ratio observed in pp (or pBe) collisions[1, 2]. The mechanism of the strangeness enhancement and K^+ distillation has been discussed theoretically in terms of quark-gluon plasma, normal hadronic processes, media effects of the hadronic gas, etc.

During the last three years (1989-92), much more refined data analysis has been done. We describe here only a few highlights:

- Anti-proton production: In the last year E802 published the first results of measurements of anti-proton production in A-A collisions[3]. These data are of intense interest to theorists as the annihilation of anti-protons in the medium is thought to be dependent strongly on the baryon densities attained during the collisions. The measured anti-proton yields change only slightly between Si+Al and Si+Au collisions. The understanding of these results is ambiguous since the lack of increase may be attributed either to a slow increase in the production with increasing target thickness or annihilation of anti-protons in the final state. Regardless, these data will constrain any model of A-A collisions that tries to describe anti-proton production.
- Bose-Einstein correlations: E802 has reported[4] the first direct measurements of this pion source size at AGS energies, as determined by Bose-Einstein correlations measured for both $2\pi^+$ and $2\pi^-$ events from Si+Al and Si+Au collisions. These data have been analyzed in a variety of ways. For example, Figure 1 shows raw and corrected correlation data analyzed in terms of q_T , the component of pair relative momentum perpendicular to the beam axis. While our results indicate that the measured transverse radii are larger than the longitudinal radii, a simple analysis has demonstrated that this results not from a geometric deformation but from the non-zero lifetime. In fact, all results are consistent at the 2σ level with a spherical source with radius and lifetime given by $R = c\tau = 2$ fm (Figure 2.)

A Physical Review Letter summarizing these findings has recently been submitted.

Figure 1: The effect of successive corrections on $C_2(q_L, q_T)$ in the region $q_L < 10$ MeV for the $^{28}\text{Si} + ^{197}\text{Au} \rightarrow 2\pi^- + X$ data set: a.) Raw ratio b.) Corrected with the Gamow factor c.) Corrected for the Gamow factor plus acceptance corrections. The result of the 2-d fit is also shown projected into the same q_L slice as the data. d.) Fully corrected data for the $^{28}\text{Si} + ^{197}\text{Au} \rightarrow \pi^-\pi^+X$ data set.

Figure 2: Confidence contours for radii and lifetimes determined by fits to the form of Eq. 2 for a.) $^{28}\text{Si} + ^{27}\text{Al} \rightarrow 2\pi^- + X$, b.) $^{28}\text{Si} + ^{197}\text{Au} \rightarrow 2\pi^- + X$, c.) $^{28}\text{Si} + ^{27}\text{Al} \rightarrow 2\pi^+ + X$, d.) $^{28}\text{Si} + ^{197}\text{Au} \rightarrow 2\pi^+ + X$.

- **High-transverse momentum spectra:** During the last year, the analysis of data taken with the E802 Cherenkov Complex was completed. The Cherenkov Complex provides identification of particles at high $p_{\perp}$ over a limited angular region of the E802 main spectrometer acceptance. These data extend the upper limit of our $p_{\perp}$ coverage near mid-rapidity from 0.8 GeV/c to 3.5 GeV/c for protons (0.5 GeV/c beyond the nucleon-nucleon kinematic limit) and to 2.5 GeV/c and 2.0 GeV/c for pions and kaons, respectively, (just beyond the nucleon-nucleon kinematic limits for both). The high- $p_{\perp}$ data, shown in Figure 3 for Si+Al and Si+Au collisions, indicate that the exponential behavior of the invariant cross-sections when plotted as a function of transverse mass ($m_{\perp} \equiv \sqrt{p_{\perp}^2 + m^2}$),

$$E \cdot d^3\sigma/dp^3 = A \cdot e^{-(m_{\perp}-m)/T},$$

observed in the E802 data at low $p_{\perp}$ [1] continues over the entire measured $p_{\perp}$ range. No deviations in the exponential behavior are seen at high- $p_{\perp}$, even beyond the nucleon-nucleon kinematic limits.

The high- $p_{\perp}$ data also show that the increase in the K^+/π^+ ratio observed in the the $p_{\perp}$ -integrated yields is present over the entire $p_{\perp}$ range. This result excludes any description of the kaon enhancement that relies on the additional kaons being produced at low $p_{\perp}$ or on absorption of pions at low $p_{\perp}$. We have investigated the centrality dependence of the transverse spectra at mid-rapidity. No significant changes in the shape of the spectra are seen with increasing centrality, though the inverse slope parameter, T , increases monotonically with increasing target thickness and increasing produced particle multiplicity.

The most peripheral Si+Al spectra are extremely similar to transverse spectra at mid-rapidity from pp collisions. Studies of pp, peripheral Si+Al, and inclusive Si+A pion spectra using an alternative variable, $p_{\perp}^2$, that is often used in analysis of resonance production, suggest that the observed simple $m_{\perp}$ dependence of the transverse spectra arises from a very complicated superposition of baryon resonance decays, meson resonance decays and direct pion production processes.

The final high- $p_{\perp}$ data are currently being prepared for publication.

2.2 AGS Experiment E859

To further study the nature of strangeness enhancement in heavy ion collisions, the E859 collaboration[5] has upgraded the E802 apparatus to provide very high statistics single-particle spectra for charged kaons produced in these reactions. Additionally, E859 has chosen to examine the space-time structure of the meson source(s) by measuring $2\pi^{\pm}$ and $2K^+$ correlations, as well as making more precise measurements of anti-proton production.

All of these highly desirable physics topics are characterized by their low production cross sections. The difficulty of triggerin on these rare events is exacerbated by the broad-band acceptance of the E802 spectrometer[6], which, while essential for good pair

Figure 3: Transverse spectra at mid-rapidity ($y = 1.7$) for pion, proton, K^+ , and K^- production in Si+Al and Si+Au collisions. The nucleon-nucleon kinematic limits are indicated on the axes.

acceptance, makes development of physics-sensitive Level I hardware triggers extremely difficult. Therefore, E859 decided to develop a Level II trigger that would determine the mass of all charged particles in the spectrometer aperture. Such a trigger must:

- Track and identify all charged particles in a "typical" event (up to roughly 5 tracks/event).
- Cover a dynamic range in momentum from $\sim 0.3 < p < \sim 2.0$ GeV/c.
- Make a trigger decision in $\sim 40\mu s$.
- Be sufficiently flexible to provide user-selectable "physics" triggers at run time.

The E859 Level II trigger was implemented using LeCroy ECLine modules to perform various read-out, table look-up, calculation and control functions[7]. The trigger was designed to be constructed in a two-stage process. The first stage, implemented for the June-90 heavy ion run, used the E802 TOF wall in conjunction with two new MWPC's to find and momentum-analyze tracks. This provided a "two-like-sign" trigger, which was used to significantly extend the E802 two-pion data, and also to obtain a large sample of two-proton events. The last day of the June-90 running period was also used to demonstrate a proof-of-principle of the second stage in the implementation of the Level II trigger, which involves using the TOF wall to provide high quality on-line particle identification. This capability was further developed and exploited in the February-91

14.6 A GeV/c Si+Au $\rightarrow$ K $^-$

Figure 4: Comparison of transverse mass spectra for K $^-$'s from inclusive events (E802) and central events (E859).

run, where we demonstrated the ability to calibrate and maintain the TOF wall to within ~ 100 ps for the entire data-taking period.

By using the full Level II trigger in the February-91 run, E859 improved the E802 statistics on single-particle K $^+$ and K $^-$ spectra by roughly an order-of-magnitude. This is illustrated in Figure 4, which compares the E802 transverse mass spectrum of inclusively-produced K $^-$'s, versus the E859 data for K $^-$'s from central events. Although central events comprise only $\sim 10\%$ of the inclusive cross-section, the improvement in the E859 statistics is evident. Similar improvements have been obtained for K $^+$ and anti-proton spectra.

A major goal of the February-91 run was measurement of the Bose correlations between two positive kaons. Despite devoting less than 25% of the total running time to this effort, we managed to obtain sufficient data to make the first measurement of $2K^+$ correlations, and to compare them to both $2\pi^+$'s and two-protons measured in the same apparatus. These results were presented at the Quark Matter '91 conference in November '91 by W.A. Zajc. A copy of that presentation, which also summarizes other results from E802 and E859, is included as Attachment #2.

E859 concluded its data-taking with a five week run in March and April of 1992. The flexibility and programmability of the Level II trigger was exploited to address the following topics:

- The extended measurements of K $^+$ and K $^-$ spectra over all available rapidities were completed.
- The measurements of π^+ and π^- spectra were extended to the puzzling low-Pt region.

- The measurements of anti-proton spectra were extended to cover a wider range of rapidities. Good evidence for anti-lambdas was observed in the best anti-proton sample (central Si + Au).
- A study of the high- p_T behavior of spectra from peripheral Si + Al collisions was completed.
- The study of two-pion correlations as a function of multiplicity (impact parameter) was extended to the Si + Al system.
- The statistics for $2K^+$ correlations were increased.
- The statistics for $2\pi^-$ and $2\pi^+$ correlation measurements were dramatically increased for central Si + Au collisions.
- A new trigger for Λ 's was commissioned and a significant sample was obtained for central Si + Au collisions.
- The rapidity coverage for the measurements of phi-mesons was increased.

In addition, two physics topics were addressed which did not require the Level II trigger: 1.) The study of baryon spectra in the target fragmentation region was completed and extended to include a third target. 2.) Detector calibrations needed for the first measurements with the gold beam were performed.

The above list indicates that there are many exciting topics to pursue in the offline analysis of the E859 data sets. At Nevis, two graduate students will take their Ph.D's from these data. O. Vossnack's thesis will contain the analysis of the $2K^+$ and $2\pi^+$ correlation data. Y. Wang will work on the measurement of the inclusive cross-section for Φ -production. (A strong signal for Φ production has been one of the "unexpected" results from our triggering on two-kaons.) Additionally, B. Cole and W. Zajc will be involved in many aspects of the overall data analysis. The extensive nature of these measurements presents a real opportunity to gain insight into the nature of strangeness production by high-density nuclear matter. We hope to exploit this opportunity to the fullest by collaborating with the authors of the ARC cascade code at Brookhaven (T. Nayak will lead this effort), and via another joint theoretical venture aimed at a deeper understanding of the nature of the observed Bose-Einstein correlations. This program is also a natural part of the NA44 experiment, and is further discussed in the corresponding section of this proposal.

2.3 AGS Experiment E858/878

The BNL-PAC approved 100 hours of running time for this experiment to search for rare negative secondaries — anti-protons, anti-nuclei and possibly strangelets — produced in heavy ion collisions. H. Crawford (UCSSL-Berkeley) is spokesman for this experiment. From Columbia, P. Stankus and S. Nagamiya joined the experiment.

The experiment was the first to use full-intensity beams of ^{28}Si at the AGS; no stable negative particles heavier than the $\bar{d}$, and none with $|Z| > 1$ were seen. The observed cross sections for π^- , K^- , $\bar{p}$, and $\bar{d}$ produced at 0° were measured. Preliminary results show that the $\bar{d}$ yield is one order of magnitude smaller than the prediction of a coalescence model, while it is two orders of magnitude larger than what a direct $d\bar{d}$ creation calculation would predict. Publication of these data is in progress.

E858 was followed by E878 in the AGS 1992 running period. E878 is an upgraded rare-particle search, and will provide a factor of 20 greater sensitivity through a combination of increased intensity, acceptance and running time (300 hours). P. Stankus was involved in the proposal, design and preliminary setup/tests of E878, and W. L. Zhan participated in the run.

2.4 CERN Experiment NA44

NA44 is a multiple particle focusing spectrometer with very high resolution in angle and momentum and extremely good particle identification. Its large rate capability allows good statistics to be obtained on the spectra and correlations of rare particles, for example anti-protons and pairs of kaons. These measurements are well-known to be of vital importance for the understanding of ultra-relativistic heavy ion collisions, particularly in the case of creation of a quark-gluon plasma (where kaons play a special role in predicted signatures) or a chiral phase transition (where the anti-proton is expected to be enhanced and display special features related to the transition).

This year has seen a number of major developments in this experiment:

- a.) The first heavy ion beams were available for this experiment last October. Data taking began at once, since the spectrometer had been commissioned and thoroughly tested in proton running. Though the ^{32}S beam intensity and reliability were not perfect, essentially all the planned data were obtained. With targets of sulfur and lead, the single particle transverse momentum spectra were obtained for π , K and p of both signs. The design of the spectrometer allows the measurement of these spectra over a wide range, including transverse momenta down to zero. Structure in the low $p_\perp$ range for the heavier and rarer particles constitutes a particularly important source of information about the dynamics of the collision. The two particle correlations were studied with large statistics, particularly for pions.
- b.) The Quark Matter '91 Conference took place near the end of the 1991 run. Since the analysis programs had already been brought to a fairly advanced state on the proton beam data, it was possible to present results at the conference based on data taken a few days earlier, illustrating the power of a focusing spectrometer, which does not require a great deal of computer time per event analyzed. (Data from proton beam runs had been presented at the Lepton-Photon Conference during the previous August.)
- c.) In March 1992, NA44 was approved for running with lead beams in the SPS. The SPSC considered that the experiment would do an adequate job on lead-lead colli-

sions with minor upgrades, but has encouraged us to proceed with certain improvements which enhance the performance, particularly in the very high event particle multiplicities characteristic of these collisions.

- d.) Our collaboration held a two week Workshop in Hiroshima in March 1992, with the goals of fixing a common set of analysis procedures and codes. It was especially valuable to have the Workshop on the home grounds of our "most distant" collaborators, to ensure a full understanding of the issues under circumstances where the collaboration as a whole could direct its attention to the circumstances of the work in a typical local group, and all of the Japanese team could be present at one time. The Workshop was considered a success, and we believe that a fully common and transparent strategy of analysis has been established.
- e.) Our second run began on the seventh of April 1992, and is scheduled to last an unprecedented six weeks. The beam intensity and reliability has been excellent during the first half of this run. The beam intensity which optimizes the rate of accumulation of data is between 5 and 10 million per burst, which is just at the limit of the allowed radiation level in this beam. We have been able to obtain samples of pairs of kaons with the order of 60,000 good events, which allows an analysis in terms of the space-time dimensions of the source in three orthogonal directions. Such an analysis, with adequate statistics, has not heretofore been reported even for pion pairs. Our samples of pion pairs are of course considerably larger.
- f.) During this run we have for the first time carried out the operation of changing to our "large angle" configuration. The whole spectrometer was rotated by 4.5 degrees in the laboratory, to allow us to carry our studies of spectra and correlations to p_t of the order of 1 GeV/c. In order to make this possible, the critical elements are mounted on a large beam which rotates about the fixed point of the target while keeping an alignment accuracy of 0.1 mm. Thanks to this considerable investment on the part of CERN, the move was carried out during the run in half a day, and the alignment of target and beam was indeed found to be within 0.1 mm. Since this is the last run planned with sulfur beams, it is fortunate that it seems that we shall obtain the full set of data we thought necessary.
- g.) An intense period of analysis is now beginning, which is expected to produce some of the main results in about three months. A key element of the analysis is the simulation of the acceptance, including effects of finite target size, focusing properties and aberrations, detector element granularity, multiple Coulomb scattering and kaon decay. This is the primarily the responsibility of Columbia, and is near the end of development for this stage of the experiment.

2.5 PHENIX Contributions

During the last three years the group has contributed largely to the design work for a RHIC experiment. The basic philosophy that our group has always tried to promote in the collaboration is the measurement of as many potential signatures of quark-gluon plasma as possible, as a function of a well defined variable which can be directly connected to the energy density of the system. Fortunately, this approach was agreed to in the collaboration called PHENIX and, thus, the Nevis group is now very actively involved in this experiment.

The PHENIX detector is designed to measure di-leptons (di-electrons in the mid-rapidity region and di-muons in the forward region), photons and hadrons. The proposed detector combines an open-geometry axial magnetic field, tracking chambers, calorimeters and various particle-identification and event characterization devices to provide high-rate capability, good particle identification, and high momentum resolution. The forward muon arm consists of muon tracking and muon identifiers.

The ambitious physics goals of PHENIX raise several technical issues: For example, the experiment will require over thousand channels of time-of-flight with sub-100 ps time resolution. The rare signals sought by PHENIX require a pipelined data acquisition system coupled to a multi-level trigger scheme. In addition, as a candidate for electron identification, the possibility of installing a TRD (transition radiation detector) was discussed. Therefore, during the past years the Nevis group has been involved in serious R&D work in these critical areas. Each is discussed in the following sub-sections.

2.5.1 TOF Tests

Development work on time-of-flight systems for PHENIX was conducted by T. Nayak, C. Y. Chi, X. H. Yang, S. Nagamiya, in collaboration with University of Tsukuba (Y. Miake, et al.). The time-of-flight wall along with beam-beam counters will be used for hadron identification over a wide p_T range. A series of beam tests were conducted both at the KEK-PS and the BNL-AGS. The goal of the tests were two-fold. One was to try to achieve a good time resolution by a scintillator hodoscope (including scintillating fiber bundles and super-scintillator, where the latter is a normal scintillator coated with a small- n material). The other was to test the performance of a newly designed Cerenkov free phototube where Cerenkov light from the phototube window was shielded by separating the photocathode from the window. In addition, efforts are being made to design the front-end electronics and develop a suitable on-tube discriminator.

The following two figures show some typical results from the tests. Figure 5 shows the comparison of light output from a scintillating fiber bundle made of 5mm round fibers of 3m long and a 1.5m long scintillator rod. The light output from the rod is larger compared to the fiber bundle, but the attenuation length is only about 80cm compared to 200cm for the fiber bundle. A scatter plot showing the time-of-flight (TOF) between the start and stop counters and the integrated charge for the case of 5mm round fiber bundles 3m long is given in the left panel of Figure 6, whereas the right panel shows the corresponding TOF distribution. The average value of TOF is taken to be zero. The

Figure 5: Light output from 5mm round scintillating fiber bundle compared to that of a scintillator rod.

Figure 6: a.) Scatter plot of TOF and ADC values after slewing correction for 5mm round fiber bundles and b.) the corresponding TOF spectrum. The solid curve is a Gaussian fit with $\sigma = 117 \text{ ps}$.

TOF resolution for this fiber bundle is shown to be 117 psec. The corresponding TOF resolution for 1.5m long scintillator rod is 77 psec. Currently, we are incorporating the knowledge gained from these tests into a major effort to design a cost effective and high resolution time-of-flight system for PHENIX.

2.5.2 TRD Tests

A TRD prototype detector, (rebuilt at Nevis from one used in experiment NA34), was tested by V. Cherniatin, C. Y. Chi, Y. F. Wang, W. Willis and two summer students at the AGS during June-91 proton run. Studies were made both on the efficiencies of various radiators and on the possibility of using dE/dx to enhance the e/π separation around 1 GeV/c. Signals from the chamber were read out through a pre-amp and a shaper, then recorded in a 40MHz 6 bit fast sampling ADC system. The electrons and pions in the test beam are tagged by two Cerenkov counters before and a lead glass detector after the TRD chamber. The TRD data were taken with seven different types of foam and one type of foil radiator. The dE/dx data were taken at beam momenta of 1.0, 1.5, and 2.0 GeV/c with the detector placed at 0, 20, and 35 degrees with respect to the beam angle. Data from these tests are currently being analyzed at Nevis, BNL and Yale University. The Nevis group took major responsibility for the readout system and online software during the run, and for analysis of the dE/dx data afterwards. The results of the analysis were presented at the IEEE nuclear instrumentation conference last year.

2.5.3 Readout system design

Nevis has taken the responsibility for the design of an analog pipeline chip for front-end signal recording in PHENIX. This chip (based on a switched-capacitor array) will be used in the time of flight system, and is also applicable to the readout of a liquid argon calorimeter (should that technology be chosen for the PHENIX EM calorimeter). This effort is headed by W. Sippach (Nevis electronics engineer), who has been closely involved with the ZEUS analog pipeline design, and will be supported by C.Y. Chi. A preliminary design has been completed and submitted to a silicon foundry. We will perform tests of the prototype chips to determine the yield, stability and timing resolution of this technology.

**DATE
FILMED
8/25/92**

11/11/11