DOE/PC/89785--3

DE93 008383

THE SINGLE ELECTRON CHEMISTRY OF COALS

Report for the Period
April 1, 1990 to June 30, 1990

John W. Larsen
and Robert A. Flowers II

Department of Chemistry
Lehigh University
Bethlehem, PA 18015

July 24, 1990

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific or aerial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE UNITED STATES DEPARTMENT OF ENERGY

MASTER

20

OBJECTIVES AND SCOPE OF WORK

Objectives

To investigate and characterize the single electron reactions of coals in order to determine the role these reactions play in the chemistry of coal.

Scope of Work

1. Attempts will be made to demonstrate that the radicals from the inertinite maceral group will initiate the polymerization of 4-vinylpyridine.
2. The molecule N,N-diphenyl-phenylenediamine will be deposited in coals to characterize their native free radicals.
3. Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) will be used to characterize the numbers and strengths of single electron donors in coals.

DETAILED DESCRIPTION OF TECHNICAL PROGRAM

Introduction

The principal organic chemical component of coals is aromatic structures. Numerous polynuclear aromatic (PNA) systems are contained in coals, including a variety of heterocyclic systems. Many of the PNAs carry oxygen, nitrogen or sulfur substituents. It is expected that many of these aromatic systems, especially the larger ones, should be able to participate readily in charge transfer interactions. Indeed, charge transfer interactions have previously been proposed as significant by Elofsan and Schulz (1976) and by Schwager and Yen (1976). In this paper, we shall present data which demonstrate that coals can serve as very effective donors in charge transfer interactions, and we suggest that this donation is inherently cooperative in nature, involves the overlap of the π systems of many PNA systems and electron acceptors, and is best described as a lattice effect. The individual PNA structures in coals are not effective as charge transfer donors in one-to-one complexes.

As the rank of coal increases, the size of the PNA systems increases only slowly, up to about 89% carbon (Solum et al., 1989 and Hirsch, 1960). The principal change is an increase in the concentration of the PNA systems in the coal. As the concentration of PNA systems increases, long range cooperative interactions become more prevalent. One need only consider the high melting point and low solubility of polynuclear aromatic hydrocarbons to conclude that the cooperative interactions between

even unsubstituted non-polar aromatic systems are quite strong. The relevance of these non-covalent forces to coal structure has been discussed by Quinga (1987). The preferred geometry for the van der Waals interactions is perpendicular for small rings ("T"-shaped) with increasing deviations from perpendicularity as the ring size increases. As the π delocalization increases, interesting optical and electronic effects occur. For example, most anthracite coals are semi-conductors (van Krevelen, 1981).

The field of electrically conducting organic materials is large and fascinating. It has been well reviewed so a detailed discussion here is unnecessary (Soos and Klein, 1975). One family of solid organic conductors is based on the charge transfer from strongly electron donating systems, for example, tetrathiofulvalene (1), to strong electron acceptors such as tetracyanoquinodimethane (TCNQ) (2). In these conductors, extended stacks of donors and acceptors with overlapping π systems are formed. These overlapping orbitals give rise to a conducting band electronic structure. Materials having band structures formed by overlapping systems may also be insulators due to electron correlation effects.

IR and Raman spectroscopies are powerful tools for investigating the structure of organic conductors. This area has been recently reviewed (Ferras and Williams, 1990). The particular technique which we draw on in this paper is to use the change in the CN stretching frequency of TCNQ when it serves as an

acceptor in a charge transfer interaction. The electron transferred to the TCNQ goes into the lowest unoccupied molecular orbital (LUMO) which is an anti-bonding orbital. Population of the LUMO weakens the overall C N interaction. Occupation of the LUMO by one electron gives rise to a 44 cm^{-1} shift in its stretching frequency. It has been shown that the shift in the stretching frequency is linear in the electron population in the LUMO. The shift in the CN stretching frequency is an accurate index of the amount of charge transferred to the TCNQ (Chappel et al., 1981). Similar relationships exist for other good electron acceptors such as tetracyanoethylene (TCNE, 3).

1

2

3

Results and Discussion

A. TCNQ Charge Transfer Complexes with Coals

TCNQ can be readily deposited in coals from pyridine solution. IR spectra of TCNQ and TCNQ in Illinois No. 6 coal are shown in Fig. 1. It is clear that the stretching frequency has been shifted by the full 44 cm^{-1} caused by the transfer of a single electron. Similar behavior has been observed with a variety of coals, including lignites, subbituminous and a range of bituminous coals.

There are two possible explanations for the observed shift. The simplest explanation is that there exist in coals structures

which are excellent single electron donors capable of transferring an electron to TCNQ in the ground state. All of the TCNQ dissolved in the coal is shifted. No uncomplexed TCNQ remains in the sample, as demonstrated by the absence of the unaltered CN stretch at 2227 cm^{-1} . The spectrum shown is for TCNQ in coal in a molar concentration equivalent to approximately 20% of the PNA systems in this coal as deduced from the NMR studies of Solum et al. (1989). It is highly unlikely that 20% of the PNA systems in coal are such good electron donors that the charge transfer complex would have an electron transferred in the ground state. The second explanation is that cooperative interactions between the TCNQ and the aromatic systems in coal have led to the formation of an extended valance band structure, that the TCNQ LUMO is part of this band structure, and that the band is half filled.

B. TCNQ Charge Transfer Complexes with Aromatic Donors

It was necessary to evaluate the charge transfer electron donor ability of aromatic systems of the sort found in coals. Abundant data are available showing that the equilibrium constants for TCNE and aromatic donors are low, no higher than about 300, and most below 100 (Frey et al., 1990 and Merrifield and Phillips, 1950). Equilibrium constants are readily available for TCNE, but rare for TCNQ because TCNE is more soluble than TCNQ. They are similar strength electron acceptors. We obtained the IR spectra of the complexes formed between TCNQ and a variety of aromatic donors. Table I contains a list of typical donors and the

observed changes in the CN stretching frequency. It is immediately apparent that none of the molecules involved is capable of transferring a significant amount of electron density to TCNQ in the ground state. This result is entirely in accord with theory.

Table I Change in the $C\equiv N$ stretching frequency (Δ, ν) of TCNQ upon complexation with aromatic donors at 25° in nitromethane solution

Donor	$\Delta, \nu (\text{cm}^{-3})$
Methoxybenzene	4
2-Methoxynaphthalene	4
9-Methylanthracene	4
9-Phenanthrol	3
Phenathridine	4
1,3,5-Trimethoxybenzene	2
Hexamethylbenzene	3

The association to form complexes is due primarily to dipolar and van der Waals interactions, not stabilization caused by the charge transfer. It is unlikely that the change in stretching frequency of TCNQ in coals is due to the formation of ground state charge transfer dimers. This observation has been confirmed by

ESR spectra which show at most small increases in the population of unshared electrons and which contain no evidence for the formation of the discrete TCNQ radical anion (Silbernagel, 1990). These results will be published elsewhere.

Conclusion

We conclude that the TCNQ interacts with the aromatic groups in the coal to form an extended band structure. The LUMO of the TCNQ is part of the band structure and the band is approximately half filled. Similar results have been obtained with tetracyanoethylene as shown by the spectra in Fig. 2. In the case of TCNE, the conducting band appears to be about 30% occupied. The structure of these complexes is under active investigation, utilizing a variety of techniques.

References

Chappel J.S., Bloch, A.N., Bryden, W.A., Maxfield, M., Poehler, T.O., Cowan, D.O. (1981), Degree of Charge Transfer in Organic Conductors by Infrared Absorption Spectroscopy, J. Am. Chem. Soc., 103, 2442-2443.

Elofson, R.M. and Schluz, K.F. (1967), The Electron Spin Resonance Spectra of Cellulose Chars Treated with Halogens, Amer. Chem. Soc., Div. Fuel Chem. Preprints, 11(2), 513-520.

Ferraro, J. R., Williams, J. M. in Practical Fourier Transform Infrared Spectroscopy, Ferraro, J.R. and Krishnan, K., Ed., Academic Press, San Diego, CA, 1990.

Frey, J.E., Andrews, A.M., Ankoviac, D.G., Beaman, D.N., Du Pont, L. E., Elsner, T.E., Lang, S.R., Oosterbaan Szart, M.A., Seagle, R. E., Torreano, L.A., (1990), Charge-Transfer Complexes of Tetracyanoethylene with Cycloalkanes, Alkenes, and Alkynes and Some of Their Aryl Derivatives, J. Org. Chem., 55, 606-624.

Hirsch, P.B., Carz, L. (1960), A Contribution to the Structure of Coals from X-Ray Diffraction Studies, Phil. Trans. R. Soc. London, Ser. A252, 557-602.

Merrifield, R.E. and Phillips, W.D., (1958), Cyanocarbon Chemistry. II. Spectroscopic Studies of the Molecular Complexes of Tetracyanoethylene, J. Amer. Chem. Soc., 80, 2778-2881.

Quinga, E.M.Y., Larsen, J.W., (1987), Noncovalent Interactions in High-Rank Coals, Energy & FUELS, 1, 300-304.

Schwager, I., Yen, T.F., (1976), Preliminary Examination of Coal Liquefaction Products, American Chemical Society Fuel Division Preprints, 21(5), 199-206.

Silbernagel, B., Gebhard, L., personal communication.

Solum, M.S., Pugmire, R. J., Grant, D.M., (1989), ¹³C Solid-State NMR of Argonne Premium Coals, Energy & Fuels, 3, 187-193.

Soos, Z.G., Klein, D.J., Charge Transfer in Solid State Complexes in Molecular Association, Vol. 1, Ed. by R. Foster, Academic Press, New York, 1975.

Van Krevelen, D.W., Coal. Typology, Chemistry, Physics, Constitution, Elsevier Scientific Publishing Company, New York, NY, 379-381, 1981.

Fig. 1 Diffuse Reflectance FTIR Spectra of a) TCNQ and b) TCNQ Deposited in Illinois No. 6 Coal from Pyridine Solution in a Molar Amount Equivalent to 20% of the PNA Systems in the Coal

Fig. 2 Diffuse Reflectance FTIR Spectra of a) TCNE and b) TCNE Deposited in Illinois No. 6 Coal from Pyridine Solution in a Molar Amount Equivalent to 20% of the PNA Systems in the Coal

END

DATE
FILMED

4 / 1 / 93