MASTER

UCID- 17926

Lawrence Livermore Laboratory

PRELIMINARY SUMMARY OF PARTICLE TRANSPORT
EFFECTS IN NON-AXISYMMETRIC TANDEM MIRRORS

D. E. Baldwin

September 15, 1978

U. S. GOVERNMENT

PRELIMINARY SUMMARY OF PARTICLE TRANSPORT
EFFECTS IN NON-AXISYMMETRIC TANDEM MIRRORS

by

D. E. Baldwin*
Lawrence Livermore Laboratory
Livermore, CA 94550

ABSTRACT

This report reviews the physical basis for the theory of enhanced transport in non-axisymmetric tandem mirror systems recently published by Ryutov, et al. For TMX and thermal ions in a reactor, the radial loss is estimated to be somewhat less than the axial loss; energetic alphas in reactors are susceptible to rapid loss. A number of variations of current magnetic field designs are suggested for reducing this transport.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

* Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore Laboratory under contract number W-7405-ENG-48.

I. Introduction

In a recent series of papers, Ryutov and coworkers have developed the theory of neo-classical and related transport in non-axisymmetric tandem mirror systems.¹⁻⁴ Because of the importance of this work in accessing the tandem mirror as a reactor concept and in designing nearer term experiments, it was deemed useful to summarize the physical basis for their results and their implications.

This report is devoted to that purpose. We first repeat their heuristic derivation of radial diffusion coefficients, without the more detailed determination of numerical coefficients. From these we estimate particle lifetimes with respect to radial transport both in TMX⁵ and for the previously published LLL preliminary conceptual tandem mirror reactor design.⁶ In summary, we find that the loss due to the worst of such transport is less than previously calculated axial losses, but for existing magnetic field designs the margin of difference is sufficiently small to warrant more careful calculations. Finally, we present some ideas for altered field design which can reduce this transport.

II Drift Surface Properties when $\omega_{\text{bounce}} \gg \omega_{\text{drift}}$

Under conditions that particles' axial bounce frequencies are high compared to their azimuthal drift frequencies, they drift on surfaces of constant

$$J = \sqrt{2} \oint ds \sqrt{\epsilon - \mu B - q\phi} \quad (1)$$

where q, ϵ, μ are the particle charge, energy, and magnetic moment, ϕ is the ambipolar potential, and s measures distance along the field line. In a system which is long and thin in the sense that field lines deviate from a straight axis only by a distance small compared to the scale of axial variation, J may be expanded near the z -axis in a Fourier series in which the odd powers of r vanish due to the basic quadrupole character of the plugs,

$$J(\epsilon, \mu, r, \theta, z) = J_0(\epsilon, \mu, r, z) + \sum_{n=1}^{\infty} r^{2n} \cos 2n\theta J_{2n}(\epsilon, \mu, r, z) \quad (2)$$

where $J_{2n} r \rightarrow 0$ constant. If the fans at the ends of the solenoid have a 90° mutual orientation, there is an important symmetry about the solenoidal midpoint ($z = 0$), i.e., $z \rightarrow -z$, $\theta \rightarrow \theta/2 - \pi$. From Eq. (2) it follows that

$$J_{2n}(\epsilon, \mu, r, -z) = (-1)^n J_{2n}(\epsilon, \mu, r, z). \quad (3)$$

In particular this means

$$J_2(\epsilon, \mu, r, z = 0) = 0, \quad (4)$$

so that the θ -dependence of drift surfaces at the midplane occurs only in order r^4 ,

$$J(\epsilon, \mu, r, \theta, 0) = J_0(\epsilon, \mu, r, 0) + r^4 J_4(\epsilon, \mu, 0, 0) \cos 4\theta + \dots, \quad (5)$$

or the curve of the interception of a drift surface with the midplane satisfies to lowest order

$$\frac{dr}{d\theta} = \frac{4r^4 J_4 \sin 4\theta}{\partial J_0 / \partial r}. \quad (6)$$

This lowest order cancellation of the θ -dependence of the drift surfaces for configurations satisfying $z \rightarrow z$, $\theta \rightarrow \theta/2 - \pi$ has been termed omnigenity by Hall and McNamara.⁷

To estimate J_4 and $\partial J_0 / \partial r$, we assume that the solenoid has a uniform midsection of length L_c and two transition regions, each of length L_{tr} , where the radius of curvature of the magnetic lines is of order $L_{tr}^2 r^{-1}$. Then we estimate

$$\left| \frac{\partial J_0}{\partial r} \right| = 0 \left| \frac{2L_{tr}}{v_{\parallel}} \frac{r_{\perp} B}{L_{tr}^2} + \frac{L_c}{v_{\parallel}} \left(\mu \frac{dB}{dr} + \frac{d\phi}{dr} \right) \right| \quad (7)$$

where the first term in parentheses originates from the finite- β depression of the solenoidal field, and the second term from EXB drift. J_4 has contribution only from the transition regions and has θ -dependence due only to the non-axisymmetric components of the field,

$$J_4 = 0 \left| \frac{L_{tr}}{v_{\parallel}} \frac{\mu B_0}{L_{tr}} \right|. \quad (8)$$

(In practice, the scale length for the axisymmetric and non-axisymmetric components may differ somewhat. Here we do not distinguish and use the common length L_{tr} .)

In strictly vacuum magnetic fields, for which only the first term of Eq. (7) contributes, Eqs. (6-8) give as an estimate for the departure of drift surfaces from circularity

$$\frac{\delta r}{r} = 0 \left(\frac{r^2}{L_{tr}^2} \right). \quad (9)$$

Shown in Fig. (1) are the δr 's for three radii calculated by J. Foote integrating guiding center particles in the numerically generated fields for one set of currents in the TMX magnet set. The cubic dependence of δr on r justifies our paraxial expansion; measurement of the slope and interpreting Eq. (9) as an equality, gives a measure of $L_{tr} = 1.1$ meters, which is entirely reasonable based on the TMX physical dimensions.

By estimating

$$\frac{dB}{dr} = 0 \left| \frac{\frac{1}{2} \beta}{1-\beta} \frac{\mu B r}{r_c^2} \right| \quad (10)$$

and

$$\frac{d\phi}{dr} = 0 \left(\frac{r T_e}{r_c^2} \right), \quad (11)$$

where r_c is the central cell radius, it may be seen that whenever the solenoid contains plasma, $\partial J_0 / \partial r$ will be dominated by these terms, leading to the much smaller estimate,

$$\frac{\delta r}{r} = 0 \left(\frac{r_c^2}{L_c L_{tr}^3} \right). \quad (12)$$

The rapid azimuthal drift induced by dB/dr or $d\phi/dr$ reduces the radial displacement, provided the two drifts do not cancel. It is convenient to introduce the parameter

$$\alpha \equiv \frac{r_c^4}{L_c^3 L_{tr}} \quad (13)$$

as a measure of the radial displacement under these conditions, ie., $\delta r/r \approx \alpha r^2/r_c^2$. For TMX, typical parameters are $r_c \approx 0.3\text{m}$, $L_c \approx 3\text{m}$, and $L_{tr} \approx 1.3\text{m}$, so that $\alpha \approx 10^{-3}$. For a reactor, typical parameters are $r_c \approx 1.0\text{m}$, $L_c \approx 100\text{m}$, $L_{tr} \approx 10\text{m}$, so that $\alpha \approx 10^{-5}$. The drift orbit diffusion resulting from such displacements is very small.

Larger displacements are suffered by those particles for which $\partial J_0 / \partial r$ vanishes locally, in which case the higher order expansion of Eq. (5) leads to the estimate

$$\frac{\delta r}{r} \approx \alpha^{1/2} \frac{r}{r_c}. \quad (14)$$

Determining the conditions for the drift cancellation, which occurs only for ions if the radial electric field is positive, requires models for $B(r)$ and $\phi(r)$.

Throughout the uniform solenoid, $B(r)$ will be determined by perpendicular pressure balance,

$$B_0^2 = B^2(r) + 8\pi p(r), \quad (15)$$

where B_0 is the vacuum field. For a parabolic pressure $p = p_0(1 - r^2/r_c^2)$ this gives

$$\frac{dB}{dr} = -B \frac{B_0}{1-B} \frac{r}{r_c^2} \quad (16)$$

where $\beta = 8\pi p/B_0^2$ and $B_0 = B(0)$.

Ryutov, et al., have argued that electrons trapped in the plugs will be non-omigenous, leading to a thorough radial mixing. This conjecture is based on electron confinement in electrostatic potentials in the plugs which are not symmetric, on the solenoid side vs. the outside. Based on this radial mixing, they argue that T_e will be global, and ϕ will be distributed radially by a Boltzmann dependence on the ion density - even in the solenoid. Adopting this as a model for ϕ , they obtain

$$\begin{aligned} |e| \frac{d\phi}{dr} &= + T_e \frac{1}{n} \frac{dn}{dr} \\ &= \frac{2T_e n_0}{n} \frac{r}{r_c^2} \end{aligned} \quad (17)$$

for the parabolic profile. Thus ions of magnetic moments

$$\begin{aligned} \mu = \mu_*(r) &= - \frac{q}{dB/dr} \frac{d\phi}{dr} \\ &= -2 \frac{T_e}{B} \frac{1-B}{B_0} \end{aligned} \quad (18)$$

have zero azimuthal drift. Ions of neighboring μ and r drift on surfaces whose midplane interceptions are banana-shaped curves centered on $\mu = \mu_*(r)$. While tracing out a banana of characteristic width $\propto r_c^{1/2}$, ions have an azimuthal velocity of order $\propto r_c^{1/2} \omega_{drift}$, so that the period for execution of a banana orbit is of order $\propto r_c^{-1/2} \omega_{drift}^{-1}$. An example of a banana orbit embedded in a family of conventional axis encircling drift orbits is sketched in Fig. 2.

III. Neo-classical Diffusion Coefficients

When $\beta \ll 1$, so that no banana orbits occur for ions in the bulk distribution, the radial diffusion coefficient becomes

$$D = \alpha^2 r_c^2 \frac{v_i}{1 + v_i^2 \tau_{dr}^2} \quad (19)$$

where $\tau_{dr} = \omega_{drift}^{-1}$ is the azimuthal drift time. (For this and other estimates of diffusion coefficients, we set $r = r_c$. However, the r -dependence of diffusion coefficients would be important in any more detailed calculations.) When banana orbits can occur, the degree of collisionality determines the diffusion regime. By expanding Eq. (7) to higher order around μ_* , it may be seen that the width in μ of a banana

is $\Delta\mu/\mu \approx \alpha^{1/2}$, so that the life-time of a particle on a banana

is $\alpha v_i^{-1} \equiv v_{eff}^{-1}$, where v_i is the ion-ion 90°-collision frequency.

Similarly the time to drift around a banana is $\tau_{ban} \approx \alpha^{-1/2} \omega_{dr}$. When $v_{eff} \ll \tau_{ban}^{-1}$, or $v_i \tau_{dr} \ll \alpha^{3/2}$, the diffusion step size is the banana width, the rate is v_{eff} , and the number participating is $\Delta\mu/\mu$, giving

$$D_{neo\ ban} \approx \left(\alpha^{1/2} r_c \right)^2 \left(\frac{v_i}{c} \right) \alpha^{1/2} \approx \alpha^{1/2} r_c^2 v_i. \quad (20)$$

When $\alpha^{3/2} < v_i \tau_{dr} < 1$, there is a wider range of μ which can scatter into a banana in one τ_{ban} , determined by

$$\left(\frac{\Delta\mu}{\mu} \right)^2 \frac{1}{v_i} \approx \tau_{dr} \frac{\mu}{\Delta\mu}$$

or

$$\left(\frac{\Delta\mu}{\mu} \right)^3 \approx \begin{cases} v_i \tau_{dr} & \text{if } v_i \tau_{dr} < 1 \\ 1 & \text{if } v_i \tau_{dr} > 1 \end{cases}. \quad (21)$$

The radial velocity is given by $\alpha \omega_{dr} r_c$, and the correlation time is

v_{eff}^{-1} , giving a diffusion coefficient

$$D \approx \left(\alpha \omega_{dr} r_c \right)^2 \left(\frac{\Delta\mu}{\mu} \right)^2 \frac{1}{v_i} \frac{\Delta\mu}{\mu}$$

or

$$\begin{aligned}
 D_{\text{neo plat}} &\approx \alpha^2 \omega_{\text{dr}}^2 r_c^2 & \alpha^{3/2} < v_i \tau_{\text{dr}} < 1 \\
 D_{\text{neo coll}} &\approx \alpha^2 \omega_{\text{dr}}^2 r_c^2 / v_i & 1 < v_i \tau_{\text{dr}}
 \end{aligned} \quad (22)$$

The banana, plateau, and collisional regimes are shown in Fig. (3). Note that in the plateau regime $D \propto D_{\text{Bohm}}$.

Both TMX and a reactor would lie in the plateau regime (for TMX $v_i \tau_{\text{dr}} \approx 1$, for a reactor $v_i \tau_{\text{dr}} \approx 10^{-3}$), so that an estimate of the lifetime against neo-classical diffusion is much longer than the lifetime against axial loss. Using the above estimates for α in each case, we obtain

$$\tau_{\text{neo plat}} \approx \frac{\tau_{\text{dr}}}{\alpha^2} \begin{cases} 10^6 v_i^{-1} & \text{TMX} \\ 10^7 v_i^{-1} & \text{Reactor.} \end{cases} \quad (23)$$

These are to be compared to lifetimes of 100 and 50 collision times for TMX and a reactor, respectively.

IV Resonant Diffusion

The foregoing neo-classical transport estimates were based on the assumption that the axial bounce frequency was sufficiently larger than the drift frequency to make J a good invariant. In fact, under the most common tandem mirror conditions, this turns out not to be the case. The ratio of these frequencies may be estimated as

$$\frac{\omega_{\text{dr}}}{\omega_{\text{Bounce}}} \approx \frac{v_{\perp}^2 L_c}{\Omega r_c^2 v_{\parallel}} \approx \frac{r_c L_c}{r_c^2} \quad (24)$$

where we neglect the pitch angle dependency and ρ_c is the Larmor radius. For both TMX and thermal ions in the LLL preliminary reactor design, this ratio of lengths is about unity. Under conditions that this ratio of frequencies equals or exceeds unity, there occurs a resonant diffusion which can lead to a substantially enhanced radial transport.

Note that, roughly speaking, this resonant transport occurs for those particles having high azimuthal drifts, whereas the largest neo-classical transport occurred for those having a small drift velocity over a limited range of μ . Both types of transport may be occurring in the same machine in different regions of phase space, but the resonant diffusion will always dominate.

To describe particle motion under conditions of fast drift, define

$$\Delta\theta \approx \frac{\rho_c L_c}{r_c^2} \quad (25)$$

as the angular deflection of a particle in one axial transit, and

$$\Delta r \approx \frac{L_{tr}}{v_{||}} \frac{v_{\perp}^2 r_c}{L_{tr}^2} \cos 2\theta \quad (26)$$

$$\approx \frac{\rho_c r_c}{L_{tr}} \cos 2\theta$$

as the radial displacement suffered upon reflection from one transition region. Δr for an 80 eV deuterium ion in TMX is shown in Fig. 1. On the $n+1$ -st completes bounce period measured from the midplane,

$$\theta_{n+1} = \theta_n + 2\Delta\theta \quad (27)$$

$$r_{n+1} = r_n + |\Delta r| \left[\cos 2\left(\theta_n + \frac{1}{2} \Delta\theta\right) - \cos 2\left(\theta_n + \frac{3}{2} \Delta\theta\right) \right].$$

When $\Delta\theta$ is small, there is a lowest order cancellation to $r_{n+1} - r_n$ which is the physical basis for omnigenity. When, locally in radius, $\Delta\theta(r)$ satisfies

$$\Delta\theta(r_k) = \kappa\pi/2$$

where $\kappa = 1, 3, 5, \dots$, successive changes in r will accumulate. Around such points, particles will drift on new banana-like orbits, called "resonant bananas". Expanding for radii near r_κ , in the continuum limit one obtains equations describing this resonant banana motion

$$\dot{r} = (-1)^{\frac{\kappa+1}{2}} \frac{2|\Delta r|}{\tau_{\parallel}} \sin 2\theta \quad (28)$$

$$\dot{\theta} = \frac{r-r_\kappa}{\tau_{\parallel}} \theta'(r_\kappa)$$

where $\theta' \equiv \partial\theta/\partial r|_{r_\kappa}$ and $\tau_{\parallel} = L_c/v_{\parallel}$. From Eqs (28) we obtain estimates for resonant banana width Δr_{res} and period τ_{res}

$$\Delta r_{\text{res}} \approx \left| \frac{r}{\theta'} \right|^{1/2} \quad (29)$$

$$\tau_{\text{res}} \approx \left| \frac{1}{\Delta r \theta'} \right|^{1/2}, \quad (30)$$

and the range of μ of particles lying on a resonance is given by

$$\frac{\Delta \mu}{\mu} \approx \frac{\Delta r_{\text{res}}}{r_c}. \quad (31)$$

Depending on the degree of collisionality, motion on these resonant bananas will lead to different types of diffusion, obtained by arguments analogous to the neo-classical case described earlier. When

$$\nu_{\text{eff}} \approx \nu_i \left(\frac{\Delta \mu}{\mu} \right)^2 < \tau_{\text{res}}^{-1}, \quad (32)$$

diffusion is of the resonant banana type; estimating $\epsilon' \approx r_c^{-1}$, this gives

$$D_{\text{res ban}} \approx (r_c \Delta r) \left(\nu_i \frac{\mu^2}{\Delta \mu^2} \right) \left(\frac{\Delta \mu}{\mu} \right) \approx r_c^2 \left(\frac{\Delta r}{r_c} \right)^{1/2} \nu_i. \quad (33)$$

When $\nu_{\text{eff}} \tau_{\text{res}} > 1$, a wider range of μ can scatter into a resonant banana in one τ_{res} determined by

$$\left(\frac{\Delta u}{u}\right)^2 \frac{1}{v_i} \approx \tau_{res} \approx \tau_{\parallel} \frac{u}{\Delta u}, \quad (34)$$

or

$$\left(\frac{\Delta u}{u}\right)^3 \approx \begin{cases} v_i \tau_{\parallel} & \text{if } v_i \tau_{\parallel} < 1 \\ 1 & \text{if } v_i \tau_{\parallel} > 1 \end{cases}, \quad (35)$$

where again we estimate $r_{c0} \approx 1$. With the radial velocity given by $\Delta r / \tau_{\parallel}$ and a correlation time by $v_i^{-1} (\Delta u / u)^2$, the diffusion becomes

$$D_{res} \approx \left(\frac{\Delta r}{\tau_{\parallel}}\right)^2 \frac{\Delta u^2}{v_i u^2} \frac{\Delta u}{u} \\ \approx \frac{(\Delta r)^2}{\tau_{\parallel}} \begin{cases} 1 & \text{if } v_i \tau_{\parallel} < 1 \\ \frac{1}{v_i \tau_{\parallel}} & \text{if } v_i \tau_{\parallel} > 1 \end{cases}. \quad (36)$$

As for the neoclassical transport, machines of interest all lie in the plateau regime. This diffusion, equal to step sizes of order Δr at a rate $\tau_{\parallel}^{-1}$, can be orders of magnitude faster than neoclassical.

Another regime not covered by the above analysis occurs when resonances "overlap", i.e., at a point in phase space a particle appears to lie on more than one banana. More precisely, this condition prevails when

$$\Delta r \frac{\partial \Delta \theta}{\partial r} > 1.$$

In this case the diffusion coefficient has the same scaling as that in the plateau regime independent of the collisionality, even in the complete absence of collisions. Because randomizing events between end reflections will not alter this result, there is no decrease in D for $v_i \tau_{\parallel} > 1$ as there is for neoclassical transport. Because this regime of stochastic diffusion usually requires $\omega_{dr} \tau_{\parallel} \gg 1$, we do not expect it to occur in machines of interest.

To compare lifetimes for TMX and a reactor against resonance transport as compared to end-loss, note that

$$\nu_i \tau_{\parallel} \approx \begin{cases} 1 & \text{TMX} \\ 10^{-3} & \text{Reactor} \end{cases}$$

(placing both in the plateau regime), so that the lifetime with respect to resonance diffusion becomes

$$\tau_{\text{res diff}} \approx \tau_{\parallel} \left(\frac{L_{\text{tr}}}{\rho_c} \right)^2$$

$$\approx \begin{cases} \nu_i^{-1} (L_{\text{tr}}/\rho_c)^2 & \text{TMX} \\ 10^{-3} \nu_i^{-1} (L_{\text{tr}}/\rho_c)^2 & \text{Reactor} \end{cases}$$

Both of these lifetimes are larger than the respective end-loss lifetimes for thermal ions by about a factor five. The small margin of safety for these lifetime estimates underlines both the potential severity of this transport and the need for careful calculations. The confinement of energetic alphas based on this result is much worse. In the next section are briefly discussed a number of ideas to reduce this transport should it prove necessary.

V. Some Means for Reducing Transport

The neoclassical and resonant transport in a tandem mirror system described above results from non-axisymmetry. In an axisymmetric system, both of the radial displacements Δr and δr vanish identically, as would like-particle diffusion. Generally speaking, reduction of transport in a non-symmetric system can be affected by reducing the degree of non-symmetry. However, because non-symmetry was originally introduced for reasons of MHD stability, each idea for reduction of transport must be examined for its MHD implications. To do so in a quantitative way is beyond the scope of this report. Here, the descriptions will be strictly qualitative, and the list is, presented partially as a call for further investigation.

Even within the currently conceived tandem mirror field designs, resonant transport of thermal ions can be reduced by increasing the transition

length L_{tr} , for which the lifetime increases as the square. Improvements by factors of two to four are conceivable. The loss of energetic alpha particles would not be sufficiently improved by this simple alteration.

Another approach applicable to current field designs makes the parameter $L_c v_c / r_c^2 \ll 1$ so that resonance cannot occur for the bulk particles. Small $v_{||}$ particles would always be resonant; but, if this condition were satisfied only over a small region of phase space, transport of the bulk would again depend upon collisions.

Several methods for stabilizing a purely symmetric system deserve to be studied. The first is some form of line-tying, either passive or active using feedback control. There is an important point to be made here: as the length of a solenoid is increased for given plug parameters, the system end-loss current increases while the bad curvature drive for instability remains constant. This is not the case for single cell mirrors, for which line tying has received most of its attention. This decoupling of the end-current from the drive does not hold for rotation, for which the drive also increases with length. However, in an axisymmetric system the radial electric field may be substantially reduced. As for any line-tying scheme, care must be taken in balancing the competing demands of good electrical contact and thermal insulation of the electrons from the end walls.

A second means for stabilizing axisymmetric systems are ECRH heated rings. These have proved very successful in stabilizing the ELMO single cell mirrors and ELMO Bumpy Torus. A disadvantage for tandem mirrors is the very high plug magnetic fields which are presently conceived; however, these designs do have that advantage of having $\omega_{pe} < \Omega_e$ as required for ECRH. Increased overall MHD stability would be improved by employing two sets of rings per end; one in the plug midplane as in ELMO, and another held in a local minimum of B at the end of the solenoid just before the increase of B on passing into the plugs. In this way, all regions of bad curvature are held by rings. The latter set would be used only if required, of course; it is suggested solely because of uncertainty as to the efficacy of the single ring in holding the solenoid. Because of known properties of these rings, such a stabilized axisymmetric tandem mirror might be constrained in some operating parameters, for example B, but the possible advantages warrant its consideration.

Short of going to a completely axisymmetric system, gains can be realized by modifications of a quadrupole-plug stabilized system. For

example in our conventional designs, if mod-B is increased in an axisymmetric step by a factor of two or so before the non-axisymmetric part of the transition, roughly one-half of the alpha particles would be confined in an axisymmetric magnetic well for their scattering time, which is enough to lose their energy to electrons. By adjusting this internal mirror ratio, these alphas can be caused to be lost at an energy of roughly $15 T_e$ so that they do not heat thermal ions in the solenoid.

A more complicated magnetic design involves plugs formed of double wells, the inner one being axisymmetric and having the density required for potential confinement of the solenoid, the outer one being minimum-B and having sufficient pressure to anchor the entire system to flute-averaged stability. The inner plug would probably have to have a very small mirror ratio.

Finally, in reapplying the arguments given earlier, it can be seen that radial displacements δr and δR are reduced for plugs having higher-than-quadrupole stabilizing bars. This follows from the fact that the internal flux surfaces of such systems have near-axisymmetry. By the same token, good magnetic curvature is realized only for the outer lines, and it is on these lines that pressure gradients must occur. Some MHD and magnetic design questions are currently being considered by the author.

VI. Summary and Conclusions

With a rather large margin of safety, conventional neoclassical drift transport does not appear to compete with axial end-loss from a tandem mirror system. Because of the absence of neoclassical banana orbits for energetic alphas, the same can be said for them.

The lack of omnigenity of electrons in the plugs suggests their good radial mixing, the concomitant uniformity of T_e , and a Boltzmann ambipolar potential distribution. The degree to which this mixing is complete in the sense that small volume plugs can totally control the radial potential profile of a large volume solenoid remains to be calculated. For now the limit of constant T_e and Boltzmann ϕ must be considered only as a limited case. The resultant radial field is important for considerations both of transport and rotation.

For systems with $L_C \rho_C / r_C^2 \geq 1$, resonant transport of thermal ions appears much more competitive with axial loss in current tandem mirror designs. Based upon the rough estimates, radial transport of thermal ions

is less than axial loss by a factor of several and can be reduced somewhat by magnetic design. Systems with $L_C \rho_C / r_C^2 < 1$ will again have transport of the bulk which is collision frequency dependent.

Resonant transport of energetic alphas is more severe because of the energy dependency of Δr . By field shaping in the solenoid, this might be eliminated for alphas with larger pitch angles, but loss of those penetrating deep into the transition regions seems unavoidable.

The feature of no resonant transport makes an axisymmetric system even more attractive than the obvious reason of simple magnets would suggest. The issue is MHD stability, and this area deserves further study. Because the conjectured Boltzmann radial potential was a feature of non-axisymmetry, the radial electric field in axisymmetric systems would presumably be much less, with a corresponding reduced drive for rotational instability.

Plugs of higher order multipole fields are attractive from the point of view of transport, but stability requires further consideration.

In terms of the parameter L_p / r^2 , TMX has the same value as the LLL preliminary reactor design, so that sensitivity to this important parameter can be studied.

References

1. D. D. Ryutov and G. V. Stupakov; Pis'ma Zh. Eksp. Teor 26, 186 (1977); trans. in Sov. Phys. JETP Lett 26, 177 (1977).
2. D. D. Ryutov and G. V. Stupakov; Fiz. Plazmy 4, 501 (1978), Lawrence Livermore Laboratory trans. 02544.
3. D. D. Ryutov and G. V. Stupakov, Dokl. Akad. Nauk, 240, (1978); Lawrence Livermore Laboratory trans. 02548.
4. M. E. Kishinevshij, et al., in Plasma Physics and Controlled Nuclear Fusion Research (7th IAEA Int. Conf., Innsbruck, 1978), paper CN-37-J3, to be published.
5. F. H. Coensgen, TMX Major Project Proposal, Lawrence Livermore Laboratory Report, Prop-148 (1977).
6. R. W. Moir, et al., Preliminary Design Study of the Tandem Mirror Reactor (TMR); Lawrence Livermore Laboratory Report UCRL-52302 (1977).
7. L. S. Hall and B. McNamara, Phys. Fluid 18, 553 (1975).

Figure Captions

- Fig. 1 δr and Δr vs. r for sample TMX vacuum magnetic field.
For Δr the ion is 80 eV deuterium.
- Fig. 2 Example of a set of banana orbits embedded in a family
of axis encircling drift orbits.
- Fig. 3 Qualitative form of neoclassical diffusion showing banana,
plateau, and collisional regimes.

Fig. 1

Fig. 2

Fig. 3-