SRI International

APR 01 1993
A-2

DOE/PC/91053--T3

DE93 013609

SRI Report No. 2847-15 - Quarterly Report No. 5
4 February 1993

FUNDAMENTAL STUDIES OF COAL LIQUEFACTION

David S. Ross
Chemistry Laboratory

Covering the Period October 1, 1992 to January 1, 1993

Prepared for:

U. S. Department of Energy
Pittsburgh Energy Technology Center
P.O. Box 10940, MS 921-165
Pittsburgh, PA 15236

Contract No. DE-AC22-91PC91053

Approved:

David M. Golden
Vice President
Physical Sciences Division

INTRODUCTION

Our diamond cell is now working properly following replacement of a cracked diamond. Mr. Sjoerd Hoogwater, who joined our program for 1 year from Prof. Johannes Penninger laboratories at Eindhoven University of Technology, has now returned to Holland. Dr. Indira Jayaweera, a post doctoral worker in our laboratory with considerable experience in studies in hydrothermal media, has replaced him.

Following repair of the cell and just before his departure, Mr. Hoogwater conducted a large number of experiments, including runs with Wyodak and Illinois No. 6 coal with both tetralin and water as media. He also conducted run control runs in argon with both coals and no liquid medium. We are carrying out analyses of our video taped record, and are at present focusing on the argon and tetralin work with the Illinois coal. Those data are discussed here.

CURRENT RESULTS

Background and Some Comments on Procedures

In earlier accounts we discussed the effects of heating coal samples composed primarily of small particles, generally no larger than around $10\mu\text{m}$ in extent. In those cases in both water and tetralin the major action appeared to be Brownian motion, most active between 150° and 350°C . However it was difficult to develop much more information because the particles were so small.

In the more recent runs we have attempted to isolate larger particles in the observation area of the cell. The procedure is a tedious one since the finer particles are considerably more mobile than the heavier ones. However with experience we have learned to manipulate coal particles in the $50\text{-}100\mu\text{m}$ range into the observation area. We have found not unexpectedly that Brownian motion is not very apparent with such large particles. The major occurrence in these cases is the direct observation of the swelling and shrinking of the particles, the size changes dependent upon the coal and the medium.

The heating rate used in the work discussed here is $10^\circ\text{C}/\text{min}$. For our analysis we have viewed the video tapes of the action in the cell through a rectangular grid fixed to the monitor screen. The particles are presumed to be elliptical, so the area occupied by the particle is simply $A = [\pi/4] \times l \times w$, where l and w are the dimensions of the rectangle circumscribing the particle. The changes in size of selected particles are then recorded as a function of time and temperature.

Illinois No. 6 - Argon

The results for the control run for a single particle are shown in Figure 1. The analysis was made difficult by the deposition of a tar-like material on the window with increasing temperature which increasingly blocked the image. That feature of the experiment will be discussed below.

The figure, which includes other thermolytic data for Illinois No. 6 coal, shows that the particle was stable in size to around 440°C. At that point it contracted very rapidly to about half its size over the next 15°. Also shown are data for the same coal from SRI's field ionization mass spectrometer (FIMS), in which case a sample of the coal is heated at 2.5°C/min from ambient to 500°C and the effluent is swept into mass spectrometer. The evolution of the coal volatiles is reflected by accumulated ion count with increasing temperature, and is presented in the figure as the fraction of total volatiles. As the figure shows, the abrupt increase in volatility corresponds very closely to the equally abrupt reduction in particle size.

The weight average molecular weight profile from the FIMS work is also in the figure, and shows that the coal experiences a simple loss of volatiles of increasing molecular weight up to around 440°C. At that point pyrolysis and fragmentation come about, reflected in the quick reversal and decline in molecular weight. The high rate of volatility and shrinkage are clearly associated with this second stage in the pyrolytic process. These results are further in line with the account of Eser et al. (1991) who found in studies of Illinois No. 6 coal in a high pressure microdilatometer that the coal contracted by about 40% in the same region of temperature at heating rates of around 20°C/min. At higher heating rates the contraction was followed by a swelling.

We have included in the figure results for Illinois No. 6 coal of Solomon, et al. (1988). They studied the thermolytically generated volatiles produced at heating rates of 30°C/min using FTIR techniques, and the water and tar data are presented in the figure. Their data show that the rates of water and tar production become significant at around the 440°C point where we see so much activity both visually and via FIMS. They continue to be produced well above 500°C.

Finally, in our cell work we observed the deposition of tar in the diamond windows starting at around 350°C. It was an unexpected observation; it preceded any apparent contraction of the coal, and continued slowly, occluding the view as it continued. It is reminiscent of the production of tar from Illinois No. 6 coal we discussed in our last quarterly report in work with the front lighted cell and with water present. We expect to reproduce this phenomenon in repeat runs and evaluate its importance to the pyrolytic process in coals.

Illinois No. 6 Coal - Tetralin

The results of work in tetralin are shown in Figure 2, which includes the profile for the argon work. The figure shows that in tetralin the coal swells beginning at around 250°C, with the swelling leveling off at 300°C. At about 390°C the coal particles appear to shrink, although it is likely we are observing the results of dissolution of portions as a result of reduction by tetralin. Qualitatively the result is not unanticipated, since we would expect that tetralin would bring about reductive conversion of the coal to soluble fragments.

However, the starting temperature of the decline is very surprising, coming about 60°C below the point at which coal undergoes pyrolytic degradation. This result is not consistent with the view that liquefaction takes place through the action of an H-donor (Don-H) quenching the free radicals generated as a result of the pyrolytic homolysis of bonds in the coal:

In this sequence the coal pyrolysis in the first step initiates the process, and the donor is the passive partner. The results shown in Figure 2, however, suggest that the tetralin may be active in initiating the conversion process. It is clearly involved in reaction with some portion of the coal structure at temperatures considerably below where significant bond scission in the coal occurs.

FUTURE WORK

With the replacement of the cracked diamond, we have added some other improvements into our cell. Accordingly our recent images are somewhat more clear than earlier. Further we have just begun processing the television images of the processes in the cell using a Macintosh personal computer equipped with a Video Spigot card which allows us to transfer the images to disc. With Adobe Photoshop we will then improve and manipulate the images, and extract portions of them for side-by-side comparison of changes in particle morphology during heating in various media. We expect now to be able to present pictures of changes in the particles in our forthcoming reports.*

* We expect that this approach in addition will allow us to present our visual results properly in these reports. At present it is difficult to integrate our images with the reports at a reasonable cost with satisfactory quality.

With the improved cell we are now conducting a full series of studies with both Illinois No. 6 and Wyodak coals in n-undecane and water. We have been in contact with our research partner, Dr. Charles Scouten at Amoco Oil Co., and in accord with our statement of work he is preparing coal samples impregnated with various catalysts. We expect then to be able to observe directly the effects the catalysts bring about in the behavior of the coals in the various media.

REFERENCES

- S. Eser, R. Jenkins, G. Wei, H. Schobert, and J. Joseph, 1991. *Fuel*, **70** (12), 1445-1455.
- P. Solomon and M. Serio, 1988. "Fundamental Studies of Retrograde Reactions in Direct Liquefaction," First Quarterly Report from Advance Fuel Research, DOE Contract No. DE-AC22-88PC88814. See also: P. Solomon, M. Serio, G. Despande, and E. Kroo, 1990. *Fuel*, **1** (4), 42-54.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Figure 1. Temperature profiles of the thermal treatment Illinois No. 6 coal in vacuum or inert gases. The points represent the size change for a single particle as observed in the cell. The curves display FIMS data from this work, and *tar* and *water* profiles from the work of Solomon, et al. (1988).

Figure 2. Temperature profiles of the thermal treatment Illinois No. 6 coal with tetralin and argon media.

END

DATE
FILMED
6/11/93