225
8-1-78

lh. 307

MASTER

**ACID RAIN RESEARCH PROGRAM
ANNUAL PROGRESS REPORT
JULY 1976 THROUGH SEPTEMBER 1977**

December 1977

**LAND AND FRESHWATER ENVIRONMENTAL SCIENCES GROUP
DEPARTMENT OF ENERGY AND ENVIRONMENT**

**BROOKHAVEN NATIONAL LABORATORY
ASSOCIATED UNIVERSITIES, INC.**

UNDER CONTRACT NO. EY-76-C-02-0016 WITH THE

UNITED STATES DEPARTMENT OF ENERGY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

ACID RAIN RESEARCH PROGRAM ANNUAL PROGRESS REPORT JULY 1976 THROUGH SEPTEMBER 1977

Lance S. Evans,* A.J. Francis,** and Gilbert S. Raynor

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

MASTER

December 1977

*Laboratory of Plant Morphogenesis, Manhattan College, Bronx, N.Y. and Consultant,
Department of Energy and Environment, Brookhaven National Laboratory.

**Department of Nuclear Energy.

LAND AND FRESHWATER ENVIRONMENTAL SCIENCES GROUP
DEPARTMENT OF ENERGY AND ENVIRONMENT

BROOKHAVEN NATIONAL LABORATORY
UPTON, NEW YORK 11973

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy (DOE), nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America
Available from

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

Price: Printed Copy \$6.50; Microfiche \$3.00

May 1978

360 copies

CONTENTS

	Page
Abstract	iv

Part I

EFFECTS OF ACIDIC CONDITIONS ON GROWTH, DEVELOPMENT AND PRODUCTIVITY OF FOREST AND CROP PLANTS

Lance S. Evans, Director	1
Study A - Foliar Response of Six Clones of Hybrid Poplar to Simulated Acid Rain	2
Study B - Effect of Buffered Solutions and Various Anions on Vegetative and Sexual Development in Gametophytes of <u>Pteridium aquilinum</u>	29
Study C - Effects of Acidic Conditions on the Sequence of Required Steps for Symbiotic Nitrogen Fixation of <u>Pisum sativum</u>	49

Part II

EFFECT OF ACID RAIN ON SOIL MICROBIOLOGICAL ACTIVITIES

A. J. Francis, Director	73
-------------------------	----

Part III

SEQUENTIAL PRECIPITATION CHEMISTRY AND ALLIED METEOROLOGICAL DATA

G. Raynor, Director	97
---------------------	----

ABSTRACT

Experiments were carried out and chemical aspects of ambient precipitation were determined using a sequential precipitation collector for the period July 1976 through September 1977. A related report provides experimental details.

In experiments with plants, experiments were aimed to document 1) the foliar response of six clones of hybrid poplar to simulated acid rain, 2) effects of buffered solutions and various anions on vegetative and sexual development of gametophytes of Pteridium aquilinum and the acid-sensitive steps of symbiotic nitrogen fixation of the garden pea (Pisum sativum). After five 6 min daily exposures to simulated rain of pH 2.7, up to 10% of the leaf area of some poplar clones was injured. Lesions developed mostly near stomata and vascular tissue as shown with other plant species. Acidic solutions have a marked effect on sperm motility and fertilization (sexual reproduction) of bracken fern. Since sexual reproduction of ferns is very sensitive to mildly acidic conditions under laboratory conditions, experiments are planned to view the response of sexual stages of other plant species. Nodulation and symbiotic nitrogen fixation in Pisum is very sensitive to nutrient solution acidity. Specific isolates of Rhizobium bacteria are used and the medium pH can be maintained rigidly.

In experiments to determine the effects of excess acidity on soil microbiological processes, the rate of denitrification may be slowed so drastically that increases of N₂O in the atmosphere may result with a subsequent reduction in soil nitrogen levels.

Part I

EFFECTS OF ACIDIC CONDITIONS ON GROWTH,
DEVELOPMENT AND PRODUCTIVITY OF FOREST AND CROP PLANTS

Lance S. Evans, Director

Study A

Foliar Response of Six Clones of Hybrid
Poplar to Simulated Acid Rain

ABSTRACT

After exposure to simulated acid rain from pH levels of 2.7 to 3.4, lesions of several types were produced on foliage of six clones of Populus hybrids. The types of adaxial leaf surface lesions were observed at low magnification, by scanning electron microscopy, and via leaf histology. On two clones, galls resulting from hyperplasia and hypertrophy of parenchyma cells predominated. In contrast, two clones exhibited neither hyperplasia nor hypertrophy after exposure to simulated acid rain. In these clones the upper epidermis, palisade parenchyma, and sponge parenchyma were injured in succession. In a third set of two clones, hyperplasia and hypertrophy were present in areas between injured and apparently uninjured tissues. In general, percent leaf area with lesions and percent leaves injured were similar among all six clones at all pH levels tested. At pH 2.7 up to 10% of the leaf area was injured after 5 daily exposures of 6 min each. Injury to about 1.0% at pH 3.4. Lesions developed mostly near stomata and vascular tissues and occurred most frequently on leaves just prior to maximum leaf enlargement. Younger and older leaves were less affected. The results support the hypothesis that the adaxial leaf surface is the most affected surface of plant foliage after exposure to simulated acid rain.

Previous experiments with simulated acid rain [4,6,8,14] or assessments of injury in the environment that result from exposure to acid rain [1,2,13] have been made. There is need for standardization of procedures to determine levels of pollutants that influence forest plant growth, development, and productivity.

The experiments outlined herein are focused upon the effects of simulated acid rain on six hybrid clones of Populus, specifically: (i) percent of leaves injured, (ii) percent of leaf area with leaf lesions, (iii) a visual description of lesion development at low magnifications and with scanning electron microscopy, and (iv) a histological description of lesion development. Some lesions produced by simulated acid rain resemble galls with abnormal cell enlargement and cell proliferation such as those produced by microorganisms and insects. A description of events during lesion and gall formations is presented.

MATERIALS AND METHODS

Stem cuttings of six clones of hybrid Poplar were obtained as a gift from Drs. Keith Jensen and Leon Dochinger as follows:

No. 8-hybrid of P. nigra L. X P. laurifolia Ledeb.,

No. 43-hybrid of P. maximowiczii Henry X P. berolinensis Dipp.,

No. 44-hybrid of P. maximowiczii Henry X P. berolinensis Dipp.

No. 201-hybrid of P. deltoides Bartr. X P. trichocarpa
Torr. and Gray.,

No. 211-hybrid of P. deltoides Bartr. X P. trichocarpa
Torr. and Gray.,

No. 327-hybrid of P. balsamifera L. cv. candicans
X P. berolinensis.

Basal ends of stem cuttings were dipped in an auxin compound (Hormodin 1, Merck Co., Rahway, N. J. 07065, 0.1% indole-3-butyric acid in talc) and placed in a soil mix in a greenhouse with activated charcoal air filters for 12 weeks. Plants were subsequently transferred to a controlled environmental chamber with activated charcoal filters 1 wk prior to experimentation. Plants were kept at 24 C, 40-50% relative humidity, with an 18-hr light period. Light intensity was 10,760 lux at plant height. Plants were fertilized once weekly. Before initiation of rain applications, plants were fitted with a plastic hood around the base of each plant to cover the soil. During experiments plants were irrigated under the plastic hoods only.

Experimental plants were sprayed in a fully enclosed chamber (3.0 x 1.5 x 1.3 meters) with light provided. Plants were placed on a turntable (1 meter diameter) which revolved at 2.5 rpm. Simulated rain was distributed from a set of nozzles 1 meter above plant height. The nozzle system consisted of two different nozzle types [5] mounted in a downward direction directly over plants to give rain and fog drop sizes simultaneously. A vibrator assured uniform delivery of drops over plant foliage.

In rain simulations, only one rain application of 6 min with an effective rainfall rate of 7.2 mm/hr was given daily. A rain solution [4] was created to simulate the chemistry of rain of northeastern United States. The pH of the rain was adjusted with 1 N H_2SO_4 or 1 N NaOH to obtain desired levels. Control plants were sprayed with a rain of pH 5.7.

Histology -- Samples were fixed in cold acrolein for 4 hrs. After fixation, samples were dehydrated through a series of cold alcohol solutions [7]. Tissues were embedded and sectioned at 20-30 μm . Sections were stained according to Shellhorn and Hull [12].

Scanning Electron Microscopy -- Leaf samples for scanning electron microscopy (SEM) were fixed in acrolein and dehydrated in ethanol. Samples were critical point dried [4]. A layer of carbon or silver was applied to each sample in a vacuum evaporator with a rotating and precessing stage. The microscope (Model 700, Materials Analysis Co., Palo Alto, Ca.) was operated at 5 kv or 10 kv.

Densities of Trichomes and Stomata -- Densities of trichomes and stomata were determined from positive leaf impressions by the silicone rubber technique of Sampson [11]. A total of eight areas (314 mm^2) were sampled randomly on each of four leaf replicas from each clone. No estimate of stomatal functional capacity was obtained.

RESULTS

The foliar response of clones of hybrid poplar was quantified by measuring lesion diameters of less than 0.25 mm, 0.25 to 1.00 mm, 1.00 to 2.00 mm, and greater than 2.00 mm, respectively [4].

After five exposures to simulated acid rain in which one rain application of 6 min was administered daily at five pH levels, a variety of lesion types was exhibited on all six clones (Table 1). No lesions were observed on leaves exposed to pH 5.7 rain (the theoretical control pH of unpolluted water in equilibrium with ambient concentrations, ≈ 300 ppm, of carbon dioxide). However, several clones exhibited some small (< 1.0 mm) lesions after exposure to simulated rain of pH 3.4. The percent leaf area with lesions and percent leaves injured both increased as the simulated rain pH decreased. At pH 2.7, the percent of leaf area injured was 4 to 6% of the leaf surface and 60 to 91% of the leaves exhibited lesions.

Leaf surface photographs (Figs. 1 through 6) document lesion development on various clones. Small lesions (< 0.25 mm) (Figs. 2,3 and 5) may develop after one 6-min exposure to simulated rain at either pH 2.7 or 2.9. These lesions initially exhibited chlorosis followed by necrosis. Chlorotic lesions exhibited a slight depression in the leaf surface (Fig. 2). Larger lesions (0.25 - 1.0 mm) exhibited necrotic tissues usually in a circular pattern. In some clones (No. 8, No. 207, No. 211 and No. 327) a red coloration developed in peripheral areas of large (< 1.0 mm) lesions, although colorations were not present on either clones No. 43 or No. 44 (Fig. 4). Fig. 5 shows a large lesion with several smaller lesions (< 0.25 mm) denoted as either chlorotic (chl) or necrotic (nec). The large lesion is about 1 mm in diameter and exhibits a darkened area between healthy and necrotic tissues. A large lesion (2.0 mm in length) (Fig. 6) encompasses several small veins that retain chlorophyll longer than interveinal areas.

Scanning electron microscopy of leaf surface also was performed on healthy and uninjured leaf samples of various clones of poplar (Fig. 7 through 16). Only clone No. 44 exhibited a low density of trichomes on both abaxial and adaxial leaf surfaces. Lesions were initially localized near stomata (Fig. 9 and 10) and were generally localized near vascular tissues. More internal leaf tissues were probably injured as estimated by depression depth (Fig. 11).

The responses of clones No. 8 and No. 327 to simulated acid rain was markedly different from the responses of the other clones. In these two clones, exposure to simulated acid rain produced gall-like outgrowths even after one or two, 6 minute exposures daily (Fig. 13 through 16). The epidermis of the gall and the tissues that surround the gall collapse (Fig. 17).

The collapse of epidermal cells around the gall form a depression on the leaf surface. This depression was present in interveinal tissues but not if galls were near or within vascular tissues (Fig. 15 and 16). On occasion, galls were observed on clones No. 207 and No. 211 but none were present on clones No. 43 and No. 44.

In a later stage of gall formation, cells of both the palisade and spongy parenchyma areas exhibited hypertrophy (abnormal cell enlargement) and hyperplasia (abnormal cell proliferation) in response to simulated acid rain after collapse of the adaxial epidermis. These cell abnormalities generally cause elevations on both adaxial and abaxial leaf surfaces (Fig. 18).

The responses of clones No. 207 and No. 211 were intermediate between (a) those of both No. 43 and No. 44 and (b) those of clones No. 8 and No. 327. Clones No. 8 and No. 327 exhibited galls with abnormal cell enlargement and cell proliferation, but clones No. 43 and No. 44 did not produce either abnormal cell enlargement or abnormal cell proliferation (Fig. 21-24).

In small lesions on leaves of clones No. 207 and No. 211 galls were infrequent. Usually the adaxial epidermis collapsed initially with some injury to the palisade parenchyma. As lesions enlarged after additional exposures to simulated acid rain, palisade cells in the lesion usually collapsed completely. However, cells of the spongy parenchyma of lesions (0.25 - 1.00 mm) exhibited hyperplasia that usually preceded the presence of hyperplasia and hypertrophy in spongy and palisade parenchyma in zones adjacent to the lesion (Fig. 21 and 22). This latter zone of hyperplasia and hypertrophy produced a ridge of tissue surrounding the lesion (Fig. 23 and 24).

All six clones of poplar exhibited a similar response at each pH level of simulated acid rain as indicated by the low standard error of

each mean (Table 2). In general, the percentage of leaf surface area with lesions increased with an increase in number of exposures at each pH level. In addition, after each rain event, the percent of total leaf area with lesions increased with an increase in acidity of simulated rain.

Since most lesions were associated with stomata, experiments were focused on the density of stomata on abaxial and adaxial leaf surfaces. Density of stomata on abaxial leaf surfaces was about three times that of adaxial surfaces (Table 3). The number of lesions per unit area on adaxial leaf surfaces after four consecutive daily simulated rain events (Table 4) was compared with the density of stomata per unit area on adaxial leaf surface (Table 5). The density of stomata and density of lesions per unit adaxial leaf area were not correlated. For example, clone No. 8 had the highest density of stomata but had the lowest density of lesions while clone No. 43 ranked second in both categories. As a result, no correlation is present between the two densities.

Experiments were focused on the response of density of lesions with stage or leaf development on clone No. 8. The increase in leaf length occurred during a rain period of 3 days in which plants were exposed to one, 6-minute simulated rain event daily for 3 consecutive days. The number of lesions per unit area was determined 1 day after the last rain application. The greatest densities of lesions occurred in leaf numbers 5,6,7 and 8 which represented leaves that had just become fully expanded or were undergoing leaf expansion (Fig. 25).

DISCUSSION

The experimental approach was aimed at a prediction of the acute and long term responses that foliage of a variety of plant species may have after exposure to acidic precipitation in the environment. The experimental

results with leaves of Phaseolus vulgaris, Helianthus annuus, and hybrid clones of Populus suggest that the density of lesions caused by acid rain is low during early leaf development. The density of lesions remains low before the stage of most rapid leaf enlargement. In poplar, the density of lesions per unit area is greatest during maximum leaf expansion. In older leaves the density of lesions declines. A similar decline in sensitivity to lesion development with foliage maturation on fronds of bracken fern had been noted (unpublished results). The presence of acidic rain during May, June and July may have the most detrimental effects on deciduous tree and crop plant foliage.

Simulated acid rain produced more lesions per unit area at a particular pH on clones of hybrid poplar compared with foliage of Phaseolus vulgaris and Helianthus annuus [4]. The sensitivities of these plants are markedly higher than that exhibited by needles of coniferous trees. Coniferous needles usually do not exhibit visual lesions above pH 2.5 (Hindawi, personal communication). Thus the foliage of predominantly deciduous forests of the Northeastern United States may be more sensitive compared with the coniferous forests of Scandinavia at similar acidity levels (3, 9, 10).

The features of deciduous tree and crop plant leaves that determine maximum sensitivity to simulated acid rain remain unknown. Among the clones of hybrid poplar used, the density of lesions was not correlated with density of stomata per unit leaf area, even though most lesions develop adjacent to or at stomata. In addition, density of lesions was not correlated with densities of stomata or trichomes even though 95% of all lesions occur at these sites in Phaseolus vulgaris [5]. Some other factors must be involved in predisposing these leaf structures and, possibly other areas of the leaf, since

maximum sensitivity to simulated acid rain is correlated with maximum leaf expansion rate in all three plants tested. This developmental stage of maximum sensitivity may be correlated with a temporary decrease in cuticle thickness in some leaf areas as a result of differential leaf growth, or may involve synthesis of different types of cutin or cell wall components for short time periods.

In two clones, No. 8 and No. 327, injured areas produced mostly galls after one or two exposures to simulated acid rain. In contrast, leaf cells of clones No. 43 and No. 44 underwent no hyperplasia and hypertrophy after exposure to simulated acid rain. In two clones, No. 207 and No. 211, hyperplasia and hypertrophy were limited to peripheral areas surrounding lesions. Both galls and the more limited response in peripheral areas of lesions were responses to exposure to simulated acid rain.

In accordance with lesion development on other plants, lesions of the hybrid clones of poplar exhibited an increase in penetration and lateral enlargement with number of exposures to simulated acid rain. With continued exposures to rain solutions of low pH, the number of lesions increased but lesion enlargement over the 6 to 7 day period of exposure reached a final diameter of several millimeters only. As a result, at any one time there was a steady state in the progression of small to large lesion diameters.

Our results coupled with results of other investigators demonstrate that simulated sulfate acid rain preferentially affects the leaf indumentum near trichomes, stomata, and vascular tissues. After initial injury, subsequent lesion development shows a progressive deterioration of the adaxial leaf surface to more abaxial structures. Because lesion frequency was not correlated with either stomatal or trichome frequency, the results suggest that epidermal cell morphology during development predisposes leaves to

lesion development. These results with simulated rain suggest that differential sensitivity of plant taxa in nature to acid rain may be influenced chiefly by characteristics of the leaf indumentum.

REFERENCES

1. Abrahamsen, G., K. Bjor, R. Horntvedt, and B. Tveite, Effects of acid precipitation on coniferous forests, in *Impact of Acid Precipitation on Forest and Freshwater Ecosystems in Norway*, ed. by F. H. Braekke, Oslo, Norway, Reclamo Printers, 1976, pp. 37-63.
2. Beamish, R. J., W. L. Lockhart, J. C. Van Loon, and H. H. Harvey, Long-term acidification of a lake and resulting effects on fishes, *Ambio* 4, pp. 98-102 (1975).
3. Cogbill, C. V., The history and character of acid precipitation in Eastern North America, in *Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem*, ed. by L. S. Dochinger and T. A. Selinga, USDA Forest Service General Technical Report NE-23, 1976, pp. 363-370.
4. Evans, L. S., N. Gmur, and F. Da Costa, Leaf surface and histological perturbations of leaves of Phaseolus vulgaris and Helianthus annuus after exposure to simulated acid rain, *Amer. J. Bot.* 64, pp. 903-913 (1977).
5. Evans, L. S., N. Gmur, and J. J. Kelsch, Perturbations of upper leaf surface structures by acid rain, *J. Environmental and Experimental Bot.* (in press).
6. Fairfax, J. A. and N. W. Lepp, Effect of simulated "acid rain" on cation loss from leaves, *Nature* 255, pp. 324-325 (1975).
7. Feder, N. and T. P. O'Brien, Plant microtechnique: Some principles and new methods, *Amer. J. Bot.* 55, pp. 123-142 (1968).
8. Ferenbaugh, R. W., Effects of simulated acid rain on Phaseolus vulgaris L. (Fabaceae), *Amer. J. Bot.* 63, pp. 283-288 (1976).
9. Likens, G., F. H. Bormann, and N. M. Johnson, Acid rain, *Environment* 14, pp. 33-44 (1972).
10. Nordø, J., Long range transport of air pollutants in Europe and acid precipitation in Norway, in *Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem*, ed. by L. S. Dochinger and T. A. Selinga, USDA Forest Service General Technical Report NE-23, 1976, pp. 87-103.
11. Sampson, J., A method of replicating dry or moist surfaces for examination by light microscopy, *Nature* 191, pp. 932-933 (1961).
12. Shellhorn, S. J. and H. M. Hull, A six-dye staining schedule for sections of mesquite and other desert plants, *Stain Technol.* 36, pp. 69-71 (1961).

13. Wiklander, L., The acidification of soil by acid precipitation, *Grundförbättring r*, pp. 155-164 (1973-74).
14. Wood, T. and F. H. Bormann, Increases in foliar leaching caused by acidification of an artificial mist, *Ambio* 4, pp. 169-171 (1974).

Table 1. Responses of six clones of poplar after exposure to simulated rain^a

Clone No.	pH of simulated rain	Percent of leaf area with various lesion diameters				Percent of leaves injured ^c
		<0.25mm	0.25-1.00mm	1.00-2.00mm	>2.00mm	
8	2.7	0.04 ^b	0.41	0.96	1.95	91
	2.9	0.02	0.20	0.16	0.65	56
	3.1	0.02	0.08	0.16	0.00	40
	3.4	0.01	0.04	0.00	0.00	1
	5.7	0.00	0.00	0.00	0.00	0
43	2.7	0.07	0.49	1.54	4.12	69
	2.9	0.18	1.14	1.54	4.12	61
	3.1	0.07	0.49	0.51	0.00	38
	3.4	0.04	0.25	0.00	0.00	8
	5.7	0.00	0.00	0.00	0.00	0
44	2.7	0.08	0.68	0.79	2.12	71
	2.9	0.03	0.22	0.52	2.12	56
	3.1	0.03	0.22	0.24	0.00	32
	3.4	0.02	0.07	0.00	0.00	5
	5.7	0.00	0.00	0.00	0.00	0

207	2.7	0.06	0.40	0.76	2.00	80
	2.9	0.02	0.16	0.15	0.67	66
	3.1	0.01	0.04	0.00	0.00	22
	3.4	0.01	0.04	0.00	0.00	6
	5.7	0.00	0.00	0.00	0.00	0
211	2.7	0.05	0.30	1.18	2.12	89
	2.9	0.04	0.30	0.47	1.06	57
	3.1	0.04	0.15	0.00	0.00	26
	3.4	0.00	0.00	0.00	0.00	0
	5.7	0.00	0.00	0.00	0.00	00
327	2.7	0.04	0.58	1.41	3.27	60
	2.9	0.04	0.22	0.28	1.64	58
	3.1	0.02	0.15	0.28	0.00	30
	3.4	0.02	0.07	0.00	0.00	7
	5.7	0.00	0.00	0.00	0.00	0

^aPlants were exposed to five exposures of simulated acid rain. One rain application was given daily for 6 min only. The plants were rated 1 day after the fifth exposure.

^bValues represent estimates of mean percent leaf area with lesions from all leaves of three plants in four separate experiments.

^cValues represent estimate of percent of leaves injured from three plants in four separate experiments.

Table 2. Injury on leaf surfaces of six clones of poplar at various simulated rain pH levels^a

pH of simulated rain	Percent leaf surface with lesions with days after beginning of exposure:			
	1	2	3	4
5.7	0 ± 0 ^b	0 ± 0	0 ± 0	0 ± 0
3.1	0.7 ± 0.0	1.8 ± 0.3	1.8 ± 0.2	2.2 ± 0.2
2.9	3.8 ± 0.2	3.9 ± 0.3	4.1 ± 0.2	4.6 ± 0.4
2.7	3.2 ± 0.4	3.5 ± 0.2	4.1 ± 0.2	4.6 ± 0.2
2.5	3.9 ± 0.2	4.4 ± 0.2	5.3 ± 0.4	5.9 ± 0.3
2.3	5.0 ± 0.3	5.1 ± 0.2	7.7 ± 0.6	8.4 ± 0.4

^aTwo plants of each clone were observed for each of six clones. Plants were exposed to one-6 minute exposure of simulated rain per day at each pH level. Plant injury was assayed 1 day after each rain application just prior to another application.

^bMean and standard error of the mean, respectively.

Table 3. Frequency of stomata on adaxial and abaxial leaf surfaces
of Populus sp. hybrids

Clone No.	adaxial surface (stomata/mm ²)	abaxial surface (stomata/mm ²)
8	1.07 ± 0.19 ^a	0.73 ± 0.14
43	0.71 ± 0.04	2.36 ± 0.18
44	0.51 ± 0.04	2.92 ± 0.17
207	0.41 ± 0.05	1.27 ± 0.06
211	0.54 ± 0.03	0.95 ± 0.04
327	0.41 ± 0.02	1.57 ± 0.14

^aEach figure represents the average and standard error of the mean from four impressions on each leaf surface from each clone. Eight areas (314 mm²) on each impression were scored.

Table 4. Number of lesions per unit area of the adaxial leaf surface of the hybrid clones of poplar^a

Clone no.	No. of lesions per 314 mm ²
8	14.9 ± 1.9 ^b
43	20.6 ± 2.6
44	21.2 ± 1.5
207	15.2 ± 3.4
211	15.9 ± 1.6
327	19.3 ± 2.8

^aPlants were exposed to one, 6 min simulated rain event daily at pH 2.7 for four consecutive days. Plants were evaluated 1 day after the last rain event. Four areas (each 314 mm²) on five leaves from five plants were evaluated per clone. Four consecutive leaves with the most lesions were selected from each plant.

^bEach figure represents the mean and standard error of the mean.

Table 5. Relationship between densities of stomata and lesions on
adaxial leaf surfaces among clones of hybrid poplar.

Clone No.	No. of stomata per 314 mm ² ^a	No. of lesions per 314 mm ² ^b	Ratio stomata: Lesions
8	336	14.9	22.6:1
43	223	20.6	10.8:1
44	160	21.2	7.6:1
207	129	15.2	8.5:1
211	170	15.9	10.7:1
327	129	19.3	6.7:1

^aData taken from Table 3.

^bData taken from Table 4.

Figure 1-6.

Figure 1-6. Photographs of leaf surfaces of several leaves of Poplar clone No. 43 that exhibit several stages of injury after exposure to simulated acid rain.

(1) Unexposed surface of a leaf. Note sharp contrast between veinal and interveinal areas.

(2) Leaf surface of clone No. 207 exposed to several rain events. A large lesion (0.25-1.00mm) (B) adjacent to several smaller lesions (< 0.25mm) (A). The light areas (*) represent initial injury symptoms. Initial lesions appear only slightly chlorotic. Note darkened areas (red colorations) between necrotic and healthy tissue of the larger lesion (B).

(3) Enlargement of several lesions in Figure 2. A larger lesion (0.25-1.00mm) (B) is shown with two lesions (< 0.25mm) (A). The darkened area between necrotic and healthy tissues of the (B) lesion (0.25-1.00mm) represents a red coloration that occurred in clones No. 8, No. 207, No. 211, and No. 327 but not in clones No. 43 and No. 44. Red pigmentation was present in tissues that exhibited hyperplasia and hypertrophy.

(4) An elongated lesion (0.25-1.00mm) (B*) occurred near a vein adjacent to a more circular lesion (0.25-1.00mm) (B) in clone No. 43. A small lesion is also present near the B* lesion. Note no darkened (red coloration) areas border necrotic and healthy tissues.

(5) A large lesion (1.00-2.00mm) (C) is present near vascular tissue in clone No. 211. Several smaller lesions (< 0.25) are shown. The smaller lesions exhibit only chlorosis (chl) while other more advanced lesions exhibit necrosis (nec.). Red colorations are also present in large lesions also.

(6) The large lesion (> 2.00mm) encompasses a large area in clone No. 207 including several veins as well as interveinal tissue. In large lesions, small veins retain chlorophyll longer than interveinal areas. Large lesions were also characterized by a very thin lamina.

Figure 7-12.

Figure 7-12. Scanning electron micrographs of the adaxial leaf surface of several clones of poplar after exposure to simulated rain at pH 2.7 unless otherwise noted. Note the absence of trichomes on all surfaces shown. VG = vascular tissue; S = stoma (ta).

(7),(8) Leaf surfaces after four applications of simulated rain pH 5.7 only. Photographs show no leaf surface perturbations. Most epidermis cells are fully turgid. Figure 7 view of clone No. 44; Figure 8 view of clone No. 8.

(9) A smaller lesion ($< 0.25\text{mm}$) surrounds a stoma on clone No. 44. About 20-30 epidermal cells within a radius of $100\text{ }\mu\text{m}$ of the stoma are flaccid and surround a depression. Palisade cells may be injured in the centrally depressed area near the stoma.

(10) A small lesion ($< 0.25\text{mm}$) after one 6 min exposure to simulated acid rain to clone No. 43. Note injury to epidermal cells near two stomata and vascular tissues. About six epidermal cells are flaccid in this view. No indication of injury to more internal cells is present.

(11) Surface of leaf with a lesion (B) (0.4mm in length) on clone No. 43. Several stomata are present within the lesion. More internal tissues may be injured as denoted by lesion depression depth.

(12) Low magnification view of a large lesion (1.00mm) on clone No. 44. A clear demarcation between injured and apparently uninjured area is present. A single 6 min exposure of simulated acid rain was given daily for 3 days.

Figure 13-16.

Figure 13-16. Scanning electron micrographs of the adaxial leaf surface of several clones of poplar after exposure to simulated rain at pH 2.7. VT = vascular tissues; G = galls produced by simulated acid rain pH 2.7.

(13) Overview of gall on the leaf surface of clone No. 327. Mound is present near vascular tissues. The gall was present on a leaf twenty-four hr after a single 6-min exposure to pH 2.7.

(14) Enlargement of gall in Figure 13. Photograph illustrates mounded nature of this small gall (< 0.25mm) with completely collapsed epidermis. Epidermal cells immediately adjacent to the mound are collapsed and bounded by apparently normal epidermal cells. The light ovals on the surface of the gall are stomata. The epidermis over the gall is discontinuous.

(15) Two galls on the surface of clone No. 327. One gall encompassed vascular tissue in addition to adjacent interveinal area. This gall exhibits a depression into the leaf tissue on one side.

(16) Enlarged view of a gall on the surface of clone No. 207. Epidermal cells on the gall are no longer distinct but several stomata are present. The gall is small and the tissue is continuous compared with the discontinuous surface of the gall in Figure 14.

Figure 17-24

Figure 17-24. Cross sections of leaves of clones No. 327, No. 207, No. 8 and No. 211 of hybrid poplar that exhibit (1) gall formation and (2) lesion development with associated cell hyperplasia and hypertrophy in response to simulated acid rain exposure of pH 2.7.

(17) Initial gall response of clone No. 207 to simulated acid rain injury. Collapse of epidermal cells with distortion of palisade parenchyma (pp) cells appears darkly stained (*). Elongation of injured palisade parenchyma cells produces a slight elevation of adaxial leaf surface (arrows). No apparent injury to spongy parenchyma (sp) cells occurs.

(18) Initial gall formation of clone No. 8. Cells of both the palisade and spongy parenchyma exhibit early stages of hypertrophy and hyperplasia in response to simulated acid rain exposure. Note the degree of elongation in altered spongy parenchyma cells (arrow). An elevation of both the adaxial and abaxial leaf surface occurs (*). Collapse of the epidermal cells occurs on the adaxial leaf surface. Normal epidermis = N.E.; Vascular tissue = V.T.

(19) Micrograph of a small gall (< 0.25mm) in clone No. 8. The leaf mesophyll consists of altered parenchyma cells in late stages of hypertrophy and hyperplasia. Note the degree of elongation in several mesophyll cells (arrows). A region of hyperplasia activity occurs in the palisade parenchyma (*). No tissue layers are distinguishable. Elevation of the leaf surface with collapse of the epidermal cells is apparent on both adaxial and abaxial leaf surfaces.

(20) Micrograph of a gall (0.25-1.00mm) in clone No. 327. Collapse of epidermal and palisade parenchyma cells occurs on the adaxial leaf surface (*). Zones of hyperplastic cell activity surround the elevated leaf surfaces (A). Huge elongated cells lie in the central zone of cell hypertrophy (B). Normal tissue = N.

(21) Cell hypertrophy in a small lesion (0.25-1.00mm) of clone No. 211. Malformations of both epidermal and palisade parenchyma cells occur in the lesion (A). Spongy parenchyma cells below the lesion exhibit cell hypertrophy (B). Cell enlargement of the spongy parenchyma cells in the area of the lesion (C).

(22) Higher magnification of cell hypertrophy surrounding a small lesion (0.25-1.00mm) of clone No. 211. Palisade and spongy parenchyma cells exhibit the hypertrophic activity responsible for the elevation of the leaf surface (arrows). Cell types are indistinguishable in the zone of hypertrophy. Epidermal and palisade parenchyma cells are destroyed in the lesion.

(23) Cell hypertrophy and hyperplasia surrounding a large lesion (1.00-2.00mm) of clone No. 207. Collapse of the epidermal, palisade and spongy parenchyma cells within the lesion results in a depression of the leaf surface (A). Mesophyll tissue surrounding the lesion is subject to cell hypertrophy and hyperplasia which results in an elevation of the leaf surface on both the adaxial and abaxial sides (B).

(24) Enlargement of Figure 23 that exhibits the zone of hyperplasia and hypertrophy surrounding a lesion (1.00-2.00mm). Note the proliferation of parenchyma cells (hyperplasia) which appears as a darkly stained band adjacent to the lesion (arrow). Hypertrophic cells lie to the left of this zone of hyperplasia. Only remnants of parenchyma cell walls are evident in the lesion proper.

Figure 25. Leaf length, increase in leaf length, and number of lesions per unit area on leaves of poplar clone No. 8. Increase in leaf length occurred during the rain period of 3 days. The number of lesions per unit area was measured one day after the last of 3, 6 min daily rain events. Measurements of the number of lesions were made on four areas per leaf. Characters represent mean and standard error of the mean from ten plants. Leaves were numbered from oldest (Number 10) to youngest (Number 1).

Study B

Effect of Buffered Solutions and Various Anions on Vegetative and Sexual Development in Gametophytes of Pteridium aquilinum

ABSTRACT

Experiments were performed to determine the effects of chloride, nitrate, and sulfate in buffered solutions on sperm motility and fertilization in gametophytes of Pteridium aquilinum. Buffered solutions with various anions simulated exposures to acidic precipitation up to 3.5 hr. The presence of each anion decreased both sperm motility and fertilization. Addition of 43.3 μM sulfate (equivalent to the sulfate concentration in rain of northeastern United States) decreased sperm motility about 60% compared with plants exposed to buffers without sulfate. Additions of 86.6 μM and 173.2 μM sulfate gave values approximately 45% and 36% of values of plants unexposed to sulfate. The reduction in sperm motility during sulfate exposure was independent of buffer pH. Additions of chloride and nitrate at pH 4.8 gave sperm motility values similar to buffers with sulfate. However, at pH 5.8, chloride ions were less harmful than nitrate or sulfate. Additions of anions had a detrimental effect upon fertilization or production of sporophyte plants. In all experiments the presence of chloride and nitrate alone or in combination with sulfate gave reductions in fertilization similar to that of sulfate alone at equivalent concentrations. The presence of these three anions exhibited additive effects with or without low pH to reduce both sperm motility and fertilization in gametophytes of Pteridium aquilinum.

INTRODUCTION

Acidic precipitation has become an important environmental problem in Northern Europe and Eastern North America the past decade that influences both forest and freshwater lake environments (Jensen and Snekvik, 1972; Likens, Bormann, and Johnson, 1972). Presently, this acidification has been attributed to the conversion of the oxides of sulfur and nitrogen to their strong acids in the upper atmosphere (Brosset, 1973; Nordø, 1976).

Recent experimental results suggest that both sperm and the process of fertilization in the fern Pteridium aquilinum (Bracken fern) were inhibited when exposed to buffered solutions below pH 5.8. Addition of sulfate ions increased the inhibition (Evans and Bozzone, 1977). The above experiments were performed to determine the response of sexual development in this species. These parameters could be tested effectively and could be used as an indicator of the effects of acid precipitation with additions of sulfate on fertilization of plants of forests in the northeastern United States.

Experiments presented here explore (a) the effectiveness of multiple inoculations on fertilization of gametophytes, (b) effect of various concentrations of sulfate, nitrate, and chloride on sperm motility at various pH levels, (c) effect of various concentrations of sulfate, nitrate and chloride on fertilization of gametophytes at various pH levels, and (d) the effects of various concentrations of sulfate, nitrate and chloride anions in combinations on fertilization of gametophytes of Pteridium aquilinum.

MATERIALS AND METHODS

The materials and methods used were reported previously (Evans and Bozzone, 1977). Spores were harvested from plants at Wave Hill Preserve, 675 West 252nd St., Riverdale, N. Y. during August 1976. Under aseptic

conditions, spores of Pteridium aquilinum were washed with aerosol OT and then with a 15% solution of fresh Clorox bleach that contained 0.80% sodium hypochlorite. Spores were sterilized for 20 seconds and then added to 50 ml of water. Spores were placed on either (a) Moore's medium with 0.9% agar or (b) sterile vermiculite with Moore's medium added in liquid form (Moore, 1903). Cultures were placed under lighted conditions (200 foot candles) at 20 C throughout the experiments (Näf, 1958).

Effect on Sperm Motility

To view the duration of sperm motility at various pH levels, spores were inoculated on Moore's medium with agar and allowed to grow for about 21 days. Sperm were obtained when prothalli were removed from culture flasks and were placed in a small petri dish. Specific amounts of glass distilled water were added so antheridia would release sperm into the water. Maximum concentrations of sperm for experiments were obtained 12 minutes after water addition. After 12 minutes, citrate-phosphate buffer (Jensen, 1962) stock solutions were added to the water to obtain a specific buffered pH level (0.01M). After the buffer stock solution was mixed with the water, 2 or 3 drops of the buffered sperm suspension were placed in the center of a ring of methyl cellulose on a microscope slide and covered with a cover slip.

Each sperm suspension was observed under phase contrast optics 2-4, 5-7, and 8-10 min. after addition of buffer stock solution. All sperm observed in each 2 minute interval were scored as either motile or non-motile dependent upon the presence or absence of flagellar movements respectively. The number of sperm counted in the two minute period varied from a minimum of 150 to a maximum 250. The methyl cellulose only reduced the velocity of

flagellar movement and sperm motility so individual sperm were much more stationary. Because slides were moved frequently and rapidly distances of about 200 μm between views, the probability of scoring the same sperm twice was minimized during any two minute interval.

Fertilization Experiments

To determine degree of fertilization at a single pH level, about 2 mm³ (Näf, 1958) spored were inoculated onto sterile vermiculite with Moore's medium in liquid form. Petri dishes were reinoculated 6 days later with additional spores to insure an abundance of sperm at time of treatment. Twenty-eight days after the first inoculation, citrate-phosphate buffers of pH 2.2, 3.2, 4.2, 5.2 and 5.8 (control buffer) were added to vermiculite medium in order to immerse gametophytes for 3.5 hours only. The buffer was then removed and a buffer of pH 5.8 was added to the dishes in the same manner. After several minutes, this control pH buffer was also removed. Dishes were then returned to culture conditions for a 14 day period after buffer treatments for development of sporophytes. Almost all fertilization was restricted to the 3.5 hour exposure to buffer solutions. Some cultures had no buffer treatments whatsoever. These cultures were used to measure the degree of spontaneous fertilization when no buffers were added.

The acids, H_2SO_4 , HNO_3 and HCl at anion concentrations of 43.3, 86.6 and 173.2 μM were added to all buffer solutions. The acids were used to produce the anion levels. The amounts of acids added did not alter the pH of the buffers either initially or at the end of the buffer exposure period.

RESULTS

Under the experimental conditions used, antheridia developed on most gametophytes 14 days after inoculation but live sperm were not present until day 19. On day 21 a maximum number of sperm were obtained in our experimental

conditions. On this day only archegonial initials were present. Mature archegonia were present 28 days after inoculation (Evans and Bozzone, 1977). Because of this lack of synchronous maturation of male and female sexual organs on an individual prothallus were provided two inoculations, 6 days apart. In this manner prothalli from the first inoculation would be 28-days-old and have mature archegonia while prothalli from the second inoculum would be 22-days-old and have mature antheridia.

The effectiveness of single and double inoculations were rated by the number of cordate gametophytes with sporophytes (Table 1). Since about 90% of the gametophytes with apical notches from the first inoculation had mature archegonia (Evans and Bozzone, 1977), and about 50% of the gametophytes in the petri dishes resulted from the first inoculation, about 45 to 50% of the notched gametophytes present could be fertilized. Since about 52% of the notched gametophytes had sporophytes, most of the notched gametophytes from the first inoculation would be fertilized.

If plants from the first and second inoculations were not present simultaneously then fertilization was reduced drastically. In the last column of Table 1, 43% of all gametophytes had sporophytes after two inoculations. However, if only 22-day-old plants were assayed, only 9% had sporophytes. In addition, if only 28-day-old plants were present, then 19% of the gametophytes had sporophytes. In both single inoculation cases, low percentages of fertilization from both single inoculation systems are added together, the resultant sum, 28.2% (19.1% plus 9.1%) is below the 43% which occurred when both inoculations were together. It should be mentioned that a slight increase in the proportion of prothalli with mature archegonia could have been present. However, this increased proportion in the combination inoculum treatment was

probably due to an excess of sperm from the second inoculum. These results suggest that the sperm contributed by the second inoculation might be instrumental in fertilizing cordate gametophytes of the first inoculum. This movement of sperm from gametophytes of the second inoculum of gametophytes of the first inoculum could occur because ample amounts of buffer were added.

Experiments were performed to determine the inhibitory effects of various concentrations of sulfate ions on sperm motility, 2-4 minutes (Fig. 1) and 5-7 minutes (Fig. 2) after exposure. After a 2-4 minute exposure, addition of sulfate ions markedly depressed motility especially at pH levels of 5.8 and 5.6 (Table 2). In general, addition of 43.3 μM sulfate decreased motility 35 to 45% at the 2-4 minute exposure time. At higher sulfate concentrations, the percent motile sperm decreased and was somewhat independent of pH levels. If the results are viewed in terms of decrease in pH with respect to pH 5.8 at all sulfate levels, motility was depressed more on a percentage basis at the lower sulfate concentrations (0 and 43.3 μM sulfate) compared with higher sulfate concentrations (86.6 μM and 173.2 μM). However, this differential response to pH at the various sulfate levels becomes more equal at lower pH levels (Table 2).

When sperm are exposed to buffer solutions for 5-7 minutes, the differential effects noted at the higher sulfate levels and lower pH levels are not as marked as at 2-4 minutes. No marked differences in motility occurred below pH 5.6 and the responses at 86.6 μM and 173.2 μM sulfate concentrations were similar at all pH levels. The only significant difference occurred at pH 5.8 at lower sulfate concentrations. In general, as the duration of exposure to lower pH levels and higher sulfate concentrations increased, longevity of motility decreased.

Since the acidity of rain in the northeastern United States may be coincident with an increase in nitrate and/or chloride concentrations, experiments were focused upon the response of sperm motility at 2-4 minutes after addition of nitrate and chloride to buffers (Table 3). The anion concentrations of 43.3, 86.6 and 173.2 μM were chosen so a comparison could be made with the response to sulfate additions (Table 2). Percent sperm with flagellar movements was higher with addition of chloride compared with the more similar responses obtained when either sulfate or nitrate were present at pH 5.8. However, at pH 4.8, the response to chloride was similar to that of nitrate. Motility may be more depressed with additions of sulfate than with additions of chloride or nitrate at pH 4.8 (Table 3). However, since sperm motility was very low at this pH level, tests of significance among the responses to the various anions were not performed.

Degree of fertilization was also used to measure the response of gametophytes to various anions. The response of gametophyte fertilization to various sulfate concentrations at various pH levels is shown in Table 4. At all pH levels used from 4.2 to 5.8, additions of sulfate produced a similar decrease in fertilization. At 43.3, 86.6 and 173.2 μM , the percent cordate gametophytes with sporophytes averaged 86.5%, 74.0% and 61.5% of dishes with no sulfate added. In a similar fashion, among the various sulfate concentrations used, the effect of pH was consistent. At 5.2, 4.8 and 4.2, fertilization was 92.3%, 75.8% and 64.5% of that obtained at pH 5.8 (experimental control).

At pH's 5.8 and 4.2, the effect of various anions on fertilization was measured. At both pH levels used, addition of chloride, nitrate, and sulfate produced similar reductions in fertilization at all anion concentrations observed (Table 5). No significant, consistent differences were observed among the three anions.

In a similar manner, the effects of two anions in combination (Table 6) were similar to the responses of twice the concentration of each anion individually (Table 5). For example, the average of the three combination treatments at 43.3 μM (35.2, 32.3 and 31.1) (Table 6) is 32.9 which is similar to the average of values at 86.6 μM , 31.5 (31.5, 30.1 and 32.8) (Table 5). The same comparison of the average of the three combination treatments at 86.6 μM (28.7) (Table 6) were similar to the average of single anions at 173.2 μM (28.6) (Table 5). These results suggest that the three anions, nitrate, chloride and sulfate have similar effects on fertilization singly or in combination at the pH levels tested.

The above results suggest that sperm motility and fertilization are sensitive to acidic solutions and acidic solutions with anions added. However, it may be questioned if acidic solutions influence development of cordate gametophytes during prothallial growth. Spores were germinated and grown at pH 5.8 for 2 weeks and then treated with buffer at pH 2.2. The pH of the vermiculite was not readjusted and plants were examined for the percentage of cordate gametophytes two weeks after treatment. Under these conditions, $39.6\% \pm 0.6\%$ (mean and standard error of the mean, respectively) of the total gametophytes were cordate. This mean was very close to the control value of 43.0% obtained previously (Evans and Bozzone, 1977) when no acidic buffer was added on day 28. These results suggest that gametophyte development is not as sensitive as fertilization to acidic conditions in Pteridium aquilinum.

DISCUSSION

The results of the present investigation plus previous results (Evans and Bozzone, 1977) suggest that fertilization and particularly, sperm motility, are very sensitive stages in the life cycle of Pteridium aquilinum. Acidic solutions applied to sexually immature prothalli have very limited influence on eventual development of sexual organs. The experimental results of all experiments show that mobility of unicellular sperm cells may be the limiting factor for completion of the life cycle in P. aquilinum.

The lack of a significant reduction in prothallus development under acidic conditions is not surprising because bracken fern inhabits bogs of low pH substrata. However, since gametophytes of P. aquilinum may inhabit acidic substrata does not suggest that all stages of the life cycle or all cells of sporophyte or gametophyte plants can survive under pH levels of bogs. One could speculate that sperm on the surface of gametophytes would be subjected to pH levels more similar to the ambient rain than pH levels of the soil or bog strata. Since the average pH of rain in the northeastern United States is about 4.1 with sulfate concentrations that range from 36 to 557 $\mu\text{eq/l}$ (Jacobson, Heller and Van Leuken, 1976), our experimental results suggest a very significant reduction in sperm motility and subsequent fertilization.

Our results were obtained when gametophytes and sperm were exposed to buffers with added anions. Buffers were utilized instead of a simulated rain solution because the pH of the buffers, although low in ionic strength (0.01 M), was adequate to maintain the appropriate pH. These experimental results show that fern gametophytes represent a plant developmental system that differentiates between the responses to vegetative and reproductive portions of the life cycle. Moreover, the results suggest that only limited fertilization

could occur with ambient levels of hydrogen ions and various anions in rain in the northeastern United States.

In general, the presence of the anions, chloride, nitrate and sulfate decreased both sperm motility and fertilization uniformly. Under all conditions, the effects of sulfate and nitrate were considered identical. In sperm motility experiments, chloride had less detrimental effects than either sulfate or nitrate at low concentrations at pH 5.8. However, when the effects of various anions were observed in fertilization experiments, no differences were considered significant.

The presence of various anions in these developmental systems facilitates further analysis of the relation that sperm motility has to actual fertilization in our experiments. Since there was a consistent decrease in both motility and fertilization with an increase in anion concentrations, this consistent decrease suggests that both techniques give a relative scale to measure sexual development in Pteridium aquilinum.

In procaryotic organisms, motility appears maximal between pH 6.8 and 9.0 (Ogiuti, 1936; Shoesmith, 1960; Adler and Templeton, 1967). As pH decreased from 7.5 to 5.0, flagellar velocity decreased 50% in Bacillus brevis. Hoeningner (1965) demonstrated a decrease in pH produced an increase in "curly" flagella. "Curly" flagella are characterized by a wavelength one-half of the normal wavelength. The presence of curly flagellae decreased motility so that at pH 5.2 motility was restricted to rotational movements only although motility appeared normal at pH 5.6. Below pH 5.0 almost all motility ceased. These results suggest that flagellar motility is very sensitive to pH.

Although assessments of environmental injury have been performed (Abrahamsen, et al., 1976; Wiklander, 1973-1974; Beamish, et al., 1975) and

experiments have been performed to document injury to plants by simulated acid rain (Wood and Bormann, 1974; Fairfax and Lepp, 1975; Ferenbaugh, 1976; Shriner, 1976; Evans, Gmur and Da Costa, 1977) no studies to date have been designed to investigate the ecological relationships between organisms with this environmental stress. In addition, many experiments with aquatic organisms indicate that acidity may result in a lower degree of species diversity in freshwater lakes of North America and Scandanavia (Wright, et al., 1976). One of the most important areas for research upon long-term changes in pH would be unicellular organisms. The experimental results of this paper indicate that fertilization may block completion of the life cycle by preferential injury to unicellular sperm cells. Since most of the organisms on earth are unicellular, and these organisms are necessary for almost all nutrient cycles, close attention should be given to these organisms.

The marked effect of buffered solutions on fertilization of P. aquilinum is similar to the influence that air pollutants exert on lichens. From experimental results of the effects of SO₂ on lichens in chambers of fixed environmental factors, surveys of lichen species in an area can be used as bioindicator of ambient air pollution levels when complex and/or expensive monitoring instrumentation is unavailable (Skye, 1968; Leblanc and De Sloover, 1970, Horntvelt, 1975).

REFERENCES

- Abrahamsen, G., K. Bjor, R. Horntvedt, and B. Tveite, Effects of acid precipitation on coniferous forests, in *Impact of Acid Precipitation on Forest and Freshwater Ecosystems in Norway*, ed. by F. H. Braekke, Oslo, Norway, Reclamo Printers, 1976, pp. 37-63.
- Adler, J. and B. Templeton, The effect on environmental conditions on the motility of Escherichia coli, *J. Gen. Microbiol.* 46, pp. 175-184 (1967).
- Beamish, R. J., W. L. Lockhart, J. C. Van Loon, and H. H. Harvey, Long-term acidification of a lake and resulting effects on fishes, *Ambio* 4, pp. 98-102 (1975).
- Brosset, C., Air-borne acid, *Ambio* 2, pp. 2-7 (1973).
- Evans, L. S. and D. M. Bozzone, Effect of buffered solutions and various anions on vegetative and sexual development in gametophytes of Pteridium aquilinum, *Amer. J. Bot.* 64(7), pp. 897-902 (1977).
- Evans, L. S., N. Gmur, and F. Da Costa, Leaf surface and histological perturbations of leaves of Phaseolus vulgaris and Helianthus annuus after exposure to simulated acid rain, *Amer. J. Bot.* 64(7), pp. 903-913 (1977).
- Fairfax, J. A. and N. W. Lepp, Effect of simulated "acid rain" on cation loss from leaves, *Nature* 255, pp. 324-325 (1975).
- Ferenbaugh, R. W., Effects of simulated acid rain on Phaseolus vulgaris L. (Fabaceae), *Amer. J. Bot.* 63, pp. 283-288 (1976).
- Hoeninger, J. F., Influence of pH on Proteus flagella, *J. Bact.*, 90, pp. 275-277 (1965).
- Horntvedt, R., Epiphytic macrolichens on Scots pine related to air pollution from industry in Odda, Western Norway, *Reports of the Norwegian Forest Research Institute*, 31, pp. 1-12 (1975).
- Jacobson, J. S., L. I. Heller, and P. Van Leuken, Acidic precipitation at a site within the northeastern conurbation, in *Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem*, ed. by L. S. Dochinger and T. A. Selinga, USDA Forest Service General Technical Report NE-23, 1976, pp. 267-279.
- Jensen, K. W. and E. Snekvik, Low pH levels wipe out salmon and trout populations in southernmost Norway, *Ambio* 1, pp. 223-225 (1972).
- Jensen, W. A., *Botanical Histochemistry*, San Francisco, Calif., Freeman & Co., 1962.
- Le Blanc, F. and J. De Sloover, Relation between industrialization and the distribution and growth of Epiphytic lichens and mosses in Montreal, *Can. J. Bot.* 48, pp. 1485-1496 (1970).

- Likens, G., F. H. Bormann, and N. M. Johnson, Acid rain, *Environment* 14, pp. 33-44 (1972).
- Moore, G. T., Methods for growing pure cultures of algae, *J. Appl. Microscopy and Lab Methods* 6, 2309 (1903).
- Näf, U., On the physiology of antheridium formation in the bracken fern Pteridium aquilinum (L) Kuhn, *Physiol. Plant* 11, pp. 728-746 (1958).
- Nordø, J., Long range transport of air pollutants in Europe and acid precipitation in Norway, in *Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem*, ed. by L. S. Dochinger and T. A. Selinga, USDA Forest Service General Technical Report NE-23, 1976, pp. 87-103.
- Ogiuti, K., Untersuchungen über die Geschwindigkeit der Eingenbewegung von Bakterien, *Jap. J. Exp. Med* 14, pp. 19-28 (1936).
- Shoesmith, J. G., The measurement of bacterial motility, *J. Gen. Microbiol* 22, pp. 528-535 (1960).
- Shriner, D. S., Effects of simulated rain acidified with sulfuric acid on host-parasite interactions, in *Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem*, ed. by L. S. Dochinger and T. A. Selinga, USDA Forest Service General Technical Report NE-23, 1976, pp. 919-925.
- Skye, E., Lichens and air pollution: A study of cryptogamic Epiphytes and environment in the Stockholm region, *Acta Phytogeographica Suecia* 52, pp. 1-123 (1968).
- Wiklander, L., The acidification of soil by acid precipitation, *Grundförbättring r*, pp. 155-164 (1973-74).
- Wood, T. and F. H. Bormann, Increases in foliar leaching caused by acidification of an artificial mist, *Ambio* 4, pp. 169-171 (1974).
- Wright, R. F., D. Torstein, E. Gjessing, G. Hendrey, A. Hendriksen, M. Johannesen, and I. Muniz, Impact of acid precipitation on freshwater ecosystems on Norway, in *Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem*, ed. by L. S. Dochinger and T. A. Selinga, USDA Forest Service General Technical Report NE-23, 1976, pp. 459-476.

Table 1

Effectiveness of First and Second Inoculations on Fertilization of Prothalli of P. aquilinum

Inoculation #1 (day 0)	Inoculation #2 (day 6)	Buffer Treatment (day 28)	Cordate gametophytes with sporophytes (percent)	Cordate gametophytes with sporophytes (percent above control)
+ ^a	+	none	8.9 ± 2.9	
+	+	pH 5.8, 3.5 hr.	51.7 ± 3.1	42.8
+	-	none	12.2 ± 1.5	
+	-	pH 5.8, 3.5 hr.	31.3 ± 1.0	19.1
-	+	none	13.3 ± 4.5	
-	+	pH 5.8, 3.5 hr.	22.4 ± 3.1	9.1

^a plus - denotes inoculation was performed, negative - denotes inoculation was omitted

Table 2

Effect of sulfate concentration on sperm motility^a of Pteridium aquilinum, 2-4 min. after exposure to buffered solutions

pH ^b	Sulfate concentration (μ m)			
	0	43.3	86.6	173.2
5.8	68.3 $\pm$ 6.9 ^c	39.5 $\pm$ 3.5 -58.0- ^e	29.3 $\pm$ 2.6 -44.2-	20.6 $\pm$ 1.7 -30.4-
5.6	46.3 $\pm$ 2.7 (64.7) ^d	30.5 $\pm$ 4.5 (77.2) -66.0-	23.7 $\pm$ 3.0 (80.9) -51.2-	17.3 $\pm$ 2.3 (84.0) -37.3-
5.2	28.8 $\pm$ 4.5 (42.3)	18.1 $\pm$ 1.6 (46.0) -62.8-	15.5 $\pm$ 3.5 (52.9) -53.8-	12.0 $\pm$ 1.7 (58.3) -41.7-
4.8	19.0 $\pm$ 1.2 (28.0)	12.3 $\pm$ 1.0 (31.1) -64.7-	7.3 $\pm$ 1.8 (21.9) -38.4-	6.3 $\pm$ 1.1 (30.5) -33.1-

^a Percent sperm with flagellar movements.

^b pH of citrate-phosphate buffer at 0.01M.

^c Values represent mean and standard error of the mean, respectively. Experiments were performed three times and are comparable to previous results (Evans and Bozzone, 1977).

^d Values surrounded by parentheses represent the percent sperm with flagellar movement within each sulfate level compared with pH 5.8 (control).

^e Values within dashes represent the percent sperm with flagellar movement within each pH level compared with the no sulfate control.

Table 3

Effect of chloride and nitrate concentrations in buffered
solutions on sperm motility in Pteridium aquilinum,
2-4 minutes after exposure

pH	Percent sperm with flagellar movement			
	0	Anion concentration (μM)		
		43.3 Chloride nitrate	86.6 Chloride nitrate	173.2 Chloride nitrate
5.8	66.5 $\pm$ 4.8	44.6 $\pm$ 2.1	36.0 $\pm$ 1.7	30.0 $\pm$ 0.7
		39.1 $\pm$ 1.4	24.5 $\pm$ 1.6	16.9 $\pm$ 0.8
4.8	22.4 $\pm$ 1.8	18.1 $\pm$ 1.2	12.7 $\pm$ 1.0	9.3 $\pm$ 0.8
		17.3 $\pm$ 1.3	13.3 $\pm$ 2.3	9.1 $\pm$ 1.2

Table 4

Effect of sulfate concentration in buffered solutions on percent cordate gametophytes with sporophytes^a in Pteridium aquilinum

pH ^c	Sulfate concentration (μ M) ^b			
	0	43.3	86.6	173.2
5.8	50.3 $\pm$ 0.9 ^d	44.5 $\pm$ 0.86	38.4 $\pm$ 3.2	31.3 $\pm$ 1.6
		-88.4- ^f	-76.3-	-62.2-
5.2	46.5 $\pm$ 2.9	39.8 $\pm$ 1.1	34.8 $\pm$ 0.9	30.3 $\pm$ 0.5
	(92.4) ^e	(89.4)	(90.6)	(96.8)
		-85.6-	-74.8-	-65.2-
4.8	38.7 $\pm$ 1.0	31.7 $\pm$ 0.2	28.3 $\pm$ 1.1	25.4 $\pm$ 0.3
	(76.9)	(71.2)	(73.7)	(81.2)
		-81.9-	-73.1-	-65.6-
4.2	34.1 $\pm$ 0.5	30.7 $\pm$ 0.9	24.4 $\pm$ 1.6	18.0 $\pm$ 1.0
	(67.8)	(69.0)	(63.5)	(57.5)
		-90.0-	-71.6-	-52.8-

^a Cordate gametophytes with sporophytes as a percentage of all cordate gametophytes.

^b Concentration of sulfate added as sulfuric acid to buffers of various pH levels. The pH of the buffer was not altered during the 3.5 hr exposure to the buffers at all sulfate levels. The sulfate concentration of 43.3 μ M is equivalent to the concentration of sulfate in rain of New York (see discussion).

^c The pH was maintained with 0.01M citrate-phosphate buffer.

^d Values represent mean and standard error of the mean of at least 4 dishes per treatment. At least 400 prothalli per dish were scored. The experiment was performed twice and gave values comparable with previous results (Evans and Bozzone, 1977).

^e Values surrounded by parentheses represent the percent cordate gametophytes with sporophytes within each sulfate level compared with pH 5.8 (control).

^f Values within dashes represent the percent cordate gametophytes with sporophytes within each pH level compared with the no sulfate controls.

Table 5

Effect of chloride, nitrate, and sulfate in buffered solutions on the percent cordate gametophytes with sporophytes in Pteridium aquilinum^a

		Cordate gametophytes with sporophytes as a percent of all cordate gametophytes			
		<u>Anion concentration (μM)</u>			
pH	0	43.3 ^b Chloride nitrate sulfate	86.6 Chloride nitrate sulfate	173.2 Chloride nitrate sulfate	
5.8	48.4 $\pm$ 2.3	38.3 $\pm$ 0.6	31.5 $\pm$ 1.9	30.8 $\pm$ 2.7	
		36.3 $\pm$ 0.8	30.1 $\pm$ 1.2	25.3 $\pm$ 2.8	
		40.0 $\pm$ 3.7	32.8 $\pm$ 5.5	29.7 $\pm$ 1.1	
4.2	32.2 $\pm$ 1.6	29.0 $\pm$ 0.6	25.2 $\pm$ 1.1	19.4 $\pm$ 1.8	
		26.7 $\pm$ 0.9	23.5 $\pm$ 1.1	17.1 $\pm$ 1.9	
		32.7 $\pm$ 1.8	27.1 $\pm$ 1.0	20.3 $\pm$ 1.7	
No Buffer Added	8.3 $\pm$ 0.7				

^a Gametophytes were exposed to buffered solutions with either HCl, HNO₃ or H₂SO₄.

^b Values represent mean and standard error of the mean of at least 5 dishes per treatment. At least 300 plants were scored per dish.

Table 6

Effect of several anions in combination of buffered solutions
at pH 5.8 on percent cordate gametophytes with
sporophytes in Pteridium aquilinum^a

Cordate gametophytes with sporophytes as a percent of total cordate gametophytes				
Concentration of each anion (μ M)				
Anion Combination	0	43.3	86.6	173.2
None	49.4 $\pm$ 1.7 ^b	-	-	-
Chloride and Nitrate	-	35.2 $\pm$ 0.8	28.2 $\pm$ 2.6	27.7 $\pm$ 2.7
Chloride and Sulfate	-	32.3 $\pm$ 1.7	29.3 $\pm$ 3.2	26.2 $\pm$ 1.1
Nitrate and Sulfate	-	31.1 $\pm$ 0.3	28.6 $\pm$ 2.8	26.5 $\pm$ 1.4
No buffer added	11.0 $\pm$ 0.4	-	-	-

^a Gametophytes were exposed to buffered solutions with combinations of anions.

^b Values represent mean and standard error of the mean of at least 5 dishes per treatment. At least 300 plants were scored per dish.

Figure 1. Sperm with flagellar movement after exposure to buffers of various pH levels for 2-4 minutes. Sperm motility values represent percent sperm with movement of the total observed in the two minute period. Sulfuric acid was added to make the various sulfate concentrations. Additions of sulfuric acid did not alter the pH. Each point represents a mean and standard error of the mean, respectively of at least five trials per experiment. The experiment was performed three times. Values are given in Table 2.

Study C

Effects of Acidic Conditions on the Sequence of Required Steps for Symbiotic Nitrogen Fixation of Pisum sativum

ABSTRACT

The research measured the limiting step(s) in nodulation of Pisum sativum by Rhizobium leguminosarum bacteria for symbiotic nitrogen fixation.

This approach represents an efficient method to measure the effects of energy generation by-products from fossil-fuels on the symbiotic relationship(s) between Rhizobium bacteria and leguminous crop plants important for food production. The program measured the responses of (i) legume root growth and development, (ii) survival and growth rate of inoculum, (iii) degree of nodulation, (iv) rates of nitrogen fixation, and plant biomass when roots of P. sativum were exposed to various pH levels that might result from prolonged exposure to acid rain.

The experimental results show that the most limiting factor of nitrogen fixation is nodule formation and N_2 reduction. Both root and bacterial growth was equal above pH 4.8 but efficient nodulation and nitrogen fixation did not occur until a pH of 5.6 was obtained. The processes required for the symbiotic relationship are about 80 times more sensitive to pH than independent growth and development.

INTRODUCTION

One of the most important problems in the world today is the production of adequate quantities of food concomitant with thrift energy consumption. In crop production, one of the limiting factors is adequate nitrogen fertilizer for cereal grain crops (Quispel, 1974). Under these conditions, supplemental nitrogen fertilizer is necessary to maximize production per unit area of tillable land. However, the availability of nitrogen fertilizers to improve grain yields requires large amounts of energy (Quispel, 1974). As a result, as the demand for food increases so will the demand for nitrogen fertilizers and energy. Unfortunately, the economic conditions of developing nations usually limit use of nitrogen fertilizers.

In contrast to cereal grain crops, legumes require little or no supplemental nitrogen fertilization because they are able to produce sufficient nitrogen derived from a symbiotic relationship with bacteria of the genus Rhizobium. This symbiosis is predicated on the fact that individual bacteria infect roots of legumes and form nodules which a) protect the bacteria from a more harsh soil environment, and b) provide the bacteria with ample carbohydrate. At the same time, the bacteria convert molecular nitrogen from the atmosphere into nitrogen compounds that plants can utilize. Table 1 from Quispel (1974) dramatizes the effect biological nitrogen fixation has on the overall nitrogen budget of the earth.

In addition to the steadily increased use of energy in order to provide adequate commercial nitrogen fertilizers for non-legume crops, by-products of energy production may influence this symbiotic relationship between legumes and Rhizobium species. Air pollutant particulates, acidic precipitation, soil surface run off, and minor anthropogenic sources of energy generation

activities have been shown to distribute high concentrations of sulfates, copper, manganese, nickel, and vanadium as well as hydrogen ions (see Table 2 taken from Heicklen, 1976; Wiklander, 1973-1974). Large accumulations of the above components may influence symbiotic nitrogen fixation relationship of legumes. The topic of this program was centered on a determination the effects of pH on (i) plant root development, (ii) proliferative capacity of a Rhizobium strain that is agronomically important, and (iii) the symbiotic relationship necessary for effective biological nitrogen fixation.

Studies on the effects various soil contaminants may elicit in solid root media are excessively complicated because actual concentrations of constituents in the soil solution per se are unknown. Because concentrations of nutrients were considered unknown in solid root media, hydroponic solutions were perfected (Hoagland and Arnon, 1950). Hydroponic solutions enabled researchers to instantaneously alter the chemical composition of the medium. These hydroponic solutions enabled researchers to effectively evaluate responses of plants to various constituents. Recently, an aeroponics system of growing plants that a) maximizes root growth rates, b) Rhizobium survival, and c) symbiotic nitrogen fixation was refined (Zobel, Del Tredici, and Torrey, 1976). This aeroponic system provides adequate growth in all three areas.

Recently, reviews of the present knowledge of biological nitrogen fixation (Quispel, 1974; Nutman, 1976) and development and function of roots (Torrey and Clarkson, 1975) have been published. However, information that relates concentrations of heavy metals and pH to the various required steps for symbiotic nitrogen fixation are needed.

Different groups of Rhizobium bacteria differ in tolerance to acid and alkaline conditions (Fred, et al., 1932). Rhizobium meliloti (Medicago, Melilotus, Trigonella group), is the most acid sensitive and generally does

not survive pH levels below 5.5. The bacterial symbiont that inhabits clovers, R. trifolii, does not survive pH levels below 4.5. Like R. trifolii, R. leguminosarum (Pea-vegetable group) will tolerate lower pH levels than R. meliloti (Fred, et al., 1932). Inocula for cowpeas (Vigna sinensis) and some other tropical legumes are commonly found in soils below pH 4.5. Growth of many rhizobia may be enhanced in acid soils by the application of lime (CaCO_3) (Vincent and Waters, 1954). However, because of competition between microbes, specific isolates may not be able to survive in native soils if the pH is not adjusted over long time periods (Mulder and Van Veen, 1960).

With regard to bacterial growth and acidic conditions, Rhizobia may be artificially separated into species that produce acids in their growth medium (R. leguminosarum, R. trifolii, R. meliloti, and R. phaseoli) and are generally considered to multiply rapidly. In contrast, other species of Rhizobium (R. lupini and R. japonicum) excrete basic substances in their medium and are considered to multiply at a slower rate than acid producers (Fred, et al., 1932; Norris, 1965). From the above discussion, production of acid is associated with fast growth while slow growth is associated with alkali production. In general, species can be placed in order of increasing acid tolerance: R. meliloti, R. phaseoli, R. trifolii, and R. leguminosarum (Graham and Parker, 1964). It should be noted that fast growing rhizobia have the Entner-Doudoroff, pentose phosphate, and Embden-Meyerhof hydrolytic pathways but slow growing species only have the Entner-Doudoroff pathway (Katznelson and Zagallo, 1957).

Many mineral elements are required by Rhizobium species for adequate growth. Vincent (1974) has recently categorized the mineral nutrient requirements of various species. Iron is probably required for the haemoprotein biosynthesis of leghaemoglobin. Magnesium, iron, and molybdenum are required

for nitrogenase activity in conjunction with ATP binding (Orme-Johnson, et al., 1972). Although Co is necessary for vitamin B12 activity the requirement is fulfilled at 10^{-8} M. Co may also be involved in prevention of nickel and copper toxicities (Lowe, et al., 1960; Lowe and Evans, 1962). Calcium is required at about 0.025 mM and abnormal wall structures of Rhizobium bacteria occur if calcium is below this threshold concentration (Humphrey and Vincent, 1962, 1965). During rhizobial infection of legume roots, calcium is required during the period of maturation of root hairs (Munns, 1970). Availability of calcium before and after the critical time period has no effect on the ability of roots to respond to infection if calcium is absent (Albrecht and Davis, 1929). Magnesium cannot substitute for calcium during this critical period (Lowther and Loneragan, 1968; Munns, 1970).

The growth of Rhizobium bacteria has been stated above. Obviously, if bacterial growth is inhibited completely the probability of effective symbiotic nitrogen fixation is limited. Heavy metals such as Hg, Zn, Ni, Cu have toxic effects at low concentrations (Vincent, 1974). As stated above cobalt tends to negate the toxic levels of nickel and copper (Fred, et al., 1932; Wilson and Reisenauer, 1970). High levels of manganese do not promote rapid bacterial growth and is thought to limit infection and effective nitrogen fixation (Holding and Lowe, 1971), but does not limit plant growth at these levels. The tolerance of plants and bacteria to high manganese levels is varied (Vose and Jones, 1963; Döbereiner, 1966; Sonto and Döbereiner, 1969). Nodulation of alfalfa and of clover was reduced by 50% in the presence of 20 ppm manganese (Kliewer, 1961; Vose and Jones, 1963). It should be reiterated that low acidity in soils limits the availability of molybdenum while manganese and aluminum are more available and uptake of these elements may be enhanced (Jensen, 1947).

Calcium and alkaline ions are also inhibitory at very high concentrations (Fred, et al., 1932). The following compounds are toxic: CaCl_2 , 0.14 M, KCl , 0.3-0.6 M; NaCl , 0.3-0.6 M; LiCl , 0.024 M (Vincent, 1974). NaNO_3 is more toxic at 0.15-0.3 M than Na_2SO_4 , 0.3-1.2 N (Vincent, 1974). The above toxic levels reflect two possibly independent inhibitory conditions; general osmotic effects (salt concentrations) (Bernstein and Ogata, 1966) and the effects of combined forms of nitrogen (Rautenberg and Kuhn, 1864; Munns, 1968a, 1968b, 1968c, 1968d).

Generally, 0.1 mM nitrate or ammonium considerably reduce root production and curling in alfalfa by Rhizobium meliloti (Thornton, 1936; Munns, 1968b). Usually, Rhizobium bacteria are less sensitive to salt than nodulation of host plant roots. For example, R. Meliloti grow in 3% NaCl but roots of alfalfa will not nodulate in 0.6% NaCl (Subba-Rao, et al., 1972).

An interaction between nitrate and copper concentrations with nodulation is present. When the nitrate level was controlled at 7 or 14 ppm nitrogen, nodulation was higher at 0.1 to 5 μM copper than at lower or higher copper levels (Hallsworth, et al., 1964). The exact relationship between nitrate and copper concentrations and various physiological parameters remain obscure, and represent a fertile area for experimentation.

Although vanadium may be a minor contaminant in acidic precipitation, no information is available on its effects on symbiotic nitrogen fixation. Becking (1962) stated that vanadium may substitute for molybdenum in nitrogenase of Azotobacter vinelandii in the absence of molybdenum (McKenna, et al., 1970; Burns, et al., 1971). More recently, Benemann, et al., (1972) observed that vanadium may either form a nitrogenase analogue which is inactive or has a limited effect in substituting for molybdenum. Whether vanadium may

substitute for molybdenum in symbiotic rhizobial relationships is unknown and whether vanadium could competitively inhibit molybdenum binding of nitrogenase at high vanadium concentration is equally intriguing.

Biological parameters measured:

1. root growth and development
 - a. mitotic cycle activity
 - b. terminal root length - length from terminal meristem to first emergent secondary root. This is an indication of the autonomy of the individual root meristem.
2. growth rate of inoculum (increase in Rhizobium bacteria) per unit time by standard bacteriology techniques.
3. degree of nodulation - number of nodules per root segment
4. rates of nitrogen fixation - production of ethylene from acetylene
5. biomass - fresh and dry masses of roots and shoots, separately.

The parameters to be assayed above represent the sequence of steps in effective biological nitrogen fixation. Items with lower numbers were required before items with higher numbers can occur. For example, root growth and growth of inoculum (items 1 and 2) are necessary in a particular milieu before infection (item 3) can occur. Our particular experimental design will determine the limiting stage(s) in the inhibition of nitrogen fixation. Since we view these processes in various concentration gradients in a defined medium, a clearer estimation of the adverse effects of the individual components can be made.

Plants of Pisum sativum (garden pea) are infected by Rhizobium leguminosarum (Vincent, 1974), a fast growing, acid producer isolate (Isolate RP-212-1) of R. leguminosarum obtained from North American Plant

Breeders Association. However, isolated roots of Pisum sativum grow well at pH 4.8 (Van't Hof and Kovacs, 1972; results below).

Choice of appropriate inocula is based on the assumption that investigations with pure strains of very effective isolates are more appropriate than mixed cultures (Munns, 1968a; Francis, personal communications). If mixed cultures are used, only one or two selected isolates may be effective at low elemental concentrations. Such a selection procedure may be of interest eventually but such a selection system would complicate the already involved analyses.

MATERIALS AND METHODS

Seeds of Pisum sativum (the garden pea) var Alaska (Asgrow Seed Co.), were surface sterilized and placed on moistened sterilized vermiculite until seedlings were produced. Seedlings were transferred to aeroponic boxes (Fig. 7 and 2) when roots were approximately 4 or 5 cm in length. The buffered solutions (0.01 M phosphate) buffer were placed in the boxes two or three days prior to availability of plants. This two or three day period was convenient for stabilization of the pH and addition of the chemical constituents to be tested. One day after placement of seedlings into the aeroponic boxes, appropriate nodulating bacteria are added (Rhizobium leguminosarum, RP 212-1). This inoculum is a pure culture and an effective symbiont obtained as a gift from Dr. Thomas Wacek, North American Plant Breeders, Princeton, Ill. On a daily basis, the pH of each box was adjusted at least once, twice daily if necessary. Under these circumstances the pH never varied more than 0.2 units.

On a weekly basis, 4 or 5 seedlings were harvested for symbiotic nitrogen fixation via the acetylene assay. This assay system was performed on 4 or 5 plants per treatment. A maximum of 6 treatments can be performed per day.

Symbiotic nitrogen fixation was assayed one hour after addition of acetylene to sealed reaction vessels. Duplicate gas samples were taken from each reaction vessel.

Other parameters such as mitotic cycle activity, terminal root length, and degree of nodulation as well as fresh and dry weights of roots and shoots were obtained after the acetylene assays.

About 30-36 seedlings were placed in one treatment box and 6 acetylene tests were performed during an experiment.

Root growth and development were assayed via 1) mitotic cycle activity, and 2) terminal root length. Seedling roots for mitotic cycle activity were fixed in ethanol-acetic acid (3:1, v:v) and squashed on microscope slides. The mitotic index (percent of cells in mitosis; Evans and Van't Hof, 1974) was assayed. The results in Fig. 4 show that the mitotic index is a good measure of the proliferation capability of the terminal during the experimental conditions. Measurements of terminal root length were also obtained. Terminal root length is the length from the meristem (root cap) to the first emergent secondary root. If the terminal root is inhibited, this distance will be short; but if the distance is long, then the meristem is less inhibited. Generally, a high mitotic index is associated with a long terminal root length while a low mitotic index is associated with a short terminal root length.

Survival and growth rates of bacteria will be evaluated with standard bacteriological techniques.

RESULTS

Results show the effects of various buffered pH levels on nitrogen fixation in P. sativum. The results shown in Fig. 3 show that the phosphate buffer (0.01 M) used was effective in limiting the pH to within 0.2 units. Generally, the pH remained within 0.1 units of the desired pH.

In tissue culture experiments, root growth was closely correlated with mitotic cycle activity (Fig. 4). Because mitotic and label indices were positively correlated with root growth rates, if either mitotic or label indices were assayed relative root growth rates would be predicted. This correlation is needed because relative root growth rates (mm/hr) in the aeroponic system were sometimes difficult to measure.

Terminal root length (determined to be the length of the root between the terminal meristem and the first secondary root) was measured as a function of pH. If the meristem is inhibited then the terminal root length is short compared with uninhibited meristems. In both Fig. 4 and 5 root growth was maximized at pH levels above 4.8.

Fig. 6 shows experimental results in which bacteria of Rhizobium leguminosarum (Isolate RP-212-1) were grown at various pH levels in citrate-phosphate buffer (0.025 M) (Jensen, 1962). The curve demonstrates that growth rate is inhibited below pH 4.8 but at 4.8 to 6.6 rates were identical.

From the results shown above rapid root growth and development and bacterial growth are similar at pH levels above 4.8. In general, the inhibitory effects on root development and bacterial growth are similar at pH levels below pH 4.8.

Number of nodules and ethylene production (an estimate of nitrogen fixation) per plant are shown in Figures 7 and 8. Below pH 5.6 few nodules and little ethylene production occurs. In general, high initial nitrogen fixation occurred only at pH 5.6, concomitant with a large number of nodules. As the duration of the aeroponic experiments increased, the effectiveness of nodules at pH 5.6 decreased after several weeks. In contrast, in roots exposed to pH 6.6, nodulation and effective nitrogen fixation were delayed until 6 weeks after the

start of the experiments. In this manner initial effective nitrogen fixation was limited to pH 5.6 in these experiments.

Mean dry weights of both roots and shoots at various pH levels is shown in Fig. 9. It is evident that dry weights of roots were similar among pH 5.6 to 4.4 but shoot dry weights were depressed below pH 5.6. In other words, the mass of roots was relatively constant among the pH levels while the inhibitory effect of low pH are not compensated in shoot growth. This effect has been shown previously (Harper and Nicholas, 1976). Representative roots from the experiments are shown in Fig. 10.

REFERENCES

- Albrecht, W. A. and F. L. Davis, *Botan. Gaz.* 88, 310 (1929).
- Becking, H. H., *Plant Soil* 16, 171 (1962).
- Benemann, J. R., C. E. McKenna, R. F. Lie, T. G. Traylor and M. Kamen, *Biochem. Biophys. Acta* 264, 25 (1972).
- Bernstein, L. and G. Ogata, *Agron. J.* 38, 25 (1966).
- Burns, R. C., W. H. Fuchsmann, and R. W. F. Hardy, *Biochem. Biophys. Res. Commun.* 42, 353 (1971).
- Döbereiner, J., *Plant Soil* 24, 153 (1966).
- Evans, L. S. and J. Van't Hof, *Amer. J. Bot.* 61, 1104 (1974).
- Fred, E. B., I. L. Baldwin, and E. McCoy, *Root Nodule Bacteria and Leguminous Plants*, Madison, Wis., Univ. of Wisconsin Press, 1932.
- Graham, P. H. and C. A. Parker, *Plant Soil* 20, 393 (1964).
- Hallsworth, E. G., E. A. N. Greenwood, and M. G. Yates, *Plant Soil* 20, 17 (1964).
- Harper, J. E. and J. C. Nicholas, *Physiol. Plant* 38, 24 (1976).
- Heicklen, J., *Atmospheric Chemistry*, New York, Academic Press, 1976.
- Hoagland, D. R. and D. I. Arnon, *Calif. Agr. Exp. Sta. Circ.* 347, 1 (1950).
- Holding, A. J. and J. F. Lowe, *Plant Soil, Spec. Vol.* 153 (1971).
- Humphrey, B. A. and J. M. Vincent, *J. Gen. Microbiol.* 29, 557 (1962).
- Humphrey, B. A. and J. M. Vincent, *J. Gen. Microbiol.* 41, 109 (1965).
- Jensen, H. L., *Proc. Linnean Soc., N.S. Wales* 72, 265 (1947).
- Jensen, W. A., *Botanical Histochemistry*, San Francisco, W. H. Freeman & Co., 1962.
- Katznelson, H. and A. C. Zagallo, *Can. J. Microbiol.* 3, 879 (1957).
- Kliwer, W. M., The effects and interactions of various combinations of Mo, Al, Mn, lime, and Rhizobium strain on growth, composition, and nodulation of several legumes, Ph.D. thesis, Cornell Univ., 1961.
- Lowe, R. H., L. S. Evans, and Shaukat-Ahmed, *Biochem. Biophys. Res. Comm.* 3, 675 (1960).
- Lowe, R. H. and H. J. Evans, *J. Bacteriol.* 83, 210 (1962).

- Lowther, W. L. and J. F. Loneragan, *Plant Physiol.* 43, 1362 (1968).
- McKenna, C. E., J. R. Benemann, and T. G. Taylor, *Biochem. Biophys. Res. Comm.* 41, 501 (1970).
- Mulder, E. G. and W. L. Van Veen, *Plant Soil* 13, 91 (1960).
- Munns, D. N., *Plant Soil* 28, 129 (1958a); 28, 246 (1968b); 29, 33 (1968c); 29, 257 (1968d).
- Munns, D. N., *Plant Soil* 32, 90 (1970).
- Norris, D. O., *Plant Soil* 22, 143 (1965).
- Nutman, P. S., Editor, *Symbiotic Nitrogen Fixation in Plants. International Biological Programme 7*, Cambridge, England, Cambridge Univ. Press,, 1976.
- Orme-Johnson, W. H., W. D. Hamilton, T. L. Jones, M. Y. W. Tso, R. H. Burris, V. K. Shah, and W. J. Brill, *Proc. Natl. Acad. Sci., U.S.* 69, 1972, p. 3142.
- Quispel, A., General Introduction, in *The Biology of Nitrogen Fixation Frontiers of Biology, North-Holland Research Monographs, Vol. 33*, Amsterdam, North-Holland Publ. Co., 1974, p. 1.
- Rautenberg, F. and G. Kuhn, *J. Landwirt.* 12, 107 (1864).
- Sonto, S. M. and J. Döbereiner, *Pesq. Agropec. Bras.* 4, 129 (1969).
- Subba Rao, N., M. Lakshimi-Kumari, C. S. Sing, and S. P. Magu, *Indian J. Agr. Sci.* 42, 384 (1972).
- Thornton, H. G., *Proc. Roy. Soc. Ser. B.* 119, 474 (1936).
- Torrey, J. G. and D. T. Clarkson, Editors, *The Development and Function of Roots*, New York, Academic Press, 1975.
- Van't Hof, J. and C. J. Kovacs, *Mitotic Cycle Regulation in the Meristem of Cultured Roots: The Principal Control Points Hypothesis*, ed. by M. W. Miller and C. C. Kuehnert, New York, Plenum Press, 1972.
- Vincent, J. M. and L. M. Waters, *Australian J. Agr. Res.* 5, 61 (1954).
- Vincent, J. M., Root-nodule symbioses with Rhizobium, in *The Biology of Nitrogen Fixation*, ed. by A. Quispel, *Frontiers of Biology Vol 33*, ed. by A. Neuberger and E. L. Tatum, Amsterdam, North-Holland Publ. Co., 1974, p. 265.
- Vose, P. B. and D. G. Jones, *Plant Soil* 18, 372 (1963).
- Wiklander, L., *Grundförbättring* 4, 155 (1973-74).
- Wilson, D. O. and H. M. Reisenauer, *Plant Soil* 32, 81 (1970).
- Zobel, R. W., P. Del Tredici, and J. G. Torrey, *Plant Physiol.* 57, 344 (1976).

Table 1

Impact of biological nitrogen fixation as a
portion of the nitrogen balance on earth*

Data for a nitrogen balance on earth
(based on Delwiche 1970; Hardy and Holsten 1972)

	Area ha x 10 ⁶	KgN ₂ fixed per ha x yr	Metric tons per yr x 10 ⁶
Biological fixation			
legumes	250	55-140	14-35
non-legumes	1,015	5	5
rice fields	135	30	4
other soils and vegetations	12,000	25-30	30-95
marine	36,100	0.3-1	10-36
Industrial fixation			30
Atmospheric fixation			7.6
Juvenile addition			0.2
Denitrification			
terrestrial	13,400	3	43
marine	36,100	1	40
Loss to sediments			0.2

* Table 1.2, page 3 of Vincent, 1974

Table 2

Mean and maximum urban particulate concentrations
of the United States by pollutant*

Arithmetic Mean and Maximum Urban Particulate Concentrations in the U.S.,
Biweekly Samplings, 1960-1965

Pollutant	Number of stations	Concentrations ($\mu\text{g}/\text{m}^3$)	
		Arith. average	Maximum
Suspended particulates	291	105	1254
Fractions			
Benzene-soluble organics	218	6.8	_____
Nitrates	96	2.6	39.7
Sulfates	96	10.6	101.2
Ammonium	56	1.3	75.5
Antimony	35	0.001	0.160
Arsenic	133	0.02	_____
Beryllium	100	<0.0005	0.010
Bismuth	35	<0.0005	0.064
Cadmium	35	0.002	0.420
Chromium	103	0.015	0.330
Cobalt	35	<0.0005	0.060
Copper	103	0.09	10.00
Iron	104	1.58	22.00
Lead	104	0.79	8.60
Manganese	103	0.10	9.98
Molybdenum	35	<0.005	0.78
Nickel	103	0.034	0.460
Tin	85	0.02	0.50
Titanium	104	0.04	1.10
Vanadium	99	0.050	2.200
Zinc	99	0.67	58.00
Gross beta radioactivity	323	(0.8 pCi/m ³)	(12.4 pCi/m ³)

* Table IV-36, page 229 of Heichlen (1976)

Figure 1. Aeroponics box set-up with pinto bean plants in an early test. Note the stirrer motor on the wood frame centrally located on the top.

Figure 2. A view of the aeroponics box shown in photograph 1 with one side of the box elevated to show roots of plants in position and the stirrer rod attached to the motor.

Figure 3. Variation in pH level of aerobionics boxes versus duration of the experiment. Each point shows pH level present in each box before pH adjustment to the desired levels of 6.6, 5.6, 5.2, 4.8, 4.4 and 3.8, respectively, daily. The pH was adjusted only if the pH was at least 0.1 units from the desired level. Position A: Plants inserted and first inoculation of *Rhizobium*; 2 ppm N as CaNO_3 added. Position B: Second inoculation of *Rhizobium*. Position C: 8 ppm N as CaNO_3 added. Position D: 10 ppm N as CaNO_3 added. To inoculate each box 30 grams of peat moss with inocula was placed in distilled water. This slurry was agitated for several hours. The peat moss was allowed to settle out and the liquid volume equivalent to 5 grams of peat moss at day 0 and day 10 after the start of the experiment.

Figure 4. Root growth during 3 days of aseptic culture, as well as mitotic and label indices of excised roots exposed to nutrient solutions for 3 days versus pH of medium. The cell cycle parameters of mitotic and label indices closely resembled the pattern of root growth under the experimental conditions. These correlations suggest that a single parameter would indicate the status of the other parameters.

Figure 5. The length of 25 lateral roots and 5 tap roots per pH is plotted with pH of buffer. Plants 29- and 43-days-old are plotted. Terminal root length is defined as the distance from the root meristem to the insertion of the first branch root. Squares represent day 29 while circles represent the day 43 harvest after the start of the experiment. Closed figures represent tap roots only while open figures denote lateral roots only.

Figure 6. Plot of bacteria counts versus the pH of buffered medium. Bacteria were maintained on slant cultures for one week at 25 C. At the end of this time, broth media (Ph.D. dissertation, A. J. Francis, Cornell Univ., 1974) was inoculated with one needle tip of *Rhizobium*. Two days later, the optical density of this stock culture was determined and a dilution series was made at 10^{-1} , 10^{-2} , 10^{-4} , 10^{-6} , and 10^{-8} . One ml of each diluted broth culture was placed in 5 ml of agar medium. Cultures were plated by the pour plate method to determine the concentration of the stock culture. Also, 0.1 ml of the stock culture was placed into sterile medium (citrate-phosphate buffer, 0.025 M) at various pH levels. Culture flasks were agitated and maintained at 25 C for 4 days. At the end of this time period cultures were diluted to 10^{-1} , 10^{-2} , 10^{-4} , 10^{-6} , and 10^{-8} with medium and plated by the pour plate method. Cultures were maintained for 14 days at 25 C. Plates were scored for the number of colonies and original colony concentrations were determined by multiplying the number of colonies by the reciprocal of the dilution factor.

Figure 7. Mean number of nodules per plants versus the duration of the aeroponics test. Few nodules are produced on roots below pH 5.6. A significant number are present on pH levels of 5.6 and 6.6 but the appearance of effective nitrogen fixation at pH 6.6 is delayed (see Figure 8). Symbols are: closed circles, pH 6.6; open circles, pH 5.6; closed triangles, 4.8; open squares, pH 4.4; closed squares, pH 3.8.

Figure 8. Plot of mean ethylene production per root system versus duration of exposure to various pH levels. Almost no ethylene production occurs in roots exposed to pH levels below 5.6. In the initial assay very high rates were produced at pH 5.6 with low rates at pH 6.6. Large amounts of ethylene production at 6.6 were produced 43 and 50 days after initiation of the experiments. This delay is correlated with the increase in nodules produced (Figure 7). Symbols: closed circles, pH 6.6; open circles, pH 5.6; closed triangles, pH 4.8; open squares, pH 4.4; closed squares, pH 3.8.

Figure 9. Mean over dry weights of roots and shoots at various pH levels. The curves represent five weekly harvest dates after the start of the experiment. Symbols: Closed triangles: harvest day - 22; open squares: May 29; open triangles: day 36; open circles: day 43; closed circles: day 50.

Figure 10. Root systems of *Pisum sativum* exposed to various pH levels are shown. Number of roots, length of roots, and number of nodules is evident. Samples were taken on day 54 after placement in aeroponic solutions.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Part II

EFFECT OF ACID RAIN ON SOIL MICROBIOLOGICAL ACTIVITIES

A. J. Francis, Director

ABSTRACT

Effect of acidity on microbial denitrification in an agricultural control soil, pH 5.2 and in a forest soil, pH 4.5 were studied by measuring the rate of N_2O production. At near neutral pH the rate of denitrification was very rapid and no N_2O was detected after a few days indicating further reduction to N_2 , the end product of denitrification. As the acidity of the soil increases, the rate of denitrification was rather slow and at greater acidities, pH 3.2, the N_2O accumulated in the incubation vessel affecting only partial denitrification.

These results suggest that significant amounts of N_2O can be generated from natural acid soils such as forest soils and from acidified soils as a result of acid rain. Thus acid precipitation may increase the N_2O concentrations in the atmosphere by affecting the biological denitrification in soils.

A preliminary study was conducted on the nitrogen fixation by free-living bacteria in a forest soil, and the data indicate that the distribution of free-living nitrogen fixing bacteria in an acidic forest soil is very low and nitrogen gains by asymbiotic bacterial fixation in such a forest ecosystem may be insignificant.

INTRODUCTION

Recent observations in the northeastern United States indicate that the pH of the rain is becoming more acidic due to the presence of increased levels of sulfur and nitrogen oxides in the atmosphere. In certain instances, a pH of 3.2 has been recorded at the laboratory. This steady increase in acidity causes a great concern, particularly of the living organisms in the ecosystem that are sensitive to increased hydrogen ion concentration.

Soil microorganisms play an important role in nature and are critical to ecosystem function, the well being of plants, animals and humans. They are responsible for transformations of various elements and occupy a significant place in carbon, nitrogen, phosphorus, and sulfur cycles. Soil is considered the ultimate sink for certain atmospheric pollutants. It has been shown that atmospheric gaseous pollutants such as carbon monoxide, ethylene, sulfur dioxide, nitrogen dioxide, and hydrocarbons are removed when reacted with soil by microbial and/or chemical means (Abeles, et al., 1971; Inman, et al., 1971). Although much is known about the chemical reactions of the pollutants, very little information is available on the effects of pollutants on soil microbial activity.

The objective of this study is to determine the effects of acid rain on soil microbial activity. As part of this study, (a) denitrification in an agricultural control soil and a forest soil were studied, and (b) a preliminary study was conducted on the nitrogen fixation by free-living bacteria in the forest soil.

DENITRIFICATION

Soil bacteria are known to volatilize nitrogen containing compounds in soil to nitrogenous gases. The conversion of nitrate to molecular nitrogen is called denitrification. Microbial denitrification and chemical volatilization of nitrogen are the two predominant processes that take place in soil. In the absence of molecular oxygen, nitrate serves as an electron acceptor in the metabolic oxidation of organic carbon by the bacteria. Bollag, et al. (1973), reported that chemical nitrite decomposition is predominant at low pH in sterile and non-sterile soils incubated under aerobic and anaerobic conditions. Little chemical decomposition of nitrite was observed in a neutral and alkaline environment (Bollag, et al., 1973). Major gases evolved during chemical nitrite volatilization are nitric oxide and nitrogen dioxide. Loss of nitrite occurred at a much greater rate under anaerobic, non-sterile conditions due to microbial activity. Nitrous oxide and molecular nitrogen were the major gaseous end products of biological denitrification.

NITROGEN FIXATION BY FREE-LIVING BACTERIA

Although an increase in nitrogen in soils by nonsymbiotic fixation is low, the ecological significance cannot be overlooked. With the advent of the acetylene reduction technique, an increased number of free-living nitrogen fixing organisms have been detected. Among bacteria Azotobacter, Beijerinckia and Clostridium have been extensively studied. Azotobacter is sensitive to high hydrogen ion concentration. Environments less than pH 6.0 contain little or no Azotobacter. Beijerinckia sp. are known to

grow and fix nitrogen from pH 3 to 9. Very little information is available on the effect of environmental factors particularly acidity on the ability of the nitrogen fixing bacteria to survive, grow and fix nitrogen in soils.

MATERIALS AND METHODS

Sample Preparation

Soil samples used in this study were collected from a control agricultural field and from a pine forest at Brookhaven National Laboratory. Soil samples 0-4" deep were collected, air dried, thoroughly mixed, and passed through a 2 mm sieve. The pH of the soil samples were determined as described by Greweling and Peech (1965).

To study the effect of acidity on soil microbial activity, the pH of the control agricultural soil was adjusted to 3.0, 3.5, 4.5, 6.5, and 7.2 by addition of 0.1N H_2SO_4 or 0.1N NaOH. To 500 g of soil, 500 ml of deionized water was added and the pH adjusted to the desired level, air dried and stored at room temperature until further use.

About 1000 g of air dried forest soil sample was adjusted to pH 3.0, 4.0, 6.0 and 7.0 by addition of 1N H_2SO_4 or $\text{Ca}(\text{OH})_2$. The pH adjusted and the natural soil samples were incubated in the dark at 26°C. The moisture content was maintained at 28% by periodic addition of distilled water. A portion of the soil sample was removed, air dried, the pH of the sample was determined and then used in the nitrogen transformation studies.

Denitrification

Twenty-five grams of native and pH adjusted soil samples were transferred to 60 ml glass bottles. The agricultural control soil sample

received 10 ml of distilled water with KNO_3 to give a final concentration of 56 ppm NO_3^- -N. The forest soil sample received 7 ml of distilled water with KNO_3 to give a final concentration of 11 ppm of NO_3^- -N. Control samples received only distilled water. Reaction bottles were fitted with serum stoppers, evacuated, and flushed with helium several times and finally filled with helium to atmospheric pressure. Triplicate samples were prepared and incubated at 26°C. A 0.2 ml head space gas sample was withdrawn with a gas tight syringe at time 0 and at regular intervals and analyzed by gas chromatography for denitrification products. Standard gas samples of N_2O and CO_2 were prepared and analyzed to obtain quantitative data.

Gas Chromatography

A Perkin-Elmer model 3920 gas chromatograph equipped with a thermal conductivity detector was used to detect N_2O and CO_2 . A 12' x 1/8" stainless steel column packed with Porapak QS (80/100 mesh) was used to separate the various components. The detector was operated at 260°C with a filament current of 175 mA. The injector temperature was 80°C. The column temperature was held initially at 70°C for one minute, and increased to 120°C at the rate of 16°/min. Helium was used as carrier gas at a flow rate of 35 ml/min.

Enumeration of Nitrate Reducing and Denitrifying Bacteria

The nitrate reducing and denitrifying bacterial population in the natural and pH adjusted soil samples were determined by the most probable number (MPN) technique (Alexander, 1965), modified by Focht and Joseph (1973).

Nitrogen Fixation

Freshly collected forest soil sample and pH adjusted moist soil sample incubated for more than four weeks were used. Fifteen grams of air dried soil samples were transferred to 20 ml serum bottles. Four milliliters of distilled water containing glucose was added to give a final concentration of 1% glucose. Unamended samples received 4 ml of distilled water. Samples were pre-incubated either aerobically or anaerobically for 96 hours. Aerobic sample bottles were fitted with foam plugs. Anaerobic sample bottles were fitted with serum caps, evacuated and flushed several times with N_2 , and finally filled with N_2 to the atmospheric pressure, and incubated at 26°C in the dark.

Acetylene Reduction Assay

After 96 hours pre-incubation, the samples were assayed for nitrogen fixation by acetylene reduction technique.

Nitrogen fixation by free-living aerobic bacteria were assayed after the foam plugs of the aerobically pre-incubated samples were replaced with serum caps. The atmosphere was replaced with He, 0.22 atm of O_2 , and 0.1 atm of acetylene.

Nitrogen fixation by anaerobic bacteria was determined by replacing the N_2 atmosphere with He and 0.1 atm of acetylene. Control samples were included to detect the presence of ethylene producing organisms in the soil.

Samples were incubated at 26°C in the dark for two hours. At the end of the incubation period, 0.2 ml gas sample was withdrawn and analyzed for ethylene by means of a gas chromatograph (Perkin-Elmer model 3920) equipped with a flame ionization detector. The gas chromatograph was fitted with a 9 ft. long, 1/8 in. diameter stainless steel column packed with

Porapak R (100-120 mesh). The column, detector, and injector temperatures were 50°, 60°, and 75°C, respectively.

RESULTS AND DISCUSSION

Denitrification

A representative chromatogram of the head space gas analysis by gas chromatography used in the denitrification studies is shown in Figure 1. An agricultural control field soil with a normal pH 5.2 produced N_2O when amended with various levels of nitrate (Table 1). Nitrous oxide was not detected in any control soil samples without nitrate. Rate of N_2O production in soils amended with 56, 112, 168, and 224 ppm nitrate-N were similar and at these levels available carbon may be limiting as these samples were not amended with any carbon source. The native carbon present in the soil must be used as the energy source by the denitrifying bacteria. There were no significant differences in the amounts of CO_2 produced between the control and the nitrate treated soils (Table 2).

Effects of pH on microbial denitrification were studied anaerobically in pH adjusted soil, amended with 56 ppm of NO_3^- -N. Samples were analyzed for N_2O production and no attempt was made to determine the N_2 gas in these experiments (Figure 2). Less N_2O was observed in soil at pH 7.2 than lower pH levels which indicates a rapid conversion of N_2O to N_2 . At pH 6.5, 1.2 μ moles of N_2O was detected at 50 hours of incubation and then it started to disappear. The rate of N_2O formation and disappearance at pH 5.2 (normal pH of the soil) and 4.5 followed more or less the same pattern. Nitrous oxide was detected at pH 3.5 and 3.0 after 3 and 6 days of incubation, respectively. At pH 3.5, maximum N_2O production (1.6 μ moles) was observed at day 17.

The effect of pH on CO_2 production in nitrate (56 ppm) amended soil is shown in Figure 3. More CO_2 was produced at pH 6.5 followed by 7.2, 5.2 and 4.5 than at pH 3.0 and 3.5. Since these soils were not enriched with organic carbon, the native organic material present in the soil served as carbon source for the denitrifying organisms. Carbon dioxide production decreased as soil acidity increased, indicating the sensitivity of denitrifying and other anaerobic bacteria to increased acidity. The change in pH of soil samples used in this experiment at the end of incubation period is shown in Table 3.

The natural and pH adjusted forest soil samples that had been pre-incubated under moist conditions for 2 weeks and 5 weeks were used to measure the rate of denitrification. Soils were amended with 11 ppm of NO_3^- -N and the nitrous oxide and CO_2 evolution were monitored. Control soils received no nitrate.

Nitrous oxide and CO_2 production by the two weeks pre-incubated soil samples are shown in Figures 4 and 5. Control soils that received no nitrate produced little nitrous oxide. The rate of N_2O production and disappearance at pH 6.6 and 5.8 was very rapid. The natural pH 4.6 and pH adjusted soil 4.3 followed the same pattern. However, at pH 3.2 N_2O production was slow and did not disappear as rapidly as in higher pH soils. The pH analyses of the soil samples are shown in Table 4.

Nitrous oxide and CO_2 production by the five weeks pre-incubated soil samples are shown in Figure 6 and 7. The pH of the soil samples changed gradually from two weeks to five weeks incubation by about 0.2 units. Nitrous oxide production at pH 6.8 and 6.1 followed the same pattern as in the previous experiment. The rate of production and disappearance of

N_2O at pH 4.5 was more rapid and after two days no N_2O was detected. The most acidic soil (pH 3.4) produced N_2O at a slow rate and it accumulated in the incubation vessel. The pH analyses of the soil samples at the termination of the experiment is shown in Table 5. Control soil samples without nitrate produced nitrous oxide. However, after 2 days no nitrous oxide was detected in these samples except pH 3.4 soil sample where nitrous oxide accumulated in the headspace as shown in Figure 8. Carbon dioxide production by the control unamended soil is shown in Figure 9 and the final pH at the end of the experiment is tabulated in Table 6.

No nitrous oxide was detected in autoclaved control soil samples (pH 3.4, 4.5, 6.1 and 6.8) with and without nitrate. These results suggest that nitrous oxide production is biological and at very acidic conditions, there is an incomplete or partial denitrification, where nitrous oxide is formed and is not further reduced to N_2 gas.

Nitrate reducing and denitrifying bacterial populations in native and pH adjusted forest soil samples used in the N_2O evolution study were enumerated. Initial and final counts of the bacterial populations are tabulated in Tables 7 and 8. In these experiments, the assay media pH was adjusted to the soil pH or to pH 7.0. At soil pH 3.4, denitrifying populations were very low, and did not increase in numbers even when incubated with nitrate. Moreover, these bacteria did not grow at pH 3.4 of the assay media which indicates their sensitivity to acidity. At pH 4.5, initial population counts were the same as pH 3.4 samples, but increased after two weeks of incubation. There was no significant difference in the denitrifying population in pH 4.5 soil samples incubated with and without nitrate and at pH 4.5 and 7.0 of the assay media. At pH 6.1 and 6.8, initial and final counts of the bacteria were higher which correlated with rapid N_2O production and reduction.

These results suggest that at acidic pH (3.4) denitrifying bacteria are unable to grow but are present in the soil. Since no N_2O was detected in the autoclaved soils suggests that N_2O formation is biological and at acidic pH, there may be microsites conducive for the bacteria for N_2O production.

Nommik (1956), Wjler and Delwiche (1954), and Bremner and Shaw (1958) have shown that the rate of denitrification increases linearly from pH 4 to pH 7-8 and declines at pH 9.5. Neutral to slightly alkaline pH ranges produce faster rates of total denitrification. Proportionally less N_2O was observed in soil as the pH rises (Nommik, 1956; Wjler and Delwiche, 1954). The rate of N_2O reduction apparently increases with pH faster than its rate of formation. In native soil samples (pH 5.2), the rate of denitrification is slower when compared to pH adjusted neutral soils. As the acidity of the soil increases, the rate of formation of N_2O and conversion to N_2 decreases. However, at greater acidities, N_2O accumulates in the incubation vessel. In an open and dynamic system, such as the natural ecosystem, it is possible that N_2O is released into the atmosphere.

If there is a change in soil pH due to acid rain, then there will be an effect on biological denitrification, especially in sandy and poorly buffered acid soils, such as soils of Long Island. Thus, acidic soils and forest soils may play a major role in the biogenic emission of nitrous oxide.

Nitrogen Fixation by Free-Living Bacteria

Acetylene reduction by free-living bacteria in a freshly collected forest soil sample (pH 4.4) is shown in Table 9. Little ethylene was produced by samples treated with glucose and incubated aerobically. Anaerobic samples without glucose produced little ethylene; whereas, glucose amended samples produced 3.2 μ moles of ethylene/g dry soil/hr.

Nitrogen fixation by free-living bacteria in pH adjusted forest soil samples that had been kept moist and incubated for several weeks in the laboratory were studied. At pH 3.6 none of the treatments showed any detectable activity (Table 10). Similar results were observed with pH 4.7 soil sample (Table 11). At pH 5.8, only samples amended with glucose and incubated under aerobic or anaerobic conditions showed activity (Table 12). Anaerobic soil samples amended with glucose at pH 6.4 had a higher activity 37.5 μ moles of ethylene formed/g dry soil/hr; whereas, other treatments did not have any measurable activity (Table 13).

These results suggest that the distribution of free-living nitrogen fixing bacteria in acidic forest soil samples are very low and nitrogen input by asymbiotic bacterial fixation in forest soils may be insignificant.

REFERENCES

- Abeles, F. B., L. E. Craker, L. E. Forrence, and G. R. Leather, Fate of air pollutants: Removal of ethylene, sulfur dioxide and nitrogen dioxide by soil, *Science* 173, pp. 914-916 (1971).
- Alexander, M., Most-probable-number method for microbial populations, in *Methods of Soil Analysis, Part 2*, ed. by C. A. Black, et al., Madison, Wis., Amer. Soc., Agron., 1965, pp. 1467-1472.
- Bollag, J. M., S. Drzymala and L. T. Kardos, Biological versus chemical nitrite decomposition in soil, *Soil Science* 116, pp. 44-50 (1973).
- Bremner, J. M. and K. Shaw, Denitrification in soil. II. Factors affecting denitrification, *J. Agric. Sci.* 51, pp. 40-52 (1958).
- Focht, D. D. and H. Joseph, An improved method for the enumeration of denitrifying bacteria, *Soil Sci. Soc. Amer. Proc.* 37, pp. 698-699 (1973).
- Greweling, T. and M. Peech, Chemical Soil Tests, Cornell University Agricultural Experimentation, Bulletin #960, 1965.
- Inman, R. E., R. B. Ingersoll, and E. A. Levy, Soil: A natural sink for carbon monoxide, *Science* 172, pp. 1229-1231 (1971).
- Nommik, H., Investigations on denitrification in soil, *Acta Agr. Scand.* 6, pp. 195-228 (1956).
- Wijler, J. and C. C. Delwiche, Investigations on the denitrifying process in soil, *Plant and Soil* 5, pp. 155-169 (1954).

TABLE 1

N₂O Production in an Agricultural Control Soil Amended with Nitrate

Nitrate-N added (ppm)	μ moles of N ₂ O Produced/g Dry Soil					
	Days					
	2	3	4	9	10	11
0 (Control)	0	0	0	0	0	0
28	0.44	0.80	0.84	.08	.08	0
56	0.48	1.16	1.56	0.76	.80	0.48
112	0.52	1.12	1.52	1.8	2.20	2.20
168	0.44	1.20	1.68	2.0	2.44	2.32
224	0.44	1.36	1.76	2.12	2.60	2.36

TABLE 2

CO₂ Production in an Agricultural Control Soil Amended with Nitrate

Nitrate-N added (ppm)	μ moles of CO ₂ Produced/g Dry Soil					
	Days					
	2	3	4	9	10	11
0 (Control)	1.2	1.5	2.2	4.4	5.2	7.2
28	1.8	2.4	3.6	4.4	5.6	6.9
56	2.0	2.6	3.8	4.5	5.2	7.0
112	1.8	2.6	3.7	5.1	5.1	6.4
168	1.9	2.6	3.9	5.2	5.1	6.9
224	1.9	2.7	4.1	5.3	5.2	7.2

TABLE 3

pH of Soil Samples Amended with Nitrate

<u>Initial</u>	<u>Final</u>
7.2	7.0
6.5	6.6
5.2	6.1
4.5	5.0
3.5	3.6
3.0	3.2

TABLE 4

pH of Forest Soil Sample Amended with Nitrate

<u>Initial</u>	<u>Final</u>
6.6	6.7
5.8	6.0
4.5	5.3
4.3	5.1
3.2	3.5

TABLE 5

pH of the Forest Soil Sample Amended with Nitrate

<u>Initial</u>	<u>Final</u>
6.8	6.7
6.1	6.3
4.5	5.2
3.4	3.6

TABLE 6

pH of the Control Forest Soil Sample

<u>Initial</u>	<u>Final</u>
6.8	6.8
6.1	6.2
4.5	5.8
3.4	3.7

TABLE 7

Population of Denitrifying Bacteria in a pH Adjusted BNL - Forest Soil

pH				Denitrifiers/g Dry Soil			
Soil		Assay Media		Soil Incubated			
Day		Day		Without NO_3^- -N		With 11 ppm NO_3^- -N	
Day		Day		Day		Day	
0	14	0	14	0	14	0	14
3.4	3.6	3.4	3.4	0	0	0	0
3.4	3.6	7.0	6.8	2.3×10^1	2	2.3×10^1	1.8×10^1
4.5*	5.2	4.5	6.7	2.3×10^1	1.3×10^2	2.3×10^1	5.5×10^2
4.5*	5.2	7.0	-	2.1×10^1	1.9×10^2	2.1×10^1	9.4×10^2
6.1	6.3	6.2	7.1	-	-	7.8×10^4	2.0×10^5
6.8	6.7	6.9	7.2	-	-	1.4×10^5	3.6×10^6

*Natural pH of the soil sample.

TABLE 8

Population of Nitrate Reducing Bacteria in a pH Adjusted BNL - Forest Soil

pH				Nitrate Reducers/g Dry Soil			
Soil		Assay Media		Soil Incubated			
Day		Day		Without NO_3^- -N		With 11 ppm NO_3^- -N	
Day		Day		Day		Day	
0	14	0	14	0	14	0	14
3.4	3.6	3.4	3.4	0	0	0	0
3.4	3.6	7.0	6.8	2.2×10^4	3.8×10^4	2.2×10^4	6.2×10^4
4.5*	5.2	4.5	6.7	2.1×10^3	2.7×10^4	2.1×10^3	1.7×10^4
4.5*	5.2	7.0	-	7.0×10^4	3.6×10^4	7.0×10^4	4.2×10^4
6.1	6.3	6.2	7.1	-	-	7.9×10^5	1.1×10^7
6.8	6.7	6.9	7.2	-	-	2.2×10^6	1.6×10^7

*Natural pH of the soil sample.

TABLE 9
Nitrogen Fixation by Free Living Bacteria
in a BNL - Forest Soil* (pH 4.4)

<u>Treatment</u>	<u>n Moles of Ethylene Formed/g Dry Soil/Hr</u>	<u>Final pH</u>
<u>Aerobic</u>		
Glucose -; C ₂ H ₂ -	0	4.5
Glucose +; C ₂ H ₂ -	0	3.5
Glucose -; C ₂ H ₂ +	0	4.5
Glucose +; C ₂ H ₂ +	.07	3.5
<u>Anaerobic</u>		
Glucose -; C ₂ H ₂ -	0	4.4
Glucose +; C ₂ H ₂ -	0	4.0
Glucose -; C ₂ H ₂ +	.01	4.4
Glucose +; C ₂ H ₂ +	3.22	4.1

* Fresh soil sample collected on 5/17/77 and analyzed on 5/23/77.

TABLE 10
Nitrogen Fixation by Free Living Bacteria in a
pH Adjusted BNL - Forest Soil (pH 3.6)

<u>Treatment</u>	<u>n Moles of Ethylene Formed/g Dry Soil/Hr</u>	<u>Final pH</u>
<u>Aerobic</u>		
Glucose -; C ₂ H ₂ -	0	3.6
Glucose +; C ₂ H ₂ -	0	3.5
Glucose -; C ₂ H ₂ +	0	3.6
Glucose +; C ₂ H ₂ +	0	3.5
<u>Anaerobic</u>		
Glucose -; C ₂ H ₂ -	0	3.6
Glucose +; C ₂ H ₂ -	0	3.6
Glucose -; C ₂ H ₂ +	0	3.6
Glucose +; C ₂ H ₂ +	0	3.6

TABLE 11

Nitrogen Fixation by Free Living Bacteria in a
pH Adjusted BNL - Forest Soil (pH 4.7)

<u>Treatment</u>	<u>n Moles of Ethylene Formed/g Dry Soil/Hr</u>	<u>Final pH</u>
<u>Aerobic</u>		
Glucose -; C ₂ H ₂ -	0	4.9
Glucose +; C ₂ H ₂ -	0	4.7
Glucose -; C ₂ H ₂ +	0	4.9
Glucose +; C ₂ H ₂ +	0	4.8
<u>Anaerobic</u>		
Glucose -; C ₂ H ₂ -	0	4.8
Glucose +; C ₂ H ₂ -	0	3.3
Glucose -; C ₂ H ₂ +	0	4.8
Glucose +; C ₂ H ₂ +	0	3.4

TABLE 12

Nitrogen Fixation by Free Living Bacteria in a
pH Adjusted BNL - Forest Soil (pH 5.8)

<u>Treatment</u>	<u>n Moles of Ethylene Formed/g Dry Soil/Hr</u>	<u>Final pH</u>
<u>Aerobic</u>		
Glucose -; C ₂ H ₂ -	0	5.8
Glucose +; C ₂ H ₂ -	0	4.0
Glucose -; C ₂ H ₂ +	0	5.8
Glucose +; C ₂ H ₂ +	0.25	4.0
<u>Anaerobic</u>		
Glucose -; C ₂ H ₂ -	0	5.8
Glucose +; C ₂ H ₂ -	0	5.1
Glucose -; C ₂ H ₂ +	0	5.8
Glucose +; C ₂ H ₂ +	0.04	5.1

TABLE 13

Nitrogen Fixation by Free Living Bacteria in a
pH Adjusted BNL - Forest Soil (pH 6.4)

<u>Treatment</u>	<u>n Moles of Ethylene Formed/g Dry Soil/Hr</u>	<u>Final pH</u>
<u>Aerobic</u>		
Glucose -; C ₂ H ₂ -	0	6.4
Glucose +; C ₂ H ₂ -	0	4.7
Glucose -; C ₂ H ₂ +	0	6.4
Glucose +; C ₂ H ₂ +	0	4.7
<u>Anaerobic</u>		
Glucose -; C ₂ H ₂ -	0	6.6
Glucose +; C ₂ H ₂ -	0	5.5
Glucose -; C ₂ H ₂ +	0	6.6
Glucose +; C ₂ H ₂ +	37.45	5.5

Figure 1. Gas chromatogram of CO₂ and N₂O analysis.

Figure 2. Effect of pH on N_2O production in an agricultural control field soil amended with 56 ppm NO_3^- -N.

Figure 3. Effect of pH on CO_2 production in an agricultural control field soil amended with 56 ppm NO_3^- -N.

Figure 4. Effect of pH on N₂O production in a BNL-forest soil amended with 11 ppm NO₃⁻-N. (Natural and acidified soil samples pre-incubated for 2 weeks under moist conditions without any other additions were used in this experiment.)

Figure 5. Effect of pH on CO₂ production in a BNL-forest soil amended with 11 ppm of NO₃⁻-N. (Natural and acidified soil samples pre-incubated for 2 weeks under moist conditions without any other additions were used in this experiment.)

Figure 6. Effect of pH on N_2O production in a BNL-forest soil amended with 11 ppm NO_3^- -N. (Natural and acidified soil samples pre-incubated for 5 weeks under moist conditions without any other additions were used in this experiment.)

Figure 7. Effect of pH on CO_2 production in a BNL-forest soil amended with 11 ppm NO_3^- -N. (Natural and acidified soil samples pre-incubated for 5 weeks under moist conditions without any other additions were used in this experiment.)

Figure 8. Effect of pH on N₂O production in a BNL-forest control soil. (Natural and acidified soil samples pre-incubated for 5 weeks under moist conditions without any other additions were used in this experiment.)

Figure 9. Effect of pH on CO₂ production in a BNL-forest control soil. (Natural and acidified soil samples pre-incubated for 5 weeks under moist conditions without any other additions were used in this experiment.)

Part III

SEQUENTIAL PRECIPITATION CHEMISTRY
AND ALLIED METEOROLOGICAL DATA

G. Raynor, Director

PRECIPITATION SAMPLES

The Automatic Sequential Precipitation Sampler described in the previous annual progress report (BNL 50575) was put in operation on June 1, 1976, and has operated continuously except for a few brief outages caused by power failure, electrical surges from nearby lightning strikes and a shorted heating tape under the funnel. A detailed report on the sampler and its operation is in press.¹

Over 700 hourly precipitation samples large enough for at least the pH measurement were collected through September 1977. Samples were taken from the collector as soon as possible after precipitation ended, transferred to 500-ml covered plastic bottles and taken to the Analytical Chemical Laboratory of the Department of Energy and Environment for analysis. All were analyzed, as far as volume permitted, by the following methods:

pH Analysis. pH is determined as soon as possible after the sample is received using an Orion pH meter.

Specific Conductivity is determined as soon as possible after the sample is received using a Markson Electromark Digital Analyzer.

$\text{NO}_3^- + \text{NO}_2^-$ are determined colorimetrically using an automated procedure and the Technicon Auto Analyzer. NO_3^- is reduced with hydrazine and the NO_2^- reacts with sulfanilamide under acidic conditions to form a diazo compound. This compound couples with N-1-naphthylethylene-diamine dihydrochloride to form a reddish-purple azo dye which is read at 530 nm. Samples are refrigerated prior to analysis to prevent degradation of the ions.

NH_4^+ is determined using the Berthelot reaction in which a blue-colored compound, indophenol blue, is formed when a solution of an ammonium salt is added to sodium phenoxide followed by the addition of sodium hypochlorite in

an alkaline medium. The indophenol blue is measured at 630 nm. The procedure is automated and run on a Technicon Auto Analyzer. Samples are refrigerated prior to this analysis to prevent degradation of the NH_4^+ .

$\text{SO}_4^{=}$ analyses are performed using an automated turbidimetric procedure and the Technicon Auto Analyzer. $\text{SO}_4^{=}$ is converted to a BaSO_4 suspension under controlled conditions and the resulting turbidity is read at 420 nm. An alternate procedure occasionally used is the methyl thymol blue colorimetric procedure. Identical results are obtained with both procedures.

CL^- is determined using an automated colorimetric procedure and the Technicon Auto Analyzer. Thiocyanate ion (SCN^-) is liberated from mercuric thiocyanate through sequestering of mercury by chloride. The liberated SCN^- in the presence of ferric ion forms a highly colored ferric thiocyanate, in concentration proportional to the original chloride. The complex is measured at 467 nm.

Na^+ is determined by atomic absorption analysis using a Perkin-Elmer 303 or Instrumentation Laboratory Atomic Absorption Spectrophotometer.

All the data were tabulated on input sheets from which cards were punched. The data were then transferred to magnetic tape for analyzing by computer. Precipitation data recorded included station, date, precipitation event number during the month, sample number during the event, time of beginning and end of each sample, duration of each sample and amount of rainfall during each sample.

Chemical data recorded were pH, conductivity and concentrations of total nitrogen, ammonia, sulfate, sodium and chloride ions. Meteorological data included precipitation type, cloud type, synoptic situation causing the precipitation, wind direction and speed at 108 m on a nearby meteorological

tower and air temperature. Storm trajectories will be determined later from appropriate surface and upper air data and added to the data base.

A comprehensive statistical analysis of the first year of data has been performed and a report on results is in press.² Chemical constituents of the samples were related to each other by correlation and regression analyses and to meteorological variables by standard statistical procedures. In addition, a number of case studies of changes during individual precipitation events have been made. Some of these and some preliminary statistical studies have been described in a paper prepared for presentation at a meeting of the American Institute of Chemical Engineers.³

As examples of the results obtained, Figure 1 shows the concentration of hydrogen ion for the four seasons and for six synoptic types. The weighted values are weighted by rainfall rate during the sample period while the unweighted values are arithmetic means for each classification. Figure 2 shows the sulfate concentrations for eight wind direction sectors and Figure 3 shows the relationship between nitrate and sulfate concentrations.

REFERENCES

1. Raynor, G. S. and J. P. McNeil, *The Brookhaven Automatic Sequential Precipitation Sampler* (in press).
2. Raynor, G. S. and J. V. Hayes, *Experimental Data from Analyses of Sequential Precipitation Samples at Brookhaven National Laboratory* (in press).
3. Raynor, G. S., *Meteorological and Chemical Relationships from Sequential Precipitation Samples*, presented at 70th Annual Meeting AIChE, New York, New York, Nov. 13-17, 1977, Brookhaven National Laboratory Report 22879.

Figure 1.

Figure 2. Sulfate concentration (ppm),
weighted mean 3.72.

Figure 3.