Project Summary:**Biochemical and Molecular Analysis of a Transmembrane Protein Kinase from *Arabidopsis thaliana***

A. B. Bleecker, Department of Botany

Receptor protein kinases on the plasma membrane of cells interact with various growth factors and hormonal agents, generating intracellular signals from these external signals. As such the receptors play a pivotal role in the ability of a cell to respond to its environment and develop. Although widely studied in animals, very little is known about receptor kinases in plants.

We have isolated genomic and cDNA clones encoding a novel receptor-like protein kinase from the higher plant *Arabidopsis thaliana*. This kinase is being studied by combining biochemical, molecular, and genetic approaches. Domain-specific antibodies immunodecorate a polypeptide with a molecular mass of 120,000 daltons in extracts of *Arabidopsis*, where it has been found in all portions of the plant examined including root, stem, leaf, flower, and silique. Cytochemical analysis and initial studies using the kinase promoter with the GUS reporter gene system also indicate that the kinase is present throughout the plant. The kinase is glycosylated, like animal receptor kinases, and has been partially purified from *Arabidopsis* by using lectin columns. The kinase has been expressed in *E. coli*, purified, and found to autophosphorylate on serine and threonine residues, but not on tyrosine residues. As such, it belongs to the small family of receptor-like kinases with serine/threonine specificity. Transgenic plants are now being produced that either overexpress or carry altered forms of the protein kinase gene. These experiments will help determine the natural role the kinase plays in a pathway of signal transduction.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

RECEIVED

JUN 01 1993

OSTI

MASTER

**Biochemical and Molecular Analysis of a Transmembrane Protein
Kinase from *Arabidopsis thaliana***

**Progress Report (January 1993)
for Grant No. DE-FG02-91ER20029**

A. B. Bleecker

Introduction:

Receptor protein kinases on the plasma membrane of cells interact with various growth factors and hormonal agents, generating intracellular signals from these external signals. As such the receptors play a pivotal role in the ability of a cell to respond to its environment and develop. Although widely studied in animals, very little is known about receptor kinases in plants.

We have isolated genomic and cDNA clones encoding a novel receptor-like protein kinase from the higher plant *Arabidopsis thaliana*. We have designated this transmembrane kinase gene TMK1. The product of the TMK1 gene has all the features expected in a receptor protein kinase: an amino-terminal signal sequence for targeting to a membrane, an extracellular receptor domain, a hydrophobic transmembrane domain, and an intracellular kinase domain. Two features of TMK1 are of particular interest. First, the extracellular domain has 11 copies of a leucine-rich repeat (LRR) sequence. Leucine rich repeats have only been found in proteins associated with the plasma membrane, and are implicated in protein/protein interactions. No protein with leucine rich repeats has previously been identified or characterized in plants. Second, the kinase domain of TMK1 contains sequences that are diagnostic for a serine/threonine specific kinase, not a tyrosine specific kinase as is typical of the animal receptor kinases. We are following biochemical, molecular, and genetic approaches in characterizing the TMK1 kinase and the role that it plays in plant signal transduction.

Progress Summary: *Reprint removed*

Enclosed Reprint. A reprint is enclosed titled "The TMK1 Gene from *Arabidopsis* Codes for a Protein with Structural and Biochemical Characteristics of a Receptor Protein Kinase." This paper was published this past year and details the isolation of clones for the TMK1 gene and structural features of the predicted polypeptide. It also describes the raising of antibodies against the TMK1 polypeptide. Using the antibodies we detect a 120 kD polypeptide in extracts of *Arabidopsis* from leaf, stem, root, and floral tissues. Finally, we describe the expression of the TMK1 gene as a fusion protein in *E. coli*. The purified fusion protein was found to autophosphorylate on serine and threonine residues, but not on tyrosine residues. This result indicates that the TMK1

polypeptide does indeed belong to the small family of receptor-like kinases with serine/threonine specificity.

Expression of the TMK1 kinase in *E. coli*. As described in the enclosed reprint, we are using an *E. coli* expression system in order to examine the protein kinase activity of the TMK1 polypeptide. The kinase portion of the TMK1 gene was cloned into the pMAL-c vector and transformed into *E. coli*. Upon induction, *E. coli* expressed a fusion of the kinase with maltose-binding protein, which could then be affinity-purified by binding to amylose resin. Our initial studies indicated that the fusion protein had kinase activity and autophosphorylated on serine and threonine residues. We are continuing our characterization of the expressed kinase in order to gain a better understanding of the biochemical nature of the TMK1 kinase and its similarities to the known receptor kinases.

Protein kinases typically require a divalent cation for activity, usually Mg^{2+} . It has been found, however, that most of the animal receptor kinases exhibit greater activity with Mn^{2+} than with Mg^{2+} . We examined the divalent cation requirement of the TMK1 fusion protein and we found that, like the animal receptor kinases, the kinase exhibited greater autophosphorylation activity with Mn^{2+} than with Mg^{2+} .

Overexpression of kinases in *E. coli*, particularly the animal tyrosine kinases, often results in promiscuous phosphorylation of many polypeptides. In contrast, we find that the TMK1 fusion kinase has a very limited substrate specificity. In crude extracts of *E. coli*, the TMK1 fusion protein autophosphorylates to a high extent, but does not appear to appreciably phosphorylate any of the *E. coli* polypeptides. We have also examined the substrate specificity of the TMK1 kinase in more detail following affinity purification. The TMK1 kinase phosphorylates typical arginine-rich substrates such as histone VIII_S, but does not phosphorylate lysine-rich substrates such as histone III_S. However, the best exogenous substrate we have found is a fusion protein containing just the carboxy-terminal region of the TMK1 polypeptide. This is a fusion of glutathione S-transferase to the terminal 99 amino acids of the TMK1 polypeptide. This carboxy-terminal fusion inhibits autophosphorylation of the TMK1 fusion protein and is also itself phosphorylated even at very low concentrations. As such it appears to be acting as a competitive substrate for the autophosphorylation sites of the TMK1 fusion kinase. Glutathione S-transferase by itself does not show these substrate characteristics.

Autophosphorylation is an important means of regulation for many of the animal receptor kinases, where it has been found to modify enzyme kinetics and mediate interactions between the kinase and other proteins. We are working to determine autophosphorylation sites of the TMK1 kinase. For this purpose the carboxy-terminal fusion is extremely useful, since it is a relatively short sequence that appears to contain some of the major phosphorylation sites of the TMK1 kinase.

Localization of the native form of the TMK1 polypeptide in *Arabidopsis*. Using antibodies, we have detected the TMK1 polypeptide in extracts of all plant tissue types examined. We are pursuing two independent methods to gain more explicit information as to what cell types express the TMK1 gene--immunocytochemical localization and the β -glucuronidase (GUS) reporter gene system.

Immunocytochemical localization at the cellular level indicates the presence of the TMK1 polypeptide in all living cells examined. Paraffin-embedded stem, leaf, and root sections all showed staining throughout the section. Controls using preimmune serum or TMK1 antibody preabsorbed with excess antigen showed no staining.

For the GUS reporter gene system, we inserted the promoter region of the TMK1 gene in front of the reporter gene GUS on a plasmid. *Arabidopsis* plants were transformed with the construct, and multiple independent transgenic lines were isolated. These were histochemically assayed for GUS expression with X-GLUC, which results in a deep blue color in expressing cells. We observe GUS expression in all cell types, consistent with the immunocytochemical data.

We are now initiating subcellular immunolocalization of the TMK1 polypeptide in *Arabidopsis* using electron microscopy. Amino acid sequence and cellular fractionation studies indicate that the TMK1 polypeptide is probably membrane-associated. Subcellular immunolocalization should allow us to clearly determine to what membrane system the TMK1 polypeptide is localized, generating a clearer understanding of its function. For this work plant tissues are fixed in 2% glutaraldehyde and 2% paraformaldehyde, and then resin-embedded. The embedded tissue is then sectioned using a diamond knife, incubated with antibody, and labeled with immunogold for ultrastructural localization.

Transformation of TMK1 into plants. In order to better understand the natural role of the TMK1 kinase in the plant, we have transformed *Arabidopsis* with several TMK1 constructs.

In one case we have put the entire TMK1 gene behind the soybean heat shock promoter, by which we hope to see what effect overexpression of the protein has upon the plant. We now have several independent transformants, and progeny are being examined for changes in response to overexpression. Plants are incubated 2 hours daily at 37 C for two weeks to induce the heat shock promoter without developmentally affecting the plants. Overexpression of TMK1 will be confirmed by RNA and protein analysis.

In a second case, we are collaborating with Dr. Mitchell Altshuler to inhibit expression of the TMK1 gene using a hairpin ribozyme. Ribozymes are RNAs capable of RNA sequence-specific recognition that then destroy the RNA substrate by a site-specific cleavage. We have designed a ribozyme capable of specific cleavage of the TMK1 RNA. The ribozyme has been tested *in vitro* and found capable of efficient and specific cleavage of the TMK1 RNA. A construct encoding this ribozyme, under control of the 35S promoter, has now been introduced into *Arabidopsis* and progeny are currently being analyzed.

Finally, we are preparing several other constructs that we also wish to transform into plants. These include just the receptor portion of TMK1, which based on studies with animal receptor kinases could serve as a dominant inhibitor of the kinase, and just the kinase portion of TMK1 which could be constitutively active.

END

**DATE
FILMED**

6 / 17 / 93