BNL # 48381
Informal Report
Limited Distribution

EFFECTS OF SELECTED THERMOPHILIC MICROORGANISMS ON
CRUDE OILS AT ELEVATED TEMPERATURES AND PRESSURES

BNL--48381
DE93 009026

QUARTERLY REPORT (07/01/92 - 09/30/92)

Prepared by

E. T. Premuzic and M. S. Lin

Department of Applied Science
Brookhaven National Laboratory
Upton, New York 11973

October 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Contract No AS-219-ECD, U.S. DOE, Division of Fossil Fuels, Bartlesville
Project Office
Contractor: Associated Universities, Brookhaven National Laboratory

* This work performed under the auspices of the U.S. DOE Contract No. DE-
AC02-76CH00016.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

RECEIVED
MAR 10 1993
071
MASTER

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor, or subcontractor thereof.

QUARTERLY REPORT

BPO EOR FY 1992 (07/01/92 - 09/30/92)

TITLE: Effects of Selected Thermophilic Microorganisms
on Crude Oils at Elevated Temperatures and
Pressures

CONTRACT NO.: AS-219-ECD

CONTRACTOR: Associated Universities, Inc., Brookhaven
National Laboratory

PRINCIPAL INVESTIGATORS: E. T. Premuzic and M. S. Lin

OBJECTIVES OF RESEARCH PROJECT

The objective of this program is to determine the chemical and physical effects of thermophilic and thermoadapted organisms on crude oils and cores at elevated temperatures and pressures. Ultimately a data base will be generated which will be used in technical and economic feasibility studies leading to field application.

SUMMARY OF PROGRESS

1. Duration of Biotreatment and Media Effects

Continuing effort at BNL in which trends in the biochemical interactions between microorganisms and crude oils are being identified has led to comparative studies of microbial effects on different types of heavy oils. These included oils which are heavy because they are immature, and oils which are heavy because they are biodegraded. We have reported (1) the effects of several thermoadapted microorganisms on Monterey crude oil which is an immature off-shore Monterey California crude. Comparative studies were carried out with another immature off-shore Monterey crude oil (A837) and an on-shore heavy crude which is biodegraded (A851). The details of these studies were recently discussed at the Fourth International Microbial Enhanced Oil Recovery Conference (2). The results are summarized in Tables 1 through 6 and Figures 1 and 2. Medium 2 and 3 were used throughout (2,3). Medium 2 contains 0.0075% of added nutrient carbon while Medium 3 contains only inorganic salts in which sulfates were replaced by chlorides. The experiments were carried out under identical experimental conditions in the presence of a detergent (e.g., tergitol) over a period of twenty days and a period of seven days. The results are consistent with previously reported observations (4,5). Biochemical interaction between the selected microorganisms and oils follow

Table 1. Monterey Oil A837 Biotreatment in Medium 3 and in the Absence of an Emulsifying Agent.

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-22	NO	Medium 3	0.489	16.875	5.0
BNL-4-23	NO	Medium 3	0.534	3.710	4.9
BNL-NZ-3	NO	Medium 3	0.482	7.265	5.7
Control, Oil	NO	Medium 3	0.486	1.205	5.0
BNL-4-22 Control	NO	Medium 3	0.000	3.550	4.9
BNL-4-23 Control	NO	Medium 3	0.000	1.825	5.0
BNL-NZ-3 Control	NO	Medium 3	0.000	2.105	5.7
Control/Control	NO	Medium 3	0.000	0.080	4.9
BNL-4-22 (65°C)	NO	Medium 3	0.496	3.565	5.5
BNL-4-23 (65°C)	NO	Medium 3	0.494	2.900	5.5
BNL-NZ-3 (65°C)	NO	Medium 3	0.589	4.515	5.5
Control, Oil (65°C)	NO	Medium 3	0.501	6.495	5.5
BNL-4-22 Control (65°C)	NO	Medium 3	0.000	3.760	5.5
BNL-4-23 Control (65°C)	NO	Medium 3	0.000	1.970	5.5
BNL-NZ-3 Control (65°C)*	NO	Medium 3	0.000	3.275	5.5
Control/Control (65°C)**	NO	Medium 3	0.000	2.915	5.5
BNL-4-22	NO	Medium 3	0.539	33.325	4.0
BNL-4-23	NO	Medium 3	0.509	35.950	4.0
Control, Oil	NO	Medium 3	0.559	39.085	4.0
BNL-4-22 Control	NO	Medium 3	0.000	14.595	4.3
BNL-4-23 Control	NO	Medium 3	0.000	5.365	4.0
Control/Control	NO	Medium 3	0.000	9.710	4.0
BNL-4-22	NO	Medium 3	0.476	40.885	5.5
BNL-4-23	NO	Medium 3	0.476	7.225	5.5
BNL-NZ-3	NO	Medium 3	0.476	5.675	5.3
Control, Oil	NO	Medium 3	0.480	5.465	5.5
BNL-4-22 Control	NO	Medium 3	0.000	9.080	5.0
BNL-4-23 Control	NO	Medium 3	0.000	3.670	5.0
BNL-NZ-3 Control	NO	Medium 3	0.000	5.105	5.5
Control/Control	NO	Medium 3	0.000	3.160	5.5

* Medium plus organism only.

** Oil plus water only

Table 2. Monterey Oil A837 Biotreatment in Medium 2 and in the Absence of an Emulsifying Agent.

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-23 Control	NO	Medium 2	0.503	380.000	5.50
BNL-NZ-3	NO	Medium 2	0.543	1700.000	4.50
A837 Control, Oil	NO	Medium 2	0.502	2000.000	4.50
BNL-4-23 Control	NO	Medium 2	0.000	79.000	5.30
BNL-NZ-3 Control	NO	Medium 2	0.000	60.500	3.70
BNL-4-23 Control	NO	Medium 2	0.000	74.500	4.50
BNL-NZ-3 Control	NO	Medium 2	0.000	65.500	3.80
BNL-4-22	NO	Medium 2	0.492	259.155	5.5
BNL-4-23	NO	Medium 2	0.483	110.245	5.5
BNL-NZ-3	NO	Medium 2	0.497	188.6	5
A837 Control	NO	Medium 2	0.486	5.19	5
BNL-4-22 Control	NO	Medium 2	0.000	342.155	5
BNL-4-23 Control	NO	Medium 2	0.000	139.235	5.1
BNL-NZ-3 Control	NO	Medium 2	0.000	6.09	5
Control/Control	NO	Medium 2	0.000	4.58	5.4

Table 3. Monterey Oil A837 Biotreatment in Medium 2 and 3 in the Presence of an Emulsifying Agent.

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-23	YES	Medium 2	0.479	2000.000	5.50
BNL-NZ-3	YES	Medium 2	0.496	1880.000	3.75
A837 Control, Oil	YES	Medium 2	0.487	255.000	3.50
BNL-4-23 Control	YES	Medium 2	0.000	95.000	4.50
BNL-NZ-3 Control	YES	Medium 2	0.000	67.000	4.30
A837 Control, Oil	YES	Medium 2	0.487	195.000	4.00
BNL-4-23 Control	YES	Medium 2	0.000	82.500	4.50
BNL-NZ-3 Control	YES	Medium 2	0.000	69.500	4.00
EMULSION AGENT		0.0075% TERGITOL			
BNL-4-22	YES	Medium 3	0.567	13.000	3.50
BNL-4-23	YES	Medium 3	0.510	9.000	3.50
Control, Oil	YES	Medium 3	0.500	7.000	3.50
BNL-4-22 Control	YES	Medium 3	0.000	26.000	3.50
BNL-4-23 Control	YES	Medium 3	0.000	15.000	3.50
Control/Control (65C)	YES	Medium 3	0.000	19.500	3.50
EMULSION AGENT		0.0075% TERGITOL			
BNL-4-22	YES	Medium 3	0.490	18.785	4.25
BNL-4-23	YES	Medium 3	0.521	10.010	4.00
Control, Oil	YES	Medium 3	0.530	6.035	4.00
BNL-4-22 Control	YES	Medium 3	0.000	51.800	4.25
BNL-4-23 Control	YES	Medium 3	0.000	61.950	4.00
Control/Control	YES	Medium 3	0.000	10.285	4.25
Control, Oil	YES	Medium 3	0.488	7.300	5.5
BNL-4-22 Control	YES	Medium 3	0.000	5.150	5.25
BNL-4-23 Control	YES	Medium 3	0.000	5.125	5.5
Control/Control	YES	Medium 3	0.000	3.465	5.5

Table 4. Biotreatment of Monterey Oil A851 in Medium 3 in the Absence of an Emulsifying Agent.

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-22	NO	Medium 3	0.505	2.500	3.5
BNL-4-23	NO	Medium 3	0.511	2.500	3.5
Control, Oil	NO	Medium 3	0.447	0.500	3.5
BNL-4-22 Control	NO	Medium 3	0.000	5.000	3.5
BNL-4-23 Control	NO	Medium 3	0.000	0.500	3.5
Control/Control	NO	Medium 3	0.000	0.000	3.5
BNL-4-22	NO	Medium 3	0.484	6.270	5.5
BNL-4-23	NO	Medium 3	0.476	3.950	5.5
BNL-4-NZ-3	NO	Medium 3	0.476	6.035	5.5
Control, Oil	NO	Medium 3	0.480	3.660	5.5
BNL-4-22 Control	NO	Medium 3	0.000	3.500	5.5
BNL-4-23 Control	NO	Medium 3	0.000	2.815	5.5
BNL-NZ-3 Control	NO	Medium 3	0.000	0.505	5.5
Control/Control	NO	Medium 3	0.000	0.031	5.3
BNL-4-22	NO	Medium 3	0.484	6.470	6.0
BNL-4-23	NO	Medium 3	0.476	3.850	5.9
BNL-NZ-3	NO	Medium 3	0.476	4.500	6.0
Control, Oil	NO	Medium 3	0.480	2.250	6.0
BNL-4-22 Control	NO	Medium 3	0.000	3.065	5.8
BNL-4-23 Control	NO	Medium 3	0.000	2.175	5.9
BNL-NZ-3 Control	NO	Medium 3	0.000	3.710	5.9
Control/Control	NO	Medium 3	0.000	0.185	6.0
AVERAGE OF 7 DAYS					
BNL-4-22	NO	Medium 3	0.491	5.508	5.0
BNL-4-23	NO	Medium 3	0.488	3.265	5.0
BNL-NZ-3	NO	Medium 3	0.476	5.275	5.8
Control Oil	NO	Medium 3	0.469	2.135	5.0
BNL-4-22 Control	NO	Medium 3	0.000	3.550	4.9
BNL-4-23 Control	NO	Medium 3	0.000	1.825	5.0
BNL-NZ-3 Control	NO	Medium 3	0.000	2.105	5.7
Control/Control	NO	Medium 3	0.000	0.080	4.9

Table 4. (Continued).

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-21 (65C)	NO	Medium 3	0.628	3.355	5.5
BNL-4-22 (65C)	NO	Medium 3	0.628	4.220	5.5
BNL-4-23 (65C)	NO	Medium 3	0.625	3.435	5.5
BNL-NZ-3 (65C)	NO	Medium 3	0.629	4.020	5.5
Control, Oil (65C)	NO	Medium 3	0.626	3.615	5.5
BNL-4-21 Control (65C)	NO	Medium 3	0.628	1.160	5.5
BNL-4-22 Control (65C)	NO	Medium 3	0.000	3.760	5.5
BNL-4-23 Control (65C)	NO	Medium 3	0.000	1.970	5.5
BNL-NZ-3 Control (65C)	NO	Medium 3	0.000	3.275	5.5
Control/Control (65C)	NO	Medium 3	0.000	2.915	5.5
BNL-6-32 (65C)	NO	Medium 949	0.476	20.310	5.3
Control, Oil (65C)	NO	Medium 949	0.480	8.530	4.5
BNL-6-32 Control (65C)	NO	Medium 949	0.000	28.265	4.9
Control/Control (65C)	NO	Medium 949	0.000	26.170	4.4
BNL-4-22	NO	Medium 3	0.477	2.335	4.8
BNL-4-23	NO	Medium 3	0.481	4.015	4.3
Control, Oil	NO	Medium 3	0.493	4.830	4.5
BNL-4-22 Control	NO	Medium 3	0.000	14.595	4.3
BNL-4-23 Control	NO	Medium 3	0.000	5.365	4.0
Control/Control	NO	Medium 3	0.000	9.710	4.0
BNL-4-22	NO	Medium 3	0.484	12.605	5.5
BNL-4-23	NO	Medium 3	0.484	14.205	5.5
BNL-NZ-3	NO	Medium 3	0.476	6.720	5.0
Control, Oil	NO	Medium 3	0.480	1.070	5.5
BNL-4-22 Control	NO	Medium 3	0.000	9.080	5.0
BNL-4-23 Control	NO	Medium 3	0.000	3.670	5.0
BNL-NZ-3 Control	NO	Medium 3	0.000	5.105	5.5
Control/Control	NO	Medium 3	0.000	3.160	5.5
BNL-4-22 Control	NO	Medium 3	0.000	5.105	5.0

Table 5. Biotreatment of Monterey Oil A851 in Medium 2 in the Absence of an Emulsifying Agent.

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-23	NO	Medium 2	0.537	2000.000	5.50
BNL-NZ-3	NO	Medium 2	0.490	2000.000	4.50
A851 Control, Oil	NO	Medium 2	0.516	1280.000	5.25
BNL-4-23 Control	NO	Medium 2	0.000	79.000	5.30
BNL-NZ-3 Control	NO	Medium 2	0.000	60.500	3.70
BNL-4-23 Control	NO	Medium 2	0.000	74.500	4.50
BNL-NZ-3 Control	NO	Medium 2	0.000	65.500	3.80
BNL-4-22	NO	Medium 2	0.482	282.310	5.00
BNL-4-23	NO	Medium 2	0.480	155.935	5.50
BNL-NZ-3	NO	Medium 2	0.479	130.510	5.50
A836 Control	NO	Medium 2	0.494	5.900	5.00
BNL-4-22 Control	NO	Medium 2	0.000	309.785	5.50
BNL-4-23 Control	NO	Medium 2	0.000	183.655	5.50
BNL-NZ-3 Control	NO	Medium 2	0.000	111.665	5.00
Control/Control	NO	Medium 2	0.000	1.685	5.00
BNL-4-22	NO	Medium 2	0.478	407.525	4.50
BNL-4-23	NO	Medium 2	0.476	116.25	5.50
BNL-NZ-3	NO	Medium 2	0.490	261.07	4.00
A836 Control	NO	Medium 2	0.484	60.98	5.50
BNL-4-22 Control	NO	Medium 2	0.000	342.155	5.00
BNL-4-23 Control	NO	Medium 2	0.000	139.235	5.10
BNL-NZ-3 Control	NO	Medium 2	0.000	6.09	5.00
Control/Control	NO	Medium 2	0.000	4.58	5.40

Table 6. Biotreatment of Monterey Oil A851 in Medium 2 and 3 in the Presence of an Emulsifying Agent.

Microorganisms	Emulsion Agent	Medium	%Oil Incubation (Days)	Extent of Emulsification in Klett Units	pH
BNL-4-23	YES	Medium 2	0.490	940.000	5.3
BNL-NZ-3	YES	Medium 2	0.511	1360.000	3.5
A851 Control	YES	Medium 2	0.494	1380.000	5.0
BNL-4-23 Control	YES	Medium 2	0.000	95.000	4.5
BNL-NZ-3 Control	YES	Medium 2	0.000	67.000	4.3
BNL-4-22 Control	YES	Medium 3	0.510	10.900	4.0
BNL-4-23 Control	YES	Medium 3	0.517	13.900	4.25
Control, Oil	YES	Medium 3	0.488	5.450	4.0
BNL-4-22 Control	YES	Medium 3	0.000	51.800	4.25
BNL-4-23 Control	YES	Medium 3	0.000	61.950	4.0
Control/Control	YES	Medium 3	0.000	10.285	4.25
Control, Oil	YES	Medium 3	0.480	58.425	5.25
BNL-4-22 Control	YES	Medium 3	0.000	5.150	5.25
BNL-4-23 Control	YES	Medium 3	0.000	5.125	5.5
Control/Control	YES	Medium 3	0.000	3.465	5.5
BNL-4-22	YES	Medium 3	0.504	19.000	3.50
BNL-4-23	YES	Medium 3	0.505	12.500	3.50
Control, Oil	YES	Medium 3	0.485	11.000	3.50
BNL-4-22 Control	YES	Medium 3	0.000	26.000	3.50
BNL-4-23 Control	YES	Medium 3	0.000	15.000	3.50
Control/Control	YES	Medium 3	0.000	19.500	3.50

Figure 1. GC-MS Fragmentograms, m/z 57, Monterey, CA, crude A837 treated with BNL-4-22, BNL-4-23, and BNL-NZ-3.

Figure 2. GC-MS Fragmentograms, m/z 57, Monterey, CA, crude A831 (1) Treated with BNL-4-22, BNL-4-23, and BNL-NZ-3 and (2) Control and pyrolyzed control and BNL-NZ-3 at 640°C.

distinct patterns. Thus, treatment of Monterey A837 with BNL-4-22, BNL-4-23, and BNL-NZ-3 leads to the conversion of heavier hydrocarbon fractions of crude oil to lighter. Furthermore, there are differences in the effects due to the presence or absence of tergitol, a C₂₈ hydrocarbon based surface active agent. For all cases in this series of experiments, a seven-day treatment appeared to be sufficient to bring about the biochemical conversion. The working hypothesis is based on the premise that the biochemical conversion starts by an attack on the complex structure, possibly on the N,S,O rich polar components of the maltene fraction derived from the asphaltene component of the crude (3). In the three crude oils considered, A836, A837, and A851, the highest concentration of this fraction is in A851. The overall redistribution of the relative amounts of heavy and light hydrocarbons is shown in Figure 1 and 2. The results also show that BNL-4-22 and BNL-4-23 are not very effective in the biotreatment of A851. BNL-NZ-3 on the other hand, is remarkably effective. Further biochemical evidence which supports a mechanism or mechanisms by which microorganisms may interact with crude oils by converting the complex N, S, O structure is offered by a comparative chromatographic analysis of organosulfur compounds present in Monterey A836, A837, and A851 before and after biotreatment. The extent of bioconversion of the crude oils varies with different microorganisms used. Biotreatment of Monterey A836 with BNL-4-22 and BNL-4-23 results in changes of organosulfur composition as reported earlier (see Premuzic and Lin, Quarterly Report July, 1992). These changes differ to those due to the biotreatment of A836 with BNL-NZ-3 as shown in Figure 3. Relative to control there is a compositional change without an overall decrease in organic sulfur content. However, biotreatment of Monterey A837 with BNL-4-22, BNL-4-23, and BNL-NZ-3 under identical experimental conditions results in a significant overall decrease in organic sulfur content as shown in Figures 4 and 5. In this series of experiments, the Monterey A851 an on-shore highly biodegraded California crude oil was considered a test case in terms of differentiation of "biotreated and "biodegraded" crude oil. Biotreatment of this crude oil further indicated the importance of "diagnostic" evaluation of the biotreatment i.e., an initial small scale evaluation of the microorganism potential for MEOR. All the experiments referred to in this discussion were run under identical experimental conditions for sample treatment, concentration, and chromatographic conditions. Thus, treatment of A851 with BNL-4-22 and BNL-4-23 had no effect as shown in Figures 2 and 6. However, treatment of A851 with BNL-NZ-3 resulted in major changes in the composition of A851 as shown in Figures 2 and 7. This thermophilic isolate breaks down A851 into a much more volatile hydrocarbon mixture (see Figure 2) with a significant increase in the lower molecular weight (< C₃₀) hydrocarbons. The FPD gas chromatographic trace shown in Figure 7 is consistent and supports further the proposed mechanism(s) that the initial biochemical reactions may occur in the heavy N, S, O portion of asphaltenes (Premuzic et al., 1992). A qualitative test for crude is offered by a comparison of the two FPD scans in Figures 7. In order to plot the FPD traces, in a manner consistent with previous analyze, the sensitivity of the chromatograph (i.e., attenuation) was set at 32. At that setting, the response was off-scale (Figure 7(1)) and the sensitivity had to be reduced to 64 or by 50% (Figure 7(2)) to get a full response on that scale. This chromatographic analysis result, indicates that the biotreatment causes a substantial breakdown and conversion of the complex structures associated with heavy crude oils.

Figure 3. Gas Chromatographic trace with Flame Photoemission Detector (FPD) of,
 A: Monterey A336 control; B: Monterey A337 treated with BNL-NZ-3.
 Medium 3, seven days, no tergitol.

Figure 4. Gas Chromatographic trace with Flame Photoemission Detector (FPD) of, A: Monterey A336 control; B: Monterey A337 treated with BNL-4-22; C: Monterey A337 treated with BNL-NZ-3. Medium 3, seven days, no tergitol.

Figure 5. Gas Chromatographic trace with Flame Photoemission Detector (FPD) of, A: Monterey A337 control; B: Monterey A337 treated with BNL-NZ-3. Medium 3, seven days, no tergitol.

Figure 6. Gas Chromatographic trace with Flame Photoemission Detector (FPD) of,
 A: Control; B: Monterey A851 + BNL-4-22; C: Monterey A851 + BNL-4-23.
 Medium 3, seven days, no tergitol.

Figure 7. Gas Chromatographic trace with Flame Photoemission Detector (FPD) of Monterey A851 crude treated with BNL-NZ-3. Medium 3, seven days, no tergitol. Figure 7(1) shows a scan at attenuation of 32 and 7(2) at attenuation 64 (see text for discussion).

Pyrolysis of A851 at 640°C produced a volatile fraction with some enrichment in the lighter ends of the crude, however, not very effective when compared to biotreatment. Pyrolysis at 640°C of the BNL-NZ-3 treated oil produced a hydrocarbon product whose GC-MS fragmentogram compares to the non-pyrolyzed one (Figure 2), however, with a loss of very lighter fraction in the 1200-1800 (C₁₀ - C₁₈) scan region. The FPD scans, Figure 8, are consistent with the GC-MS results.

Comparative experiments are continuing. Analysis by XANES spectrometry has also been expanded to a number of other oils.

2. Construction of Core-flooding Systems

The first generation of a core-flooding apparatus has been designed and constructed. The core bioreactor was constructed from a 304 stainless 1-1/4" x 12" tube with a thickness of 0.065 inches, and can be operated up to a pressure of 2600 psi. The tube can be treated under controlled conditions and various pressures. Cores made of Berea sandstone cut to appropriate size are introduced into the tube and a tight fit achieved by means of an epoxy treatment of the core. Several core-flooding experiments are underway. Details of the construction, the flow diagram and preliminary results will be reported upon completion of this series of experiments.

3. Microscopic Comparison of Reaction Mixtures (RMs)

Earlier we have discussed that the effects on crude oils due to the "biotreatment" of the crude oils as described in preceding paragraphs are different from the effects due to biodegradation which may occur in the reservoir over geological periods of time. Results obtained from comparative studies of interaction between different crude oils and microorganisms deliberately introduced for the purposes of biotreatment suggested that the process(es) may be driven by the introduced and not indigenous microorganisms. Concurrent with chemical analysis, a microscopic analysis of RMs was initiated. Preliminary results from these analyses obtained for two Monterey and an Arkansas crude oil are given in Figures 9, 10, and 11. Note, in all three sets of experiments, Medium 2 was used (Premuzic et al., 1992), i.e., a medium which contains 0.0075% of added organic nutrient (other than the crude oil). This was done deliberately to encourage the growth of indigenous microorganisms. In all cases, the microscopic characteristics change drastically upon the introduction of the test organisms. Continuing comparative and microscopic studies should ultimately lead to a better understanding of the reaction mixtures (RMs, see refs.) components and their influence on the formation of products.

Figure 8. Chromatographic trace with Flame Photoemission Detector (FPD) of Monterey A851 crude treated with BNL-NZ-3 and then analyzed by Pyrolysis-Gas Chromatography (PY-GC). Medium 3, seven days, no tergitol.

Figure 9. Microscopic images of control and biotreated Arkansas crude oil. Medium 2, no tergitol, seven days.

Figure 10. Microscopic images of control and biotreated Monterey A836 crude oil. Medium 2, no tergitol, seven days.

Figure 11. Microscopic images of control and biotreated Monterey A837 crude oil. Medium 2, no tergitol, seven days.

REFERENCES

1. Premuzic, E. T., and Lin, M.S. Effects of selected thermophilic microorganisms on crude oils at elevated temperatures and pressures. Quarterly Report (01/01/92-03/30/92) BNL 47555, May 1992.
2. Premuzic, E. T., and Lin, M. S. Prospects for thermophilic microorganisms in microbial enhanced oil recovery (MEOR). Proc. 1990 International Conference on Microbial Enhancement of Oil Recovery - Recent Advances, Norman, May 1990, E. C. Donaldson, Editor, Developments in Petroleum Science, Vol. 31, Chapter R-18, pp. 277-96, Elsevier Sci. Publ., Amsterdam 1991. (BNL 44684).
3. Premuzic, E. T., and Lin, M. S. Prospects for thermophilic microorganisms in microbial enhanced oil recovery (MEOR)--Part II. Proc. International Symposium on Oil Field Chemistry, Feb. 1991, SPE 21015, pp. 143-48, Society of Petroleum Engineers, Anaheim 1991. (BNL 45650).
4. Zajic, J. E. and Mahomedy, A. Y. Biosynthesis of surface active agents. R. M. Atlas, Editor, Petroleum Microbiology, Chapter 6, pp. 221-97, Macmillan Publ. Co., New York 1984.
5. E. T. Premuzic, Lin, M. S., Racaniello, L. K., and B. Manowitz. "Chemical Markers of Induced Microbial Transformation in Crude Oils," Fourth International Conference on Microbial Enhanced Oil Recovery, eds. E. T. Premuzic and A. Woodward (Elsevier Publishing Company, Amsterdam, 1992) in preparation.

END

**DATE
FILMED**

4 / 30 / 93