Final Progress Report: Gene Transcription and Electromagnetic Fields
DE-FG01-89CE34023

Prepared by: Ann S. Henderson, Ph.D., Hunter College.

DOE/CE/34023--T1

DE93 010854

Rationale. Our research has focused on how cells are affected by electromagnetic field exposure. This area of research is particularly important in view of the increasing public awareness that extremely low frequency (elf) electric and/or magnetic fields (EM fields), in both the home and in the work place, may have detrimental effects on human health. Despite many epidemiological studies implicating exposure to EM fields in human cancer, and an substantial body of basic research designed to understand the interaction of these fields with cells and tissues, no plausible interaction mechanism(s) between the cell and EM fields has emerged. This is due to a lack of information regarding which specific parameters of the field the cell is responding to during exposure. The result is both skepticism concerning possible health risks from EM fields and inadequate funding for research at the cell, animal or human levels.

Hypothesis. Our overall aim is to obtain sufficient information to allow us to ultimately determine whether ELF EM field exposure is an initiating factor in neoplastic transformation and/or if exposure can mimic characteristics of the "second-step" counterpart in neoplastic disease. This aim is based on our previous findings that levels of some transcripts are increased in cells exposed to EM fields. While the research is basic in nature, the ramifications have bearing on the general safety of exposure to EM fields in industrial and everyday life.

A large array of diverse biological effects are reported to occur as the result of exposure to elf EM fields, suggesting that the cell response to EM fields is at a basic level, presumably initiated by molecular and/or biophysical events at the cell membrane ^(rev. in 1,2,11,12). The hypothesized route is a signal transduction pathway involving membrane calcium fluxes. Information flow resulting from signal transduction can mediate the induction of regulatory factors in the cell, and directly affect how transcription is regulated.

Research. The focus of our research since 1983 has been to determine whether exposure of cells to extremely low frequency (elf) electric and magnetic (EM) fields [<150 Hz, <570 μ T (rms)] affects transcription and translation. We have concentrated on fields at 60 Hz, initiating experiments with B-fields that are about 2x ambient fields [B (magnetic) field = 0.57 μ T (rms); E (electric) field = ~ 1 μ V/m] in the laboratory, and using log increments in the E and B-fields up to approximately 2000x. The most effective exposure condition, relative to transcript increase, is 20X ambient laboratory levels using exposure time of 20 minutes. The majority of the experiments have used dipteran salivary gland cells and human HL-60 cells, but some experimental data has been obtained from yeast, as well as other human cells lines, including HeLa and HAL cells. The use of dipteran salivary gland cells in conjunction with transcription autoradiography allows us to determine the location of nascent RNA chains directly on the giant polytene chromosomes. This strongly suggest that one effect of elf EM fields is directly at the

MASTER
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

js

transcriptional level, rather than in the processing or storage of mRNAs or increased RNA stability^(4-6,9). Short exposures (less than 60 mins) caused a change in the expected pattern of transcription over specific regions of salivary gland chromosomes. These studies have been extended to map specific regions of the salivary gland chromosomes in *Drosophila* where known genes are present. Other research has used two-dimensional gel electrophoresis to determine that transcriptional changes are reflected in altered polypeptide synthetic patterns ^(manuscript in preparation).

The exposure of human cultured cells to elf EM fields also induces an increase in some mRNA levels^(1,2,5,7,8,12). The magnitude of the response, however, is dependent on the signal used. The presence of increased transcript levels in cells exposed to EM fields has been confirmed in reports by Czerska *et al.* in Daudi cells and by Phillips *et al.* in CEM-CM3 human T-lymphoblastoid cells. An increase in *c-myc* transcript levels following short exposures to EM fields has also been shown to be coordinate with an increase in intracellular calcium in thymocytes stimulated with Con A by Liburdy *et al.*

Experiments in our laboratories show that the response of the cell to elf EM fields is not due to some unexplained general activation of transcription. Increased transcript levels were not observed for all genes tested. As expected, in experiments using HL-60 cells, the level of α -globin transcripts was not significant in either exposed or unexposed cells⁽¹⁾. This gene is not normally expressed in HL-60 cells, but these experiments allowed us to rule out the outside possibility that an unexpressed gene would be affected by EM fields. No hybridization on Northern blots was detected in the expected region of migration of globin mRNA. In other experiments, β 2-microglobulin cDNA was used as a probe. This gene is expressed in HL-60 cells, but is unaffected by agents normally used for induction of differentiation of HL-60 cells. Levels of β 2-microglobulin transcripts were unaltered by elf EM field exposure⁽⁵⁾. Phillips *et al.* also showed that there was no effect on transcription of a group of genes. One possible conclusion is that the response of cells to EM fields is restricted to a subset (presently unidentified) of genes normally expressed in the cells.

Recent findings from our laboratories on increased transcripts in cells exposed to EM fields are summarized below (see attached reprints).

►(1) Some RNA transcripts in cells are increased in the presence of 60 Hz continuous wave sinusoidal fields with increases that are magnetic field and frequency dependent ^(1,2,7,8). These experiments initially used "dot blot" hybridizations in concert with Northern blots. More recently, we have used quantitative RT-PCR to confirm that increased transcripts for *c-myc*, *c-fos*, *c-jun*, and p53 are present in cells exposed to 60 Hz EM fields ^(16 and in manuscript).

►(2) The response of the cell is modulated at specific field strengths ^(5,7,8), exposure times ⁽⁸⁾, and culture conditions⁽⁸⁾. The effect on the cell is not related to either time or field strength; *i.e.*, there is no obvious dose effect.

►(3) Four minutes is the earliest time point after exposure that an increase in *c-myc*

transcripts can be measured^(7,8). This rapid response is in agreement with proposed mechanisms that EM field effects are initiated at the cell surface.

►(4) Cells can adapt and/or accommodate to the presence of EM fields⁽⁸⁾. The transcription of *c-myc* and *c-fos* is rapidly modified in many cells responding to various types of induction. It is known that induction of *c-fos* can be observed within 5 minutes after treatment with serum; the transcriptional activation peaks at about 15 minutes and decreases to basal levels within 2 hours. This is the type of pattern that we have measured following exposure of cells to elf EM fields. As an example, at the field strengths used in many of our experiments (8 or 80 μ T), an increase in *c-myc* transcripts above control levels can be detected as early as 4 to 8 minutes, with a peak at 20 minutes. If exposures are continued, however, the transcript levels approach those of the controls at about 2 hours. The effect is the same when cells are exposed for 20 minutes and then removed from the field. Thus, the response is typical of induction—an initial response is observed, but then a return to normal levels occurs at a time characteristic of the type of induction, and the gene involved. Preliminary data, however, show that exposing the cells to a different field (57 μ T rather than 5.7 μ T magnetic fields) after about 2 hours cause the transient increase in transcript levels to reoccur. These data suggest that experimental strategies used in studying induction in general can be applied to studies which pursue the mechanism of interaction of cells with elf EM fields. EM field effects are within the time frame for those observed for other types of induction (*e.g.*, heat shock or TPA).

►(5) Although hsp70 mRNA is increased (as possibly other heat shock proteins on the basis of 2-D gel electrophoresis) in the presence of EM fields, the reaction is not heat shock *per se* ^(manuscript in preparation). There is no increase in temperature in the media during exposures, and none expected at the cellular level based on exposure geometry. There is over-expression of the gene hsp70 in all cells (yeast, *Drosophila*, human) exposed to EM fields; the peak in transcript level for hsp70 invariably coincides with transcript levels for *c-myc*. [The research in human cells was prompted by observations from transcription autoradiography of *Drosophila* salivary gland chromosomes. Analysis of chromosome transcription showed increased autoradiographic labelling of several heat shock genes, including hsp70, in isolated glands exposed to elf EM fields ^(4,5,13)].

►(6) The cell membrane is important in the response to EM fields⁽¹⁶⁾. Our experiments have tested whether plasma membrane fluctuations influence the response of the cell to EM fields as measured by changes in the levels of β -actin and *c-fos* transcripts. Treatment of cells with colchicine is known to disturb microtubules and decrease plasma membrane fluctuations. Cytochalasin treatment influences microfilament structure and increases local membrane fluctuations. HL-60 cells were exposed to 60 Hz EM fields in the presence of either colchicine or cytochalasin B and transcript levels for β -actin and *c-fos* were determined by RT-PCR. Treatment of cells with colchicine or cytochalasin B produce opposite effects in the response of the cell to electromagnetic field exposure. The stimulatory effect of EM fields on the expression of β -actin and *c-fos* is abolished in the presence of colchicine. The levels for both β -actin and *c-fos*, however, are enhanced in the presence of cytochalasin B. Based on this data, we conclude that there is a

correlation between the amplitude of plasma membrane fluctuations and the response of the cell to electromagnetic field exposure.

►(7) In preliminary experiments CAT expression is increased in cells exposed to EM fields following transfection of portions of the *c-myc* promoter linked to a CAT reporter gene (manuscript in preparation). These experiments were designed to determine if the *c-myc* promoter region *per se* was responsive to EM fields. *C-myc* promoter regions were transfected into both mouse and human cells as CAT constructs. The presence of the 2.3 kb domain upstream of the human gene results in increased expression of CAT following 20 minute exposure of cells to 60 Hz elf EM fields. The transfection of a similar region from the mouse *c-myc* promoter showed increased expression in mouse cells. Specific portions of the promoter region were deleted from the human promoter construct and introduced into cells to determine the responsive region. The region responsive to EM fields is less than 800 bp and is located between -353 and -1257 relative to the P₁ promoter. The experiments will be continued to identify a more precise region.

Publications (since 1991)

1. Goodman, R. and A. Henderson, 1991. Transcription in cells exposed to extremely low frequency electromagnetic fields: a review. *Bioelectrochem. Bioenerget* 25, 335-355.
2. Goodman, R., J. Bumann, L-X Wei, J-C Xu and A. Shirley-Henderson, 1991. Transcriptional changes in cells exposed to extremely low frequency electromagnetic fields. *In Electromagnetics in Biology and Medicine* (Carl T. Brighton and Solomon R. Pollack, eds), pp. 127-132, San Francisco Press, San Francisco.
3. Hubbard-Smith, K., P. Patsalis, J. Pardinias, J. Jha, A. Henderson and Ozer, H., 1992. Altered chromosome 6 in immortal human fibroblasts. *Mol. Cell Biol.*, 12, 2273-2281.
4. Goodman, R., D. Weisbrot, A. Uluc and A. Henderson, 1992. Transcription in *Drosophila melanogaster* salivary gland cells is altered following exposure to low frequency electromagnetic fields: Analysis of chromosome 3R. *Bioelectromagnetics* 13: 111-118.
5. Blank M., L. Soo, H. Lin, A. Henderson and R. Goodman, 1992. Changes in transcription in HL-60 cells following exposure to alternating current from electric fields. *Bioelectrochem. Bioenerget.* 28:301-309.
6. Goodman, R., D. Weisbrot, A. Uluc and A. Henderson, 1992 Transcription in *Drosophila melanogaster* salivary gland cells is altered following exposure to low frequency electromagnetic fields: Analysis of chromosomes 3L and X. *Bioelectrochem. Bioenerget.*, 28: 311-318.
7. Goodman, R., J. Bumann, L-X Wei and A. Shirley-Henderson, 1992. Exposure of human cells to electromagnetic fields: Effect of time and field strength on transcript levels. *Electro- and Magnetobiology* 11: 19-28.

8. Goodman, R., L-X Wei, J. Bumann and Ann S. Henderson, 1992. Exposures to electric and magnetic (EM) fields increases transcripts in HL-60 cells; does adaptation to EM fields occur? *Bioelectrochem Bioenergetics* 29: 185-192.
9. Goodman, R., Weisbrot, D., Uluc, A. and Henderson, A. (1992) Transcription in *Drosophila melanogaster* salivary gland cells is altered following exposure to low frequency electric and magnetic fields: analysis of chromosomes 2R and 2L. *Bioelectrochem. Bioenerget.*, in press.
10. Gold, S., R. Goodman and A. Henderson, 1992. Exposure of human cells to electric and magnetic fields results in increased levels of T-antigen mRNA and protein. *Bioelectromagnetics*, in press.
11. Reba Goodman, Yuri Chizmadzhev and Ann Shirley-Henderson, 1993. Electromagnetic fields and cells. *J. Cellular Biochemistry* 52: 1-6.
12. Goodman, R. and Henderson, A., 1992. Effects of electric and magnetic fields on gene expression. *IN Biological effects of electromagnetic fields* (D. Carpenter, ed.) Academic Press, Orlando, FLA, in press
13. Weisbrot, D.R., O. Khorkova, H.Lin, A.Henderson and Reba Goodman, 1992. The effect of low frequency electric and magnetic fields on gene expression in *Saccharomyces cerevisiae*. *Bioelectrochem. Bioenerget.*, in press.
14. Hadjiargyrou, M., K. McDowell and A. Henderson, 1993. A transfected human ribosomal RNA gene is active in nucleolus formation in human cells. Submitted to *Am. J. Hum. Genetics*.
15. McDowell, K., M. Hadjiargyrou, P. Patsalis and A. Henderson, 1993. A partial human rDNA repeat functions as a nucleolar organizer region. Submitted to *Am. J. Hum. Genetics*.
16. Broude, N., R. Karabakhtsian, N. Shalts, R. Goodman and A. S. Henderson, 1993. Response of human HL60 cells to electromagnetic fields is altered by treatment with colchicine and cytochalasin B. *Bioelect. Bioenerget.* , in press

Abstracts and Presentations:

- a. R. Goodman, L-X Wei and A. Henderson, 1991. Relationship of fields strength and time to induction of increased transcripts by elf EM fields. RCMI Symposium, Atlanta.
- b. R. Goodman and A. Henderson, 1991. Transcriptional changes in cells exposed to extremely low frequency electromagnetic fields. FASEB, Atlanta,
- c. Goodman, R., J. Bumann and A. Henderson (1991) Defining signal parameters which induce transcriptional/translational alterations in HL-60 cells. Presented at 13th Ann Bioelectromagnetics Soc. Salt Lake City, B-2-3.

- d. Patsalis, P., K. Hubbard-Smith, J. Pardinas, K. Jha, A. Henderson and H. Ozer, 1991. Deletion of 6q is involved in immortalization of SV-40 transformed human fibroblasts. 8th International Congress of Human Genetics, Washington.
- e. Goodman, R., L-X Wei, S. Gold and A. Henderson, 1991. Transcriptional and translational alterations in human cells are induced by low frequency electromagnetic fields: Relationship to time and exposure parameters. American Society of Cell Biology, Boston.
- f. Goodman, R., L-X Wei and A. Henderson, 1991. An increase in mRNA and protein levels is induced in human cells exposed to electromagnetic fields: relationship to time of exposure, post-exposure time and frequency. DOE/Contractor's Review. Milwaukee, WI, A-18.
- g. Camargo, E., S. Ogiste, R. Goodman and A. Henderson, 1991. Effect of 60 Hz EM fields on cell division in human HL-60 cells. RCMI symp, Washington, 1991.
- h. Goodman, R., L-X Wei, J. Bumann and A. Henderson, 1991. Relationship of field strength and time on increased levels of transcripts induced by elf EM fields. DOE/Contractor's Review. Milwaukee, WI,
- i. Goodman, R., L-X Wei, S. Gold and A. Henderson, 1992. An increase in messenger RNA and protein levels is induced in cells exposed to electromagnetic fields: the relationship of time and exposure conditions. World Congress, Bioelectromagnetics, Orlando.
- j. Goodman, R., J. Bumann, L-X Wei and A. Henderson, 1991. Increased transcripts in HL-60 cells exposed to EM fields: comparison of exposure geometries. The Ann. Rev. of Research on Biological Effects on 50 and 60 Hz Electric and Magnetic Fields, Milwaukee, WI, P-13.
- k. Bumann, J., R. Goodman and A. Henderson, 1991. Die Wirkung Niederfrequenter Elektromagnetischer Wellen auf die Gentranskription, Stuttgart
- l. Blank, M., L. Soo, H. Lin, A. Henderson and R. Goodman, 1991. Stimulation of transcription in HL-60 cells by alternating currents from electric fields. DOE/Contractors Review, Milwaukee, WI, A-19
- m. Goodman, R., L-X Wei and A. Henderson, 1992. An increase in mRNA and protein levels is induced in human cells exposed to electromagnetic fields: Relationship to time and exposure conditions. First World Congress for Electricity and Magnetism in Biology and Medicine, Orlando, FL.
- n. Goodman, R., L-X Wei and A. Henderson, 1992. Changes in translational patterns are induced in human HL-60 cells at specific times and field strengths. First World Congress for Electricity and Magnetism in Biology and Medicine, Orlando, FL.
- o. Goodman, R., O. Khorkova, A. Henderson and D. Weisbrot, 1992. The effect of low frequency electric and magnetic fields on gene expression in *Saccharomyces cerevisiae*. DOE Contractor's Meeting in press.
- p. Goodman, R. M. Blank, O. Khorkova, D. Weisbrot, and A. Henderson, 1992. The heat shock hsp70 gene is induced in a variety of cell types exposed to elf EM fields or ac current. Does the cell respond to EM field exposure as cellular stress? DOE Contractor's Meeting, on press
- q. Gold, S., R. Goodman and A. Henderson, 1992. Increased levels of T-antigen mRNA and protein in human cells exposed to 60 Hz sinusoidal field. DOE Contractor's Meeting, in press.

r. H. Lin, R. Goodman and A. Henderson. EM field exposure affects the *c-myc* promotor. DOE Contractor's Meeting, in press
s. Patsalis, P., H. Ozer and A. Henderson, 1992. A breakpoint on chromosome 6q is involved in cell immortalization. Chromosomes in Cancer, Phoenix, Az.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

DATE
FILMED

5 / 21 / 93