Shemya AFB, Alaska 1992 IRP FIELD INVESTIGATION REPORT

Volume 3 of 4
APPENDIXES B, C, AND D

FINAL

February 1993

Prepared for

U.S. Air Force

Elmendorf AFB, Alaska

11th Air Control Wing

11th Civil Engineering Operations Squadron

Under Contract DEU-91-06

Prepared by

CH2M HILL

P.O. Box 8748

Boise, Idaho 83707

For

Environmental Management Operations

Under a Related Services Agreement

with the U.S. Department of Energy

Environmental Management Operations

Richland, Washington 99352

MASTER

CONTENTS

APPENDIX B - GEOPHYSICAL CONTOUR MAPS AND PROFILE PLOTS	B.1
INTRODUCTION	B.1
CONTOUR MAPS AND PROFILE PLOTS	
APPENDIX C - HUMAN HEALTH RISK ASSESSMENT-PRELIMINARY RISK LEVELS	C.1
INTRODUCTION	
SELECTION OF CHEMICALS OF POTENTIAL CONCERN	
EXPOSURE ASSESSMENT	
DEVELOPMENT OF PRELIMINARY RISK LEVELS	
RISK CHARACTERIZATION	
REFERENCES	
APPENDIX D - ECOLOGICAL RISK ASSESSMENT	D.1
INTRODUCTION	
IDENTIFICATION OF ECOLOGICAL CHEMICALS OF POTENTIAL CONCERN	
ECOLOGICAL EXPOSURE ASSESSMENT	
DEVELOPMENT OF PRELIMINARY REMEDIATION GOALS	
ECOLOGICAL TOXICITY ASSESSMENT	
ECOLOGICAL RISK CHARACTERIZATION	
UNCERTAINTY ANALYSIS	
REFERENCES	

CONTENTS

APPENDIX B - GEOPHYSICAL CONTOUR MAPS AND PROFILE PLOTS	B.1
INTRODUCTION	B.1
CONTOUR MAPS AND PROFILE PLOTS	

Appendix B
Geophysical Contour Maps and Profile Plots

CONTENTS

APPENDIX B - GEOPHYSICAL CONTOUR MAPS AND PROFILE PLOTS	B.1
INTRODUCTION	B.1
CONTOUR MAPS AND PROFILE PLOTS	

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Appendix B

Geophysical Contour Maps and Profile Plots

Introduction

Organization of Technical Memorandum

This technical memorandum (TM) summarizes the geophysical investigations performed at Shemya Air Force Base as part of the Installation Restoration Program. The geophysical investigations consisted of electromagnetic (EM) ground conductivity surveys performed over 16 different areas at the base. The areas investigated were the following:

- FT-1 Lightning Strike Fire Training Area
- FT-2 Aircraft Mock-up Fire Training Site
- FT-3 Structural Fire Training Area
- PS-1B Old Cobra Dane
- PS-6 Refueling Vehicle Maintenance Shop
- PS-7 Vehicle Maintenance
- PS-8 Old White Alice
- PS-9 Asphaltic Drum Storage Area
- Water Gallery
- SW-4 Old Barrel Storage Area
- SW-5 Hospital Lake
- SW-10 Barrel Bay
- SW-12 Scrap Metal Dump
- SW-14 Scrap Metal Landfill

In addition to the above listed sites, reconnaissance surveys were performed at the following locations:

- PS-3 West End Oil/Water Separator Ponds
- PS-83 Old Bulk Fuel Storage Area

This TM describes the procedures and equipment used in the acquisition of the data and describes data handling and interpretation procedures.

This TM is organized into two sections. The remainder of this introductory section presents the purpose and scope of the investigation. The remaining section presents the data handling and interpretation procedures.

Purpose and Scope

The EM conductivity surveys were performed to aid in locating buried metal, wastes, or utilities at each of the sites investigated. The data will be used to help plan locations for monitoring wells by identifying potential problem areas for drilling as well as by delineating the extent of buried wastes or fill. Detailed surveys were performed over all sites except for PS-83, the old fuel storage area, and PS-3, the west end oil separation ponds where reconnaissance surveys were performed. Detailed surveys consisted of recording measurements at known locations and subsequently plotting and analyzing the data. The reconnaissance surveys were performed by observed conductivity values directly on the instrument while walking along the traverses. The reconnaissance investigations were performed to provide a basis for planning future work at the area but not to actually delineate any waste areas at the time.

Procedures

Data Acquisition

The EM conductivity investigations were performed between July 15 and July 29, 1992. Detailed surveys were performed over approximately 49 acres and reconnaissance surveys were performed over approximately 20 acres.

Prior to acquisition of geophysical data, a grid was established over each site by surveyors. The grid spacing was either 200 feet, 100 feet, or 50 feet, depending on the area to be covered. Additional grid markers needed to perform the geophysical investigations were placed by the geophysical crew by measuring between the civil survey stakes.

The EM conductivity data were collected using a Geonics EM31 ground conductivity meter. Data were collected at 5-foot intervals along parallel grid lines. A line spacing of 25 feet was used over the solid waste sites. A 10-foot line spacing was used over the fire training areas and some of the petroleum spill sites, where the additional detail was desired.

Measurements consisted of conductivity and in-phase values and were stored along with the grid coordinates for each measurement location in a digital data logger. Conductivity values were also recorded periodically (approximately every 100 feet) in the field logbook along with observations of features that may have affected the measurements and observations of features, such as roads, cliffs, etc., which were useful in preparing base maps. Data were transferred daily from the data logger to a personal computer. The data manually recorded in the logbook were used to verify that the information transferred from the data logger was correct.

Data Processing

The raw EM data consist of computer files in X, Y, Z format. The X and Y refer to the grid location and the Z refers to the measurement. Several data processing steps were performed prior to interpretation of the data:

1. The data were edited to remove unwanted data, such as that from sections of lines that were duplicated.
2. Data were input into a spreadsheet program and rearranged into a grid format. Profile plots of the data were generated using the spreadsheet program.
3. Contour maps of the data were generated using the edited X, Y, Z files.

Interpretation

Both the profile plots of the data and the contour maps were used to interpret the data. The data profile plots were used to identify anomalies and determine their location along the survey liens. The anomaly positions were transferred to a base map. Contour maps of the conductivity and in-phase values were used to help establish spatial relationships between anomalies on adjacent lines. Anomalies, or groups of anomalies, on adjacent lines that appear to be related to the same source have been grouped together and shown as a single area on the interpretive maps.

Buried metal can often be mapped with EM conductivity methods because of the characteristic response produced by the metal. If enough metal is present and relatively shallow, both conductivity and in-phase values will be negative. With lesser amounts of metal or deeper burial, the EM data may exhibit values less than background without going negative. If the fill contains scattered metal rather than closely packed metal, the amplitude of the EM data may not be particularly anomalous. Instead, variations in values along the survey lines indicate the presence of metal. (In this case, the data look "noisy.") Fill material containing little or no metal can often be identified by elevated conductivity and/or in-phase measurements.

Several types of features have been interpreted from the data based on the appearance of the data as described above. Although the distinction between these interpreted features is somewhat subjective, there has been an attempt to maintain consistency in categorizing the features interpreted in the different areas.

Isolated Buried Metal. An anomaly was observed in the data that indicates buried metal. Corresponding anomalies on adjacent lines were not observed.

Possible Isolated Buried Metal. A relatively weak anomaly that does not exhibit all the characteristics of buried metal was observed in the data. Corresponding anomalies were not observed on adjacent lines.

Buried Utility. If narrow metal or possible metal responses were observed on several adjacent lines and if they lined up in a straight line, this was interpreted as a utility line and may be either a buried pipe or cable.

Area of Buried Metal

Areas of buried metal represent large areas of buried metal. Anomalies observed on a single traverse indicated multiple buried metal objects or wide continuous sections of buried metal. Similar anomalies were observed on adjacent lines, indicating extensive areas of metal.

Area of Scattered Buried Metal

This category is similar to Areas of Buried Metal, except that the buried metal is more scattered. Metal objects are close together enough to make delineation of individual sources unwarranted, however.

Area of Waste or Fill with Little or No Metal

These areas are defined by anomalously high conductivities but there are few in any of the characteristic metal-type responses. Similar anomalies are observed on adjacent lines, indicating some extent to these features.

The results of this investigation are presented in Chapter 3 of Volume 1 of this report along with the interpretive maps.

This Appendix presents the geophysical contour maps, both in-phase and conductivity, from each site where a complete geophysical investigation was performed on Shemya in 1992. Table B.1 lists the sites investigated and the order in which the maps are presented. The profile plots from each site follow the contour maps in the same order as given in Table B.1. Both the contour maps and profile plots were used to develop the interpretive maps presented in the main body of the report.

Table B.1.
Geophysical Investigations in 1992

FT-1	Lightning Strike Fire Training Site
FT-2	Aircraft Mock-up Fire Training Site
FT-3	Structural Fire Training Site
PS-1B	Old Cobra Dane
PS-6	Refueling Vehicle Maintenance Shop
PS-7	Vehicle Maintenance Shop
PS-8	Old White Alice
PS-9	Asphaltic Drum Storage Area
Water Gallery	
SW-4	Old Barrel Storage Area
SW-5	Hospital Lake
SW-10	Barrel Bay
SW-12	Scrap Metal Dump
SW-14	Scrap Metal Landfill

FIGURE B.1 FT-1 Geophy

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.2 FT-1 Geophy-

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.3 FT-2 Geophy

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.4 FT-2 Geophysical

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.5 FT-3 Geophy

- LEGEND
- GEOPHYSICAL SURVEY LINE
- Ⓜ METAL OBSERVED AT SURFACE

FIGURE B.6 FT-3 Geophysical

- LEGEND
- GEOPHYSICAL SURVEY LINE
- Ⓜ METAL OBSERVED AT SURFACE

FIGURE B.7 PS-1B Geophysical Contour Map - In Phase

18 January 1993

FIGURE B.8 PS-1B Geophysical Contour Map - Conductivity

18 January 1993

SURVEY MONUMENT
J-3 111V

FIGURE B.9 PS-6 Geoph

LEGEND
 — GEOPHYSICAL SURVEY
 LINE

FIGURE B.10 PS-6 Geophys

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.11 PS-7 Geophy

LEGEND

- ⊙ MANHOLE
- GEOPHYSICAL SURVEY LINE

FIGURE B.12 PS-7 Geophysi

LEGEND

⊙ MANHOLE

— GEOPHYSICAL SURVEY
LINE

FIGURE B.13 PS-8 Geophy

LEGEND

- ⊙ MANHOLE
- GEOPHYSICAL SURVEY LINE
- Ⓜ METAL OBSERVED AT SURFACE

FIGURE B.14 PS-8 Geophys

LEGEND

⊙ MANHOLE

— GEOPHYSICAL SURVEY LINE

Ⓜ METAL OBSERVED AT SURFACE

FIGURE B.15 PS-9 Geophy:

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.16 PS-9 Geophysical

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.17 WG Geophy

FIGURE B.18 WG Geophy

al Contour Map - Conductivity

18 January 1993

FIGURE B.19 SW-4 Geoph

sical Contour Map - In Phase

18 January 1993

FIGURE B.20 SW-4 Geophy:

LEGEND

— GEOPHYSICAL SURVEY
LINE

FIGURE B.21 SW-5 Geophysical Contour Map - In Phase

18 January 1993

LEGEND

— GEOPHYSICAL SURVEY LINE

FIGURE B.22 SW-5 Geophysical Contour Map - Conductivity

18 January 1993

11°59'35"

LEGEND

- GEOPHYSICAL SURVEY LINE
- Ⓜ METAL OBSERVED AT SURFACE
- ⌒ STEEP BANK

FIGURE B.23 SW-10 Geop

11°59'35"

LEGEND

—|— GEOPHYSICAL SURVEY LINE

(M) METAL OBSERVED AT SURFACE

⌋ STEEP BANK

OCEAN

FIGURE B.24 SW-10 Geophy:

FIGURE B.25 SW-12 Geop

LEGEND

- STEEP BANK
- UTILITY SERVICE BOX & POSTS
- BUILDING
- GEOPHYSICAL SURVEY LINE
- METAL OBSERVED AT SURFACE
- MARKED UNDERGROUND UTILITY LINE

FIGURE B.26 SW-12 Geoph

- LEGEND**
- STEEP BANK
 - UTILITY SERVICE BOX & POSTS
 - BUILDING
 - GEOPHYSICAL SURVEY LINE
 - METAL OBSERVED AT SURFACE
 - MARKED UNDERGROUND UTILITY LINE

FIGURE B.27 SW-14 Geophy

FIGURE B.28 SW-14 Geophys

LEGEND

— GEOPHYSICAL
SERVEY LINE

Ⓜ METAL OBSERVED
AT SURFACE

— DITCH

TTTT STEEP BANK

al Contour Map - Conductivity

18 January 1993

FT 1
LINE 390

FT 1
LINE 400

FT 1
LINE 410

FT 1
LINE 420

FT 1
LINE 430

FT 1
LINE 440

FT 1
LINE 450

— CONDUCTIVITY — IN-PHASE

FT 1
LINE 460

FT 1
LINE 470

— CONDUCTIVITY — IN-PHASE

FT 1
LINE 480

FT 1
LINE 490

FT 1
LINE 500

FT 2
LINE 100

FT 2
LINE 110

FT 2
LINE 120

FT 2
LINE 130

FT 2
LINE 140

— CONDUCTIVITY — IN-PHASE

FT 2
LINE 150

FT 2
LINE 160

FT 2
LINE 170

FT 2
LINE 180

FT 2
LINE 190

FT 2
LINE 200

FT 2
LINE 210

— CONDUCTIVITY — IN-PHASE

FT 2
LINE 220

FT 2
LINE 230

FT 2
LINE 240

FT 2
LINE 250

FT 2
LINE 260

FT 2
LINE 270

FT 2
LINE 280

— CONDUCTIVITY — IN-PHASE

FT 2
LINE 290

FT 2
LINE 300

FT 3
LINE 10 N

FT 3
LINE 20 N

FT 3
LINE 30 N

FT 3
LINE 40 N

FT 3
LINE 50 N

FT 3
LINE 60 N

FT 3
LINE 70 N

FT 3
LINE 80 N

FT 3
LINE 90 N

FT 3
LINE 100 N

OLD COBRA DANE (PS-1B)
LINE 10 N

OLD COBRA DANE (PS-1B)
LINE 20 N

OLD COBRA DANE (PS-1B)
LINE 30 N

OLD COBRA DANE (PS-1B)
LINE 40 N

OLD COBRA DANE (PS-1B)
LINE 50 N

OLD COBRA DANE (PS-1B)
LINE 60 N

OLD COBRA DANE (PS-1B)
LINE 70 N

OLD COBRA DANE (PS-1B)
LINE 80 N

OLD COBRA DANE (PS-1B)
LINE 90 N

OLD COBRA DANE (PS-1B)
LINE 100 N

OLD COBRA DANE (PS-1B)
LINE 110 N

OLD COBRA DANE (PS-1B)
LINE 120 N

OLD COBRA DANE (PS-1B)
LINE 130 N

PS 6
LINE 25 N

PS 6
LINE 0 N

— CONDUCTIVITY — IN-PHASE

PS 6
LINE 25 S

PS 6
LINE 50 S

PS 6
LINE 75 S

PS 6
LINE 100 S

PS 6
LINE 125 S

PS 6
LINE 150 S

PS 6
LINE 175 S

PS 6
LINE 200 S

PS 6
LINE 225 S

PS 7
LINE 100 W

PS 7
LINE 125 W

PS 7

LINE 150 W

PS 7
LINE 175 W

PS 7
LINE 200 W

PS 7
LINE 225 W

PS 7
LINE 250 W

PS 7
LINE 275 W

PS 8
LINE 25 N

PS 8
LINE 50 N

PS 8
LINE 75 N

PS 8
LINE 100 N

PS 8
LINE 125 N

PS 8
LINE 150 N

PS 8
LINE 175 N

PS 9
LINE 25 E

PS 9
LINE 50 E

PS 9
LINE 75 E

PS 9
LINE 100 E

PS 9
LINE 150 E

PS 9
LINE 175 E

PS 9
LINE 200 E

PS 9
LINE 125 E

PS 9
LINE 225 E

— CONDUCTIVITY — IN-PHASE

PS 9
LINE 250 E

PS 9
LINE 25 E

PS 9
LINE 50 E

PS 9
LINE 75 E

PS 9
LINE 100 E

— CONDUCTIVITY — IN-PHASE

PS 9
LINE 125 E

PS 9
LINE 150 E

— CONDUCTIVITY — IN-PHASE

PS 9
LINE 175 E

PS 9
LINE 200 E *metal*

PS 9
LINE 225 E

PS 9
LINE 250 E

— CONDUCTIVITY — IN-PHASE

WATER GALLERY LINE 25

WATER GALLERY LINE 50

— CONDUCTIVITY — IN-PHASE

WATER GALLERY LINE 75

WATER GALLERY LINE 100

— CONDUCTIVITY — IN-PHASE

WATER GALLERY
LINE 125

WATER GALLERY LINE 150

— CONDUCTIVITY — IN-PHASE

WATER GALLERY LINE 175

— CONDUCTIVITY — IN-PHASE

WATER GALLERY LINE 200

WATER GALLERY LINE 225

WATER GALLERY LINE 250

— CONDUCTIVITY — IN-PHASE

WATER GALLERY LINE 275

**WATER GALLERY
LINE 300**

WATER GALLERY LINE 325

WATER GALLERY LINE 350

— CONDUCTIVITY — IN-PHASE

**WATER GALLERY
LINE 375**

WATER GALLERY LINE 400

— CONDUCTIVITY — IN-PHASE

WATER GALLERY LINE 425

WATER GALLERY LINE 450

SW 4
LINE 300

SW 4
LINE 325

SW 4
LINE 350

SW 4
LINE 375

SW 4
LINE 400

SW 4
LINE 425

SW 4
LINE 450

SW 4
LINE 475

SW 4
LINE 500

SW 4
LINE 525

SW 4
LINE 550

SW 4
LINE 575

SW 4
LINE 600

SW 4
LINE 625

SW 4
LINE 650

SW 4

LINE 675

SW 4
LINE 700

SW 4
LINE 725

SW 4
LINE 750

SW 4
LINE 775

SW 4
LINE 800

SW 4
LINE 825

SW 4
LINE 850

SW 4
LINE 875

SW 4
LINE 900

SW 4
LINE 925

SW 4
LINE 950

SW 4
LINE 975

SW 4
LINE 1000

SW 4
LINE 1025

SW 4
LINE 1050

SW 4
LINE 1075

SW 4
LINE 1100

SW 4
LINE 1125

SW 4
LINE 1150

SW 4
LINE 1175

SW 4
LINE 1200

SW 4
LINE 1225

SW 4
LINE 1250

SW 4
LINE 1275

SW 4
LINE1300

SW 4
LINE 1325

SW 4

LINE 1350

— CONDUCTIVITY — IN-PHASE

SW 4
LINE 1375

SW 4
LINE 1400

SW 4
LINE 1425

SW 4
LINE 1450

SW 4
LINE 1475

SW 4
LINE 1500

SW 4
LINE 1525

SW 4
LINE 1550

HOSPITAL LAKE

LINE 500 N

HOSPITAL LAKE
LINE 525 N

— CONDUCTIVITY — IN-PHASE

HOSPITAL LAKE

LINE 550 N

HOSPITAL LAKE

LINE 575 N

HOSPITAL LAKE
LINE 600 N

— CONDUCTIVITY — IN-PHASE

HOSPITAL LAKE

LINE 625 N

HOSPITAL LAKE

LINE 650 N

HOSPITAL LAKE

LINE 675 N

HOSPITAL LAKE

LINE 700 N

HOSPITAL LAKE LINE 725 N

HOSPITAL LAKE

LINE 750 N

HOSPITAL LAKE LINE 775 N

HOSPITAL LAKE

LINE 800 N

— CONDUCTIVITY — IN-PHASE

HOSPITAL LAKE

LINE 825 N

HOSPITAL LAKE LINE 850 N

HOSPITAL LAKE

LINE 875 N

HOSPITAL LAKE

LINE 900 N

HOSPITAL LAKE
LINE 925 N

HOSPITAL LAKE LINE 950 N

— CONDUCTIVITY — IN-PHASE

HOSPITAL LAKE LINE 975 N

HOSPITAL LAKE LINE 1000 N

— CONDUCTIVITY — IN-PHASE

SW 10
LINE 4025

SW 10
LINE 4050

SW 10
LINE 4075

SW 10
LINE 4100

SW 10
LINE 4125

SW 10
LINE 4150

SW 10

LINE 4175

SW 10
LINE 4200

SW 10
LINE 4225

SW 10
LINE 4250

SW 10
LINE 4275

SW 10
LINE 4300

SW 10

LINE 4325

— CONDUCTIVITY — IN-PHASE

SW 10
LINE 4350

SW 10

LINE 4375

SW 10
LINE 4400

SW 10
LINE 4425

SW 10
LINE 4450

SW 10
LINE 4475

SW 10
LINE 4500

SW 10
LINE 4525

SW 10
LINE 4550

SW 10

LINE 4575

SW 10
LINE 4600

SW 10
LINE 4625

SW 10

LINE 4650

SW 10
LINE 4675

SW 10
LINE 4700

— CONDUCTIVITY — IN-PHASE

SW 10
LINE 4725

SW 10
LINE 4750

SW 10
LINE 4775

SW 10
LINE 4800

SW 10
LINE 4825

SW 10
LINE 4850

SW 10
LINE 4875

SW 10
LINE 4900

SW 10
LINE 4925

SW 10
LINE 4950

— CONDUCTIVITY — IN-PHASE

SW 10
LINE 4975

SW 10
LINE 5000

SW 10
LINE 5025

SW 10
LINE 5050

SW 10
LINE 5075

SW 10
LINE 5100

SW 12
LINE 4350 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4375 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4400 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4425 N

SW 12
LINE 4450 N

SW 12
LINE 4475 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4500 N

SW 12
LINE 4525 N

SW 12
LINE 4550 N

SW 12
LINE 4575 N

SW 12
LINE 4600 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4625 N

SW 12
LINE 4650 N

SW 12
LINE 4675 N

SW 12
LINE 4700 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4725 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4750 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4775 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4800 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4825 N

— CONDUCTIVITY — IN-PHASE

SW 12
LINE 4850 N

SW 12
LINE 4875 N

SW 12
LINE 4900 N

SW 14
LINE 100 N

SW 14
LINE 125 N

SW 14
LINE 150 N

SW 14
LINE 175 N

SW 14
LINE 200 N

SW 14
LINE 225 N

SW 14
LINE 250 N

SW 14
LINE 275 N

SW 14
LINE 300 N

SW 14
LINE 325 N

SW 14
LINE 350 N

SW 14
LINE 375 N

SW 14
LINE 400 N

SW 14
LINE 425 N

SW 14
LINE 450 N

SW 14
LINE 475 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 500 N

SW 14
LINE 525 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 550 N

SW 14
LINE 575 N

SW 14
LINE 600 N

SW 14
LINE 625 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 650 N

SW 14
LINE 675 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 700 N

SW 14
LINE 725 N

SW 14
LINE 750 N

SW 14
LINE 775 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 800 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 825 N

SW 14
LINE 850 N

SW 14
LINE 875 N

SW 14
LINE 900 N

— CONDUCTIVITY — IN-PHASE

SW 14
LINE 925 N

SW 14
LINE 950 N

SW 14
LINE 975 N

SW 14
LINE 1000 N

SW 14
LINE 1025 N

SW 14
LINE 1050 N

SW 14
LINE 1075 N

SW 14
LINE 1100 N

SW 14
LINE 1125 N

SW 14
LINE 1150 N

SW 14
LINE 1175 N

SW 14
LINE 1200 N

Appendix C
Human Health Risk Assessment
Preliminary Risk Levels

CONTENTS

APPENDIX C - HUMAN HEALTH RISK ASSESSMENT-PRELIMINARY	
RISK LEVELS	C.1
1.0 INTRODUCTION	C.1.1
1.1 PURPOSE	C.1.1
1.2 SITE DESCRIPTION	C.1.1
1.3 ORGANIZATION OF RISK ASSESSMENT PROTOCOLS	C.1.2
2.0 SELECTION OF CHEMICALS OF POTENTIAL CONCERN	C.2.1
3.0 EXPOSURE ASSESSMENT	C.3.1
3.1 IDENTIFICATION OF POTENTIAL RECEPTORS	C.3.1
3.2 POTENTIAL EXPOSURE PATHWAYS	C.3.1
3.3 CONTAMINANT SOURCES	C.3.2
3.4 CONTAMINANT FATE AND TRANSPORT	C.3.2
3.5 EXPOSURE POINT	C.3.4
3.6 EXPOSURE ROUTES	C.3.6
3.7 EXPOSURE PARAMETERS	C.3.6
4.0 DEVELOPMENT OF PRELIMINARY RISK LEVELS	C.4.1
4.1 FEDERAL OR STATE ACTION LEVELS TO BE CONSIDERED FOR HUMAN HEALTH PRLS	C.4.1
4.2 RISK-BASED CALCULATIONS FOR HUMAN HEALTH PRLS	C.4.2
5.0 RISK CHARACTERIZATION	C.5.1
6.0 REFERENCES	C.6.1

TABLES

	After Page
C.2.1 Chemicals of Potential Concern	C.2.2
C.3.1 IRP Sites	C.3.2
C.3.2 Exposure Parameters for the Construction Worker Exposure Scenario	C.3.6
C.4.1 Shemya Groundwater ARARs	C.4.3
C.4.2 Risk-based Concentrations for Carcinogenic Effects Based on Ingestion of Soil and Sediment by Construction Workers	C.4.7
C.4.3 Risk-based Concentrations for Noncarcinogenic Effects Based on Ingestion of Soil and Sediment by Construction Workers	C.4.10
C.4.4 Risk-based Concentrations for Carcinogenic Effects Based on Ingestion/Inhalation of Groundwater by AFB Personnel	C.4.13
C.4.5 Risk-based Concentrations for Noncarcinogenic Effects Based on Ingestion of Groundwater by AFB Personnel	C.4.16
C.5.1 Selected Soil and Sediment Preliminary Risk Levels	C.5.2
C.5.2 Selected Groundwater Preliminary Risk Levels	C.5.5
C.5.3 COPCs that Exceed Preliminary Risk Levels	C.5.7

Appendix C

Human Health Risk Assessment— Preliminary Risk Levels

1.0 Introduction

The U.S. Air Force is currently conducting a site investigation and closure activities under the Installation Restoration Program (IRP) at the Shemya Air Force Base (AFB), Shemya Island, Alaska. This Appendix presents the Preliminary Risk Levels (PRLs) for the Shemya AFB based on human health risk assessment methods.

1.1 Purpose

The objective of the risk assessment for the Shemya AFB is to compare risk-based action levels that are adequately protective of human health to available analytical results. Risk assessment methods developed by U.S. EPA (EPA, 1989 and EPA, 1991) will be used to calculate PRLs for soil, surface water, sediment, and groundwater. PRLs could be used as cleanup goals and: 1) are protective of human health and the environment; and 2) comply with regulatory standards, criteria, or action levels (e.g., MCLs). Chemical concentrations greater than the PRL concentration may present potential adverse health effects to human receptors at the site. These criteria are preliminary as they may be changed through negotiation between the Air Force and regulatory agencies. The concentrations of positively detected chemicals at 11 sites across the island will be compared to the most stringent PRLs. Sites with chemicals exceeding a PRL may be considered for remedial action.

1.2 Site Description

The Shemya AFB occupies the entire 3.5-mile-long by 1.5-mile-wide Shemya Island. The island is located at the western tip of the Aleutian Archipelago, 1,500 miles southwest of Anchorage, Alaska. The primary mission of the 5073rd Air Base Group based at Shemya AFB is as an early warning radar installation. The island was first developed as an airfield in 1943 and has served in that capacity, in one form or another, for the past 50 years.

A records search conducted in September 1984 identified the following classifications for areas on the island as potentially containing hazardous materials from past activities:

- petroleum, oil, and lubricants spill sites (PS)

- solid waste disposal sites (SW)
- fire training sites (FT)

Site data were collected from 10 IRP sites during the 1992 field season. PRLs for these 10 areas or sites are developed: FT-1, FT-2, SW-4, SW-5, SW-10, SW-12, PS-1A, PS-1B, PS-8, and PS-10. Contamination has also been detected at the water gallery. The water gallery is a near-surface water collection system that collects potable water for the base from a shallow aquifer in the eastern watershed.

1.3 Organization of Risk Assessment Protocols

The following risk assessment tasks are addressed in this Appendix:

- **Selection of Chemicals of Potential Concern** identifies the chemicals evaluated in the risk assessment.
- **Exposure Assessment** describes how receptors could come into contact with contaminants from the site.
- **Development of PRLs** describes how the PRLs are developed from the site.
- **Risk Characterization** involves comparison of site concentrations to PRLs.

2.0 Selection of Chemicals of Potential Concern

Chemicals of Potential Concern (COPC) were identified on the basis of analytical results for soil, sediment, surface water, and groundwater samples collected in 1992. Samples were analyzed using onsite Close Support Laboratory (CSL) facilities and offsite laboratories using standard EPA methodology. PRLs are developed for each COPC. If a chemical was positively detected in a site-related soil, groundwater, surface water, or sediment sample, that chemical is considered a COPC.

Background samples were collected as part of the 1992 field investigation: two sediment samples (BSSD-126 and BSSD-127); two soil samples (BS-124 and BS-125); and one groundwater sample (BG-BW01). Concentrations from each of these background samples are compared to the PRL. If a background concentration and the associated site-related concentration exceeds its PRL, the site may not require remediation as the chemical also exists in nature at a high enough concentration to exceed a PRL. This method was used because of the lack of background samples to perform a more rigorous statistical analysis.

Table C.2.1 contains the soil, groundwater, surface water, and sediment COPCs. The chemicals listed in Table C.2.1 were detected at least once in the respective media at the IRP sites included in the 1992 field investigation. PRLs have been developed for these chemicals using the methods described below. However, it was not possible to develop PRLs for all COPCs based on the available methods and information. Additional effort not covered by the scope of this assessment would be necessary to complete PRLs for all COPCs.

Table C.2.1
Chemicals of Potential Concern

Organics	Inorganics
Soil	
Volatiles:	
1,1,1-Trichloroethane 4-Methyl-2-pentanone Benzene Carbon disulfide Chlorobenzene Chloroform Ethylbenzene Methylene chloride Tetrachloroethylene Toluene Trichloroethylene Xylene	Arsenic Beryllium Cadmium Chromium Copper Cyanide Lead Mercury Zinc
Semivolatiles:	
Acenaphthene Anthracene Benzo(a)anthracene Benzo(a)pyrene Benzo(b)fluoranthene Benzo(g,h,i)perylene Benzo(k)fluoranthene bis(2-ethylhexyl)phthalate Butylbenzylphthalate Carbazole di-n-Butylphthalate di-n-Octylphthalate Dibenz(a,h)anthracene Dibenzofuran Diethylphthalate Fluoranthene	

Table C.2.1
Chemicals of Potential Concern

		(cont.)
Organics	Inorganics	

Soil (Continued)

Semivolatiles (Continued):

2-Methylnaphthalene
4-Methylphenol
Chrysene
Dioxins/Furans (Total)
Fluorene
Indeno(1,2,3-cd)pyrene
Naphthalene
Phenanthrene
Phenol
Polychlorinated dibenzo
Pyrene

Pesticides/PCBs:

4,4'-DDE, DDT, DDD
Endrin ketone

TPH:

Total Petroleum Hydrocarbons

Table C.2.1
Chemicals of Potential Concern

(cont.)

Organics	Inorganics
Groundwater	
Volatiles:	
1,1,1-Trichloroethane	Aluminum
Acetone	Arsenic
Benzene	Barium
Carbon disulfide	Beryllium
Chloroform	Cadmium
Ethylbenzene	Chromium
Methylene chloride	Cobalt
Toluene	Copper
Trichloroethylene	Manganese
Xylene	Molybdenum
Semivolatiles:	Nickel
2-4-Dimethylphenol	Silver
2-Methylnapthalene	Thallium
2-Methylphenol	Vanadium
4-Methylphenol	Zinc
bis(2-ethylhexyl)phthalate	
Dibenzofuran	
Diethylphthalate	
Fluorene	
Naphthalene	
Phenol	
TPH:	
Total Petroleum Hydrocarbons	

Table C.2.1
Chemicals of Potential Concern

(cont.)

Organics	Inorganics
Sediment	
Volatiles:	
2-Butanone	Antimony
Benzene	Arsenic
Ethylbenzene	Cadmium
Toluene	Chromium
Xylene	Copper
	Lead
	Zinc
Semivolatiles:	
bis(2-ethylhexyl)phthalate	
Fluoranthene	
Pyrene	
Pesticides/PCBs:	
4,4'-DDD, DDE, DDT	
Aroclor 1254	
Endrin ketone	
TPH:	
Total Petroleum Hydrocarbons	

Table C.2.1
Chemicals of Potential Concern

(cont.)

Organics	Inorganics
Surface Water	
Volatiles:	Copper Zinc
Benzene Ethylbenzene Toluene Xylene	
Semivolatiles	
bis(2-ethylhexyl)phthalate	
General Chemistry:	
Nitrate/Nitrite	

3.0 Exposure Assessment

3.1 Identification of Potential Receptors

The potential receptors at Shemya Island are the military and civilian service personnel stationed at Shemya AFB and the DOJ and civilian contractors who are involved in construction work on the island. The AFB personnel normally spend a year on the island, and the contractors return every year for many years. All receptors potentially may be exposed to contaminated soil (surface and subsurface), surface water and sediment, and drinking water from the water gallery and inland sites (e.g., FT-2).

The Air Force is allowed to use the island as long as they need it. For this reason, and because of the island's remote location and harsh climate, the presence of human receptors, besides those mentioned above (i.e., trespassers) is highly unlikely.

The exposure scenario discussed in this assessment involves the potential exposure of construction workers to contaminants found on the island. The assessment of this scenario follows the standard EPA guidance for a commercial/industrial worker scenario (EPA, 1989). This scenario was chosen because it is more conservative than a scenario involving base personnel or a trespasser to the island. The base personnel are on the island only one year compared to a possible 15 to 20 years for a construction worker.

3.2 Potential Exposure Pathways

An exposure pathway is the means by which a person (receptor) may come into contact with (be exposed to) chemical contaminants in environmental media. A complete pathway has five elements:

- source of contamination
- mechanisms for contaminant release
- environmental transport medium
- exposure point
- feasible route of exposure

If one of these five elements does not exist, the exposure pathway is incomplete and exposure will not occur and there is no risk of adverse health effects.

3.3 Contaminant Sources

This assessment will address 11 sites on Shemya Island (see Table C.3.1). These sites are included in the installation restoration program based on historical land use and disposal activities. Constituents found at each historical site on the island and a more detailed description of each site can be found in the Field Investigation Plan (CH2M HILL, 1991) and the body of this report.

Table C.3.1
IRP Sites

Site	Site Name
FT-1	Lightning Strike
FT-2	Aircraft Mockup
SW-4	Barrel Dump Site
SW-5	Hospital Lake
SW-10	Barrel Bay
SW-12	Scrap Metal Disposal Site
WG	Water Gallery
PS-1A	Transformer Oil Spills at Cobra Dane (New Cobra Dane)
PS-1B	Old Cobra Dane
PS-8	Old White Alice
PS-10	JP-4 Spill at Base Operations Terminal

3.4 Contaminant Fate and Transport

Three important factors affecting the fate of chemicals in the environment are environmental partitioning, persistence, and bioaccumulation. Several chemical and physical properties can influence the rates and extent of chemical partitioning between soil, water, and air. These include water solubility, vapor pressure, Henry's Law Constant, and soil-water partition coefficients. Bioaccumulation is the partitioning between environmental media and living organisms and is measured by the property of bioconcentration. Persistence is affected by susceptibility to degradation reactions and the properties previously mentioned. The following paragraphs discuss the partitioning, persistence, and bioaccumulation characteristics of the classes of chemicals found at Shemya.

Petroleum Hydrocarbons

Petroleum hydrocarbons (reported as TPH) are composed of a mixture of several chemical constituents: normal and branched (aliphatic hydrocarbons) alkanes, cyclic (aromatic hydrocarbons) alkanes, alkenes, and olefins. Transport and transformation of a TPH spill or plume will vary depending on the physical, chemical, and biological properties of the constituents. Some constituents will dissolve more quickly, be sorbed less strongly on soils and other solids, and may be more or less susceptible to degradation. Hydrocarbons found in JP-4 jet fuel mixture are expected to be relatively mobile and nonpersistent in most soil and water systems (AFSC, 1987). Persistence in deep soils and groundwater may be higher. Volatilization, photooxidation, and biodegradation are important fate processes. Surface spills are expected to be weathered by evaporation and photooxidation. Downward migration of weathered JP-4 fuel spills represents a potential threat to the groundwater system. Biodegradation of JP-4 fuel is expected to be significant under environmental conditions favorable to microbial oxidation (AFSC, 1987).

Straight chain aliphatic hydrocarbons are usually sparingly soluble in water. The solubility of hexane is approximately 10 mg/L; this solubility decreases by a factor of 3 or 4 for every carbon added. The branched aliphatic hydrocarbons and aromatic hydrocarbons tend to be more soluble in water. Overall, JP-4 has a solubility in water of 300 mg/L, a log octanol-water partition coefficient of 3 to 7, and Henry's Law Constant of 1×10 to 1×10^6 Pa m³/mol; and a bioconcentration factor of 50 to 50,000 based on a range for typical components (AFSC, 1987).

PAHs

Polynuclear aromatic hydrocarbons (PAHs) (e.g., benzo(a)anthracene, benzo(a)pyrene, anthracene) are formed under conditions when a fuel is burned without sufficient oxygen and are present in exhausts from engines and power plants. PAHs typically are slightly soluble in water and tend to partition to organic matter in soil and groundwater systems; PAHs tend to be rated as having low mobility to being immobile. PAHs have a typical solubility in water of 0.01 to 30 mg/L, an organic carbon-water partition coefficient of 1,300 to 2,000,000, and Henry's Law Constant of 9×10^{-2} to 3.5×10^1 Pa m³/mol; and a bioconcentration factor of 50 to 50,000 based on a range for typical components.

Chlorinated Hydrocarbons

If chlorine is substituted for a hydrogen in a hydrocarbon (e.g., methylene chlorine or 1,1,1-trichloroethane) the resulting compound tends to be less soluble and more persistent in the environment. Chlorinated hydrocarbons tend to penetrate the tissues of organisms more easily. Chlorinated hydrocarbons have a typical solubility in water of

200 to 13,000 $\mu\text{g/L}$, a log octanol-water partition coefficient of 1 to 2.5, and Henry's Law Constant of 0.09 to 0.8 $\text{Pa m}^3/\text{mol}$; and a bioconcentration factor of 4 to 4.5 based on a range for typical components (Mackay, 1992).

PCBs, Chlorobenzenes, Dioxins, and Pesticides

PCBs and chlorobenzenes are very persistent due to desirable properties of stability and have a tendency to partition out of air and water into receptor organisms. Chlorobenzenes have low to moderate mobility in a soil-water system and PCBs are immobile (Roy and Griffin 1985). Dioxins and furans are similar to PCBs in their relative environmental persistent and partitioning tendencies. Aroclor 1260 has a solubility in water of 3 to 14 $\mu\text{g/L}$, a log octanol-water partition coefficient of 4 to 7, and Henry's Law Constant of 10 to 1000 $\text{Pa m}^3/\text{mol}$; and a bioconcentration factor of 1,000 to 1,000,000 based on a range for typical components (Mackay, 1992). Chlorobenzenes have a typical solubility in water of 25 to 150 mg/L , a log octanol-water partition coefficient of 3 to 4, and Henry's Law Constant of 120 to 470 $\text{Pa m}^3/\text{mol}$; and a bioconcentration factor of 500 to 1000 based on a range for typical components (Mackay, 1992). Dioxins have a typical solubility in water of 0.2 to 0.00008 $\mu\text{g/L}$, a log octanol-water partition coefficient of 6 to 8, and Henry's Law Constant of 0.002 to 2 $\text{Pa m}^3/\text{mol}$; and a bioconcentration factor of 100 to 10,000 based on a range for typical components (Mackay, 1992).

Pesticides such as DDT, DDD, and DDE are known for their persistence in the environment. These pesticides are not soluble in water and are immobile in a soil-water system. DDT and its degradation products, DDD and DDE, have a typical solubility in water of 0.01 to 0.002 mg/L and a organic carbon-water partition coefficient of 155,000 to 238,000 (Roy and Griffin, 1985).

Metals

Metals have a complex environmental chemistry that varies considerably between compounds depending on their ionic state. The chemical properties discussed above are not as readily applied to inorganic compounds because of this complex chemistry. This assessment assumes the detected metals are available to the receptor in the form in which toxicity is measured.

3.5 Exposure Point

Construction workers may be exposed to contaminated soil or sediment during excavation and construction activities via incidental ingestion of soil and/or dermal absorption of contaminated soil. In this preliminary assessment of risk it is assumed the receptor may be exposed to soil or sediment at any depth due to excavation activities. Construction workers may also ingest contaminated groundwater at a site that would

support a producing well. However, there is no indication that any sites, other than the water gallery, would produce enough potable water for a complete exposure pathway. Receptors may also be exposed to chemicals volatilizing from groundwater into residences or via dermal absorption of chemicals in water (i.e., showering, bathing, preparing food, washing dishes). These receptors may also be exposed to contaminated surface water by incidental ingestion and dermal absorption while working in a lake or stream. Workers are not expected to be exposed via the incidental ingestion of surface water (EPA, 1989). This point of exposure would not occur at Shemya Island at a sufficient frequency to significantly impact a receptor's total exposure to contaminants. (For example, using EPA standard parameters, if a worker swam in Hospital Lake for 8 hours per day for 180 days in a year, the total amount of water intake through incidental ingestion would be 10 times less than similar exposure to groundwater.) Given this information and that high concentrations of contaminants have not been found in surface water and the climate on the island, this exposure point will not be quantified in this assessment. Dermal absorption is a complex activity controlled by many factors; it is difficult to predict exposures from this route. Therefore, dermal absorption is not considered in this assessment.

It is not anticipated that there will be any dust emissions. This is based on the following considerations:

- many sites have been covered by soil or crushed rock
- dust emissions are not a problem because of the high number of days of precipitation (approximately 330 per year), the low evaporation due to low temperature and high humidity, and the cover material
- volatilization of the organic COCs from contaminated soil is not considered a complete pathway for purposes of development of the PRLs. The low temperatures at Shemya Island, the tendency for organic chemicals to sorb to the organic rich soil and peat, the depth of contamination, and the turbulent wind conditions all combine to decrease the importance of this exposure pathway. The low temperature at Shemya Island may not eliminate volatilization entirely, since diffusion may continue even if the ground is frozen (Lyman, 1982). Further, the temperature at Shemya does fluctuate from 28.5°F to 48.5°F (the mean annual temperature = 38.5°F). However, low temperatures do greatly reduce diffusion and volatilization of organic compounds from soil (Dragun 1988). Further, the wind would carry any volatilized organic compound away from the island due to a high wind velocity (the average year-round velocity is 20 mph) and the tendency for high wind gusts (up to 63 mph) to occur throughout the year. Snow cover also reduces volatilization of organic compounds (Dragun, 1988). Neglecting this pathway in this assessment may introduce uncertainty (i.e., underestimate risk). The potential completeness of

volatilization of organics from soil and subsequent transport to a receptor may warrant evaluation in the future to address this uncertainty.

Because of these factors, exposure to air emissions is not considered to be a complete exposure pathway.

3.6 Exposure Routes

Based on the discussion in the previous section, the following exposure routes will be assessed:

- incidental ingestion of surface soil
- incidental ingestion of subsurface soil
- ingestion of groundwater
- inhalation of chemicals volatilizing from contaminated water

3.7 Exposure Parameters

Table C.3.2 lists the exposure parameters for calculation of the human health PRLs based on a construction worker scenario.

Table C.3.2
Exposure Parameters for the Construction Worker Exposure Scenario

Exposure Parameter	Value	Reference
Soil/Sediment Ingestion Rate (IR_s)	50 mg/day	EPA 1991
Groundwater Ingestion Rate (IR_w)	2 L/day	EPA 1991
Groundwater Inhalation Rate (IR_g)	15 m ³ /day	EPA 1991
Volatilization Factor (K)	0.5 L/m ³	EPA 1991
Body Weight (BW)	70 kg	EPA 1991
Exposure Frequency (EF)	180 days/year	Based on available days per year to perform construction activities on the island
Exposure Duration (ED)	20 years	Based on a reasonable maximum exposure duration for AFB personnel or contractors
Averaging Time (AT)	70 years for carcinogens; equal to ED for noncarcinogens	EPA 1989

4.0 Development of Preliminary Risk Levels

The Preliminary Risk Levels (PRLs) consist of medium-specific goals (i.e., groundwater, surface water, sediment, and soil) for protecting human health. The PRLs discussed here are chemical-specific and are established using readily available information such as reference doses, slope factors (used in the development of risk-based PRLs) and federal or state action levels (e.g., MCLs or water quality criteria).

4.1 Federal or State Action Levels to be Considered for Human Health PRLs

The following chemical-specific federal or state standards, criteria, or action levels were considered as PRLs for protection of human-health:

- Safe Drinking Water Act and the National Primary Drinking Water Regulations (40 CFR 141)
- RCRA Groundwater Protection Standards (40 CFR 264.94)
- Clean Water Act Water Quality Criteria
- Alaska Drinking Water Regulations (18 AAC 80)
- Alaska Water Quality Standards (18 AAC 70)

Safe Drinking Water Act and the National Primary Drinking Water Regulations (40 CFR 141)

These regulations establish enforceable maximum contaminant levels (MCLs) for inorganic and organic compounds in public drinking water systems. Because groundwater is the primary drinking water source at Shemya AFB, MCLs are potentially applicable standards.

Clean Water Act Water Quality Criteria

The water quality criteria for human health protection may be relevant and appropriate since surface waters at the island are highly accessible.

Alaska Drinking Water Regulations (18 AAC 80)

These regulations establish maximum contaminant concentrations (MCCs) for organic and inorganic contaminants in public water systems. These concentrations are equivalent to the SDWA MCLs with the exception of fluoride which has been set at a lower limit of 2.4 mg/L.

Alaska Water Quality Standards (18 AAC 70)

These standards apply to ground and surface waters and establish criteria for protected water use classes. In addition, Alaska has adopted a nondegradation policy for waters of higher quality than the criteria. Chemical-specific water quality criteria are listed in Table C.4.1. General water quality criteria are not addressed in this assessment but are found in the Field Investigation Plan (CH2M HILL, 1991).

Table C.4.1 lists the ARARs used in this assessment.

4.2 Risk-Based Calculations for Human Health PRLs

Risk-based human health PRLs were derived for this assessment using the exposure assumptions discussed in Section 3 of this Appendix and equations presented in Part B of Risk Assessment Guidance for Superfund (EPA, 1991). There will be up to two risk-based PRLs for each exposure scenario due to the type of toxicological effect the chemical may exhibit: noncarcinogenic or carcinogenic. Risk-based PRLs for carcinogenic chemicals are calculated by setting the total risk for carcinogenic effects at a target risk level and then solving for the concentration variable. Target risk levels used in this assessment will include excess lifetime risk estimates of 10^{-4} , 10^{-5} , and 10^{-6} for carcinogenic chemicals. A cumulative lifetime risk of 10^{-6} is the National Oil and Hazardous Substances Pollution Contingency Plan (NCP) point of departure for the analysis for remedial alternatives and according to EPA (EPA, 1991), a cumulative lifetime risk of 10^{-4} is the point at which action is generally warranted at a site. The risk-based PRLs for chemicals that caused noncarcinogenic effects are set at a target hazard index of one for each COPC with a reference dose (RfD).

The detection limit used by the laboratory may be higher than the PRL. This introduces the uncertainty of not detecting contamination between zero and the detection limit that exceeds a PRL. This is a source of uncertainty not addressed in this assessment and may tend to underestimate risk.

PRLs for chemicals with carcinogenic or noncarcinogenic effects in soil and groundwater (based on soil ingestion and groundwater ingestion exposures) can be estimated by the following procedures.

Table C.4.1
Shemya Groundwater ARARs

Chemical	CWA Water Quality Criteria for Protection of Human Health Water & Fish Ingestion (mg/L)	SDWA MCLs (mg/L)
<u>Organics</u>		
Acetone	-	-
Benzene	0.00066	0.005
bis(2-ethylhexyl)phthalate	-	0.004-p
Carbon disulfide	-	-
Chloroform	0.00019	-
Dibenzofuran	-	-
1,2-Dichloroethene (total)	-	0.1
Diethylphthalate	350	-
Ethylbenzene	1.4	0.7
Fluorene	0.0000031 (Total PAHs)	-
Methylene chloride	0.00019(total halomethanes)	0.005-p
2-Methylnaphthalene	-	-
2-Methylphenol	-	-
4-Methylphenol	-	-
2,4-Dimethylphenol	-	-
Naphthalene	-	-
Nitrate (As NO3)	-	10
Phenol	3.5	-
Toluene	15	1
1,1,1-Trichloroethane	19	0.2
Trichloroethylene	0.0028	0.005
Total Xylenes	-	10
<u>Inorganics</u>		
Aluminum	-	-
Antimony	0.146	0.01
Arsenic	0.000025	0.05
Barium	-	1
Beryllium	0.0000039	0.001
Cadmium	0.005	0.01
Chromium (Total)	0.05 (hexavalent)	0.1
Cobalt	-	-
Copper	-	1.3-p
Cyanide	0.2	0.2-p
Iron	-	-
Lead	-	-
Magnesium	-	-
Manganese	-	-
Mercury	0.01	-
Molybdenum	-	-
Nickel	0.0154	.1-p
Selenium	0.01	0.01
Silver	0.05	0.05
Thallium	0.0178	.002-p
Vanadium	-	-
Zinc	-	-

Notes:

p = proposed MCL

Carcinogenic Effects

Soil Ingestion. Target concentrations (TC) for individual carcinogens are developed by estimating the exposure required to equal a designated risk level. This requires estimating the concentration needed in the exposure medium under the assumed exposure condition to achieve that exposure level.

The TC for carcinogens from ingested soil can be calculated (using Equation C-1) as follows:

$$TC_s = (TRL \times AT \times 365 \text{ days/yr}) / (SF \times 10^{-6} \text{ mg/kg} \times EF \times IF) \quad (C-1)$$

where:

TC_s = Target chemical concentration for soil ($\mu\text{g/kg}$)
 TRL = Target risk level
 BW = Body weight (kg)
 AT = Averaging time (years)
 SF = Slope factor (mg/kg-day^{-1})
 IR = Ingestion rate (mg/day)
 EF = Exposure frequency (days/year)
 ED = Exposure duration (years)
 IF = Age-adjusted soil ingestion factor (mg-yr/kg-day)
 (using Equation C-2)

$$IF_{adult} = (IR_{adult} \times ED_{adult}) / BW_{adult} \quad (C-2)$$

Groundwater Exposure. The TC (potential PRLs) for carcinogens from groundwater use can be calculated (using Equation C-3) as follows:

$$TC_{gw} = \frac{(TRL \times BW \times AT \times 365 \text{ days/yr})}{EF \times ED \times [(SF_i \times K \times IR_3) + (SF_o \times IR_w)]} \quad (C-3)$$

where:

TC_{gw} = Target chemical concentration in groundwater (mg/l)
 TRL = Target risk level
 BW = Body weight (kg)
 AT = Averaging time (years)
 EF = Exposure frequency (days/year)
 ED = Exposure duration (years)
 SF_i = Inhalation slope factor ($\text{mg/kg} \cdot \text{day}^{-1}$)

- SF_o = Oral slope factor (mg/kg - day)⁻¹
 IR_3 = Daily inhalation while showering intake rate (L/day)
 IR_w = Daily water ingestion rate (L/day)
 K = Volatilization Factor (L/m³)

Noncarcinogenic Effects

Target concentrations for chemicals with noncarcinogenic effects (such as having reference dose concentrations) are developed by estimating the exposure required to equal the reference dose for the particular chemical. This requires estimating the concentration needed in the exposure medium under the assumed exposure condition to achieve that exposure level. The following assumes exposure to a single chemical. Exposure to multiple chemicals may require weighing factors to obtain resulting risk levels associated with an RfD and are not considered in this assessment.

Soil Ingestion. The TC (potential PRLs) for chemicals with noncarcinogenic effects for soil ingestion can be estimated (using Equation C-4) as follows:

$$TC_s = (THI \times AT \times 365 \text{ days/yr}) / (1/RfD \times 1 \times 10^6 \text{ kg/mg} \times EF \times IF) \quad (C-4)$$

where:

- TC_s = Target concentration for soil (μg/kg)
 THI = Target hazard index
 RfD = Reference dose (mg/kg-day)
 BW = Body weight (kg)
 AT = Averaging time (year)
 IR = Ingestion rate (g/day)
 EF = Exposure frequency (days/year)
 ED = Exposure duration (years)
 IF = Age-adjusted soil ingestion factor (mg-yr/kg-day)
 (using Equation C-5)

$$IF_{adult} = (IR_{adult} \times ED_{adult}) / BW_{adult} \quad (C-5)$$

Groundwater Exposure. The TC (potential PRLs) for chemicals with noncarcinogenic effects for exposure to groundwater via ingestion and inhalation can be estimated (using Equation C-6) as follows:

$$TC_{gw} = \frac{(THI \times BW \times AT \times 365 \text{ days/yr})}{EF \times ED \times (1/RfD_i \times K \times IR_3) + (1/RfD_o \times IR_w)} \quad (C-6)$$

where:

TC_{gw}	=	Target concentration for groundwater ($\mu\text{g/l}$)
THI	=	Target hazard index
BW	=	Body weight (kg)
AT	=	Averaging time (years)
EF	=	Exposure frequency (days/year)
ED	=	Exposure duration (years)
RfD_i	=	Inhalation chronic reference dose (mg/kg-day)
RfD_o	=	Oral chronic reference dose (mg/kg-day)
IR_3	=	Daily inhalation while showering intake rate (L/day)
IR_w	=	Daily water ingestion rate (L/day)
K	=	Volatilization Factor (L/m^3)

Tables C.4.2 and C.4.4 summarize the risk-based concentrations for individual chemical carcinogens based on soil and sediment exposure and groundwater and surface water exposure, respectively. Tables C.4.3 and C.4.5 summarize target concentrations for individual chemical noncarcinogens based on soil and sediment exposure and groundwater and surface water exposure.

The risk-based PRGs use reference doses (RfD) and carcinogenic slope factors from the most recent version of IRIS (EPA, 1992a) and HEAST (EPA, 1992b). Tables C.4.2, C.4.3, C.4.4, and C.4.5 list the RfDs and slope factors used in this assessment. Reference doses and carcinogenic slope factors were not developed for chemicals lacking a toxicity value. The toxicity value for 2,3,7,8-Tetrachlorodibenzodioxin was used for total dioxins (PCDD) and total furans (PCDF). This is a very conservative assumption as no 2,3,7,8 isomer was detected.

A provisional RfD for a JP-4 fuel mixture was used to quantify the toxicity of total petroleum hydrocarbons (TPH). The provisional RfD for JP-4 of 0.08 mg/kg-day was derived by the EPA Superfund Health Risk Technical Support Center (EPA, 1992c). JP-4 and diesel were given a provisional cancer classification of D (not classifiable as a carcinogen due to lack of supporting evidence). The RfD has an uncertainty factor of 10,000 and an overall low level of confidence. In addition to the low levels of uncertainty associated with the derivation of an RfD, there is additional uncertainty because of the difference in composition between the original fuel mixture and the mixture contacted by the receptor. Differential volatilization, water solubility, and biodegradation will alter the composition of the fuel mixture in the environment. Also, further uncertainty exists in that the TPH detected at Shemya Island is assumed composed entirely of JP-4 fuel.

The Alaska Department of Environmental Conservation has issued interim guidance for non-underground storage tank soil cleanup levels of petroleum hydrocarbons (ADEC, 1991). These levels are guidance and not promulgated regulations. The cleanup level

Table C.4.2
Risk-based Concentrations for Carcinogenic Effects
Based on Ingestion of Soil and Sediment by Construction Workers

Chemical	U.S. EPA Carcinogen Classification	Oral Slope Factor (kg-day/mg)	Risk-Based Concentration/Carcinogenic Effects		
			Target Risk: 1E-04 (mg/kg)	Target Risk: 1E-05 (mg/kg)	Target Risk: 1E-06 (mg/kg)
<i>Organics</i>					
1,1,1-Trichloroethane	D	-	-	-	-
1,1-Dichloroethane	-	-	-	-	-
1,2-Dichloroethane	B2	9.0E-02	1.1E-04	1.1E-03	1.1E-02
1,2-Dichloroethene (total)	-	-	-	-	-
2,3,7,8-Tetrachlorodibenzodioxin	B2	1.5E+05	6.6E-03	6.6E-04	6.6E-05
2-Butanone	D	-	-	-	-
2-Methylnaphthalene	-	-	-	-	-
4,4'-DDD	B2	2.4E-01	4.1E+03	4.1E+02	4.1E+01
4,4'-DDE	B2	3.4E-01	2.9E+03	2.9E+02	2.9E+01
4,4'-DDT	B2	3.4E-01	2.9E+03	2.9E+02	2.9E+01
4-Methyl-2-pentanone	-	-	-	-	-
4-Methylphenol	C	-	-	-	-
Acenaphthene	-	-	-	-	-
Acetone	-	-	-	-	-
Anthracene	D	-	-	-	-
Benzene	A	2.9E-02	3.4E+04	3.4E+03	3.4E+02
Benzo(a)anthracene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
Benzo(a)pyrene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
Benzo(b)fluoranthene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
Benzo(g,h,i)perylene	D	-	-	-	-
Benzo(k)fluoranthene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
bis(2-ethylhexyl)phthalate	B2	1.4E-02	7.1E+04	7.1E+03	7.1E+02
Butylbenzylphthalate	C	-	-	-	-
Carbazole	B2	2.0E-02	5.0E+04	5.0E+03	5.0E+02
Carbon disulfide	-	-	-	-	-
Chlorobenzene	D	-	-	-	-
Chloroform	B2	6.1E-03	1.6E+05	1.6E+04	1.6E+03
Chrysene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
Di-n-butylphthalate	D	-	-	-	-
Dibenzo(a,h)anthracene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
Dibenzofuran	D	-	-	-	-
Diethylphthalate	D	-	-	-	-
Endrin ketone	-	-	-	-	-
Ethylbenzene	D	-	-	-	-
Fluoranthene	D	-	-	-	-
Fluorene	D	-	-	-	-
Indeno(1,2,3-cd)pyrene	B2	7.3E+00	1.4E+02	1.4E+01	1.4E+00
Methylene chloride	B2	7.5E-03	1.3E+05	1.3E+04	1.3E+03
Naphthalene	D	-	-	-	-
PCB-1248 (Aroclor 1248)	B2	7.7E+00	1.3E+02	1.3E+01	1.3E+00
PCB-1254 (Aroclor 1254)	B2	7.7E+00	1.3E+02	1.3E+01	1.3E+00

Table C.4.2
Risk-based Concentrations for Carcinogenic Effects
Based on Ingestion of Soil and Sediment by Construction Workers

Chemical	U.S. EPA Carcinogen Classification	Oral Slope Factor (kg-day/mg)	Risk-Based Concentration/Carcinogenic Effects		
			Target Risk: 1E-04 (mg/kg)	Target Risk: 1E-05 (mg/kg)	Target Risk: 1E-06 (mg/kg)
PCB-1260 (Aroclor 1260)	B2	7.7E+00	1.3E+02	1.3E+01	1.3E+00
Phenanthrene	D	-	-	-	-
Phenol	D	-	-	-	-
Pyrene	D	-	-	-	-
Tetrachloroethylene	B2	5.1E-02	1.9E+04	1.9E+03	1.9E+02
Toluene	D	-	-	-	-
Total Xylenes	-	-	-	-	-
TPH-JP4	-	-	-	-	-
Trichloroethylene	B2	1.1E-02	9.0E+04	9.0E+03	9.0E+02
<i>Inorganics</i>					
Aluminum	-	-	-	-	-
Antimony	-	-	-	-	-
Arsenic	A	1.8E+00	5.7E+02	5.7E+01	5.7E+00
Barium	-	-	-	-	-
Beryllium	B2	4.3E+00	2.3E+02	2.3E+01	2.3E+00
Cadmium	-	-	-	-	-
Calcium	-	-	-	-	-
Chromium (Total)	-	-	-	-	-
Cobalt	-	-	-	-	-
Copper	D	-	-	-	-
Cyanide	D	-	-	-	-
Iron	-	-	-	-	-
Lead	B2	-	-	-	-
Magnesium	-	-	-	-	-
Manganese	D	-	-	-	-
Mercury	D	-	-	-	-
Nickel	-	-	-	-	-
Selenium	D	-	-	-	-
Silver	D	-	-	-	-
Thallium	D	-	-	-	-
Vanadium	-	-	-	-	-
Zinc	D	-	-	-	-

Table C.4.2
Risk-based Concentrations for Carcinogenic Effects
Based on Ingestion of Soil and Sediment by Construction Workers

Chemical	U.S. EPA Carcinogen Classification	Oral Slope Factor (kg-day/mg)	Risk-Based Concentration/Carcinogenic Effects		
			Target Risk: 1E-04 (mg/kg)	Target Risk: 1E-05 (mg/kg)	Target Risk: 1E-06 (mg/kg)

EXPOSURE ASSUMPTIONS:

TRL: Target Risk Level:	1E-04	1E-05	1E-06
IR: soil intake rate (mg/day)	50	50	50
AT: Averaging Time (yrs)	70	70	70
BW: Body Weight(kg)	70	70	70
EF: Exposure Frequency (days/year)	180	180	180
ED: Exposure Duration(yrs)	20	20	20
SF: Slope factor (mg/kg-day) ⁻¹			

Information Sources:

Integrated Risk Information System. U.S. EPA 1992a.

Health Effects Assessment Summary Tables, 1992b.

Risk Assessment Guidance for Superfund: Part B, 1991.

Notes:

TPH-JP4 = Total Petroleum Hydrocarbons; assumes TPH is 100% JP-4 fuel mixture and uses provisional RfD (EPA, 1992c).

Table C.4.3
Risk-based Concentrations for Noncarcinogenic Effects
Based on Ingestion of Soil and Sediment by Construction Workers

Chemical	Oral Chronic Reference Dose (mg/kg/day)	Risk-based Concentration Noncarcinogenic Effects Target Hazard Index: 1 (mg/kg)
<i>Organics</i>		
1,1-Dichloroethane	1.0E-01	2.8E + 05
1,2-Dichloroethane	-	-
1,2-Dichloroethene (total)	9.0E-03	2.6E + 04
2,3,7,8-Tetrachlorodibenzodioxin	0.0E + 00	-
2-Butanone	0.0E + 00	-
2-Methylnaphthalene	0.0E + 00	-
4,4'-DDD	0.0E + 00	-
4,4'-DDE	0.0E + 00	-
4,4'-DDT	5.0E-04	1.4E + 03
4-Methyl-2-pentanone	5.0E-02	1.4E + 05
4-Methylphenol	5.0E-02	1.4E + 05
Acenaphthene	6.0E-02	1.7E + 05
Acetone	1.0E-01	2.8E + 05
Anthracene	3.0E-01	8.5E + 05
Benzene	0.0E + 00	-
Benzo(a)anthracene	0.0E + 00	-
Benzo(a)pyrene	0.0E + 00	-
Benzo(b)fluoranthene	0.0E + 00	-
Benzo(g,h,i)perylene	0.0E + 00	-
Benzo(k)fluoranthene	0.0E + 00	-
bis(2-ethylhexyl)phthalate	2.0E-02	5.7E + 04
Butylbenzylphthalate	2.0E-01	5.7E + 05
Carbazole	0.0E + 00	-
Carbon disulfide	1.0E-01	2.8E + 05
Chlorobenzene	2.0E-02	5.7E + 04
Chloroform	1.0E-02	2.8E + 04
Chrysene	0.0E + 00	-
Di-n-butylphthalate	1.0E-01	2.8E + 05
Dibenzo(a,h)anthracene	0.0E + 00	-
Dibenzofuran	0.0E + 00	-
Diethylphthalate	8.0E-01	***
Endrin ketone	0.0E + 00	-
Ethylbenzene	1.0E-01	2.8E + 05
Fluoranthene	4.0E-02	1.1E + 05
Fluorene	4.0E-02	1.1E + 05
Indeno(1,2,3-cd)pyrene	0.0E + 00	-
Methylene chloride	6.0E-02	1.7E + 05
Naphthalene	4.0E-02	1.1E + 05
PCB-1248 (Aroclor 1248)	-	-
PCB-1254 (Aroclor 1254)	-	-
PCB-1260 (Aroclor 1260)	-	-

C.4.10

Table C.4.3
Risk-based Concentrations for Noncarcinogenic Effects
Based on Ingestion of Soil and Sediment by Construction Workers

Chemical	Oral Chronic Reference Dose (mg/kg/day)	Risk-based Concentration Noncarcinogenic Effects Target Hazard Index: 1 (mg/kg)
Phenanthrene	0.0E+00	-
Phenol	6.0E-01	***
Pyrene	3.0E-02	8.5E+04
Tetrachloroethylene	1.0E-02	2.8E+04
Toluene	2.0E-01	5.7E+05
Total Xylenes	2.0E+00	***
TPH-JP4	8.0E-02	2.3E+05
Trichloroethylene	-	-
<i>Inorganics</i>		
Aluminum	-	-
Antimony	4.0E-04	1.1E+03
Arsenic	3.0E-04	8.5E+02
Barium	7.0E-02	2.0E+05
Beryllium	5.0E-03	1.4E+04
Cadmium	5.0E-04	1.4E+03
Calcium	-	-
Chromium (Total)	5.0E-03	1.4E+04
Cobalt	-	-
Copper	-	-
Cyanide	2.0E-02	5.7E+04
Iron	-	-
Lead	-	-
Magnesium	-	-
Manganese	5.0E-03	1.4E+03
Mercury	3.0E-04	8.5E+02
Nickel	2.0E-02	5.7E+04
Selenium	5.0E-03	1.4E+04
Silver	5.0E-03	1.4E+04
Thallium	8.0E-05	2.3E+02
Vanadium	7.0E-03	2.0E+04
Zinc	3.0E-01	8.5E+05

Table C.4.3
Risk-based Concentrations for Noncarcinogenic Effects
Based on Ingestion of Soil and Sediment by Construction Workers

Chemical	Oral Chronic Reference Dose (mg/kg/day)	Risk-based Concentration Noncarcinogenic Effects Target Hazard Index: 1 (mg/kg)
----------	---	--

EXPOSURE ASSUMPTIONS:

IR: Soil/Sediment Ingestion Rate (mg/day)	50
AT: Averaging Time (years)	20
BW: Body Weight (kilograms)	70
EF: Exposure Frequency (days/year)	180
ED: Exposure Duration (years)	20
THI: Target hazard index	1

Information Sources:

Integrated Risk Information System. U.S. EPA 1992a.
 Health Effects Assessment Summary Tables, 1992b.
 Risk Assessment Guidance for Superfund: Part B, 1991.

Notes:

TPH-JP4 = Total Petroleum Hydrocarbons; assumes TPH is 100% JP-4 fuel mixture and uses provisional RfD (EPA, 1992c).

*** = Target Concentration exceeds 100%; chronic exposure may not produce adverse health effects.

Table C.4.4
Risk-based Concentrations for Carcinogenic Effects
Based on Ingestion/Inhalation of Groundwater by AFB Personnel

Chemical	U.S. EPA Carcinogen Classification (oral)	Oral Slope Factor (kg-day/mg)	U.S. EPA Carcinogen Classification (inhalation)	Inhalation Slope Factor (kg-day/mg)	Risk-based Concentration/Carcinogenic Effects		
					Target Risk: 1E-04 (mg/L)	Target Risk: 1E-05 (mg/L)	Target Risk: 1E-06 (mg/L)
Organics							
Acetone	D	-	D	-	-	-	-
Benzene	A	2.900E-02	A	2.9E-02	2.0E-01	2.0E-02	2.0E-03
bis(2-ethylhexyl)phthalate	B2	1.400E-02	B2	-	3.5E+00	3.5E-01	3.5E-02
Carbon disulfide	-	-	-	-	-	-	-
Chloroform	B2	6.100E-03	B2	8.1E-02	8.1E-02	8.1E-03	8.1E-04
Dibenzofuran	D	-	D	-	-	-	-
1,2-Dichloroethene (total)	-	-	-	-	-	-	-
Diethylphthalate	D	-	D	-	-	-	-
Ethylbenzene	D	-	D	-	-	-	-
Fluorene	D	-	D	-	-	-	-
Methylene chloride	B2	7.500E-03	B2	1.6E-03	2.5E+00	2.5E-01	2.5E-02
2-Methylnaphthalene	-	-	-	-	-	-	-
2-Methylphenol	C	-	C	-	-	-	-
4-Methylphenol	C	-	C	-	-	-	-
2,4-Dimethylphenol	-	-	-	-	-	-	-
Naphthalene	D	-	D	-	-	-	-
Nitrate (As NO3)	-	-	-	-	-	-	-
Phenol	D	-	D	-	-	-	-
Toluene	D	-	D	-	-	-	-
TPH-JP4	-	-	-	-	-	-	-
1,1,1-Trichloroethane	D	-	D	-	-	-	-
Trichloroethylene	B2	1.1E-02	B2	1.7E-02	3.6E-01	3.6E-02	3.6E-03
Total Xylenes	D	-	D	-	-	-	-
Inorganics							
Aluminum	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-
Arsenic	A	1.750E+00	A	-	2.8E-02	2.8E-03	2.8E-04
Barium	-	-	-	-	-	-	-

C.4.13

Table C.4.4
Risk-based Concentrations for Carcinogenic Effects
Based on Ingestion/Inhalation of Groundwater by AFB Personnel

Chemical	U.S. EPA Carcinogen Classification (oral)	Oral Slope Factor (kg-day/mg)	U.S. EPA Carcinogen Classification (inhalation)	Inhalation Slope Factor (kg-day/mg)	Risk-based Concentration/Carcinogenic Effects		
					Target Risk: 1E-04 (mg/L)	Target Risk: 1E-05 (mg/L)	Target Risk: 1E-06 (mg/L)
Beryllium	B2	4.3E+00	-	-	1.2E-02	1.2E-03	1.2E-04
Cadmium	-	-	B1	6.3E+00	1.1E-03	1.1E-04	1.1E-05
Calcium	-	-	-	-	-	-	-
Cobalt	-	-	-	-	-	-	-
Chromium (Total)	-	-	A	4.2E+01	1.6E-04	1.6E-05	1.6E-06
Copper	D	-	D	-	-	-	-
Cyanide	D	-	-	-	-	-	-
Iron	-	-	-	-	-	-	-
Lead	B2	-	-	-	-	-	-
Magnesium	-	-	-	-	-	-	-
Manganese	D	-	D	-	-	-	-
Molybdenum	-	-	-	-	-	-	-
Nickel	-	-	-	-	-	-	-
Selenium	D	-	-	-	-	-	-
Silver	D	-	D	-	-	-	-
Thallium	D	-	D	-	-	-	-
Vanadium	-	-	-	-	-	-	-
Zinc	D	-	D	-	-	-	-

EXPOSURE ASSUMPTIONS:

TRL: Target Risk Level:

1E-04 1E-05 1E-06

IRw: Daily water ingestion rate (l/day)

IR3: Daily indoor inhalation rate (m3/day)

AT: Averaging Time (yrs.)

1	1	1
15	15	15
70	70	70

Table C.4.4
Risk-based Concentrations for Carcinogenic Effects
Based on Ingestion/Inhalation of Groundwater by AFB Personnel

Chemical	U.S. EPA Carcinogen Classification (oral)	Oral Slope Factor (kg-day/mg)	U.S. EPA Carcinogen Classification (inhalation)	Inhalation Slope Factor (kg-day/mg)	Risk-based Concentration/Carcinogenic Effects		
					Target Risk: 1E-04 (mg/L)	Target Risk: 1E-05 (mg/L)	Target Risk: 1E-06 (mg/L)
BW: Body weight (kilograms)					70	70	70
EF: Exposure Frequency (days/year)					180	180	180
ED: Exposure Duration (yrs.)					20	20	20
SFi: Inhalation cancer slope factor (mg/kg-day) ⁻¹							
SFo: Oral cancer slope factor (mg/kg-day) ⁻¹							
K: Volatilization factor (L/m ³)					0.5	0.5	0.5

Information Sources:

Integrated Risk Information System. U.S. EPA 1992a.
 Health Effects Assessment Summary Tables, 1992b
 Risk Assessment Guidance for Superfund: Part B, 1991.

C.4.15

Table C.4.5
Risk-based Concentrations for Noncarcinogenic Effects
Based on Ingestion of Groundwater by AFB Personnel

Chemical	Oral Chronic Reference Dose (mg/kg-day)	Inhalation Chronic Reference Dose (mg/kg-day)	Risk-based Concentration Noncarcinogenic Effects Target Hazard Index: 1 (mg/L)
<i>Organics</i>			
Acetone	1.000E-01	-	1.4E+01
Benzene	-	-	-
bis(2-ethylhexyl)phthalate	2.000E-02	-	2.8E+00
Carbon disulfide	1.000E-01	-	1.4E+01
Chloroform	1.000E-02	-	1.4E+00
Dibenzofuran	-	-	-
1,2-Dichloroethene (total)	9.000E-03	-	3.8E-01
Diethylphthalate	8.000E-01	-	1.1E+02
Ethylbenzene	1.000E-01	2.857E-01	3.9E+00
Fluorene	4.000E-02	-	5.7E+00
Methylene chloride	6.000E-02	8.600E-01	5.6E+00
2-Methylnaphthalene	-	-	-
2-Methylphenol	5.000E-02	-	7.1E+00
4-Methylphenol	5.000E-02	-	7.1E+00
2,4-Dimethylphenol	2.000E-02	-	2.8E+00
Naphthalene	4.000E-02	-	5.7E+00
Nitrate (As NO3)	1.600E+00	-	2.3E+02
Phenol	6.000E-01	-	8.5E+01
Toluene	2.000E-01	1.143E-01	2.0E+00
1,1,1-Trichloroethane	9.000E-02	2.857E-01	3.8E+00
Trichloroethylene	-	-	-
Total Xylenes	2.000E+00	-	2.8E+02
<i>Inorganics</i>			
Aluminum	-	-	-
Antimony	4.000E-04	-	5.7E-02
Arsenic	3.000E-04	-	4.3E-02
Barium	7.000E-02	-	9.9E+00
Beryllium	5.000E-03	-	7.1E-01
Cadmium	5.000E-04	-	7.1E-02
Calcium	-	-	-
Chromium (Total)	5.000E-03	-	7.1E-01
Cobalt	-	-	-
Copper	-	-	-
Cyanide	2.000E-02	-	2.8E+00
Iron	-	-	-
Lead	-	-	-
Magnesium	-	-	-
Manganese	5.000E-03	-	7.1E-01
Mercury	3.000E-04	-	4.3E-02

Table C.4.5
Risk-based Concentrations for Noncarcinogenic Effects
Based on Ingestion of Groundwater by AFB Personnel

Chemical	Oral Chronic Reference Dose (mg/kg-day)	Inhalation Chronic Reference Dose (mg/kg-day)	Risk-based Concentration Noncarcinogenic Effects Target Hazard Index: 1 (mg/L)
Molybdenum	5.000E-03	-	7.1E-01
Nickel	2.000E-02	-	2.8E+00
Nitrite	1.000E-01	-	1.4E+01
Selenium	5.000E-03	-	7.1E-01
Silver	5.000E-03	-	7.1E-01
Thallium	8.000E-05	-	1.1E-02
Vanadium	7.000E-03	-	9.9E-01
Zinc	3.000E-01	-	4.3E+01

EXPOSURE ASSUMPTIONS:

THI: target hazard index	1
RfDo: oral chronic reference dose (mg/kg-day)	
RfDi: inhalation chronic reference dose (mg/kg-day)	
BW: body weight (kilograms)	70
AT: averaging time (yr)	20
EF: Exposure Frequency (days/year)	180
ED: Exposure Duration (yr)	20
IR3: Daily indoor inhalation rate (m3/day)	15
IRw: daily water ingestion rate (L/day)	1
K: Volatilization factor (L/m3)	0.5

Information Sources:

Integrated Risk Information System. U.S. EPA 1992a.
 Health Effects Assessment Summary Tables, 1992b.
 Risk Assessment Guidance for Superfund:Part B, 1991.

recommended by this guidance for Shemya AFB is 200 mg/kg (3 orders of magnitude less the derived PRL of 230,000 mg/kg). If the more conservative exposure parameters of a residential scenario are used, a target concentration of 20,000 mg/kg (EPA 1992c) may be used as a PRL. Six samples have TPH levels exceeding 20,000 mg/kg: FT2-128 (29,360 mg/kg), FT2-129 (29,883 mg/kg), FT2-130 (28,034 mg/kg), FT2-134 (20,107 mg/kg), FT2-B3 (23,649 mg/kg), and FT2-B6 (20,573 mg/kg). Each of these exceedances occurs at IRP site FT-2, Aircraft Mockup.

5.0 Risk Characterization

Of the several human health PRLs calculated or available from literature, one was selected for comparison to concentrations of chemicals detected at each site. The lowest risk-based soil and sediment PRL concentration was selected from the two effects (i.e., carcinogenic or noncarcinogenic). Table C.5.1 lists the most stringent PRL for cleanup of soil and sediment.

If an MCL or a proposed MCL exists for a chemical, it was first used as the selected PRL. If no MCL exists, the lowest concentration (i.e., most stringent and most protective) of the following was used as the selected PRL:

- risk-based concentration (noncarcinogenic effects)
- risk-based concentration (carcinogenic effects)
- ambient water quality criteria

Table C.5.2 lists the selected PRL used for cleanup of groundwater.

Every detected chemical from the 1992 field investigation samples (using both onsite and offsite laboratory analytical results) was compared to the most stringent PRL (Tables C.5.1 and C.5.2). Table C.5.3 list those samples, and the sites where they are found, that exceed the selected PRL. Chemicals not found in Table C.5.3 do not exceed the most stringent PRL.

Table C.5.1
Selected Soil and Sediment Preliminary Risk Levels

Page 1 of 3

Chemical	Most Stringent Preliminary Risk Level (mg/kg)	Source
<i>Organics</i>		
Acenaphthene	170,000	NC
Acetone	280,000	NC
Anthracene	850,000	NC
Benzene	340	C
Benzo(a)anthracene	1.4	C
Benzo(a)pyrene	1.4	C
Benzo(b)fluoranthene	1.4	C
Benzo(k)fluoranthene	1.4	C
Benzo(g,h,i)perylene	-	-
bis(2-Ethylhexyl)phthalate	710	C
2-Butanone	-	-
Butylbenzylphthalate	570,000	NC
Carbazole	500	C
Carbon disulfide	280,000	NC
Chlorobenzene	57,000	NC
Chloroform	28,000	NC
Chrysene	1.4	C
4,4'-DDE	41	C
4,4'-DDT	29	C
4,4'-DDD	29	C
Dibenzofuran	-	-
Dibenzo(a,h)anthracene	14	C
1,1-Dichloroethane	280,000	NC
1,2-Dichloroethane	110	C
1,2-Dichloroethene (total)	26,000	NC
Di-n-butylphthalate	280,000	NC
Diethylphthalate	***	NC
2,3,7,8-Tetrachlorodibenzodioxin	7E-05	C
Ethylbenzene	280,000	NC
Endrin ketone	-	-
Fluoranthene	110,000	NC
Fluorene	110,000	NC
Indeno(1,2,3-cd)pyrene	1.4	C
Methylene chloride	1300	C

C.5.2

Table C.5.1
Selected Soil and Sediment Preliminary Risk Levels

Page 2 of 3

Chemical	Most Stringent Preliminary Risk Level (mg/kg)	Source
<i>Organics (Continued)</i>		
2-Methylnaphthalene	-	-
4-Methylphenol	140,000	NC
4-Methyl-2-pentanone	140,000	NC
Naphthalene	110,000	NC
Phenanthrene	-	-
Phenol	***	NC
Pyrene	85,000	NC
PCB-1260 (Aroclor 1260)	1.3	C
PCB-1254 (Aroclor 1254)	1.3	C
PCB-1248 (Aroclor 1248)	1.3	C
PCB-1248 (Aroclor 1248)	1.3	C
Tetrachloroethylene	190	C
Toluene	570,000	NC
1,1,1-Trichloroethane	260,000	NC
Trichloroethylene	900	C
TPH-JP4	230,000	
Total Xylenes	***	NC
<i>Inorganics</i>		
Aluminum	-	-
Antimony	1100	NC
Arsenic	5.7	C
Barium	2.0×10^{-5}	NC
Beryllium	2.3	C
Cadmium	1400	NC
Calcium	-	-
Chromium (Total)	14,000	NC
Cobalt	-	-
Copper	-	-
Cyanide	57,000	NC
Iron	-	-
Lead	-	-
Magnesium	-	-
Manganese	1.4×10^{-3}	NC

C.5.3

Table C.5.1
Selected Soil and Sediment Preliminary Risk Levels

Page 3 of 3

Chemical	Most Stringent Preliminary Risk Level (mg/kg)	Source
<i>Inorganics (Continued)</i>		
Mercury	850	NC
Nickel	5.7×10^{-4}	NC
Selenium	1.4×10^{-4}	NC
Silver	1.4×11^{-4}	NC
Thallium	2.3×10^{-2}	NC
Vanadium	2.0×10^{-4}	NC
Zinc	570,000	NC

Notes:

MCL = Safe Drinking Water Act Maximum Contaminant Level

MCL-p = Proposed Safe Drinking Water Act Maximum Contaminan

WQC = Ambient Water Quality Criteria

NC = Risk-based Concentration/Noncarcinogenic Effects

C = Risk-based Concentration/Carcinogenic Effects

*** = Target Concentration exceeds 100%, chronic exposure may not produce adverse effects.

Table C.5.2
Selected Groundwater Preliminary Risk Levels

Page 1 of 2

Chemical	Most Stringent Preliminary Risk Level (mg/L)	Source
<i>Organics</i>		
Acetone	14	NC
Benzene	0.005	MCL
bis(2-Ethylhexyl)phthalate	0.004	MCL-p
Carbon disulfide	14	NC
Chloroform	0.00019	WQC
Dibenzofuran	-	-
1,2-Dichloroethene (total)	0.1	MCL
Diethylphthalate	110	NC
Ethylbenzene	0.7	MCL
Fluorene	0.0000031	WQC (total PAHs)
Methylene chloride	0.00019	QC (total halomethanes)
2-Methylnaphthalene	0.0000031	WQC (total PAHs)
2-Methylphenol	7.1	NC
4-Methylphenol	7.1	NC
2,4-Dimethylphenol	2.8	NC
Naphthalene	5.7	NC
Nitrate (As NO3)	230	NC
Phenol	3.5	WQC
Toluene	2	NC
1,1,1-Trichloroethane	0.2	MCL
Trichloroethylene	0.005	MCL
Total xylenes	10	MCL
<i>Inorganics</i>		
Aluminum	-	-
Antimony	.010	MCL-p
Arsenic	0.05	MCL
Barium	1	MCL
Beryllium	0.0000039	WQC
Cadmium	0.005	MCL
Calcium	-	-
Chromium (Total)	0.05	MCL
Cobalt	-	-
Copper	-	-

C.5.5

Table C.5.2
Selected Groundwater Preliminary Risk Levels

Page 2 of 2

Chemical	Most Stringent Preliminary Risk Level (mg/L)	Source
<i>Inorganics (Continued)</i>		
Cyanide	0.2	MCL-p
Iron	-	-
Magnesium	-	-
Manganese	0.71	NC
Mercury	0.043	NC
Molybdenum	0.71	NC
Nickel	0.1	MCL-p
Selenium	.010	MCL
Silver	0.05	MCL
Thallium	0.002	MCL-p
Vanadium	0.99	NC
Zinc	42	NC

Notes:

MCL = Safe Drinking Water Act Maximum Contaminant Level

MCL-p = Proposed Safe Drinking Water Act Maximum Contaminant Level

WQC = Ambient Water Quality Criteria

NC = Risk-based Concentration/Noncarcinogenic Effects

C = Risk-based Concentration/Carcinogenic Effects

Table C.5.3
COPCs that Exceed Preliminary Risk Levels (PRLs)

Page 1 of 2

Sample No.	Depth (ft.)	Chemical	Concentration	Qualifier	Laboratory	PRL Concentration	PRL Source
GROUNDWATER COPCs			(ug/L)			(ug/L)	
<i>FT-2: Aircraft Mockup</i>							
FT2-ER		Chloroform	2	J	OFFSITE	0.19	WQC
FT2-W1		Benzene	512		ONSITE	5	MCL
FT2-W1DL		Benzene	710	D	OFFSITE	5	MCL
FT2-W1DL		Methylene Chloride	420	BDJ2	OFFSITE	0.19	WQC
<i>Water Gallery</i>							
CM1		Trichloroethylene	7.2		ONSITE	5	MCL
SO2		Trichloroethylene	5.1		ONSITE	5	MCL
WGW4		Trichloroethylene	24.9		ONSITE	5	MCL
WG4		Trichloroethylene	16.4		ONSITE	5	MCL
WGW4		Trichloroethylene	22.2		ONSITE	5	MCL
WG7		Trichloroethylene	17.6		ONSITE	5	MCL
WGW7		Trichloroethylene	18.2		ONSITE	5	MCL
A-WGW3		Antimony	31.1		WC	10	MCL-p
A-WGW3		Beryllium	0.75		WC	0.0039	WQC
A-WGW3		Chromium	53.4		WC	50	MCL
A-WGW3		Manganese	2170		WC	710	NC
A-WGW4		Antimony	31.1		WC	10	MCL-p
A-WGW4		Beryllium	0.5		WC	0.0039	WQC
A-WGW4		Manganese	892		WC	710	NC
A-WGW4		Trichloroethene (TCE)	15		WC	5	MCL
<i>Well 410</i>							
A-W410		Chloroform	17		WC	0.19	WQC
<i>Background Samples (BG-BW01)</i>							
A-BKGD1		Antimony	31.1		WC	10	MCL-p
A-BKGD1		Beryllium	0.5		WC	0.0039	WQC
A-BKGD1		Manganese	803		WC	710	NC

C.5.7

Table C.5.3
COPCs that Exceed Preliminary Risk Levels (PRLs)

Page 2 of 2

Sample No.	Depth (ft.)	Chemical	Concentration	Qualifier	Laboratory	PRL Concentration (ug/kg)	PRL Source
SURFACE SOILS COPCs			(ug/kg)			(ug/kg)	
<i>FT-1: Lightning Strike</i>							
FT1T21.0	1	PCDD	0.3		OFFSITE	0.066	C
FT1T310.5	0.5	PCDD	4.9		OFFSITE	0.066	C
FT1T310.5	0.5	PCDF	1.1		OFFSITE	0.066	C
FT1T41.2	1.2	PCDD	0.9		OFFSITE	0.066	C
FT1T41.2	1.2	PCDF	0.1		OFFSITE	0.066	C
FT1T41.2D	1.2	PCDD	15		OFFSITE	0.066	C
FT1T41.2D	1.2	PCDF	1.1		OFFSITE	0.066	C
<i>SW10: Barrel Bay</i>							
SW10-114	0.5	Arsenic	6000		OFFSITE	5700	C
SW10-116	0.5	Arsenic	6900		OFFSITE	5700	C
<i>SW-12: Scrap Metal Disposal Site</i>							
SW12-117	0.5	Arsenic	10300		OFFSITE	5700	C
SW12-118	0.5	Arsenic	6100		OFFSITE	5700	C
SW12-120	0.5	Arsenic	8300		OFFSITE	5700	C
<i>WG: Water Gallery</i>							
WG-109	0.5	Crysene	35000		OFFSITE	1400	C
WG-109	0.5	Benzo(a)anthracene	43000		OFFSITE	1400	C
WG-109	0.5	Benzo(b)fluoranthene	61000		OFFSITE	1400	C
WG-109	0.5	Benzo(k)fluoranthene	21000		OFFSITE	1400	C
WG-109	0.5	Benzo(a)pyrene	36000		OFFSITE	1400	C
WG-109	0.5	Indeno(1,2,3-cd)pyrene	20000		OFFSITE	1400	C
WG-109	0.5	Arsenic	6800		OFFSITE	5700	C
<i>PS1B: West Dock JP-4 Spill (Old Cobra Dane)</i>							
O-A-PS1B	2	Arsenic	6100		WC	5700	C
SEDIMENT COPCs			(ug/kg)			(ug/kg)	
<i>Background Samples</i>							
BSSD-126	0.5	Arsenic	8300		OFFSITE	5700	C

Notes:

MCL = Safe Drinking Water Act Maximum Contaminant Level

MCL-p = Proposed Safe Drinking Water Act Maximum Contaminant Level

WQC = Ambient Water Quality Criteria

NC = Risk-based Concentration/Noncarcinogenic Effects

C = Risk-based Concentration/Carcinogenic Effects

ONSITE = Close Support Laboratory (on Shemya Island)

OFFSITE = Contract Laboratory Program Methodology (samples sent to offsite lab)

WC = Woodward-Clyde Offsite Data

J = estimated value

D = dilution

B = blank contamination

C.5.8

6.0 References

40 CFR 264.94. "RCRA Groundwater Protection Standards." U.S. Code of Federal Regulations.

40 CFR 141. "Safe Drinking Water Act and the National Primary Drinking Water Regulations." U.S. Code of Federal Regulations.

Alaska Administrative Code (AAC). "Alaska Water Quality Standards." 18 AAC 70.

Alaska Administrative Code (AAC). "Alaska Drinking Water Regulations." 18 AAC 80.

Air Force Systems Command (AFSC). 1987. The Installation Restoration Program Toxicity Guide. Volume 4. Air Force Systems Command, Aerospace Medical Division, Wright-Patterson Air Force Base, Ohio. 1987.

Alaska Department of Environmental Conservation (ADEC). 1991. Alaska Department of Environmental Conservation Interim Guidance for NON-UST Contaminated Soil Cleanup Levels. July 17, 1991.

CH2M HILL. 1991. Installation Restoration Program (IRP) Field Investigation Plan: Shemya Air Force Base, Alaska. October 4, 1991.

Dragun, J. 1988. The Soil Chemistry of Hazardous Materials. Hazardous Materials Control Research Institute.

Lyman, W. J., W. F. Reehl, and D. H. Rosenblatt. 1982. Handbook of Chemical Property Estimation Methods: Environmental Behavior of Organic Compounds.

Mackay. 1992. Illustrated Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals. Lewis Publishers. 1992.

Roy, W. R., and R. A. Griffin. 1985. "Mobility of Organic Solvents in Water-saturated Soil Materials." *Environmental Geology and Water Sciences*, 7(1985):241-247.

U.S. Environmental Protection Agency (EPA). 1989. Risk Assessment Guidance for Superfund: Volume I—Human Health Evaluation Manual (Part A). Publication EPA/540/1-89/002. December 1989.

U.S. Environmental Protection Agency (EPA). 1991. Risk Assessment Guidance for Superfund: Volume I--Human Health Evaluation Manual (Part B, Development of Risk-Based Preliminary Remediation Goals). Publication 9285.7-01B. December 1991.

U.S. Environmental Protection Agency (EPA). 1992a. Integrated Risk Information System: U.S. EPA IRIS Data Base. Office of Research and Development. (92-12).

U.S. Environmental Protection Agency (EPA). 1992b. Health Effects Summary Tables, FY 92. Office of Research and Development, DERR 9200.6-303 (92-1).

U.S. Environmental Protection Agency (EPA). 1992c. Memorandum by Joan S. Dollarhide, Associate Director, Superfund Health Risk Technical Support Center, Chemical Mixtures Assessment Branch. Dated March 24, 1992 on "Oral Reference Dose and Oral Slope Factors for JP-4, JP-5, Diesel Fuel, and Gasoline (AVGAS)." [McChord AFB (Wash Rack/Treatment), Tacoma, WA]

Appendix D
Ecological Risk Assessment

CONTENTS

APPENDIX D—ECOLOGICAL RISK ASSESSMENT	D.1.1
1.0 INTRODUCTION	D.1.1
1.1 PURPOSE	D.1.1
1.2 ECOLOGICAL RISK ASSESSMENT GUIDANCE	D.1.1
1.3 SITE DESCRIPTION AND ECOLOGICAL SETTING	D.1.2
1.4 SCOPE AND ORGANIZATION OF THE ECOLOGICAL RISK ASSESSMENT	D.1.2
2.0 IDENTIFICATION OF ECOLOGICAL CHEMICALS OF POTENTIAL CONCERN	D.2.1
2.1 DATA SUMMARY	D.2.1
3.0 ECOLOGICAL EXPOSURE ASSESSMENT	D.3.1
3.1 CHARACTERIZATION OF THE EXPOSURE SETTING AND IDENTIFICATION OF POTENTIAL RECEPTORS	D.3.1
3.1.1 FT-1 Lightning Strike	D.3.2
3.1.2 FT-2 Aircraft Mockup	D.3.2
3.1.3 SW-4 Barrel Dump Site	D.3.2
3.1.4 SW-5 Hospital Lake	D.3.11
3.1.5 SW-10 Barrel Bay	D.3.11
3.1.6 SW-12 Scrap Metal Dump	D.3.11
3.1.7 Water Gallery	D.3.11
3.1.8 PS-1A New Cobra Dane	D.3.11
3.2 POTENTIAL ECOLOGICAL EXPOSURE PATHWAYS	D.3.12
3.2.1 Terrestrial Receptor Exposure Pathways	D.3.13
3.2.2 Aquatic Receptor Exposure Pathways	D.3.16
3.3 AREA OF CONCERN CONTAMINANT SOURCES AND EXPOSURE POINT CONCENTRATIONS	D.3.16

CONTENTS (Continued)

4.0	DEVELOPMENT OF PRELIMINARY RISK LEVELS	D.4.1
4.1	PRELIMINARY RISK LEVELS PROTECTIVE OF TERRESTRIAL LIFE	D.4.1
4.1.1	Surface Soil and Sediment	D.4.1
4.1.2	Surface Soil	D.4.19
4.1.3	Surface Water and Groundwater	D.4.20
4.1.4	Sediment	D.4.20
4.2	PRELIMINARY RISK LEVELS PROTECTIVE OF AQUATIC LIFE ..	D.4.23
4.2.1	Surface Water and Groundwater	D.4.23
4.2.2	Sediment	D.4.23
5.0	ECOLOGICAL TOXICITY ASSESSMENT	D.5.1
5.1	TOXICITY ASSESSMENT OF INORGANIC ECOLOGICAL CHEMICALS OF POTENTIAL CONCERN	D.5.1
5.2	TOXICITY ASSESSMENT OF ORGANIC CHEMICALS	D.5.3
5.2.1	Polychlorinated Biphenyls	D.5.3
5.2.2	Polycyclic Aromatic Hydrocarbons	D.5.4
6.0	ECOLOGICAL RISK CHARACTERIZATION	D.6.1
6.1	METHODS FOR TERRESTRIAL RECEPTOR RISK CHARACTERIZATION	D.6.1
6.2	METHODS FOR AQUATIC RECEPTOR RISK CHARACTERIZATION	D.6.3
6.3	SITE-SPECIFIC RISK CHARACTERIZATIONS	D.6.4
6.3.1	FT-1	D.6.4
6.3.2	FT-2	D.6.4
6.3.3	SW-4	D.6.10
6.3.4	SW-5	D.6.10
6.3.5	SW-10	D.6.10
6.3.6	SW-12	D.6.18

CONTENTS (Continued)

6.3.7	Water Gallery	D.6.18
6.3.8	PS-1A	D.6.27
6.3.9	PS-1B	D.6.27
7.0	Uncertainty Analysis	D.7.1
8.0	References	D.8.1

TABLES

D.2.1	Data Summary of FT-1 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.3
D.2.2	Data Summary of FT-2 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.5
D.2.3	Data Summary of SW-4 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.7
D.2.4	Data Summary of SW-5 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.8
D.2.5	Data Summary of SW-10 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.9
D.2.6	Data Summary of SW-12 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.12
D.2.7	Data Summary of WG Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.15
D.2.8	Data Summary of PS-1A Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.19
D.2.9	Data Summary of PS-1B Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.21
D.2.10	Data Summary of PS-2 Site-related Ecological Chemicals of Potential Concern by Matrix	D.2.23
D.3.1	Potential Ecological Receptors for Shemya Island	D.3.3
D.3.2	Water and Food Intake Values for Terrestrial Organisms Used in Establishing Acceptable Water Concentrations	D.3.14
D.4.1	Toxicity Values for the Ecological Chemicals of Potential Concern	D.4.2
D.4.2	Uncertainty Factors Used in Establishing Adjusted Lowest Observable Adverse Effect Level Values for Soil and Sediment Dose Comparisons	D.4.6

TABLES (Continued)

D.4.3	Uncertainty Factors Used in Establishing No Adverse Effect Water Criteria Concentrations	D.4.7
D.4.4	Toxicity Criteria Used for the Calculation of Preliminary Risk Levels for the Protection of Terrestrial Life	D.4.8
D.4.5	Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern	D.4.13
D.4.6	Preliminary Risk Levels for the Protection of Terrestrial Receptors Exposed to Surface Water and Groundwater	D.4.21
D.4.7	Sediment Criteria for Inorganic Ecological Chemicals of Potential Concern	D.4.24
D.4.8	Sediment Criteria for Organic Ecological Chemicals of Potential Concern	D.4.25
D.4.9	Sediment Standards for the State of Washington	D.4.27
D.6.1	Comparison of FT-1 Soil Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.5
D.6.2	Comparison of FT-1 Groundwater Ecological Chemicals of Potential Concern Terrestrial Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.7
D.6.3	Comparison of FT-2 Soil Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.8
D.6.4	Comparison of FT-2 Groundwater Ecological Chemicals of Potential Concern Terrestrial Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.9

TABLES (Continued)

D.6.5	Comparison of SW-4 Soil Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.11
D.6.6	Comparison of SW-5 Sediment Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.12
D.6.7	Comparison of SW-5 Surface Water Ecological Chemicals of Potential Concern Terrestrial Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.13
D.6.8	Comparison of SW-5 Surface Water and Sediment Ecological Chemicals of Potential Concern Aquatic Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.14
D.6.9	Comparison of SW-10 Soil and Sediment Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.15
D.6.10	Comparison of SW-10 Groundwater Ecological Chemicals of Potential Concern Terrestrial Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.17
D.6.11	Comparison of SW-10 Surface Water and Sediment Ecological Chemicals of Potential Concern Aquatic Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values . . .	D.6.19
D.6.12	Comparison of SW-12 Soil and Sediment Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.20
D.6.13	Comparison of SW-12 Groundwater Ecological Chemicals of Potential Concern Terrestrial Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.22

TABLES (Continued)

D.6.14	Comparison of WG Soil Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.23
D.6.15	Comparison of WG Groundwater Ecological Chemicals of Potential Concern Terrestrial Receptor Exposure Concentrations to Applicable Preliminary Risk Level Values	D.6.26
D.6.16	Comparison of PS-1A Soil Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.28
D.6.17	Comparison of PS-1B Soil Ecological Chemicals of Potential Concern Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable Preliminary Risk Level Values	D.6.30

Appendix D Ecological Risk Assessment

1.0 Introduction

The Appendix presents results of the baseline ecological risk assessment (BERA) for the Shemya Air Force Base Installation Restoration Program (IRP) sites. The purpose of the BERA is to provide ecologically-based preliminary risk levels (PRLs) for site-related contaminants at the IRP sites that will be used to help determine the need for remedial action (or no action) at the sites. By using contaminant exposure levels considered to have no observable adverse effects (NOAELs), with water quality criteria and sediment quality criteria as the ecological end points, biota should not suffer any adverse effects such as behavioral, reproductive, or survivability changes if exposed to chemical concentrations at these levels.

This BERA focuses on potential exposure to ecological receptors attributable to the sites at Shemya in the absence of remedial action (the no-action alternative) (EPA, 1989a). Living receptors at or near the sites are identified and characterized, and the potential adverse effects resulting from exposure of these receptors to the ecological chemicals of potential concern (ECOPCs) are evaluated.

1.1 Purpose

This BERA addresses the potential exposure at the sites at Shemya under current land uses. The no-action alternative assumes that no corrective actions have taken place. Evaluation of this no-action alternative is required under Section 300.430(d) of the NCP (EPA, 1990).

1.2 Ecological Risk Assessment Guidance

This BERA was performed in accordance with the following guidance:

- Risk Assessment Guidance for Superfund—Volume II, Environmental Evaluation Manual (EPA, 1989a)
- Framework for Ecological Risk Assessment (EPA, 1992a)
- Peer Review Workshop Report on a Framework for Ecological Risk Assessment (EPA, 1992b)
- Summary Report on Issues in Ecological Risk Assessment (EPA, 1991)

- Ecological Assessment of Hazardous Waste Sites—A Field and Laboratory Reference (EPA, 1989b)

1.3 Site Description and Ecological Setting

A comprehensive description of the physical, geological, hydrologic, and topographic characteristics of Shemya Air Force Base is presented in Sections 1-3 of this report. In addition, past disposal practices and potential sources of contamination are reviewed in Sections 1-3. For the purposes of this document, only those components of the Shemya sites that are relevant to the BERA are addressed.

1.4 Scope and Organization of the Ecological Risk Assessment

The purpose of this BERA is to provide PRLs for contaminants of concern that would protect ecological receptors (EPA, 1989a; 1992a; 1992b). There is no standard approach for conducting a BERA, and BERAs may vary widely from site to site.

Surface soils, surface water, groundwater, and sediment data are evaluated in this assessment. In addition, only direct exposure pathways such as ingestion are addressed. Exposure resulting from bioaccumulation and biomagnification (biotic fate and transport), as well as abiotic fate and transport of contaminant-transfer in various media, were not evaluated.

Exposure to ecological receptors is addressed at each site and is predominantly of a quantitative nature. Where possible, exposure estimates are calculated. Because of the limitations imposed by a lack of literature on contaminant effects to specific species, the risk characterization of the BERA is qualitatively addressed by comparing observed concentrations of ECOPCS to PRL values.

The overall objectives of the exposure assessment (EA) include:

- Identifying existing and potential animal and plant populations on Shemya Island
- Evaluating exposure pathways from contaminated media
- Conservatively evaluating onsite exposure point concentrations
- Assessing if exposure to ECOPCs could constitute a potential adverse effect to terrestrial and aquatic populations based on qualitative comparisons to literature-derived toxicity values as identified in the toxicity assessments

- Identifying areas of uncertainty involved with the development of the EA

The BERA was completed with the following limitations:

- The BERA does **not** separate natural variability of exposure or effects between or within species
- The BERA is **not** able to account for mechanical influences (environmental and site-specific disturbances) that contribute to variability in exposure
- Data comparisons to toxicity values may **not** be species-specific comparisons because of a lack of literature values
- Applied uncertainty estimates may **not** encompass the entire uncertainty associated with the quantitative assessment

The components of this BERA include:

- Ecological setting description
- Identification of the actual and potential ecological receptors that can become exposed to site-related contamination
- Identification of ECOPCs at each IRP site
- Characterization of the ecological exposure pathways
- Exposure concentrations and calculation of possible intake rates (for terrestrial organisms only)
- Discussion of uncertainty

The BERA is based on the following assumptions:

- No remedial actions have been taken.
- For the purposes of exposure and risk calculation, contaminant concentrations will not and have not changed over time.
- The media of concern include soil, sediment, groundwater, and surface water.

- All detections of organic analytes are assumed to be caused by site-related contamination.
- The maximum observed concentration of ECOPCs was used as the potential ecological exposure point concentration, and observed maximum concentrations were assumed to be 100 percent bioavailable for organism uptake.
- Terrestrial, freshwater aquatic, and marine aquatic organisms were considered as potential ecological receptors of concern.
- Ingestion rates from available literature for terrestrial species were used for the quantification of exposure to incidental soil ingestion, surface water ingestion, and incidental sediment ingestion.
- Aquatic organism exposure was assessed by comparing observed maximum concentrations of chemicals of concern in surface water and groundwater to established federal water quality criteria.
- Aquatic organism risk posed by exposure to contaminated sediment was assessed by comparing observed maximum concentrations of sediment ECOPCs to sediment quality criteria derived from ambient water quality criteria.

The BERA is based on data collected on the media of concern during the 1992 field investigation. These media include surface soils, surface water, groundwater, and sediment.

Uncertainties and lack of standard procedures necessitate an overall qualitative approach to establish ecological risk for the areas of concern. The BERA addresses the surface soil, surface water, groundwater, and sediment ECOPCs and identifies those chemicals that may contribute the majority of adverse effects resulting from exposure to ecological receptors.

2.0 Identification of Ecological Chemicals of Potential Concern

Wildlife species are very transient organisms and exposure can potentially occur throughout the island. It was assumed in this BERA that receptor exposure could occur throughout a site, but not between sites. Since most terrestrial organisms do not limit their activity to a defined area, it is assumed that an organism can be exposed to surface soil, surface water, groundwater, and sediment, where available, throughout a site. Therefore, the data for each media were not analyzed for spatial or temporal distribution patterns within individual sites for this BERA. Instead, all data points were combined for each media and analyzed within each site for frequency of detection and toxicity criteria applicable to the selection of ECOPCs. The maximum observed concentration was used as the exposure point concentration for conservative purposes in the quantitative evaluation of exposure.

In this investigation, gaseous chemicals are not regarded as ECOPCs. Because of the lack of inhalation toxicity data for ecological organisms in the literature, qualitative or quantitative determinations of exposure associated with gaseous chemicals would be highly uncertain. In addition, total petroleum hydrocarbons (TPH), common ions, and essential nutrients were not evaluated due to the lack of applicable toxicological data. Instead, individual hydrocarbon data points (i.e., xylene, toluene, etc.) were quantitatively addressed where available. The common ions and essential nutrients that were excluded from the evaluation include sodium, calcium, potassium, and magnesium. These elements were excluded due to their common occurrence and low toxicity.

Data evaluation protocols discussed as part of the data evaluation task explicitly consider data uses relative to performing the exposure assessment (EA). Surface soil, surface water, groundwater, and sediment data are screened to identify data of acceptable quality for use in the BERA and to eliminate low-quality data points that are unacceptable for risk calculations.

2.1 Data Summary

Initial screening of data to derive ECOPCs was dependent upon the frequency of detection. Chemicals that were infrequently detected may be anomalies in the data because of sampling errors, analytical errors, or other problems and, therefore, may not be site-related (EPA, 1989a). For this BERA, a detection frequency of 1 has been selected as the level required to retain a chemical as an ECOPC.

Data summaries for those chemicals that exceeded a detection frequency of 1 are summarized by area of concern and presented in Tables D.2.1 through D.2.10. These tables present the minimum, and maximum concentrations for organic and inorganic analytes at the RP sites by matrix (surface water, groundwater, sediment, and soil).

Table D.2.1
Data Summary of FT-1 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 2

Contaminant	Min.	Max.
Groundwater	($\mu\text{g/l}$)	($\mu\text{g/l}$)
1,1,1-Trichloroethane	-	15
bis(2-ethylhexyl)phthalate	-	3
Toluene	-	1
Copper	-	72
Zinc	-	163
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
2-Methylnaphthalene	-	130
Anthracene	-	98
Benzene	12	1,600
Benzo(a)anthracene	76	220
Benzo(a)pyrene	63	200
Benzo(b)fluoranthene	62	280
Benzo(k)fluoranthene	58	84
bis(2-ethylhexyl)phthalate	50	790
2-Butanone	-	1,400
Butylbenzylphthalate	69	130
Chlorobenzene	-	4
Chrysene	140	440
Dibenzofuran	-	92
Di-n-butylphthalate	89	710
Di-n-octylphthalate	-	68
Ethylbenzene	1	8,300

Table D.2.1
Data Summary of FT-1 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 2

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Fluoranthene	73	490
Fluorene	-	78
Indeno(1,2,3-cd)pyrene	-	69
2-Methylnaphthalene	-	130
Naphthalene	46	500
Phenanthrene	-	630
Polychlorinated Dibenzodioxins	0.3	15
Polychlorinated Dibenzofurans	0.1	1.1
Pyrene	40	460
Toluene	48	51,000
Total Petroleum Hydrocarbons	15.7	2,300,000
Xylene (Total)	5	130,000
Cadmium	2000	11,000
Chromium	22300	87,300
Copper	66700	238,000
Lead	41300	671,000
Zinc	76200	1,320,000
Total Organic Carbon	650000	31000000

Table D.2.2
Data Summary of FT-2 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 2

Contaminant	Min.	Max.
Groundwater	(ug/l)	(ug/l)
2-Methylnaphthalene	-	1
2-Methylphenol	-	18
2,4-Dimethylphenol	-	31
4-Methylphenol	-	31
Benzene	512	710
Chloroform	-	2
Ethylbenzene	511	680
Methylene Chloride	-	420
Naphthalene	-	32
Phenol	7	11
Toluene	896	1,400
Xylene (Total)	1.5	3,600
Chromium	13.9	15.6
Zinc	59.0	87.6
Surface Soil	(µg/kg)	(µg/kg)
2-Methylnaphthalene	5,700	45,000
4-Methyl-2-pentanone	-	15
Benzene	10.9	33,000
bis(2-ethylhexyl)phthalate	700	2,200
Chlorobenzene	5	18

Table D.2.2
Data Summary of FT-2 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 2

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Dibenzofuran	1,200	1,700
Ethylbenzene	13.1	130,000
Fluoranthene	-	1,500
Fluorene	1,200	1,600
n-Nitrosodiphenylamine	-	950
Methylene Chloride	3,300	3,400
Acetone	-	5,400
2-Butanone	-	8,600
Tetrachloroethene	3,300	8,600
Styrene	-	22
Fluorene	760	2,200
Naphthalene	3,500	27,000
Phenanthrene	940	1,900
Pyrene	-	1,300
Toluene	31.7	610,000
Xylene (Total)	19	799,000
Chromium	-	23,300
Total Petroleum Hydrocarbons	18,700	29,883,000
Total Organic Carbon	6,330,000	15,600,000
Lead	2,000	2,300

Table D.2.3
Data Summary of SW-4 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
2-Butanone	-	82
4,4'-DDD	2.5	5.6
Aroclor 1260	59	120
Arsenic	-	25,000
Benzo(a)pyrene	-	43
bis(2-ethylhexyl)phthalate	54	91
Cadmium	-	2,000
Chromium	17,700	22,800
Copper	48,300	66,100
Di-n-butylphthalate	110	160
Indeno(1,2,3-cd)pyrene	-	45
Lead	17,000	534,000
Methylene Chloride	53	54
Nickel	-	96,500
PCE (Tetrachloroethylene)	-	2
Total Petroleum Hydrocarbons	6,800	371,000
Toluene	-	38
Xylene (Total)	39.8	42.3
Zinc	48,400	143,000

Table D.2.4
Data Summary of SW-5 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Contaminant	Min.	Max.
Surface Water	($\mu\text{g/l}$)	($\mu\text{g/l}$)
bis(2-ethylhexyl)phthalate	-	1
Copper	-	48,800
Nitrate/nitrite	38	1410
Total Petroleum Hydrocarbons	60	170
Zinc	-	77,200
Sediment	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
2-Butanone	-	18
4,4'-DDD	-	5.8
Barium	-	94,700
bis(2-ethylhexyl)phthalate	95	130
Chromium	-	34,100
Copper	-	55,300
Endrin Ketone	-	21
Fluoranthene	-	46
Fluorene	-	78
Lead	-	36,100
Pyrene	-	88
Total Petroleum Hydrocarbons	51,000	274,000
Zinc	58,700	101,000

Table D.2.5
Data Summary of SW-10 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 3

Contaminant	Min.	Max.
Groundwater	($\mu\text{g/l}$)	($\mu\text{g/l}$)
2-Methylnaphthalene	-	12
Acetone	-	11
Benzene	3	5.1
bis(2-ethylhexyl)phthalate	-	2
Carbon Disulfide	-	3
Dibenzofuran	-	1
Diethylphthalate	-	2
Ethylbenzene	-	1.6
Fluorene	-	2
Naphthalene	-	8
Toluene	1.7	2.9
Xylenes (Total)	0.75	6.2
Chromium	25	48.3
Copper	71.8	202
Zinc	147	700
Sediment	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
bis(2-ethylhexyl)phthalate	-	110
Arsenic	-	6,900
Chromium	-	17,800
Lead	-	23,800
Zinc	-	59,500

Table D.2.5
Data Summary of SW-10 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 3

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
2-Butanone	590	1,000
4',4'-DDD	2.1	15
4',4'-DDE	-	3.2
4'4'-DDT	2.5	12
Anthracene	-	68
Aroclor 1248	-	150
Aroclor 1254	23	110
Aroclor 1260	-	630
Benzo(a)anthracene	-	130
Benzo(a)pyrene	-	330
Benzo(b)fluoranthene	-	300
Benzo(g,h,i)perylene	-	270
Benzo(k)fluoranthene	-	270
bis(2-ethylhexyl)phthalate	46	5,300
Carbon Disulfide	-	1
Chloroform	-	2
Chrysene	-	390
Di-n-butylphthalate	-	46
Endrin ketone	-	2.1
Ethylbenzene	-	50.2
Fluoranthene	45	68

Table D.2.5
Data Summary of SW-10 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 3 of 3

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Arsenic	5,200	53,100
Barium	47,200	561,000
Cadmium	1,800	5,800
Chromium	18,300	81,300
Copper	39,900	892,000
Indeno(1,2,3-cd)pyrene	-	300
Lead	7,700	1,740,000
Methylene Chloride	13	14
Nickel	33,800	57,300
Phenanthrene	-	54
Pyrene	44	80
Toluene	2	62.8
Total Petroleum Hydrocarbons	6,400	16,244,000
Total Organic Carbon	3,480,000	5,550,000
Xylenes (Total)	-	170.9
Zinc	36,400	2,110,000

Table D.2.6
Data Summary of SW-12 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 3

Contaminant	Min.	Max.
Groundwater	($\mu\text{g/l}$)	($\mu\text{g/l}$)
Acetone	-	21
Aroclor 1254	-	87
Benzene	1.6	1.9
bis(2-ethylhexyl)phthalate	-	1
Carbon Disulfide	2	4
Toluene	-	0.5
Arsenic	62.8	66.9
Chromium	19.2	80.7
Copper	52.9	255
Zinc	163	369
Sediment	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
4',4'-DDD	-	5.3
4',4'-DDE	-	2.6
4',4'-DDT	-	17
bis(2-ethylhexyl)phthalate	-	390
Antimony	-	8,700
Arsenic	-	5,500
Cadmium	-	2,100
Chromium	-	31,800
Copper	-	371,000
Lead	-	381,000
Zinc	-	567,000

Table D.2.6
Data Summary of SW-12 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 3

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
4,4'-DDD	-	15
4,4'-DDE	-	2.7
4,4'-DDT	2.5	34
Acetone	-	21
Aroclor 1254	-	87
Aroclor 1260	36	89
Benzene	9	49
bis(2-ethylhexyl)phthalate	40	780
Carbon Disulfide	-	2
Chloroform	-	1
Diethylphthalate	58	130
Di-n-butylphthalate	46	5,000
Ethylbenzene	-	780
Methylene Chloride	-	19
Toluene	19	990
Total Petroleum Hydrocarbons	7,800	18,484,000
Xylenes (Total)	5	2,680
Arsenic	6,100	18,000
Antimony	-	8,700
Barium	65,300	331,000
Chromium	20,400	59,300
Copper	41,300	13,7000

Table D.2.6
Data Summary of SW-12 Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 3 of 3

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Lead	12,900	335,000
Nickel	32,600	62,400
Zinc	44,400	419,000
Total Organic Carbon	3,850,000	4,970,000

Table D.2.7
Data Summary of WG Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 4

Contaminant	Min.	Max.
Groundwater	($\mu\text{g/l}$)	($\mu\text{g/l}$)
1,2 Dichloroethene	-	1.0
Aluminum	1,610	30,400
Antimony	-	31.10
Arsenic	-	2.00
Barium	24.8	135
Benzene	-	1.8
Beryllium	0.5	0.75
Cadmium	-	2.7
Chromium	-	53.4
Cobalt	8.8	19.8
Copper	21.6	200
Ethylbenzene	1	3.7
Iron	1,370	17,900
Lead	6.2	8.8
Manganese	892	2,170
Mercury	-	0.2
Nickel	19.4	32.2
Selenium	-	2.9
Silver	1.5	2.0
Thallium	1.6	1.6
Trichloroethene	1.4	24.9
Toluene	1.0	16.3

Table D.2.7
Data Summary of WG Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 4

Contaminant	Min.	Max.
Groundwater	($\mu\text{g/l}$)	($\mu\text{g/l}$)
Vanadium	6.3	56
Xylenes (Total)	4.3	18.9
Zinc	18.3	150
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
2-Butanone	-	200
2-Methylnaphthalene	-	1,200
4-Methylphenol	-	1,000
Acenaphthene	-	15,000
Acetone	-	920
Anthracene	-	38,000
Aroclor 1254	-	36
Benzo(a)anthracene	-	43,000
Benzo(a)pyrene	-	36,000
Benzo(b)fluoranthene	-	61,000
Benzo(g,h,i)perylene	-	13,000
Benzo(k)fluoranthene	-	21,000
bis(2-ethylhexyl)phthalate	140	2,100
Carbazole	-	13,000
Chloromethane	-	57
Chrysene	-	35,000
Di-n-butylphthalate	130	860
Dibenzo(a,h)anthracene	-	4,800

Table D.2.7
Data Summary of WG Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 3 of 4

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Dibenzofuran	-	12,000
Endrin Ketone	-	3.4
Ethylbenzene	2.1	41
Fluoranthene	59	96,000
Fluorene	-	21,000
Indeno(1,2,3-cd)pyrene	-	20,000
Methylene Chloride	-	17
Naphthalene	-	1,600
Phenanthrene	-	110,000
Phenol	-	160
Pyrene	-	79,000
Toluene	3	98.7
Total Petroleum Hydrocarbons	19,100	3,203,000
Xylenes (Total)	1.5	347.4
Aluminum	9,500,000	41,900,000
Antimony	7,600	24,800
Arsenic	3,300	6,800
Barium	18,800	60,500
Beryllium	140	400
Cadmium	1,400	2,200
Chromium	11,200	38,700
Cobalt	750	9,100

Table D.2.7
Data Summary of WG Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 4 of 4

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Copper	24,700	122,000
Cyanide	610	2,000
Iron	11,500,000	28,200,000
Lead	1,700	137,000
Manganese	147,000	379,000
Mercury	110	400
Nickel	15,300	23,000
Selenium	710	2,000
Silver	420	690
Thallium	390	1,300
Vanadium	42,500	216,000
Zinc	30,000	86,300

Table D.2.8
Data Summary of PS-1A Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 2

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Aluminum	-	24,400,000
Anthracene	-	120
Antimony	-	6,800
Arsenic	-	2,700
Barium	-	99,400
Benzo(a)anthracene	-	380
Benzo(a)pyrene	-	390
Benzo(b)fluoranthene	-	170
Benzo(g,h,i)perylene	-	360
Beryllium	-	220
bis(2-ethylhexyl)phthalate	-	300
Cadmium	-	2,200
Carbazole	-	82
Chromium	-	264,000
Chrysene	-	410
Cobalt	-	27,300
Copper	-	133,000
Cyanide	-	550
Dibenzo(a,h)anthracene	-	79
Fluoranthene	-	1,100
Indeno(1,2,3-cd)pyrene	-	440

Table D.2.8
Data Summary of PS-1A Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 2

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Iron	-	59,600,000
Lead	-	29,100
Manganese	-	1,160,000
Mercury	-	170
Nickel	-	23,600
Phenanthrene	-	530
Pyrene	-	600
Silver	-	1,600
Thallium	-	570
Toluene	-	1
Total Petroleum Hydrocarbons	-	34,000
Vanadium	-	213,000
Zinc	-	95,900

Table D.2.9
Data Summary of PS-1B Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 1 of 2

Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
1,1-Dichloroethane	-	65
1,1-Dichloroethene	-	97
1,1,1-Trichloroethane	4	600
1,2-Dichloroethane	-	31
4,4'-DDD	-	34
Aluminum	-	41,300,000
Antimony	-	16,300
Arsenic	-	6,100
Barium	-	453,000
Beryllium	-	620
bis(2-ethylhexyl)phthalate	-	88,000
Cadmium	-	28,900
Chromium	-	62,600
Cobalt	-	7,700
Copper	-	2,630,000
Cyanide	-	3,300
Iron	-	28,200,000
Lead	-	1,430,000
Manganese	-	210,000
Mercury	-	5,400
Nickel	-	672,000
Selenium	-	16,100
Silver	-	497,000
Thallium	-	1,600

Table D.2.9
Data Summary of PS-1B Site-Related Ecological
Chemicals of Potential Concern by Matrix

Page 2 of 2

Contaminant	Min.	Max.
Surface soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Total Petroleum Hydrocarbons	21,000,000	129,000,000
Trichloroethene	10	2,200
Vanadium	-	43,430
Zinc	-	6,110,000

Table D.2.10 Data Summary of PS-2 Site-Related Ecological Chemicals of Potential Concern by Matrix		
Contaminant	Min.	Max.
Surface Soil	($\mu\text{g/kg}$)	($\mu\text{g/kg}$)
Total Petroleum Hydrocarbons	48,000	439,000

3.0 Ecological Exposure Assessment

An exposure assessment is a qualitative and/or quantitative determination of the exposure magnitude, frequency, duration, and route. The magnitude of exposure is determined by measuring or estimating the amount of an agent available at the exchange boundaries (i.e., the lungs, intestines, and skin). Exposure can occur when contaminants migrate from the site to an exposure point (a location where receptors can come into contact with contaminants) or when a receptor comes into direct contact with waste or contaminated media at the site.

The site description, receptor identification, and physical characterization of the individual sites are presented in Subsections 3.1 and 3.2.

This section presents a summary of the following information:

- Characterization of the exposure setting
- Identification of ecological receptors
- Identification of ecological exposure pathways
- Exposure point concentrations

3.1 Characterization of the Exposure Setting and Identification of Potential Receptors

Terrestrial and aquatic ecological receptors can use sites as habitat and be exposed to site-related contamination through everyday activities. For terrestrial organisms, locating and procuring food and water can lead to the ingestion of contaminated surface water and biotic food sources (from bioconcentration and bioaccumulation from contaminated sources) as well as the incidental ingestion of contaminated soils and sediment. In addition, onsite activities of terrestrial organisms can result in their inhalation of dusts and dermal contact with soil, sediment, and surface water. Aquatic organisms can become exposed to surface water, groundwater seeps into surface water, and sediment through direct contact.

Because of the transient nature of wildlife species (i.e., large mammals and birds), their distribution and habitat range can be extensive. Small mammals occurring on Shemya (occupying somewhat restricted areas for their home range) can become exposed to onsite contamination through various media. Therefore, those species known to occur close to the sites can also become receptors to contamination from natural feeding or migrational activity.

A list of species believed to occur in the area of the Shemya IRP sites is presented in Table D.3.1. Not all of these species have been observed on Shemya Island. The species list presents possible or potential ecological receptors that could occur on the island because of the presence of certain habitat types. The following sections provide an overview of the ecological setting and the receptors that have been observed or could occur within and outside of the IRP sites investigated during the 1992 field season at Shemya.

3.1.1 FT-1 Lightning Strike

FT-1 is located on the southwest corner of Shemya Island, adjacent to the coast. The area is a pit with steep sides, exposed soil, and sparse vegetation. Pictures taken in 1991 show the area strewn with scrap lumber. Although no wildlife data are available for this specific site, Asian ducks and emperor geese are known to roost on the beach and rocks along the southern side of the island, just east of FT-1. The FT-1, SW-10, and SW-12 locations are within the same bay area.

3.1.2 FT-2 Aircraft Mockup

FT-2 is located near the western coast of Shemya Island, approximately 600 feet from Upper Lake. The area is characterized by exposed rocky soils and little or no vegetation. Recent photographs show scrap wood and metal strewn around the area, as well as large above-ground storage tanks adjacent to the site. Although no wildlife data are available for this area, it is important to note that Upper Lake provides habitat for migrating Aleutian geese during the periods April-June and August-October.

3.1.3 SW-4 Barrel Dump Site

SW-4 is located on the northern edge of the island, adjacent to the coast. The area is somewhat rocky with areas of exposed soils and a vegetation community of grasses and forbes characteristic of the Aleutian tundra. There is a low-lying wet area near the site, and the soil exhibits an iron-stained horizon. The northern coast of the island where SW-4 is located has been documented as important habitat for migrating Asiatic songbirds during the periods May-June and September-November. Pelagic and red-faced cormorants maintain nesting colonies on the bluffs and islets west of the site, and harbor seals utilize habitat provided by offshore rocks along the coastline. Small numbers of tufted and horned puffins have also been observed in the immediate vicinity of SW-4.

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 1 of 8

Common Name	Scientific Name
Birds	
Aleutian Tern	<i>Sterna aleutica</i>
American Wigeon	
Ancient Murrelet	<i>Synthliboramphus antiquus</i>
Arctic Loon	<i>Garvia arctica</i>
Arctic Tern	<i>Sterna paradisaea</i>
Arctic Warbler	<i>Phylloscopus borealis borealis</i>
Baird's Sandpiper	<i>Calidris bairdii</i>
Bald Eagle	
Bank Swallow	
Bar-tailed Godwit	<i>Limosa lapponica (baueri)</i>
Barn Swallow	<i>Hirundo rustica (gutturialis)</i>
Barrow's Goldeneye	
Bean Goose	<i>Anser fabalis (serrirostris)</i>
Black Scoter	<i>Melanitta nigra (americana)</i>
Black Oystercatcher	
Black-backed Wagtail	
Black-bellied Plover	<i>Pluvialis squatarola</i>
Black-footed Albatross	
Black-headed Gull	<i>Larus ridibundus</i>
Black-legged Kittiwake	<i>Rissa tridactyla</i>
Black-tailed Godwit	<i>Limosa limosa melanuroides</i>
Black-winged Stilt	<i>Himantopus himantopus</i>
Blue-winged Teal	
Bohemian Waxwing	
Brambling	<i>Fringilla montifringilla</i>
Brant	
Bristle-thighed Curlew	<i>Numenius tahitiensis</i>

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 2 of 8

Common Name	Scientific Name
Broad-billed Sandpiper	<i>Limicola falcinellus sibirica</i>
Brown Shrike	<i>Lanius cristatus lucionensis</i>
Buff-breasted Sandpiper	<i>Tryngites subruficollis</i>
Bufflehead	<i>Bucephala albeola</i>
Bullfinch	<i>Pyrrhula pyrrhula cassinii</i>
Canada Goose	<i>Branta canadensis (leucopareia)</i>
Canvasback	<i>Aythya valisineria</i>
Cassin's Auklet	
Common Buzzard	<i>Buteo buteo</i>
Common Cuckoo	
Common Eider	<i>Somateria mollissima (v.-nigrum)</i>
Common Goldeneye	<i>Bucephala clangula (ssp.)</i>
Common Loon	<i>Gavia immer</i>
Common Merganser	<i>Mergus merganser merganser</i>
Common Murre	<i>Uria aalge (inornata)</i>
Common Pochard	<i>Aythya ferina</i>
Common Raven	<i>Corvus corax (principalis)</i>
Common Redpoll	<i>Acanthis flammea (flammea)</i>
Common Ringed Plover	
Common Rose-finch	<i>Carpodacus erythrinus (grebnitskii)</i>
Common Sandpiper	<i>Actitis hypoleucos</i>
Common Snipe	<i>Gallinago gallinago gallinago</i>
Common Tern	<i>Sterna hirundo longipennis</i>
Cook's Petrel	
Crested Auklet	<i>Aethia cristatella</i>
Curlew Sandpiper	<i>Calidris ferruginea</i>
Dotterel	<i>Eudromias morinellus</i>
Double-crested Cormorant	

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 3 of 8

Common Name	Scientific Name
Dunlin	<i>Calidris alpina (sakhulina)</i>
Dusky Thrush	<i>Turdus naumanni eunomus</i>
Dusky Warbler	<i>Phylloscopus fuscatus fuscatus</i>
Emperor Goose	<i>Anser canagicus</i>
Eurasian Kestrel	<i>Falco tinnunculus interstinctus</i>
Eurasian Wigeon	<i>Anas penelope</i>
Eye-browed Thrush	<i>Turdus obscurus</i>
Falcated Teal	<i>Anas falcata</i>
Far Eastern Curlew	<i>Numenius madagascariensis</i>
Fork-tailed Storm-Petrel	<i>Oceanodroma furcata (furcata)</i>
Gadwall	<i>Anas strepera (strepera)</i>
Garganey	<i>Anas querquedula</i>
Glaucous Gull	<i>Larus hyperboreus (ssp.)</i>
Glaucous-winged Gull	<i>Larus glaucescens</i>
Grasshopper-warbler	<i>Locustella (sp.)</i>
Gray Bunting	<i>Emberiza variabilis</i>
Gray Wagtail	<i>Motacilla cinerea robusta</i>
Gray-cheeked Thrush	<i>Catharus minimus (minimus)</i>
Gray-crowned Rosy-Finch	<i>Leucosticte arctoa (griseonucha)</i>
Gray-spotted Flycatcher	<i>Muscicapa griseisticta</i>
Gray-tailed Tattler	<i>Heteroscelus brevipes</i>
Great Knot	<i>Calidris tenuirostris</i>
Greater Scaup	<i>Aythya marila</i>
Greater White-fronted Goose	
Greater Yellowlegs	<i>Tringa melanoleuca</i>
Green Sandpiper	<i>Tringa oceropus</i>
Green-winged Teal	<i>Anas crecca (nimia)</i>
Greenshank	<i>Tringa nebularia</i>

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 4 of 8

Common Name	Scientific Name
Gyr Falcon	
Harlequin Duck	<i>Histrionicus histrionicus</i>
Hawfinch	
Herring Gull	<i>Larus argentatus vegae</i>
Hoary Redpoll	<i>Acanthis hornemanni (exilipes)</i>
Hooded Merganser	
Horned Grebe	<i>Podiceps auritus (auritus)</i>
Horned Lark	<i>Eremophila alpestris flava</i>
Horned Puffin	<i>Fratercula corniculata</i>
Indian Tree Pipit	<i>Anthus hodgsoni yunnanensis</i>
King Eider	
Kittlitz's Murrelet	
Lapland Longspur	<i>Calcarius lapponicus (alascensis)</i>
Laysan Albatross	<i>Diomedea immutabilis</i>
Leach's Storm-petrel	
Least Auklet	
Least Sandpiper	
Lesser Golden Plover	<i>Pluvialis dominica (fulva)</i>
Lesser Scaup	
Lesser Yellowlegs	
Little Bunting	<i>Emberiza pusilla</i>
Long-billed Dowitcher	<i>Limnodromus scolopaceus</i>
Long-tailed Jaeger	<i>Stercorarius longicaudus</i>
Long-toed Stint	<i>Calidris subminuta</i>
Mallard	<i>Anas platyrhynchos (platyrhynchos)</i>
Marbled Murrelet	
Marsh Hawk	<i>Circus cyaneus (ssp.)</i>
Merlin	

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 5 of 8

Common Name	Scientific Name
Mew Gull	<i>Larus canus kamtschatschensis</i>
Mongolian Plover	<i>Charadrius mongolus (stegmanni)</i>
Northern Fulmar	<i>Fulmarus glacialis</i>
Northern Harrier	
Northern Phalarope	<i>Phalaropus lobatus</i>
Northern Shoveler	<i>Anas chyeata</i>
Northern Shrike	
Oldsquaw	<i>Clangula hyemalis</i>
Oriental Greenfinch	<i>Carduelis sinica kawarahiba</i>
Pacific Loon	
Parakeet Auklet	
Parasitic Jaeger	<i>Stercorarius parasiticus</i>
Pectoral Sandpiper	<i>Calidris melanotos</i>
Pelagic Cormorant	<i>Phalacrocorax pelagicus (pelagicus)</i>
Peregrine Falcon	<i>Falco peregrinus (pealei)</i>
Pigeon Guillemot	<i>Cephus columba (kaiurka)</i>
Pin-tailed Snipe	
Pintail	<i>Anas acuta (acuta)</i>
Polynesian Tattler	<i>Heteroscelus brevipes</i>
Pomarine Jaeger	<i>Stercorarius pomarinus</i>
Red Knot	<i>Calidris canutus (canutus)</i>
Red Phalarope	<i>Phalaropus fulicarius</i>
Red-breasted Flycatcher	<i>Ficedula parva albicilla</i>
Red-breasted Merganser	<i>Mergus serrator</i>
Red-faced Cormorant	<i>Phalacrocorax urile</i>
Red-legged Kittiwake	<i>Rissa brevirostris</i>
Red-necked Grebe	<i>Podiceps grisegena (holbollii)</i>
Red-necked Phalarope	

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 6 of 8

Common Name	Scientific Name
Red-throated Loon	<i>Gavia stellata</i>
Red-throated Pipit	<i>Anthus cervinus</i>
Reed Bunting	<i>Emeriza schoeniclus pyrrhulina</i>
Ring-necked Duck	
Rock Ptarmigan	
Rock Sandpiper	<i>Calidris ptilocnemis (quarta)</i>
Rock Sandpiper	<i>Calidris ptilocnemis (couesi)</i>
Ross' Gull	<i>Rhodostethia rosea</i>
Rosy Finch	<i>Leucosticte arctoa</i>
Rough-legged Hawk	<i>Buteo lagopus (ssp.)</i>
Ruddy Turnstone	<i>Arenaria interpres (interpres)</i>
Ruff	<i>Philomachus pugnax</i>
Rufous-necked Sandpiper	<i>Calidris ruficollis</i>
Rufous-necked Stint	<i>Calidris ruficollis</i>
Rustic Bunting	<i>Emberiza rustica latifascia</i>
Sabine's Gull	
Sanderling	<i>Calidris alba</i>
Sandhill Crane	
Savanna Sparrow	<i>Ammodramus sandwichensis anthinus</i>
Semipalmated Plover	<i>Charadrius semipalmatus</i>
Semipalmated Sandpiper	<i>Calidris pusilla</i>
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>
Short-eared Owl	<i>Asio flammeus (flammeus)</i>
Short-tailed Shearwater	<i>Puffinus tenuirostris</i>
Siberian Accentor	<i>Prunella montanella (ssp.)</i>
Siberian Rubythroat	<i>Erithacus calliope</i>
Skylark	<i>Alauda arvensis pekinensis</i>
Slaty-backed Gull	<i>Larus schistisagus</i>

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 7 of 8

Common Name	Scientific Name
Smew	<i>Mergus albellus</i>
Snow Bunting	<i>Plectrophenax nivalis (townsendi)</i>
Snowy Owl	<i>Nyctea scandiaca</i>
Song Sparrow	<i>Melospiza melodia maxima</i>
Sooty Flycatcher	<i>Muscicapa sibirica sibirica</i>
Sooty Shearwater	
Spot-billed Duck	
Spotted Redshank	<i>Tringa erythropus</i>
Steller's Eider	<i>Polysticta stelleri</i>
Surf Scoter	
Temminck's Stint	<i>Calidris temminckii</i>
Terek Sandpiper	<i>Xenus cinereus</i>
Thick-billed Murre	<i>Uria lomvia (arra)</i>
Townsend's Warbler	<i>Dendroica townsendi</i>
Tree Swallow	
Tufted Duck	<i>Aythya fuligula</i>
Tufted Puffin	<i>Lunda cirrhata</i>
Tundra Swan	
Violet-green Swallow	<i>Tachycineta thalassina lepida</i>
Wandering Tattler	<i>Heteroscelus incanus</i>
Water Pipit	<i>Anthus spinoletta (rubescens)</i>
Water Pipit	<i>Anthus spinoletta japonicus</i>
Western Grebe	
Western Sandpiper	<i>Calidris mauri</i>
Wheatear	<i>Oenanthe oenanthe (oenanthe)</i>
Whimbrel	<i>Numenius phaeopus (variegatus)</i>
Whiskered Auklet	
White Wagtail	<i>Motacilla alba lugens</i>

Table D.3.1
Potential Ecological Receptors for Shemya Island

Page 8 of 8

Common Name	Scientific Name
White-rumped Swift	<i>Apus pacificus pacificus</i>
White-winged Crossbill	
White-winged Scoter	<i>Melanitta fusca (deglandi)</i>
Whooper Swan	
Winter Wren	<i>Troglodytes troglodytes (meligerus)</i>
Wood Sandpiper	<i>Tringa glareola</i>
Wood Warbler	<i>Phylloscopus sibilatrix</i>
Yellow Wagtail	<i>Motacilla flava simillima</i>
Yellow-billed Loon	<i>Gavia adamsii</i>
Mammals	
Harbor Seal	<i>Phoca vitulina</i>
Steller's or Northern Sea Lion	<i>Eumetopias jubatus</i>

3.1.4 SW-5 Hospital Lake

Hospital Lake is a small interior lake located in the northeastern corner of Shemya Island. The lake is approximately 100 to 200 feet wide with lushly vegetated banks and a very small littoral zone.

3.1.5 SW-10 Barrel Bay

SW-10 is located on the southern coast of the island, just east of FT-1. The area is highly disturbed, very rocky, and sparsely vegetated. Scraps of metal, wire, wood, and rubber litter the area. Slightly inland of the coast, the cover material on the dump supports a thin cover of grasses and forbes. The beaches and rocky areas immediately east of this site have been documented to provide habitat for Asian ducks and emperor geese.

3.1.6 SW-12 Scrap Metal Dump

SW-12 is located just east of SW-10 on the southern shore of the island. The site is similar to SW-10, with rocky, sparsely vegetated ground and areas of exposed soils. The coastal side of SW-12 exhibits steep, rocky slopes. As with SW-10, the beaches and rocky areas near SW-12 provide habitat for Asian ducks and emperor geese.

3.1.7 Water Gallery

The water gallery is located in the southeastern interior of the island, near the airstrip. The site is characterized by densely vegetated grass/sedge areas and little or no exposed soil. A paved road and hanger are situated immediately upgradient of the water gallery. No wildlife data are available for this specific site; however, large numbers of glaucous-winged gulls have been observed resting on the nearby airstrip during fall months, and the densely vegetated wet areas could potentially provide important habitat for birds and other wildlife.

3.1.8 PS-1A New Cobra Dane

PS-1A is located near the northern coast on the highest point of the island (approximately 300 feet above sea level). The PS-1A area includes a building and gravel parking lot surrounded by a fence. The adjacent areas support a vegetation community characteristic of Aleutian tundra and provide important habitat for migrating Asiatic songbirds during the periods May-June and September-November.

3.2 Potential Ecological Exposure Pathways

An exposure assessment cannot be conducted unless the following components have been identified:

- A contaminant source
- An exposure pathway
- A receptor organism

In several ecological exposure scenarios at the IRP sites, the above components may not exist.

Potential receptor groups (aquatic, terrestrial, vegetation, microfauna, etc.) have been addressed for the exposure assessment. Only those routes related to terrestrial organism ingestion of contaminated media (surface soil, surface water, groundwater, and sediment) and aquatic organism contact with contaminated surface water, and sediment were quantitatively addressed for the following reasons:

- Dermal and inhalation exposure cannot be quantitatively addressed because of the lack of applicable literature.
- The lack of applicable literature for comparing vegetation and microflora responses to certain chemical exposures minimizes the ability to compare chemical concentrations and determine effects.
- Site-specific species lists do not exist; therefore, certain receptors may not reside at the IRP sites.
- Groundwater seeps into aquatic habitats were not identified.

Potential exposure to terrestrial ecological receptors is created by exposure to contaminated surface soil, surface water, groundwater, and sediment. Potential exposure routes for terrestrial and aquatic receptors occurring within the sites include:

- Incidental ingestion or dermal adsorption of chemicals from surface soil, surface water, groundwater seeps, and sediment.
- Inhalation of volatilized chemicals and wind-eroded particulate matter (dust) originating from the surface soil, surface water, and sediment.
- Ingestion of surface water by animals, including subsequent bioaccumulation to predator species.

- Uptake of surface water by plants and subsequent exposure to herbivores ingesting contaminated plants.
- Uptake of chemicals associated with bottom litter or sediment by microbia through the natural process of decay.
- Terrestrial organism incidental ingestion or dermal adsorption of chemicals from groundwater seeps.
- Aquatic organism (freshwater and marine) incidental contact and exposure via all routes to contaminated surface water, groundwater seeps, and sediments (where applicable).

The relative importance of exposure routes depends primarily on land use and the receptors that use the area. Media interactions that influence exposure were not examined in detail in this assessment. Only those exposures that result from direct contact (via ingestion) to a particular matrix were addressed in this assessment.

Under the no-action definition for current land use conditions, the following assumptions were made for the BERA:

- Land use conditions are presented as they were at the time of the evaluation.
- Interim remedial actions are of no consequence for the exposure evaluation.

Based upon these assumptions, the following sections summarize the potential exposure pathways to terrestrial and aquatic receptors.

3.2.1 Terrestrial Receptor Exposure Pathways

3.2.1.1 Incidental Surface Soil Ingestion. The possibility of terrestrial organisms becoming exposed to soil-related contaminants through incidental ingestion may exist. Because of the wide variety of species with widely varying feeding habits, soil ingestion is highly variable. Animals that burrow or prey on burrowing species, such as earthworms or voles, are more likely to ingest greater quantities of soil than those that prey on species with little ground contact. Birds ingest grit and any accompanying soil for use in the gizzard. Very little research has been done to quantify soil ingestion by wildlife species. For this reason, soil ingestion was addressed for mice and rats whose soil intake rates were derived from research conducted by Garten (1980).

In order to conservatively estimate exposure to contaminated soil, the following assumptions were made:

- All surface soil consumed was derived from a contaminated source.
- The maximum observed concentration represents the potential exposure concentration.

It was assumed that the potential future exposure scenarios of receptor organisms exposed to soil would be the same as the current scenarios.

3.2.1.2 Surface Water and Groundwater Ingestion. Organisms are directly exposed to contaminants by ingestion of surface water and groundwater seeps (where appropriate). The surface water pathway is important for animals such as small mammals, waterfowl, and birds that might use surface water at the IRP sites as a drinking water source. Bioconcentration, as defined for aquatic organisms, is not applicable to terrestrial organisms because tissue concentrations are not a direct function of water concentration. However, uptake of contaminants from surface water consumption can occur, with accumulation rates depending on the amount of water ingested daily and the concentration of contaminants in the water supply. Daily water consumption rates for certain terrestrial animals are reported in Table D.3.2.

Table D.3.2 Water and Food Intake Values for Terrestrial Organisms Used in Establishing Acceptable Water Concentrations		
Species	Daily Water Consumption (l/kg bw/day)	Daily Soil Consumption (mg soil/kg bw/day)
Rat (adult)	0.125	280 ^a
Mouse	0.2	740 ^b
Rabbit (adult)	0.165	NA
^a Derived by Garten (1980) for the hispid cotton rat (<i>Sigmodon hispidus</i>). ^b Derived by Garten (1980) for the white-footed mouse (<i>Peromyscus leucopus</i>). Notes: l/kg bw/day = liters/kilogram body weight/day. g/kg bw/day = grams/kilogram body weight/day. NA = Not available in the literature reviewed. Source: Sax, 1992; Garten, 1980.		

Many small mammals are adapted to a semiarid environment and do not consume surface water on a regular basis. For example, cottontail rabbits obtain most of their water needs from metabolic water (Turkowski, 1975). Therefore, criteria developed from water ingestion rates for laboratory animals represent a conservative estimate because many small mammals at Shemya do not consume surface water on a regular basis, and many animals that consume surface water could have access to uncontaminated water supplies.

Water consumption data for laboratory rats, mice, and rabbits were used to represent small mammals at the sites that may consume surface water (Sax, 1992). Assuming toxicity from dietary exposure is similar to toxicity resulting from ingesting contaminated water, a water concentration associated with no observable adverse effects is derived from toxicity data for dietary intake and water ingestion rates for a similar species.

These values are estimates of the total water requirements; thus, since some of these water needs are met with food intake, the water ingestion rates are probably sufficiently conservative to represent most species. In addition, mobile species have access to multiple sources of drinking water and thus effectively dilute the exposure concentrations. To determine conservative estimates of exposure, the following assumptions were made:

- All water consumption was from a contaminated source.
- All water consumption was from a surface water source and not a groundwater-supplied source.
- The maximum observed concentration represents the potential exposure concentration.

3.2.1.3 Incidental Sediment Ingestion. Organisms that forage and hunt in the riparian zone or within the beach areas can be exposed to sediment contamination through the dermal and incidental ingestion of sediment. In addition, the possibility of incidental ingestion of sediment can occur from sediment disturbances while the animal is drinking. Although the onsite existence of waterfowl and riparian organisms is limited, the possibility of certain terrestrial animals ingesting sediment is addressed quantitatively in this assessment using the same techniques for quantifying soil ingestion.

To determine conservative estimates of exposure to contaminated sediment, the following assumptions were made:

- All sediment consumed was derived from a contaminated source.
- The maximum observed concentration represents the potential exposure concentration.

It was assumed that the possible future exposure scenarios of receptor organisms exposed to sediment would be the same as at present.

3.2.2 Aquatic Receptor Exposure Pathways

3.2.2.1 Direct Contact with Surface Water and Groundwater Seeps. Aquatic organisms (freshwater and marine) can be exposed to water-borne contamination through a variety of routes. Direct contact exposure routes include ingestion, dermal (integument) contact, and inhalation. There is also the potential at some sites for groundwater to seep into surface water. For this reason, aquatic organisms may come in contact with groundwater-related contamination. Aquatic organism exposure to groundwater was not addressed since this exposure pathway is not complete. Further investigation of groundwater transport needs to be completed before this pathway is identified.

3.2.2.2 Direct Contact with Sediments. Aquatic organisms can be exposed to contaminated sediments by incidentally (or purposely) ingesting sediment, inhaling sediment particulates, and contacting the sediment with the integument (dermal areas). All of these routes are categorized as "direct contact" in this assessment.

3.3 Area of Concern Contaminant Sources and Exposure Point Concentrations

Exposure concentrations for terrestrial and aquatic receptors were based on the data derived from surface soils, surface water, groundwater, and sediment. The quantitative determination of exposure concentrations was based on the following assumptions:

- Data represent current conditions at the IRP sites at Shemya.
- The maximum concentration is representative of the potential exposure concentration to ecological receptors.

- Maximum concentrations of the various chemicals of concern in the media of concern represent exposure point concentrations for acute and chronic exposure durations.
- Assessment of risk was ultimately addressed for only those species for which applicable NOAEL, LOAEL, or LD50 values have been derived (derived from ingestion studies exposing rats, mice, or rabbits).

Assumptions used to estimate specific chemical terrestrial organism intake rates that are derived for surface soil ECOPCs, sediment ECOPCs, and water quality criteria include:

- Organisms are exposed to the maximum contaminant level in the various matrices.
- The observed concentrations of contaminants in the various matrices are 100 percent bioavailable.
- Organisms ingest surface soil, surface water, groundwater, and sediment strictly from the affected area with the observed maximum chemical concentration.

Terrestrial and aquatic organism exposure and risk are addressed for each site, but, only for those exposure routes considered feasible.

4.0 Development of Preliminary Risk Levels

Preliminary risk levels (PRLs) for the protection of ecological receptors are not widely established. Ambient water quality criteria provided by state and federal agencies are enforced for the protection of marine and freshwater organisms. These criteria were used as the basis for establishing aquatic life PRLs. PRLs for terrestrial organisms were based upon literature-derived toxicity values from laboratory and field investigations. There is a great deal of uncertainty involved with using these concentrations as action levels because of intra- and inter-species differences, extrapolation from laboratory to environmental conditions, and extrapolation from acute to chronic exposure scenarios.

The following sections describe how terrestrial and aquatic life PRL values were established for this ecological risk assessment.

4.1 Preliminary Risk Levels Protective of Terrestrial Life

For the purposes of the terrestrial organism PRL derivation, the ingestion exposure route to surface soil, surface water, and sediment was quantified. Literature toxicity information for the ECOPCs was evaluated to develop criteria for chemical concentrations in surface water that would result in no adverse effect and for incidental ingestion doses of surface soil and sediment that would result in no adverse effect to terrestrial organisms. If the observed maximum concentrations of chemicals exceed the derived surface water criteria, surface soil dose, and sediment dose, adverse effects to the exposed receptors could potentially occur.

A toxicity assessment approach was used for risk characterization. Toxicological information was used to evaluate the potential adverse effects of biota exposed to the IRP site contaminants and to determine contaminant levels in the abiotic environment that should have no adverse effect on biota. The toxicity-concentration screening step allows the BERA to focus on the compounds that are most likely to contribute to exposure levels that may cause an adverse effect for the various exposure scenarios.

4.1.1 Surface Soil and Sediment

Toxicity values (LOAELs) presented in Table D.4.1 were used as comparative values to assess soil and sediment doses derived from the maximum observed concentrations of the ECOPCs (based upon incidental soil ingestion rates for small mammals). Incidental soil ingestion rates used to calculate soil and sediment doses are presented in Table D.3.2. The toxicity values were divided by appropriate uncertainty factors

Table D.4.1
Toxicity Values (LOAELs) for the Ecological Chemicals of Potential Concern

Page 1 of 3

ECOPC	Laboratory Species	LOAEL (mg/kg bw/day)	Uncertainty Factor
Inorganic ECOPCs			
Aluminum	Rat	14	100
Antimony	Rat	0.35	10
Arsenic	Rat	6.4	10
Barium	Rat	0.25	100
Beryllium	Rat	0.85	10
Cadmium	Rat	0.014	10
Chromium (Total)	Rat	2,160 ³	1,000
Cobalt	Rat	0.5	100
Copper	Mouse	4.2	10
Cyanide	Rat	30	10
Iron	Rat	900	1,000
Lead	Rat	2.8	10
Manganese	Rat	930	10
Mercury	Rat	0.015	10
Nickel	Rat	50	10
Selenium	Rat	0.25	10
Silver	Mouse	18.1	100
Thallium	Rat	0.7	100
Vanadium	Rat	2.8	10
Zinc	Mouse	38	10
Organic ECOPCs			
1,1-Dichloroethane	Mouse	543	1,000
1,1-Dichloroethene	Rat	9	10
1,1,1-Trichloroethane	Rat	750	10
1,2-Dichloroethane	Rat	47	10
2-Butanone (MEK)	Rat	1,080	1,000
2-Methylnaphthalene	Rat	4,360	1,000
2-Methylphenol	Rat	150	100
2,3,7,8-Tetrachlorodibenzodioxin	Rat	0.001	1,000
4-Methyl-2-pentanone	Rat	250	100
4-Methylphenol	N/A	N/A	N/A
4,4-DDT	Rat	0.25	10
4,4-DDD	Rat	10	10
Acenaphthene	Mouse	350	100
Anthracene	Mouse	1,000	1,000
Aroclor 1260	Rat	841	1,000
Aroclor 1254	Rat	0.5	1,000
Aroclor 1248	Rat	0.8	1,000
Benzene	Rat	17.65	1,000

D.4.2

Table D.4.1
Toxicity Values (LOAELs) for the Ecological Chemicals of Potential Concern

Page 2 of 3

ECOPC	Laboratory Species	LOAEL (mg/kg bw/day)	Uncertainty Factor
Benzo(a)anthracene	Mouse	10 ^c	100
Benzo(a)pyrene	Mouse	10	100
Benzo(b)fluoranthene	Mouse	10 ^c	100
Benzo(g,h,i)perylene	Mouse	10 ^c	100
Benzo(k)fluoranthene	Mouse	10 ^c	100
Benzo(k)fluoranthene	Mouse	10 ^c	100
bis(2-ethylhexyl)phthalate	Rat	200	10
Butylbenzylphthalate	N/A	N/A	N/A
Carbazole	N/A	N/A	N/A
Carbon Disulfide	Rat	3,188	1,000
Chlorobenzene	Rat	2,290 ^a	1,000
Chloroform/Chloromethane	Rat	908	100 ^b
Chrysene	Mouse	10 ^c	100
DDE	N/A	N/A	N/A
Di-n-butylphthalate	Rat	8,000	1,000
Di-n-octylphthalate	N/A	N/A	N/A
Dibenzo(a,h)anthracene	Mouse	10 ^c	100
Dibenzofuran	Rat	6	1,000
Diethylphthalate	N/A	N/A	N/A
Endrin Ketone	Rat	3	1,000
Ethylbenzene	Rat	291	10
Fluoranthene	Mouse	250	100
Fluorene	Mouse	250	100
Indeno(1,2,3-cd)pyrene	Mouse	10 ^c	100
Methylene Chloride	Rat	52.58	10
N-Nitrosodiphenylamine	N/A	N/A	N/A
Naphthalene	Mouse	300	1,000
Phenanthrene	Mouse	40	1,000
Phenol	Rat	120	1,000
Polychlorinated Dibenzofurans	Rat	6	1,000
Polychlorinated Dibenzodioxins	Rat	0.001	1,000
Pyrene	Mouse	250	100
Styrene	N/A	N/A	N/A
Tetrachloroethylene	Mouse	71	1,000
Toluene	Rat	446	100

D.4.3

Table D.4.1
Toxicity Values (LOAELs) for the Ecological Chemicals of Potential Concern

Page 3 of 3

ECOPC	Laboratory Species	LOAEL (mg/kg bw/day)	Uncertainty Factor
Xylenes (Total)	Rat	357	10
<p>^aLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.</p> <p>^bUncertainty factor encompasses intra and interspecies extrapolations (dog ~ rat).</p> <p>^cToxicity values for various PAHs were unavailable in literature renewed. Values presented represent values derived from benzo(a)pyrene studies.</p> <p>Notes: ECOPC = Ecological chemical of potential concern. LOAEL = Lowest observable adverse effect level.</p>			

D.4.4

(Tables D.4.2 and D.4.3) to adjust for the level of uncertainty involved with these data comparisons. If the calculated incidental ingestion dose exceeded the adjusted toxicity value, it was determined to be a final ECOPC.

Small mammals ingest soil during feeding, grooming, and burrowing activities. A mean soil ingestion rate of 0.000873 grams soil/gram bw/day was estimated from data reported by Garten (1980) for hispid cotton rats, white-footed mouse, and eastern chipmunk, and from body weight data reported by Hoogland (1981) and Linder (1988). Soil consumption data are presented in Table D.3.2. A summary of the NOAEL and LOAEL values, as well as the reference sources are provided in Table D.4.5.

Chronic exposure derived LOAEL values were preferably chosen for the no adverse effect water criteria calculations. However, where chronic LOAEL values were absent, other toxicity values (subchronic/acute LOAELs or oral LD50s) were used with an uncertainty factor. A summary of the NOAEL, LOAEL, and LD50 values used for the exposure assessment calculations is presented in Table D.4.4. The applied uncertainty factors for these criteria are presented in Tables D.4.2 and D.4.3, and the applied water ingestion rates used for the calculation of the no adverse effect water criteria are presented in Table D.3.2. A summary of the NOAEL and LOAEL values, as well as the reference sources, are provided in Table D.4.5.

Toxicity values (NOAELs) presented in Table D.4.4 were used to calculate water criteria based upon the oral ingestion rate of drinking water for small mammals. The derived criteria is a concentration that corresponds to the NOAEL level. Therefore, an observed concentration that fell below this criteria should not result in adverse effects to the exposed receptor, while concentrations exceeding this criteria could potentially cause adverse effects.

"No effect" criteria were developed by evaluating and assessing the toxicity and effects of each contaminant on wildlife populations. Pertinent regulatory documents and the general literature were used as sources of information in developing and selecting appropriate criteria from toxicity values (i.e., EPA Ambient Water Quality Criteria [AWQC], Health Advisories, IRIS data base, and Health Effects Assessment Documents).

NOAEL and LOAEL values are literature-derived toxicity values based upon toxicological investigations using laboratory or wild animal species. Data related to chronic oral exposure studies (gavage, drinking water, or diet) were used in preference to data derived from intraperitoneal or intramuscular exposure studies. Inhalation toxicity data were not included in the toxicity assessments. For similar reasons, the potential for dermal exposure was not addressed in this document. Criteria were not estimated because of the uncertainty in correlating dermal toxicity under laboratory conditions

Table D.4.2
Uncertainty Factors Used in Establishing Adjusted LOAEL
Values for Soil and Sediment Dose Comparisons

Health Effects	Factor Used to Convert Effect to a Chronic LOAEL	Factor Applied for Interspecific Variation	Total Uncertainty Factor
Chronic LOAEL	--	10	10
Subchronic LOAEL	10	10	100
Acute LOAEL, LD50	100	10	1,000
Source: IRIS, 1992; EPA, 1992c.			

Table D.4.3
Uncertainty Factors Used in Establishing
No Adverse Effect Water Criteria Concentrations

Health Effects	Factor Used to Convert Effect to a Chronic NOAEL	Factor Applied for Interspecific Variation	Total Uncertainty Factor
Chronic NOAEL	--	10	10
Chronic LOAEL	5	10	50
Subchronic NOAEL	10	10	100
Subchronic LOAEL	50	10	500
Acute NOAEL	100	10	1,000
Acute LOAEL, LD50	1,000	10	10,000
Source: IRIS, 1992; EPA, 1992c.			

Table D.4.4
Toxicity Criteria Used for the Calculation of
Preliminary Risk Levels for the Protection of Terrestrial Life

Page 1 of 5

Ecological Chemical of Potential Concern (ECOPCs)	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	LD50^{a,b}
Inorganic ECOPCs			
Aluminum	19	14	--
Antimony	N/A	0.35	--
Arsenic	1.2	6.4	--
Barium	0.25	N/A	54 ^a
Beryllium	0.54	0.85	--
Cadmium	0.004	0.014	--
Chromium (Total)	0.46	N/A	2,160 ^a
Cobalt	0.05	0.5	--
Copper	12.5	4.2	--
Cyanide	10.8	30	--
Iron	N/A	900	30,000 ^a
Lead	0.45	2.8	--
Manganese	290	930	--
Mercury	0.003	0.015	--
Molybdenum	--	--	6,050
Nickel	5	50	--
Nitrate	N/A	N/A	--
Selenium	0.5	0.25	--
Silver	181.2	18.1	--
Thallium	0.2	0.7	--

Table D.4.4
Toxicity Criteria Used for the Calculation of
Preliminary Risk Levels for the Protection of Terrestrial Life

Page 2 of 5

Ecological Chemical of Potential Concern (ECOPCs)	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	LD50^{a,b}
Vanadium	0.7	2.8	--
Zinc	98.3	38	--
Organic ECOPCs			
1,1-Dichloroethane	543	N/A	--
1,1-Dichloroethene	10	9	--
1,1,1-Trichloroethane	500	750	--
1,2-Dichloroethane	25	47	--
2-Butanone	173	1,080	--
2-Methylnaphthalene	N/A	N/A	4,360 ^a
2-Methylphenol	50	150	--
2,3,7,8-Tetrachlorodibenzodioxin	0.0001	0.001	--
2,4-Dimethylphenol	50	250	--
4 Methylphenol	N/A	N/A	N/A
4-Methyl-2-pentanone	50	250	--
4,4-DDD	2.4	N/A	--
4,4-DDE	N/A	N/A	N/A
4,4-DDT	0.05	N/A	--
Acenaphthene	175	350	--
Acetone	100 ^c	500 ^c	--
Anthracene	1,000	N/A	3,300 ^d
Benzene	17.65	N/A	3,306 ^a
Benzo(a)anthracene	N/A	10	--

Table D.4.4
Toxicity Criteria Used for the Calculation of
Preliminary Risk Levels for the Protection of Terrestrial Life

Page 3 of 5

Ecological Chemical of Potential Concern (ECOPCs)	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	LD50 ^{a,b}
Benzo(a)pyrene	N/A	10	50 ^d
Benzo(b)fluoranthene	N/A	10	40 ^d
Benzo(k)fluoranthene	N/A	10	72 ^d
Benzo(g,h,i)perylene	N/A	10	--
bis(2-ethylhexyl)phthalate	65	200	--
Butylbenzylphthalate	N/A	N/A	--
Carbon Disulfide	11 (Rat) 11 (Rabbit)	N/A N/A	3,188 ^a --
Carbazole	N/A	N/A	500
Chlorobenzene	N/A	N/A	2,290
Chloroform	60 30	N/A 15	908 ^a --
Chrysene	N/A	10	99
Di-N-Butylphthalate	N/A	N/A	8,000 ^a
Dibenzo(a,h)anthracene	N/A	10	0.006 ^d
Dibenzofuran	1	6	--
Diethylphthalate	750	3,160	8,600
Endrin Ketone	N/A	N/A	3
Ethyl Benzene	97.1	291	--
Fluoranthene	125	250	--
Fluorene	125	250	--
Indeno(1,2,3-cd)pyrene	N/A	10	72 ^d

Table D.4.4
Toxicity Criteria Used for the Calculation of
Preliminary Risk Levels for the Protection of Terrestrial Life

Page 4 of 5

Ecological Chemical of Potential Concern (ECOPCs)	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	LD50^{a,b}
Methylene Chloride	5.85	52.58	--
Naphthalene	41	300	--
n-Nitrosodiphenylamine	N/A	N/A	N/A
Phenol	60	120	--
Phenanthrene	10	40	--
Polychlorinated Dibenzodioxins	0.0001	0.001	--
Polychlorinated Dibenzofurans	1 6	6 --	-- --
PCB (Aroclor 1248)	N/A	N/A	0.8 ^c
PCB (Aroclor 1254)	N/A	N/A	0.5 ^c
PCB (Aroclor 1260)	N/A	N/A	841 ^b
Pyrene	125	250	--
Styrene	14	71	5,000
Tetrachloroethylene	14	71	--
Trichloroethylene	18.4	393	--

Table D.4.4
Toxicity Criteria Used for the Calculation of
Preliminary Risk Levels for the Protection of Terrestrial Life

Page 5 of 5

Ecological Chemical of Potential Concern (ECOPCs)	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	LD50^{a,b}
Toluene	223	446	--
Xylenes (Total)	179	357	--

^aLD50 values utilized only when NOAEL and LOAEL values could not be located in the literature (units: mg/kg bw). NOAEL and LOAEL values were used in preference to LD50 values. (Sources: Sax, 1992; USFWS, 1984.)

^bSource = USFWS, 1984.

^cLiterature NOAELs and LOAELs do not apply for the purposes of this report, since they were derived from inhalation exposures.

^dSource = Eisler, 1987b.

^eSource = Eisler, 1986.

Notes: N/A = Not available in the literature reviewed.

NOAEL = No observable adverse effect level.

LOAEL = Lowest observable adverse effect level.

Source: Sax, 1992; USFWS, 1984.

Table D.4.5
Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern

Table D.4.5 Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern											Page 1 of 6
Ecological Chemical of Potential Concern (ECOPCs)	No Observable Adverse Effect Level (NOAEL)					Lowest Observable Adverse Effect Level (LOAEL)					
	NOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b	LOAEL mg/kg bw/day	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b	
Inorganic ECOPCs											
Aluminum	19	mouse	chronic	W	Ondreich et al., 1966	14	rat	subchronic	G	Paternain et al., 1988	
Ammonia	6.4 ^c	rat	subchronic	I	Broderick et al., 1976	17.4 ^c	rat	subchronic	I	Broderick et al., 1976	
Antimony	NA				IRIS, 1992	0.35	rat	chronic	W	Schroeder et al., 1970	
Arsenic	1.2	dog	chronic	F	Byron et al., 1967	6.4	rat	chronic	F	Byron et al., 1967	
Barium	0.25 0.825	rat mouse	chronic chronic	W W	Schroeder and Mitchner, 1975 ^{a,b}	NA					
Beryllium	0.54	rat	chronic	W	Schroeder and Mitchner, 1975 ^{a,b}	0.85	rat	chronic	W	Schroeder and Mitchner, 1975 ^a	
Boron	8.8	dog	chronic	F	Weir and Fisher, 1972	29	dog	chronic	F	Weir and Fisher, 1972	
Cadmium	0.004 3.5	rat rat	chronic chronic	W W	Kopp et al., 1982 Shaikh et al., 1989	0.014	rat	chronic	W	Kopp et al., 1982	
Chromium (Total)	0.46	rat	chronic	W	Schroeder et al., 1965	NA				IRIS, 1992	
Cobalt	0.05	rat	subchronic	G	Krasovskii and Fridlyand, 1971	0.5	rat	subchronic	G	Krasovskii and Fridlyand, 1971	
Copper	12.5	rat	subchronic	F	Murphy et al., 1981	4.2	mouse	chronic	W	Massie and Aiello, 1984	
Cyanide	10.8	rat	chronic	F	Philbrick et al., 1979	30	rat	chronic	F	Philbrick et al., 1979	
Iron	NA				IRIS, 1992	900	rat	acute	W	Venugopol and Luckey, 1978	
Lead	0.45	rat	chronic	W	Perry et al., 1988	2.5 2.8	dog rat	chronic chronic	W F	Carmignani et al., 1988 Azar et al., 1973	
Manganese	290	rat	chronic	F	Hejmanick et al., 1987 ^{a,b}	930	rat	chronic	F	Hejmanick et al., 1987 ^{a,b}	
Mercury	0.003	rat	chronic	F	Fitzhugh et al., 1950	0.015	rat	chronic	F	Fitzhugh et al., 1950	
Nickel	5	rat	chronic	F	Ambrose et al., 1976	50	rat	chronic	F	Ambrose et al., 1976	
Selenium	0.5	rat	chronic	F	Nelson, et al., 1943	0.25	rat	chronic	F	Nelson et al., 1943	
Silver	181.2	rat	acute	W	Walker, 1971	18.1	mouse	subchronic	W	Rungby and Danscher, 1984	

Page 2 of 6

Table D.4.5
Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern

Ecological Chemical of Potential Concern (ECOPCs)	No Observable Adverse Effect Level (NOAEL)					Lowest Observable Adverse Effect Level (LOAEL)				
	NOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b	LOAEL mg/kg bw/day	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b
Tin	20-30	rat	subchronic	F	DeGroot et al., 1973	100-150	rat	chronic	F	WHO, 1980
Thallium	0.2	rat	subchronic	G	Stoltz et al., 1986	0.7	rat	subchronic	W	Formigli et al., 1986
Vanadium	0.7	rat	chronic	W	Schroeder et al., 1970	2.8	rat	chronic	W	Kowalska, 1988
Zinc	98.3	rat	subchronic	F	Drinker et al., 1927	38	mouse	chronic	W	Aughey et al., 1977
Organic ECOPCs										
1,1-Dichloroethane	543	mouse	subchronic	G	Klaunig et al., 1986	NA				IRIS, 1992
1,1-Dichloroethylene	10	rat	chronic	W	Rampy et al., 1977	9	rat	chronic	W	Quast et al., 1983
1,1,1-Trichloroethane	500	rat	chronic	G	Maltoni et al., 1986	750	rat	chronic	G	Maltoni et al., 1986
1,2-Dichloroethane	25	rat	chronic	F	Alumot et al., 1976	47	rat	chronic	G	NCI, 1978
1,2-Dichloroethylene ^d	17 ^e	mouse	subchronic	W	Barnes et al., 1985	175 ^d	mouse	sub-chronic	W	Barnes et al., 1985
2-Butoleone (MFK)	173	rat	subchronic	G	Rabison et al., 1985	1,000	rat	acute	G	Brown and Hewitt, 1984
2-Butene	1,500 488	rat rat	acute subchronic	G G	Hewitt et al., 1980 Eben et al., 1979	600 100	guinea pig hen	subchronic subchronic	W G	Abdel-Rahman et al., 1978 Abou-Donia et al., 1983
2-Methylnaphthalene	NA				IRIS, 1992	NA				IRIS, 1992
2-Methylphenol	50	rat	subchronic	G	EPA, 1986 ^b , 1987	150	rat	subchronic	G	EPA, 1986 ^b , 1987
2,3,7,8-Tetrachlorodibenzodioxin	0.0001	rat	acute	G	Gupta et al., 1973	0.001	rat	acute	G	Gupta et al., 1973
2,4-Dichlorophenol	0.3	rat	subchronic	W	Exon and Koller, 1985	3.0	rat	subchronic-chronic	W	Exon and Koller, 1985
2,4-Dimethylphenol	50	mouse	subchronic	G	EPA, 1989 ^c	250	mouse	subchronic	G	EPA, 1989 ^c
4-Chloraniline	NA				IRIS, 1992	12.5	rat	chronic	F	NCI, 1979
4-Methylphenol	NA				IRIS, 1992	NA				IRIS, 1992
4-Methyl-2-pentene (MIBK)	50	rat	subchronic	G	Microbiological Associates, 1986	250	rat	subchronic	G	Microbiological Associates, 1986

Table D.4.5
Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern

Page 3 of 6

Ecological Chemical of Potential Concern (ECOPCs)	No Observable Adverse Effect Level (NOAEL)					Lowest Observable Adverse Effect Level (LOAEL)				
	NOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b	LOAEL mg/kg bw/day	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b
4,4-DDE	NA				IRIS, 1992	NA				IRIS, 1992
4,4-DDD	2.4	mouse	chronic (15 mo.)	Feeding	Wolf et al., 1979	10	rat	chronic	Feeding	Ottoboni, 1969
4,4-DDT	0.05 1.0	rats rats	chronic chronic	F F	Laug et al., 1950 Ottoboni, 1969	0.25 0.26	rats mouse	chronic chronic	F F	Laug et al., 1950 Tomatis et al., 1972
Acenaphthene	175	mouse	chronic	F	EPA, 1989 ^c	350	mouse	subchronic	oral	US EPA, 1989 ^c
Acetone	100 ^c	rat	subchronic	G	EPA, 1986 ^a	500 ^c	rat	subchronic	G	EPA, 1986 ^a
Aniline	19 ^c	rat	acute	I	duPont de Nemours, 1982	64.7 ^c	rat	acute	I	duPont de Nemours, 1982
Anthracene	1,000	mouse	subchronic	G	EPA, 1989 ^f	NA				IRIS, 1992
Benzene	2.35 17.65	rat rat	acute subchronic	G G	Diechman et al., 1963 EPA, 1980 ^a					
Benzo(a)anthracene						10 ^h	mouse	chronic	F	Mackenzie and Angevine, 1981
Benzo(a)pyrene						10	mouse	chronic	F	Mackenzie and Angevine, 1981
Benzo(b)fluoranthene						10 ^h	mouse	chronic	F	Mackenzie and Angevine, 1981
Benzo(k)fluoranthene						10 ^h	mouse	chronic	F	Mackenzie and Angevine, 1981
Benzo(g,h,i)perylene						10 ^h	mouse	chronic	F	Mackenzie and Angevine, 1981
Benzoic Acid	250 175	rabbit mouse	subchronic subchronic	G G	FDRL, 1972 FDRL, 1972	80	mouse	chronic	G	Ignatiev, 1965
bis(2-ethylhexyl)phthalate	65	rat	chronic	F	Carpenter et al., 1953	200	rat	chronic	F	Carpenter et al., 1953
Butylbenzylphthalate	NA				IRIS, 1992	NA				IRIS, 1992
Carbazole	NA				IRIS, 1992	NA				
Carbon Disulfide	11	rats and rabbits	subchronic	I	Hardin et al., 1981	NA				IRIS, 1992
Chlorobenzene	NA				IRIS, 1992	NA				IRIS, 1992

Table D.4.5 Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern										
Ecological Chemical of Potential Concern (ECOPCs)	No Observable Adverse Effect Level (NOAEL)					Lowest Observable Adverse Effect Level (LOAEL)				
	NOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b	LOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b
Chloroform	60 30	rat dog	chronic chronic	G C	Palmer et al., 1979 Heywood et al., 1979	15	dog	chronic	C	Heywood et al., 1979
Chrysene	NA				IRIS, 1992	10 ^b	mouse	chronic	F	Mackenzie and Augerine, 1981
Dibenz(a,h)anthracene	NA				IRIS, 1992	10 ^b	mouse	chronic	F	Mackenzie and Augerine, 1981
Dibenzofuran	NA				IRIS, 1992	NA				IRIS, 1992
Diethylphthalate	750	rat	subchronic	Oral	Brown et al., 1978	3,160	rat	subchronic	Oral	Brown et al., 1978
Di-n-butylphthalate	62	rat	intermed.	Feeding	Nikonow et al., 1973	125	rat	intermed.	Feeding	Killinger et al., 1988
Di-n-octylphthalate	NA				IRIS, 1992	NA				IRIS, 1992
Endrin ketone	NA				IRIS, 1992					IRIS, 1992
Ethylbenzene	97.1	rat	subchronic	G	Wolf et al., 1956	291	rat	subchronic	G	Wolf et al., 1956
Fluoranthene	125	mouse	subchronic	G	EPA, 1988 ^b	250	mouse	subchronic	G	EPA, 1988 ^b
Fluorene	125	mice	subchronic	G	EPA, 1989 ^d	250	mouse	subchronic	G	EPA, 1989 ^d
Heptachlorodibenzofuran ^f	1 6	rat mouse	acute acute	G G	Moore et al., 1976 Moore et al., 1976	6	rat	acute	G	Moore et al., 1976
Hexachlorodibenzofuran ^f	1 6	rat mouse	acute acute	G G	Moore et al., 1976 Moore et al., 1976	6	rat	acute	G	Moore et al., 1976
Heptachlorodibenzo-p-dioxin ^g	0.0001	rat	acute	G	Gupta et al., 1973	0.001	rat	acute	G	Gupta et al., 1973
Hexachlorodibenzodioxin ^g	0.0001	rat	acute	G	Gupta et al., 1973	0.001	rat	acute	G	Gupta et al., 1973
Indeno(1,2,3-cd)pyrene	NA				IRIS, 1992	10 ^b	mouse	subchronic	F	Mackenzie and Augerine, 1981
Methoxy (1)thiophene	5.85 6.47	male rats female rats	chronic chronic	G G	NCA, 1982 NCA, 1982	52.58 58.32	male rats female rats	chronic chronic	G G	NCA, 1982 NCA, 1982
Naphthalene	41	rat	chronic	F	Schmahl, 1955	300	mouse	acute	G	Plaster et al., 1985
Octachlorodibenzodioxin ^g	NA				IRIS, 1992	NA				IRIS, 1992

Table D.4.5 Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern											
Ecological Chemical of Potential Concern (ECOPCs)	No Observable Adverse Effect Level (NOAEL)						Lowest Observable Adverse Effect Level (LOAEL)				
	NOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b		LOAEL mg/kg bw/day	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b
Octachlorodibenzodioxins ^g	0.0001	rat	acute	G	Gupta et al., 1973		0.001	rat	acute	G	Gupta et al., 1973
Octachlorodibenzofurans ^f	1 6	rat mouse	acute acute	G G	Moore et al., 1976 Moore et al., 1976		6	rat	acute	G	Moore et al., 1976
PCB (Aroclor 1248)	NA				IRIS, 1992		NA				IRIS, 1992
PCB (Aroclor 1254)	NA				IRIS, 1992		NA				IRIS, 1992
PCB (Aroclor 1260)	NA				IRIS, 1992		NA				IRIS, 1992
Pentachlorodibenzodioxins ^g	0.0001	rat	acute	G	Gupta et al., 1973		0.001	rat	acute	G	Gupta et al., 1973
Phenanthrene	10 ^h	mouse	acute	G	MacKenzie and Angervine, 1981		40 ^h	mouse	acute	G	MacKenzie and Angervine, 1981
Phenol	60	rat	acute	G	NTP, 1983		120	rat	acute	G	NTP, 1983
Polychlorinated Dibenzodioxins	0.0001	rat	acute	G	Gupta et al., 1973		0.001	rat	acute	G	Gupta et al., 1973
Polychlorinated Dibenzofurans	1 6	rat mouse	acute acute	G G	Moore et al., 1976 Moore et al., 1976		6	rat	acute	G	Moore et al., 1976
Pyrene	125	mouse	subchronic	G	EPA, 1988 ^b		250	mouse	subchronic	G	EPA, 1988 ^b
Styrene	NA				IRIS, 1992		NA				IRIS, 1992
Tetrachloroethylene	14	mouse	acute	G	Buben and O'Flaherty, 1985		71	mouse	acute	G	Buben and O'Flaherty, 1985
Trichloroethylene	18.4 17.9	male mouse female mouse	subchronic subchronic	G G	NCI, 1976 NCI, 1976		393	mouse	subchronic	G	NCI, 1976
Toluene	223	rat	acute	G	NTP, 1989		446	rat	subchronic	G	NTP, 1989

Page 6 of 6

Table D.4.5
Literature Derived Oral Toxicity Values for the Ecological Chemicals of Potential Concern

Ecological Chemical of Potential Concern (ECOPCs)	No Observable Adverse Effect Level (NOAEL)				Lowest Observable Adverse Effect Level (LOAEL)					
	NOAEL (mg/kg bw/day)	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b	LOAEL mg/kg bw/day	Laboratory Animal Species	Exposure Duration	Exposure Route ^a	References ^b
Vinyl Chloride	0.17	rat	chronic	G	Tile et al., 1983	1.7	rat	chronic	G	Tile et al., 1983
Xylenes (Total)	179	rat	chronic	G	NTP, 1986	357	rat	chronic	G	NTP, 1986

^aG = gavage, F = diet/food, W = diet/drinking water, C = diet/capsule, I = Inhalation.

^bFor complete reference citation, please refer to References section.

^cOral exposure data through ingestion not available. NOAEL and LOAEL values derived for inhalation exposure—units are mg/cu. m.

^dDerived for Trans-1,2 dichloroethylene.

^eDerived from inhalation exposure data.

^fToxicological data for individual Dibenzofuran isomers were unavailable. Values represented within the table have been derived from investigations utilizing a composite of dibenzofurans for exposures.

^gToxicological data for individual polychlorinated dibenzodioxins were unavailable. Values represented within the table have been derived from investigations utilizing composites of 2,3,7,8-tetrachlorinated dibenzo-p-dioxin.

^hDerived from studies involving oral exposure to benzo(a)pyrene.

Note:

NA = Not available

(concentrated solutions, shaved skin of test animals) with toxicity under field conditions (generally dilute concentrations mixed with soil or water, contact with various body surfaces that can be covered with hair or are calloused).

Toxicity values chosen for this BERA are media- and species-specific. Laboratory toxicity values extracted from the literature were used as a basis of data comparison for ECOPC determination. Analytes with little or no toxicity literature values did not have criteria for ECOPC selection established.

The criteria presented in this section were used to estimate concentrations of ECOPCs in the abiotic environment and to then evaluate potential adverse ecological effects resulting from exposure to site contaminants. These values are used in this document as a basis for evaluating potential adverse effects to species and ecosystems where the effects may be subtle or difficult to detect by direct field observation alone. Further evaluation of these concentrations during the exposure characterization process leads to the development of criteria based on the same approach used to calculate the exposure concentrations resulting in no adverse effect.

4.1.2 Surface Soil

Soil dose levels were developed in each toxicity assessment to the extent that appropriate data were available. A soil ingestion dose is derived by multiplying the oral incidental soil ingestion rate (of the applicable species) to the maximum observed soil concentration of an ECOPC (in mg/mg soil). The resulting value is compared to an adjusted literature derived LOAEL with an applied uncertainty factor.

$$\text{Soil ECOPC Dose (mg/kg/bw/day)} = \text{Incidental Soil Ingestion Rate (mg soil/kg bw/day)} \times \text{maximum observed concentration (mg/mg soil)} \quad (1)$$

If the derived soil dose of an ECOPC exceeds the adjusted literature LOAEL value (LOAEL/applicable uncertainty factor), it is considered to potentially pose a risk to exposed organisms. If the derived incidental soil dose of an ECOPC falls below the adjusted LOAEL value, it is not considered to present a risk. A summary of the applied uncertainty factors, comparable species, and toxicity values used for the estimation of soil doses is presented in Table D.4.1.

4.1.3 Surface Water and Groundwater

Information pertaining to contaminated surface water ingestion (and incidental groundwater seep ingestion) by terrestrial organisms was evaluated to assess potential toxicity due to onsite conditions. By assuming that toxicity via oral ingestion would be similar regardless of the carrier, the most sensitive NOAEL or LOAEL was divided by both water intake and the appropriate uncertainty factor to estimate water concentrations associated with no adverse effect (water criteria).

Where both LOAEL and NOAEL values were available, the NOAEL was selected as the preferred value. Chronic data were used in preference to subchronic or acute values because there is more applicability and less uncertainty involved with this estimate. Appropriate NOAEL, LOAEL, and LD50 values used for the water criteria were presented in Table D.4.4.

The equation used to derive water criteria is as follows:

$$\frac{\text{Water Criteria}^*}{\mu\text{g/L}} = 1,000^c \times \left[\left(\frac{\text{Chronic NOAEL}^a \text{ (mg/kg bw/day)}}{\text{Water Ingestion Rate (L/kg bw/day)}} \right) \times \text{UncertaintyFactor}^b \right] \quad (2)$$

*Water Criteria is equal to a concentration of an ECOPC in surface water that would be associated with no observable adverse effect.

^{a,b}Chronic exposure derived NOAEL values were used preferably for water criteria calculations. However, where chronic LOAEL values were absent in the literature, other toxicity values (subchronic NOAEL, Acute LOAEL, Chronic-Subchronic-Acute LOAEL or LD50 values) with the appropriately applied uncertainty factor^b.

^cMultiplying factor of 1,000 will convert water criteria from mg/l to $\mu\text{g/l}$.

If the maximum observed concentration of the potential ECOPC exceeded the corresponding water criteria for that chemical, the chemical potentially represents a risk to exposed terrestrial populations. If the maximum observed concentration did not exceed the established water criteria, the chemical was not regarded as a potential risk to exposed populations. A summary of the applied uncertainty factors for the derivation of the water criteria is presented in Table D.4.6.

4.1.4 Sediment

Selection of sediment ECOPCs that contributed significant risk was conducted in the same technique as surface soil ECOPC selection. An incidental sediment ingestion dose was calculated for small mammals with the assumption that the organisms would

Table D.4.6
Preliminary Risk Levels for the Protection of Terrestrial Receptors Exposed to Surface Water and Groundwater

Page 1 of 2

Ecological Chemical of Potential Concern (ECOPCs)	Applicable Species	Uncertainty Factor ^a	Water Criteria ^b (µg/L)
Inorganic ECOPCs			
Aluminum	Mouse	10	9,500
Arsenic	Dog	10	2,400
Barium	Rat/Mouse	10	200-412
Beryllium	Rat	10	432
Cadmium	Rat	10	3.2
Chromium (Total)	Rat	10	368
Cobalt	Rat	100	4
Copper	Rat	100	1,000
Manganese	Rat	10	232,000
Molybdenum	Rat	1,000	48,400
Nickel	Rat	10	400
Nitrate	d	d	d
Silver	Rat	1,000	1,450
Thallium	Rat	0.2	16
Vanadium	Rat	10	560
Zinc	Rat	100	7,864
Organic ECOPCs			
1,1-Dichloroethene	Rat	10	8,000
1,1,1-Trichloroethane	Rat	10	400,000
2-Methylnaphthalene	Rat	10,000	3.48
2-Methylphenol	Rat	100	4,000
2,4-Dimethylphenol	Mouse	100	2,500
4-Methyl-2-pentanone	Rat	100	4,000
4-Methylphenol	NA	NA	NA
Acetone	NA ^c	100	NA ^c
Antimony	Rat	100	28
Aroclor 1254	Rat	1,000	4
Benzene	Rat	100	1,412

Table D.4.6
Preliminary Risk Levels for the Protection of Terrestrial Receptors Exposed to Surface Water and Groundwater
Page 2 of 2

Ecological Chemical of Potential Concern (ECOPCs)	Applicable Species	Uncertainty Factor ^a	Water Criteria ^b (µg/L)
bis(2-ethylhexyl)phthalate	Rat	10	52,000
Carbon Disulfide	Rat	100	880
Chloroform	Rat	10	48,000
Dibenzofuran	Rat	1,000	8
Diethylphthalate	Rat	1,000	68,800
Ethyl Benzene	Rat	100	7,768
Fluorene	d	d	d
Iron	Rat	1,000	7,200
Lead	Rat	10	360
Mercury	Rat	10	2.4
Methylene Chloride	Rat	10	4,680
Naphthalene	Rat	10	32,800
Phenol	Rat	1,000	480
Selenium	Rat	10	400
Toluene	Rat	1,000	1,784
Trichloroethylene	Mouse	100	920
Xylenes (Total)	Rat	10	143,200

^aUncertainty factors applied for derivation of criteria appear in Table D.4.2.

^bWater criteria were derived using the equation:

$$\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) + \text{The Uncertainty Factor} \right]$$

^cWater criteria were not derived for acetone, since all reviewed toxicological literature was derived from inhalation exposure.

^dCriteria were not developed due to lack of available toxicity data.

Final units are in µg/L.

ingest sediment at the same rate as they ingest soil. The equation for the calculation of a soil ingestion dose that would correlate to the literature derived LOAEL was also applied to the sediment ECOPC selection (refer to Equation 1.). A summary of the sediment exposure point concentrations, applied LOAEL value, and applied uncertainty factor, is presented in Table D.4.1.

4.2 Preliminary Risk Levels Protective of Aquatic Life

4.2.1 Surface Water and Groundwater

The Clean Water Act requires the Environmental Protection Agency (EPA) to publish and periodically update Ambient Water Quality Criteria (AWQC). These criteria are developed for the protection of both human health and aquatic life and are non-enforceable guidelines that set concentrations of pollutants considered adequate to protect surface waters. ECOPCs were not identified in surface water and groundwater at the Shemya IRP sites because of the incomplete pathway. These will need to be developed at a later date.

4.2.2 Sediment

Currently, the EPA has not approved of an established method for developing sediment quality criteria, in part due to issues regarding bioavailability and exposure point concentrations. However, several approaches appear in the scientific literature (Burton, 1992). To assess sediment quality and thus, potential effects of exposure to sediment ECOPCs at Shemya, it was necessary to compare ECOPC concentrations in sediment to established adverse effect levels. The following sections outline the procedures used to make these comparisons.

The affinity of a chemical to persist in sediment is proportional to the amount of organic carbon present in the sediment. This relationship is best described by the persistence coefficient, K_{oc} , which is an equilibrium constant that measures the partitioning of a chemical between organic carbon and water. Using the K_{oc} , an absorption coefficient, K_d , can then be determined for a particular chemical by multiplying the K_{oc} by the actual amount of organic carbon present in the sediment or soil.

The first step in assessing sediment quality was to calculate K_d for the individual ECOPCs. These values were obtained by multiplying both the minimum and maximum values detected for organic carbon content at Shemya by the corresponding K_{oc} value. The sediment quality criteria were then developed by multiplying the K_d value by the appropriate Ambient Water Quality Criteria for a given ECOPC. The derived criteria are presented in Tables D.4.7 and D.4.8. Several K_d values have been previously

Table D.4.7
Sediment Criteria for Inorganic ECOPCs

Sediment COCs	Ambient Water Quality Criteria ^a				K _d ^b (unitless)	Sediment Criteria			
	Freshwater (µg/l)		Marine (µg/l)			Freshwater (µg/kg)		Marine (µg/kg)	
	Acute	Chronic	Acute	Chronic		Acute	Chronic	Acute	Chronic
Antimony	/p/88	/p/30	/p/1500	/p/500	45	3960	1350	67,500	22,500
Arsenic(III)	360	190	69	36	200	72,000	38,000	13,800	7200
Arsenic(V)	850*	--	2319*	--	200	170,000	--	463,800	--
Chromium(III)	1700	210+	10,300*	--	850	1,445,000	178,000	8,755,000	--
Chromium(VI)	16	11	1100	50	850	13,600	9350	935,000	42,500
Copper	18+	12+	2.9	--	35	630	420	102	--
Cadmium	3.9+	1.1+	43	9.3	6.5	25.4	7.2	280	60.5
Lead	83+	3.2+	220	8.5	900	74,700	2880	198,000	7650
Zinc	120+	110+	95	86	40	4800	4400	3800	3440

/p/ - proposed criterion

* - Insufficient data to develop criterion. Value presented is the LOEL - Lowest Observed Effect Level Inorganic Chemicals
+ - Hardness dependent criteria (100 mg/l CaCO₃ used)

^aUSEPA, Ambient Water Quality Criteria, 1993.

^bK_d values: CH2M HILL, 1990. K_d Document for Organic and Inorganic Chemicals

10010997 BC/mmm

Table D.4.8
Sediment Criteria for Organic ECOPCs

Sediment COCs	Ambient Water Quality Criteria ^a				Koc ^b (unitless)	Kd (unitless)		Sediment Criteria							
	Freshwater (µg/l)		Marine (µg/l)			Min. TOC (0.00105)	Max. TOC (0.031)	Freshwater (µg/kg)			Marine (µg/kg)				
	Acute	Chronic	Acute	Chronic				Acute (min.-max. TOC)	Chronic (min.-max. TOC)	Acute (min.-max. TOC)	Chronic (min.-max. TOC)				
bis(2-ethylhexyl)- phthalate	p/400	p/360	p/400	p/360	87,400	92	2709	36,800	1,116,000	33,120	975,240	36,800	1,116,000	33,120	975,240
2-Butanone	--	--	--	--	4.5	0.005	0.140	--	--	--	--	--	--	--	--
4,4-DDD	0.6*	--	3.6*	--	770,000	809	23,870	485	14,322	--	--	2912	85,932	--	--
4,4-DDT	1.1	0.001	0.13	0.001	243,000	4620	136,400	5082	150,040	4.62	136.4	601	17,732	4.62	136.4
4,4-DDE	1050*	--	14	--	4,400,000	255	7533	267,750	7,909,650	--	--	3570	105,462	--	--
Endrin Ketone	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Fluoranthene	3980*	--	40*	16*	38,000	40	1178	159,200	4,688,440	--	--	1600	47,120	640	18,848
PCB-1254	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Pyrene ^c	--	--	300*	--	38,000	40	1178	--	--	--	--	12,000	353,400	--	--

TOC - Total Organic Carbon

/p/ - proposed criterion

* - Insufficient data to develop criterion. Value presented is the LOEL - Lowest Observed Effect Level

Sources:

^aUSEPA, Ambient Water Quality Criteria.

^bUSEPA, 1987. Superfund Public Health Evaluation Manual. EPA 540/1-86/060.

^cBased on total PAHs.

established by the EPA for inorganic analytes. Therefore, the K_d and resulting sediment criteria do not have minimum and maximum values based upon organic matter content from Shemya.

In addition to these derived sediment criteria, sediment criteria established by the state of Washington were used as a second source of PRL values for the protection of aquatic life exposed to sediment. However, these criteria were derived for Washington specifically and, therefore, will introduce an amount of uncertainty to this assessment. The summarized Washington state sediment criteria are presented in Table D.4.9.

**Table D.4.9
Sediment Standards
for the State of Washington**

Page 1 of 3

Chemical Parameter	Sediment Quality Standards	Sediment Cleanup Standards
Metals (mg/kg dry weight)		
Arsenic	57	93
Cadmium	5.1	6.7
Chromium	260	270
Copper	390	390
Lead	450	530
Mercury	0.41	0.59
Silver	6.1	6.1
Zinc	410	960
Nonpolar Organic Compounds (mg/kg organic carbon)^a		
Aromatic Hydrocarbons		
LPAH ^b	370	780
Naphthalene	99	170
Acenaphthylene	66	66
Acenaphthene	16	57
Fluorene	23	79
Phenanthrene	100	480
Anthracene	220	1,200
2-Methylnaphthalene	38	64
HPAH ^c	960	5,300
Fluoranthene	160	1,200
Pyrene	1,000	1,400
Benz(a)anthracene	110	270
Chrysene	110	460

**Table D.4.9
Sediment Standards
for the State of Washington**

Page 2 of 3

Chemical Parameter	Sediment Quality Standards	Sediment Cleanup Standards
Total benzo(a)fluoranthenes ^d	230	450
Benzo(a)pyrene	99	210
Indeno(1,2,3-c,d)pyrene	34	88
Dibenz(a,h)anthracene	12	33
Benzo(g,h,i)perylene	31	7
Chlorinated Benzenes		
1,2-Dichlorobenzene	2.3	2.3
1,4-Dichlorobenzene	3.1	9
1,2,4-Trichlorobenzene	0.81	1.8
Hexachlorobenzene	0.38	2.3
Phthalate Esters		
Dimethyl phthalate	53	53
Dimethyl phthalate	61	110
Di-n-butyl phthalate	220	1,700
Butyl (benzyl) phthalate	4.9	64
bis(2-ethylhexyl) phthalate	47	78
Di-n-octyl phthalate	58	4,500
Miscellaneous		
Dibenzofuran	15	58
Hexachlorobutadiene	3.9	6.2
N-nitrosodiphenylamine	11	11
Total PCBs	12	65

**Table D.4.9
Sediment Standards
for the State of Washington**

Page 3 of 3

Chemical Parameter	Sediment Quality Standards	Sediment Cleanup Standards
Ionizable Organic Compounds ($\mu\text{g/kg}$ dry weight; parts per billion)		
Phenol	420	1,200
2-Methylphenol	63	63
4-Methylphenol	670	670
2,4-Dimethylphenol	29	29
Pentachlorophenol	360	690
Benzyl alcohol	57	73
Benzoic acid	650	650

^aThe listed chemical parameter criteria represent concentrations in parts per million, "normalized," or expressed, on a total organic carbon basis. To normalize to total organic carbon, the dry-weight concentration for each parameter is divided by the decimal fraction representing the percent total organic carbon content of the sediment.

^bThe LPAH criterion represents the sum of the following "low-molecular-weight polynuclear aromatic hydrocarbon" compounds: naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, and anthracene. The LPAH criterion is not the sum of the criteria values for the individual LPAH compounds as listed.

^cThe HPAH criterion represents the sum of the following "high-molecular-weight polynuclear aromatic hydrocarbon" compounds: fluoranthene, pyrene, benz(a)anthracene, chrysene, total benzo(a)fluoranthenes, benzo(a)pyrene, indeno(1,2,3-c,d)pyrene, dibenz(a,h)anthracene, and benzo(g,h,i)perylene. The HPAH criterion is not the sum of the criteria values for the individual HPAH compounds as listed.

^dThe total benzo(a)fluoranthenes criterion represents the sum of the concentrations of the "B," "J," and "K" isomers.

Source: Ginn and Pastorok, 1992.

5.0 Ecological Toxicity Assessment

A variety of inorganic and organic ECOPCs were identified in surface soil, surface water, groundwater, and sediment. A brief toxicity summary of some of the major groups of chemicals detected and identified as ECOPCs is presented in the following sections.

In order to utilize literature values when evaluating toxicity assessment data, the following assumptions were made regarding data quality and applicability:

- Chronic values were preferred to acute values
- Similar species data were preferred to nonrelated species data
- Only data derived from oral route exposure investigations were utilized to be consistent with the exposure assessment
- Only nonhuman data were considered

5.1 Toxicity Assessment of Inorganic ECOPCs

The toxicity of a number of inorganic elements found at Shemya has been reviewed in relation to aquatic and terrestrial organisms by various authors (see Venugopol and Luckey, 1978; and Eisler, 1985; 1987a and b; 1988a; 1988b).

A number of inorganic elements are essential in small amounts for animal nutrition because they are an integral part of at least one enzyme. These include arsenic, calcium, chlorine, chromium, copper, iron, magnesium, manganese, potassium, sodium, and zinc. The animal body has developed a variety of homeostatic mechanisms with regard to these essential macro and trace elements, so they are less likely to produce toxic effects than are the nonessential elements such as barium, cadmium, and lead. Nevertheless, those control mechanisms can be overwhelmed or circumvented and some essential elements can produce toxic effects.

Exposure of birds and mammals to various elements can be measured by analyzing their food, water, and air, or by analyzing selected tissues from their bodies. The toxicity of many elements is greatly influenced by the chemical state in which they occur in the water or air (or in soil or groundwater, which are not ingested in large volumes), which generally are not good indicators of potentially toxic exposure; concentrations in food are usually the most important. Because potentially toxic trace elements occur in many different forms, even the total concentration of some elements in the diet may not be a good predictor of toxicity.

Toxicity of inorganic elements in the diet varies so widely that it is not practical to list specific concentrations that cause adverse effects under conditions of acute or chronic exposure. Furthermore, the reported concentrations in soils or groundwater cannot reasonably be extrapolated to the concentrations that might result in foods of birds and mammals living near the site. The following general summary of the relative toxicity of inorganic elements is based primarily on reviews by Venugopal and Luckey (1978), Sax (1984), and Eisler (1985; 1987a and b; 1988a; 1988b).

The various elements are divided into bond valence groups as depicted on the periodic table of elements. General characteristics of those elements that belong to a particular group are described below.

Group I. Potassium and sodium are essential elements that are considered to be harmless and generally nontoxic. Copper is an essential trace metal that stimulates growth when moderately high levels are fed to mammals, but it is highly toxic to aquatic organisms.

Group II. Magnesium and calcium are also essential elements for animals, and barium is stimulatory but not essential. All three are considered relatively nontoxic at physiologic levels, but definitely toxic at higher levels. Cadmium is nonessential, but it is stimulatory at very low doses and highly toxic at relatively low levels. Homeostatic mechanisms maintain normal levels of magnesium and calcium, and to a certain extent barium, but cadmium is not controlled by homeostasis. Barium and cadmium are retained in the tissues and the body levels of these metals increase with age. Freshwater biota are the most sensitive animals to cadmium; concentrations of 0.7 to 5 $\mu\text{g/l}$ were associated with sublethal effects. Zinc is an essential trace metal that is relatively nontoxic because of efficient homeostatic mechanisms that maintain a proper balance within the body.

Group III. Aluminum is not known to be an essential element in animal nutrition. Intestinal absorption is generally very poor, and its toxicity is low in comparison to many other metals.

Group IV. Lead is a nonessential metal that is readily absorbed from the digestive tract. Organic compounds are more toxic than inorganic salts because of their greater lipid solubility, stability in biological fluids, and penetration into tissues such as the brain and lodging in the central nervous system. Adverse effects on aquatic biota have been reported at waterborne lead concentrations of 1 to 5 $\mu\text{g/l}$.

Group V. In general, inorganic arsenic compounds are more toxic than organic compounds, and trivalent compounds (in the form of soluble arsenites) are far more toxic than pentavalent compounds (arsenates). Arsenic is a teratogen and carcinogen that produces death and malformations in many species of mammals.

Group VI. Chromium is an essential trace metal for mammals, but it is toxic at high doses. Hexavalent chromium is the most biologically active form, although little is known about the properties of organochromium compounds, water-soluble species, or their interactions in complex mixtures. However, in mammals hexavalent chromium is chemically reduced in the acid fluid of the stomach. Chromium concentrations are usually highest at the lowest trophic levels, and no biomagnification has been observed in food chains.

Group VII. Manganese salts appear to be the least toxic of the essential metals. An efficient homeostatic mechanism prevents manganese accumulation in tissues, and toxic effects of manganese are not clearly manifested.

Group VIII. Iron is an essential metal that is generally not considered to be toxic to animals.

5.2 Toxicity Assessment of Organic Chemicals

5.2.1 Polychlorinated Biphenyls

Polychlorinated biphenyls (PCBs) are organic compounds commercially produced by chlorination of a biphenyl (BP) with anhydrous chlorine in the presence of iron filings or ferric chloride as the catalyst. The purified product is a complex mixture of chlorobiphenyls containing 18 to 79 percent chlorine; the precise composition depends on the conditions under which chlorination occurred (EPA, 1980c). Ten possible degrees of chlorination of the biphenyl molecule give rise to ten PCB congener groups: mono-, di-, tri-, tetra-, penta-, hexa-, octa-, and decabiphenyl (Eisler, 1986).

Toxic materials that exist as impurities in PCBs include polychlorinated dibenzofurans (PCDFs) in some domestic and foreign mixtures. Sometimes, PCBs used in electrical capacitors and transformers are converted under the action of heat or electrical arcing to form PCDFs, including 2,3,7,8-tetrachlorobenzofuran and 2,3,4,7,8-pentachlorobenzofuran (Eisler, 1986).

PCBs are extremely stable compounds and slow to chemically degrade under environmental conditions. Microbial degradation of PCBs depends on the degree of chlorination and the position of the chlorine atom on the biphenyl molecule; lower-chlorinated BPs are readily transformed by bacteria, but not the higher-chlorinated compounds (NAS, 1979). Higher chlorobiphenyls (those with five or more chlorines) are more persistent in the environment than those with three or less (EPA, 1980).

In mammals, PCBs are readily absorbed through the gut, respiratory system, and skin. Initially, PCBs concentrate in liver, blood, and muscle; eventually, accumulations are highest in adipose tissue and skin. Phenolic derivatives of dihydrodioxins are the major metabolites, but susceptibility of the individual PCB isomers to metabolism is a function of the number of chlorine atoms present on the biphenyl rings and their arrangement (Eisler, 1986).

The number and position of the chlorine atoms on the biphenyl rings affects the biological properties of the compound. For example, PCBs with hydrogen atoms on two adjacent carbon atoms in one or both rings are more readily metabolized than those with hydrogen atoms adjacent only to chlorines (EPA, 1980). Furthermore, PCBs with chlorines in the 2 and 6 (ortho) positions are easily metabolized by humans while those with chlorines in the 4 and 4' (para) positions tend to be biologically active and well-retained in tissues (EPA, 1980).

PCBs are usually taken up by animals and stored in lipids under circumstances of increasing lipid content in organs. However recent studies with marine fishes indicate that PCB components remain mobilizable from organs whose lipid contents increased (Boon et al., 1984). The degree of mobilization appeared to be related to polar lipid components such as phospholipids and glycerols. In other aquatic species, the role of diet and tissue-specific sites are important.

Biological availability and uptake of individual PCBs from aqueous solution are influenced primarily by two factors: the partition coefficient (Kow) based on the solubilities of compounds in N-octanol/water; and steric factors resulting from different patterns of chlorine substitution. Log Kow values for various isomers of Aroclor 1242, 1254, and 1260 are high, varying from 4.0 to 9.35, indicating high biological uptake potential. Steric effect coefficients are based on the number of chlorine atoms in the biphenyl molecule and their arrangement (Shaw and Connell, 1982). The product of log Kow and the steric effect coefficient seem to be directly related to bioaccumulation (Shaw and Connell, 1982). Thus, maximum uptake was found with penta and hexachlorobiphenyls predominant in Aroclor 1254, which have high values for log Kow and for steric effect coefficients. Comparatively less uptake was found with di-, tri-, and tetrachlorobiphenyls typical of Aroclor 1242, which have lower values for log Kow, and with hepta and octachlorobiphenyls predominant in Aroclor 1260, which have lower steric effect coefficients (Eisler, 1986).

5.2.2 Polycyclic Aromatic Hydrocarbons (PAHs)

Polycyclic aromatic hydrocarbons (PAHs) consist of hydrogen and carbon arranged in the form of two or more fused benzene rings. There are thousands of PAH compounds, each differing in the number and position of the aromatic rings and in the position of substituents on the basic ring system (Eisler, 1987). Environmental concern

has focused on PAHs that range in molecular weight from 128.16 (naphthalene 2-ring structure) to 300.36 (coronene, 7-ring structure). Unsubstituted lower molecular weight PAH compounds containing 2 or 3 rings exhibit significant, acute toxicity and other adverse effects to some organisms, but are noncarcinogenic; the higher molecular weight PAHs containing 4 to 7 rings compounds are demonstrably carcinogenic, mutagenic, or teratogenic to a wide variety of organisms including fish, other aquatic life, amphibians, birds, and mammals (Eisler, 1987). In general, PAHs show little tendency to biomagnify in food chains despite their high lipid solubility probably because most PAHs are rapidly metabolized. Inter and intraspecies responses to individual PAHs are quite variable and are significantly modified by many inorganic and organic compounds including other PAHs (Eisler, 1987).

PAHs are ubiquitous in nature and result from natural processes such as forest fires, microbial synthesis, and volcanic activities. Concern about PAHs in the environment reflects their persistence and the fact that some are known to be potent mammalian carcinogens, although environmental effects of most noncarcinogenic PAHs are poorly understood (Neff, 1985). Prior to 1900, a natural balance existed between the production and degradation of PAHs. Synthesis of PAHs by microorganisms and volcanic activity and production by man-made high temperature paralytic reactions and open burning seemed to be balanced by PAH destruction via photodegradation and microbial transformation. With increased industrial development and increased emphasis of fossil fuels as energy sources, the balance has been disturbed to the extent that PAH production and introduction into the environment greatly exceeds known PAH removal processes (Suess, 1976; Sims and Overcash, 1983). When released into the atmosphere, PAH compounds become associated with particulate materials. Their residence time in the atmosphere and transport to different geographic locations are governed by particle size, meteorological conditions, and atmospheric physics. Highly reactive PAHs decompose readily in the atmosphere by reaction with ozone and various oxidants; degradation times range from several days to six weeks for PAHs adsorbed onto particulates $<1\ \mu\text{m}$ in diameter to less than 1 day to several days for those adsorbed to larger particles (Suess, 1976).

In water, PAHs may either evaporate, disperse into the water column, become incorporated into bottom sediments, concentrate in aquatic biota, or experience chemical oxidation and biodegradation (Suess, 1976). The most important degradative processes for PAHs in aquatic systems are photooxidation, chemical oxidation, and biological transformation by bacteria and animals (Neff, 1979). Most PAHs in aquatic environments are associated with particulate materials; only about 33 percent are in dissolved form (Lee and Grant, 1981). PAHs in dissolved form will probably degrade rapidly through photooxidation (EPA, 1980) and degrade most rapidly at higher concentrations at elevated temperatures, at elevated oxygen levels, and at a higher incidence of solar radiation (Suess, 1976; Eisler, 1987).

PAHs in aquatic sediment degrade slowly. The ultimate fate of those PAHs that accumulate in the sediments is believed to be biotransformation and biodegradation by benthic organisms (EPA, 1980). PAHs will degrade rather slowly in the absence of penetrating radiation and oxygen and may persist indefinitely in oxygen-poor basins or in anoxic sediments. PAH degradation in aquatic environments occurs at a slower rate than that in the atmosphere (Suess, 1976).

6.0 Ecological Risk Characterization

Risk to ecological receptors has been established by comparing observed maximum concentrations of site-specific ECOPCs to derived water criteria, soil and sediment dose levels (for terrestrial organisms), and to federal ambient water quality criteria and derived sediment criteria (for aquatic organisms). The following sections describe the methods for risk characterization and the risk associated with the individual IRP sites.

6.1 Methods for Terrestrial Receptor Risk Characterization

The assessment of terrestrial receptor ecological risk is based upon:

- Comparison of the maximum observed concentration of each of the chemicals in surface water to derived water criteria based upon a no observable adverse effect level (NOAEL)
- Comparison of the maximum observed concentration of the chemicals in surface soil to an oral incidental soil ingestion dose derived from a lowest observable adverse effect level (LOAEL) from the literature
- Comparison of the maximum observed concentration of the chemicals in sediment to an oral incidental sediment ingestion dose derived from a lowest adverse effect level from the literature

The NOAEL and LOAEL values deemed appropriate for this assessment are presented in Table D.4.1.

The purpose of the ecological exposure characterization is to compare the results of the exposure assessment with the results of the toxicity assessment. The results are used to characterize the probability of adverse effects resulting from exposure to site-related contaminants.

Ecological exposure characterization was performed by estimating incidental ingestion doses of surface soil and sediment, ECOPCs, and water criteria (resulting in no adverse effect) from the exposure concentrations and literature-derived toxicity values provided in the toxicity assessments. The utilized toxicity values are the NOAEL or LOAEL values reported for a particular study. In the event that the lowest or no effect levels were not given, toxicity data were reviewed and the lowest value reported as toxic (i.e., LD50) was used as the comparative toxicity value with an applied uncertainty factor.

The values chosen for each ECOPC are reported in Tables D.4.4 and D.4.5. Often, data were unavailable for certain species. For instance, laboratory mammal studies were common, but few data were available specifically for wild mammals or cattle. The toxicological data base is even more sparse for birds. For chemicals lacking toxicity information for a species, quantification of risk was not made specifically for that group because of the uncertainty involved in extrapolating between taxonomic Kingdom, Phyla, or Class. For example, if toxicity data for plants were unavailable, risk for plants or crops was not quantified from data for rats. Although extrapolation within a Class (i.e., rats to cows) is also uncertain, values for mammals, fish, or birds were considered to be representative within a class because of the lack of available toxicological data for most of the ECOPCs. For related chemicals like the dibenzodioxins and dibenzofurans, toxicity data were insufficient to represent the different chemical forms; thus, one toxicity value was selected for the chemical group.

Exposure was not quantified in the manner of human health. Instead, exposure point concentrations were used to calculate intake doses in mg/kg bw/day and compared to the literature derived toxicity values. When the observed maximum concentration of a chemical exceeded these values it was an indication that potential exposure resulting in adverse effect to ecological health may exist, and further evaluation of ecological exposure is appropriate for these media of concern.

Each of the exposure pathways was evaluated for rats and mice, since literature information encompasses laboratory species only. Each media of concern was addressed qualitatively and quantitatively. Although the exposure estimates are uncertain, they are believed to be conservative because:

- The maximum observed concentration was used to represent the exposure point concentration.
- The contaminated source is assumed to be the total source of intake.
- The contaminant concentrations were assumed to represent acute and chronic exposure point concentrations.
- The measured concentration of the ECOPC is equivalent to the amount of chemical available for uptake (it was assumed that the chemical and physical properties of the ECOPC would not influence chemical availability and uptake).

No EPA-derived toxicity reference values exist for terrestrial organisms exposed to contaminated soil. As an alternative, the calculation of exposure point dose concentrations are compared with toxicity values from literature sources. Where literature sources are lacking, the potential hazard cannot be readily determined.

Exposure to sediment-related chemicals was quantitatively addressed in a similar manner to soil exposure. Little is known about the risk posed to receptor populations due to sediment-related contamination. It was assumed that terrestrial organisms will incidentally ingest the same amount of sediment as they do soil. This is probably an overly conservative assumption, since contact with sediment occurs less frequently than contact with soil. Based upon these results it was determined that incidental oral ingestion of sediment by small mammals may result in adverse effects.

Results of the soil dose comparisons to literature LOAEL values revealed few final ECOPCs in the surface soil. This is in part because of the small percentage of soil ingestion in the diet. These calculations do not take into consideration the possibility of multiple exposure routes or exposure sources. Organisms occurring onsite can become exposed to soil-related contamination by inhalation of dusts, dermal contact with the soils, and incidental ingestion. Combined exposures could present additional risk to the receptor organisms.

Concentrations of trace inorganic elements in soils at the site are not likely to be directly toxic to terrestrial organisms. However, some metals and other elements may influence biological and biochemical processes in the soil, limiting the decomposition rate of vegetative matter and thereby limiting metal concentrations in soil to a few times the background level (Beijer and Jernelov, 1979). Certain metals are taken up by plant roots and subsequently distributed to stems, leaves, and seeds whereby herbivores can be exposed through bioaccumulation.

Uptake of certain metals is affected by such factors as soil organic matter content, pH, and the presence of other metals that can act antagonistically or synergistically. With the presence of a diversity of metals within the Shemya Site surface soil it is difficult to assess their toxicity.

Based upon the results of this BERA, the potential toxicity of site contaminants requires further investigation to assess the potential adverse effects posed to receptor organisms.

6.2 Methods for Aquatic Receptor Risk Characterization

Aquatic organisms exposure to soil-related contamination was not addressed in this assessment. It is unknown whether significant amounts of soil-related contaminants would be transported to areas in which aquatic organisms could be exposed; therefore, this source of exposure was not assessed.

Comparison of criteria to the observed concentrations addressed only those exposures associated with single chemical exposure. Since a variety of chemicals are present that could act synergistically or antagonistically towards each other, the true toxicity posed by exposure to surface water is unknown.

6.3 Site-Specific Risk Characterizations

Risks posed to aquatic and terrestrial receptors exposed to various media within the areas of concern are described below. Suggestions for further investigation are also included.

6.3.1 FT-1

6.3.1.1 Terrestrial Risk Characterization. Results comparing derived soil doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.1. Observed maximum concentrations of ECOPCs in the water sources are compared to appropriate water criteria in Table D.6.2. Results of these comparisons indicate that terrestrial exposure to site soils will not provide a significant risk. In addition, terrestrial exposure to water resources within this area would not contribute significant risk.

6.3.1.2 Aquatic Risk Characterization. Because of the lack of appropriate media data (surface water and/or sediment), aquatic organism exposure and risk was not assessed.

6.3.2 FT-2

6.3.2.1 Terrestrial Risk Characterization. Results comparing derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.3. Observed maximum concentrations of ECOPCs in the water sources are compared to appropriate water criteria in Table D.6.4. Results of the comparisons indicate potential risk to terrestrial organisms exposed to site soil can result from the presence of benzene and lead. Water resources within the area are not a potential risk to exposed terrestrial organisms.

6.3.2.2 Aquatic Risk Characterization. Because of the lack of appropriate media data, aquatic organism exposure and risk was not assessed for this area of concern.

Table D.6.1
Comparison of FT-1 Soil ECOPC Terrestrial Receptor Incidental Ingestion
Dose Concentrations to Applicable PRL Values

Page 1 of 2

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Risk Level (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Inorganic ECOPCs						
Cadmium	11×10^{-6}	3.08×10^{-3}	Rat	0.014	10	No
Chromium (Total)	87×10^{-6}	2.4×10^{-2}	Rat	2,160 ^b	1,000	No
Copper	0.24×10^{-3}	1.78×10^{-1}	Mouse	4.2	10	No
Lead	0.67×10^{-3}	1.88×10^{-1}	Rat	2.8	10	No
Zinc	1.32×10^{-3}	9.78×10^{-1}	Mouse	38	10	No
Organic ECOPCs						
2-Butanone	1.4×10^{-6}	3.9×10^{-4}	Rat	1,080	1,000	No
2-Methylnaphthalene	1.3×10^{-7}	3.64×10^{-5}	Rat	4,360	1,000	No
Anthracene	98×10^{-9}	7.25×10^{-5}	Mouse	1,000	100	No
Benzene	$1,600 \times 10^{-9}$	4.48×10^{-4}	Rat	2.35	100	No
Benzo(a)anthracene	220×10^{-9}	1.63×10^{-4}	Mouse	10	100	No
Benzo(a)pyrene	200×10^{-9}	1.48×10^{-4}	Mouse	10	100	No
Benzo(b)fluoranthene	280×10^{-9}	2.1×10^{-4}	Mouse	10	100	No
Benzo(k)fluoranthene	84×10^{-9}	6.23×10^{-5}	Mouse	10	100	No
bis(2-ethylhexyl)phthalate	790×10^{-9}	2.2×10^{-4}	Rat	200	10	No
Butylbenzylphthalate	130×10^{-9}	NA	NA	NA	NA	?
Chlorobenzene	4×10^{-9}	1.12×10^{-6}	Rat	4	1,000	No
Chrysene	440×10^{-9}	3.26×10^{-4}	Mouse	10	100	No
Dibenzofuran	92×10^{-9}	2.58×10^{-5}	Rat	92	1,000	No
Di-n-butylphthalate	710×10^{-9}	1.98×10^{-4}	Rat	8,000	1,000	No
Di-n-octylphthalate	68×10^{-9}		NA	NA	NA	?
Ethyl benzene	$8,300 \times 10^{-9}$	2.34×10^{-3}	Rat	291	100	No
Fluoranthene	490×10^{-9}	3.6×10^{-4}	Mouse	250	100	No
Fluorene	78×10^{-9}	5.7×10^{-5}	Mouse	250	100	No
Indeno(1,2,3-cd)pyrene	69×10^{-9}	1.9×10^{-5}	Rat	72	1,000	No

D.6.5

26 February 1993

Table D.6.1
Comparison of FT-I Soil ECOPC Terrestrial Receptor Incidental Ingestion
Dose Concentrations to Applicable PRL Values

Page 2 of 2

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Risk Level (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Naphthalene	500×10^{-9}	3.7×10^{-4}	Mouse	300	1,000	No
Phenanthrene	630×10^{-9}	4.66×10^{-4}	Mouse	40	1,000	No
Polychlorinated Dibenzodioxins	15×10^{-9}	4.2×10^{-6}	Rat	0.001	1,000	No
Polychlorinated Dibenzofurans	1.1×10^{-9}	3.08×10^{-7}	Rat	6	1,000	No
Pyrene	460×10^{-9}	3.4×10^{-4}	Mouse	250	100	No
Toluene	5.1×10^{-5}	1.4×10^{-2}	Rat	446	100	No
Xylene	1.3×10^{-4}	3.6×10^{-2}	Rat	357	10	No

^aDose = soil ingestion rate (mg soil/kg bw/day) $\times$ maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992, Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern

LOAEL = Lowest observable adverse effect level

D.6.6

26 February 1993

109A0.B01jp

Table D.6.2
Comparison of FT-1 Groundwater ECOPC Terrestrial
Receptor Exposure Concentrations to Applicable PRL Values

Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Applicable Species	Preliminary Remediation Goal Water Criteria ^a (µg/L)	Uncertainty Factor ^b	Potential Risk (yes or no)
Inorganic ECOPCs					
Copper	72	Rat	1,000	100	No
Zinc	163	Rat	7,864	100	No
Organic ECOPCs					
bis(2-ethylhexyl)phthalate	3	Rat	52,000	10	No
1,1,1-Trichloroethane	15	Rat	400,000	10	No
Toluene	1	Rat	1,784	1,000	No

^aWater criteria were derived using the equation:

$$\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) + \text{The Uncertainty Factor} \right]$$

Final units are in µg/L.

^bUncertainty factors applied for derivation of criteria appear in Table D.4.2.

Table D.6.3 Comparison of FT-2 Soil ECOPC Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable PRL Values						
ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Inorganic ECOPCs						
Chromium (Total)	2.3×10^{-5}	6.5×10^{-3}	Rat	2,160 ^b	1,000	No
Lead	$2,300 \times 10^{-9}$	6.4×10^{-4}	Rat	2.8	10	Yes
Organic ECOPCs						
2-Butanone	$8,600 \times 10^{-9}$	2.41×10^{-3}	Rat	1,080	1,000	No
2-Methylnaphthalene	4.5×10^{-5}	1.2×10^{-2}	Rat	4,360	1,000	No
4-Methyl-2-pentanone	1.5×10^{-8}	4.2×10^{-6}	Rat	250	100	No
Acetone	$5,400 \times 10^{-9}$	N/A	N/A	N/A	N/A	?
Benzene	3.3×10^{-5}	9.24×10^{-3}	Rat	235	1,000	Yes
bis(2-ethylhexyl)phthalate	2.2×10^{-6}	6.0×10^{-4}	Rat	200	10	No
Chlorobenzene	18×10^{-9}	5.04×10^{-6}	Rat	2,290	1,000	No
Dibenzofuran	1.7×10^{-6}	4.76×10^{-4}	Rat	6	1,000	No
Ethylbenzene	130×10^{-6}	3.64×10^{-2}	Rat	291	100	No
Fluoranthene	1.5×10^{-6}	1.1×10^{-3}	Mouse	250	100	No
Fluorene	$1,600 \times 10^{-9}$	4.5×10^{-4}	Rat	250	100	No
Methylene Chloride	$3,400 \times 10^{-9}$	9.52×10^{-4}	Rat	52.58	10	No
n-Nitrosodiphenylamine	950×10^{-9}	N/A	N/A	N/A	N/A	?
Naphthalene	27×10^{-6}	1.9×10^{-2}	Mouse	300	1,000	No
Phenanthrene	1.9×10^{-6}	1.4×10^{-3}	Mouse	40	1,000	No
Pyrene	1.3×10^{-6}	9.6×10^{-4}	Mouse	250	100	No
Styrene	22×10^{-9}	6.16×10^{-6}	Rat	5,000	1,000	No
Tetrachloroethene	8.6×10^{-6}	6.4×10^{-3}	Mouse	71	1,000	No
Toluene	6.1×10^{-4}	1.7×10^{-1}	Rat	446	100	No
Xylene (Total)	7.9×10^{-4}	2.2×10^{-1}	Rat	357	10	No

^aDose = soil ingestion rate (mg soil/kg bw/day) × maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern.
LOAEL = Lowest observable adverse effect level.

Table D.6.4 Comparison of FT-2 Groundwater ECOPC Terrestrial Receptor Exposure Concentrations to Applicable PRL Values					
Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Applicable Species	Preliminary Remediation Goal Water Criteria ^a (µg/L)	Uncertainty Factor ^b	Potential Risk (yes or no)
Inorganic ECOPCs					
Chromium (Total)	15.6	Rat	368	10	c
Zinc	87.6	Rat	7,864	100	No
Organic ECOPCs					
2-Methylnaphthylene	1	Rat	3.48	10,000	No
2-Methylphenol	18	Rat	4,000	100	No
2,4-Dimethylphenol	31	Mouse	2,500	100	No
4-Methylphenol	31	NA	NA	NA	No
Benzene	710	Rat	1,412	100	No
Chloroform	2	Rat	48,000	10	No
Ethyl Benzene	680	Rat	7,768	100	No
Methylene Chloride	420	Rat	4,680	10	No
Naphthalene	32	Rat	32,800	10	No
Phenol	11	Rat	480	1,000	No
Toluene	1,400	Rat	1,784	1,000	No
Xylenes (Total)	3,600	Rat	143,200	10	No
^a Water criteria were derived using the equation: $\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) + \text{The Uncertainty Factor} \right]$ Final units are in µg/L. ^b Uncertainty factors applied for derivation of criteria appear in Table D.4.2. ^c Although the maximum observed concentrations fall below the water criteria, these ECOPCs may be of concern--due to their toxicity.					

6.3.3 SW-4

6.3.3.1 Terrestrial Risk Characterization. Results comparing derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.5. Results of these comparisons do not indicate a potential risk to terrestrial organisms.

6.3.3.2 Aquatic Risk Characterization. Because of the lack of appropriate media data, aquatic risk was not identified for this area of concern.

6.3.4 SW-5

6.3.4.1 Terrestrial Risk Characterization. Results comparing derived sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.6. Results of these comparisons indicate that the observed concentrations of barium and copper in sediment may pose a potential risk to terrestrial organisms exposed to the sediment through incidental ingestion. However, due to the overconservative nature of these calculations, the actual risk is uncertain and requires further investigation. Observed maximum concentrations of ECOPCs in the water sources are compared to appropriate water criteria in Table D.6.7. Results of these comparisons indicate that the presence of copper and zinc in water resources at the site poses a potential risk to terrestrial receptors. It is unknown whether the presence of nitrate will pose an adverse effect on ecological receptors.

6.3.4.2 Aquatic Risk Characterization. Observed maximum concentrations of water and sediment related ECOPCs are compared to applicable ambient water quality criteria and sediment quality criteria in Table D.6.8. Results of these comparisons indicate a potential risk to aquatic receptors exposed to sediment and surface water due to the presence of copper, chromium, lead, and zinc. This potential risk could be further investigated with the possible use of either in-situ or laboratory bioassays. In addition, it is recommended that further sediment and surface water samples be collected to fully characterize the ecological setting.

6.3.5 SW-10

6.3.5.1 Terrestrial Risk Characterization. Results comparing derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.9. Observed maximum concentrations of ECOPCs in the water sources are compared to appropriate water criteria in Table D.6.10. Results of these comparisons indicate a potential risk is posed to terrestrial organisms exposed to soils resulting from cadmium, lead, and barium contamination. Results of the water

Table D.6.5
Comparison of SW-4 Soil ECOPC Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable PRL Values

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Inorganic ECOPCs						
Arsenic	2.5×10^{-5}	7.0×10^{-3}	Rat	6.4	10	No
Cadmium	2×10^{-6}	5.6×10^{-7}	Rat	0.014	10	No
Chromium (Total)	1.07×10^{-4}	2.9×10^{-2}	Rat	2,160 ^b	1,000	No
Copper	6.61×10^{-5}	4.8×10^{-2}	Mouse	4.2	10	No
Lead	5.34×10^{-4}	1.49×10^{-1}	Rat	2.8	10	No
Nickel	9.65×10^{-5}	2.7×10^{-2}	Rat	50	10	No
Zinc	1.43×10^{-4}	1.05×10^{-1}	Mouse	38	10	No
Organic ECOPCs						
2-Butanone	8.2×10^{-8}	2.3×10^{-5}	Rat	1,080	1,000	No
4,4'-DDD	5.6×10^{-9}	1.57×10^{-6}	Rat	10	10	No
Benzo(a)pyrene	4.3×10^{-8}	3.2×10^{-5}	Mouse	10	10	No
bis(2-ethylhexyl)phthalate	9.1×10^{-8}	2.5×10^{-5}	Rat	200	10	No
Di-n-butylphthalate	1.6×10^{-7}	4.5×10^{-5}	Rat	8,000	1,000	No
Indeno(1,2,3-cd)pyrene	4.5×10^{-8}	3.3×10^{-5}	Mouse	10	100	No
Methylene chloride	5.4×10^{-8}	1.5×10^{-5}	Rat	52.58	10	No
PCB (Aroclor 1260)	1.2×10^{-7}	3.4×10^{-5}	Rat	841	1,000	No
Tetrachloroethylene	2×10^{-9}	1.48×10^{-6}	Mouse	71	1,000	No
Toluene	3.8×10^{-8}	1.06×10^{-5}	Rat	446	100	No
Xylene	4.23×10^{-8}	1.18×10^{-5}	Rat	357	10	No

^aDose = soil ingestion rate (mg soil/kg bw/day) x maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern.

LOAEL = Lowest observable adverse effect level.

Comparison of SW-5 Sediment ECOPC Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable PRL Values						
ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Inorganic ECOPCs						
Barium	9.47×10^{-5}	2.65×10^{-2}	Rat	0.25	100	Yes
Chromium (Total)	3.4×10^{-5}	9.55×10^{-2}	Rat	2,160 ^b	1,000	No
Copper	5.53×10^{-5}	4.1×10^{-2}	Mouse	4.2	10	No
Lead	3.6×10^{-5}	1.0×10^{-2}	Rat	2.8	10	No
Zinc	1.01×10^{-4}	7.4×10^{-2}	Mouse	38	10	No
Organic ECOPCs						
2-Butanone	1.8×10^{-8}	5.04×10^{-6}	Rat	1,080	1,000	No
4,4'-DDD	5.8×10^{-9}	1.6×10^{-6}	Rat	10	10	No
bis(2-ethylhexyl)phthalate	1.3×10^{-7}	3.6×10^{-5}	Rat	200	10	No
Endrin Ketone	2.1×10^{-8}	5.8×10^{-6}	Rat	3	1,000	No
Fluoranthene	4.6×10^{-8}	3.4×10^{-5}	Mouse	250	100	No
Fluorene	7.8×10^{-8}	5.77×10^{-5}	Mouse	250	100	No
Pyrene	8.8×10^{-8}	6.5×10^{-5}	Mouse	250	100	No

^aDose = soil ingestion rate (mg soil/kg bw/day) $\times$ maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern.

LOAEL = Lowest observable adverse effect level.

Table D.6.7 Comparison of SW-5 Surface Water ECOPC Terrestrial Receptor Exposure Concentrations to Applicable PRL Values					
Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Applicable Species	Preliminary Remediation Goal Water Criteria ^a (µg/L)	Uncertainty Factor ^b	Potential Risk (yes or no)
Inorganic ECOPCs					
Copper	48,800	Rat	1,000	100	Yes
Zinc	77,200	Rat	7,864	100	Yes
Organic ECOPCs					
bis(2-ethylhexyl)phthalate	1	Rat	52,000	10	No
Nitrate	1,410	NA	NA	NA	?
^a Water criteria were derived using the equation: $\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) + \text{The Uncertainty Factor} \right]$ Final units are in µg/L. ^b Uncertainty factors applied for derivation of criteria appear in Table D.4.2.					

Table D.6.8
Comparison of SW-5 Surface Water and Sediment ECOPC Aquatic Receptor Exposure
Concentrations to Applicable PRL Values

Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/kg or µg/L)	Sediment Criteria			Water Criteria (µg/L)	Potential Risk (yes or no)
		Freshwater	Marine	Freshwater Acute		
Inorganic ECOPCs						
Barium	94,700	NA	NA	NA	NA	?
Chromium (Total)	34,100	9,350	42,500	NA	NA	Yes
Copper (SW)	48,800	NA	NA	12	2.4	Yes
Copper (Sediment)	55,300	420	102	NA	NA	Yes
Lead	36,100	2,880	198,000	NA	NA	Yes
Zinc (SW)	77,200	NA	NA	110	86	Yes
Zinc (Sediment)	101,000	4,400	3,440	NA	NA	Yes
Organic ECOPCs						
2-Butanone	18	NA	NA	NA	NA	?
4,4'-DDD	5.8	485	2,912	NA	NA	No
bis(2-ethylhexyl)phthalate (SW)	1	NA	NA	360	360	No
bis(2-ethylhexyl)phthalate (Sediment)	130	33,120	33,120	NA	NA	No
Endrin Ketone	21	NA	NA	NA	NA	?
Fluoranthene	46	159,200	640	NA	NA	No
Fluorene	78	NA	NA	NA	NA	?
Pyrene	88	NA	12,000	NA	NA	No
Final units are in µg/L. NA = Water quality criteria are not available.						

Comparison of SW-10 Soil and Sediment ECOPC Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable PRL Values							Page 1 of 2
Table D.6.9							
ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)	
Inorganic ECOPCs							
Arsenic (soil)	5.3 x 10 ⁻⁵	1.48 x 10 ⁻²	Rat	6.4	10	No	
Arsenic (sediment)	6.9 x 10 ⁻⁶	1.9 x 10 ⁻³	Rat	6.4	10	No	
Barium	5.6 x 10 ⁻⁴	1.56 x 10 ⁻¹	Rat	0.25	100	Yes	
Cadmium	5.8 x 10 ⁻⁶	1.6 x 10 ⁻³	Rat	0.014	10	Yes	
Chromium (total) (soil)	8.13 x 10 ⁻⁵	2.3 x 10 ⁻²	Rat	2,160 ^b	1,000	No	
Chromium (sediment)	1.78 x 10 ⁻⁵	4.9 x 10 ⁻³	Rat	2,160 ^b	1,000	No	
Copper (soil)	8.92 x 10 ⁻⁴	6.6 x 10 ⁻¹	Mouse	4.2	10	No	
Lead (soil)	1.74 x 10 ⁻³	4.76 x 10 ⁻¹	Rat	2.8	10	Yes	
Lead (sediment)	2.4 x 10 ⁻⁵	6.6 x 10 ⁻³	Rat	2.8	10	No	
Nickel	5.73 x 10 ⁻⁵	1.6 x 10 ⁻²	Rat	50	10	No	
Zinc (soil)	2.11 x 10 ⁻³	1.554 x 10 ⁻¹	Mouse	38	10	No	
Zinc (sediment)	5.95 x 10 ⁻⁵	4.4 x 10 ⁻²	Mouse	38	10	No	
Organic ECOPCs							
2-Butanone	1.0 x 10 ⁻⁶	2.8 x 10 ⁻⁴	Rat	1,080	1,000	No	
4',4'-DDD	1.5 x 10 ⁻⁸	4.2 x 10 ⁻⁶	Rat	10	10	No	
4',4'-DDE	3.2 x 10 ⁻⁹	NA	NA	NA	NA	?	
4',4'-DDT	1.2 x 10 ⁻⁸	3.36 x 10 ⁻⁶	Rat	0.25	10	No	
Anthracene	6.8 x 10 ⁻⁸	1.9 x 10 ⁻⁵	Rat	1,000	100	No	
Benzo(a)anthracene	1.3 x 10 ⁻⁷	9.62 x 10 ⁻⁵	Mouse	10	100	No	
Benzo(a)pyrene	3.3 x 10 ⁻⁷	2.4 x 10 ⁻⁴	Mouse	10	10	No	
Benzo(b)fluoranthene	3.0 x 10 ⁻⁷	2.22 x 10 ⁻⁴	Mouse	10	100	No	
Benzo(g,h,i)berylene	2.7 x 10 ⁻⁷	1.9 x 10 ⁻⁴	Mouse	10	100	No	
Benzo(k)fluoranthene	2.7 x 10 ⁻⁷	1.9 x 10 ⁻⁴	Mouse	10	100	No	
bis(2-ethylhexyl)phthalate (sediment)	1.1 x 10 ⁻⁷	3.1 x 10 ⁻⁵	Rat	200	10	No	
bis(2-ethylhexyl)phthalate (soil)	5.3 x 10 ⁻⁶	1.5 x 10 ⁻³	Rat	200	10	No	

D.6.15

26 February 1993

100109AC.B01/jp

Table D.6.9 Comparison of SW-10 Soil and Sediment ECOPC Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable PRL Values							Page 2 of 2
ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)	
Carbon disulfide	1.0 x 10 ⁻⁹	2.8 x 10 ⁻⁷	Rat	3,188	1,000	No	
Chloroform	2.0 x 10 ⁻⁹	5.6 x 10 ⁻⁷	Rat	908	100 ^c	No	
Chrysene	3.9 x 10 ⁻⁷	2.8 x 10 ⁻⁴	Mouse	10	100	No	
Di-n-butylphthalate	4.6 x 10 ⁻⁸	1.3 x 10 ⁻⁵	Rat	8,000	1,000	No	
Endrin ketone	2.1 x 10 ⁻⁹	5.8 x 10 ⁻⁷	Rat	3	1,000	No	
Ethyl benzene	5.02 x 10 ⁻⁸	1.4 x 10 ⁻⁵	Rat	291	100	No	
Fluoranthene	6.8 x 10 ⁻⁸	5.0 x 10 ⁻⁵	Mouse	250	100	No	
Indeno(1,2,3-cd)pyrene	3.0 x 10 ⁻⁷	2.22 x 10 ⁻⁴	Mouse	10	100	No	
Methylene chloride	1.4 x 10 ⁻⁸	3.9 x 10 ⁻⁶	Rat	52.58	10	No	
PCB-1248	1.5 x 10 ⁻⁷	4.2 x 10 ⁻⁵	Rat	0.8	1,000	No	
PCB-1254	1.1 x 10 ⁻⁷	3.08 x 10 ⁻⁵	Rat	0.5	1,000	No	
PCB-1260	6.3 x 10 ⁻⁷	1.7 x 10 ⁻⁴	Rat	1.3	1,000	No	
Phenanthrene	5.4 x 10 ⁻⁸	3.99 x 10 ⁻⁵	Mouse	40	1,000	No	
Pyrene	8.0 x 10 ⁻⁸	5.92 x 10 ⁻⁵	Mouse	250	100	No	
Toluene	6.28 x 10 ⁻⁸	1.75 x 10 ⁻⁵	Rat	446	100	No	
Xylenes	1.71 x 10 ⁻⁷	4.78 x 10 ⁻⁵	Rat	357	10	No	

^aDose = soil ingestion rate (mg soil/kg bw/day) x maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

^cUncertainty factor encompasses intra and interspecies extrapolations (dog - rat).

Notes: ECOPC = Ecological chemical of potential concern.
LOAEL = Lowest observable adverse effect level.

^aDose = soil ingestion rate (mg soil/kg bw/day) × maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Lucky, 1978.

^cUncertainty factor encompasses intra and interspecies extrapolations (dog ~ rat).

Notes: ECOPC = Ecological chemical of potential concern.

LOAEL = Lowest observable adverse effect level.

Table D.6.10
Comparison of SW-10 Groundwater ECOPC Terrestrial
Receptor Exposure Concentrations to Applicable PRL Values

Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Applicable Species	Preliminary Remediation Goal Water Criteria ^a (µg/L)	Uncertainty Factor ^b	Potential Risk (yes or no)
Inorganic ECOPCs					
Chromium (Total)	48.3	Rat	368	10	No
Copper	202	Rat	1,000	100	No
Zinc	700	Rat	7,864	100	No
Organic ECOPCs					
Acetone	11	N/A ^c	N/A ^c	100	d
Benzene	5.1	Rat	1,412	100	No
bis(2-ethylhexyl)phthalate	2	Rat	52,000	10	No
Carbon Disulfide	3	Rat	880	100	No
Dibenzofuran	1	N/A	N/A	N/A	d
Diethylphthalate	2	N/A	N/A	N/A	d
Ethyl Benzene	1.6	Rat	7,768	100	No
Flourene	2	N/A	N/A	N/A	d
2-Methylnaphthalene	12	Rat	3.48	10,000	Yes
Naphthalene	8	Rat	32,800	10	No
Toluene	2.9	Rat	1,784	1,000	No
Xylenes (Total)	6.2	Rat	143,200	10	No

^aWater criteria were derived using the equation:

$$\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) + \text{The Uncertainty Factor} \right]$$

Final units are in µg/L.

^bUncertainty factors applied for derivation of criteria appear in Table D.4.2.

^cWater criteria were not derived for acetone, since all reviewed toxicological literature was derived from inhalation exposure.

^dAlthough the maximum observed concentrations fall below the water criteria, these ECOPCs may be of concern—due to their toxicity.

ingestion analysis indicate a potential risk to terrestrial life exists from the presence of methyl-naphthalene. It is suggested that these media be further investigated to assess in-situ toxicity.

6.3.5.2 Aquatic Risk Characterization. Observed maximum concentrations of water and sediment-related ECOPCs are compared to applicable ambient water quality criteria and sediment quality criteria in Table D.6.11. Results of these comparisons indicate a potential risk to aquatic organisms exists from the presence of chromium, lead, and zinc.

6.3.6 SW-12

6.3.6.1 Terrestrial Risk Characterization. Results comparing of derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.12. Observed maximum concentrations of ECOPCs in the water sources are compared to appropriate water criteria in Table D.6.13. Results of these comparisons indicate there is risk posed to terrestrial organisms exposed to barium and nickel in the soil.

6.3.6.2 Aquatic Risk Characterization. Because of the lack of appropriate media data, aquatic organism risk and exposure could not be addressed.

6.3.7 Water Gallery

6.3.7.1 Terrestrial Risk Characterization. Results comparing derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.14. Observed maximum concentrations of ECOPCs in the water sources are compared to appropriate water criteria in Table D.6.15. Results of these comparisons indicate a potential risk to terrestrial receptors exposed to soil from the presence of Indeno(1,2,3-cd) pyrene, phenanthrene, and antimony at the site. Antimony, aluminum, cobalt, and iron in the water source could pose a risk to terrestrial receptors.

6.3.7.2 Aquatic Risk Characterization. Because of the lack of appropriate media data, aquatic organism exposure and risk was not assessed.

Table D.6.11 Comparison of SW-10 Surface Water and Sediment ECOPC Aquatic Receptor Exposure Concentrations to Applicable PRL Values				
Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Sediment Criteria		Potential Risk (yes or no)
		Freshwater	Marine	
Inorganic ECOPCs				
Arsenic	6,900	170,000	463,800	No
Chromium (Total)	17,800	9,350	42,500	Yes
Lead	23,800	2,880	7,650	Yes
Zinc	59,500	4,400	3,440	Yes
Organic ECOPCs				
bis(2-ethylhexyl)phthalate	110	33,120	33,120	No

Table D.6.12
Comparison of SW-12 Soil and Sediment ECOPC Terrestrial Receptor
Incidental Ingestion Dose Concentrations to Applicable PRL Values

Page 1 of 2

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
4,4'-DDD (sediment)	5.3×10^{-9}	1.4×10^{-6}	Rat	10	10	No
4,4'-DDD (soil)	1.5×10^{-6}	4.2×10^{-6}	Rat	10	10	No
4,4'-DDE (sediment)	2.6×10^{-9}	N/A	Rat	N/A	N/A	?
4,4'-DDE (soil)	2.7×10^{-9}	N/A	Rat	N/A	N/A	?
4,4'-DDT (soil)	3.4×10^{-8}	9.5×10^{-6}	Rat	0.25	10	No
4,4'-DDT (sediment)	1.7×10^{-8}	4.7×10^{-6}	Rat	0.25	10	No
Acetone	2.1×10^{-8}	NA	NA	NA	NA	?
Antimony	8.7×10^{-6}	2.4×10^{-3}	Rat	0.35	10	No
Aroclor 1260	8.9×10^{-8}	2.5×10^{-5}	Rat	1.3	1,000	No
Aroclor 1254	8.7×10^{-8}	2.4×10^{-5}	Rat	0.5	1,000	No
Arsenic (sediment)	5.5×10^{-6}	1.5×10^{-3}	Rat	6.4	10	No
Arsenic (soil)	1.8×10^{-5}	5.0×10^{-3}	Rat	6.4	10	No
Barium	3.3×10^{-4}	.092	Rat	0.25	100	Yes
Benzene	4.9×10^{-8}	1.37×10^{-5}	Rat	3,306	1,000	No
bis(2-ethylhexyl)phthalate (soil)	7.8×10^{-7}	2.1×10^{-4}	Rat	200	10	No
bis(2-ethylhexyl)phthalate (sediment)	3.9×10^{-7}	1.1×10^{-4}	Rat	200	10	No
Cadmium (sediment)	2.1×10^{-6}	5.8×10^{-4}	Rat	0.014	10	No
Carbon Disulfide	2×10^{-9}	5.6×10^{-7}	Rat	3,188	1,000	No
Chloroform	1×10^{-9}	2.8×10^{-7}	Rat	15	100	No
Chromium (sediment)	3.1×10^{-5}	8.6×10^{-3}	Rat	2,160	1,000	No
Chromium (Total) (soil)	5.9×10^{-5}	1.6×10^{-2}	Rat	2,160 ^b	1,000	No
Copper (soil)	1.4×10^{-4}	1.04×10^{-1}	Mouse	4.2	10	No
Copper (sediment)	3.7×10^{-4}	2.7×10^{-1}	Mouse	4.2	10	No
Di-n-butylphthalate	5.0×10^{-6}	1.4×10^{-3}	Rat	125	100	No
Diethylphthalate	1.3×10^{-7}	3.6×10^{-5}	Rat	3,160	100	No

Table D.6.12
Comparison of SW-12 Soil and Sediment ECOPC Terrestrial Receptor
Incidental Ingestion Dose Concentrations to Applicable PRL Values

Page 2 of 2

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Ethyl Benzene	7.8×10^{-7}	2.2×10^{-4}	Rat	291	10	No
Lead (sediment)	3.8×10^{-4}	1.06×10^{-1}	Rat	2.8	10	No
Lead (soil)	3.35×10^{-4}	9.4×10^{-2}	Rat	2.8	10	No
Methylene Chloride	1.9×10^{-8}	5.3×10^{-6}	Rat	52.58	10	No
Nickel	6.24×10^{-5}	1.74×10^{-2}	Rat	50	10	No
Toluene	9.9×10^{-7}	2.7×10^{-4}	Rat	446	100	No
Xylenes	2.6×10^{-6}	7.3×10^{-4}	Rat	357	10	No
Zinc (soil)	4.19×10^{-4}	3.1×10^{-1}	Mouse	38	10	No
Zinc (sediment)	5.6×10^{-4}	4.14×10^{-1}	Mouse	38	10	No

^aDose = soil ingestion rate (mg soil/kg bw/day) $\times$ maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern.

LOAEL = Lowest observable adverse effect level.

D.6.21

26 February 1993

Table D.6.13
Comparison of SW-12 Groundwater ECOPC Terrestrial
Receptor Exposure Concentrations to Applicable PRL Values

Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Applicable Species	Preliminary Remediation Goal Water Criteria ^a (µg/L)	Uncertainty Factor ^b	Potential Risk (yes or no)
Acetone	21	N/A ^c	N/A ^c	100	^d
Aroclor 1254	87	Rat	4	1,000	No
Arsenic	66.9	Dog	2,400	10	No
Benzene	1.9	Rat	1,412	100	No
bis(2-ethylhexyl)phthalate	1	Rat	52,000	10	No
Carbon Disulfide	4	Rat	880	10	No
Chromium (Total)	80.7	Rat	368	10	No
Copper	255	Rat	1,000	100	No
Toluene	0.5	Rat	1,784	1,000	No
Zinc	369	Rat	7,864	100	No

^aWater criteria were derived using the equation:

$$\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) \cdot \text{The Uncertainty Factor} \right]$$

Final units are in µg/L.

^bUncertainty factors applied for derivation of criteria appear in Table D.4.2.

^cWater criteria were not derived for ammonia and acetone, since all reviewed toxicological literature was derived from inhalation exposure.

^dAlthough the maximum observed concentrations fall below the water criteria, these ECOPCs may be of concern—due to their toxicity.

Table D.6.14
Comparison of WG Soil ECOPC Terrestrial Receptor Incidental
Ingestion Dose Concentrations to Applicable PRL Values

Page 1 of 3

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
2-Butanone	2×10^{-7}	5.7×10^{-5}	Rat	1,080	1,000	No
2-Methylnaphthalene	1.2×10^{-6}	3.4×10^{-4}	Rat	4,360	1,000	No
4-Methylphenol	1.0×10^{-6}	2.8×10^{-4}	Rat	150	100	No
Acenaphthene	1.5×10^{-5}	1.11×10^{-2}	Mouse	350	100	No
Acetone	9.2×10^{-7}	2.6×10^{-4}	Rat	500	100	No
Aluminum	4.19×10^{-2}	11.7	Rat	14	100	No
Anthracene	3.8×10^{-5}	2.8×10^{-2}	Mouse	1,000	1,000	No
Antimony	2.48×10^{-5}	6.9×10^{-3}	Rat	0.35	100	Yes
Antimony	8.7×10^{-6}	2.4×10^{-3}	Rat	0.35	10	No
Aroclor 1254	3.6×10^{-8}	1.0×10^{-5}	Rat	0.5	1,000	No
Arsenic	6.8×10^{-6}	1.9×10^{-3}	Rat	6.4	10	No
Barium	6.05×10^{-5}	1.7×10^{-2}	Rat	54	1,000	No
Benzo(a)anthracene	4.3×10^{-5}	3.2×10^{-2}	Mouse	10	100	No
Benzo(a)pyrene	3.6×10^{-5}	2.6×10^{-2}	Mouse	10	10	No
Benzo(b)fluoranthene	6.1×10^{-5}	4.5×10^{-2}	Mouse	10	100	No
Benzo(g,h,i)perylene	1.3×10^{-5}	9.6×10^{-3}	Mouse	10	100	No
Benzo(k)fluoranthene	2.1×10^{-5}	1.55×10^{-2}	Mouse	10	100	No
Beryllium	4.0×10^{-7}	1.12×10^{-4}	Rat	0.85	10	No
bis(2-ethylhexyl)phthalate	2.1×10^{-6}	5.8×10^{-4}	Rat	200	10	No
Cadmium	2.2×10^{-6}	6.16×10^{-4}	Rat	0.014	10	No
Carbazole	1.3×10^{-5}	3.6×10^{-3}	Rat	500	1,000	No
Chloromethane	5.7×10^{-8}	1.6×10^{-5}	Rat	60	100	No
Chromium (Total)	3.9×10^{-5}	1.08×10^{-2}	Rat	2,160 ^b	1,000	No
Chrysene	3.5×10^{-5}	2.59×10^{-2}	Mouse	10	100	No
Cobalt	9.1×10^{-6}	2.5×10^{-3}	Rat	0.5	100	No

D.6.23

26 February 1993

100109B2.ROI/jp

Table D.6.14
Comparison of WG Soil ECOPC Terrestrial Receptor Incidental
Ingestion Dose Concentrations to Applicable PRL Values

Page 2 of 3

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal(LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Copper	1.2×10^{-4}	8.8×10^{-2}	Mouse	4.2	10	No
Cyanoide	2.0×10^{-6}	5.6×10^{-4}	Rat	30	10	No
Dibenzo(a,h)anthracene	4.8×10^{-6}	3.5×10^{-3}	Mouse	10	100	No
Dibenzofuran	1.2×10^{-5}	3.36×10^{-3}	Rat	6	1,000	No
Di-n-butylphthalate	8.6×10^{-7}	2.4×10^{-4}	Rat	125	100	No
Endrin Ketone	3.4×10^{-9}	9.5×10^{-7}	Rat	3	1,000	No
Ethyl Benzene	4.1×10^{-8}	1.14×10^{-3}	Rat	291	100	No
Fluoranthene	9.6×10^{-5}	7.1×10^{-2}	Mouse	250	100	No
Fluorene	2.1×10^{-5}	1.5×10^{-2}	Mouse	250	100	No
Indeno(1,2,3-cd)pyrene	2.0×10^{-5}	1.5×10^{-2}	Mouse	10	100	Yes
Iron	2.82×10^{-2}	7.8	Rat	900	1,000	Yes
Lead	1.4×10^{-4}	3.8×10^{-2}	Rat	2.8	10	No
Manganese	3.79×10^{-4}	1.06×10^{-1}	Rat	930	10	No
Mercury	4.0×10^{-7}	1.12×10^{-4}	Rat	0.015	10	No
Methylene Chloride	1.7×10^{-8}	4.8×10^{-6}	Rat	52.58	10	No
Naphthalene	1.6×10^{-6}	1.18×10^{-3}	Mouse	300	1,000	No
Nickel	2.3×10^{-5}	6.4×10^{-3}	Rat	50	10	No
Phenanthrene	1.1×10^{-4}	8.14×10^{-2}	Mouse	40	1,000	Yes
Phenol	1.6×10^{-7}	4.48×10^{-5}	Rat	120	1,000	No
Pyrene	7.9×10^{-5}	5.8×10^{-2}	Mouse	250	100	No
Selenium	2.0×10^{-6}	5.6×10^{-4}	Rat	0.25	10	No
Silver	6.9×10^{-7}	5.1×10^{-4}	Mouse	18.1	100	No
Thallium	1.3×10^{-6}	3.6×10^{-4}	Rat	0.7	100	No
Toluene	9.8×10^{-8}	2.78×10^{-5}	Rat	446	100	No
Vanadium	2.16×10^{-4}	6.05×10^{-2}	Rat	2.8	10	No

Table D.6.14

Comparison of WG Soil ECOPC Terrestrial Receptor Incidental
Ingestion Dose Concentrations to Applicable PRL Values

Page 3 of 3

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal(LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Xylene	3.5×10^{-7}	9.8×10^{-5}	Rat	357	10	No
Zinc	8.63×10^{-5}	6.38×10^{-2}	Mouse	38	10	No

^aDose = soil ingestion rate (mg soil/kg bw/day) x maximum observed concentration (mg/mg soil).
^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern.
LOAEL = Lowest observable adverse effect level.

D.6.25

26 February 1993

100109B2.B01/jp

Table D.6.15
Comparison of WG Groundwater ECOPC Terrestrial
Receptor Exposure Concentrations to Applicable PRL Values

Ecological Chemical of Potential Concern (ECOPCs)	Maximum Observed Concentration (µg/L)	Applicable Species	Uncertainty Factor ^a	Preliminary Remediation Goal Water Criteria ^b (µg/L)	Potential Risk (yes or no)
1,1-Dichloroethene	1.0	Rat	10	8,000	No
Aluminum	30,400	Mouse	10	9,500	Yes
Antimony	31.10	Rat	100	28	Yes
Arsenic	2.0	Dog	10	2,400	No
Barium	135	Rat/Mouse	10	200 - 410	No
Benzene	1.8	Rat	100	1,412	No
Beryllium	0.75	Rat	10	432	No
Cadmium	2.7	Rat	10	3.2	No
Chromium	53.4	Rat	10	368	No
Cobalt	19.8	Rat	100	4	Yes
Copper	200	Rat	100	1,000	No
Ethyl Benzene	3.7	Rat	100	7,768	No
Iron	17,900	Rat	1,000	7,200	Yes
Lead	8.8	Rat	10	360	No
Manganese	2,170	Rat	10	232,000	No
Mercury	0.2	Rat	10	2.4	No
Nickel	32.2	Rat	10	400	No
Selenium	2.9	Rat	10	400	No
Silver	2.0	Rat	1,000	1,450	No
Thallium	1.6	Rat	0.2	16	No
Toluene	16.3	Rat	1,000	1,784	No
Trichloroethylene	24.9	Mouse	100	920	No
Vanadium	56	Rat	10	560	No
Xylenes (Total)	18.9	Rat	10	143,200	No
Zinc	150	Rat	100	7,864	No

^aUncertainty factors applied for derivation of criteria appear in Table D.4.2.

^bWater criteria were derived using the equation:

$$\frac{\text{Water Criteria}}{\mu\text{g/L}} = \left[\left(\frac{\text{NOAEL (mg/kg bw/day)}}{\text{Water Intake Rate (L/kg bw/day)}} \right) + \text{The Uncertainty Factor} \right]$$

Final units are in µg/L.

6.3.8 PS-1A

6.3.8.1 Terrestrial Risk Characterization. Results comparing of derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.16. Results of these comparisons indicate there is a potential risk posed to terrestrial organisms exposed to site soil due to the presence of aluminum, cobalt, and iron.

6.3.8.2 Aquatic Risk Characterization. Because of the lack of appropriate media data, aquatic organism exposure and risk was not assessed.

6.3.9 PS-1B

6.3.9.1 Terrestrial Risk Characterization. Results comparing derived soil and sediment doses (based on the maximum observed concentration) to NOAELs for soil are presented in Table D.6.17. Results of these comparisons indicate that potential risk to terrestrial organisms does exist because of the presence of cadmium, aluminum, antimony, iron, and silver.

6.3.9.2 Aquatic Risk Characterization. Because of the lack of appropriate media data, aquatic organism risk and exposure could not be addressed.

Table D.6.16
Comparison of PS-1A Soil ECOPC Terrestrial Receptor Incidental
Ingestion Dose Concentrations to Applicable PRL Values

Page 1 of 2						
ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Aluminum	2.4×10^{-2}	6.72	Rat	14	100	Yes
Anthracene	1.2×10^{-7}	8.8×10^{-5}	Mouse	1,000	100	No
Antimony	6.8×10^{-6}	1.9×10^{-3}	Rat	0.35	100	No
Arsenic	2.7×10^{-6}	7.56×10^{-4}	Rat	6.4	10	No
Barium	9.94×10^{-5}	2.78×10^{-2}	Rat	54	1,000	No
Benzo(a)anthracene	3.8×10^{-7}	2.8×10^{-4}	Mouse	10	100	No
Benzo(a)pyrene	3.9×10^{-7}	2.8×10^{-4}	Mouse	10	10	No
Benzo(b)fluoranthene	1.7×10^{-7}	1.3×10^{-4}	Mouse	10	10	No
Benzo(g,h,i)perylene	3.6×10^{-7}	2.6×10^{-4}	Mouse	10	100	No
Beryllium	2.2×10^{-7}	6.16×10^{-5}	Rat	0.85	10	No
bis(2-ethylhexyl)phthalate	3.0×10^{-7}	8.4×10^{-5}	Rat	200	10	No
Cadmium	2.2×10^{-6}	6.2×10^{-4}	Rat	0.014	10	No
Carbazole	8.2×10^{-8}	2.3×10^{-5}	Rat	500	1,000	No
Chromium (Total)	2.6×10^{-5}	5.0×10^{-3}	Rat	2,160 ^b	1,000	No
Chrysene	4.1×10^{-7}	3.03×10^{-4}	Mouse	10	100	No
Cobalt	2.73×10^{-5}	7.6×10^{-3}	Rat	0.5	100	Yes
Copper	1.33×10^{-4}	4.8×10^{-2}	Mouse	4.2	10	No
Cyanide	5.5×10^{-7}	1.5×10^{-4}	Rat	30	10	No
Dibenzo(a,h)anthracene	7.9×10^{-8}	5.8×10^{-5}	Mouse	10	100	No
Fluoranthene	1.1×10^{-6}	8.14×10^{-4}	Mouse	250	100	No
Indeno(1,2,3-cd)pyrene	4.4×10^{-7}	3.3×10^{-4}	Mouse	10	100	No
Iron	5.96×10^{-2}	16.68	Rat	900	1,000	Yes
Lead	2.9×10^{-5}	8.2×10^{-3}	Rat	2.8	10	No
Manganese	1.16×10^{-3}	0.0325	Rat	930	10	No
Mercury	1.7×10^{-7}	4.76×10^{-5}	Rat	0.015	10	No

Table D.6.16
Comparison of PS-1A Soil ECOPC Terrestrial Receptor Incidental
Ingestion Dose Concentrations to Applicable PRL Values

Page 2 of 2

ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)
Nickel	2.36×10^{-5}	6.61×10^{-3}	Rat	50	10	No
Phenanthrene	5.3×10^{-7}	3.9×10^{-4}	Mouse	40	1,000	No
Pyrene	6.0×10^{-7}	4.4×10^{-4}	Mouse	250	100	No
Silver	1.6×10^{-6}	1.18×10^{-3}	Mouse	18.1	100	No
Thallium	5.7×10^{-7}	1.59×10^{-4}	Rat	0.7	100	No
Toluene	1×10^{-9}	2.8×10^{-7}	Rat	446	100	No
Vanadium	2.13×10^{-4}	5.96×10^{-2}	Rat	2.8	10	No
Zinc	9.6×10^{-5}	7.1×10^{-2}	Mouse	38	10	No

^aDose = soil ingestion rate (mg soil/kg bw/day) $\times$ maximum observed concentration (mg/mg soil).

^bLOAEL values were not available in the literature reviewed. Values reported are LD50 oral toxicity values for rats (units mg/kg bw/day). Source: Sax, 1992; Venugopol and Luckey, 1978.

Notes: ECOPC = Ecological chemical of potential concern.
 LOAEL = Lowest observable adverse effect level.

D.6.29

26 February 1993

100109B6.B01/jp

Table D.6.17 Comparison of PS-1B Soil ECOPC Terrestrial Receptor Incidental Ingestion Dose Concentrations to Applicable PRL Values							Page 1 of 2
ECOPC	Maximum Observed Concentration (mg/mg soil)	Dose ^a (mg/kg bw/day)	Laboratory Species	Preliminary Remediation Goal (LOAEL) (mg/kg bw/day)	Uncertainty Factor	Potential Risk (Yes or No)	
1,1-Dichloroethane	6.5 x 10 ⁻⁷	4.81 x 10 ⁻⁴	Mouse	543	100	No	
1,1-Dichloroethene	9.7 x 10 ⁻⁷	2.7 x 10 ⁻⁴	Rat	9	10	No	
1,1,1-Trichloroethane	6.0 x 10 ⁻⁷	1.68 x 10 ⁻⁴	Rat	750	10	No	
1,2-Dichloroethane	3.1 x 10 ⁻⁷	8.7 x 10 ⁻⁵	Rat	47	10	No	
4,4'-DDD	3.4 x 10 ⁻⁸	9.5 x 10 ⁻⁶	Rat	10	10	No	
Aluminum	4.13 x 10 ⁻²	11.5	Rat	14	100	Yes	
Antimony	1.63 x 10 ⁻⁴	4.6 x 10 ⁻²	Rat	0.35	100	Yes	
Arsenic	6.1 x 10 ⁻⁶	1.71 x 10 ⁻³	Rat	5.4	10	No	
Barium	4.53 x 10 ⁻⁴	1.26 x 10 ⁻³	Rat	54	1,000	No	
Beryllium	5.2 x 10 ⁻⁷	1.4 x 10 ⁻⁴	Rat	0.85	10	No	
bis(2-ethylhexyl)phthalate	8.8 x 10 ⁻⁵	2.5 x 10 ⁻²	Rat	200	10	No	
Cadmium	2.9 x 10 ⁻⁵	8.1 x 10 ⁻³	Rat	0.014	10	Yes	
Chromium (Total)	3.5 x 10 ⁻⁵	9.8 x 10 ⁻³	Rat	2,160 ^b	1,000	No	
Cobalt	7.7 x 10 ⁻⁶	2.15 x 10 ⁻³	Rat	0.5	100	No	
Copper	1.45 x 10 ⁻⁴	1.07 x 10 ⁻¹	Mouse	4.2	10	No	
Cyanide	2.2 x 10 ⁻⁶	6.2 x 10 ⁻⁴	Rat	30	10	No	
Iron	2.82 x 10 ⁻²	7.89	Rat	900	1,000	Yes	
Lead	5.2 x 10 ⁻⁵	1.44 x 10 ⁻²	Rat	2.8	10	No	
Manganese	2.1 x 10 ⁻⁴	5.8 x 10 ⁻²	Rat	930	10	No	
Mercury	5.4 x 10 ⁻⁶	1.5 x 10 ⁻³	Rat	0.015	10	No	
Nickel	6.72 x 10 ⁻⁴	1.8 x 10 ⁻¹	Rat	50	10	No	
Selenium	1.61 x 10 ⁻⁵	4.5 x 10 ⁻³	Rat	0.25	10	No	
Silver	4.97 x 10 ⁻⁴	3.6 x 10 ⁻¹	Mouse	18.1	100	Yes	
Thallium	1.6 x 10 ⁻⁶	4.5 x 10 ⁻⁴	Rat	0.7	100	No	
Trichloroethane	2.2 x 10 ⁻⁶	1.6 x 10 ⁻³	Mouse	393	100	No	

7.0 Uncertainty Analysis

Assumptions and conclusions made within this BERA are believed to be conservative compared to the true conditions within Shemya Island. However, because of the chemical and physical properties of the chemicals detected within the media of concern, it is difficult to ascertain the true toxicologic potential posed to receptors. This assessment addressed the toxicity posed by individual chemicals in separate media. Organisms onsite can become exposed to all chemical constituents by a combination of exposure routes. Because of the difficulty of addressing multiple route/multiple contaminant exposure toxicity, the total exposure created by exposure to contaminated media cannot be determined.

A number of assumptions were made in determining soil dose concentrations, sediment dose concentrations, and water criteria. With the incorporation of these assumptions and the final selection of ECOPCs by media, an amount of uncertainty was incorporated into the EA.

Sources of uncertainty include:

- Media disturbances between the time of sample collection and the time of this evaluation.
- Alterations in concentrations of chemicals of potential concern because of natural and man-made influences causing either the increase or decrease in ECOPC concentrations.
- Physical and chemical characteristics of the ECOPCs and the media of concern can influence the availability and subsequent toxicity of the ECOPCs.
- Exposure point concentrations were assumed to represent acute and chronic exposure levels.
- Observed detected concentrations of ECOPCs were assumed to be 100 percent bioavailable.
- Physical habitat disturbances cause species composition alterations. New opportunistic species could occupy the areas while others less tolerant to site disturbances could no longer exist.
- Chronic exposure-derived toxicity values were preferably used for dose and criteria calculations. However, when these values were unavailable, subchronic, or acute values were used.

- Between species and within species extrapolations for the toxicity assessment. Many research investigations utilize standard laboratory organisms (Sprague Dawley rats, mice, and specific strains of rabbits). Correlations between these species and the actual receptor species have significant uncertainty (an intraspecies uncertainty factor was applied to every calculation).
- Calculations of surface soil dose, sediment dose, and water criteria associated with no observable adverse effect assume that exposure is strictly confined to surface water and surface soil containing the highest detections of the ECOPCs.
- Uncertainty factors were applied with every calculation. Selection of applicable uncertainty factors was based upon rationale applied in human health assessments and the professional judgement of the author.
- Calculations of soil dose, sediment dose, and water criteria associated with no observable adverse effect do not encompass the possibility of multiple exposure routes to multiple contaminant sources.
- Assessments of soil and sediment dose and water criteria involve species that may not be the most sensitive to contaminant exposure.
- Chronic exposure was assessed in preference to acute; however, certain chemicals can be more acutely toxic than chronically toxic.

Use of the maximum observed concentrations is highly conservative. However, it also adds considerable uncertainty since there was no consideration given to the likely decrease of observed contamination with time because of fate and transport processes such as biodegradation or photolysis.

Several potential ECOPCs did not have ingestion-derived NOAEL and LOAEL values because of their chemical nature. For instance, acetone and benzene have been studied via the inhalation exposure route because of their volatility. The resulting inhalation NOAEL and LOAEL values could not be used to estimate ingestion because of considerable extrapolation between these two exposure routes.

Those chemicals that could not be quantitatively addressed because of the lack of comparable literature values were retained as potential ECOPCs in the media in which they were detected. Using these chemicals as ECOPCs incorporates a great deal of uncertainty since their potential toxicity is unknown.

8.0 References

Abdel-Rahman, M. S., J. J. Saladin, and C. E. Bohman. 1978. *The Effect of 2-hexanone and 2-hexanone Metabolites on Pupillomotor Activity and Growth*. American Industrial Hygiene Association Journal 39:94-99.

Abou-Donia, M. B. 1983. *Interaction between Neurotoxicities Induced by Organophosphorus and Long-Chain Hexacarbon Compounds*. Neurotoxicity 4:117-135.

Agency for Toxic Substances and Disease Registry (ATSDR). 1989a. *Selenium*. United States Department of Health and Human Services. Public Health Service. December 1989.

Agency for Toxic Substances and Disease Registry (ATSDR). 1989c. *Zinc*. United States Department of Health and Human Services. Public Health Service. December 1989.

Agency for Toxic Substances and Disease Registry (ATSDR). 1989d. *1,1-Dichloroethene*. United States Department of Health and Human Services. Public Health Service. December 1989.

Agency for Toxic Substances and Disease Registry (ATSDR). 1989e. *Trichloroethylene*. United States Department of Health and Human Services. Public Health Service. October 1989.

Agency for Toxic Substances and Disease Registry (ATSDR). 1990a. *Polycyclic Aromatic Hydrocarbons*. United States Department of Health and Human Services. Public Health Service. December 1990.

Agency for Toxic Substances and Disease Registry (ATSDR). 1990b. *Copper*. United States Department of Health and Human Services. Public Health Service. December 1990.

Agency for Toxic Substances and Disease Registry (ATSDR). 1990c. *Silver*. United States Department of Health and Human Services. Public Health Service. December 1990.

Agency for Toxic Substances and Disease Registry (ATSDR). 1990d. *Naphthalene and 2-methylnaphthalene*. United States Department of Health and Human Services. Public Health Service. December 1990.

Agency for Toxic Substances and Disease Registry (ATSDR). 1990e. *1,1,1-Trichloroethane*. United States Department of Health and Human Services. Public Health Service. December 1990.

Agency for Toxic Substances and Disease Registry (ATSDR). 1992a. *Manganese*. United States Department of Health and Human Services. Public Health Service. July 1992.

Agency for Toxic Substances and Disease Registry (ATSDR). 1992b. *2-Butanone*. United States Department of Health and Human Services. Public Health Service. July 1992.

Agency for Toxic Substances and Disease Registry (ATSDR). 1992c. *Aluminum*. United States Department of Health and Human Services. Public Health Service. July 1992.

Agency for Toxic Substances and Disease Registry (ATSDR). 1992d. *Cobalt*. United States Department of Health and Human Services. Public Health Service. July 1992.

Alumot, E., G. Poli, and A. Tomasi. 1976. *Tolerance and Acceptable Daily Intake of Chlorinated Fumigants in the Rat Diet*. Food Cosmet. Toxicol. 14:105-110.

Ambrose, A. M., P. S. Larson, and J. F. Borzelleca. 1976. *Long Term Toxicologic Assessment of Nickel in Rats and Dogs*. Journal Food Science Technology. 13:181-187.

Ammerman, C. B., K. R. Flick, S. L. Hansards, and S. H. Miller. 1973. *Toxicity of Certain Minerals to Domestic Animals: A Review*. National Feed Ingredients Association. Des Moines, IA.

Aughey, E., L. Grant, and B. L. Furrman. 1977. *The Effects of Oral Zinc Supplementation in the Mouse*. Journal Complete Pathology 87:1-14.

Azar, A., H. J. Trochimowicz, and M. E. Maxfield. 1973. *Review of Lead Studies in Animals Carried Out at Haskell Laboratory: Two-Year Feeding Study and Response to Hemorrhage Study*. In: Proceedings of the International Symposium on Health Aspects of Lead. pp. 199-210. D. Barth, A. Berlin, and R. Engel, eds. October 1972, Amsterdam, The Netherlands. Luxembourg, Commission of the European Communities.

Bard, C. C., J. J. Murphy, D. L. Stone, and C. J. Terhaar. 1976. *Silver in Photoprocessing Effluents*. Journal of the Water Pollution Control Federation 489(2):389-394.

Barnes, D. W., V. M. Sanders, and K. L. White, Jr. 1985. *Toxicology of trans-1,2-Dichloroethylene in the Mouse*. Drug Chem. Toxicol. 8:373-392.

Beijer, K., and A. Jernelov. 1979. *General Aspects and Specific Data on Ecological Effects of Metals*. Handbook on the Toxicology of Metals. pp. 197-209. L. Friberg (ed.). Elsevier/North Holland Biomedical Press, New York.

Birdsall, C. W., C. E. Grue, and A. Anderson. 1986. *Lead Concentrations in Bullfrog (Rana Catesbeiana) and Green Frog (R. Clamitans) Tadpoles Inhabiting Highway Drainages*. Environmental Pollution. 40A:223-247.

Bock, F.G., and M. Burnham. 1960. *The Effect of Experimental Conditions Upon the Concentration of Hydrocarbon in Mouse Skin After Cutaneous Application*. Cancer Res. 21:510-515.

Bodek, I., W. J. Lyman, W. F. Reehl, and D. H. Rosenblatt, eds. 1988. *Environmental Inorganic Chemistry: Properties, Processes and Estimation Methods*. pp. 4.2-1 through 4.3-13, Pergamon Press, Inc.

Bomhard, E., E. Loser, A. Dornemann, and B. Schilde. 1982. *Subchronic Oral Toxicity and Analytical Studies on Nickel Rutile Yellow and Chrome Rutile Yellow with Rats*. Toxicology Letters 14:189-194.

Boon, J. P., R.C.H. Oudejans, and J. C. Duinker. 1984. *Kinetics of Polychlorinated Biphenyl (PCB) Components in Juvenile Sole (Solea solea) in Relation to their Concentrations in Food and to Lipid Metabolism*. Comp. Biochem. Physiol. 79C:131-142.

Bradley, W. R., and W. G. Fredrick. 1941. *The Toxicity of Antimony*. Ind. Med. Ind. Hyg. Section 2:15-22.

Broderson, J. R., J. R. Lindsey, and J. E. Crawford. 1976. *The Role of Environmental Ammonia in Respiratory Mycoplasmosis of Rats*. American Journal of Pathology 85:115-130.

Brown, D., K. R. Butterworth, I. F. Gaunt, P. Grasso, and S. D. Gangolli. 1978. *Short-term Oral Toxicity Study of Diethyl Phthalate in the Rat*. Food Cosmet. Toxicol. 16:415-422.

Brown E., and W. Hewitt. 1984. *Dose-Response Relationships in Ketone-Induced Potentiation of Chloroform Hepato- and Nephrotoxicity*. Toxicol. Appl. Pharmacol. 76:437-454.

Buben, J. A., and E. J. O'Flaherty. 1985. *Delineation of the Role of Metabolism in the Hepatotoxicity of Trichloroethylene and Perchloroethylene: A Dose-Effect Study*. Toxicol. Appl. Pharmacol. 78:105-122.

Burton, G.A. 1992. ed. *Sediment Toxicity Assessment*. Lewis Publishers, Inc. Chelsea, MI.

Byron, W. R., G. W. Bierbower, and J. B. Brouwer. 1967. *Pathological Changes in Rats and Dogs from Two-year Feeding of Sodium Arsenite or Sodium Arsenate*. Toxicol. Appl. Pharmacol. 10:132-147.

Carmignani, M. P. Boxcolo, and P. Preziosi. 1988. *Cardiovascular Actions of Lead in Rats as Related to the Level of Chronic Exposure*. Arch. Toxicol. Suppl. 12:449-452.

Carpenter, C. P., C. S. Weil, and H. F. Smyth. 1953. *Chronic Oral Toxicity of DEHP for Rats, Guinea Pigs, and Dogs*. AMA Arch. Ind. Hyg. 8:219-226.

CH2M HILL. 1990. *Water Quality Modeling and Standard Default Values for the EPA Hazard Ranking System*. Denver, CO. (In review).

Clarkson, T. W. 1979. *Effects-General Principles Underlying the Toxic Action of Metals*. Handbook on Toxicology of Metals, pp. 99-117. ed. L. Gribert et al. Elsevier/North-Holland Biomedical Press, New York.

Cousins, R. J., A. K. Barber, and J. R. Trout. 1973. *Cadmium Toxicity in Growing Swine*. Journal of Nutrition 103:964-972.

DeGroot, A. P., V. J. Feron, and H. P. Til. 1973. *Short-Term Toxicity Studies on Some Salts and Oxides of Tin in Rats*. Food Cosmet. Toxicol. 11:19-30.

Deichman, W. B., W. E. MacDonald, and E. Bernal. 1963. *The Hemopoietic Toxicity of Benzene Vapors*. Toxicol. Appl. Pharmacol. 5:201-224.

Deutch-Wnzel, R. P., H. Brunce, O. Grimmer, G. Dettbarn, and J. Misfeld. 1983. *Experimental Studies in Rat Lungs on the Carcinogenicity and Dose-response Relationships of Eight Frequently Occurring Environmental Polycyclic Aromatic Hydrocarbons*. J. Natl. Cancer Inst. 71:539-544.

Doherty, P. C., R. M. Barlow, and K. W. Angus. 1969. *Spongy Changes in the Brains of Sheep Poisoned by Excess Dietary Copper*, Research in Veterinary Science 10:303-304.

Doyle, J. J., W. H. Pfander, S. E. Grebing, and J. O. Pierce II. 1974. *Cadmium Absorption and Cadmium Tissue Levels in Growing Lambs*. Journal of Nutrition 104:160-166.

Drinker, K. R., P. K. Thompson, and M. Marsh. 1927. *An Investigation of the Effect Upon Rats of Long-Continued Ingestion of Zinc Compounds, with Special Reference to the Relation of Zinc Excretion to Zinc Intak*. American Journal Physio. 81:284-306.

duPont deNemours and Company, Inc. 1982. *Subacute Inhalation Toxicity Study of Aniline in Rats*. OTS No. 878220240, Fiche No. 0215025, U.S. Environmental Protection Agency, Washington, DC.

Eben, A., W. Flucke, and F. Mihail. 1979. *Toxicological and Metabolic Studies of methyl n-butylketone, 2,5-hexanedione and 2,5-hexaneol in Male Rats*. Ecotoxicol. Environmental Safety 3:204-217.

Edens, F. W., and J. D. Garlich. 1983. *Lead Induced Egg Production Decrease in Leghorn and Japanese Quail*. Poultry Science 62:1757-1763.

Eisler, R. 1985. *Cadmium Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review*. United States Fish and Wildlife Service, Biological Report 85 (1.2). Washington, DC.

Eisler, R. 1986. *Polychlorinated Biphenyl Hazards to Fish, Wildlife and Invertebrates: A Synoptic Review*. U.S. Fish and Wildlife Service, Biological Report 85(1.7), U.S. Department of the Interior, Washington, DC.

Eisler, R. 1987a. *Mercury Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review*. United States Fish and Wildlife Service, Biological Report 85(1.10), U.S. Department of the Interior, Washington, DC.

Eisler, R. 1987b. *Polycyclic Aromatic Hydrocarbon Hazards to Fish, Wildlife and Invertebrates: A Synoptic Review*. U.S. Fish and Wildlife Service, Biological Report 85(1.11). U.S. Department of the Interior, Washington, DC.

Eisler, R. 1988a. *Arsenic Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review*. United States Fish and Wildlife Service. Biological Report 85(1.12). U.S. Department of the Interior, Washington, DC.

Eisler, R. 1988b. *Lead Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review*. United States Fish and Wildlife Service. Biological Report 85(1.14). U.S. Department of the Interior, Washington, DC.

Environmental Protection Agency (EPA). 1980a. *Ambient Water Quality Criteria Document for Benzene*. EPA 440/5-80-018. Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office (Cincinnati, OH) and Carcinogen Assessment Group (Washington, DC), and the Environmental Research Labs (Corvallis, OR) for the Office of Water Regulations and Standards, Washington, DC.

Environmental Protection Agency (EPA). 1980b. *Ambient Water Quality Criteria for Naphthalene*. EPA 440/5-80-059. Office of Water Regulations and Standards Criteria and Standards Division. Washington, DC.

Environmental Protection Agency (EPA) 1980c. *Ambient Water Quality Criteria for Polychlorinated Biphenyls*. EPA 440/5-80-068, Washington, DC.

Environmental Protection Agency (EPA) 1980d. *Ambient Water Quality Criteria for Cadmium*. EPA 440/55-80-025, Office of Water Regulations and Standards, Criteria and Standards Division, Washington, DC.

Environmental Protection Agency (EPA). 1980e. *Ambient Water Quality Criteria for Polynuclear Aromatic Hydrocarbons*. EPA 440/5-80-069. Washington, DC.

Environmental Protection Agency (EPA) 1984. *Ambient Water Quality Criteria for Cadmium*. EPA 440/5-84-032. Office of Water Regulations and Standards, Criteria and Standards Division, Washington, DC.

Environmental Protection Agency (EPA). 1986a. *Ninety-day Gavage Study in Albino Rats using Acetone*. Office of Solid Waste. Washington, DC.

Environmental Protection Agency (EPA). 1986b. *o,m,p-Cresol. 90-day Oral Subchronic Toxicity Studies in Rats*. Office of Solid Waste. Washington, DC.

Environmental Protection Agency (EPA). 1987. *o,m,p-Cresol. 90-day Oral Subchronic Neurotoxicity Study in Rats*. Office of Solid Waste. Washington, DC.

Environmental Protection Agency (EPA). 1988a. *Guidance for Conducting Remedial Investigations and Feasibility Studies under CERCLA*. Interim Final. Office of Emergency and Remedial Response. Washington, DC.

Environmental Protection Agency (EPA). 1988b. *Thirteen-week Mouse Oral Subchronic Toxicity Study*. Prepared by Toxicity Research Laboratories, Ltd., Muskegon, MI for the Office of Solid Waste. Washington, DC.

Environmental Protection Agency (EPA). 1989a. *Risk Assessment Guidance for Superfund Volume II: Environmental Evaluation Manual*. Interim Final. EPA/540/1-89/001. Office of Emergency and Remedial Response. Washington, DC, March 1989.

Environmental Protection Agency (EPA). 1989b. *Ecological Assessment of Hazardous Waste Sites: A Field and Laboratory Reference*. EPA/600/3-89/013. Office of Environmental Research-Laboratory. Corvallis, OR. EPA/540/1-89/001. Office of Emergency and Remedial Response. Washington, DC.

Environmental Protection Agency (EPA). 1989c. *Mouse Oral Subchronic Study with Acenaphthene*. Study conducted by Hazelton Laboratories America, Inc., for the Office of Solid Waste, Washington, DC.

Environmental Protection Agency (EPA) 1989d. *Mouse Oral Subchronic Toxicity Study*. Prepared by Toxicity Research Laboratories, LTD., Muskegon, MI for the Office of Solid Waste, Washington, DC.

Environmental Protection Agency (EPA). 1989e. *Ninety-day Gavage Study in Albino Mice Using 2,4-dimethylphenol*. Study No. 410-2831. Prepared by Dynamic Corporation, Rockville, MD, for the Office of Solid Waste and Emergency Response. Washington, DC.

Environmental Protection Agency (EPA). 1989f. *Subchronic Toxicity in Mice with Anthracene*. Final Report. Prepared by Hazelton Laboratories America, Inc., for the Office of Solid Waste, Washington DC.

Environmental Protection Agency (EPA). 1990a. *National Oil and Hazardous Substances Pollution Contingency Plan (NCP)* 40 CFR Part 300; 53 Federal Register 51394.

Environmental Protection Agency (EPA). 1990b. *Reducing Risk: Setting Priorities and Strategies for Environmental Protection*. Science Advisory Board SAB-EC-90-021, Washington, DC.

Environmental Protection Agency (EPA). 1991. *Summary Report on Issues in Ecological Risk Assessment*. EPA/625/3-91/018. Risk Assessment Forum. Washington, DC.

Environmental Protection Agency (EPA). 1992a. *Framework for Ecological Risk Assessment*. EPA/630/R-92/001. Risk Assessment Forum. Washington DC.

Environmental Protection Agency (EPA). 1992b. *Peer Review Workshop Report on a Framework for Ecological Risk Assessment*. EPA/625/3-91/022. Risk Assessment Forum. Washington, DC.

Environmental Protection Agency (EPA). 1992c. *Health Effects Assessment Summary Tables, FY-92 Annual*. Office of Research and Development. OERR 9200.6-303 (92-1 January).

Environmental Protection Agency (EPA). 1993. *Ambient Water Quality Criteria*. Electronic Bulletin Database. February 10, 1993.

Exon, J. H. and L. D. Koller. 1985. *Toxicity of 2-chlorophenol, 2,4-dichlorophenol and 2,4,6-trichlorophenol*. In: *Water Chlorination: Chemistry, Environmental Impact and Health Effects*. pp. 307-330. eds. Jolley et al. Lewis Publishers. Chelsea, MI.

Fitzhugh, O. G., A. A. Nelson, and E. P. Laug. 1950. *Chronic Oral Toxicities of Mercuri-Phenyl and Mercuric Salts*. Arch. Ind. Hyg. Occup. Med. 2:433-442.

Food and Drug Research Laboratories, Inc. (FDRL). 1972. *Teratologic Evaluation of FDA 71-37 (sodium benzoate)*. pp. 75-79.

Formigli, L., Scelsi, R., and P. Poggi. 1986. *Thallium-induced Testicular Toxicity in the Rat*. Environ. Res. 40:531-539.

Garten, C. T. 1980. *Ingestion of Soil by Hispid Cotton Rats, White Footed Mice and Eastern Chipmunks*. Journal of Mammalogy. 61:136-137.

Ginn, T. C. and R. A. Pastorok. 1992. *Assessment and Management of Contaminated Sediments in Puget Sound*. In: *Sediment Toxicity Assessment*, pp. 371-397. ed. G. Allen Burton, Jr. Lewis Publishers, Boca Raton, FL.

Gipp, W. F., W. G. Pond, F. A. Kallfelz, J. B. Tasker, D. R. Van Campen, L. Krook, and J. W. Visek. 1974. *Effect of Dietary Copper, Iron and Ascorbic Acid Levels on Hematology, Blood and Tissue Copper, Iron and Zinc Concentrations, and Copper and Iron Metabolism in Young Pigs*. Journal of Nutrition 104:532-541.

Gupta, B. N., J. G. Vos, J. A. Moore, J. G. Zinkl, and B. C. Bullock. 1973. *Pathologic Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin in Laboratory Animals*. Environmental Health Perspective 5:125-140.

Hardin, B. D., G. P. Bond, M. R. Sikor, F. D. Andrew, R. P. Beliles, and R. W. Nieheir. 1981. *Testing of Selected Work Place Chemicals for Teratogenic Potential*. Scand. J. Work Environmental Health 7:66-75.

Hecht, S. S., W. E. Bondinell, and D. Hoffmann. 1974. *Chrysene and methylchrysenes: Presence in Tobacco Smoke and Carcinogenicity*. J. Nat. Cancer Inst. 53:1121-1133.

Hejtmancik, M., A. C. Peters and J. D. Toft. 1987a. *The Chronic Study of Manganese Sulfate Monohydrate in F344 Rats*. Report to National Toxicology Program, Research Triangle Park, NC by Battelle's Columbus Laboratories, Columbus.

Hejtmancik, M., A. C. Peters and J. D. Toft. 1987b. *The Chronic Study of Manganese Sulfate Monohydrate in B6C3F1 Mice*. Report to National Toxicology Program, Research Triangle Park, NC by Battelle's Columbus Laboratories. Columbus.

Hewitt, W. R., H. Miyajima, and M. G. Cote. 1980. *Acute Alteration of Chloroform-Induced hepato-and nephrotoxicity by n-hexane, methyl n-butyl ketone and 2,5-hexanedione*. Toxicol. Appl. Pharmacology 53:230-248.

Heywood, R., R. J. Sortwell, and P. R. B. Noel. 1979. *Safety Evaluation of Toothpaste Containing Chloroform*. III. Long-Term Study in Beagle Dogs. Journal Environmental Pathology Toxicology 2:835-851.

Hoogland, J. L. 1981. *The Evolution of Coloniality in White-tailed and Black-tailed Prairie Dogs*. Ecology 62:252-272.

Horton, A. W. and G. M. Christian. 1974. *Cocarcinogenic Versus Incomplete Carcinogenic Activity Among Aromatic Hydrocarbons: Contrasts Between Chrysenes and Benzo[b]triphenylene*. J. Natl. Cancer Inst. 53:1017-1020.

HSDB. 1988. *Hazardous Substances Data Bank*.

Ignat'ev, A. D. 1965. *Experimental Information Contributing to a Hygienic Characterization of the Combined Effect Produced by Some Food Presentations*. vop. Pitan. 24:61-68.

Integrated Risk Information System (IRIS). 1992. *Chemical Files Office of Research and Development*. Cincinnati.

International Agency for Research on Cancer (IARC). 1980. *Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man*. Volume 23. Some Metals and Metallic Compounds. World Health Organization. Lyon, France.

Killinger, J. M., A. H. Basaran, and R. L. Persing. 1988. *Maximum Perinatal Dose Feed Study of Dibutylphthalate in Fischer 344 Rats*. Report to National Toxicology Program, Research Triangle Park, NC by Battelle, Columbus.

Klaunig, J. E., R. J. Ruch, and M. A. Pereira. 1986. *Carcinogenicity of Chlorinated Methane and Ethane Compounds Administered in Drinking Water to Mice*. Environmental Health Perspective 69:89-95.

Klein, M. 1963. *Susceptibility of Strain B6AF/J Hybrid Infant Mice to Tumorigenesis with 1,2-benzanthracene, Deoxycholic acid and 3-methylcholanthrene*. Cancer Res. 23:1701-1707.

Kopp, S. J., T. Glonek, and H. M. Perry. 1982. *Cardiovascular Actions of Cadmium at Environmental Exposure Levels*. Science 217:837-839.

Kowalska, M. 1988. *The Effect of Vanadium on Lung Collagen Content and Composition in Two Successive Generations of Rats*. Toxicology Letter 41:203-208.

Krasovskii, G. N., and S. A. Fridlyand. 1971. *Experimental Data for the Validation of the Maximum Permissible Concentration of Cobalt in Water Bodies*. Hygiene Sanitary 36:277-279.

Lai, J. C. K., T. K. C. Leung, and L. Lim. 1981. *Brain Regional Distribution of Glutamic Acid Decarboxylase, Choline Acetyl Transferase, and Acetylcholinesterase in the Rat: Effects of Chronic Manganese Chloride Administration After Two Years*. Journal of Neurochemistry 36:1443-1448.

Lai, J. C. K., T. K. C. Leung, J. F. Guest, A. N. Davison and L. Lim. 1982. *The Effects of Chronic Manganese Chloride Treatment Expressed as Age Dependent, Transient Changes in Rat Brain Synaptosomal Uptake of Amines*. Journal of Neurochemistry. 38:844-847.

Laug, E.P., A.A. Nelson, O.G. Fitzhugh, and F.M. Kunze. 1950. *Liver Cell Alterations and DDT Storage in the Fat of the Rat Induced by Dietary Levels of 1-50 ppm DDT*. Pharmacol. Exp. Therap. 98:268-273.

Lee, S.D., and L. Grant (eds). 1981. *Health and Ecological Assessment of Polynuclear Aromatic Hydrocarbons*. Pathotex Publishing, Park Forest South, Illinois.

Leung, T. K. C., J. C. K. Lai, and L. Lim. 1981. *The Regional Distribution of Monoamine Oxidase Activities Towards Different Substrates: Effects in Rat Brain of Chronic Administration of Manganese Chloride and of Aging*. Journal of Neurochemistry 36:2037-2043.

Lewis, P. K., W. G. Hoekstra, and R. H. Grummer. 1957. *Restricted Calcium Feeding Versus Zinc Supplementation for the Control of Parakeratosis in Swine*. Journal of Animal Science 16:578-588.

Linder, G. 1988. *NSI/Environmental Sciences*. Corvallis, OR. Personal communication. November 30, 1988.

Mackenzie, K. M., and D. M. Angevine. 1981. *Infertility in Mice Exposed in utero to benzo(a)pyrene*. *Bio. Reprod.* 24:183-191.

Maltoni, C., G. Cotti, and V. Patella. 1986. *Results of Long-Term Carcinogenicity Bioassays on Sprague-Dawley Rats of Methyl Chloroform, Administered by Ingestion*. *ACTA. Oncol.* 7:101-117.

Massie, H. R., and V. R. Aiello. 1984. *Excessive Intake of Copper: Influence on Longevity and Cadmium Accumulation in Mice*. *Mech. Aging Dev.* 26:195-203.

Microbiological Associates. 1986. *Subchronic Toxicity of Methyl Isobutyl Ketone in Sprague-Dawley Rats*. Preliminary Report. Research Triangle Park. North Carolina. Research Triangle Park Institute (Study No. 5221. 04).

Ming-Dean Luh, R. A., and D. E. Henley. 1973. *Arsenic Analysis and Toxicity—A Review*. *The Science of the Total Environment*. 2:1-12.

Moore, J. A., B. N. Gupta, and J. G. Vos. 1976. *Toxicity of 2,3,7,8-tetrachlorodibenzofuran—Preliminary Results*. In: *Proceedings from the National Conference on Polychlorinated Biphenyls*. November 19-21, 1975, Chicago, Illinois. Research Triangle Park. North Carolina. Research Triangle Park Institute.

Murphy, R. L., S. Lal, and D. K. Saxena. 1981. *Effect of Manganese and Copper Interaction on Behavior and Biogenic Amines in Rats fed a 10 Percent Casein Diet*. *Chemical Biological Interaction* 37:299-308.

National Academy of Science (NAS). 1977. *Arsenic*. National Academy of Science, Washington DC.

National Academy of Sciences (NAS). 1979. *Polychlorinated Biphenyls*. National Academy of Science, Washington, DC.

National Cancer Institute (NCI). 1976. *Carcinogenesis Bioassay of Trichloroethylene*. Case No. 79-01-6, NCI-CG-TR-2. Department of Health, Education, and Welfare. National Institutes of Health Publication 76-802.

National Cancer Institute (NCI). 1978. *Bioassay of Technical Grade 1,2-dichloroethane for Possible Carcinogenicity*. Bethesda, MD: NCI-CG-TR-55.

National Cancer Institute (NCI). 1979. *Bioassay of p-chloraniline for Possible Carcinogenicity*. NCI Carcinogenesis Technical Report Serial No. 189. NTIS PB 295896.

National Coffee Association (NCA). 1982. *Twenty-four Month Chronic Toxicity and Oncogenicity Study of Methylene Chloride in Rats*. Final Report. Prepared by Hazleton Laboratories, America Inc., Vienna, VA.

National Toxicology Program (NTP). 1983. *Teratologic Evaluation of Phenol in CD Rats and Mice*. Report prepared by Research Triangle Institute. Research Triangle Park, NC. NTIS PB83-247726. Gov. Rep. Announce Index. 83:6247.

National Toxicology Program (NTP). 1986. *Toxicology and Carcinogenesis Studies of Xylenes (mixed) in F344/N Rats and B6C3F1 Mice (Gavage Studies)*. NTP TR 327 NIH PB No. 86-2583.

National Toxicology Program (NTP). 1989. *Toxicology and Carcinogenesis Studies of Toluene in F344/N Rats and B6C3F1 Mice*. Technical Report Series No. 371. Research Triangle Park, NC.

Neff, J. M. 1979. *Polycyclic Aromatic Hydrocarbons in the Aquatic Environment*. Applied Science Publ. Ltd., London.

Neff, J. M. 1985. *Polycyclic Aromatic Hydrocarbons*. pp. 416-454. In: *Fundamentals of Aquatic Toxicology*, eds. G. M. Rand and S. R. Petrocelli. Hemisphere Publishing Corporation, New York.

Nelson, A. A., O. G. Fitzhugh, and H. O. Calvery. 1943. *Liver Tumors Following Cirrhosis Caused by Selenium in Rats*. *Cancer Research* 3:230-236.

Nikonorow, M., H. Mazur, and H. Piekacz. 1973. *Effect of Orally Administered Plasticizers and Polyvinyl Chloride Stabilizers in the Rat*. *Toxicol. Appl. Pharmacol.* 26:253-259.

Nomiyama, K., C. Sato, and A. Yamamoto. 1973. *Early Signs of Cadmium Intoxication in Rabbits*. *Toxicology and Applied Pharmacology* 24:624-635.

O'Connor, D. J., and S. W. Nielson. 1980. *Environmental Survey of Methylmercury Levels in Wild Mink and Otter from the Northeastern United States and Experimental Pathology of Methylmercurialism in the Otter*. pp. 1728-1745. In: *Proceedings Worldwide Furbearers Conference*, eds. J. A. Chapman and D. Pursley.

- Olcott, C. T. 1948. *Experimental Argyrosis. IV. Morphologic Changes in the Experimental Animal*. American Journal of Pathology 24:813-833.
- Ondreicka, R. E. Ginter, and J. Kortus. 1966. *Chronic Toxicity of Aluminum in Rats and Mice and its Effects on Phosphorus Metabolism*. British Journal of Industrial Medicine 23:305-312.
- Ott, E. A., W. H. Smith, R. B. Harrington, and W. M. Beeson. 1966a. *Zinc Toxicity in Ruminants. I. Effects of High Levels of Dietary Zinc on Gains, Feed Consumption and Feed Efficiency of Lambs*. Journal of Animal Science 25:414-418.
- Ott, E. A., W. H. Smith, R. B. Harrington, and W. M. Beeson. 1966b. *Zinc Toxicity in Ruminants. II. Effects of High Levels of Dietary Zinc on Gains, Feed Consumption and Feed Efficiency of Beef Cattle*. Journal of Animal Science 25:419-423.
- Ottoboni, A. 1969. *Effect of DDT on Reproduction in the Rat*. Toxicol. Appl. Pharmacol. 14:74-81.
- Palmer, A. K., A. E. Street, and J. C. Roe. 1979. *Safety Evaluation of Toothpaste Containing Chloroform. II. Long-term Studies in Rats*. Journal of Environmental Pathology Toxicology 2:821-833.
- Paternain, J. L., J. L. Domingo, and J. M. Llobet. 1988. *Embryotoxic and Teratogenic Effects of Aluminum Nitrate in Rats upon Oral Administration*. Teratology 38:253-257.
- Perry, H. M., S. J. Kopp, M. W. Erlanger and E. F. Perry. 1983. *Cardiovascular Effects of Chronic Barium Ingestion*. In: Proceedings of the University of Missouri 17th Annual Conference on Trace Substances in Environmental Health, ed. D. D. Hemphill. University of Missouri Press, Columbia, MO.
- Perry, H. M., M. W. Erlanger, and E. F. Perry. 1988. *Increase in the Blood Pressure of Rats Chronically Fed Low Levels of Lead*. Environmental Health Perspective 78:107-111.
- Pershagan, G., B. Lind, and N. E. Bjorkund. 1982. *Long Retention and Toxicity of Some Inorganic Arsenic Compounds*. Environmental Research 29:425-434.
- Pershagan, G., G. Nordberg, B. Lind, and N. E. Bjorkund. 1984. *Carcinomas of the Respiratory Tract in Hamsters Given Arsenic Trioxide and/or Benzo(a)pyrene by Pulmonary Route*. Environmental Research 34:227-241.

Philbrick, D. J., J. B. Hopkins, D. C. Hill, J. C. Alexander, and R. G. Thomson. 1979. *Effects of Prolonged Cyanide and Thiocyanate Feeding in Rats*. Journal of Toxicology Environmental Health 5:579-592.

Plasterer, P., I. Schmeltz, J. Tosk, and E. L. Wynder. 1985. *Bioassays of Naphthalene and Alkyl naphthalenes*. Carcinog. 3:47-60.

Quast, J. F., C. G. Humiston, and C. E. Wade. 1983. *A Chronic Toxicity and Oncogenicity Study in Rats and Subchronic Toxicity Study in Dogs on Ingested Vinylidene Chloride*. Fundamental Appl. Toxicology 3:55-62.

Ralston, W., R. Hilderbrand, and D. Uddin. 1985. *Potentiation of 2,5-hexanedione Neurotoxicity by Methyl Ethyl Ketone*. Toxicol. App. Pharmacol. 81:319-327.

Rampy, L. W., J. F. Quast, and C. G. Huiston. 1977. *Interim Results of Two-year Toxicological Studies in Rats of Vinylidene Chloride Incorporated in the Drinking Water or Administered by Repeated Inhalation*. Environmental Health Perspective 21:33-43.

Rungby, J., and G. Danscher. 1984. *Hypoactivity in Silver Exposed Mice*. Acta. Pharmacol. Toxicol. 55:398-401.

Sax, I. 1984. *Dangerous Properties of Industrial Materials*. VanNostrand Reinhold Company. New York.

Sax, N. I. 1992. *Dangerous Properties of Industrial Materials, Sixth Edition*. Van Nostrand Reinhold Company, New York.

Schmahl, D. 1955. *Examination of the Carcinogenic Action of Naphthalene and Anthracene in Rats*. Z. Krebsforsch. 60:697-710.

Schroeder, H. A., J. J. Balassa, and W. H. Vinton, Jr. 1965. *Chromium, Cadmium and Lead in Rats: Effects on Life Span Tumors and Tissue Levels*. Journal of Nutrition 86:51-66.

Schroeder, H. A., I. H. Tipton, and A. P. Nason. 1972. *Trace Metals in Man: Strontium, and Barium*. J. Chronic. Dis. 25:491-517.

Schroeder, H. A., M. Mitchener, and A. P. Nason. 1970. *Zirconium, Niobium, Antimony Vanadium and Lead in Rats Life Term Studies*. Journal of Nutrition. 100:59-68.

Schroeder, H. A., and M. Mitchener. 1975a. *Life-Term Effects of Mercury, Methyl Mercury, and Nine Other Trace Elements on Mice*. Journal of Nutrition 105:452-458.

- Schroeder, H. A., and M. Mitchener. 1975b. *Life-Term Studies in Rats: Effects of Aluminum, Barium, Beryllium, and Tungsten*. Journal of Nutrition 105:421-427.
- Shaikh, Z. A., K. M. Harnett, and S. A. Perlin. 1989. *Chronic Cadmium Intake Results in Dose-Related Excretion of Metallothionein in Urine*. Experientia 45:146-148.
- Shaw, G.R., and D.W. Connell. 1982. *Factors Influencing Polychlorinated biphenyls in Organisms from an Estuarine Ecosystem*. Aust. J. Mar. Freshwater Res. 33:1057-1070.
- Sims, R.C., and R. Overcash. 1983. *Fate of Polynuclear Aromatic Compounds in Soil-Plant Systems*. Residue Rev. 88:1-68.
- Stoltz, M.L., M.A. Stedham, and L.K. Brown. 1986. *Subchronic (90-day) Toxicity of Thallium (I) Sulfate in Sprague-Dawley Rats*. Prepared by Midwest Research Institute, Kansas City, MO, for the U.S. Environmental Protection Agency, Office of Solid Waste, Washington, DC.
- Suess, M. J. 1976. *The Environmental Load and Cycle of Polycyclic Aromatic Hydrocarbons*. Sci. Total Environ. 6:239-250.
- Suttle, N. F., and C. F. Mills. 1966. *Studies of the Toxicity of Copper to Pigs: 1. Effect of Oral Supplements of Zinc and Iron Salts on the Development of Copper Toxicosis: 2. Effect of Protein Source and Other Dietary Components on the Response to High and Moderate Intakes of Copper*. British Journal of Nutrition 20:135-149.
- Sutton, W. R., and V. E. Nelson. 1937. *Studies on Zinc*. Proceedings from the Society of Experimental Biology and Medicine 36:211-214.
- Syed, I. B., and F. Hosain. 1972. *Toxicity of Lesser-known Metals to Laboratory Animals*. 22:150-152.
- Tardiff, R. G., M. Robinson, and N. S. Ulmer. 1980. *Subchronic Oral Toxicity of BaCl₂ in Rats*. Journal of Environmental Pathology and Toxicology 4:267-275.
- Tile, H. P., H. R. Immel, and V. J. Feron. 1983. *Lifespan Oral Carcinogenicity Study of Vinyl Chloride in Rats*. Final Report. Report No. V 93. 285/291-99. Civo Institutes, TNO.
- Tomatis, L., V. Turusov, and N. Day. 1972. *The Effect of Long-term Exposure to DDT on CF-1 Mice*. Int. J. Cancer 10:489-506.
- Turkowski, F. J. 1975. *Dietary Adaptability of the Desert Cottontail*. Journal of Wildlife Management 39:748-756.

United States Fish and Wildlife Service (USFWS). 1984. *Handbook of Toxicity of Pesticides to Wildlife*. Publication 153. U.S. Department of the Interior, Fish and Wildlife Service Resource.

Van Duuren, B.L., L. Langseth, B.M. Goldschmidt and L. Orris. 1967. *Carcinogenicity of Epoxides, Lactones, and Peroxycompounds*. VI. Structure and carcinogenic activity. *J. Nat. Cancer Inst.* 39:1217-1228.

Venugopol, B. and T. D. Luckey. 1978. *Metal Toxicity in Mammals. 2: Chemical Toxicity of Metals and Metalloids*. Plenum Press, New York.

Walker, F. 1971. *Experimental Argyria: A Model for Basement Membrane Studies*. *British Journal of Experimental Pathology* 52:589-593.

Walters, M., and F. Roe. 1965. *A Study of Effects of Zinc and Tin Administered Orally to Mice Over a Prolonged Period*. *Food Cosmet. Toxicology* 3:221-276.

Weir, R. J., Jr., and R. S. Fisher. 1972. *Toxicologic Studies on Borax and Boric Acid*. *Toxicol. Appl. Pharmacol.* 23:351-364.

Wolf, J., R. Esher, and K. Ronbinson. 1979. *Lethal and Reproductive Effects of Dietary Mirex and DDT on Old-field Mice *Peromyscus polionotus**. *Bull. Environ. Contam. Toxicol.* 21:397-402.

Wolf, M. A., V. K. Rowe, D. D. McCollister, R. L. Hollinsworthm and F. Oyen. 1956. *Toxicological Studies of Certain Alkylated Benzenes and Benzene*. *Arch. Ind. Health.* 14:387-398.

World Health Organization (WHO). 1980. *Tin and Organotin Compounds: A Preliminary Review*. Environmental Health Criteria 15. World Health Organization. Finland.

World Health Organization (WHO). 1989. *Environmental Health Criteria 85: Lead-Environmental Aspects*. World Health Organization. Geneva.

World Health Organization (WHO). 1989. *Polychlorinated dibenzo-para-dioxins and dibenzofurans*. Environmental Health Criteria 88. World Health Organization. Geneva.

Wren, C. D. 1986. *A Review of Metal Accumulation and Toxicity in Wild Mammals. I. Mercury*. *Environmental Research* 40:210-244.

Wynder, E. L., and D. Hoffman. 1959. *A Study of Tobacco Carcinogenesis. VII. The Role of Higher Polycyclic Hydrocarbons*. *Cancer* 12:1079-1086.

Wynder, E. L., and D. Hoffmann. 1959. *The Carcinogenicity of Benofluoranthenes*. Cancer 12:1194-1199.

Zahid, Z. R., Z. S. Al-Hakkak, and A. H. H. Kaahim. 1990. *Comparative Effects of Trivalent and Hexavalent Chromium on Spermatogenesis of the Mouse*. Toxicology Environment Chem. 25:131-136.

**DATE
FILMED**

7 / 19 / 93