DE/PC/91289--TL

APR 26 1993

Technical Report

December 1, 1992 through February 28, 1993

Contract No.: DE FG-22-91PC-91289

Project Title: Characterization of the Organic-Sulfur-Degrading Enzymes

Principal Investigator: Nancy W.Y. Ho
LORRE
Purdue University
1295 Potter Building
West Lafayette, IN 47907-1295

RECEIVED
JUN 14 1993
OSTI

OBJECTIVES

The immediate objective of this project is to characterize and purify the enzymes involved in degrading organic sulfur in coal from two well characterized organic sulfur degrading strains, IGTS8 and K3B. We believe that characterization and purification of these enzymes may provide valuable information that will lead to developing or isolating better strains for desulfurization of coal. Our original plan was first to characterize these enzymes, followed by purification of these enzymes, particularly the IGTS8 enzymes and cloning and overexpressing the genes encoding these enzymes in IGTS8 or other better organisms.

Concurrently we have another project engaged in the development of gene cloning systems for microorganisms that can better endure the harsh conditions of coal desulfurization. With the development of these gene cloning systems, microorganisms will be able to serve as the surrogate hosts to overproduce the desulfurization enzymes and create improved systems for coal desulfurization.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ya

RD

4-4
1

Nevertheless, our recent results imply that the IGTS8 enzymes are firmly attached to the cell wall, most likely that they are covalently bonded to the cell wall. These results have made us change our overall approach in solving the coal desulfurization problem. We believe that for coal desulfurization, it is better to have a microorganism that can secrete its desulfurization enzymes into the medium. The enzymes can penetrate into coal hundreds of times better than a bacterial cell. Thus we have changed the priority of this project. We believe that the isolation of a mutant of IGTS8 which can release the desulfurization enzymes into the cultural medium will be able to develop a much improved microbial process for coal desulfurization. Furthermore, the isolation of such a mutant will be able to shed some light on how the wild type IGTS8 desulfurization enzyme(s) are attached to the cell wall and make the purification of these enzymes much easier to accomplish.

Research Performed

In this period, on the one hand we tried to establish the best procedure for the screening of the IGTS8 mutants that can secrete their desulfurization enzymes in the medium. On the other hand, we used the already established method to disrupt the cell wall of IGTS8. It is possible that some of the intact desulfurization enzymes might become soluble while the cell wall is disrupted.

The methods we used include (1) shaking the cells with glass beads in the presence or absence of polyethylene glycol, (2) treating the cells with lysozyme in the presence or absence of polyethylene glycol, and (3) suspending the cells in water or in very low salt

buffer. We found that each of these methods can release a small amount of the intact desulfurization activity. Nevertheless, none of them are totally reliable.

Proper concentration of polyethylene glycol, sucrose, and sorbitol can all prevent protoplasts of any cell to lyse in the absence of its cell wall. However, we found that only polyethylene glycol is suitable to be used for keeping IGTS8 cells intact in the absence of cell wall. This is because the other two reagents will react with Gibbs reagent and produce blue colored substance. Since Gibbs' test is the easiest method to assay whether IGTS8 produces enzyme that can degrade DBT, this makes it impossible to use sucrose or sorbitol to stabilize the protoplasts of IGTS8.

We also tried to analyze the proteins released by these methods. We found that they are similar to those we obtained earlier as shown in Figure 1.

During this period we also analyzed the products produced by IGTS8 from degrading DBT. According to previous researchers (1), the major product should be 2-hydroxybiphenol. However, analyzing the products present in the cultural supernatant by HPLC, we found that 2-hydroxybiphenol was not a major product. There was a major product, as shown in Figure 2, with retention time far less than any of the possible products proposed by other researchers (1). However, the culture supernatant turned blue after treating it with the Gibbs reagent (2) which indicate the presence of a phenolic product in the cultural supernatant. Among the proposed possible degradation products from DBT only 2-hydroxybiphenol reacts with Gibbs' reagent and produces a blue colored product. Other researchers (1) relied heavily on Gibbs' test to analyze the products produced by IGTS8 degradation of DBT. This might be the reason that they concluded 2-

hydroxybiphenol to be the major product from IGTS8 degradation of DBT. According to our results, it is possible that 2-hydroxybiphenol has been further degraded to produce other products including one that can react with the Gibbs' reagent to form the characteristic blue color. It seems highly desirable, to use other sophisticated techniques to further characterize the degradation product(s) resulted from IGTS8 digestion of DBT.

References

1. Kilbane, J. J., "Microbial removal of organic sulfur from coal and petroleum," *Proceedings: 1991 Second International Symposium on the Biological Processing of Coal*, Electric Power Research Institute, Palo Alto, CA, pp. 5-3--5-17 (1991).
2. "For identification of 4-unsubstituted and 4-alkylphenols," *Anal. Chem.*, 56, 813 (1984).

Figure 1 Polyacrylamide gel electrophoresis analysis of proteins released from IGTS8 during various cell wall disruption treatments.

Lane 1: Protein molecular weight standards.

Lane 2: Proteins released from the cell surface of IGTS8.

Flow rate = 1.0 ml/min.
Mobile = 45% CH₃CN
UV = 254

CH. 1 C.5 2.50 ATT 8 OFFS 10 09/02/92 15:03

- Mixture in 45% CH₃CN
- ①. Benzoic Acid
 - ②. DBT sulfoxide
 - ③. DBT sulfone
 - ④. O,O'-Biphenol (lower layer)
 - ⑤. DBT

09/02/92 15:03

D-2500

METHOD: JING LU TAG: 27 CH: 1

FILE: 1 CALC-METHOD: EXT-STD TABLE: 8 CONC: AREA

NAME

NO.	RT	AREA	HEIGHT	6/DL	BC
1	1.95	11638	368	*****	BU
2	2.88	10830684	761227	*****	UU
3	3.46	3403	790	3403.000	TBB
4	3.76	2998	353	2998.000	TBB
5	4.35	4396	445	4396.000	TBB
6	4.82	2339520	215770	*****	UU
7	6.13	1134694	81048	*****	UB
8	8.32	10106496	524694	*****	BB
9	30.87	1269591	21355	*****	BB

Figure 2A. HPLC analysis of standards of DBT (dibenzothiophene). DBT sulfoxide, DBT-sulfone, 2-hydroxybiphenol, and benzoic acid. Benzoic acid was not one of the proposed degradation products from DBT by IGTS8 but the other three were

Retention time for benzoic acid: 2.89
DBT sulfoxide: 4.82
DBT sulfone: 6.13
2-hydroxybiphenol: 8.32
DBT: 30.87.

DBT with 16TS8 (9/15)

up layer

Figure 2B

DBT with 16TS8 (9/25)

low layer

Figure 2C.

Figure 2B HPLC analysis of organic chemicals present in culture supernatant of IGTS8, 20 ml of culture supernatant of IGTS8, cultured for 48 to 72 hr with cell density ~ 1.00 OD₆₀₀ mμ, was concentrated to dry and extracted with 1 ml 45% CH₃CN. It separated into two layers. Top layer with a volume ~ 0.2 ml was analyzed by HPLC with the same solvent system used for the standards. The results are shown in Figure 2B. The bottom layer with a volume ~ 0.8 ml was also analyzed by HPLC and the results are shown in Figure 2C.

Retention time for an unknown compound:	2.66-3.00
DBT sulfoxide:	4.16
DBT sulfone:	5.92
2-hydroxybiphenol:	---
DBT:	17.13

END

DATE
FILMED

7 / 13 / 93