AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

DOE Continuation Proposal

Coal Liquefaction Model Compounds

Joseph J. Gajewski and Kevin E. Gilbert, Co-Principal Investigators

Progress Report/Proposals:

Tetralin Pyrolysis-

Tetralin has been pyrolyzed at low concentrations (1-3 torr) in a well-conditioned static reactor at 450 degrees C. Hydrogen loss products, dehydronaphthalene and naphthalene, account for 75% ; C2 loss products, styrene and ethylbenzene, account for 12%; and neophyl radical rearrangement products, methylindenes and indanes, account for 12%. Tetramethylbutane initiates the reaction, and the product distribution is unchanged. In the absence of initiator, the rate of reaction is approximately first order in tetralin. The rate is roughly 0.5 order in initiator. In the liquid phase at neat concentrations, the reaction is still first order but is roughly three times faster but gives mostly 1-methylindane with the remainder being dehydrogenation product.

This data stands in contrast to the report of Bergman and Moore who performed pulsed laser pyrolysis of the material and Tsang who did shock tube pyrolysis and kinetics. Both groups found retro Diels-Alder product, benzocyclobutene. Tsang reported initiation studies with tetramethylbutane and reported many of the same products found here. Trahanovsky reported that the flash vacuum pyrolysis of tetralin lead to some of the same products reported by Bergman and Moore, but also formed were substantial amounts of dehydrogenation product. The liquid phase pyrolysis of tetralin has long been known to result in ring contraction to indane products.

In effect, our data provides the bridge between all previous studies at low and high concentrations, and

MASTER

radical chain mechanisms are clearly involved in dehydrogenation and ring contraction. What is remarkable is the first order kinetics in the absence of initiators. No mechanistic scheme considered by us thus far can rationalize the kinetics, particularly when the initiator is unknown.

Proposal. . . .

Further work in this area can only focus on the kinetics in the presence of external initiators. Unknown initiators in carefully purified tetralin obscured any attempt to explain the kinetics or the details of mechanism. Thus it should be possible to examine the kinetics and product distribution in the presence of tetramethylbutane in the gas phase and in solution. Studies such as this were performed in a preliminary way previously, but rigorous attempts to determine the reaction orders at low conversions have yet to be performed.

Chroman Pyrolysis—

Gas phase pyrolysis of chroman gives *o*-cresol, benzofuran, and styrene in a 4:2:1 ratio at 413 degrees C. In the presence of 2-butene, a moderately stereospecific 4+2 reaction occurs with the likely intermediate, *o*-benzoquinone methide. The addition of 2-butene increases the amount of reaction and decreases the extent of formation of benzofuran and styrene. Addition of hydrogen gas increases the amount of hydrodealkoxylation product, styrene, and gives more of its reduction product, ethylbenzene, at the expense of benzofuran. Only a slight increase in the amount of *o*-cresol was observed upon addition of hydrogen gas. No dehydrochroman was observed in the reactions, and pyrolysis of 4H- and 2-H-benzopyran gives no *o*-cresol, but instead gives styrene, benzofuran, *i*-indanone, and some chroman. The 2-butene trapping experiments suggest that the *o*-benzoquinone methide is formed reversibly. And the presence of 2-butene retards formation of the radical chain products. This work has been submitted to the Journal of Organic Chemistry.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Solvent Effects on Reactions–

A solvent rate effect equation which use four physical properties of the solvent for parameters has been applied with good success to solvolysis reactions, solvent Et values, Claisen Rearrangements, Diels–Alder reactions, endo–exo ratios in Diels–Alder reactions, azo–Cope rearrangements, and enol–ketone equilibria of the dimedone and pyridone variety. The coefficients of the parameters, which are the Kirkwood–Onsager function, the solvent cohesive energy density, the specific solvation of chloride ion, and the specific solvation of potassium ion, lead to differences in dipole moments, in volume differences between initial and final states, and in the extent of specific solvation of anionic and cationic sites relative to the solvation of chloride ion and potassium ion, respectively. This work has been published in the Journal of Organic Chemistry 1992, 57, 5500.

Proposal

Efforts are continuing to better parameterize the specific solvation of chloride and potassium ion by excluding the dielectric and solvent cohesive energy density. Examination of the previously analyzed reactions is being pursued. Nucleophilic displacement reactions of the SN2 variety are also be examined with inclusion of a solvent polarizability term. The importance of solvent polarizability is being examined in general. If anything, this work suggests that efforts to characterize polar solvent effects by a continuum dielectric model should not be successful. A cohesive energy term and specific solvation terms in the form

of hydrogen bond donors and acceptors are necessary.

Pi Conjugated Biradicals–

Previous, published work in our laboratory demonstrated that the cross-conjugated biradical *m*-xylene could be generated by addition of a 2,2-dimethylvinylidene to 6,6-dimethylfulvene. The product that resulted was octamethyl-[2,2]-*meta*-cyclophane, the dimer of the biradical. Interestingly, no CIDNP signals were observed that would suggest a triplet state was formed and reacted despite the prediction that such biradicals should be triplets. The prediction that these biradicals are triplets explained our concern for these species. Magnetism is a direct result of unpaired electrons in molecules. So these biradicals may be organic magnets.

Recently, in an effort to prevent the *meta*-xylene from dimerizing, the 2,2-dimethylvinylidene was added to 6-methyl-6-*tert*-butylfulvene. The product distribution is still being characterized.

Proposal

We hope to not only characterize the product distribution from the reaction just described, but also study the effect of oxygen on the original reaction which generated the octamethyl [2,2]-*meta*-cyclophane.

Oxygen trapping of triplets states has become a standard protocol for characterization of triplet states.

We also hope to generate higher multiplet states based on *meta*-xylene. For instance, [4]calixarene,

if converted to the tetraaradical, is predicted to be a quintet which is much more stable than the corresponding singlet state. Further, the materials might stack appropriately without intermolecular bonding to give an organic magnet or perhaps a conductor. Generation of sterically congested radical sites should decrease the extent of intermolecular interactions.

Molecular Mechanics of Organometallic systems–

We have developed a molecular mechanics force field based on Allinger's MM2 program with VESCF pi calculation routines from MM1. The initial approach focused on parameters for small unsaturated molecules.

Here it was found that acetylene bending constants were much too large in MM2 to account for the geometry and relatively small strain energy of cyclooctyne. It was further found that the out of plane bending constants for sp² carbon were much too small to account for the very large strain of pyramidalized olefins. This latter concern impacts on the strain energy of molecules like C₆₀ where the 600+ kcal/mol of strain comes substantially from this factor. With the current parameterization, the structure and stability of many of the model compounds and C₆₀ itself are reasonably reproduced in a program that runs quickly on a 386 or 486 Intel-based PC.

One of the early concerns was the geometry and relative stabilities of organometallic compounds. So an attempt was made to develop a force field which was general to all metals, ligands, and geometries. The force field is based on a Urey–Bradley force field at the metal (the steric size and electrostatics of the ligands determine the geometry at the metal, i.e. no bending constants at the metal) and 25 iron–ligand

distances are used as parameters which are adjusted by the covalent radius of the metal regardless of oxidation state. The stretching force constants are set to 2. mdyne/angstrom**2 except for metal CO bonds whose force constants are 4. mdyne/angstrom**2. Pi coordinated ligands are treated like a van der Waals attraction except that the attractive potential is proportional to $1/r^{**2}$, and the optimum distance is also proportional to the covalent radius of the metal. Further the well-depth is roughly 12 kcal/mol and has an off-p orbital axis correction factor of $1-(1-\cos^{**2})$ which is a relatively "soft" angular dependence. Initial studies revealed that the model worked reasonably well for coordinately saturated metals, but for coordinately unsaturated metals, the pi ligand-metal distance had to be increased by .15 angstroms and ligands with lone pairs of electrons had to have their distances to the metal decreased by .1 angstroms. Recent evaluation of the model using 276 structures from the Cambridge Crystallographic Database indicated a 0.022 angstrom mean deviation between calculated and experimental bond distances, however, the standard deviation is 0.178 angstroms.

Proposal

We have performed a Simplex optimization of the 25 metal-ligand distances and the covalent radii of the metals represented in the data base in order to reduce the standard deviations - all keeping the force constants the same as previously. In the course of this it also became clear that ligands like CO, carbene, carbyne, oxide, and nitride, required separate atom designations to distinguish them from groups which would use the atom types 4(sp C), 30(C+), 4(sp C), 7(=O), and 10(sp N), respectively. Further, the metal-phosphine force constant was too high to account for bond distance changed with phosphine having large cone angles.

Even more recently, JJG found, while visiting the laboratories of Prof. R.G. Bergman at Berkeley, that the pi-coordination potential was not deep or steep enough to account for the geometries and relative stabilities of some iridium compounds. Thus the attractive term has become proportional to $1/r^{**6}$ with a deeper well.

Other changes incorporated recently include a substantial reduction (to 0.0) in out of plane bending constants for pi ligands when attached to metals. Further, the two-fold barrier term in the torsion potential

for bonded pi ligands has been reduced from 15.0 kcal/mol to 3.9 kcal/mol. These changes allow pi ligands to undergo reasonable distortion in and out of the plane perpendicular to the coordination axis to the metal. Other changes which are being considered include a recognition that pi coordination to very electron deficient metals is stronger (with substantially shorter distances) than to the same metals in not very positive oxidation states.

All of these changes should be appropriate for the wide spectrum of organometallic compounds, but needs to be tested with the Cambridge Data Set.

Joseph J. Gajewski, ER13616

Publications for the past funding period

1. J.J. Gajewski, "A Simitheoretical Multiparameter Approach to Correlate Solvent Effects on Reactions and Equilibria," *J. Org. Chem.* 57, 1992 (1992).
2. G.C. Paul, J.J. Gajewski, "Thermal Isomerization of a Vinylcyclobutene to a Cyclohexadiene," *J. Org. Chem.* 57, 1970 (1992).

**DATE
FILMED**

8 / 5 / 93

END