1 of 1

DOE/ER/60759-T1

DE-FG03-84ER60754

POSTED

Ecology for a Changing Earth

James H. Brown, University of New Mexico
Jonathan Roughgarden, Stanford University

Final Report, February 6, 1990

Abstract

To forecast the ecological impact of global change, research initiatives are needed on the explicit role of humans in ecological systems, and on how ecological processes functioning at different spatial and temporal scales are coupled. Furthermore, to synthesize the results of ecological research for Congress, policymakers, and the general public, a new agency, called the United States Ecological Survey (USES) is urgently required. Also, a national commitment to environmental health, as exemplified by establishing a National Institutes of the Environment (NIE), should be a goal.

RECEIVED
JUN 01 1993
OSTI

1 Introduction

Our earth is changing. It has always been changing, but what is happening now matches the rare perturbations that produced huge effects such as the extinction of dinosaurs. The previous cataclysms were driven by astronomical processes such as continental drift, shifts in the earth's orbit, impact with comets, and fluctuations in the sun's brightness—but today the changes are caused by a single species, our own *Homo sapiens*.

We know that humans are changing the climate, the distribution of organisms and habitats, and the composition of air, water, and soil. Yet we lack a clear vision of how these changes will impact our own species, or the millions of others with whom we share the earth. While we know that a nuclear world war will make the earth uninhabitable, we are just beginning

to appreciate that the cumulative effect of our day-to-day activities might be just as severe.

In early December, 1988, a group of scientists met in Santa Fe, New Mexico to consider what ecological research is necessary to understand and to predict how we should cope with global change. The workshop, comprised of 23 scientists and 14 observers from government agencies, was supported jointly by grants from the National Science Foundation and the Department of Energy. This article is a report of that conference—it notes recent progress in the ability of ecologists to forecast results of ecological processes; it identifies several topics that urgently need research; and it recommends that Congress create a new single agency to determine the ecological state of the nation.

2 The Problem

At a glance, the problem seems clear. *Homo sapiens* has become the dominant species of the planet. Humans have converted about 11% of the earth's land area, and more than 90% of certain productive habitats, such as prairies and river valleys, to agricultural fields. Most of the remaining land, together with coastal waters and lakes, is used to harvest natural resources. Two recent studies estimate that *Homo sapiens* uses 20–40% of the terrestrial primary production of the earth (Vitousek et al. 1986, Wright 1987).

By burning fossil fuels and cutting forests, humans have altered the balance of carbon dioxide production and consumption, causing a dramatic increase in the concentration of CO₂ in the atmosphere. The resulting "greenhouse" effect is almost certainly changing the earth's climate, causing a global warming predicted to average 2°C by the middle of the next century (Jaeger 1988).

By converting forests to range and fields, and grasslands to deserts, humans have transformed the size and configuration of the earth's major habitats. Some species, such as domestics and those categorized as "pests" and "weeds," have benefitted, but most have not. The great majority of the earth's species are now confined to small, isolated fragments of once widespread habitats, and species are going extinct at a rate matched only by the few cataclysmic mass extinctions recorded in the fossil record. The biological diversity of the earth is declining daily.

By releasing toxic wastes, humans are changing the earth in ways lacking historical precedent. Until recently, severe effects of pollution were local, restricted to air, rivers, and lakes near population centers or dumping sites. But today pollutants, including acid rain and fluorohydrocarbons in the atmosphere, and organic toxins and heavy metals in groundwater, lakes, rivers, and coastal oceans, impact large geographic regions. What is not clear is what effects these changes are having, either for the earth, or for us. (Committee on Global Change 1988).

To illustrate the state of contemporary ecology, we note briefly some high points of recent research. This will indicate the limitations of existing knowledge and the need for research initiatives.

3 Advances in Ecological Research

Ecology is a basic science that determines how organisms interact with each other and with their physical environment. Ecological science shows how these interactions determine when species are common and rare, and where species are found. Ecological science uses controlled, manipulative experiments in field situations to test hypotheses; constructs mathematical models to forecast how ecological systems change; and applies technology to follow the passage of energy and matter through organisms and their environments. Although these activities were initiated much earlier, by the 1960s they had coalesced to make ecology the coherent modern science it is today.

Experimental advances trace to Connell's (1961) study of barnacles on rocks along the seashore, which showed how experimental techniques could be taken out of the laboratory and into a natural setting. This study demonstrated the roles of competition, predation, and physical disturbance on the abundance and distribution of species. Other experiments in the marine intertidal zone (Paine 1966, Lubchenco and Menge 1978), and increasingly in freshwater (Hall et al. 1970, Wilbur 1972, Werner and Hall 1976) and terrestrial (Brown and Davidson 1977, Pacala and Roughgarden 1985) systems, provided further evidence that species interactions and physical disturbance affect abundance and distribution (see Diamond and Case 1986, Kikkawa and Anderson 1986, Carpenter 1987, Gee and Giller 1988).

Contemporary mathematical theory traces largely to the late Robert MacArthur's contributions in three areas: interspecific interactions and com-

munity structure (MacArthur and Levins 1967, MacArthur 1972), the composition and dynamics of island biotas (MacArthur and Wilson 1963), and the theory of optimal foraging (MacArthur and Pianka 1966.) Other modeling efforts have helped to understand how natural selection, in concert with ecological processes, shape the life histories of organisms (Roughgarden 1971, Charlesworth 1980), what combinations of physical conditions permit organisms to survive ("climate spaces" Porter and Gates 1969, Porter and Tracy 1983), how food webs are organized (Cohen 1978, Pimm 1982), how forests regenerate after disturbance (Botkin et al. 1972, Shugart 1984), and how parasites and disease spread (May and Anderson 1979). The broad spectrum of contemporary mathematical theory in ecology has been recently reviewed (Roughgarden et. al. 1989).

Ecosystem ecologists have used automated analytical methods to quantify the levels and fluxes of energy and critical elements in entire marshes (Teal 1962), lakes (Edmondson 1979, Riemann and Sondergaard 1986), and forests (Borman and Likens 1979). Radioactive labels and natural isotopic signatures enable ecologists to follow organic matter and inorganic nutrients through food webs, and to compare rates of physical transport, primary production, consumption, and decomposition in different ecosystems. An important message of many recent ecosystem studies is the essential role of particular species or functional groups of organisms in biogeochemical processes (Carpenter and Kitchell 1988, Naiman et al. 1988, Pastor et al. 1988). And theoretical progress is being made in representing the hierarchical structure of ecosystems (O'Neill 1989).

4 Confronting Global Change

Despite the encouraging accomplishments of the last two decades, new ways of doing science are needed to address problems related to global change. In the past, most investigators have worked alone or in collaboration with only a few colleagues or students having similar backgrounds and interests. Experimentalists have conducted their own manipulations and collected their own data. But the scale and complexity of global problems requires collaboration among scientists in different disciplines, and the analysis of large data sets assembled by many investigators. Moreover, alternative methods of scientific inference, such as the "reconstructional logic" of earth scientists, will have

to be relied upon.

The workshop identified three major inadequacies in ecological expertise: the failure to consider humans as integral components of ecological systems; insufficient understanding of how processes that operate at vastly different spatial and temporal scales combine with one another; and lack of nation-wide coordination in obtaining, analyzing, and disseminating basic data on organisms and the environment.

4.1 Humans as an Ecological Component

Ecology, which derived from natural history studies of the last century, has traditionally defined “natural” as non-human. In searching for study sites, the emphasis has been on uncut forest, unplowed prairie, and unpolluted lake, stream, or ocean shore.

Human impacts are now found in all habitats, including the most remote deserts. Studies, even in “natural habitats,” that fail to take these human effects into account will necessarily be incomplete. Moreover, heavily human-modified systems, such as agricultural fields and urban areas, present special opportunities for ecological research—they are readily accessible, economically important; and their simple geometry, low species diversity, and sometimes well-documented perturbations facilitate study.

Today we know only roughly how humans interact with the environment. Clearly, they directly destroy some habitats while they create others; they change the configuration of landscapes, and release toxins and other substances that affect organisms and biogeochemical processes. Humans also directly affect other organisms by preying on, or competing with, wild species; by propagating domesticated plants and animals; and by defending themselves against parasites and diseases. Still, some of the most important impacts of humans may be indirect—caused by effects on other species that play key roles in ecological processes, such as nitrogen fixation or control of insect populations. Thus, the ecological role of humans is simply not well described.

Moreover, the ecological role of humans is shaped by social, economic, and political forces, as well as by strictly ecological conditions. What humans do depends on ecological factors, such as the supply of food and energy, and the suitability of the climate for human habitation, together with social attributes, such as attitudes about family size, and the style and size of

the economy. Conducting ecological research that includes humans calls for increased collaboration among ecologists and social scientists.

4.2 Scale—An Ecological Obstacle

A review of the ecological experiments published from January 1980 to January 1987 by Kariëva and Anderson (1988) found that 50% of all studies were done on plots less than 1 *m* in diameter, and 25% used plots less than 25 *cm* in diameter! Furthermore, most experiments were deliberately designed with homogeneous study sites to minimize spatial variation between experimental replicates, and were conducted over short periods to reduce the possibly confusing effects of seasonal or year-to-year variation. Indeed, a survey of the literature by Tilman (1989) showed that 40% of ecological experiments lasted less than one year (generally one field season) and only 7% lasted five or more years. In a few instances where studies have focused on the influence of scale *per se*, the answers to particular questions depended on the spatial and temporal scale used in the experiment (Wiens 1986).

Similarly, ecological models, as simplifications of complex systems, consist mostly of closed, homogeneous, scale-free formulations. Moreover, insensitivity to the significance of scale may underlie some academic disagreements in ecology, such as those concerning the importance of competition among species, and the extent to which systems exhibit steady-state properties (Caswell 1976, Sale 1977, Anderson et al. 1981).

Yet studies increasingly reveal that ecological systems are open; that is, their properties are determined in part by what happens within them, but also by processes that operate in the larger system within which they are embedded. For example, the composition of ecological communities along rocky shores is determined not only by competition and predation among the barnacles, mussels, snails, and starfish that live there, but also by the role of offshore water currents in transporting the larvae of these animals (Roughgarden et al. 1988). Similarly, the present day distribution of terrestrial species has been caused by continental drift and variation in Pleistocene climate, as well as by local species interactions and responses to today's abiotic environment. (Roughgarden et al. 1987, Myers and Giller 1988, Brown and Kurzius in press). Thus, ecological systems have to be conceived, both theoretically and experimentally, as open systems coupled to and driven by larger scale processes than most studies have taken into account.

Predicting the ecological impact of global change requires that we grapple with scale now. Data on past climates from geological sources, predictions of future climates from mathematical models, and data on human habitat modification from satellite photographs all pertain to scales much larger than those used in traditional ecological studies. These data cannot be used as inputs to ecological problems without understanding how traditional ecological systems are coupled to such scales. Conversely, a clearer understanding of the natural scales of ecological processes will improve the design of remote sensing systems.

Understanding ecological scale will also help in synthesizing information across the various subdisciplines of ecology. These subdisciplines, organismal, population, community, and ecosystem ecology, are each centered about a characteristic "level of organization," and a corresponding spatial and temporal scale. But information must be transferred across these disciplines. For example, the dynamics of populations reflect the abilities of individual organisms. These abilities, however, may be adaptations that have evolved by natural selection—which is a population process. Similarly, experiments show that ecosystem properties, such as productivity and the cycling of certain nutrients, are often controlled by particular species, sometimes dubbed "key-stone species" (Carpenter and Kitchell 1988). Thus, contemporary studies show that the subdisciplines of ecology are interdependent. Nonetheless, the transfer of information among the subdisciplines remains sporadic. When, and how much, does an ecosystem ecologist need to consult a community ecologist, and *vice versa*? Understanding the scales of different processes, and the mechanisms that cause coupling among different scales, should provide guidelines for increasing synthesis across ecological subdisciplines.

4.3 Ecological Data and the Public

The ecological condition of the United States is virtually unknown. There is no accurate assessment of how many species of mites, nematodes, fungi, and bacteria occur within the country. Even for relatively well-studied groups, such as vertebrates and vascular plants, the most basic information about the distribution and abundance of species is dispersed among the data of ecologists, the collections of museums, and the records of government agencies. The abiotic environment has been almost equally poorly monitored. Only data on weather and selected aspects of air and water quality have

been collected nationwide on a systematic, long-term basis. Obviously, there is little basis for assessing the ecological consequences of present and future environmental change, if there is no convenient central source of good data on how the environment is changing and how the organisms are responding. Contemplating the impact of global change without a comprehensive data base on the environment is analogous to discussing economic policy without knowing the GNP, the value of the dollar, and the rate of unemployment.

No existing organization has a sufficiently broad mission, or enough resources, to assemble the necessary data base. The results of academic research are published in openly accessible peer-reviewed journals. But academia has no inherent mechanism for compiling and synthesizing all of this information, or for applying it to national needs—even to such an urgent problem as whether the earth will remain habitable. Valuable data on distributions of species are contained in museum collections. Some museums have computerized their collections, but there is no national network to make such data readily accessible. Government agencies with specific environmental missions, such as the Forest Service, National Marine Fisheries Service, Environmental Protection Agency, Fish and Wildlife Service, and National Aeronautics and Space Administration have data on particular kinds of organisms and certain environmental variables. But these are currently too compartmentalized to be widely useful for ecological research.

A new agency is, therefore, urgently needed to monitor the ecological state of the nation, and to provide synthetic products to scientists, the general public, and the legislative and executive branches of the government. An administrative model is the United States Geological Survey (USGS), and this new agency—the United States Ecological Survey—might similarly be dubbed “USES,” as discussed in more detail below.

5 Recommendations

5.1 Research Initiatives

The workshop identified two areas that are recommended for immediate funding initiatives by the National Science Foundation and the Department of Energy. The appropriate sections of these agencies to sponsor the initiatives are the Ecology Program in the Division of Biotic Systems and Resources at

the NSF, and the Ecological Research Division in the Office of Energy Research at the DOE. "Research initiative" is used here to mean a substantial increment of funds to support basic research proposals on a designated topic. These new funding priorities related to global change must not reduce the already inadequate funding for traditional ecological research. The subjects of the recommended initiatives are:

5.1.1 Humans as an Integral Component of Ecological Systems

This initiative should support experimental and theoretical studies on the basic ecology of moderately to heavily human-impacted systems, including forest, pastoral, agricultural, suburban, and urban systems. There is a need for studies of direct and indirect interactions between modern humans and wild species, including patterns and processes of human-caused extinctions, spread and impacts of invading species, effects of human landscapes on the abundance and distribution of organisms, responses of different kinds of organisms to predicted scenarios of climatic and sea level change, and the structure and function of human-assembled communities. There should be support for collaborative studies among ecologists and social scientists on topics such as the growth and resource use of human populations; the ecological basis of sustainable economic development; the effects of different social, economic, and political systems on landscapes and biological diversity; and the reciprocal relationships among human activities, the organization of ecological communities, and biogeochemical ecosystem processes. There should be support for the addition of suburban and urban sites, and incorporation of more population and community studies in the LTER (Long Term Ecological Research) network.

5.1.2 The Analysis of Ecological Scale

This initiative should fund theoretical and experimental studies that will characterize the natural scales of various processes, and show how processes that function at different scales are coupled to one another. Theoretical research for the initiative may involve quantitative system-specific models, simplified qualitative representations of particular systems, and abstract ideamodels. Results from models should be compared with remote sensing data, geographic data bases, and projections from global climate and physical-

oceanographic models. Analysis of the fossil record and collaboration with paleobiologists will help to describe the biological impact and ecological implication of current changes in relation to past changes. Statistical research on the relation of experimental design to spatial analysis is also needed. Artificial intelligence techniques may aid in integrating information across various scales. Experimental studies of scale and coupling will be critical, especially in conjunction with theoretical research. Nonmanipulative studies, as carried out by astronomers, climatologists, and geologists, may be appropriate for processes whose scales are too large for conventional experiments, but it is also important to search for experimental approaches that can be “scaled up.”

5.2 A New Government Agency

The workshop concluded that a new government agency, to be called the United States Ecological Survey (USES), should be formed. This would acquire, analyze, and interpret data on the ecological state of the environment. It would conduct a long-term, nationwide ecological inventory. It would bring together theoretical and experimental scientists who, together with its permanent staff, would produce syntheses of the data, and other products. The agency would be responsible for making the data available through computer networks, and for distributing the products made from the data. The agency could make forecasts of upcoming problems. It would have the capability of responding rapidly to catastrophies, such as the Mt. Saint Helens eruption, or an accident at a nuclear power plant. The agency would also facilitate the integration of ecological data with other large-scale and long-term data bases, such as those pertaining to climate, geology, oceanography, and paleontology. Its products would be available to government, industry, educational institutions and the general public.

6 Further Observations

The workshop observed that USES, with its primary focus on ecology, addresses only a piece of a larger issue—the absence of a major national commitment to research on “environmental health,” analogous to its commitment to research on personal health *via* the National Institutes of Health. Yet, en-

Environmental problems are at least as threatening to our welfare as traditional medical problems. Moreover, many medical problems including cancer, allergies, and microbial-caused diseases have, in part, an environmental cause. Therefore, the formation of a multidisciplinary agency, to be called the National Institutes of the Environment (NIE), seems warranted now. The NIE could have five institutes: the U. S. Ecological Survey, mentioned above, an Institute of Basic Ecology, an Institute of Applied Environmental Sciences, an Institute of Human Demography, and an Institute of Sustainable Economics. The latter two institutes recognize that nearly all problems of environmental health are inextricably linked to human population growth and economic development. The creation of NIE would be a bold step, requiring the leadership of both the executive and legislative branches of the Federal Government. It would signal a national commitment to address the plight of our changing earth.

7 Conclusion

Humans are causing Herculean changes to the earth. Yet there is time to understand what our activities are leading to and, with that knowledge, surely, we can continue to inhabit this planet. The exciting advances from ecological research during the last two decades can be built upon to help ensure the possibility of a future for our descendents.

8 Acknowledgements

We gratefully acknowledge support from the Ecological Research Division, Department of Energy (DE-FG03-89ER60759) and the General Ecology Program, National Science Foundation (BSR-8821458).

9 References

Anderson, G., A. Ehrlich, P. Ehrlich, J. Roughgarden, B. Russell, and F. Talbot. 1981. The community structure of coral reef fishes. *American Naturalist* 117:476-495.

Borman, F. H., and G. E. Likens. 1979. Pattern and process in a forested ecosystem. Springer-Verlag, New York.

Botkin, D.B., J. F. Janak, and J. R. Wallis. 1972. Some ecological consequences of a computer model of forest growth. *Journal of Ecology* 60:849-872.

Brown, J. H., and D. W. Davidson. 1977. Competition between seed-eating rodents and ants in desert ecosystems. *Science* 196:880-882.

Brown, J. H., and M. Kurzius. 198-. Spatial and temporal variation in guilds of North American desert rodents. In Z. Abramsky, B. J. Fox, D. W. Morris, and M. R. Willig, eds., *Ecology of small mammal communities*. Special Publications of the Museum, Texas Tech University (in press).

Carpenter, S. R. (ed.) 1987. Complex interactions in lakes. Springer-Verlag, New York.

Carpenter, S. L., and J. F. Kitchell. 1988. Consumer control of lake productivity. *BioScience* 38:764-769.

Carpenter, S.R., J. F. Kitchell, and J. R. Hodgson. 1985. Cascading trophic interactions and lake productivity. *BioScience* 35:634-639.

Caswell, H. 1976. Community structure: a neutral model analysis. *Ecological Monographs* 46:327-354.

Charlesworth, B. 1980. Evolution in age-structured populations. Cambridge University Press, Cambridge, England.

Cohen, J. E. 1978. Food webs and niche space. Princeton University Press, Princeton, NJ.

Committee on Global Change. 1988. Toward an understanding of global change. National Academy Press, Washington, DC.

Connell, J. H. 1961. The influence of interspecific competition and other factors on the distribution of the barnacle *Chthamalus stellatus*. *Ecology*

42:710-723.

Edmondson, W. T. 1979. Lake Washington and the predictability of limnological events. *Arch. Hydrobiol. Bieh.* 13:234-241.

Diamond, J., and T. J. Case. 1986. *Community ecology*. Harper and Row, New York.

Gates, D. M. 1980. *Biophysical ecology*. Springer-Verlag, New York.

Ghee, J. H. R., and P. S. Giller. 1987. *Organization of communities: past and present*. Blackwell Scientific Publications, Oxford.

Hall, D. J., W. E. Cooper, and E. E. Werner. 1970. An experimental approach to the production dynamics and structure of freshwater animal communities. *Limnology and Oceanography* 15:839-928

Jaeger, J. 1988. Developing policies for responding to climate change. *World Climate Program Impact Studies*. WMO/TD No. 225, Stockholm.

Kareiva, P. and M. Andersen. 1988. Spatial aspects of species interactions: the wedding of models and experiments. In A. Hastings, ed., *Community Ecology*, pp.38-54. Springer Verlag, New York

Kikkawa, J., and D. J. Anderson. 1986. *Community ecology: pattern and process*. Blackwell Scientific Publications, Melbourne.

Lubchenco, J., and B. A. Menge. 1978. Community development and persistence in a low rocky intertidal zone. *Ecological Monographs* 48:67-94.

MacArthur, R. H. 1972. *Geographical ecology*. Harper and Row, New York.

MacArthur, R. H., and R. Levins. 1967. The limiting similarity, convergence and divergence of coexisting species. *American Naturalist* 101:377-385.

MacArthur, R.H., and E. R. Pianka. 1966. On optimal use of a patchy environment. *American Naturalist* 100:603-609.

- MacArthur, R. H., and E. O. Wilson. 1967. The theory of island biogeography. Princeton University Press, Princeton, NJ.
- May, R. M., and R. M. Anderson. 1979. Population biology of infectious diseases. *Nature* 280:455-461.
- Myers, A. A., and P. S. Giller. 1988. Analytical biogeography. Chapman and Hall, London.
- Naiman, R. J., C. A. Johnston, and J. C. Kelley. 1988. Alteration of North American streams by beaver. *BioScience* 38:753-762.
- O'Neill, R. V. 1989. Perspectives in hierarchy and scale. In J. Roughgarden, R. May, and S. Levin, eds., *Perspectives in Ecological Theory*, pp. 140-156. Princeton University Press.
- Pacala, S. and J. Roughgarden. 1985. Population experiments with *Anolis* lizards of St. Maarten and St. Eustatius. *Ecology* 66:129-141.
- Paine, R. T. 1966. Food web complexity and species diversity. *American Naturalist* 100:65-75.
- Pastor, J., R. Naiman, B. Dewey, and P. MacInnes. 1988. Moose, microbes, and the boreal forest. *BioScience* 38:770-777.
- Pimm, S. L. 1982. Food webs. Chapman and Hall, London.
- Porter, W. P., and D. M. Gates. 1969. Thermodynamic equilibrium of animals with environment. *Ecological Monographs* 39:245-270.
- Porter, W. P., and C. R. Tracy. 1983. Biophysical analyses of energetics, time-space utilization, and distributional limits of lizards. In E. R. Pianka and T. W. Schoener, eds., *Lizard ecology: studies of a model organism*, pp. 55-83. Harvard University Press, Cambridge, MA.
- Riemann, B., and M. Sondergaard. 1986. Carbon dynamics of eutrophic, temperate lakes. Elsevier, Amsterdam.

- Roughgarden, J. 1971. Density-dependent natural selection. *Ecology* 52:453-468.
- Roughgarden, J., S. Gaines, and S. Pacala. 1987. Supply side ecology: the role of physical transport processes. In J. Gee and P. Giller, eds., *Organization of Communities Past and Present*, pp 491-518. Blackwell Scientific Publications, Oxford England.
- Roughgarden, J., S. Gaines, and H. Possingham. 1988. Recruitment dynamics in complex life cycles. *Science* 241:1397-1560.
- Roughgarden, J., R. May, and S. Levin. 1989. *Perspectives in ecological theory*. Princeton University Press, Princeton New Jersey.
- Sale, P. 1977. Maintenance of high diversity in coral reef fish communities. *American Naturalist* 111:337-359.
- Shugart, H. H. 1984. *A theory of forest dynamics*. Springer-Verlag, New York.
- Teal, J. M. 1962. Energy flow in a salt marsh ecosystem of Georgia. *Ecology* 43:614-624.
- Tilman, D. 1989. Ecological experimentation: strengths and conceptual problems. In G. E. Likens, ed., *Long-Term Studies in Ecology. Approaches and Alternatives*, pp. 136-157. Springer-Verlag, New York.
- Vitousek, P.M., P.R. Ehrlich, A.H. Ehrlich, and P.A. Matson. 1986. Human Appropriation of the Products of Photosynthesis. *BioScience* 36:368.
- Werner, E. E., and D. J. Hall. 1976. Niche shifts in sunfishes: experimental evidence and significance. *Science* 191:404-406.
- Wiens, J. A. 1986. Spatial scale and temporal variation in studies of shrub-steppe birds. In J. Diamond and T. J. Case, eds., *Community ecology*, pp. 154-172. Harper and Row, New York.

Wilbur, H. M. 1972. Competition, predation, and the structure of the *Ambystoma-Rana sylvatica* community. *Ecology* 53:3-20.

Wright, D.H. 1987. Estimating Human Effects on Global Extinction. *International Journal of Biometeorology* 31:293.

10 Workshop Personnel

10.1 Organizers

Dr. James Brown, University of New Mexico

Dr. Jonathan Roughgarden, Stanford University

10.2 Participants

Dr. Timothy Allen, San Jose State University

Dr. Stephen Carpenter, University of Notre Dame

Dr. Hal Caswell, Woods Hole Oceanographic Institution

Dr. Peter Chesson, Ohio State University

Dr. Joel Cohen, Rockefeller University

Dr. Jared Diamond, UCLA Medical Center

Dr. Stuart Fisher, Arizona State University

Dr. Robert Holt, University of Kansas

Dr. Robert Howarth, Cornell University

Dr. Jennifer Kitchell, University of Michigan

Dr. Philippe Martin, NCAR

Dr. Joseph McAuliffe, University of Nevada

Dr. Bruce Menge, Oregon State University

Dr. Bruce Milne, University of New Mexico

Dr. Stephen Pacala, University of Connecticut

Dr. John Pastor, University of Minnesota

Dr. Thomas Schoener, UC Davis

Dr. David Strayer, Cary Arboretum

Dr. C. Richard Tracy, Colorado State University

Dr. S. David Webb, University of Florida

Dr. John Wiens, Colorado State University

10.3 Invited Observers

Dr. Fairley Barnes, Los Alamos National Laboratories
Dr. Walter Conley, Department of Energy
Dr. Diana Freckman, National Science Foundation
Dr. Charles Keller, Los Alamos National Laboratories
Dr. James Kercher, Lawrence Livermore National Laboratory
Dr. William Parton, National Science Foundation
Dr. Garth Redfield, National Science Foundation
Dr. John Rummel, NASA Headquarters
Dr. Edward Rykiel, Jr., Department of Energy
Dr. Michael Simmons, Santa Fe Institute
Dr. Michael H. Smith, Savanna River Ecology Laboratory
Dr. Stephen Threlkeld, National Science Foundation
Dr. Ron Walters, Los Alamos National Laboratory
Dr. Patrick Webber, National Science Foundation

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DATE
FILMED**

8 / 4 / 93

END