AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

Lynda Shapiro
University of Oregon

RECEIVED

JUN 11 1993

OSTI

SIX MONTH PROGRESS REPORT

DOE OCEAN MARGINS PROGRAM: GRANT #DE-FG06-92ER61423

'Controls on marine carbon fluxes via phytoplankton-mesoplankton interactions in continental shelf waters' - Lynda Shapiro, Barry F. Sherr, Evelyn B. Sherr, P.I.'s

Overall goals: The project is an in-depth evaluation of the phytoplankton --> phagotrophic protist trophic link. The principal goals of the first year are to develop methods for the second phase of the Ocean Margins Program: investigations in the field. Our project is focused on:

1) Impact of grazing by phagotrophic protists on phytoplankton, particularly on phototrophic cells < 5 μm in size which are not effectively grazed by metazooplankton.

2) Impact of grazing by phagotrophic protists on bacterioplankton.

3) Taxon-specific growth rates of phytoplankton in situ, particularly of < 5 μm sized cells, as they are affected by phagotrophy rates.

Progress to date:

During the first six months of the first year, research focused on:

1) Impact of protistan grazing on phytoplankton: We have worked on several approaches to investigation of the protist-phytoplankton grazing link. Our first studies dealt with differential grazing pressures on different phytoplankton taxa within natural coastal phytoplankton assemblages. We added live cells from laboratory cultures of several species of phytoplankton, including the coccoid cyanobacterium, Synechococcus, a 1.5 μm eukaryote, Micromonas pusilla, and several other eukaryotic algae < 8 μm in size: Emiliania huxleyi, Pelagococcus subviridis, Pycnococcus provasolii, and Thalassiosira pseudonana. Preference experiments, in which in situ phagotrophic protists were offered two different taxonomic types of phytoplankton of similar size, were carried out. Protist ingestion of added phytoplankton cells was determined by inspection of protist food vacuole contents and by following decrease in numbers of phytoplankton cells over time. During the fall, protists appeared to be omnivorous, ingesting both the prokaryote Synechococcus and the eukaryote Pycnococcus. However, in the winter, there was a strong preference among in situ protists for the eukaryote Micromonas over the prokaryote Synechococcus.

MASTER

sp

The biomass of herbivorous protists was approximately equally divided between ciliates and flagellates, and a wide size spectrum of sizes and taxonomic types of protists were observed to ingest the added phytoplankton.

We also began evaluating a new approach for quantifying in situ grazing rates of protists on specific taxonomic types or size classes of phytoplankton: the Product Release Assay. This approach is based on adding a tracer amount of radiolabeled phytoplankton cells to a seawater sample, and then following over time the cumulative amount of radiolabel that appears in the dissolved fraction due to grazing activity. We have carried out one Product Release experiment using radiolabeled *E. huxleyi* cells, and found that significant release of label into the dissolved fraction only occurred in the whole water fraction, not in $< 20 \mu\text{m}$ filtered water (microzooplankton removed) or in $< 0.8 \mu\text{m}$ filtered water (all grazers removed). Based on the data, we calculated a grazing rate, g , of 0.024 h^{-1} , or a $1/2$ decay time of 29.4 hours.

2) Impact of protist grazing on bacterioplankton: development of a enzyme-based method for estimating instantaneous rates of protistan bacterivory (with Juan Gonzalez): Quantifying in situ rates of consumption of bacteria by protists is difficult. We evaluated a novel approach to the problem of estimating protistan bacterivory: quantifying the concentration in the particulate fraction of the enzyme, lysozyme, with optimal activity at acid pH. Lysozyme specifically degrades peptidoglycan, a major component of bacterial cell walls; protist food vacuoles are acidic (pH of 4-5). We used a fluorochrome-linked artificial substrate: MUF-triacetylchitotriose, which is an analogue of peptidoglycan. The fluorochrome, MUF, only fluoresces when cleaved from the substrate, and is detectable in nanomolar amounts using a standard fluorometer with appropriate filter sets. The approach offers a 'snapshot' of recent bacterivory, with no incubations required. We found a peak of lysozyme activity at pH 4.5 in a variety of protist cultures and coastal water samples. The acid enzyme activity values were tightly correlated with rate of uptake of fluorescently labeled bacteria (FLB) measured in separate aliquots of the same samples. The description of the method has been submitted for publication. Digestive enzyme assays appear to be a promising new approach to investigating in situ protist grazing activity.

3) Cell-specific growth rates via immunostaining and DNA analysis: We have made good progress with quantitative analysis of DNA for cell cycle analysis and growth rate determinations. Fluorescent polycarbonate beads with a diameter of 1.54 microns (s.d. = .015) were used to examine the errors associated with this image analysis system. These beads are the approximate size of the *E. huxleyi* (92A) nucleus and thus represent a good comparison for purposes of calibration. Our analyses using fluorescent beads have shown a fluorescence CV of 9.64% (comparable to previously published reports of C.V. = 6.12% for

beads approximately 5X the size used in our calibration). We have assessed the appropriate DAPI staining concentrations and times for the two E. huxleyi clones (92A and 92D), and are adjusting existing protocols to extract chlorophyll from the E. huxleyi clones without altering cell shape.

4) Enumeration of phytoplankton and bacteria: We have developed a cell counting program for the Bioscan OPTIMAS image analysis software package. This program works as a macro within the OPTIMAS environment allowing the user to interactively select and measure cell specific morphometric characteristics, (area, greatest x, greatest y), sort cells into categories (dependent on user selected morphometrics), and tally the number of cells within those classes. This program will also calculate and tally volumes of cells within user defined categories. Approximately 1000 cells may be counted in 5 minutes. At present this system operates best with cell monocultures or phytoplankton assemblages with distinct size separations between species.

Overall, the work proposed in the project has been going as planned, or in the case of the new method for bacterivory, has branched out in very promising new directions. We are on target and making good progress.

Presentations and submitted mss based on this project:

Sherr, E.B., B.F. Sherr, and L.P. Shapiro. Grazing on pico-phytoplankton by coastal protistan assemblages. ASLO Aquatic Sciences Meeting, Santa Fe, New Mexico, Feb 9-14, 1992.

Gonzalez, J.M., B.F. Sherr, and E.B. Sherr. Digestive enzyme activity as a quantitative measure of protistan grazing: the acid lysozyme assay for bacterivory. ms submitted to Mar. Ecol. Prog. Ser.

In preparation:

Keller, M.D., L.P. Shapiro, E.B. Sherr, B.F. Sherr, E.M. Haugen and T.L. Cucci. Phagotrophy of fluorescently-labelled bacteria by an oceanic phytoplankter.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DATE
FILMED**

7 / 27 / 93

END