AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

Mechanisms and Genetic Control of Interspecific Crossing Barriers in *Lycopersicon*

Second year progress report: 3/27/93

Grant Number: DOE-FG02-91ER20037

PI: Martha A. Mutschler, Cornell University, Ithaca, NY

Shiela McCormick USDA Plant Gene Expression Center, Albany, CA

The goal of this program is to use *Lycopersicon esculentum* and *L. pennellii* as a model system to study the interspecific reproductive barriers unilateral incongruity (UI), hybrid breakdown and interspecific aberrant ratio syndrome (IARS). The specific experiments proposed and progress to date in each category listed in the original proposal are as follows.

A. Unilateral incongruity:

1) Determine the functional basis of UI including the timing of the failure of incongruous crosses, the developmental step(s) interrupted by UI, the tissue and genomes involved in UI.

The work for this section regarding the characterization of pollen tube growth and fertilization in congruous crosses and on the tissues and genomes involved in UI was largely completed and described in the year 1 report. The remaining work in this section involves an examination of second mitotic division in congruous versus incongruous crosses. As reported in the 8/1/92 progress report, we have narrowed unilateral incongruity in this cross to be due to failure in early pollen tube development. Unfortunately, the previously published techniques to observe second mitotic division *in vivo* have not proved effective with our material. Therefore, we are resorting to the use of the clearing technique developed for use in mixed pollen pollinations (see section A.2) and phase contrast microscopy to complete this section. Crosses for various early time points have been made and processed through the clearing protocol and are being analyzed presently. We are also preparing a manuscript describing the work to date on the nature of UI. This manuscript will include the information on UI in section A.1 and A.2 in the year 1 and 2 progress reports and on the study of second mitotic division.

2) Study male/female and direct or indirect male/male interaction in mixed pollen pollinations using pollen from plants homozygous for a pollen expressed GUS transformants, and determine whether effects of male/female interaction may be associated with UI.

We proposed to test for possible interactions between the male gametophyte and the female sporophyte or gametophyte. These tests involve a series of pollinations with mixed pollen, in which one pollen type will be produced by a plant with a pollen-specific promoter/GUS construct, allowing visual discrimination of the two pollen types in studies of pollen germination and pollen tube growth in congruous and incongruous crosses. Technical difficulties were encountered in 1) observing GUS stained pollen tubes, due to interference from chlorophyll in the pistil; and 2) the inability to combine staining pollen tubes with aniline blue and various methods to improve visibility of GUS activity. To overcome this, we have developed a protocol to stain for GUS, clear the pistil and observe specimens in phase contrast microscopy or Nomarski interference optics. Unfortunately the method uses a controlled substance, chloral hydrate, and tests of uncontrolled substances indicate that there is no satisfactory substitute for the chloral hydrate. Use of the controlled substance necessitated testing the protocol in a licensed lab and obtaining legal access to the compound to do the proposed experiment. As of January 1993, we are able to obtain the compound and are in the midst of making pollinations and processing them through the protocol. Since the UI response occurs relatively early in pollen tube growth, this experiment will be simpler than initially devised. There is no need to examine later stages in pollen tube growth, as these are eliminated in incongruous crosses. A publication on the technique has been submitted.

MASTER
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

3) Study the genetic control of UI.

Initially we were going to eliminate the work in this section due to the reduced level of funding. However, we have started this experiment since the information derived from it is critical to the program as a whole. In our initial results reported in progress report 8/1/92, we mentioned only four crosses ($Le \times F_2$, $Lp \times F_2$, $F_2 \times Le$, $F_2 \times Lp$), however, due to fertility fecundity problems and to determine what is happening in the F_1 , we have added three additional types of crosses ($F_1 \times F_2$, $F_2 \times F_1$, and F_2 selfed). The materials are currently being processed and analyzed.

B. Hybrid Breakdown:

1) determine the functional basis of non-fecundity, including the stage in reproduction affected, the role of pollen tube growth in non-fecundity, and whether female sterility or a non-reciprocal barrier such as UI is associated with non-fecundity,

This work is largely done and described in the report of year 1. The remaining part of the work involves an examination of the events immediately before/during/after double fertilization in fecund and non-fecund plants from F_2^{LP} and F_2^{Le} to determine the time and nature of the barrier in female fecundity in hybrid breakdown. This work was delayed until the technique discussed in A.1) and A.2) above was developed. Plants are currently being grown so that this work can be done.

2) determine the effects of cytoplasm on the reproductive parameters affected by hybrid breakdown, and whether these effects are direct or indirect, by its affect on IARS.

See report of year 1 for description of work pertaining to the first half of this question. The second part of the question posed in this section involves whether the effects of the cytoplasm on reproductive parameters are direct or indirect, through the effect of cytoplasm on IARS. To answer this question, the completion of a 20cM maps using the twin populations, population 3 (F_2^{Le}) and population 4 (F_2^{LP}) was required. Ninety-seven RFLP markers (From Dr. S. D. Tanksley) and three isozyme markers were used to construct genetic maps. Data of population 3, population 4, and pooled data from both populations were analyzed using the MAPMAKER computer program. Map units (centimorgan, cM) were calculated using Kosambi function. Genetic maps constructed from all three data sets resulting in similar linkage groups and marker orders, except that data from population 4 showed no segregation in the top region of chromosome 1. The control data of the F_1 plant producing population 4 (F_2^{LP}) and the series plants producing the NY^{LP} plant indicated that the NY^{LP} plant used was heterozygous in this top region of chromosome 1, and the F_1 plant was homozygous for *L. pennellii* in this top region of chromosome 1. For the remainder of the genome, the genetic map from the pooled data had highest LOD and agrees with the tomato linkage map published by Dr. S. D. Tanksley except that two markers on chromosome 2 were inverted in order compared to the Tanksley map. Since quantitative traits loci (QTLs) analysis is under the hypothesis that the genetic map is known and fixed, we decided to use the genetic map constructed from the pooled data and the top section of chromosome 1 constructed by population 3 data for QTLs analysis. The final map contains 87 intervals covering 1350 cM with average spacing between markers of approximately 15.5 cM (Fig. 1).

A similar map is under construction using a much larger F_2^{LP} population (population 6) to provide a better opportunity of mapping genomic regions associated with nonfertility and non-fecundity. Nearly all of the 97 RFLP markers mapped in population 3 and 4 have been used in mapping population 6. The framework map, with average spacing between markers of approximately 20 cM, is currently being constructed.

3) determine the number and location of chromosomal regions associated with reduction in male fertility and non-fecundity

The relationship between reproductive parameters and genotypes was examined for each interval bounded by pairs of neighboring markers on the top of chromosome 1 using the computer program HYPERGENE. For each of the intervals examined in population 3, the subpopulation selected as being homozygous *L. esculentum* genotypes showed similar mean values and standard errors to that of the entire population for the reproductive parameters. Most of the population 3 subpopulations selected as being homozygous *L. pennellii* also showed similar mean values and standard errors to that of the entire population for the reproductive parameters. Furthermore, for most of the intervals, the subpopulations selected as being homozygous *L. pennellii* genotypes from population 3 still had lower mean values than that of population 4. Therefore, we conclude that there is no evidence that the homozygous *L. pennellii* genotype in population 4 for the top of chromosome 1 is responsible for the differences in the effects of hybrid breakdown observed between population 3 and population 4.

The association of markers with the reproductive parameters has been tested using several procedures. Percent pollen stainability data from population 3 (F_2^{Le}) and population 4 (F_2^{LP}) were applied as normal distributed quantitative data, tested for single marker by one-way ANOVA and regression using MINITAB software, and analyzed by interval mapping using MAPMAKER/QTL software. In population 3 (F_2^{Le}) 8 regions had markers which showed significance by the single marker test, of which one region was also significant on chromosome 4 by interval mapping. In population 4 (F_2^{LP}) 6 regions had markers which were significant for single marker test, of which one on chromosome 1 showed significance by interval mapping. Of all the related regions, there are 4 that indicated the same regions from both population 3 (F_2^{Le}) and population 4 (F_2^{LP}), and all 4 were indicated by single marker analysis.

Distribution of other reproductive parameters were extremely non-normal. Fruit set data from population 3 (F_2^{Le}) and population 4 (F_2^{LP}) has been examined with different transformations for interval mapping, and converted to categorical data for independence test. Several genomic regions were identified as being associated with fruit set by the independent tests, but the interval mapping failed to identify significant regions. Further analyses are needed including single marker analysis and fitting a cumulative logit model using categorical reproductive index.

C. Interspecific Aberrant Ratio Syndrome (IARS):

1) determine the functional basis of IARS including the stage of reproductive development affected by IARS, what layers of the female sporophyte are involved, whether both male and female gametes are affected and whether gamete competition/selection is involved in IARS,

We have available five transgenic *L. esculentum* cultivar 'VF36' (from Dr. McCormick's laboratory, each came from independent transformation) [in proposal I.A.3], two plants carry the chimeric gene LAT59-GUS and three plants carry the chimeric gene LAT52-GUS (Twell et al., 1989; McCormick et al., 1989). After pollinating the transgenic *L. esculentum* with *L. pennellii* accession LA716, two or three interspecific F_1 plants for each of the five transgenic plants were planted. GUS assay was done for pollen grains from three flowers of each F_1 plant. No evidence suggested that the ratio of GUS positive: GUS negative pollen deviated significantly from 1:1. GUS assay was done for immature seed from green tomato fruits resulting from manual self pollinations of the F_1 plants. Goodness-of-fit tests were done for a 3:1 (GUS positive: GUS negative) ratio for each fruit and each plant. There was no absolute consistency about which F_1 plant always produced fruits with significant deviation, but three F_1 plants, one with the chimeric gene LAT59-GUS and two with LAT52-GUS, were identified as skewers favoring *L. pennellii* genotype. Further investigation of *in vivo* pollen germination and

observation of double fertilization will answer the question about whether the GUS positive pollen tubes (*L. esculentum* genotype) was stopped in the style or not be able to proceed fertilized.

2) study the effect of cytoplasm on IARS, does exchange of cytoplasm affect IARS at chromosomal regions other than those near PGI-1 and PGM-2, and do the effects at these regions involve reversal of direction or elimination of IARS.

To detect segregation distortion for single locus, data from RFLP and isozyme mapped markers were analyzed using G^2 (likelihood-ratio chi-squared) tests (Table 1). For each marker a goodness-of-fit test for monogenic segregation (1:2:1 for homozygous *L. esculentum* genotype (EE): heterozygous *L. esculentum* and *L. pennellii* genotype (EP): homozygous *L. pennellii* genotype (PP)) was performed. This genotypic frequencies test with 2 degree of freedom was partitioned to two tests with 1 degree of freedom, the allele frequencies test and the allele based genotypic frequencies test. The allele frequencies test was a goodness-of-fit test for single locus allele segregation (1:1 for *L. esculentum* allele (E) and *L. pennellii* allele (P)). The allele based genotypic frequencies test was a goodness-of-fit test for random fertilization ($e^2:2ep:p^2$ where e and p are the frequencies of E allele and P allele, respectively, in the population for tested marker). If the segregation distortion was due to selection at prezygotic stage, the G^2 tests would show significance for the genotypic frequencies test and allele frequencies test but not for the allele based genotypic frequencies test. If the segregation distortion was due to some selection a postzygote stage and affects heterozygous genotype only, the G^2 tests would show significance for the genotypic frequencies test and allele based genotypic frequencies test but not for allele frequencies test. Results of the G^2 tests showed differences in the number and location of markers with skew in allelic frequencies. In population 3 (F_2^{Le}) and population 4 (F_2^{Lp}), there are 20 and four, respectively, markers of allele frequencies skew, of which all favor the *L. pennellii* allele. In population 3 (F_2^{Le}), there are 16 markers with 1:2:1 genotypic frequencies skew, 11 of which are due to allelic skew by favoring *L. pennellii* allele. In contrast, there are only eight markers with 1:2:1 genotypic frequencies skew in population 4 (F_2^{Lp}), four of which are due to allelic skew by favoring *L. pennellii* allele.

Presentations

Liu, S.-C. and M.A. Mutschler. An F₂ derived from *L. esculentum* x *L. pennellii* and possessing the cytoplasm of *L. pennellii* exhibits increased pollen fertility and fecundity and alterations of nuclear segregation ratios. XII International Congress on Sexual Plant Reproduction

Mutschler, M.A., S.C. Lui and B.E. Liedl. Unilateral incongruity is overcome by recurrent F₁ bridging permitting sexual transfer of *L. pennellii* cytoplasm to tomato (*L. esculentum*). XII International Congress on Sexual Plant Reproduction

Liedl, B.E., and M.A. Mutschler. Unilateral incongruity is distinct from self-incompatibility in expression, timing and location in interspecific crosses in the genus *Lycopersicon*", XII International Congress on Sexual Plant Reproduction

Published abstracts

Liedl, B.E., and M.A. Mutschler. (1990) The unilateral interspecific crossing barrier in *Lycopersicon* is not self incompatibility. HortScience 25:1079.

Liu, S.C. and M.A. Mutschler. (1990) The nature and means to overcome hybridbreakdown in the F₂ of an interspecific cross in *Lycopersicon*. HortScience 25:1078.

Manuscripts Submitted

Mutschler, M.A., and B.E. Liedl. Interspecific Crossing Barriers in *Lycopersicon* and their Relationship to Self-Incompatibility. In: "Genetic Control of Self-incompatibility and Reproductive Development in Flowering Plants, E.G. Williams, R.B. Knox and A.E. Clarke (eds).

Liedl, B.E., S. McCormick, and M.A. Mutschler. A clearing technique for the study of GUS activity in pollen tubes and ovules of *Lycopersicon*. Submitted to Plant Molecular Biology Reporter

Manuscripts in Preparation

Mutschler, M. A., E. D. Cobb, S.-C. Liu, and B. E. Liedl. Unilateral incongruity is overcome by recurrent F₁ bridging, permitting sexual transfer of *Lycopersicon pennellii* cytoplasm to tomato (*L. esculentum*).

Liu, S.-C., B. E. Liedl and M. A. Mutschler. Pollen fertility and fecundity are increased and the aberrant segregation of molecular markers is altered in the interspecific F₂ of *Lycopersicon esculentum* x *L. pennellii* by using *L. esculentum* containing *L. pennellii* cytoplasm.

Liedl, B.E., S. McCormick, and M.A. Mutschler. Unilateral incongruity is distinct from self-incompatibility in expression, timing and location in crosses involving *Lycopersicon pennellii* and *L. esculentum*.

Table 1. Molecular makers with segregation ratios deviating significantly from the expected ratios in population 3 (F_2^{Le}) and population 4 (F_2^{Lp}).

marker	chromosome	F_2^{Le} population data and G^2 tests						F_2^{Lp} population data and G^2 tests					
		EE	EP	PP	1:2:1 ^a	1:1 ^b	e ² :2ep:p ^{2c}	EE	EP	PP	1:2:1	1:1	e ² :2ep:p ²
TG224	1	5	32	12	*		*	0	0	53	- ^d	-	-
TG19	1	5	27	7	*		*	13	25	15			
TG430B	1	11	21	8				18	18	17			*
CD33A	2	6	23	12				9	33	9			*
TG469	2	8	26	12				5	32	10	*		**
TG48	2	6	24	17		*		6	35	14	*		*
TG141	2	13	29	10				8	32	10			*
TG130	3	8	21	18		*		6	35	14	*		*
TG235	3	6	22	22	**	***		3	31	19	****	***	*
CT90A	3	4	15	14	*	*		5	24	14			
TG134	3	7	25	20	*	*		8	27	13			
TG94	3	10	34	9			*	13	28	13			
TG272B	4	3	19	12	*	*		10	26	14			
CD64	5	11	12	15			*	13	23	15			
TG379	5	13	14	15			*	6	26	10			
TG318	5	14	16	17			*	10	31	13			
TG470	6	11	21	11				5	32	14	*		*
CT90B	6	5	24	7			*	9	28	13			
TG115	6	12	25	13				17	19	17			*
TG166	7	5	22	15		*		9	31	11			
TG61	7	6	20	17		*		9	26	14			
TG272A	7	5	24	6			*	8	27	10			
TG18	9	3	25	15	**	**		13	25	16			
TG223	9	7	22	17		*		12	27	14			
TG35	9	6	25	18	*	*		10	29	14			
TG230	10	1	19	21	****	****		3	20	21	****	****	
CT234	10	2	27	18	****	****	*	3	27	25	****	****	
TG43	10	5	25	18	*	**		8	21	23	**	***	
CD72	10	3	17	22	****	****		7	29	15			
TG63	10	8	14	21	***	***		10	30	11			
TG194	11	8	21	21	*	**		10	20	16			
CT211A	12	6	35	8	**		***	8	26	13			
Pgi-1	12	8	25	20		*		11	32	15			
TG111	12	7	15	18	*	*		8	27	15			
TG28A	12	7	24	17		*		8	32	9			*

Significance level: $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.005$ (***), $p < 0.001$ (****).

^a The genotypic frequencies G^2 test.

^b The allele frequencies test G^2 test.

^c The allele based genotypic frequencies G^2 test.

^d The marker did not segregate in population 4 (F_2^{Lp}).

**DATE
FILMED**

7 / 29 / 93

END