LEGIBILITY NOTICE

A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.

Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.

Received by OSTI

ORNL/RASA-88/20

MAR 20 1989

**OAK RIDGE
NATIONAL
LABORATORY**

MARTIN MARIETTA

**RESULTS OF THE
RADIOLOGICAL SURVEY
AT
266 EAST SPRING VALLEY
AVENUE (MJ024),
HACKENSACK, NEW JERSEY**

**R. D. Foley
J. W. Crutcher
R. F. Carrier
L. M. Floyd**

**OPERATED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represent that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

HEALTH AND SAFETY RESEARCH DIVISION

**Nuclear and Chemical Waste Programs
(Activity No. AH 10 05 00 0; ONLWCO1)**

**RESULTS OF THE RADIOLOGICAL
SURVEY AT 236 EAST SPRING VALLEY AVENUE (MJ024),
HACKENSACK, NEW JERSEY**

R. D. Foley, J. W. Crutcher, R. F. Carrier, and L. M. Floyd

Date of Issue - February 1989

Investigation Team

**R. E. Swaja - Measurement Applications and Development Manager
W. D. Cottrell - FUSRAP Project Director
R.D. Foley - Field Survey Supervisor**

Survey Team Members

**A. C. Butler
P. F. Tiner**

**Work performed by the
MEASUREMENT APPLICATIONS AND DEVELOPMENT GROUP**

**Prepared by the
CAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6285
operated by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U. S. DEPARTMENT OF ENERGY
under Contract No. DE-AC05-84OR21400**

MASTER

CONTENTS

LIST OF FIGURES	v
LIST OF TABLES	vii
ACKNOWLEDGMENTS	ix
INTRODUCTION	1
SURVEY METHODS	2
SURVEY RESULTS	2
Surface Gamma Radiation Levels	2
Systematic and Biased Soil Samples	2
SIGNIFICANCE OF FINDINGS	3
REFERENCES	4

LIST OF FIGURES

1	Gamma radiation levels ($\mu\text{R/h}$) measured on the surface at 266 East Spring Valley Avenue, Hackensack, New Jersey (MJ024)	5
2	Diagram showing locations of soil samples taken at 266 East Spring Valley Avenue, Hackensack, New Jersey (MJ024)	6

LIST OF TABLES

1	Applicable guidelines for protection against radiation	7
2	Background radiation levels for the northern New Jersey area	7
3	Concentrations of radionuclides in soil at 266 East Spring Valley Avenue, Hackensack, New Jersey (MJ024)	8

ACKNOWLEDGMENTS

Research for this project was sponsored by the Division of Facility and Site Decommissioning Projects, U.S. Department of Energy, under Contract No. DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc. The authors wish to acknowledge the support of J. E. Baublitz, Deputy Director, Office of Remedial Action and Waste Technology; J. J. Fiore, Director, Division of Facility and Site Decommissioning Projects; and members of their staff. The authors also appreciate the contributions of B. C. Littleton and L. J. Jeffers of IR&A Publications Office, B. S. Ellis, D. A. Roberts, and T. R. Stewart of the Measurement Applications and Development Group, and W. H. Shinpaugh of Don Stone Associates for participation in the collection, analyses, and reporting of data for this survey.

RESULTS OF THE RADIOLOGICAL SURVEY AT 266 EAST SPRING VALLEY AVENUE (MJ024), HACKENSACK, NEW JERSEY*

INTRODUCTION

From 1916 to 1956, process wastes and residues associated with the production and refining of thorium and thorium compounds from monazite ores were generated by the Maywood Chemical Works (MCW), Maywood, New Jersey. During the latter part of this period, MCW supplied rare earth metals and thorium compounds to various government agencies. In the 1940s and 1950s, MCW produced thorium and lithium, under contract, for the Atomic Energy Commission (AEC). These activities ceased in 1956, and approximately three years later, the 30-acre real estate was purchased by the Stepan Company. The property is located at 100 Hunter Avenue in a highly developed area in Maywood and Rochelle Park, Bergen County, New Jersey.

During the early years of operation, MCW stored wastes and residues in low-lying areas west of the processing facilities. In the early 1930s, these areas were separated from the rest of the property by the construction of New Jersey State Highway 17. The Stepan property, the interim storage facility, and several vicinity properties have been designated for remedial action by the Department of Energy (DOE).

The waste produced by the thorium extraction process was a sandlike material containing residual amounts of thorium and its decay products, with smaller quantities of uranium and its decay products. During the years 1928 and 1944 to 1946, area residents used these process wastes mixed with tea and cocoa leaves as mulch in their lawns and gardens. In addition, some of the contaminated wastes were apparently eroded from the site into Lodi Brook and carried downstream.

As a result of the Energy and Water Appropriations Act of Fiscal Year 1984, the property discussed in this report and properties in its vicinity contaminated with residues from the former MCW were included as a decontamination research and development project under the DOE Formerly Utilized Sites Remedial Action Program. As part of this project, DOE is conducting radiological surveys in the vicinity of the site to identify properties contaminated with residues derived from the MCW. The principal radionuclide of concern is thorium-232. The radiological survey discussed in this report is part of that effort and was conducted, at the request of DOE, by members of the Measurement Applications and Development Group of Oak Ridge National Laboratory.

A radiological survey of the private, residential property at 266 East Spring Valley Avenue, Hackensack, New Jersey, was conducted during 1987. The survey and sampling of the ground surface and subsurface were carried out on April 23, 1987.

*The survey was performed by members of the Measurement Applications and Development Group of the Health and Safety Research Division at Oak Ridge National Laboratory under DOE contract DE-AC05-84OR21400.

SURVEY METHODS

The radiological survey of the property included a gamma scan of the entire property outdoors and collection of surface and subsurface soil samples. No indoor survey measurements were performed.

Using a portable gamma scintillation meter, ranges of measurements were recorded for areas of the property surface. Systematic soil samples were then obtained at randomly selected locations irrespective of gamma exposure rates. In addition, biased soil samples were collected in areas of elevated gamma levels. These survey methods followed the plan outlined in Reference 1. A comprehensive description of the survey methods and instrumentation is presented in *Procedures Manual for the ORNL Radiological Activities (RASA) Program*, Oak Ridge National Laboratory, ORNL/TM-8600 (April 1987).²

SURVEY RESULTS

Applicable federal guidelines are summarized in Table 1.³ The normal background radiation levels for the northern New Jersey area are presented in Table 2. These data are provided for comparison with survey results presented in this section. All direct measurement results presented in this report are gross readings; background radiation levels have not been subtracted. Similarly, background concentrations have not been subtracted from radionuclide concentrations measured in environmental samples.

Surface Gamma Radiation Levels

Gamma radiation levels measured during a gamma scan of the surface of the property are given in Fig. 1. Gamma exposure rates over the major portion of the property ranged from 7 to 15 $\mu\text{R/h}$. Elevated gamma levels occurred near a brick and gravel area at the front of the house (17 to 25 $\mu\text{R/h}$) and at the back of the house near a metal shed (15 to 17 $\mu\text{R/h}$) and near the pool (11 to 17 $\mu\text{R/h}$).

Systematic and Biased Soil Samples

Six systematic soil samples (S1A-D and S2A-B) and five biased soil samples (B1A-C and B2A-B) were taken for radionuclide analyses from depths of 0 to 60 cm (systematic samples) and 0 to 45 cm (biased samples), at increments of 15 cm. Locations of the samples are shown in Fig. 2, with results of laboratory analyses provided in Table 3. Concentrations of ^{226}Ra and ^{232}Th in the systematic samples ranged from 0.74 to 1.3 pCi/g and from 0.86 to 3.5 pCi/g, respectively. In the biased samples, concentrations ranged from 0.75 to 1.6 pCi/g for ^{226}Ra and from 1.3 to 6.8 pCi/g for ^{232}Th . All areas on the site are in compliance with the general criteria of 5 and 15 pCi/g for surface and subsurface soil, respectively, averaged over 100 m². The average concentrations of ^{232}Th in the contaminated areas are 1.8 pCi/g for the area near the pool (8.7 m²) and 1.2 pCi/g for the area in the front yard (4.4 m²). These concentrations are well within the "hot spot" criteria shown in Table 1. Radium-226 concentrations in some samples were slightly elevated above

background for the northern New Jersey area (Table 2), but all concentrations were well below DOE guidelines.

SIGNIFICANCE OF FINDINGS

Measurements taken at 226 East Spring Valley Avenue indicate that the property contained radioactive contamination primarily from the ^{232}Th decay chain, with some contamination from ^{226}Ra . These radionuclide distributions are typical of the type of material originating from the processing operations at MCW. However, all areas of this property conform to the radiological guidelines established for the Maywood, New Jersey, remedial action plan.

REFERENCES

1. W. D. Cottrell, ORNL, to A. J. Whitman, DOE/HQ, correspondence, "Radiological Survey of Private Properties in Lodi, New Jersey" (August 15, 1984).
2. T. E. Myrick, B. A. Berven, W. D. Cottrell, W. A. Goldsmith, and F. F. Haywood, *Procedures Manual for the ORNL Radiological Survey Activities (RASA) Program*, Oak Ridge National Laboratory, ORNL/TM-8600 (April 1987).
3. U.S. Department of Energy, *Guidelines for Residual Radioactivity at Formerly Utilized Sites, Remedial Action Program and Remote Surplus Facilities Management Program Sites* (Rev. 2, March 1987).
4. T. E. Myrick and B. A. Berven, *State Background Radiation Levels: Results of Measurements Taken During 1975-1979*, Oak Ridge National Laboratory, ORNL/TM-7343 (November 1981).

Fig. 1. Gamma radiation levels ($\mu\text{R/h}$) measured on the surface at 266 East Spring Valley Avenue, Hackensack, New Jersey (MJ024).

266 E Spring Valley Ave

Fig. 2. Diagram showing locations of soil samples taken at 266 East Spring Valley Avenue, Hackensack, New Jersey (MJ024).

Table 1. Applicable guidelines for protection against radiation^a

Mode of exposure	Exposure conditions	Guideline value
Radionuclide concentrations in soil	Maximum permissible concentration of the following radionuclides in soil above background levels averaged over 100 m ² area ²³² Th ²³⁰ Th ²²⁸ Ra ²²⁶ Ra	5 pCi/g averaged over the first 15 cm of soil below the surface; 15 pCi/g when averaged over 15-cm thick soil layers more than 15 cm below the surface
Guidelines for nonhomogeneous contamination (used in addition to the 100 m ² guideline) ^b	Applicable to locations meeting the above criterion but ≤25 m ² with significantly elevated concentrations of radionuclides	Concentration limits for application to "hot spots" varying in size as follows: (m ²) (pCi/g) ^c <1 50 1- <3 30 3- <10 15 10-25 10

^aFrom Reference 3.^b"Every reasonable effort shall be made to identify and remove any source which has a concentration exceeding 30 times the guideline value, irrespective of area."^cThese guideline values are applicable to surface concentrations of ²³²Th, ²³⁰Th, ²²⁸Ra, and ²²⁶Ra only; for other radionuclides and subsurface values, see Reference 3.**Table 2. Background radiation levels for the northern New Jersey area**

Type of radiation measurement or sample	Radiation level or radionuclide concentration
Concentration of radionuclides in soil (pCi/g)	
²³² Th	0.9 ^a
²³⁸ U	0.9 ^a
²²⁶ Ra	0.9 ^a

^aReference 4.

Table 3. Concentrations of radionuclides in soil at 266 East Spring Valley Avenue, Hackensack, New Jersey (MJ024)

Sample ^a	Depth (cm)	Radionuclide concentration (pCi/g)	
		²²⁶ Ra ^b	²³² Th ^b
<i>Systematic samples^c</i>			
S1A	0–15	0.95 ± 0.06	0.97 ± 0.4
S1B	15–30	0.74 ± 0.1	0.86 ± 0.09
S1C	30–45	1.0 ± 0.1	1.0 ± 0.1
S1D	45–60	1.1 ± 0.07	1.1 ± 0.2
S2A	0–15	1.3 ± 0.1	3.5 ± 0.4
S2B	15–30	1.2 ± 0.02	1.3 ± 0.04
<i>Biased samples^d</i>			
B1A	0–15	1.6 ± 0.07	5.7 ± 0.3
B1B	15–30	1.3 ± 0.06	2.2 ± 0.2
B1C	30–45	1.3 ± 0.2	1.3 ± 0.2
B2A	0–15	1.5 ± 0.1	6.8 ± 0.5
B2B	15–30	0.75 ± 0.02	1.6 ± 0.04

^aLocations of soil samples are shown on Fig. 2.

^bIndicated counting error is at the 95% confidence level ($\pm 2\sigma$).

^cSystematic samples are taken at grid locations irrespective of gamma exposure.

^dBiased samples are taken from areas shown to have elevated gamma exposure rates.

INTERNAL DISTRIBUTION

1. B. A. Berven	12. L. M. Floyd
2. R. F. Carrier	13. S. V. Kaye
3. W. D. Cottrell	14. P. T. Owen
4. A. G. Croff	15-17. R. E. Swaja
5. J. W. Crutcher	18. J. K. Williams
6. J. T. Ensminger	19. IR&A Publications Office
7-11. R. D. Foley	20. Laboratory Records - RC

EXTERNAL DISTRIBUTION

- 21. J. D. Berger, Oak Ridge Associated Universities, P. O. Box 117, Oak Ridge, TN 37831
- 22. R. W. Doane, Eberline, Inc., 800 Oak Ridge Turnpike, Oak Ridge, TN 37831
- 23. J. J. Fiore, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
- 24-26. P. J. Gross, U.S. Department of Energy, P. O. Box E, Oak Ridge, TN 37831
- 27-29. G. K. Hovey, Bechtel National, Inc., 800 Oak Ridge Turnpike, Oak Ridge, TN 37831
- 30. L. R. Levis, Roy F. Weston, Inc., 20030 Century Blvd., Germantown, MD 20874
- 31. G. P. Turi, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
- 32. J. W. Wagoner, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
- 33-35. Andrew Wallo III, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
- 36. Office of Assistant Manager, Energy Research and Development, Oak Ridge Operations Office, Oak Ridge, TN 37831
- 37-38. Office of Scientific and Technical Information, DOE, Oak Ridge, TN 37831