INTERSTITIAL SITE OCCUPATION IN $ZrNiH^*$

D. G. Westlake, H. Shaked**, P. R. Mason†, B. R. McCart†† and M. H. Mueller

Materials Science Division
Argonne National Laboratory
Argonne, Illinois 60439 (USA)

and

T. Matsumoto and M. Amano
National Research Institute for Metals, Tokyo. 153 (JAPAN)

The submitted manuscript has been authored by a contractor of the U. S. Government under contract No. W-31-109-ENG-38. Accordingly, the U. S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U. S. Government purposes.

DECEMBER 1981

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

*Work supported by the U.S. Department of Energy.

**Permanent address: Nuclear Research Center-Negev, POB 9001 and Ben-Gurion University of the Negev, POB 653, Beer Sheva, Israel

†Permanent address: Rose Hulman Institute of Technology, Dept. of Physics, 5500 Wabash Avenue, Terre Haute, IN 47903

††Permanent address: Dept. of Physics, Augustana College, Rock Island, IL 61201

To be submitted to International Symposium on the Properties and Applications of Metal Hydrides-II, Toba, Japan, May 30-June 4, 1982. Proceedings to be published in The Journal of Less-Common Metals.

MASTER

NOTICE

PORTIONS OF THIS REPORT ARE ILLLEGIBLE. It has been reproduced from the best available copy to permit the broadest possible availability.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

INTERSTITIAL SITE OCCUPATION IN ZrNiH *

D. G. Westlake, H. Shaked**, P. R. Mason† B. R. McCart†† and M. H. Mueller

Materials Science Division,
Argonne National Laboratory
Argonne, Illinois 60439 (USA)

and

T. Matsumoto and M. Amano
National Research Institute for Metals, Tokyo 153 (Japan)

ABSTRACT

In-situ x-ray diffraction measurements during step-wise hydrogenation of the intermetallic compound ZrNi have confirmed the existence of a stable compound ZrNiH . Neutron diffraction measurements on ZrNiD have allowed a determination of the site occupied by D. This site is tetrahedrally coordinated by four Zr atoms as predicted earlier from a model based solely on geometric considerations. Occupation of this site causes the lattice to distort from an orthorhombic to a triclinic structure. The trihydride is formed when two completely different types of sites are filled, to the exclusion of the sites occupied originally, causing reversion of the lattice to an orthorhombic structure. Other confirmed predictions of the geometric model are discussed.

*Work supported in part by the U.S. Department of Energy

**Permanent address:Nuclear Research Center-Negev, POB 9001 and Ben-Gurion University of the Negev, POB 653, Beer Sheva, Israel

†Permanent address:Rose-Hulman Institute of Technology, Dept. of Physics, 5500 Wabash Avenue, Terre Haute, IN 47803

††Permanent address:Dept. of Physics, Augustana College, Rock Island, IL 61201

INTERSTITIAL SITE OCCUPATION IN ZrNiH^*

D. G. Westlake, H. Shaked**, P. R. Mason†, B. R. McCart†† and M. H. Mueller

Materials Science Division,
Argonne National Laboratory
Argonne, Illinois 60439 (USA)

and

T. Matsumoto and M. Amano
National Research Institute for Metals, Tokyo 153 (Japan)

Introduction

A model, based on hole sizes and H-H distances, was recently proposed by Westlake (1) to rationalize the observed (2-5) stoichiometry and H-site occupation in ZrNiH_3 . Further, he predicted that a stable compound ZrNiH should exist and that, in the monohydride, H would most probably be found in those sites tetrahedrally coordinated by four Zr atoms. Pressure-composition isotherms of the ZrNi-H system (2) gave some indication of the existence of a monohydride. The present collaborative investigation was undertaken to confirm or deny its existence by the use of "in-situ" x-ray diffraction and to test the H-site prediction (1) by the use of neutron diffraction.

Experimental

Westinghouse crystal bar Zr was rolled to 6mm diameter and given a zone refining treatment of four passes. The resulting rod contained 200 at. ppm Hf and only 8 at. ppm Fe. A Ni rod (Marz grade) of the same diameter

*Work supported in part by the U.S. Department of Energy

**Permanent address: Nuclear Research Center-Negev, POB 9001 and Ben-Gurion University of the Negev, POB 653, Beer Sheva, Israel

†Permanent address: Rose-Hulman Institute of Technology, Dept. of Physics, 5500 Wabash Avenue, Terre Haute, IN 47803

††Permanent address: Dept. of Physics, Augustana College, Rock Island, IL 61201

was obtained from Materials Research Corporation. The intermetallic compound ZrNi was prepared by arc-melting with care to achieve homogeneity. Combining analyses of the two metals, we conclude that the interstitial impurity content of the intermetallic compound could not exceed 2000 at. ppm.

ZrNiD was prepared at ANL by reaction of the ZrNi with a measured quantity of D₂ gas while cooling from 1075 K to 375 K. Purified D₂, obtained by thermal decomposition of UD₃, was used in this reaction. We determined gravimetrically that the desired stoichiometry had been achieved to within less than 1%. This sample was powdered under an argon atmosphere and sealed in a vanadium capsule for neutron diffraction experiments, which were performed with the pulsed neutron source ZING P' (6) at ANL.

At NIRM, another sample of ZrNi, from the same arc-melted button, was reacted with gaseous H₂ at 533 K and cooled to room temperature. The specimen was pulverized under an argon atmosphere. The resulting powder (under 250 mesh) was dehydrogenated and residual stress was relieved by heating at 1025 K in a vacuum of 1.3×10^{-4} Pa. An anneal at 575 K had failed to remove all of the hydrogen, and even an anneal at 775 K did not completely relieve the residual strain. The powdered and stress-relieved sample of ZrNi was mounted on a specially designed sample holder in an argon atmosphere. The "in-situ" x-ray diffraction measurements, with $\lambda = 1.54178$ Å, were made under various H₂ pressures at 533 K. Measurements at room temperature were not possible because the equilibrium H₂ pressure for ZrNiH is too low; pressures lower than 6.6 kPa could not be controlled properly in the autoclave, which has been described elsewhere (7).

Results and Crystallography

A. Lattices of the Hydrides of ZrNi

The pulverized specimen of ZrNi was studied with x-rays at several steps

of its hydrogenation. The spectra corresponding to ZrNi , ZrNiH and ZrNiH_3 are shown in Fig. 1, and the respective H_2 pressures are given. At a pressure of 0.152 MPa, the patterns of both ZrNiH and ZrNiH_3 were apparent, but between 0.223 and 5.0 MPa only the pattern of ZrNiH_3 could be detected. For both ZrNiH and ZrNiH_3 , changing the pressure caused virtually no changes in positions of the lines. The spectra of ZrNi and of ZrNiH_3 were indexed according to orthorhombic lattices (Table I) in agreement with previously published results (3,4,8). The ZrNiH spectrum, on the other hand, exhibited splitting of the orthorhombic lines, which indicates that it has a triclinic lattice.

The specimen of ZrNiD was studied by neutron scattering. Part of the observed time-of-flight spectrum is shown in Fig. 2. The neutron spectrum of ZrNiD and the X-ray spectrum of ZrNiH were indexed according to a triclinic lattice (Table I). In both spectra, reflections of residual ZrNi(H) or ZrNi(D) (i.e., solid solution of H or D in α ZrNi) were observed and indexed according to the orthorhombic lattice (Table I). The resolution of the x-ray spectrum was insufficient to allow any conclusion regarding an isotope effect on the volume expansion of the lattice due to solute H(D).

B. The Positions of D Atoms in ZrNiD

In a previous publication (1), five inequivalent interstitial sites in ZrNi were considered on the basis of hole size and H-H distance for possible occupancy by hydrogen (Fig. 3). In the present work, the structure factors for neutron scattering by ZrNiD were calculated for the five alternative sites of the D atoms (Table II). C_{2v} symmetry (9) was assumed for ZrNiD in these calculations. This assumption is justified as an approximation, because the triclinic distortion is not large and the split lines (e.g., 130, $1\bar{3}0$) are of approximately equal intensities. The jF^2 were calculated (from the structure

factors of Table II) for the five alternative sites and compared with the corresponding observed jF^2 (Table III): Obviously, there is good agreement with the calculated jF^2 for the first alternative and extremely poor agreement for the other four alternatives. Variations of the position parameter within a physically meaningful range cannot make any of these four alternatives agree with the observed data. The conclusion of the present study is, therefore, that the D atoms occupy the first alternative site (Table II), as predicted (1).

Discussion

By obtaining x-ray diffraction spectra at 533 K under various H_2 pressures, we have unequivocally established the existence of a stable phase, $ZrNiH$, that can coexist either with the solid solution of H in $ZrNi$ or with $ZrNiH_3$. Three other observations of the present study are also in accord with the predictions made earlier (1):

First, there were no shifts in positions of x-ray peaks as a function of pressure for either $ZrNiH$ or $ZrNiH_3$, which seems to indicate very low (or no) solubility for H in both these phases. The reported (8) observation of $ZrNiH_{3.4}$ at $T = 298$ K with $P_{H_2} = 0.4$ MPa cannot be attributed, therefore, to a hyperstoichiometric trihydride. Furthermore, one of the present experiments contradicts this report of $ZrNiH_{3.4}$. With $T = 296$ K and $P_{H_2} = 5$ MPa, our observed lattice parameters did not exceed those expected for $ZrNiH_3$. It seems no longer necessary, therefore, to postulate (1) a substoichiometric $ZrNiH_4$ to explain the reported (8), but unsubstantiated, $ZrNiH_{3.4}$.

Second, the dihydride, $ZrNiH_2$, does not exist.

Finally, D was found by neutron diffraction to occupy all the interstitial sites in the lattice of $ZrNiD$ that are tetrahedrally coordinated by four Zr atoms.

The lattice of NiZr is shown schematically in Fig. 3. It is known, now, that all sites of the type shown as diamonds are occupied by D in ZrNiD. This set of sites has the largest hole radius, $0.6\text{\AA}$, and the greatest H-H distance, $2.76\text{ \AA}$. Analysis of the combined x-ray and neutron results has revealed that occupation of this site by H(D) causes a distortion of the lattice to a triclinic structure.

During transformation of the monohydride to the trihydride, both kinds of sites represented by triangles (both base-up and base-down) in Fig. 3 become filled (4,5). This cannot be accomplished without emptying all the "diamond" sites, because, otherwise, the H-H distances would be far too short (1). The present results confirm that when the "diamond" sites are emptied and both sets of sites represented by triangles are filled the lattice reverts to an orthorhombic structure.

Conclusion

X-ray and neutron diffraction experiments confirm the predictions of a model (1) based on geometric considerations regarding both stoichiometries and sites occupied by H in hydrides of the intermetallic compound ZrNi. Application of this simple model to other compounds with different structures and subsequent experimental testing of the results would seem to be warranted.

References

1. D. G. Westlake, J. Less Comm. Met., 75 (1980) 177.
2. G. G. Libowitz, H. F. Hayes and T.R.P. Gibb, Jr., J. Phys. Chem., 62 (1958) 76.
3. W. L. Korst, J. Phys. Chem., 66 (1961) 370.
4. S. W. Peterson, V. N. Sadana, and W. L. Korst, J. Phys., 25 (1964) 451.
5. A. V. Irodova, V. A. Somenkov, S. Sh. Shil'shtein, L. N. Padurets and A. A. Chertkov, Sov. Phys. - Crystallogr., 23 (1978) 591.
6. J. D. Jorgensen and F. J. Rotella, J. Appl. Cryst. in press 1981. See also G. A. Lager, F. K. Rose, F. J. Rotella and J. D. Jorgensen, J. Appl. Cryst., 14 (1981) 137.
7. T. Hirata, T. Matsumoto, M. Amano and Y. Sasaki, J. Phys. F: Met. Phys., 11 (1981) 521.
8. J. L. Baron, A. Virost and J. Delaplace, J. Nucl. Mat., 83 (1979) 286.
9. F. M. Henry and K. Lonsdale (editors), International Tables for X-ray Crystallography, The Kynoch Press (1952).

FIGURE CAPTIONS

- Fig. 1. X-ray ($\text{CuK}\alpha$) spectra taken at 533 K. a) in vacuum, b) hydrogen pressure = 0.113 MPa, c) hydrogen pressure = 0.223 MPa. The reflections marked * belong to the sample holder. The reflections marked α belong to residual solid solution of H in ZrNi, ZrNi(H).
- Fig. 2. Part of the time of flight spectrum of neutrons scattered by ZrNiD. Scattering angle is 160° and the counts were accumulated over 1.64×10^6 pulses.
- Fig. 3. Orthorhombic lattice of ZrNi. Zr, $\bigcirc$; Ni, $\square$. Examples of the interstitial sites corresponding to the numbers 1 through 5 in Table II are shown as $\blacklozenge$, $\blacktriangle$, $\blacktriangledown$, $\bullet$ and $\star$, respectively. Pentagonal symbols denote face centers.

Table I. Lattice parameters for the hydrides of ZrNi

COMPOUND	$a_0(\text{\AA})$	$b_0(\text{\AA})$	$c_0(\text{\AA})$	VOLUME ($\text{\AA}^3$)
	$[\alpha(^{\circ})]$	$[\beta(^{\circ})]$	$[\gamma(^{\circ})]$	
ZrNi	3.272 ± 0.005	9.965 ± 0.005	4.115 ± 0.005	134.2 ± 0.4
ZrNi(H) } ZrNi(D) }	3.28 ± 0.01	10.12 ± 0.01	4.05 ± 0.01	134.6 ± 0.7
ZrNiH } ZrNiD }	3.367 ± 0.002 [89.18 ± 0.03]	10.313 ± 0.004 [89.60 ± 0.05]	4.063 ± 0.002 [89.43 ± 0.03]	141.1 ± 0.2
ZrNiH ₃	3.53 ± 0.01	10.48 ± 0.02	4.30 ± 0.02	159.1 ± 1.5

Table II. Input for the calculation of the structure factor of ZrNiD for the five alternative sites of D

ELEMENT	b	ALTERN	SITE	x	y	z	F
Zr	0.70	-	4c	0	-0.140	0.25	$4b\cos 2\pi(ky + \frac{\ell}{4})$
Ni	1.03	-	4c	0	-0.429	0.25	
		1	4c	0	0.2152	0.25	
		2	4c	0	0.067	0.25	$4b\cos 2\pi(ky + \frac{\ell}{4})\cos 2\pi\ell(z - \frac{1}{4})$
D	0.67	3	8f	0	-0.312	0.509	
		4	8f	0	0.4283	0.3833	
		5	8e	0.2275	0	0	$4b\cos \pi \frac{\ell}{2} \cos 2\pi \ell x$

Table III. Calculated and observed structure factors of ZrNiD for the five alternative sites of D. A is a scale factor such that $AjF_{131}^2 = 1000$.

hkℓ	j	A _j F ²					OBSERVED
		C A L C U L A T E D					
		1	2	3	4	5	
110	4	16	17	148	727	42	45 ± 8
021	4	27	384	2	35	4	
040	2	14	454	101	326	6	20 ± 5
111	8	31	1026	491	765	539	35 ± 5
130	4	227	427	15	608	159	285 ± 20
041	4	80	3790	462	423	437	135 ± 15
131	8	1000	1000	1000	1000	1000	1000 ± 30
002	2	412	2839	154	853	803	440 ± 20
022	4	1	941	267	139	392	0 ± 10
112	8	33	34	29	220	84	50 ± 20
200	2	412	2930	785	1777	165	360 ± 20
A		2.233	15.90	4.259	9.643	4.354	0.3189

Fig. 1. X-ray (CuK α) spectra taken at 533 K. a) in vacuum, b) hydrogen pressure = 0.113 MPa, c) hydrogen pressure = 0.223 MPa. The reflections marked * belong to the sample holder. The reflections marked α belong to residual solid solution of H in ZrNi, ZrNi(H).

Fig. 2. Part of the time of flight spectrum of neutrons scattered by ZrNiD. Scattering angle is 160° and the counts were accumulated over 1.64×10^6 pulses.

Fig. 3. Orthorhombic lattice of ZrNi. Zr, $\bigcirc$; Ni, $\square$. Examples of the interstitial sites corresponding to the numbers 1 through 5 in Table II are shown as $\blacklozenge$, $\blacktriangle$, $\blacktriangledown$, $\bullet$ and $\star$, respectively. Pentagonal symbols denote face centers.