Conf-830418--2

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Presented at the International Workshop on
Gross Properties of Nuclei and Nuclear Excitation,
Hirschegg, Austria, January 17-22, 1983

DISCOVERY OF BETA-DELAYED TWO-PROTON
RADIOACTIVITY IN ^{22}Al AND ^{26}P

M.D. Cable, J. Honkanen, R.F. Parry, S.H. Lu,
Z.Y. Zhou, and Joseph Cerny

January 1983

MASTER

Discovery of Beta-delayed Two-proton Radioactivity in ^{22}Al and $^{26}\text{p}^*$

M. D. Cable, J. Honkanen[†], R. F. Parry, S. H. Zhou[‡], Z. Y. Zhou^{*},
and Joseph Cerny

Department of Chemistry and Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720, USA

January 1983

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE**PORTIONS OF THIS REPORT ARE ILLEGIBLE.**

It has been reproduced from the best available copy to permit the broadest possible availability.

*This work was supported by the Director, U. S. Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics, and by Nuclear Sciences of the Basic Energy Sciences Program of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

Discovery of Beta-delayed Two-proton Radioactivity in ^{22}Al and $^{26}\text{p}^*$

M. D. Cable, J. Honkanen[†], R. F. Parry, S. H. Zhou[‡], Z. Y. Zhou^{*},
and Joseph Cerny

Department of Chemistry and Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720, USA

Among the highly proton-rich nuclides are candidates for two-proton radioactivity, originally proposed by Gol'danskii¹⁾, as well as candidates for beta-delayed two-proton radioactivity. Gol'danskii has also discussed the latter decay mode²⁾ and has suggested a number of nuclides that might provide its initial examples, among them the light, odd-odd, $T_z = -2$ isotopes. The recent discovery³⁾ of the first known member of this series, ^{22}Al , and the subsequent observation⁴⁾ of the next higher member, ^{26}p (both as beta-delayed, isospin-forbidden, single proton emitters), made these two nuclides ideal candidates for a search for this new mode of decay. These two nuclides have now been observed also to decay by beta-delayed two-proton radioactivity (more detailed results on ^{22}Al have been submitted for publication⁵⁾).

$^{22}\text{Al}(\tau_{1/2} \sim 70 \text{ ms})$ and $^{26}\text{p}(\tau_{1/2} \sim 20 \text{ ms})$ were produced via the ($^3\text{He}, p4n$) reaction on ^{24}Mg and ^{28}Si targets, respectively, using 110 MeV $^3\text{He}^{+2}$ beams of 3-7 μA from the 88-inch Cyclotron at the Lawrence Berkeley Laboratory: the effective cross section for their observation as high energy (7-8 MeV), beta-delayed single proton emitters was $\sim$ a few nb. A helium jet system and a high geometry, three-element particle telescope ($\Delta E1, \Delta E2, E$) were used to search for two-proton decay events. Symmetrically divided, circular " $\Delta E1$ " (24 μm for ^{22}Al , 14 μm for ^{26}p) and " $\Delta E2$ " (155-170 μm) detectors were employed to observe coincident, low-energy protons with each telescope subtending 4.5% of 4π sr. A standard 500 μm E detector placed behind and in coincidence with either of these two-counter telescopes observed the high energy groups from beta-delayed single proton decay as a monitor of the reaction.

Figure 1(a) presents the two-dimensional proton-proton coincidence spectrum obtained following the decay of ^{22}Al and figure 1(b) shows the summed proton energy spectrum (with peaks at 4.139 ± 0.020 MeV and 5.636 ± 0.020 MeV). Although the exact corresponding center of mass

XBL 82B-4028

Fig. 2. Proposed partial decay scheme for ^{22}Al . Fig. 1. (a). A proton-proton coincidence spectrum following the beta decay of ^{22}Al (E_β vs. $E_{\beta'}$). (b). A summed energy spectrum for the two-proton coincidences in (a): ($E_\beta + E_{\beta'}$).

energies depend on the mechanism of two-proton emission, these peaks can be shown to correspond to transitions from the $T=2$ analog state in ^{22}Mg (fed by the superallowed beta-decay of ^{22}Al) to the ground and first excited state of ^{20}Ne . Figure 2 illustrates the various observed decay modes of ^{22}Al ; all open particle-decay channels from the $^{22}\text{Mg}(4+)$, $T=2$ state are isospin forbidden, making this state relatively narrow.

Two of the possible mechanisms available for the emission of two protons in this decay include A) single step ^2He emission^{1,2)} (a proton pair coupled to a 1S_0 configuration) and B) a sequential two-step process proceeding through an intermediate state (or states) in ^{21}Na . Considering the stronger decay branch, ^2He emission to the ^{20}Ne (1.634 MeV) state should occur within relative laboratory angles of $\leq 40^\circ$ (see ref. 5). Since the telescope pair could detect angles of $0-70^\circ$, it could observe coincident protons arising from either mechanism. ^2He emission should show a distribution centered about $E_p^L = E_p^R$ with a shape determined by the detector configuration and the final state interaction. Sequential emission would also be (trivially) symmetric about $E_p^L = E_p^R$ but would be expected to show distinct proton groups corresponding to transitions through the intermediate state(s) in ^{21}Na . The data of figure 1(a), with their limited statistics, cannot conclusively establish the mechanism; the observed variation in energies and yields could result from either a ^2He type distribution or sequential decay through several states in ^{21}Na or both. See reference 5 for a more complete discussion.

Differences from the above arising in the possible beta-delayed two-proton decay mechanism of ^{26}P , however, can help clarify our general understanding of these results. ^{26}P should have a 3^+ , $T=2$ ground state, so that, following its superallowed beta-decay to the 3^+ , $T=2$ state in ^{26}Si , only sequential proton decay to the ^{24}Mg ground state is allowed (spin-parity conservation forbids ^2He emission). A decay scheme for ^{26}P is presented in figure 3. Figure 4(a) shows the summed, coincident two-proton energies arising from the decay of nuclides produced in the 110 MeV ^3He bombardment of ^{28}Si : both ^{26}P and ^{22}Al are observed. By setting a gate on the transitions corresponding to ^{26}P decay to the ^{24}Mg ground state (via ^{26}Si), one obtains the proton-proton coincidence spectrum shown in figure 4(b). Strong evidence for sequential decay is observed: the great majority of the coincident events

correspond to a single decay channel involving a 3.7 MeV proton associated with a 1.2 MeV proton (preliminary values). Further experiments on the decay of ^{26}P are in progress: its weaker decay to the ^{24}Mg first excited state can occur by both decay mechanisms and so is of substantial interest for comparative purposes.

Additional studies on both ^{22}Al and ^{26}P are necessary to obtain A) higher statistics with this detector configuration and B) other angular correlation data to observe the changes in yield, peak shape, and laboratory energy of the emitted coincident protons. Although the beta-delayed two-proton decay of ^{26}P to the ^{24}Mg ground state appears clearly to occur by the expected sequential proton emission, the two-proton decay mechanism(s) following the beta-decay of ^{22}Al is uncertain. Should ^{22}Al decay have a component of ^2He emission, fascinating studies of a new form of radioactivity would become possible.

Footnotes and References

*This work was supported by the Director, OER, Div. Nucl. Physics of OHEMP, and by Nucl. Sciences of BES of the U.S. Department of Energy under contract DE-AC03-76SF00098.

[†]On leave from: Dept. of Physics, University of Jyväskylä, Finland.

*On leave from: Institute of Atomic Energy, Beijing, China.

*On leave from: Dept. of Physics, Nanking University, China.

1. V. I. Gol'danskii, Usp. Fiz. Nauk 87, 255 (1965); Sov. Phys. Uspekhi 8, 770 (1966).
2. V. I. Gol'danskii, Pisma Zh. Eksp. Teor. Fiz 32, 572 (1980); JETP Lett. 32, 554 (1980)
3. M. D. Cable, J. Honkanen, R. F. Parry, H. M. Thierens, J. M. Wouters, Z. Y. Zhou, and J. Cerny, Phys. Rev. C26, 1778 (1982).
4. M. D. Cable, J. Honkanen, R. F. Parry, S. H. Zhou, Z. Y. Zhou, and J. Cerny, Lawrence Berkeley Laboratory Rept. No. LBL-15454, to be published.
5. M. D. Cable, J. Honkanen, R. F. Parry, S. H. Zhou, Z. Y. Zhou and J. Cerny, Lawrence Berkeley Laboratory Rept. No. LBL-15453, to be published.