NOTICE

PORTIONS OF THIS REPORT ARE ILLISIBLE.

It has been reproduced from the best available copy to permit the broadest possible availability.

LBL-18314

CONF-840794-3

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

**Accelerator & Fusion
Research Division**

Presented at the NRL-NSWC Workshop on Local and
Global Methods of Dynamics, Silver Spring, MD,
July 23-26, 1984

**THE COVARIANT LIE-TRANSFORMED PLASMA
ACTION PRINCIPLE**

A.N. Kaufman

September 1984

The Covariant Lie-Transformed Plasma Action Principle*

Allan N. Kaufman

Physics Department and Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

September 1984

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

12/1/84

- * This research was supported by the Office of Energy Research of the U.S. Department of Energy.

Handwritten signature

THE COVARIANT LIE-TRANSFORMED PLASMA ACTION PRINCIPLE

Allan N. Kaufman
 Physics Department and Lawrence Berkeley Laboratory
 University of California
 Berkeley, CA 94720

The Lie Transform is a systematic technique [1] for obtaining the nonlinear quasi-static effects of high-frequency phenomena. We consider, in particular, a plasma of charged particles in a self-consistent electromagnetic wave. Each particle oscillates about an "oscillation center" which undergoes non-oscillatory motion [2]. We obtain the so-called "ponderomotive" Hamiltonian for the oscillation center, in relativistically covariant form, and demonstrate, via the action principle, that it determines the dielectric susceptibility and wave propagation.

We begin with the (eight) canonical variables $z = (r, p)$, which satisfy the covariant Hamiltonian equations:

$$dr^\mu/d\tau = \partial H/\partial p_\mu, \quad dp_\mu/d\tau = -\partial H/\partial r^\mu, \quad (1)$$

with the invariant Hamiltonian function $H(z; A)$, a functional of the Maxwell potential $A(x)$. The correct particle evolution equations are obtained by the choice ($c=1$):

$$H = (p - eA(r))^2/2m, \quad (2)$$

noting that

$$A_\mu(r) = \int d^4x \delta^4(x-r) A_\mu(x). \quad (3)$$

Consider the family of all phase-space trajectories $z(\tau)$, each parameterized (smoothly) by its proper time τ . The seven-dimensional time-like surface $\tau = 0$ will be called the "initial-condition" surface. We introduce arbitrary coordinates η on this surface, and let $g(\eta)d^7\eta$ denote the number of particles (of a given species) in $d^7\eta$. With the trajectory $z(\tau; \eta)$ considered as an eight-dimensional field on the eight-dimensional phase-space (τ, η) [it's actually a mapping of phase space onto itself], we construct the action functional for the system:

$$S[z(\tau, \eta), A(x)] = \int d^7\eta g(\eta) \int d\tau [p_\mu(\tau, \eta) dr^\mu(\tau, \eta)/d\tau - H(p, r; A)] \\ - \int d^4x F_{\mu\nu} F^{\mu\nu}/16\pi, \quad (4)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the Maxwell field.

Variation of S with respect to $z(\tau; \eta)$ yields the Hamiltonian equations (1).

while variation with respect to $A(x)$ yields the Maxwell equation

$$F^{\mu\nu},_{\nu} = 4\pi j^{\mu}, \quad (5)$$

with

$$j^{\mu}(x) = \int d^7n \, g(n) \int d\tau \, j^{\mu}(x; z) \quad (6)$$

and

$$j^{\mu}(x; z) = -\delta H(z)/\delta A_{\mu}(x). \quad (7)$$

Evaluation of (7), by (2), yields

$$j^{\mu}(x; z) = (e/m) \delta^4(x-r) (p^{\mu} - eA^{\mu}). \quad (8)$$

We now define the (invariant) Vlasov distribution

$$f(\bar{z}) = \int d^7n \, g(n) \int d\tau \, \delta^8(\bar{z} - z(\tau, n)), \quad (9)$$

and derive the Ignatiev equation [3],

$$\{f(z), H(z)\} = 0, \quad (10)$$

in terms of the covariant canonical Poisson bracket:

$$\{a, b\} = (\partial a / \partial r^{\mu})(\partial b / \partial p_{\mu}) - (\partial a / \partial p_{\mu})(\partial b / \partial r^{\mu}) \quad (11)$$

To derive (10), we introduce the intermediate distribution

$$\bar{g}(\bar{z}; \tau) = \int d^7n \, g(n) \delta^8(\bar{z} - z(\tau, n)) \quad (12)$$

and obtain, by (1),

$$\partial \bar{g} / \partial \tau = -\{\bar{g}, H\}. \quad (13)$$

Integration over τ then yields (10), since $g = 0$ at infinite τ .

We now restrict the Maxwell potential to represent a single wave of eikonal form:

$$A_{\mu}(x) = \bar{A}_{\mu}(x) \exp i\theta(x)/\epsilon + c.c. \quad (14)$$

where the amplitude $\bar{A}$ and the gradient of the phase $k_{\mu}(x) = \partial_{\mu} \theta(x)$ are slowly varying fields. We substitute (14) into (2), and then invoke the Lie Transform to eliminate rapidly oscillating terms.

The new Hamiltonian K is related to the old H by the formula [1]

$$K = [\exp i \{w, \cdot\}] H, \quad (15)$$

where $w(z)$ is a suitably chosen generating function. Expanding both H and K in powers of the amplitude $\bar{A}$, we have $K^{(0)} = H^{(0)}$, and

$$K^{(1)} = H^{(1)} + \{w, H^{(0)}\}. \quad (16)$$

We can require $K^{(1)}$ to vanish, if we ignore the problems of resonant denominators. Solving (16) for $w(z)$, and proceeding to second order, we obtain $K^{(2)}(z)$, which we denote by $\Psi(z; A)$, the relativistically invariant ponderomotive potential:

$$\Psi(z; A) = \int d^4x A_\mu^*(x) \Psi_\mu^v(x; z) A^v(x); \quad (17)$$

where

$$\Psi_\mu^v(x; z) = \delta^4(x-r) (e^2/m) (p \cdot k)^{-2} [k^2 p^\mu p_\nu + \delta_\nu^\mu (k \cdot p)^2 - (k \cdot p) (p^\mu k_\nu + k^\mu p_\nu)]. \quad (18)$$

This may be expressed more concisely as

$$\Psi = m |d\tilde{u}/d\tau|^2 / (d\Theta(r)/d\tau)^2, \quad (19)$$

and reduces, in the rest-frame, to the familiar

$$\Psi = e^2 |\tilde{E}|^2 / m\omega^2. \quad (20)$$

Invariance of the phase-space Lagrangian under a canonical transformation:

$$\int d\tau [p_\mu dr^\mu/d\tau - H(p, r)] = \int d\tau [\bar{p}_\mu d\bar{r}^\mu/d\tau - K(\bar{p}, \bar{r})], \quad (21)$$

where the overbar denotes oscillation-center variables, converts the action to

$$S = \int d^7\eta g(\eta) \int d\tau [\bar{p}_\mu d\bar{r}^\mu/d\tau - \bar{p}^2/2m - \Psi(z; A)] - \int d^4x F_{\mu\nu}^* F^{\mu\nu}/8\pi. \quad (22)$$

Defining the invariant oscillation-center distribution $F(z) = \int d^7\eta g(\eta) \int d\tau \delta^8(z - \tilde{z}(\tau, \eta))$, we obtain, in analogy to the steps leading to (10), the corresponding Ignatiev equation:

$$\{F(z), p^2/2m + \Psi(z; A)\} = 0. \quad (23)$$

In order to vary S with respect to $\tilde{A}_\mu(x)$ and $\Theta(x)$, we first substitute (17) into (22), obtaining (for the terms bilinear in A)

$$S^{(2)} = \int d^4x \tilde{A}_\mu^*(x) D_\nu^\mu(x; F) \tilde{A}^\nu(x), \quad (24)$$

where the dielectric matrix D_ν^μ is the sum of its vacuum part

$$D_{\text{vac}}^\mu{}_\nu(k(x)) = (k^\mu k_\nu - k^2 \delta_\nu^\mu)/4\pi \quad (25)$$

and the susceptibility

$$\chi_\nu^\mu(x; F) = - \int d^8z F(z) \Psi_\nu^\mu(x; z). \quad (26)$$

The relation (26) is the "K-X theorem" [4], which is seen to be the essential ingredient of the action functional, coupling $F(z)$ to $A_\mu(x)$.

We now express the dielectric matrix in terms of its local eigenvalues $D_\alpha(x)$ and eigenvectors $e_\alpha(x)$:

$$D_\nu^\mu(x) = \sum_\alpha D_\alpha(x) e_\alpha^\mu(x) e_\alpha^{*\nu}(x), \quad (27)$$

whence

$$S^{(2)} = \int d^4x D_\alpha(x) |A_\alpha(x)|^2, \quad (28)$$

with $A_\alpha(x) = e^{\alpha_\nu} \star(x) \tilde{A}^\nu(x)$, the projection of A on the α eigenvector.

We now vary $S^{(2)}$ with respect to $A_\alpha(x)$, obtaining the eikonal equation for the phase:

$$D_\alpha(x, k = \partial\theta/\partial x) = 0, \quad (29)$$

associated with polarization e_α . This yields the covariant ray equations:

$$dx^\mu/d\sigma = -\partial D_\alpha/\partial k_\mu, \quad dk_\mu/d\sigma = \partial D_\alpha/\partial x^\mu. \quad (30)$$

Variation with respect to $\theta(x)$ yields the wave-action conservation law [5] $\partial J^\mu(x)/\partial x^\mu = 0$, where the wave-action density four-vector is

$$J^\mu(x) = -A_\alpha(x)^2 \partial D_\alpha(x, k)/\partial k_\mu \quad (31)$$

Since the eigenvalues $D_\alpha(x)$ are functionals of the oscillation-center distribution $F(z)$, by (26), we have thus obtained a closed self-consistent set of coupled equations for $F(z)$ and the wave amplitude and phase.

D. D. Holm was instrumental in the formulation of the principles presented here, at the Aspen Center for Physics. This research was supported by the Office of Energy Research of the U.S. Department of Energy.

Reference

- [1] J. R. Cary, Physics Reports 79 (1981) 131.
- [2a] R. L. Dewar, Phys. Fluids 16 (1973) 1102.
- [2b] R. L. Dewar, Aust. J. Phys. 30 (1977) 533.
- [3] Yu. G. Ignatiev, Zh. Eksp. Teor. Fiz. 81 (1981) 3 [Sov. Phys. JETP 54 (1981) 1].
- [4a] J. R. Cary and A. N. Kaufman, Phys. Rev. Lett. 39 (1977) 402.
- [4b] S. Johnston and A. N. Kaufman, Phys. Rev. Lett. 40 (1978) 1266.
- [4c] J. R. Cary and A. N. Kaufman, Phys. Fluids 24 (1981) 1238.
- [4d] A. N. Kaufman and B. M. Boghosian, Contemporary Math 28 (1984) 169.
- [5] R. L. Dewar, Ap. J. 174 (1972) 301.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.