53
4-27-93 JSD

CONF 920963--2

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2893
UC-426

PPPL-2893

X-RAY DIAGNOSTICS OF TOKAMAK PLASMAS

BY

M. BITTER, H. HSUAN, K.W. HILL AND M. ZARNSTORFF

APRIL, 1993

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced from the best available copy.
Available in paper copy and microfiche.

Number of pages in this report: 34

DOE and DOE contractors can obtain copies of this report from:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
(615) 576-8401.

This report is publicly available from the:

National Technical Information Service
Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4650

X-Ray Diagnostics of Tokamak Plasmas

M. Bitter, H. Hsuan, K. W. Hill, and M. Zarnstorff

*Princeton University
Plasma Physics Laboratory
Princeton, NJ 08543*

*Presented at the 4th International Colloquium on Atomic Spectra and Oscillator
Strengths for Astrophysical and Laboratory Plasmas, Septmeber 14 - 17, 1992
National Institute of Standards and Technology (NIST)
Gaithersburg , Md 20899 ,USA*

MASTER

REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

Abstract

The paper describes the X-ray diagnostics of central plasma parameters of present large tokamak fusion experiments by means of Johann type curved crystal spectrometers. These spectrometers have a very high spectral resolution, $\lambda/\Delta\lambda > 10000$, and are mainly used to determine the central ion temperature from Doppler broadening measurements of the 1s-2p resonance lines of high-Z heliumlike ions. The full diagnostic potential of the spectra of heliumlike ions is demonstrated by the detailed analysis of FeXXV satellite spectra from a particular high temperature tokamak discharge with reactor-like conditions. This analysis is based on least squares fit comparisons of the experimental data with synthetic spectra, which are constructed from theoretical predictions. It is found that the satellite spectra which are associated with the heliumlike resonance lines are in excellent agreement with the theoretical predictions. The satellite spectra can therefore be used for electron temperature measurements, measurements of the dielectronic recombination rate coefficients and the determination of relative ion charge-state abundances. Discrepancies between experiment and theory are, however, observed for the intensity ratios of the heliumlike lines. Also discussed are atomic physics data needs and the potential of the X-ray diagnostic for future large tokamaks, such as ITER.

I. Introduction

X-ray diagnostics of tokamak plasmas have become very important during the last decade due to the fact that these plasmas have achieved fusion-reactorlike experimental conditions with high temperatures of the order of 10 keV. Heliumlike ions of high-Z elements can be produced in the hot core of these plasmas, and spectra of ArXVII^{1,2}, TiXXI³⁻⁵, CrXXIII^{6,7}, FeXXV^{8,9} and NiXXVII¹⁰⁻¹³ are now widely used to determine the central plasma parameters. The main diagnostic application is the measurement of the central ion temperature, $T_i(0)$, from the Doppler widths of the emitted X-ray lines, which requires crystal spectrometers of very high spectral resolution, $\lambda/\Delta\lambda \geq 10000$. The obtained high-resolution spectra are well suited for a detailed comparison with theoretical predictions due to the fact that tokamak plasmas are in steady state for durations of several seconds and are well-diagnosed by independent measurements of plasma parameters, such as the electron density and the electron temperature. Since the spectra of heliumlike ions are also used for the study of solar flares, very accurate theoretical calculations of these spectra have been performed for the elements with $Z \leq 45$.^{3, 14-19} The spectra consist of the heliumlike lines w: $1s^2\ ^1S_0 - 1s2p\ ^1P_1$, x: $1s^2\ ^1S_0 - 1s2p\ ^3P_2$, y: $1s^2\ ^1S_0 - 1s2p\ ^3P_1$, and z: $1s^2\ ^1S_0 - 1s2s\ ^3S_1$ and the associated dielectronic and 'collisionally excited' satellites which are due to lithiumlike transitions of the type $1s^2nl - 1s2pnl$ with $n \geq 2$. The intensity ratios of the satellites and the resonance line w provide important additional diagnostic information. They can be used for measurement of the central electron temperature and determination of the ion charge state distribution. We point out that the experimental verification of theoretical predictions for the spectra of the heliumlike ions is important to establish a solid base for the theoretical calculations of the more complex ions. It is the purpose of this paper to review the atomic physics results from present tokamak experiments and to discuss the prospects of X-ray diagnostics on future large tokamaks.

The paper is organized as follows: Section II describes the experimental parameters and X-ray crystal spectrometers of the Tokamak Fusion Test Reactor (TFTR), which is one of the present large tokamak experiments. Section III presents the analysis of two spectra of heliumlike iron from a particular TFTR discharge to demonstrate the importance of heliumlike spectra for the diagnostic of tokamak plasma parameters and to show a comparison of the observed spectra with theoretical predictions. In section IV we review atomic physics results from present tokamak experiments, and in section V we discuss the potential of x-ray spectroscopy for the diagnostics of future large tokamaks. Conclusions are presented in section VI.

II. Crystal Spectrometers on Tokamak Experiments

The objective of the present large tokamak experiments is to study deuterium and deuterium-tritium plasmas under reactorlike conditions. The major tokamak experiments are the Joint European Torus (JET), the Japanese Tokamak JT-60, the French Tore Supra, and the US Tokamak

Fusion Test Reactor (TFTR). A schematic (one half of a vertical cross-section) of TFTR is shown in Fig. 1. The plasma is magnetically confined by toroidal and poloidal fields, and it is ohmically heated by an induced toroidal current which is in the range from 0.5 to 2.2 MA. Since ohmic heating is not efficient at high temperatures, auxiliary plasma heating is provided by the injection of 12 energetic (120 keV) neutral deuterium beams with a total power up to 35 MW and by radio frequency (ion cyclotron) heating with input powers up to 12 MW. Table I lists the experimental parameters of TFTR. The plasma contains small amounts (10^{-5} - 10^{-3} N_e , where N_e is the electron density) of metal impurities, such as Ti, Cr, Fe and Ni, which are ionized to different charge states. The radiation emitted from these metal impurity ions at the higher concentrations can represent a significant fraction of the total radiative power loss from the plasma. Therefore, every effort is made to reduce the amount of these impurities. On the other hand, the line radiation of these metal ions provides invaluable diagnostic information.

Figure 1 also shows the experimental arrangement of six crystal spectrometers which are used for Doppler broadening measurements.^{20,21} These spectrometers are Johann type curved crystal spectrometers with a high spectral resolution, $\lambda/\Delta\lambda \geq 10000$. The crystal curvature radii are in the range from 3.8 to 12 m. The spectrometers have one horizontal sightline in the midplane and five nearly vertical sightlines at different major radii to measure radial profiles of the ion temperature. The detectors have to be shielded against neutron and gamma radiation. The Johann configuration offers the advantages that (1) the heliumlike lines and the associated satellites can be simultaneously observed and (2) each spectral line is reflected from the entire crystal. Thus, since the resonance line and the satellites are observed under the same experimental conditions, their intensity ratios can be determined very accurately. The statistical errors of the spectral data are small due to the high throughput of the instruments.

Although plasma diagnostics generally rely on indigenous metal impurities, such as Ti, Cr, Fe and Ni, other elements have been introduced into tokamak plasmas by laser evaporation or by gas puffs. These techniques were used by the TFR group to study the Z-dependent wavelength shifts and satellite line intensities in the spectra of the heliumlike ions. Figure 2 presents spectra of argon, scandium, vanadium, chromium and manganese, obtained from these experiments.²² The spectra show a characteristic charge-dependent wavelength shift which is ascribed to the Z^4 dependence of the spin orbit interaction energy. Further documentation of these effects has been obtained from TFTR experiments by observation of the heliumlike spectra of titanium, chromium, iron and nickel.²¹ We note that the spectral features are well separated in the heliumlike spectra of ions with $Z = 24 - 26$, and that these spectra are therefore most appropriate for diagnostic applications. Figure 3 represents the fractional ion charge state abundances of iron as a function of the electron temperature for coronal equilibrium conditions.²³ Heliumlike iron is the dominant charge state for electron temperatures in the range from 1.5 to 7 keV and is therefore well suited for the diagnostics of central TFTR plasma parameters.

III. Analysis of the satellite spectra of heliumlike ions

In this section we describe the detailed analysis of two FeXXV spectra which were obtained from a TFTR deuterium plasma with ohmic heating of 1 MW and auxiliary heating of 20 MW by injection of neutral deuterium beams. The analysis is based on a detailed comparison of the observed spectra with synthetic spectra which were constructed from theoretical predictions.^{15,18} The data were recorded by the TFTR horizontal crystal spectrometer, which collects 128 FeXXV satellite spectra per discharge with a time resolution of 50 ms. Figure 4 shows the ion temperature results from Doppler broadening measurements of the 1s-2p resonance line. The ion temperature increased from 2.5 to 28 keV during the period of the deuterium beam injection from 3 to 4 sec. Figure 5 presents the observed time evolution of the central electron temperature and electron density from measurements of the electron cyclotron emission and by far-infrared laser interferometry, respectively. Both the ions and the electrons were heated very efficiently for the given experimental conditions. This type of tokamak discharge also provides high yields ($\sim 5 \times 10^{16} \text{ sec}^{-1}$) of 2.5 MeV neutrons from D-D fusion reactions and will most likely be used for the final D-T experiments on TFTR.

Figures 6 and 7 present two observed FeXXV satellite spectra and synthetic spectra, which represent least squares fits of theoretical predictions to the experimental data. The observed spectra are accumulated spectra covering periods of steady state conditions, i. e. the time of ohmic heating from 2.5 to 3 sec and the time from 3.5 to 4 sec with the injection of deuterium beams, when the parameters n_e , T_e , and T_i , had again reached steady state conditions (see Figs. 4 and 5). Due to the accumulation of data over these extended periods, the statistical errors are substantially reduced. The synthetic spectra consist of the heliumlike lines w, x, y and z, and the lithiumlike (and berylliumlike) satellites due to transitions $1s^2nl - 1s2pnl$ with $n \geq 2$. The main spectral features have been identified by key letters following Gabriel's notation¹⁴ and are explained in Table II. The heliumlike lines are mainly produced by electron impact excitation from the heliumlike ground state, whereas the satellites can be produced in the process of dielectronic recombination of the heliumlike ions and/or by electron impact (inner-shell) excitation from the ground state of the lithiumlike (and berylliumlike) ions. The relative contributions from these two processes are different for different satellites. The satellites k and j are produced by dielectronic recombination, whereas the so-called 'collisional' satellites q, r, and t with transitions involving the lithiumlike ground state, have strong contributions from the process of inner-shell excitation. In addition to the features listed in Table II, the synthetic spectra also include numerous $n \geq 3$ satellites which are of pure dielectronic nature.

Expressions for the emissivities, ϵ , of the different spectral components are given in an abbreviated form below (see Reference 15 for a detailed description):

(1) *heliumlike lines* (w, x, y, z):

$$\epsilon_{w,x,y,z} = \langle \sigma v \rangle_{w,x,y,z} N_e N_{He}$$

(2) *'collisional' satellites* :

$$\epsilon_{c.s.} = \langle \sigma v \rangle_{c.s.} N_e N_{Li} \text{ (or } N_{Be})$$

(3) *'dielectronic' satellites*:

$$\epsilon_{d.s.} = F_1(T_e) F_2(s) N_e N_{He}$$

N_e , N_{He} , N_{Li} and N_{Be} are, respectively, the electron density and the densities of the heliumlike, lithiumlike and berylliumlike charge states. $\langle \sigma v \rangle$ is the excitation rate coefficient for electron impact excitation for the case of a Maxwellian electron energy distribution. $F_1(T_e) = (2\pi h/mkT_e)^{3/2} \exp(-E_s/T_e)$, where E_s is the energy difference between the excited lithiumlike state, $1s2pnl$, and the $1s^2$ heliumlike ground state. $F_2(s) = g_s/g_1 A_r^{sf} A_a^{si} (\sum A_a^{si'} + \sum A_r^{sf'})$ is the satellite line strength; g_s and g_1 are the statistical weights of the excited lithiumlike state and the heliumlike ground state, respectively; and A_a^{si} and A_r^{sf} are the probabilities for autoionization and radiative transitions from the $1s2pnl$ state.

It follows from the expressions (1), (2) and (3), that the intensity ratios of the 'collisional' satellites and the heliumlike resonance line w are directly proportional to the relative abundances, N_{Li}/N_{He} and N_{Be}/N_{He} , since $\langle \sigma v \rangle_w$ and $\langle \sigma v \rangle_{c.s.}$ have approximately the same T_e dependence¹⁵, and that the intensity ratios of the 'dielectronic' satellites and the resonance line w are only a function of the electron temperature. The synthetic spectra depend therefore on only four parameters: T_e , N_{Li}/N_{He} , N_{Be}/N_{He} , and the ion temperature T_i , which determines the width of the spectral lines. The synthetic spectra shown in Figs. 6 and 7 represent least squares fits of the theoretical predictions to the experimental data with respect to these four parameters. The results are listed in Table III. The details of the least squares fitting procedure, which involves several steps, have been described earlier.¹³

For an experimental verification of the theory it is necessary to compare the parameter values of T_e , N_{Li}/N_{He} and N_{Be}/N_{He} in Table III with results from independent electron-temperature measurements and coronal-equilibrium calculations. In the spectrum shown in Fig. 6, the contributions from the various spectral components are clearly distinguishable due to the fact that the satellite intensities are appreciable for electron temperatures of 3 keV and that the Doppler width for an ion temperature of 2.5 keV is smaller than the separation of the main spectral features. The errors obtained for the parameters T_e , N_{Li}/N_{He} and N_{Be}/N_{He} from the least squares fit method are therefore small. The value for the electron temperature, $T_e = 3.24 \pm 0.04$ keV, in Table III is in good agreement with the electron temperature measurements during the ohmic heating phase (see Fig. 5). However, the relative abundances, $N_{Li}/N_{He} = 0.376 \pm 0.012$ and $N_{Be}/N_{He} = 0.144 \pm 0.013$, in Table III are larger than the coronal-equilibrium values $N_{Li}/N_{He} =$

0.186 and $N_{\text{Be}}/N_{\text{He}} = 0.022$ (see Fig. 3). The spectrum shown in Fig. 7 seems, at first sight, to have less structural detail than the spectrum shown in Fig. 6. This is due to the fact that the satellite intensities are very much reduced for electron temperatures of 8 keV and that the spectral features are smeared out as a result of the large Doppler broadening obtained for ion temperatures of 20 keV. The least squares fit method made it nevertheless possible to detect the very small intensity contributions of the dielectronic satellites and to derive from these contributions an electron temperature value of $T_e = 8.2 \text{ keV} \pm 0.4 \text{ keV}$, which is in excellent agreement with the central electron temperature obtained from the laser Thomson scattering diagnostic (see Fig. 8). This agreement indicates that the observed radiation originates mainly from the hot core of the plasma, even though the data represent chord-integrated measurements along a horizontal sightline through the plasma (see Fig. 1). We may therefore conclude that the measured ion temperature value is the central ion temperature $T_i(0)$. It is evident from Fig. 7, that the 'collisional' satellites are much stronger than the dielectronic satellites. In fact, the relative abundances, $N_{\text{Li}}/N_{\text{He}} = 0.260 \pm 0.008$ and $N_{\text{Be}}/N_{\text{He}} = 0.018 \pm 0.005$, in Table III are much larger than the coronal-equilibrium values, $N_{\text{Li}}/N_{\text{He}} = 0.047$ and $N_{\text{Be}}/N_{\text{He}} < 0.01$ for an electron temperature of 8 keV. This enhancement of $N_{\text{Li}}/N_{\text{He}}$ and $N_{\text{Be}}/N_{\text{He}}$ is ascribed to charge-exchange recombination of the FeXXV and FeXXIV ions with the injected deuterium atoms. The intensity ratios, x/w , y/w and z/w , of the heliumlike lines are found to be larger than the ratios obtained from Eq. (1) which only accounts for the direct excitation from the heliumlike ground state.¹⁵ In our least squares fit procedure, we introduced the quantities x_a , y_a and z_a , which measure the enhancement of x/w , y/w and z/w over the predictions from Eq. (1). The observed ratios of the heliumlike lines will be further discussed in the next section.

IV Review of experimental Results.

In this section we review the atomic physics results from the X-ray spectroscopy of tokamak plasmas and discuss the contributions from the different spectral components separately.

A. *Measurements of Dielectronic recombination rate coefficients.*

Data on dielectronic recombination rate coefficients are needed for plasma modeling calculations to determine the ionization equilibria of high-Z ions in solar flares and in fusion plasmas. However, it is very difficult to measure these rate coefficients. Standard techniques which use crossed-beam methods suffer from the lack of suitable ion beams and are limited to low-ionization states. Data on rate coefficients for high-Z ions were therefore mainly obtained from plasma experiments by indirect methods based on observations of the line radiation from successive charge states. The interpretation of these experimental data depends on extensive modeling of the charge state distribution, so that the uncertainties for the derived rate coefficients are often large. Only recently, due to the development of Electron Beam Ion Traps (EBIT), has it become possible to perform reliable measurements of rate coefficients of high-Z ions, such as

NiXXVII.²⁴ It is thus fortunate that the dielectronic satellite spectra of the 1s-2p resonance lines offer an opportunity to determine the dielectronic rate coefficients for high-Z heliumlike and hydrogenlike ions. The method, which was first proposed by Bely-Dubau et al.²⁵, is independent of plasma modeling and involves only the single-charge state of interest. Following Bely-Dubau et al., the total dielectronic recombination rate coefficient associated with the 1s - 2p core transition of heliumlike ions can be obtained from measurements of the sum of the intensity ratios of the dielectronic satellites and the resonance line, $\Sigma I_d/I_w$, using the expression

$$(4) \quad \alpha_d = \langle \sigma v \rangle_w \Sigma I_d/I_w$$

where the collisional excitation rate coefficient for the resonance line w, $\langle \sigma v \rangle_w$, is assumed to be known. Figure 9 shows the results for the dielectronic recombination rate coefficients of FeXXV, which were obtained by this method from tokamak experiments.²⁶ The data are in good agreement with the theory of Bely-Dubau et al.^{15,25}, as one may expect from the discussion of the dielectronic satellite spectra given in the previous section. Further results on the dielectronic rate coefficients of heliumlike and hydrogenlike Ti, Cr and Ni are given in Reference 27. We note that experimental results on the dielectronic recombination of heliumlike nickel NiXXVII from tokamak experiments are in good agreement with those obtained by Knapp et al..²⁴

B Comparison of the ion charge state distributions with coronal equilibrium calculations

As discussed in the previous section, it is possible to determine the relative abundances, N_{Li}/N_{He} and N_{Be}/N_{He} , of the heliumlike, lithiumlike and berylliumlike charge states from the intensity ratios of the 'collisional' satellites and the resonance line w. If radial ion transport is negligible, these relative ion abundances are expected to be in agreement with coronal equilibrium. This is due to the fact that the densities in tokamak plasmas, which are of the order of $N_e \sim 10^{13} - 10^{14} \text{ cm}^{-3}$, correspond to the so-called zero density limit for ions with $Z > 20$, so that the high-Z ions are in their ground states. Moreover, the ionization and recombination times are small compared with the times of steady state conditions, so that the charge state distribution reaches an ionization equilibrium. Effects of radial ion transport can, however, lead to a shift of the ionization equilibrium in the hot core of tokamak plasmas towards lower charge states. These effects can thus be determined by measuring the deviations from coronal equilibrium. A quantitative analysis requires, in general, detailed modeling calculations which include radial profile effects of the electron temperature and density and which must primarily be based on reliable coronal equilibrium predictions. Improved coronal equilibrium predictions for the ion charge states of nickel have recently been provided by Zastrow et al.¹² and have been compared with experimental data from JET¹² and TFTR.¹³ The JET data were obtained from a great number of discharges with very different electron density and electron temperature profiles. These data were analyzed by

comparing the observed line intensities with modeling calculations of chord-integrated line emissivities along the spectrometer sightline through the plasma. It was found that the observed line intensities were within the bounds of the coronal equilibrium predictions for the given range of radial profiles. On the other hand, the TFTR data were obtained from a special series of ohmically heated helium plasmas with very flat radial profiles of the electron density and electron temperature. The observed spectra were therefore essentially emitted from plasmas with well-defined single values of the electron density and electron temperature. Thus, it was justified to neglect radial profile effects and to compare the observed spectra with synthetic spectra for single values of the parameters T_e , T_i , N_{Li}/N_{He} and N_{Be}/N_{He} , applying the least squares fit analysis described in the previous section. The results obtained from the TFTR experiments are shown in Figs 10 and 11. Due to the least squares fit analysis, it is possible to determine the relative charge state abundances very accurately with error bars. We note that the deduced abundance ratios are larger than the coronal equilibrium predictions. As shown in Figs 10 and 11 by data points for three different electron temperatures, a more involved analysis which includes chord-integrals assuming a coronal equilibrium distribution for the ion charge state distribution cannot explain the observed deviations. Based on the present coronal equilibrium calculations, particle diffusion coefficients, D , in the range from 1 to 2.5 m²/s are needed to explain the TFTR results.

C *Intensity ratios of the heliumlike lines*

The R-Ratio

$$R(ne) = z / (x+y) ,$$

and G-Ratio,

$$G(Te) = (x + y + z)/w ,$$

of the heliumlike lines w , x , y and z have important diagnostic applications for electron density measurements²⁸ in solar flares^{29,30} and in laser-produced high-density plasmas, as well as for measurements of non-thermal electron energy distributions in tokamak plasmas with auxiliary RF heating.³¹⁻³³ However, with exception of the measurements of SiXXIII³⁴ and MgXI³⁵ from DITE, the data from tokamak experiments for the heliumlike line ratios are not in agreement with the theoretical predictions.^{5,8,12,13,36} As shown by the analysis of the two FeXXV spectra in section III, the ratios x/w , y/w and z/w are found to be larger (see Table III) than the predictions for direct electron impact excitation, which is the principal excitation mechanism. Additional processes, such as the excitation of the lines w , x , y , and z by recombination of hydrogenlike ions and inner-shell ionization of lithiumlike ions, which contributes to the line z , cannot, in general, explain the observed ratios. In fact, the largest deviations occur for low electron temperatures,^{5,13} when the concentration of hydrogenlike ions is negligible. Similar deviations from the theoretical predictions have recently been found for the satellite spectrum of CrXXIII³⁷ from the Frascati

Tokamak as shown in Fig. 12. The experimental data are compared with a synthetic spectrum (solid line) based on the theoretical predictions of J. Dubau for the dielectronic satellites and data of R. Mewe and J. Schrijver³⁸ for the relative intensities of the heliumlike lines w, x, y, and z. The electron and ion temperatures were $T_e = 2.3$ keV and $T_i = 2.0$ keV. The predictions³⁸ for the intensities of w, x, y, and z are represented by the hatched areas. The synthetic spectrum also includes contributions to the line z from the inner-shell ionization of lithiumlike ions, which were calculated for the density ratio $N_{CrXXII} / N_{CrXXIII} = 0.45$ derived from the intensity ratio q/w . We point out that recent measurements of the TiXXI line ratios on EBIT were in good agreement with theory.³⁹ Thus, the tokamak results for the heliumlike lines are still puzzling and need further investigation.

V. X-ray spectroscopy on future large tokamaks

In this section we discuss the potential diagnostic use of X-ray spectroscopy on future large tokamaks, such as ITER, the International Thermonuclear Experimental Reactor. The objectives of ITER are (1) to demonstrate controlled ignition and extended burn of D-T plasmas, (2) to validate the design concepts and engineering components for a fusion power reactor, and (3) to serve as a test facility for neutronics, blanket modules, tritium production and advanced technologies.⁴⁰ The conceptual design parameters of ITER are listed in Table IV. Comparing these parameters with those of present-day large tokamaks (see Table I), we note that crucial plasma parameters, such as the electron density and the ion temperature, $n_e = 2 \times 10^{20} \text{ m}^{-3}$ and $T_i = 10 - 30$ keV, are already obtained in the present experiments, but that other ITER parameters, such as the plasma volume (500 m^3), the stored energy (600 MJ), the energy yield from fusion products (~ 1 GW), the radiated power (~ 100 MW) and the pulse lengths (> 1000 sec), are by orders of magnitude larger than the parameters of the present tokamaks. A major problem for the operation of ITER will be the control of the plasma-wall interaction to maintain the required plasma purity of $Z_{\text{eff}} = 1.7$ ($Z_{\text{eff}} = 1$ corresponds to a pure deuterium-tritium plasma). The solution of this problem depends not only on a choice of appropriate wall and divertor materials, but also on the control of the energy release from the plasma. To control the energy release from the plasma and to reduce the heat load on the divertor under quasi-continuous operation with pulse lengths greater than 1000 sec, it has recently been proposed⁴¹ to inject high-Z impurities into the plasma which can provide radiative cooling of the plasma edge region at a rate of 100 MW. Thus, contrary to present-day tokamaks, where every effort is made to reduce the level of plasma impurities, ITER may require the injection of impurities for its operation. The most likely candidate will be krypton ($Z = 36$), since it is chemically inert and since it can be easily introduced in controllable quantities. Krypton can solve two problems for ITER: (1) the radiation from the low krypton ion charge states can provide the required radiative cooling and (2) the charge state of heliumlike krypton, which will be the dominant state of ionization in the hot core of the plasma for the projected central electron temperatures of 30 keV, can be used for Doppler broadening measurements to determine the central

ion temperature $T_i(0)$. We point out that ion temperature measurements by the techniques of charge exchange recombination spectroscopy, which are now widely used on tokamaks and which require the injection of neutral hydrogen beams, may not be applicable since neutral deuterium beams will not penetrate to the core of the plasma. However, the development of efficient high-resolution crystal spectrometers for Doppler broadening measurements at wavelengths of $0.95 \text{ \AA}$ will be an experimental challenge. Furthermore, it is important to establish a reliable atomic physics data base on high-Z elements, such as krypton, in order to predict by plasma simulation calculations the effects of radiative cooling for the different experimental conditions on ITER. The atomic data base for the ITER experiment must include other elements, which may be constituents of the plasma facing components like carbon, iron, nickel, molybdenum, tungsten, beryllium, and boron or contaminants like oxygen. Especially important are data on rate coefficients for ionization, recombination and line excitation for all the ion charge states of these elements. Obviously, atomic physics contributions are essential to solve the many critical problems related to the control of the energy release from the plasma, the diagnostics of plasma parameters, issues of the plasma-wall interaction and the removal of the 'helium ash' due to DT fusion reactions from the plasma. Experiments are now under way, on TFTR and EBIT, to study the spectra of the different krypton ion charge states and to investigate the effects of krypton on the plasma performance.

VI. Conclusion

High-resolution crystal spectrometers have been very important for the X-ray diagnostics of central plasma parameters in present tokamak fusion experiments. In particular, the Doppler broadening and Doppler shift measurements of the $1s-2p$ resonance lines of high-Z heliumlike ions have been invaluable for the determination of the central ion temperature and the central plasma motion in tokamak experiments with auxiliary heating by neutral beam injection and ion cyclotron heating. In fact, these measurements were essential for the discovery and characterization of new tokamak operating regimes with enhanced particle and energy confinement. Since tokamak plasmas are well-diagnosed by independent measurements of the electron temperature and the electron density, high-resolution spectra of heliumlike ions obtained from these plasmas are well suited for a detailed comparison with atomic theories. It is found that the spectra of dielectronic and 'collisional' satellites which are associated with the heliumlike resonance lines are accurately described by only four parameters, T_i , T_e , N_{Li}/N_{He} and N_{Be}/N_{He} , in excellent agreement with the theoretical predictions. The intensity ratios of the dielectronic satellites and the heliumlike resonance line can therefore be used for electron temperature measurements and measurements of the dielectronic recombination rate coefficients of heliumlike ions. The relative ion charge-state abundances, N_{Li}/N_{He} and N_{Be}/N_{He} , which are obtained from the intensity ratios of the 'collisional' satellites and the resonance line, are larger than the predictions for coronal equilibrium. The observed deviations can be ascribed to effects of radial ion transport in tokamak plasmas; however, uncertainties in the theoretical data for ionization and recombination rate coefficients,

which are used for the coronal equilibrium calculations, are still a concern for a quantitative determination of the transport coefficients. The ratios of the heliumlike lines x/w , y/w and z/w are larger than predicted by the processes of direct excitation, recombination of hydrogenlike ions, and inner-shell ionization of lithiumlike ions. Heliumlike ions of elements with still higher Z , like krypton, will be important for the diagnostic of the central plasma parameters in future large tokamaks, such as ITER. The development of efficient high-resolution crystal spectrometers for Doppler broadening measurements at wavelengths $\lambda \leq 1 \text{ \AA}$ will be an experimental challenge.

Acknowledgements. We gratefully acknowledge fruitful discussions with R. Barnsley, F. Bombarda and P. Platz, who also provided us with spectral data from the DITE, the FTU and the TFR tokamak experiments. We thank S. Cohen for valuable information on the ITER related physics issues. We also gratefully acknowledge the continuing support of R. J. Hawryluk, and K. M. Young as well as the technical assistance of J. Gorman, R. Such and the TFTR operating crew. This work was supported by the U.S. Department of Energy, Contract No. DE-AC02-76-CHO-3073.

Table I: Parameters of the Tokamak Fusion Test Reactor (TFTR) Experiments

Plasma Major Radius, R	2.65 m
Plasma Minor Radius, a	0.95 m
Toroidal Magnetic Field, B_T	5.0 Tesla
Plasma Current, I_p	0.6-3.0 MA
Auxiliary Heating:	
Neutral Beam Heating, NBI	35 MW
Ion Cyclotron Heating, ICH	12 MW
Electron Density, n_e	$0.1-3 \times 10^{20} \text{ m}^{-3}$
Electron Temperature, T_e	1-10 keV
Ion Temperature, T_i	1-35 keV
Stored Energy, E	1-5 MJ
D-D neutron yields	$\leq 6 \times 10^{16} \text{ sec}^{-1}$
Pulse Length	6 sec
$n \tau T_i$ ($10^{20} \text{ m}^{-3} \text{ sec keV}$)	4.3

TABLE II: Theoretical wavelengths for the heliumlike transitions w, x, y and z of FeXXV and the associated main $n = 2$ satellites. Also listed are the satellite line strengths, $F_2(s)$.

Key	Transition	Wavelength (Å)	$F_2(s)$ (10^{13} s^{-1})
w	$1s^2 \text{ } ^1S_0 - 1s2p \text{ } ^1P_1$	1.8498	—
x	$1s^2 \text{ } ^1S_0 - 1s2p \text{ } ^3P_2$	1.85503	—
t	$1s^2 2s \text{ } ^2S_{1/2} - 1s2p2s(^3P)^2P_{1/2}$	1.8566	10.9
y	$1s^2 \text{ } ^1S_0 - 1s2p \text{ } ^3P_1$	1.8590	—
q	$1s^2 2s \text{ } ^2S_{1/2} - 1s2p2s(^1P)^2P_{3/2}$	1.8605	0.003
k	$1s^2 2p \text{ } ^2P_{1/2} - 1s2p^2 \text{ } ^2D_{3/2}$	1.8626	35.0
r	$1s^2 2s \text{ } ^2S_{1/2} - 1s2p2s(^1P)^2P_{1/2}$	1.8631	6.73
j	$1s^2 2p \text{ } ^2P_{3/2} - 1s2p^2 \text{ } ^2D_{5/2}$	1.8654	51.3
z	$1s^2 \text{ } ^1S_0 - 1s2s \text{ } ^3S_1$	1.8676	—
β	$1s^2 2s^2 \text{ } ^1S_0 - 1s2p2s^2 \text{ } ^1P_1$	1.87003	0.86

Table III: Parameter values obtained from least squares fits of synthetic spectra to the observed spectra shown in Figs. 6 and 7, for the case with ohmic heating (Column A) and the case with additional neutral beam heating (Column B).

Parameters	A	B
T_i (keV)	3.2 ± 0.1	20.7 ± 0.3
T_e (keV)	3.29 ± 0.04	8.2 ± 0.4
n_{Li}/n_{He}	0.376 ± 0.012	0.260 ± 0.008
n_{Be}/n_{He}	0.144 ± 0.013	0.018 ± 0.005
xa	1.59 ± 0.05	2.49 ± 0.08
ya	1.61 ± 0.04	1.80 ± 0.04
za	1.65 ± 0.04	2.29 ± 0.06

Table IV: Conceptual Design Parameters of the International Thermonuclear
Experimental Reactor (ITER)

Plasma Major Radius, R	6.0 m
Plasma Minor Radius, a	2.15 m
Toroidal Magnetic Field, B_T	4.85 Tesla
Plasma Current, I_p	22 MA
Auxiliary Heating:	115 MW
Electron Density, n_e	$2 \times 10^{20} \text{ m}^{-3}$
Z_{eff}	1.7
Electron Temperature, T_e	10 - 30 keV
Ion Temperature, T_i	10 - 30 keV
Stored Thermal Energy, E	600 MJ
Fusion Products, P_{fusion}	1000 MW
Alpha Particles, P_a	200 MW
Radiated Power, P_{rad}	100 MW
Pulse Length	400 - 2500 sec
$n \tau T_i$ ($10^{21} \text{ m}^{-3} \text{ sec keV}$)	4 - 8

Figure Captions

Figure 1: Schematic of the Tokamak Fusion Test Reactor (TFTR) showing one half of a vertical cross-section. The dotted lines represent the vertical axis of the torus and the horizontal midplane. Also shown is the arrangement of the TFTR Horizontal and Vertical X-ray Crystal Spectrometers for Doppler broadening measurements of the ion temperature.

Figure 2: Satellite spectra of heliumlike ions from Reference 22.

Figure 3: Fractional abundances of iron ion charge states as a function of the electron temperature for coronal equilibrium conditions after Reference 23.

Figure 4: Ion temperature results from Doppler broadening measurements of the FeXXV 1s-2p resonance line w . The data were recorded from a TFTR deuterium plasma with 20 MW of auxiliary heating by deuterium neutral beam injection (NBI) from 3 to 4 sec using the TFTR Horizontal Crystal Spectrometer.

Figure 5: Time evolutions of the electron temperature and electron density from the same discharge as in Fig. 4.

Figure 6: Satellite spectrum of FeXXV. The experimental data (circles) were recorded from the same discharge as in Fig. 4 during the period of ohmical heating from 2.5 to 3 sec. Also shown is a synthetic spectrum with its various spectral components (solid lines) which was constructed from a least squares fit of theoretical predictions of References 15 and 18 to the experimental data.

Figure 7: Satellite spectrum of FeXXV recorded from the same discharge as in Fig. 4 during the period of neutral beam injection from 3.5 to 4 sec. The synthetic spectrum was constructed as described in caption of Fig. 6.

Figure 8: Electron temperature profile measured by the laser Thomson scattering diagnostic at 3.5 sec from the same discharge as in Fig. 4.

Figure 9: Dielectronic recombination rate coefficient of FeXXV as a function of the electron temperature from Reference 26.

Figure 10: Comparison of the relative abundances of lithiumlike and heliumlike nickel with coronal equilibrium predictions from Reference 13.

Figure 11: Comparison of the relative abundances of berylliumlike and heliumlike nickel with coronal equilibrium predictions from Reference 13.

Figure 12: Satellite spectrum of CrXXIII from the Frascati tokamak experiment. The solid lines represent theoretical predictions from References 6 and 38.

References

- (1) E. Källne, J. Källne, A. Dalgarno, E. S. Marmar, J. E. Rice, and A. Pradhan, Phys. Rev. Lett. **52**, 2245 (1984)
- (2) TFR Group, F. Bombarda, F. Bely-Dubau, P. Faucher, M. Cornille, J. Dubau, and M. Loulergue, Phys. Rev. A **32**, 2374 (1985)
- (3) F. Bely-Dubau, P. Faucher, L. Steenman-Clark, M. Bitter, S. von Goeler, K. W. Hill, C. Camhy-Val, and J. Dubau, Phys. Rev. A **26**, 3459 (1982)
- (4) P. Lee, A. J. Lieber, and S. S. Wojtowicz, Phys. Rev. A **31**, 3996 (1985)
- (5) M. Bitter, K. W. Hill, M. Zarnstorff, S. von Goeler, R. Hulse, L. C. Johnson, N. R. Sauthoff, S. Sesnic, K. M. Young, M. Tavernier, F. Bely-Dubau, P. Faucher, M. Cornille, and J. Dubau, Phys. Rev. A **32**, 3011 (1985)
- (6) TFR group, J. Dubau, and M. Loulergue, J. Phys. B **15**, 1007 (1981)
- (7) M. L. Apicella, R. Bartiromo, F. Bombarda, and R. Giannella, Phys. Lett. **98A**, 174 (1983)
- (8) M. Bitter, K. W. Hill, N. R. Sauthoff, P. C. Efthimion, E. Meservey, W. Roney, S. von Goeler, R. Horton, and W. Stodiek, Phys. Rev. Lett. **43**, 129 (1979)
- (9) M. Bitter, S. von Goeler, K. W. Hill, R. Horton, D. Johnson, W. Roney, N. R. Sauthoff, E. Silver and W. Stodiek, Phys. Rev. Lett. **47**, 921 (1981)
- (10) H. Hsuan, M. Bitter, K. W. Hill, S. von Goeler, B. Grek, D. Johnson, L. C. Johnson, S. Sesnic, C. P. Bhalla, K. R. Karim, F. Bely-Dubau, and P. Faucher, Phys. Rev. A **35**, 4280 (1987)
- (11) F. Bombarda, R. Giannella, E. Källne, G. J. Tallents, F. Bely-Dubau, P. Faucher, M. Cornille, J. Dubau, and A. H. Gabriel, Phys. Rev. A **37**, 504 (1988)
- (12) K.-D. Zastrow, E. Källne, and H. P. Summers, Phys. Rev. A **41**, 1427 (1990)
- (13) M. Bitter, H. Hsuan, V. Decaux, B. Grek, K. W. Hill, R. Hulse, L. A. Kruegel, D. Johnson, S. von Goeler, and M. Zarnstorff, Phys. Rev. A **44**, 1796 (1991)

- (14) A. H. Gabriel, Mon. Not. R. Astron. Soc. **160**, 99 (1972)
- (15) F. Bely-Dubau, J. Dubau, P. Faucher, A. H. Gabriel, Mon. Not. R. Astron. Soc., **198**, 239 (1982)
- (16) L. A. Vainshtein and U. I. Safronova, At. Data Nucl. Data Tables **21**,49 (1978)
- (17) L. A. Vainshtein and U. I. Safronova, At. Data Nucl. Data Tables **25**,311 (1980)
- (18) U.I. Safronova, A. M. Urnov and L. A. Vainshtein, Proc. P. N. Lebedev Phys. Inst. [Acad. Sci. USSR] **119**, 13 (1980).
- (19) L. A. Vainshtein and U.I. Safronova, P. N. Lebedev Physical Institute, Moscow, and Institute of Spectroscopy, Troizk, USSR Academy of Science, Moscow, USSR, Rep. **3** (1985)
- (20) H. Hsuan, M. Bitter, J. E. Rice, K. W. Hill, L. Johnson, S. L. Liew, S. D. Scott and S. von Goeler, Rev. Sci. Instrum. **59**, 2127 (1988)
- (21) M. Bitter, H. Hsuan, J. E. Rice, K. W. Hill, M. Diesso, B. Grek, R. Hulse, D. W. Johnson, L. C. and S. von Goeler, Rev. Sci. Instrum. **59**, 2131 (1988)
- (22) TFR group, M. Cornille, J. Dubau, and M. Loulergue, Phys. Rev. A **32**, 3000 (1985)
- (23) C. Breton, C. De Michelis, M. Finkenthal, and M. Mاتيoli, Fontenay-aux-Roses Laboratory Report No. EUR-CEA-FC-948, 1978 (unpublished)
- (24) D. A. Knapp, R. E. Marrs, M. A. Levine, C. L. Bennett, M. H. Chen, J. R. Henderson, M. B. Schneider, and J. H. Scofield, Phys. Rev. Lett. **62**, 2104 (1989)
- (25) F. Bely-Dubau., A. H. Gabriel, and S. Volonté, Mon. Not. R. Astron. Soc. **186**, 405 (1979)
- (26) F. Bely-Dubau., M. Bitter, J. Dubau, P. Faucher, A. H. Gabriel, K. W. Hill, S. von Goeler, N. Sautjhoff, and S. Volonté, Phys. Lett. **93A**, 189 (1983)
- (27) M. Bitter, H. Hsuan, S. von Goeler K. W. Hill, B. Grek, and D. W. Johnson, Physica Scripta **T37**, 66 (1991)

- (28) R. W. P. McWhirter in *Course on Plasma Diagnostics and Data Acquisition Systems, Varenna - Villa Monastero - Italy, September 3-11, 1975*, edited by H. Eubank and E. Sindoni (Editrice Compositori - Bologna, 1975)
- (29) G. A. Linford and C. J. Wolfson, *Astrophys. J.* **331**, 1036 (1988)
- (30) D. L. McKenzie, *Astrophys. J.* **322**, 512 (1987)
- (31) P. Lee, A. J. Lieber, R. P. Chase, and A. K. Pradhan, *Phys. Rev. Lett.* **55**, 386 (1985)
- (32) P. Lee, A. J. Lieber, A. K. Pradhan, and Yueming Xu, *Phys. Rev. A* **34**, 3210 (1986)
- (33) R. Bartiromo, F. Bombarda and R. Giannella, *Phys. Rev. A* **32**, 531 (1985)
- (34) F. P. Keenan, S. M. McCann, R. Barnsley, J. Dunn, K. D. Evans, and N. J. Peacock, *Phys. Rev. A* **39**, 4092 (1989)
- (35) F. P. Keenan, S. M. McCann, A. E. Kingston, R. Barnsley, J. Dunn, and N. J. Peacock, *Phys. Rev. A* **44**, 3831 (1991)
- (36) Takako Kato, Shigeru Morita and Kuniaki Masai, *Phys. Rev. A.* **36**, 795 (1987)
- (37) F. Bombarda, private communication
- (38) R. Mewe and J. Schrijver, *Astr. Ap.* **87**, 55 (MSc) (1980)
- (39) S. Chantrenne, P. Beiersdorfer, R. Cauble, and M. B. Schneider, *Phys. Rev. Lett.* **69**, 265 (1992)
- (40) ITER Documentation Series, No. 23, *ITER Operations and Research Programme*, International Atomic Energy Agency, Vienna, 1991
- (41) C. J. Cummings, S. A. Cohen, R. Hulse, D. E. Post, and M. Redi, *Journal of Nuclear Materials* **176&177**, 916 (1990)

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 9

Fig. 10

Fig. 11

PPPL#93X0005

Fig. 12

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-Europeen, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Boudiqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sakaneke, Instituto Fisica, BRAZIL
 Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James., Univ. of Alberta, CANADA
 Dr. P. Lukáč, Komenského Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique., FRANCE
 C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinni, Univ. of Ioannina, GREECE
 Dr. T. Mui, Academy Bibliographic Ser., HONG KONG
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theo Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Director, Japan Atomic Energy Research Inst., JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Instit. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 S. Mori, Technical Advisor, JAERI, JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Uley, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Dr. O. Petrus, AL I CUZA Univ., ROMANIA
 Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 Dr. L. Stanflo, Univ. of UMEA, SWEDEN
 Library, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY
 Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

**DATE
FILMED**

6 / 30 / 93