Secondary Production of Benthic Insects in Three Cold-Desert Streams

W. L. Gaines

July 1987

**Prepared for the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830**

**Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government of any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, or Battelle Memorial Institute.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America
Available from
National Technical Information Service
United States Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161

NTIS Price Codes
Microfiche A01

Printed Copy

Pages	Price Codes
001-025	A02
026-050	A03
051-075	A04
076-100	A05
101-125	A06
126-150	A07
151-175	A08
176-200	A09
201-225	A010
226-250	A011
251-275	A012
276-300	A013

3 3679 00051 1172

SECONDARY PRODUCTION OF BENTHIC INSECTS
IN THREE COLD-DESERT STREAMS

W. L. Gaines

July 1987

Prepared for
the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Richland, Washington 99352

SUMMARY

I studied aquatic insect production in three cold-desert streams in eastern Washington (Douglas Creek, Snively Springs, and Rattlesnake Springs). The size-frequency method was applied to individual taxa to estimate total insect production. Production was also assessed for functional groups and trophic levels in each stream.

Optioservus sp. (riffle beetles) and Baetis sp. (mayflies) accounted for 72% of the total insect numbers and 50% of the total biomass in Douglas Creek. Baetis sp. accounted for 42% of the total insect numbers and 25% of the total biomass in Snively Springs. Simulium sp. (blackflies) and Baetis sp. comprised 74% of the total insect numbers and 55% of the total biomass in Rattlesnake Springs.

Grazer-scrappers (49%) and collectors (48%) were the most abundant functional groups in Douglas Creek. Collectors were the most abundant functional group in Snively Springs and Rattlesnake Springs. Herbivores and detritivores were the most abundant trophic level in Snively Springs and Rattlesnake Springs.

Dipterans (midges and blackflies) were the most productive taxa within the study streams, accounting for 40% to 70% of the total community production. Production by collectors and detritivores was the highest of all functional groups and trophic levels in all study streams.

Insects with rapid development times and multiple cohorts were very important in cold-desert streams, contributing significantly to the total community production. Total community production rates were generally higher than in northern temperate streams, lower than in southeastern blackwater streams, and much lower than in Sonoran Desert streams. Much of the production in cold-desert streams can be attributed to insects feeding upon detritus, and detritus was obtained by collecting (filtering and gathering). I suggest that production, not

density or biomass, is the most accurate and meaningful way to assess the roles of organisms in aquatic ecosystems, and that the procedure given by Benke et al. (1984) should be followed to calculate production.

ACKNOWLEDGMENTS

I am grateful to Dr. Stamford D. Smith and Dr. Colbert E. Cushing for their support and advice throughout this research project. I am especially grateful for their constructive comments on this manuscript. I would like to thank Dr. William Coffman for identifying the chironomids and Dr. Pat Schefter for identifying the caddisflies. Dr. Lee Rogers kindly provided space for laboratory and office work.

This research was originally presented in a master's thesis for Central Washington University. The research was performed at Pacific Northwest Laboratory during a Northwest College and University Association for Science (NORCUS) Fellowship (University of Washington), funded under Contract DE-AM06-76-RL02225. This work was supported by the U.S. Department of Energy under Contract DE-AC06-76RLO 1830. Pacific Northwest Laboratory is operated for the U.S. Department of Energy by Battelle Memorial Institute.

CONTENTS

SUMMARY	iii
ACKNOWLEDGMENTS	v
INTRODUCTION	1
LITERATURE REVIEW	3
STUDY AREA	5
DOUGLAS CREEK	5
SNIVELY SPRINGS	5
RATTLESNAKE SPRINGS	7
METHODS	9
RESULTS	11
INSECT DENSITY AND BIOMASS	11
Douglas Creek	11
Snively Springs	11
Rattlesnake Springs	11
FUNCTIONAL GROUP DENSITY AND BIOMASS	11
Douglas Creek	11
Snively Springs	13
Rattlesnake Springs	16
TROPIC LEVEL DENSITY AND BIOMASS	18
Douglas Creek	18
Snively Springs	19
Rattlesnake Springs	19
INSECT PRODUCTION	20
Douglas Creek	20

Snively Springs	24
Rattlesnake Springs	26
FUNCTIONAL GROUP PRODUCTION	27
TROPHIC LEVEL PRODUCTION	29
DISCUSSION	31
ESTIMATING SECONOARY PRODUCTION	31
COMPARISON BETWEEN STUDY STREAMS	31
Insect Faunas	31
Functional Groups and Trophic Levels	32
COMPARISON WITH OTHER STREAMS	33
CONCLUSIONS	34
REFERENCES	37

FIGURES

1	Temperature Regime for Douglas Creek	6
2	Temperature Regime for Snively Springs	7
3	Temperature Regime for Rattlesnake Springs	8

TABLES

1	Substrate Composition of Douglas Creek, Snively Springs, and Rattlesnake Springs	6
2	Annual Mean Density and Biomass of Insects from Douglas Creek	12
3	Annual Mean Density and Biomass of Insects from Snively Springs	13
4	Annual Mean Density and Biomass of Insects from Rattlesnake Springs	14
5	Results of Qualitative Gut Content Analysis and Subsequent Functional Group and Trophic Level Classification for Insects from Douglas Creek	15
6	Results of Qualitative Gut Content Analysis and Subsequent Functional Group and Trophic Level Classification of Insects from Snively Springs	16
7	Results of Qualitative Gut Content Analysis and Subsequent Functional Group and Trophic Level Classification of Insects from Rattlesnake Springs	17
8	Annual Mean Density and Biomass of Insect Functional Groups in Douglas Creek	18
9	Annual Mean Density and Biomass of Insect Functional Groups in Snively Springs	18
10	Annual Mean Density and Biomass of Insect Functional Groups in Rattlesnake Springs	19
11	Annual Mean Density and Biomass of Insect Trophic Levels in Douglas Creek	19
12	Annual Mean Density and Biomass of Insect Trophic Levels in Snively Springs	20
13	Annual Mean Density and Biomass of Insect Trophic Levels in Rattlesnake Springs	20
14	Annual Production of Insects from Douglas Creek, July 1985 to June 1986	22
15	Annual Production of Insects from Snively Springs, July 1985 to June 1986	25

16	Annual Production of Insects from Rattlesnake Springs, July 1985 to June 1986	28
17	Annual Production of Insect Functional Groups in Douglas Creek, Snively Springs, and Rattlesnake Springs	29
18	Annual Production of Insect Trophic Levels in Douglas Creek, Snively Springs, and Rattlesnake Springs	30

INTRODUCTION

Secondary production is defined as the rate of animal tissue elaboration over time regardless of the fate (carnivory, emergence, etc.) of that production (Benke and Wallace 1980). Estimating secondary production within a stream provides an assessment of the roles of animals within the ecosystem (Benke and Wallace 1980) while gaining insight into ecosystem dynamics.

Studies dealing only with density and biomass values may not accurately describe the roles of aquatic organisms within the stream. Benke et al. (1984) provide an example where the importance of gathering-collector invertebrates is underestimated by biomass analysis and overestimated by numerical analysis. Waters (1977) states that production is important to understanding ecosystem dynamics because it is the means by which energy is made available to higher trophic levels.

While most secondary production studies have focused on one or a few species within a stream (Benke and Wallace 1980, Waters and Hokenstrom 1980, O'Hop et al. 1984), more recent studies have estimated secondary production of the entire benthic fauna (Krueger and Waters 1983, Benke et al. 1984, Smock et al. 1985). This community-level approach provides the greatest insight into the ecology of stream ecosystems.

Community-level production studies, however, have been performed on very few kinds of streams. In particular, very little is known about the secondary production of the benthic invertebrate community in desert streams. The only publications on the subject to date are those of Fisher and Gray (1983) and Jackson and Fisher (1986) on Sycamore Creek, Arizona.

I estimated the production of aquatic insects in three cold-desert streams in Washington State: Rattlesnake Springs, Snively Springs, and Douglas Creek. Ecosystem dynamics in cold-desert streams will be more fully understood by estimating the secondary production of each species, functional group, trophic level, and community.

LITERATURE REVIEW

In his textbook on limnology, Welch (1935) defined his discipline as that branch of science that deals with the biological productivity of inland waters. Subsequently, scientists attempted to measure production rates of all trophic levels within an ecosystem in attempts to quantify ecosystem dynamics. Production is an important pathway of energy flow in ecosystems and is thus of great interest (Waters 1977).

Until recently, estimates of secondary production lagged behind those made of primary production. Reasons for the lag are: 1) the organisms are very diverse and some have complex life histories; 2) large numbers of samples are needed to reduce sample variation caused by a great variation in the distribution of organisms; and 3) separating, sorting, and analyzing samples are tedious and time-consuming, thus costly, tasks. Hynes (1970) reviewed the literature through 1966 and found only four attempts at directly estimating benthic invertebrate production in streams.

Several methods have been developed to estimate secondary production in streams. The Removal-Summation, Increment-Summation, Instantaneous Growth Rate, Allen Curve, and Hynes method have all been used. The Hynes method and its modifications have stood the test of time and is the most popular secondary production estimator in use today.

Originally proposed by Hynes and Coleman (1968), the Hynes method was designed as a shortcut approach to estimate production of entire benthic faunas. The method was unique in that it was not necessary to identify individual cohorts, thus making it easier to apply (Benke 1979). Hamilton (1969) modified the method by introducing the term "average cohort" when describing the size-frequency distribution that is necessary to make Hynes production estimates. Hamilton (1969) also pointed out three assumptions of the Hynes method that are difficult to justify: 1) all species in the samples are univoltine, 2) all species can potentially grow to maximum length used in the size-frequency distribution, and 3) all species require the same amount of time to grow

through each length class. These assumptions are difficult to meet when the Hynes method is applied to species with very different life histories. Benke and Waide (1977), however, showed that many of these assumptions can be met when the method is applied to single or similar species. Another problem that plagued the Hynes method was how to accurately estimate production of semivoltine or multivoltine populations. Benke (1979) addressed this problem by introducing the Cohort Production Interval (CPI). He suggested that the Hynes production value be multiplied by $365/\text{CPI}$, where CPI is the time in days from hatching to attainment of the largest aquatic size class. A final modification was proposed by Waters (1977) and confirmed by Hynes (1980) to change the name to the size-frequency (SF) method. This method is now commonly used by secondary production biologists and is the method I used in this study.

Recently the SF method has been used to assess secondary production of entire benthic communities (Benke et al. 1984, Jackson and Fisher 1986). Community production estimates are calculated by summing the species estimates and, in this way, accounting for variations in life histories (Benke et al. 1984). As life history data on greater numbers of species are obtained, the SF method will be more widely used and will become more accurate.

STUDY AREA

I studied three streams in the cold-desert Physiographic Province of Eastern Washington. This shrub-steppe region is characterized by the climatic climax community of big sage (Artemesia tridentata)-bluebunch wheatgrass (Agropyron spicatum). Mean annual precipitation is approximately 14 cm.

DOUGLAS CREEK

Douglas Creek is a spring-fed stream located in Douglas County, Washington, near the small community of Palisades, Washington. The stream drains an area of 530 km². The lower reaches of Douglas Creek, which are used extensively for irrigation, cease to flow during dry summers. For this reason, I located all study sites in the upper reaches where flow is permanent. Riparian vegetation is dominated by water birch (Betula occidentalis) and peachleaf willow (Salix amygdaloides). Populations of crayfish (Pacifasticus lenisculus) and rainbow trout (Salmo gairdneri) inhabit this stream.

Douglas Creek is the largest of the three streams studied. The average width was 4.0 m and average depth was 31 cm during base flow. The average discharge during the study period was 0.6 m³/s, and ranged from 0.5 m³/s in late summer to 0.9 m³/s in the early spring. Dissolved oxygen concentrations measured at midday ranged from a low of 9.6 mg/l in late summer to a high of 14.0 mg/l in midwinter. The temperature regime for Douglas Creek during the study period is shown in Figure 1. Table 1 shows the substrate composition of each study site.

SNIVELY SPRINGS

Snively Springs is a small spring-stream located on the U.S. Department of Energy's Hanford Site. It drains an area of approximately 40 km². During the summer portions of the spring-stream dry up, leaving about 3.6 km of perennial flow. Riparian vegetation is dominated by

FIGURE 1. Temperature Regime for Douglas Creek

TABLE 1. Substrate Composition of Douglas Creek, Snively Springs, and Rattlesnake Springs

Stream	Substrate Size (%)				
	Boulder (>255 mm)	Cobble (64-256 mm)	Pebble (2-16 mm)	Gravel (2-16 mm)	Sand/Silt (<2 mm)
Rattlesnake	0	0.9	7.3	10.8	81.0
Snively	7.4	20.4	25.3	10.5	36.4
Douglas (Site 1)	24.0	26.0	20.0	22.0	4.0
Douglas (Site 2)	26.0	44.0	16.0	12.0	2.0
Douglas (Site 3)	14.0	18.0	38.0	16.0	24.0

dominated by cattails (*Typha latifolia*) along the upper and lower reaches, and willow (*Salix* sp.) and wild rose (*Rosa* sp.) along the midreaches where it flows through a canyon. Watercress (*Rorippa nasturtium-aquaticum*) grows extensively within the spring-stream. A small population of crayfish [*Pacifastacus lenisculus* (*trowbridgii*)] inhabits Snively Springs.

The average width and depth of Snively Springs during base flow was 1.3 m and 10 cm, respectively. Average discharge during the study period was 0.04 m³/s, and ranged from 0.02 m³/s in the late spring to 0.05 m³/s in the winter. Dissolved oxygen concentrations measured at

midday ranged from 8.6 mg/l during the summer to 12.3 mg/l during the winter. The temperature regime for Snively Springs during the study period is shown in Figure 2, and the substrate composition is shown in Table 1.

RATTLESNAKE SPRINGS

Rattlesnake Springs is a small spring-stream also located on the U.S. Department of Energy's Hanford Site. It drains an area of 350 sq km (Cushing et al. 1980). Portions of the lower reaches of the spring-stream dry up during the summer, leaving about 2.5 km of perennial flow. This spring-stream is subject to severe flash flooding during the late winter about every four years (Cushing and Wolf 1982). Riparian vegetation is dominated by peachleaf willow (Salix amygdaloides). Watercress (Rorippa nasturtium-aquatica) is the dominant in-stream autotroph, although periphyton productivity exceeded that of watercress in 1969 and 1970 (Cushing and Wolf 1984).

The average width and depth of Rattlesnake Springs during base flow was 1.7 m and 5 cm, respectively. Average discharge during the study period was 0.05 m³/s, and ranged from 0.03 m³/s in the spring to

FIGURE 2. Temperature Regime for Snively Springs

0.07 m³/s during the winter. Midday dissolved oxygen concentrations ranged from 8.2 mg/l during the summer to 10.1 mg/l in midwinter. The temperature regime for Rattlesnake Springs during the study period is shown in Figure 3, and stream substrate composition is shown in Table 1.

FIGURE 3. Temperature Regime for Rattlesnake Springs

METHODS

I used a stratified random sampling regime (as recommended by Cummins 1962) in each stream to assure that all habitats were sampled. One study site was established in each of Snively Springs and Rattlesnake Springs, with each study site being large enough to include all habitat types. Three study sites were established in Douglas Creek: a shallow riffle (Site 1), a midsize pool and riffle (Site 2), and a large slow pool (Site 3).

Samples were taken at monthly intervals from July 1985 through June 1986. I collected three Surber samples ($0.093 \text{ m}^2/\text{sample}$) from Snively Springs and Rattlesnake Springs on each sampling date. A Surber sampler was used because these spring-streams are very shallow. A Portable Invertebrate Box Sampler ($0.1 \text{ m}^2/\text{sample}$) was used at Douglas Creek to collect one sample from Site 1, two samples from Site 2, and two samples from Site 3 on each sampling date. Fewer samples were collected from Site 1 because it was located in the upper reaches where streamflow was reduced. Douglas Creek is larger than the other study streams so the more efficient Portable Invertebrate Box Sampler was used. All samples were taken to a minimum depth of 10 cm and preserved in 70% ETOH for later lab analysis.

I separated insects from organic debris by sugar flotation (Anderson 1959) and sorted them into taxa. Insects were identified to the species level when possible, counted, and body length measured to the nearest 1 mm using a microscope equipped with an ocular micrometer. The trophic status of each species was determined by reference to Merritt and Cummins (1984) and examination of gut contents. Biomass [dry wt (DW)] was determined for all size classes after insects were dried at 60°C for 24 hours and weighed to the nearest 0.1 mg. I calculated the mean coefficient of variation (CV) and the mean standard error (SE) for all density and biomass estimates.

The SF method (Hynes and Coleman 1968, Hamilton 1969, Hynes 1980, Waters and Hokenstrom 1980) was used to estimate secondary production of

the most common benthic insect species. An average size-frequency distribution was determined from monthly sample sets that represented the survivorship curve of an "average cohort" (Hamilton 1969, Benke and Waide 1977). Production was estimated by calculating the difference in biomass between successive size classes and then multiplying this difference by the number of size classes (Hynes and Coleman 1968, Hamilton 1969). Production estimates were refined by multiplying by 365/CPI (Cohort Production Interval) (Benke 1979). The CPI (Benke et al. 1984) was estimated from life history data, field observations, and in situ growth studies.

Production/biomass (P/B) ratios were used to estimate secondary production for less abundant taxa. These P/B ratios were either species-specific values derived from one of the other two study streams, or an assumed cohort P/B value of 5 (Waters 1977, Benke et al. 1984). These taxa were not present in sufficient numbers to provide an accurate size-frequency distribution curve that is necessary to compute size-frequency production estimates.

RESULTS

INSECT DENSITY AND BIOMASS

Douglas Creek

Optioservus sp. (riffle beetles) and Baetis sp. (mayflies) were by far the most abundant insects found in Douglas Creek. These taxa accounted for 72% of the total insect numbers and 50% of the total biomass. Annual mean density and biomass of Optioservus sp. were 4322/m² and 606.7 mg DW/m², respectively. Annual mean density and biomass of Baetis sp. were 2416/m² and 263.7 mg DW/m², respectively (Table 2).

Snively Springs

Baetis sp. (mayflies) were the most numerous insects in Snively Springs, accounting for 42% of the total insect numbers and 25% of the total biomass. Annual mean density was 1388/m² and the annual mean biomass was 185.4 mg DW/m² (Table 3).

Rattlesnake Springs

The most abundant insects in Rattlesnake Springs were Simulium sp. (blackflies) and Baetis sp. (mayflies). These taxa accounted for 74% of the total insect numbers and 55% of the total biomass. Annual mean density and biomass of Simulium sp. were 1777/m² and 212.3 mg DW/m², respectively. Annual mean density and biomass of Baetis sp. were 1336/m² and 47.3 mg DW/m², respectively (Table 4).

FUNCTIONAL GROUP DENSITY AND BIOMASS

Results of gut content analysis and subsequent functional group classification for insects from Douglas Creek, Snively Springs, and Rattlesnake Springs are shown in Tables 5, 6, and 7, respectively.

Douglas Creek

The most abundant functional groups in Douglas Creek were grazer-scrappers (49%) and collectors (48%). These groups accounted for 97% of

TABLE 2. Annual Mean Density and Biomass of Insects from Douglas Creek
(SE = standard error, CV = coefficient of variation)

Taxa	Density (no./m ²)	SE/ $\bar{x}$	CV	Biomass (mg/m ²)	SE/ $\bar{x}$	CV
Diptera						
<u>Simulium</u> sp.	41	0.75	168.6	31.2	0.72	136.1
<u>Brillia flavifrons</u>	12	0.25	55.0	0.9	0.26	57.4
<u>Chironomus</u> sp.	753	0.71	152.3	60.7	0.69	153.8
<u>Parametriocnemus</u> sp.	196	0.44	98.0	10.4	0.46	101.9
<u>Thienemannimyia</u> sp.	11	0.81	180.5	0.9	0.83	185.4
<u>Chaetocladius</u> sp.	115	0.57	127.8	3.5	0.66	129.4
<u>Polypedilum</u> sp.	33	0.69	154.5	2.2	0.78	129.1
<u>Heleniella</u> sp.	141	0.52	116.4	4.5	0.54	116.5
<u>Phaenospectra</u> sp.	60	0.07	15.5	4.9	0.07	15.0
Tabanidae	51	0.48	106.6	27.8	0.48	107.5
Tipulidae	37	0.37	82.5	82.1	0.48	103.1
Empididae	1	0.22	50.0	0.1	0.18	40.0
(TOTAL)	(1451)			(229.2)		
Trichoptera						
<u>Leucotrichia</u>						
<u>pictipes</u>	95	0.63	139.7	7.7	0.68	153.2
<u>Hydropsyche</u> sp.	445	0.57	127.1	413.5	0.65	145.8
<u>Cheumatopsyche</u> sp.	156	0.53	118.3	84.1	0.60	135.0
(TOTAL)	(696)			(505.3)		
Coleoptera						
<u>Optioservus</u> sp.	4322	0.37	83.5	606.7	0.36	80.0
Ephemeroptera						
<u>Leucrocuta</u> sp.	160	0.47	104.0	51.4	0.51	104.0
<u>Baetis</u> sp.	2416	0.41	92.4	263.7	0.41	91.9
<u>Paraleptophlebia</u> sp.	225	0.35	78.5	48.1	0.38	85.4
<u>Tricorythodes</u> sp.	6	0.80	159.2	1.7	0.67	151.0
(TOTAL)	(2807)			(364.9)		
Odonata						
<u>Argia tibialis</u>	30	0.46	103.9	8.9	0.49	110.3
Plecoptera						
<u>Isoperla</u> sp.	77	0.58	129.4	42.8	0.58	113.9
(GRAND TOTAL)	(9383)			(1757.8)		

TABLE 3. Annual Mean Density and Biomass of Insects from Snively Springs
(SE = standard error, CV = coefficient of variation)

Taxa	Density (no./m ²)	SE/ $\bar{X}$	CV	Biomass (mg/m ²)	SE/ $\bar{X}$	CV
Diptera						
<u>Simulium</u> sp.	276	0.70	121.3	34.3	0.82	142.6
<u>Chaetocladius</u> sp.	92	0.63	108.3	2.7	0.69	120.2
<u>Chironomus</u> sp.	412	0.54	93.2	17.1	0.58	99.8
<u>Heleniella</u> sp.	381	0.40	69.2	9.2	0.37	64.7
<u>Polypedilum</u> sp.	123	0.56	96.2	3.2	0.52	89.1
<u>Thienemannimyia</u> sp.	18	0.42	72.3	1.1	0.33	57.3
Dixidae	21	0.55	95.9	1.3	0.65	111.5
Tabanidae	52	0.47	81.5	10.5	0.50	86.4
Tipulidae	25	0.60	103.8	219.2	0.50	87.4
Empididae	4	0.15	26.6	0.6	0.12	32.1
(TOTAL)	(1404)			(299.2)		
Trichoptera						
<u>Cheumatopsyche</u> sp.	433	0.41	83.0	200.9	0.51	86.9
Ephemeroptera						
<u>Baetis</u> sp.	1388	0.62	104.7	185.4	0.55	96.3
<u>Paraleptophlebia</u> sp.	54	0.27	47.5	15.5	0.28	48.2
(TOTAL)	(1442)			(200.9)		
Odonata						
<u>Argia tibialis</u>	22	0.61	106.6	27.8	0.68	118.6
(GRAND TOTAL)	(3301)			(728.8)		

the total insect numbers and 90% of the total biomass. Annual mean density and biomass of grazer-scrapers was 4637/m² and 670.7 mg DW/m², respectively. Annual mean density and biomass of collectors was 4494/m² and 917.9 mg DW/m², respectively (Table 8).

Snively Springs

Collectors were by far the most abundant functional group in Snively Springs, accounting for 93% of the total insect numbers and

TABLE 4. Annual Mean Density and Biomass of Insects from Rattlesnake Springs (SE = standard error, CV = coefficient of variation)

Taxa	Density (no./m ²)	SE/ $\bar{X}$	CV	Biomass (mg/m ²)	SE/ $\bar{X}$	CV
Diptera						
<u>Simulium</u> sp.	1777	0.73	125.8	212.3	0.73	127.5
<u>Chironomus</u> sp.	192	0.50	87.3	7.0	0.58	100.8
<u>Thienemannimyia</u> sp.	114	0.55	94.9	3.3	0.55	95.2
<u>Chaetocladius</u> sp.		0.73	126.4	0.4	0.56	97.7
<u>Heleniella</u> sp.	352	0.51	89.0	5.4	0.51	88.4
<u>Polypedilum</u> sp.	13	0.62	108.2	0.6	0.46	78.9
Misc. Chironomidae	18	0.29	50.1	0.8	0.38	66.3
Dixidae	2	0.28	64.7	0.1	0.29	50.0
Tabanidae	34	0.51	85.6	15.9	0.64	111.0
Tipulidae	3	0.21	35.9	2.0	0.26	44.3
Empididae	8	0.39	68.3	0.4	0.23	39.8
(TOTAL)	(2572)			(248.2)		
Trichoptera						
<u>Cheumatopsyche</u> sp.	140	0.69	118.9	48.6	0.78	134.5
<u>Limnephilus</u> sp.	52	0.45	76.9	22.0	0.38	66.3
<u>Parapsyche</u> sp.	10	0.24	41.7	26.8	0.25	43.4
(TOTAL)	(202)			(98.3)		
Coleoptera						
<u>Hydaticus</u> sp.	4	0.50	87.4	1.2	0.35	60.1
Hydrophilidae	1	0.27	47.6	0.3	0.25	43.1
(TOTAL)	(5)			(1.5)		
Ephemeroptera						
<u>Baetis</u> sp.	1336	0.61	107.2	47.3	0.58	104.0
<u>Tricorythodes</u> sp.	1	0.05	8.3	0.3	0.07	12.2
(TOTAL)	(1337)			(47.6)		
Odonata						
<u>Argia tibialis</u>	67	0.72	124.1	74.3	0.78	134.9
(GRAND TOTAL)	(4183)			(469.9)		

TABLE 5. Results of Qualitative Gut Content Analysis and Subsequent Functional Group and Trophic Level Classification for Insects from Douglas Creek (N = number of guts examined)

Taxa	N	Gut Content	Functional Group	Trophic Level
Diptera				
<u>Simulium</u> sp.	14	detritus	coll-gath	detritivore
<u>Brillia flavifrons</u> (a)			shredder	detritivore
<u>Chironomus</u> sp.	21	detritus/ some algae	coll-gath	detritivore
<u>Parametriochnemus</u> sp.	14	detritus/ some algae	coll-gath	detritivore
<u>Thienemannimyia</u> sp. (a)			predator	carnivore
<u>Chaetocladius</u> sp.	10	detritus	coll-gath	detritivore
<u>Polypedilum</u> sp. (a)			shredder	herbivore
<u>Heleniella</u> sp.	10	detritus	coll-gath	detritivore
<u>Phaenospectra</u> sp. (a)			scraper	herbivore
<u>Tabanidae</u> (a)			predator	carnivore
<u>Tipulidae</u> (a)			shredder	detritivore
<u>Empididae</u> (a)			predator	carnivore
Trichoptera				
<u>Leucotrichia pictipes</u>	17	algae/some detritus	scraper	herbivore
<u>Hydropsyche</u> sp.	45	detritus	coll-filt	detritivore
<u>Cheumatopsyche</u> sp.	16	detritus	coll-filt	detritivore
Coleoptera				
<u>Optioservus</u> sp.	50	algae/some detritus	scraper	herbivore
Ephemeroptera				
<u>Leucrocuta</u> sp.	44	algae/some detritus	scraper	herbivore
<u>Baetis</u> sp.	51	detritus/ some algae	coll-gath	detritivore
<u>Paraleptophlebia</u> sp.	44	detritus/ some algae	coll-gath	detritivore
<u>Tricorythodes</u> sp.	7	detritus/ some algae	coll-gath	detritivore
Odonata			predator	carnivore
<u>Argia tibialis</u> (a)				
Plecoptera				
<u>Isoperla</u> sp.	26	animal- tissue	predator	carnivore

(a) Classifications based upon literature only (Merritt and Cummins 1984).

TABLE 6. Results of Qualitative Gut Content Analysis and Subsequent Functional Group and Trophic Level Classification of Insects from Snively Springs (N = number of guts analyzed)

<u>Taxa</u>	<u>N</u>	<u>Gut Content</u>	<u>Functional Group</u>	<u>Trophic Level</u>
Diptera				
<u>Simulium</u> sp.	43	detritus	coll-filt	detritivore
<u>Chaetocladius</u> sp.(a)			coll-gath	detritivore
<u>Chironomus</u> sp.	14	detritus/ some algae	coll-gath	detritivore
<u>Heleniella</u> sp.	14	detritus/ some algae	coll-gath	detritivore
<u>Polypedilum</u> sp.(a)			shredder	herbivore
<u>Thienemannimyia</u> sp.(a)			predator	carnivore
<u>Dixidae</u> (a)			coll-gath	detritivore
<u>Tabanidae</u> (a)			predator	carnivore
<u>Tipulidae</u> (a)			shredder	detritivore
<u>Empididae</u> (a)			predator	carnivore
Trichoptera				
<u>Cheumatopsyche</u> sp.	41	detritus	coll-filt	detritivore
Ephemeroptera				
<u>Baetis</u> sp.	36	detritus	coll-gath	detritivore
<u>Paraleptophlebia</u> sp.	10	detritus	coll-gath	detritivore
Odonata				
<u>Argia tibialis</u>	11	animal- tissue	predator	carnivore

(a) Classifications based upon literature only (Merritt and Cummins 1984).

64% of the total biomass. Annual mean density was 3057/m² and the annual mean biomass was 466.4 mg DW/m². No grazer-scrappers were found in Snively Springs (Table 9).

Rattlesnake Springs

As in Snively Springs, collectors were the most abundant functional group in Rattlesnake Springs, accounting for 93% of the total insect

TABLE 7. Results of Qualitative Gut Content Analysis and Subsequent Functional Group and Trophic Level Classification of Insects from Rattlesnake Springs (N = number of guts analyzed)

Taxa	N	Gut Content	Functional Group	Trophic Level
Diptera				
<u>Simulium</u> sp.	47	detritus	coll-filt	detritivore
<u>Chironomus</u> sp.	14	detritus	coll-gath	detritivore
<u>Thienemannimyia</u> sp.(a)			predator	carnivore
<u>Chaetocladius</u> sp.	7	detritus/ some algae	coll-gath	detritivore
<u>Heleniella</u> sp.	13	detritus/ some algae	coll-gath	detritivore
<u>Polypedilum</u> sp.(a)			shredder	herbivore
Misc. Chironomidae(a)			coll-gath	detritivore
Dixidae(a)			coll-gath	detritivore
Tabanidae(a)			predator	carnivore
Tipulidae(a)			shredder	detritivore
Empididae(a)			predator	carnivore
Trichoptera				
<u>Cheumatopsyche</u> sp.	37	algae/some detritus	coll-filt	detritivore
<u>Limnephilus</u> sp.	8	detritus	shredder	detritivore
<u>Parapsyche</u> sp.	7	detritus/ animal- tissue	coll-filt	detritivore
Coleoptera				
<u>Hydaticus</u> sp.(a)		predator	carnivore	
<u>Hydrophilidae</u> (a)			predator	carnivore
Ephemeroptera				
<u>Baetis</u> sp.	36	detritus	coll-gath	detritivore
<u>Tricorythodes</u> sp.(a)			coll-gath	detritivore
Odonata				
<u>Argia tibialis</u>	42	animal- tissue	predator	carnivore

(a) Classifications based upon literature only (Merritt and Cummins 1984).

TABLE 8. Annual Mean Density and Biomass of Insect Functional Groups in Douglas Creek

Functional Group	Density (no./m ²)	Composition (%)	Biomass (mg/m ²)	Composition (%)
Grazer-scraper	4637	49.4	670.7	38.2
Collector (TOTAL)	(4494)	(47.9)	(917.9)	(52.2)
Gatherer	3852	41.1	389.1	22.1
Filterer	642	6.8	528.8	30.1
Shredder	82	0.9	86.1	4.9
Predator	<u>170</u>	<u>1.8</u>	<u>83.1</u>	<u>4.7</u>
(GRAND TOTAL)	(9383)	(100.0)	(1757.8)	(100.0)

TABLE 9. Annual Mean Density and Biomass of Insect Functional Groups in Snively Springs

Functional Group	Density (no./m ²)	Composition (%)	Biomass (mg/m ²)	Composition (%)
Collector (TOTAL)	(3057)	(92.6)	(466.4)	(64.0)
Gatherer	2348	71.1	231.2	31.7
Filterer	709	21.5	235.2	32.3
Shredder	148	4.5	222.4	30.5
Predator	<u>96</u>	<u>2.9</u>	<u>40.0</u>	<u>5.5</u>
(GRAND TOTAL)	(3301)	(100.0)	(728.8)	(100.0)

numbers and 74% of the total biomass. Annual mean density and biomass of collectors was 3887/m² and 349.0 mg DW/m², respectively. Grazer-scrappers were absent from Rattlesnake Springs (Table 10).

TROPHIC LEVEL DENSITY AND BIOMASS

Trophic level classifications for insects from Douglas Creek, Snively Springs, and Rattlesnake Springs are shown in Tables 5, 6, and 7, respectively.

Douglas Creek

Herbivores (50%) and detritivores (48%) were the most abundant trophic levels in Douglas Creek. These trophic levels accounted for

TABLE 10. Annual Mean Density and Biomass of Insect Functional Groups in Rattlesnake Springs

Functional Group	Density (no./m ²)	Composition (%)	Biomass (mg/m ²)	Composition (%)
Collector (TOTAL)	(3887)	(92.9)	(349.0)	(74.3)
Gatherer	1960	46.9	61.3	13.1
Filterer	1927	46.1	287.7	61.2
Shredder	68	1.6	25.5	5.4
Predator	228	5.5	95.4	20.3
(GRAND TOTAL)	(4183)	(100.0)	(469.9)	(100.0)

98% of the total insect numbers and 95% of the total biomass. Annual mean density and biomass of herbivores was 4670/m² and 672.9 mg DW/m², respectively. The annual mean density and biomass of detritivores was 4543/m² and 1004.4 mg DW/m², respectively (Table 11).

Snively Springs

Detritivores were by far the most abundant trophic level in Snively Springs, accounting for 93% of the total insect numbers and 94% of the total biomass. Annual mean density was 3082/m² and the annual mean biomass was 685.6 mg DW/m² (Table 12).

Rattlesnake Springs

Detritivores were the most abundant trophic level in Rattlesnake Springs. Detritivores comprised 94% of the total insect numbers and 80% of the total biomass. Annual mean density was 3942/m² and the annual mean biomass was 373.9 mg DW/m² (Table 13).

TABLE 11. Annual Mean Density and Biomass of Insect Trophic Levels in Douglas Creek

Trophic Level	Density (no./m ²)	Composition (%)	Biomass (mg/m ²)	Composition (%)
Herbivore	4670	49.8	672.9	38.3
Detritivore	4543	48.4	1004.4	57.1
Carnivore	170	1.8	80.5	4.6
(GRAND TOTAL)	(9383)	(100.0)	(1757.8)	(100.0)

TABLE 12. Annual Mean Density and Biomass of Insect Trophic Levels in Snively Springs

<u>Trophic Level</u>	<u>Density (no./m²)</u>	<u>Composition (%)</u>	<u>Biomass (mg/m²)</u>	<u>Composition (%)</u>
Herbivore	123	3.7	3.2	0.4
Detritivore	3082	93.4	685.6	94.1
Carnivore	<u>96</u>	<u>2.9</u>	<u>40.0</u>	<u>5.5</u>
(GRAND TOTAL)	(3301)	(100.0)	(728.8)	(100.0)

TABLE 13. Annual Mean Density and Biomass of Insect Trophic Levels in Rattlesnake Springs

<u>Trophic Level</u>	<u>Density (no./m²)</u>	<u>Composition (%)</u>	<u>Biomass (mg/m²)</u>	<u>Composition (%)</u>
Herbivore	13	0.3	0.6	0.1
Detritivore	3942	94.2	373.9	79.6
Carnivore	<u>228</u>	<u>5.5</u>	<u>95.4</u>	<u>20.3</u>
(GRAND TOTAL)	(4183)	(100.0)	(469.9)	(100.0)

INSECT PRODUCTION

Douglas Creek

Diptera--Previous studies have addressed the difficulties in obtaining accurate field estimates of blackfly larvae (Simuliidae) and midge larvae (Chironomidae) CPIs, and thus production estimates (Benke et al. 1984, Behmer and Hawkins 1986). Their small size, rapid turnover rate, high density, and diversity make accurate species-specific CPI estimates difficult. These same characteristics, however, make midges and blackflies very important to a stream community. In many streams these insects contribute a large percentage of the total community production because of their rapid development and high turnover rates.

Simulium sp. (blackflies) from Douglas Creek were not present in sufficient numbers to calculate a size-frequency (SF) production estimate. A P/B ratio of 54.0 was used to estimate production. This P/B ratio was calculated by using the SF method to average the P/B

ratios obtained for Simulium sp. in Snively Springs and Rattlesnake Springs. Blackfly production in Douglas Creek was 1638.2 mg DW/m²/yr (Table 14).

Accurate CPI estimates for Chironomidae (midges) could not be obtained from field observations or SF distributions. Benke et al. (1984) provide an extensive review of literature concerning midge development times. Using these data and Mackey's (1977) laboratory growth studies, I feel that reasonable CPI estimates were derived.

Mackey (1977) reported larval development times of 21 days for Chironomus sp., 13 days for Polypedilum convictum, and 36 days for Phaenospectra flavipes at 15°C. To compensate for slightly lower average temperatures (13°C) in Douglas Creek, and environmental stress, I estimated CPIs of 25 days for Chironomus, 20 days for Polypedilum, and 40 days for Phaenospectra. For the remaining midge species I assumed a CPI of 25 days. Production estimates ranged from 67.5 mg DW/m²/yr by Brillia flavifrons to 4922.8 mg DW/m²/yr by Chironomus. Annual P/B ratios ranged from 45.0 by Phaenospectra to 121.7 by Chaetocladius. These P/B ratios seem high but are comparable to those reported in other studies when short CPIs were utilized (Benke et al. 1984, Jackson and Fisher 1986).

Tabanidae and Tipulidae were assumed to be univoltine with a development time of 1 year. This is consistent with Gray's (1981) estimate of a 1-year development time for Tabanus dorsifer (Tabanidae) in Sycamore Creek, Arizona. Empididae grew to a maximum size that was similar to many of the midges, therefore a CPI of 25 days was assumed. Table 14 shows production data for all dipterans in Douglas Creek.

Trichoptera--Leucotrichia pictipes (microcaddisfly) was univoltine, and as SF distributions and field observations indicated the larvae overwintered as a late instar and emerged as adults in the spring. This observation is supported by McAuliffe's (1982) studies on L. pictipes in Owl Creek, Montana. Production and the annual P/B ratio of Leucotrichia sp. in Douglas Creek were 32.0 mg DW/m²/yr and 4.2, respectively.

TABLE 14. Annual Production of Insects from Douglas Creek, July 1985 to June 1986

Taxa	365/CPI	Calculation Method ^(a,b)	Production (mg/m ² /yr)	Annual P/B
Diptera				
<u>Simulium</u> sp.	12	PB	1638.2	54.0 ^(c)
<u>Brillia flavifrons</u>	15	PB	67.5	75.0 ^(d)
<u>Chironomus</u> sp.	15	PB	4922.8	81.1 ^(c)
<u>Parametriocnemus</u> sp.	15	SF	874.5	84.1
<u>Thienemanniomyia</u> sp.	15	PB	75.2	83.6 ^(c)
<u>Chaetocladius</u> sp.	15	SF	426.0	121.7
<u>Polypedilum</u> sp.	18	SF	160.9	73.1
<u>Heleniella</u> sp.	15	SF	423.0	94.0
<u>Phaenospectra</u> sp.	9	PB	220.5	45.0 ^(c)
Tabanidae	1	PB	130.0	5.0 ^(c)
Tipulidae	1	PB	410.5	5.0 ^(c)
Empididae	15	PB	7.5	75.0 ^(c)
(TOTAL)			(9365.6)	
Trichoptera				
<u>Leucotrichia pictipes</u>	1	SF	32.0	4.2
<u>Hydropsyche</u> sp.	1	SF	1700.0	4.1
<u>Cheumatopsyche</u> sp.	2	SF	817.5	9.7
(TOTAL)			(2549.5)	
Coleoptera				
<u>Optioservus</u> sp.	1	SF	2162.7	3.6
Ephemeroptera				
<u>Leucrocuta</u> sp.	1	SF	237.8	4.6
<u>Baetis</u> sp.	6	SF	8316.5	31.5
<u>Paraleptophlebia</u> sp.	1	SF	249.4	5.2
<u>Tricorythodes</u> sp.	9	PB	76.5	45.0 ^(c)
(TOTAL)			(8880.2)	
Odonata				
<u>Argia tibialis</u>	1	PB	44.3	5.0 ^(c)
Plecoptera				
<u>Isoperla</u> sp.	1	SF	182.8	4.3
(GRAND TOTAL)			(23185.1)	

(a) PB = production calculated by an assumed P/B ratio.

(b) SF = production calculated by the size-frequency method.

(c) Assumed cohort P/B of 5.

(d) Assumed annual P/B is the same as derived by SF for this species in one of the other study streams.

Field observations and SF distributions of Hydropsyche sp. and Cheumatopsyche sp. indicated CPIs of 1 year and 6 months, respectively. Production and the annual P/B ratio of Hydropsyche sp. were 1700.0 mg DW/m²/yr and 4.1, respectively. Production of Cheumatopsyche sp. was 817.5 mg DW/m²/yr and the annual P/B ratio was 9.7. Total caddisfly production was 2549.5 mg DW/m²/yr (Table 14).

Coleoptera--An accurate CPI estimate for Optioservus sp. (riffle beetles) was difficult to make because few data are available concerning their development times. Field data, though not clearcut, indicated a CPI of 1 year. Using this CPI estimate, production of riffle beetles was 2162.7 mg DW/m²/yr and the annual P/B ratio was 3.6 (Table 14).

Ephemeroptera--Mayflies typically exhibit widely varied larval development times (Clifford 1982). I assumed Leucrocuta sp. (Heptageniidae) to be univoltine with a CPI of 1 year. Clifford (1982) examined life-cycle data of 85 species of Heptageniidae and found >90% had at least one univoltine life cycle. Production of Leucrocuta sp. was 237.8 mg DW/m²/yr and the annual P/B ratio was 4.6.

Field data for Baetis sp. from Douglas Creek provided little clarification of their CPI. Based upon field observations of Baetis sp. from the other study streams, and a growth study in Rattlesnake Springs (Gaines, unpublished data), I estimated a CPI of 60 days. Temperature regimes were similar between study streams, making this estimation reasonable. Production of Baetis sp., 8316.5 mg DW/m²/yr, was the highest of all insects from Douglas Creek. The annual P/B ratio was 31.5 (Table 14).

Paraleptophlebia sp. are generally univoltine, having either a summer or winter cycle (Clifford 1982). In Douglas Creek, however, seasonal cycles could not be distinguished. Since all Paraleptophlebia species have at least one univoltine cycle (Clifford 1982) and were present in Douglas Creek throughout the study year, I assumed a CPI of 1 year. Production was 249.4 mg DW/m²/yr and the annual P/B ratio was 5.2 (Table 14).

Because of low numbers of Tricorythodes sp., field data provided little indication of their CPI. McCullough et al. (1979) reported a 34-day larval development time for Tricorythodes minutus grown in the field at 18°C. I estimated a CPI of 40 days for Tricorythodes sp. because of lower stream temperatures in Douglas Creek. A cohort P/B ratio of 5 was assumed to estimate production of 76.5 mg DW/m²/yr (Table 14).

Odonata--Argia tibialis (damselflies) were univoltine and field data indicated a CPI of 1 year. I assumed a cohort P/B ratio of 5.0 to estimate production of 44.3 mg DW/m²/yr (Table 14).

Plecoptera--A CPI estimate for Isoperla sp. (stoneflies) could not be made from field data. I assumed a CPI of 1 year because stoneflies grew to a maximum size that was similar to damselflies. Production was 182.8 mg DW/m²/yr and the annual P/B ratio was 4.3 (Table 14).

Snively Springs

Diptera--Field data and observations indicated a larval development time (CPI) of 30 days for Simulium sp. (blackflies) in Snively Springs. Becker (1973) reported a larval development time of 13 days for S. vittatum grown in the laboratory at 17°C. A 30-day CPI was a reasonable estimate when lower stream temperatures and environmental stress are considered. On this basis, production was 1878.7 mg DW/m²/yr and the annual P/B ratio was 54.8 (Table 15).

The larval development times (CPIs) of Chironomidae (midges) in Snively Springs were estimated as they were in Douglas Creek. I assumed CPIs of 20 days for Polypedilum sp. and 25 days for all other midges. Production estimates ranged from 92.0 mg DW/m²/yr by Thienemannimyia sp. to 1386.0 mg DW/m²/yr by Chironomus sp. The P/B ratios ranged from 60.3 by Heleniella sp. to 83.6 by Thienemannimyia sp.

I assumed Tabanidae and Tipulidae to be univoltine with a CPI of 1 year (Gray 1981). Dixidae and Empididae reached maximum sizes that were very similar to many of the midges, and I assumed a CPI of 25 days. Table 15 shows the production data of all dipterans in Snively Springs.

TABLE 15. Annual Production of Insects from Snively Springs, July 1985 to June 1986

Taxa	365/CPI	Calculation Method ^(a,b)	Production (mg/m ² /yr)	Annual P/B
Diptera				
<u>Simulium</u> sp.	12	SF	1878.7	54.8
<u>Chaetocladius</u> sp.	15	SF	210.0	77.8
<u>Chironomus</u> sp.	15	SF	1386.0	81.1
<u>Heleniella</u> sp.	15	SF	550.0	60.3
<u>Polypedilum</u> sp.	18	SF	219.6	68.6
<u>Thienemannimyia</u> sp.	15	PB	92.0	83.6 ^(c)
Dixidae	15	PB	97.5	75.0 ^(d)
Tabanidae	1	PB	52.5	5.0 ^(d)
Tipulidae	1	PB	1096.0	5.0 ^(d)
Empididae	15	PB	45.0	75.0 ^(d)
(TOTAL)			(5627.3)	
Trichoptera				
<u>Cheumatopsyche</u> sp.	2	SF	1297.9	6.5
Ephemeroptera				
<u>Baetis</u> sp.	6	SF	7011.6	37.8
<u>Paraleptophlebia</u> sp.	1	SF	67.0	4.3
(TOTAL)			(7078.6)	
Odonata				
<u>Argia tibialis</u>	1	PB	139.1	5.0 ^(d)
(GRAND TOTAL)			(14142.9)	

(a) PB = production calculated by an assumed P/B ratio.

(b) SF = production calculated by the size-frequency method.

(c) Assumed cohort P/B of 5.

(d) Assumed annual P/B is the same as derived by SF for this species in one of the other study streams.

Trichoptera--Field observations and SF data indicated a bivoltine life cycle and a CPI of 6 months for Cheumatopsyche sp., the only caddisfly in Snively Springs. Production was 1297.9 mg DW/m²/yr and the annual P/B ratio was 6.5 (Table 15).

Ephemeroptera--Field observations and SF data indicated a CPI of 60 days for Baetis sp. in Snively Springs. Gray (1981) reported a larval development time of 20 days for Baetis guilleri in Sycamore Creek, Arizona. Due to lower stream temperatures, however, Baetis sp. developed more slowly in all of the streams in this study. Production by Baetis sp. was 7011.6 mg DW/m²/yr and the annual P/B ratio was 37.8 (Table 15).

In Snively Springs, Paraleptophlebia sp. were present only during the summer. Thus, I used only data collected during the summer to calculate production. Production of Paraleptophlebia sp. was 67.0 mg DW/m²/yr and the annual P/B ratio was 4.3 (Table 15).

Odonata--Argia tibialis were not present in sufficient numbers to make an SF production estimate. I assumed a cohort P/B ratio of 5 and a CPI of 1 year to estimate production of 139.1 mg DW/m²/yr (Table 15).

Rattlesnake Springs

Diptera--Several Simulium sp. larvae were isolated in growth chambers in Rattlesnake Springs to estimate larval development time. These data and field observations indicated a CPI of 30 days. Production by blackflies, 11174.8 mg DW/m²/yr, was the highest of all species in Rattlesnake Springs, and the annual P/B ratio was 52.6.

I estimated larval development times (CPIs) of Rattlesnake Springs Chironomidae as I did in the other study streams. CPIs of 20 days for Polypedilum sp. and 25 days for all other midges were assumed. Production estimates ranged from 30.0 mg DW/m²/yr by Chaetocladius sp. to 489.0 mg DW/m²/yr by Chironomus sp. Annual P/B ratios ranged from 68.6 by Polypedilum sp. to 83.6 by Thienemannimyia sp.

Tabanidae and Tipulidae were assumed to be univoltine with CPIs of 1 year (Gray 1981). Dixidae and Empididae grew to maximum sizes similar to many of the midges and CPIs of 25 days were assumed. Table 16 shows the production data of all dipterans from Rattlesnake Springs.

Trichoptera--I isolated several Cheumatopsyche sp. larvae in growth chambers in Rattlesnake Springs to estimate larval development time. These data indicated a bivoltine life cycle and a CPI of 6 months. Production by caddisflies in Rattlesnake Springs was 486.0 mg DW/m²/yr and the annual P/B ratio was 10.0 (Table 16).

Because of low densities, field data gave no indication of the CPIs of Limnephilus sp. or Parapsyche sp. To estimate their production, I assumed a CPI of 1 year and a cohort P/B ratio of 5. Production was 114.6 mg DW/m²/yr by Limnephilus sp. and 133.8 mg DW/m²/yr by Parapsyche sp. (Table 16).

Coleoptera--Field data provided little indication of the CPIs of beetles (coleopterans) because of low numbers. To estimate production, I assumed a CPI of 1 year and a cohort P/B ratio of 5. Production by Hydaticus sp. was 6.0 mg DW/m²/hr (Table 16). Production by Hydrophilidae was 1.5 mg DW/m²/yr (Table 16).

Ephemeroptera--I isolated several Baetis sp. larvae in growth chambers in Rattlesnake Springs to estimate larval development time. These data and field observations indicated a CPI of 60 days. Production by Baetis sp. was 2542.3 mg DW/m²/yr and the annual P/B ratio was 53.8 (Table 16).

Tricorythodes sp. were not present in sufficient numbers for an SF production estimate. I assumed a cohort P/B ratio of 5 and a CPI of 40 days (McCullough et al. 1979) to calculate production of 13.5 mg DW/m²/yr (Table 16).

Odonata--Field observations of Argia tibialis indicated a CPI of 1 year, and I assumed a cohort P/B ratio of 5 to calculate production. Production by damselflies in Rattlesnake Springs was 371.6 mg DW/m²/yr (Table 16).

FUNCTIONAL GROUP PRODUCTION

Production by collectors was the highest of all functional groups in all study streams. Collector production was highest in Douglas Creek

TABLE 16. Annual Production of Insects from Rattlesnake Springs,
July 1985 to June 1986

Taxa	365/CPI	Calculation Method ^(a,b)	Production (mg/m ² /yr)	Annual P/B
Diptera				
<u>Simulium</u> sp.	12	SF	11174.8	52.6
<u>Chironomus</u> sp.	15	SF	489.0	69.9
<u>Thienemannimyia</u> sp.	15	SF	279.0	83.6
<u>Chaetocladius</u> sp.	15	SF	30.0	75.0 ^(c)
<u>Heleniella</u> sp.	15	SF	480.0	88.9
<u>Polypedilum</u> sp.	18	PB	41.2	68.6 ^(d)
Miscellaneous	15	PB	60.0	75.0 ^(c)
Chironomidae				
Dixidae	15	PB	7.5	75.0 ^(c)
Tabanidae	1	PB	79.5	5.0 ^(c)
Tipulidae	1	PB	10.0	5.0 ^(c)
Empididae	15	PB	30.0	75.0 ^(c)
(TOTAL)			(12681.0)	
Trichoptera				
<u>Cheumatopsyche</u> sp.	2	SF	486.0	10.0
<u>Limnephilus</u> sp.	1	PB	114.6	5.0 ^(c)
<u>Parapsyche</u> sp.	1	PB	133.8	5.0 ^(c)
(TOTAL)			(734.4)	
Coleoptera				
<u>Hydaticus</u> sp.	1	PB	6.0	5.0 ^(c)
<u>Hydrophilidae</u>	1	PB	1.5	5.0 ^(c)
(TOTAL)			(7.5)	
Ephemeroptera				
<u>Baetis</u> sp.	6	SF	2542.3	53.8
<u>Tricorythodes</u> sp.	9	PB	13.5	45.0 ^(c)
(TOTAL)			(2555.8)	
Odonata				
<u>Argia tibialis</u>	1	PB	371.6	5.0 ^(c)
(GRAND TOTAL)			(16350.3)	

(a) PB = production calculated by an assumed P/B ratio.

(b) SF = production calculated by the size-frequency method.

(c) Assumed cohort P/B of 5.

(d) Assumed annual P/B is the same as derived by SF for this species in one of the other study streams.

at 19444.4 mg DW/m²/yr, accounting for 84% of the total annual production. In Snively Springs, collector production was 12498.7 mg DW/m²/yr, comprising 88% of the total annual production. Production of collectors in Rattlesnake Springs was 15416.9 mg DW/m²/yr, accounting for 94% of the total annual production. Table 17 shows the annual production by functional groups in each study stream.

TROPHIC LEVEL PRODUCTION

Detritivore production was the highest of all trophic levels in each study stream. In Douglas Creek, detritivore production was 19922.4 mg DW/m²/yr, accounting for 86% of the total annual production. In Snively Springs detritivore production was 13594.7 mg DW/m²/yr, accounting for 96% of the total annual production. Detritivore production in Rattlesnake Springs was 15541.5 mg DW/m²/yr, accounting for 95% of the total annual production. Table 18 shows the annual production by trophic levels in each study stream.

TABLE 17. Annual Production of Insect Functional Groups in Douglas Creek, Snively Springs, and Rattlesnake Springs

Functional Group	Douglas Creek (mg/m ² /yr) (%)		Snively Springs (mg/m ² /yr) (%)		Rattlesnake Springs (mg/m ² /yr) (%)	
Grazer-Scraper	2653.0	11.4	0.0	0.0	0.0	0.0
Collector						
(TOTAL)	(19444.4)	(83.9)	(12498.7)	(88.4)	(15416.9)	(94.3)
Gatherer	15288.7	66.0	9322.1	65.9	3622.3	22.2
Filterer	4155.7	17.9	3176.6	22.5	11794.6	72.1
Shredder	638.9	2.8	1315.6	9.3	165.8	1.0
Predator	448.8	1.9	328.6	2.3	767.6	4.7
(GRAND TOTAL)	(23185.1)	(100.0)	(14142.9)	(100.0)	(16350.3)	(100.0)

TABLE 18. Annual Production of Insect Trophic Levels in Douglas Creek, Snively Springs, and Rattlesnake Springs

Trophic Level	Douglas Creek (mg/m ² /yr) (%)		Snively Springs (mg/m ² /yr) (%)		Rattlesnake Springs (mg/m ² /yr) (%)	
Herbivore	2813.9	12.2	219.6	1.6	41.2	0.3
Detritivore	19922.4	85.9	13594.7	96.1	15541.5	95.0
Carnivore	<u>448.8</u>	<u>1.9</u>	<u>328.6</u>	<u>2.3</u>	<u>767.6</u>	<u>4.7</u>
(GRAND TOTAL)	(23185.1)	(100.0)	(14142.9)	(100.0)	(16350.3)	(100.0)

DISCUSSION

ESTIMATING SECONDARY PRODUCTION

Benke et al. (1984) attempted to describe a methodological framework for other investigators to follow when estimating secondary production. They stressed the importance and difficulty of accurately estimating the CPI of each taxon within a benthic community. They suggested using laboratory growth data and life history data when direct field data were not available. I followed their framework when estimating secondary production. Many of my CPI estimates are based upon a review of the literature because of a lack of direct field data. These estimates may be refined as the body of information grows concerning larval development times of aquatic invertebrates. I have stated all assumptions clearly to aid investigators in this process.

Benke et al. (1984) also stressed the importance and value of estimating secondary production. They stated that production analysis, more than density or biomass, is the most meaningful way to analyze functional feeding groups in natural ecosystems. They then provided an example where the importance of gathering-collector invertebrates was underestimated by biomass analysis and overestimated by density analysis (e.g. in Douglas Creek, grazer-scraper importance was overestimated and collector importance was underestimated). My work provides another example of the importance of estimating secondary production to accurately describe the roles of organisms in streams and to understand ecosystem dynamics.

COMPARISON BETWEEN STUDY STREAMS

Insect Faunas

Similarities occurred between the aquatic insect faunas in Douglas Creek, Snively Springs, and Rattlesnake Springs. Dipterans, midges, and blackflies were very important in all study streams and accounted for 40% to 70% of the total annual community production. Also, several insect taxa had short development times and multiple cohorts in each of

the study streams. Gray (1981) suggested that rapid development may be advantageous in streams subject to flash floods. Flash floods occur in all of the study streams [approximately every four years in Rattlesnake Springs (Cushing and Wolfe 1982)].

Functional Groups and Trophic Levels

The River Continuum Concept (RCC) provides a template to which various physical factors of a stream can be compared for predicting biotic responses (Vannote et al. 1980). An important factor affecting the stream biota is the amount of sunlight reaching the substrate for primary production. The amount of light is influenced by shading from riparian vegetation and instream autotrophs. The biota in Snively Springs and Rattlesnake Springs are greatly influenced by this interaction, as riparian vegetation and watercress shade much of the substrate. The substrate of Snively Springs and Rattlesnake Springs is mainly sand/silt (see Table 1), limiting the area available for algae to attach and grow. Based upon RCC assumptions, a lack of grazer-scrappers that rely on algae and periphyton as a food source would be expected. This occurred in Snively Springs and Rattlesnake Springs where grazer-scrappers were completely absent.

Douglas Creek, however, is larger and shading from riparian vegetation affected a smaller portion of the stream cross section. Watercress is limited to the stream banks, leaving more water open to receive sunlight. The substrate is rocky (see Table 1), which provides habitat for algae. As suggested by the RCC, grazer-scrappers are an important functional group.

The RCC predicts a significant proportion of shredders to be present when the riparian vegetation covers the stream and contributes large amounts of allochthonous detritus. However, shredders were of little importance in any of the streams I studied. The reasons for this are unclear and require further investigation. Cushing (in press) found the amount of allochthonous detritus entering these streams to be low, indicating a lack of food for shredders. Fisher and Gray (1983) found

that shredders were virtually absent from the Sonoran Desert stream, Sycamore Creek, as was their food source of large particulate organic matter.

Detritivores were dominant in all study streams, indicating a large dependence upon detritus as a food source. Collecting (filtering and gathering) was the major feeding strategy used to obtain detritus, and collectors generally consume Fine Particulate Organic Matter (FPOM) (Merritt and Cummins 1984). The conclusion can then be inferred that detritus in the form of FPOM was the major food source of the insects in these study streams. This conclusion is supported by the results of my analyses of gut contents and the results of Cushing and Rader (1982).

COMPARISON WITH OTHER STREAMS

Annual P/B ratios ranged from 3.6 to 121.7 for insects from the three study streams. The high annual P/B ratios are attributed to insects with rapid development and multiple cohorts (e.g., many Chironomidae). The annual P/B ratios that I calculated for cold-desert streams are generally lower than those reported by Jackson and Fisher (1986) for Sonoran Desert stream insects and by Benke *et al.* (1984) for southeastern blackwater stream insects. Their study streams are warmer and insect development occurs faster, resulting in a greater number of cohorts. The annual P/B ratios in my study were generally higher, however, than those reported by Krueger and Waters (1983) for northern temperate streams. Northern temperate streams are generally cooler and insect development occurs at a slower rate with fewer cohorts.

Community production rates ranged from 14 to 23 g DW/m²/yr in my three study streams. These rates are lower than the 50 to 70 g DW/m²/yr in southeastern blackwater streams (Nelson and Scott 1962, Benke *et al.* 1984), and much lower than in the Sonoran Desert stream, Sycamore Creek, where community production exceeded 120 g DW/m²/yr (Fisher and Gray 1983, Jackson and Fisher 1986). However, the rates reported here are

generally higher than the 5 to 15 g DW/m²/yr reported for northern temperate streams (Fisher and Likens 1973, Neves 1979, Krueger and Waters 1983).

CONCLUSIONS

In conclusion, I want to stress the most important findings of my investigation.

- 1) Taxa with short development times and multiple cohorts, such as midges and blackflies, are very important to cold-desert streams in terms of total production. This life history strategy may be advantageous in streams such as the ones I studied, where flash floods occur.
- 2) Insects in cold-desert streams depend largely on detritus as a food source and obtain it mainly by collecting (filtering and gathering).
- 3) Community production rates in cold-desert streams are generally higher than in northern temperate streams, lower than in southeastern blackwater streams, and much lower than in Sonoran Desert streams. Stream temperature is an important factor (maybe the most important) in controlling production rates and causing rates to differ across physiographic regions.
- 4) Investigators should calculate secondary production, not density or biomass, to accurately assess the roles of organisms in aquatic ecosystems. In Douglas Creek, for example, collector and grazer-scraper importance was misrepresented by density and biomass analysis.
- 5) The framework given by Benke et al. (1984) to estimate secondary production provides a workable process, and if followed in other studies, will make comparisons between streams more meaningful.

More research needs to be conducted on cold-desert streams before their ecology is fully understood. This study provides a significant piece to the complex puzzle of these unique ecosystems.

REFERENCES

- Anderson, R. O. 1959. A modified flotation technique for sorting bottom fauna samples. *Limnology and Oceanography* 4:223-225.
- Becker, C. D. 1973. Development of Simulium vittatum Zett. (Diptera: Simuliidae) from larvae to adults at thermal increments from 17.0 to 27.0 C. *American Midland Naturalist* 89:246-251.
- Behmer, D. J. and C. P. Hawkins. 1986. Effects of overhead canopy on macro-invertebrate production in a Utah stream. *Freshwater Biology* 16:287-300.
- Benke, A. C. and J. B. Waide. 1977. In defense of average cohorts. *Freshwater Biology* 7:61-63.
- Benke, A. C. 1979. A modification of the Hynes method for estimating secondary production with particular significance for multivoltine populations. *Limnology and Oceanography* 24:168-171.
- Benke, A. C. and J. B. Wallace. 1980. Trophic basis production among net-spinning caddisflies in a southern Appalachian stream. *Ecology* 61:108-118.
- Benke, A. C., T. C. VanArsdall Jr. and D. M. Gillespie. 1984. Invertebrate productivity in a subtropical blackwater river: importance of habitat and life history. *Ecological Monographs* 54:25-63.
- Clifford, H. F. 1982. Life cycles of mayflies (Ephemeroptera), with special reference to voltinism. *Quaestiones Entomologicae* 18:15-90.
- Cummins, K. W. 1962. An evaluation of some techniques for the collection and analysis of benthic samples with special emphasis on lotic waters. *American Midland Naturalist* 67:477-504.
- Cushing, C. E., C. D. McIntire, J. R. Sedell, K. W. Cummins, G. W. Minshall, R. C. Peterson and R. L. Vannote. 1980. Comparative study of physical-chemical variables of streams using multivariate analysis. *Archiv fur Hydrobiologie* 89:343-352.
- Cushing, C. E. and R. T. Rader. 1982. A note on the food of Callibaetis (Ephemeroptera:Baetidae). *Great Basin Naturalist* 41:431-432.
- Cushing, C. E. and E. G. Wolf. 1982. Organic energy budget of Rattlesnake Springs, Washington. *American Midland Naturalist* 107:404-407.
- Cushing, C. E. and E. G. Wolf. 1984. Primary production in Rattlesnake Springs, a cold-desert spring-stream. *Hydrobiologia* 114:229-236.
- Cushing, C. E. Allochthonous detritus input to a small, cold desert spring-stream. *Verhandlungen Internationale Vereinigung Limnologie* 23:(in press).

- Fisher, S. G. and G. E. Likens. 1973. Energy flow in Bear Brook, New Hampshire: an integrative approach to stream ecosystem metabolism. *Ecological Monographs* 43:421-439.
- Fisher, S. G. and L. J. Gray. 1983. Secondary production and organic matter processing by collector macroinvertebrates in a desert stream. *Ecology* 64:1217-1224.
- Gray, L. J. 1981. Species composition and life histories of aquatic insects in a lowland Sonoran desert stream. *American Midland Naturalist* 106:229-242.
- Hamilton, A. L. 1969. On estimating annual production. *Limnology and Oceanography* 14:771-782.
- Hynes, H. B. N. and M. J. Coleman. 1968. A simple method of assessing the annual production of stream benthos. *Limnology and Oceanography* 13:569-573.
- Hynes, H. B. N. 1970. "The Ecology of Running Waters." University of Toronto Press, Ontario, Canada.
- Hynes, H. B. N. 1980. A name change in the secondary production business. *Limnology and Oceanography* 25:778.
- Jackson, J. K. and S. G. Fisher. 1986. Secondary production, emergence, and export of aquatic insects of a Sonoran desert stream. *Ecology* 67:629-638.
- Krueger, C. C. and T. F. Waters. 1983. Annual production of macroinvertebrates in three streams of different water quality. *Ecology* 64:840-850.
- Mackey, A. P. 1977. Growth and development of larval Chironomidae. *Oikos* 28:270-275.
- McAuliffe, J. R. 1982. Behavior and life history of Leucotrichia pictipes (Banks) (Trichoptera: Hydroptilidae) with special emphasis on case reoccupancy. *Canadian Journal of Zoology* 60:1557-1561.
- McCullough, D. A., G. W. Minshall and C. E. Cushing. 1979. Bioenergetics of a stream "collector" organism, Tricorythodes minutus (Insecta: Ephemeroptera). *Limnology and Oceanography* 24:45-58.
- Merritt, R. W. and K. W. Cummins. 1984. "An Introduction to the Aquatic Insects of North America." 2nd Ed. Kendall/Hunt Publishing Company, Dubuque, Iowa.
- Nelson, D. J. and D. C. Scott. 1962. Role of detritus in the productivity of a rock out-crop community in a Piedmont stream. *Limnology and Oceanography* 7:396-413.
- Neves, R. J. 1979. Secondary production of epilithic fauna in a woodland stream. *American Midland Naturalist* 102:209-224.

- O'Hop, J., J. B. Wallace and J. D. Haefer. 1984. Production of a stream shredder, Peltoperla maria (Plecoptera: Peltoperlidae) in disturbed and undisturbed hardwood catchments. *Freshwater Biology* 14:13-21.
- Smock, L. A., E. Gilinsky and D. L. Stoneburner. 1985. Macroinvertebrate production in a southwestern United States blackwater stream. *Ecology* 66:1491-1503.
- Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell and C. E. Cushing. 1980. The River Continuum Concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-137.
- Waters, T. F. 1977. Secondary production in inland waters. *Advances in Ecological Research* 10:91-164.
- Waters, T. F. and J. C. Hokenstrom. 1980. Annual production and drift of the stream amphipod Gammarus pseudolimnaeus in Valley Creek, Minnesota. *Limnology and Oceanography* 25:700-710.
- Welch, P. S. 1935. "Limnology." McGraw-Hill, New York and London.

APPENDIX

INSECTS FOUND IN DOUGLAS CREEK, SNIVELY SPRINGS, AND RATTLESNAKE SPRINGS

APPENDIX

INSECTS FOUND IN DOUGLAS CREEK, SNIVELY SPRINGS, AND RATTLESNAKE SPRINGS

DOUGLAS CREEK

Diptera

Chironomidae

Brilla flavifrons sp.

Chaetocladius sp.

Chironomus sp.

Heleniella sp.

Parametriocnemus sp.

Phaenospectra sp.

Polypedilum sp.

Thienemannimyia sp.

Empididae

Simuliidae

Simulium sp.

Tabanidae

Tipulidae

Trichoptera

Hydroptilidae

Leucotrichia pictipes

Hydropsychidae

Cheumatopsyche sp.

Hydropsyche sp.

Coleoptera

Elmidae

Optioservus sp.

Ephemeroptera

Baetidae

Baetis sp.

Leptophlebiidae

Paraleptophlebia sp.

Heptageniidae

Leucrocuta sp.

Tricorythidae

Tricorythodes sp.

Odonata

Coenagrionidae

Argia tibialis

Plecoptera

Perlodidae

Isoperla sp.

SNIVELY SPRINGS

Diptera

Chironomidae

Chaetocladius sp.

Chironomus sp.

Heleniella sp.

Polypedilum sp.

Thienemannimyia sp.

Dixidae

Empididae

Simuliidae

Simulium sp.

Tabanidae

Tipulidae

Trichoptera

Hydropsychidae

Cheumatopsyche sp.

Ephemeroptera

Baetidae

Baetis sp.

Leptophlebiidae

Paraleptophlebia sp.

Odonata

Coenagrionidae

Argia tibialis

RATTLESNAKE SPRINGS

Diptera

Chironomidae

Chaetocladius sp.

Chironomus sp.

Heleniella sp.

Polypedilum sp.

Thienemannimyia sp.

Dixidae

Empididae

Simuliidae

Simulium sp.

Tabanidae

Tipulidae

Trichoptera

Hydropsychidae

Cheumatopsyche sp.

Parapsyche sp.

Limnephilidae

Limnephilus sp.

Coleoptera

Dytiscidae

Hydaticus sp.

Hydrophilidae

Ephemeroptera

Baetidae

Baetis sp.

Tricorythidae

Tricorythodes sp.

Odonata

Coenagrionidae

Argia tibialis

DISTRIBUTION

<u>No. of Copies</u>		<u>No. of Copies</u>
	<u>OFFSITE</u>	
30	DOE Technical Information Center	H. F. Clifford Department of Zoology University of Alberta Edmonton, Alberta CANADA T6G 2E3
	J. D. Allan Department of Zoology University of Maryland College Park, MD 20742	W. Coffman Department of Life Sciences University of Pittsburgh Pittsburgh, PA 15260
	N. Anderson Department of Entomology Oregon State University Corvallis, OR 97331	S. Cooper Department of Biological Sciences University of California Santa Barbara, CA 93106
	B. Barker Department of Biology Central Washington University Ellensburg, WA 98926	A. Covich Zoology Department University of Oklahoma Norman, OK 73019
	J. Barnes Department of Zoology Brigham Young University Provo, UT 84602	K. Cummins Appalachian Environmental Laboratory University of Maryland Frostburg, MD 21532
	A. C. Benke Department of Biology University of Alabama Tuscaloosa, AL 35486	D. Dudgeon Department of Zoology University of Hong Kong HONG KONG
	E. F. Benfield Department of Biology VPI and State University Blacksburg, VA 24061	P. Dumas Department of Biology Central Washington University Ellensburg, WA 98926
	A. Brown Zoology Department University of Arkansas Fayetteville, AR 72701	J. W. Elwood Environmental Sciences Division Oak Ridge National Laboratory Building 2001 Oak Ridge, TN 37830
	D. A. Bruns Environmental Sciences Section P.O. Box 1625 Idaho Falls, ID 83415	

No. of
Copies

S. Fisher
Department of Zoology
Arizona State University
Tempe, AZ 85281

10 W. L. Gaines
P.O. Box 428
Winthrop, WA 98862

S. Gregory
Department of Fisheries and
Wildlife
Oregon State University
Corvallis, OR 97331

M. Gurtz
Water Resources Division
U.S. Geological Survey
P.O. Box 2857
Raleigh, NC 27602

D. Hart
Academy of Natural Sciences
19th and The Parkway
Philadelphia, PA 19103

R. Hauer
Flathead Lake Biological
Station
East Shore
Bigfork, MT 59911

C. Hawkins
Department of Fisheries and
Wildlife
Utah State University - UMC52
Logan, UT 84322

N. Hedges
Bureau of Land Management
Wenatchee, WA 98801

B. Hilger
Research Institute for Nature
Management
P.O. Box 46
3956 ZR Leersum
THE NETHERLANDS

No. of
Copies

H. B. N. Hynes
Department of Biology
University of Waterloo
Waterloo, Ontario
CANADA N2L 3G1

S. Jackson
639 Crestview Dr.
Moses Lake, WA 98837

P. S. Lake
Department of Zoology
Monash University
Clayton
Victoria 3168
AUSTRALIA

G. Lamberti
Department of Fisheries
and Wildlife
Oregon State University
Corvallis, OR 97331

R. Mackay
Department of Zoology
University of Toronto
Toronto, Ontario
CANADA M5S 1A1

G. R. Marzolf
Division of Biology
Kansas State University
Manhattan, KS 66506

R. Merritt
Department of Entomology
Michigan State University
East Lansing, MI 48824

G. W. Minshall
Department of Biology
Idaho State University
Pocatello, ID 83209

M. Oswood
Department of Biology
University of Alaska
Fairbanks, AK 99701

No. of
Copies

B. Peckarsky
Department of Entomology
Cornell University
Ithaca, NY 14853

I. Pettman, Librarian
Freshwater Biological
Association
The Ferry House
Ambleside
Cumbria LS22 0LP
ENGLAND

R. Petersen
Department of Limnology
University of Lund
S-22003 Lund
SWEDEN

S. R. Reice
Department of Zoology
University of Northern Carolina
Chapel Hill, NC 27514

V. Resh
Department of Entomology
University of California
Berkeley, CA 94720

J. Sedell
Forest Sciences Laboratory,
USFS
Oregon State University
Corvallis, OR 97331

A. L. Sheldon
Department of Zoology
University of Montana
Missoula, MT 59812

10 S. Smith
Department of Biology
Central Washington University
Ellensburg, WA 98926

No. of
Copies

J. Stanford
Flathead Lake Biological
Station
East Shore
Bigfork, MT 59911

B. Statzner
Zoolog. Inst. d. Univ. (T.H.)
Kornblumenstr. 13
Postfach 6380
7500 Karlstuhel
FEDERAL REPUBLIC OF GERMANY

B. Sweeney
Stroud Water Research Center
RD #1, Box 512
Avondale, PA 19311

R. Vannote
Stroud Water Research Center
RD #1, Box 512
Avondale, PA 19311

J. B. Wallace
Department of Entomology
University of Georgia
Athens, GA 30602

G. M. Ward
Department of Biology
University of Alabama
Tuscaloosa, AL 35486

J. Ward
Department of Zoology and
Entomology
Colorado State University
Fort Collins, CO 80523

T. Waters
Department of Entomology,
Fisheries and Wildlife
University of Minnesota
St. Paul, MN 55455

No. of
Copies

J. Webster
Department of Biology
VPI and State University
Blacksburg, VA 24061

C. Wiberg
Department of Biology
Central Washington University
Ellensburg, WA 98926

M. Winterbourn
Department of Zoology
University of Canterbury
Christchurch
NEW ZEALAND

ONSITE

DOE Richland Operations Office

J. J. Sutey

48 Pacific Northwest Laboratory

C. D. Becker
C. E. Cushing (25)

No. of
Copies

Pacific Northwest Laboratory
(contd)

D. D. Dauble
D. W. Dragnich
D. H. Fickeisen
M. J. Graham
J. M. Hales
P. C. Hays
P. J. Mellinger
D. A. Neitzel
R. L. Newell
T. L. Page
T. M. Poston
W. H. Rickard
L. E. Rogers
J. A. Stottlemire
R. E. Wildung
Publishing Coordination (2)
Technical Report Files (5)