Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA-UR--90-3395

DE91 002307

TITLE: LABORATORY SPECTRA OF AMORPHOUS AND CRYSTALLINE OLIVINE:
AN APPLICATION TO COMET HALLEY IR SPECTRUM

AUTHOR(S): A. Blanco, V. Orofino, E. Bussoletti, S. Fonti, L. Colangeli,
and J. R. Stephens

SUBMITTED TO: Proceedings of the 126th IAU Colloquium "Origin and Evolution
of Interplanetary Dust", August 27-30, 1990, Kyoto, Japan

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

OSTI
1990

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

LABORATORY SPECTRA OF AMORPHOUS AND CRYSTALLINE OLIVINE: AN APPLICATION TO COMET HALLEY IR SPECTRUM

A. BLANCO, V. OROFINO

Department of Physics, University of Lecce,
C. P. 193, 73100 Lecce, Italy

E. BUSSOLETTI, S. FONTI

Via Acton 38, 80133, Napoli, Italy

L. COLANGELI

Engineering Facility, University of Cassino
Via Zamosch 43, 03043, Cassino, Italy

J. R. STEPHENS

Los Alamos National Laboratory
Los Alamos, NM 87545 USA

ABSTRACT. Among the various silicates proposed as components of cosmic dust grains, olivine is considered one of the most likely materials. In this work we present the infrared spectra of three different types of olivine grains: crystalline, amorphous, and synthetic (also amorphous). While the first and second sample derive from the same natural mineral, the third one has been prepared in laboratory according to the relative cosmic abundances of the elements. The experiment data are used to fit the emission feature observed in the comet Halley spectrum between 8 and 13 μm . Satisfactory results are obtained by using synthetic olivine mixed with a small amount (5%) of crystalline grains.

1 Introduction

Silicates have been identified as components of interstellar and interplanetary dust on the basis of agreement in the position of observed bands near 10 and 20 μm (for a recent review see Knacke, 1989).

The IR spectra of glassy silicates seem to provide a closer match to astronomical observations than crystalline minerals (Knacke, 1978; Kratschmer and Huffman, 1979). We have undertaken a research program to measure the optical properties of glassy silicates having composition similar to minerals predicted in theoretical condensation sequences (Stephens et al., 1989). Among the samples examined, one of the most "popular" is olivine, which has also been recently proposed as an important component of cometary dust (Bregman et al., 1987; Campins and Ryan, 1989). In this work we fit the broad emission feature present in the comet Halley spectrum between 8 and 13 μm , using laboratory spectral data of natural (both glassy and crystalline) and synthetic amorphous olivine with chemical composition reflecting the cosmic abundances of the various elements.

2 Sample preparation and IR spectra

The crystalline samples for IR spectroscopy have been prepared by grinding the natural minerals and embedding the resulting submicron particles in KBr pellets. Amorphous samples have been obtained by vaporizing the same materials by means of a LAMBDA PHYSIK LPX 315i excimer laser operating at a wavelength of 308 nm. The repetition rate was 9 Hz with an energy of 400 mJ/pulse. The condensed smokes were directly collected on KRS-5 substrates inside a sample chamber which was filled O_2 at a pressure of 10 mbar (this work) or 1 atmosphere (Stephens, 1980). No significant changes were, however, detected by varying the gas in the chamber.

We prepared synthetic samples because available olivine crystals, although they have a metal to silicon ratio very close to that found in typical O-rich stars, exhibit a ratio $\text{Mg}/\text{Fe} = 9$. This ratio is about an order of magnitude larger than the cosmic value, and the composition (including the metal atoms) and the metal to silicon ratio are among the most important factors affecting the IR spectra. We produced and analyzed two synthetic olivine-type silicates with the appropriate cosmic elemental content:

In spite of the presence of Al, Ca, and Na in the second sample, their IR spectra are essentially coincident. The targets for the laser were prepared starting from the following reagent grade materials:

<u>Element</u>	<u>Beginning Compound</u>	<u>Compound</u>
Mg	MgCO ₃	MgO
Fe	Fe ₂ O ₃	Fe ₂ O ₃
Si	Silicic Acid	SiO ₂
Ca	CaCO ₃	CaO
Na	NaCO ₃	NaCO ₃
Al	Gamma-Al ₂ O ₃	Gamma-Al ₂ O ₃

Each material was heated in successive steps up to 1000 C to convert the material to the oxide. NaCO₃ was not heated, and Fe₂O₃ and Al₂O₃ were heated only to dry the materials. Mixtures of powders for the synthetic samples were pressed in air (10 ton). Each pellet was then heated in a radio frequency furnace in an Ar atmosphere to temperatures up to 1400 C until the sample just melted (approximately 1 minute). The IR spectra of the three different olivine particles are shown in Fig. 1.

3 An application to comet Halley

The above laboratory work has been used to fit the 8-13 μm spectrum of comet Halley. Its broad emission feature has already been attributed to olivine (Bregman et al., 1987; Campins and Ryan, 1989).

The model we used to fit the cometary flux is an adaptation of that originally applied to circumstellar envelopes (Orofino et al., 1987; Blanco et al., 1989). The free parameters are: radius and temperature of the nucleus (R_n, T_n), radius and temperature of the dust envelope (R_d, T_d) and the total optical thickness (t) of the dust at 8 μm . The values of some of these parameters are available in literature. From our preliminary results, shown in Fig. 2, it seems that the Halley 10 μm feature can be satisfactory fitted by a mixture of amorphous and crystalline olivine. We note however that a better fit is obtained by using the synthetic olivine

with cosmic elemental composition. Moreover, in this case, a far smaller amount of crystalline component is needed. This can be important in the study of cometary evolution and its connections with interplanetary dust particles (Greenberg, 1990; Campins and Ryan, 1989; Sandford, 1989). It is worthwhile to note that laboratory IR spectra of crystalline olivine are usually taken by embedding dust samples in a matrix. This experimental procedure may change both shape and position of spectral features (Bohren and Huffman, 1983; Dorschner et al., 1977). At present laboratory work is in progress to account for this effect.

REFERENCES

Blanco, A., Borghesi, A., Bussoletti, E., Colangeli, L., Fonti, S., and Orofino, V.: 1988, in *Experiments on Cosmic Dust Analogues*, eds. E. Bussoletti, C. Fusco, G. Longo (Kluwer Academic Publishers, Dordrecht) p. 167.

Bohren, C. F. and Huffman, D. R.: 1983, *Absorption and Scattering of Light by Small Particles* (Wiley, New York).

Bregman, J. D., Campins, H., Witteborn, F. C., Wooden, D. H., Rank, D. M., Allamandola, L. J., Cohen, M. and Tielens, A. G. G. M.: 1987, *Astron. J.*, 187, 616.

Campins, H. and Ryan E. V.: 1989, *Ap. J.*, 341, 1059.

Dorschner, J., Friedemann, C. and Gurtler, J.: 1977, *Astron. Nachr.*, 298, 2798.

Greenberg, J. M.: 1990 (private communication).

Knacke, R. F.: 1978, in *Protostars and Planets*, ed. T. Gehrels (Univ. Arizona Press) p. 112.

Knacke, R. F.: 1989, in *Interstellar Dust*, eds. L. J. Allamandola and A. G. G. M. Tielens (Kluwer Academic Publishers, Dordrecht) p. 415.

Dorschner, W. and Huffman, D. R.: 1979, *Astrophys. Space Sci.*, 61, 195.

Orofino, V., Colangeli, L., Bussoletti, E., and Strafella, F.: 1987, *Astrophys. Space Sci.*, 138, 127.

Sandford, S. A. and Walker, R. M.: 1985, *Ap. J.* 291, 838.

Sandord, S. A.: 1989, in *Interstellar Dust*, eds. L. J. Allamandola and A. G. G. M. Tielens (Kluwer Academic Publishers, Dordrecht) p. 403.

Stephens, J. R.: 1980, *Ap. J.*, 237, 450.

Stephens, J. R., Blanco, A. Borghesi, Al, Fonti, S. and Bussoletti, E.: 1989, in *Interstellar Dust Contributed Papers*, eds. A. G. G. M. Tielens and L. J. Allamandola, NASA CP-3036, p. 375.

Figure 1:

Extinction efficiencies of the three different types of olivine examined in the present work: crystalline (solid line), amorphous (dotted line), and synthetic (dashed line). The curves are normalized at their maximum while a is the particle radius.

Figure 2:

Best fit of the observed (Campins and Ryan, 1989) Halley spectrum (dots) with laboratory data. The dashed curve is obtained for a 50% mixture of amorphous and crystalline olivine (best fit parameters are: $T_n = 20$ K, $T_d = 380$ K, $R_d/R_n = 1000$, $A_c = 1.086t = 0.5$ mag). The solid curve is obtained by using synthetic olivine with only 5% of crystalline olivine. Best fit parameters are the same except that $A_c = 1.2$

Fig. 1

λ (μm)

Fig. 2.

END

DATE FILMED

11 / 30 / 90