AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

48
6-8-93 JS (2)

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2907
UC-421,427

PPPL-2907

THEORY OF KINETIC BALLOONING MODES EXCITED BY
ENERGETIC PARTICLES IN TOKAMAKS

BY

S. TSAI AND L. CHEN

MAY, 1993

PRINCETON
PLASMA PHYSICS
LABORATORY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced from the best available copy.
Available in paper copy and microfiche.

Number of pages in this report: 20

DOE and DOE contractors can obtain copies of this report from:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
(615) 576-8401.

This report is publicly available from the:

National Technical Information Service
Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4650

Theory of Kinetic Ballooning Modes Excited by Energetic Particles in Tokamaks

Shih-Tung Tsai*

and

Liu Chen

Princeton University, Plasma Physics Laboratory

and

Department of Astrophysical Sciences

P.O. Box 451

Princeton, N.J. 08543

Abstract

We have analyzed theoretically the resonant excitations of kinetic ballooning modes (KBM) by the energetic ions/alpha particles in tokamaks. Our theory includes finite-size orbit effects of both circulating and trapped particles. With energetic-particle contributions suppressed in the singular inertial layer, an analytic dispersion relation can then be derived via an asymptotic matching analysis. The dispersion relation, in particular, demonstrates the existence of two types of modes; that is, the magnetohydrodynamic (MHD) gap mode and the energetic-particle continuum mode. Specific expressions for real frequencies, growth rates and threshold conditions are also derived for a model slowing-down beam ion distribution function.

*permanent address: Inst. of Physics, Chinese Academy of Sciences, Beijing 100080, PRC

I. Introduction

The dynamics of energetic ions/ α particles is an important issue for fusion research and has attracted, in recent years, increasing experimental¹⁻³ as well as theoretical⁴⁻¹³ attention. There exists a considerable amount of works devoted to the study of how the energetic particles affect the stability of MHD modes. In particular, it has been demonstrated that MHD modes can be driven unstable via resonances with energetic particles.⁷⁻¹³ Most of previous theoretical analyses have assumed that the energetic particles have negligible orbit sizes. In the presence of sharply localized singular-layer structures, this assumption is, at best, dubious. Only recently, the stabilizing effects of finite-size orbits have been included in the analyses of toroidal Alfvén eigenmodes (TAE).^{14,15}

In this paper, we examine finite-size orbit effects of energetic particles on KBM instabilities. Noting that KBM consists of a singular inertial and a regular ideal regions, the finite-size orbits then render energetic particles effective only in the ideal region. Employing an asymptotic matching analysis, a general dispersion relation can then be derived analytically for both circulating and trapped particles; including corresponding various wave-particle resonances. This dispersion relation has the generic form of the “fishbone” dispersion relation,⁷⁻⁹ and demonstrates, in particular, the existence of two types of modes.⁸ One is the MHD gap mode which has a frequency in the order of ion diamagnetic drift frequency and can be destabilized by the energetic particles only when the plasma is ideal MHD unstable. The other is the energetic-particle continuum mode; which, however, has a threshold condition and the frequency corresponds to that of energetic-particle characteristic frequency.

Section II presents the theoretical model and the relevant eigenmode equations. The corresponding dispersion relation is then obtained via asymptotic matching in Sec. III. In Sec. IV, we analyze the general stability properties of KBM and derive contributions due to circulating and trapped energetic particles. Specific results of a model slowing-down beam ion distribution function are given in Sec. V. Finally, Sec. VI contains a brief summary and discussions.

II. Theoretical Model and Eigenmode Equations

We consider a large-aspect ratio (i.e. $\epsilon = a/R_0 \ll 1$) tokamak plasma consisting of a warm core (C) component and an energetic (E) component. Here, a and R_0 are, respectively, the minor and major radii. In terms of the smallness parameter ϵ , we adopt the following formal orderings: $\beta_C \equiv 8\pi P_C/B^2 \sim \epsilon$, $\beta_E \sim \epsilon^2$, $T_C/T_E \sim \epsilon^2$, $n_E/n_C \sim \epsilon^3$, $k_\theta \rho_{iC} \sim \epsilon^{3/2}$, $k_\theta \rho_E \sim \epsilon^{1/2}$, and $\omega/\omega_A \sim \epsilon$. Here, $\omega_A = V_A/qR_0$, q is the safety factor and the rest of the notations is standard. We note that the present orderings are adopted such that effects of finite-size orbits, while suppress energetic-particle dynamics in the inertial layer, affect the core component negligibly. Furthermore, to make analytical calculations of the KBM with finite $k_\theta \rho_E$ possible, we adopt the Connor-Hastie-Taylor analytical model equilibrium of circular magnetic surfaces.¹⁶

Within the present model, the ideal MHD approximation can be shown to remain valid; i.e. $\delta E_\parallel \simeq 0$. Meanwhile, perpendicular pressure balance gives

$$B\delta B_\parallel \simeq 4\pi n_i e \omega_{*p} \frac{\delta \phi}{\omega}, \quad (1)$$

where $n_i e \omega_{*p} \equiv (c/B)(\mathbf{k} \times \mathbf{e}_\parallel) \cdot \nabla P$, and $\delta \phi$ is the scalar potential. The eigenmode equation then corresponds to the following vorticity equation:

$$\left[\frac{d}{d\theta} f \frac{d}{d\theta} + \Omega^2 f + \alpha g \right] \delta \phi - \sum_j \eta_j \langle g \Omega_d J_0(k_\perp \rho) \delta G \rangle_j = 0, \quad (2)$$

where we have used the well-known ballooning mode representation. Thus, θ is the extended coordinate along the magnetic field $\mathbf{B}$, $\Omega^2 \equiv \omega(\omega - \omega_{*pi})/\omega_A^2$, $f(\theta) = 1 + (s\theta - \alpha \sin \theta)^2$, $s = r q'/q$, $\alpha = q^2 R_0 \beta'$, $g(\theta) = \cos \theta + (s\theta - \alpha \sin \theta) \sin \theta$, $\eta_j = 4\pi e_j q^2 R_0^2 \omega/c^2$, $\omega_{dj}(\theta) = k_\theta \Omega_{dj} g(\theta)$, $\Omega_{dj} = (v_\perp^2/2 + v_\parallel^2)/\omega_{cj} R_0$, and $\langle \dots \rangle_j \equiv \int d^3 v (\dots)_j$ with j being the relevant species. The function δG_j corresponds to the nonadiabatic gyrokinetic response and is governed by the following gyrokinetic equation:

$$\left[\omega_t \frac{d}{d\theta} - i(\omega - \omega_d) \right]_j \delta G_j = i \left(\frac{e}{m} \right)_j Q F_{oj} g \Omega_{dj} J_0(\lambda_\rho) \frac{\delta \phi}{\omega}, \quad (3)$$

where $\omega_t = v_\parallel/qR_0$, $Q F_{oj} = (\omega \partial_\epsilon + \hat{\omega}_*) F_{oj}$, $\epsilon = v^2/2$, $\lambda_\rho = k_\perp \rho$, and $\hat{\omega}_* F_{oj} = (\mathbf{k} \times \mathbf{e}_\parallel / \omega_{cj}) \cdot \nabla F_{oj}$. Note that, in Eq. (3), the left-hand side stands

for the guiding-center propagator and, hence, determines the wave-particle resonance conditions, QF_{oj} corresponds to the free energy in both configuration and velocity spaces, and $\Omega_{dj}\delta\phi$ indicates energy exchange via $V_{dj} \cdot \delta E$ with V_{dj} being the magnetic drift.

For the present analysis, it is more convenient to let $\delta\psi = f^{1/2}\delta\phi$ and rewrite Eq. (2) in the following standard form:

$$\left[\frac{d^2}{d\theta^2} + \Omega^2 + \frac{\alpha \cos \theta}{f} - \frac{(s - \alpha \cos \theta)^2}{f^2} \right] \delta\psi - f^{-1/2} \sum_j \eta_j < g\Omega_d J_0(\lambda_\rho) \delta G >_j = 0. \quad (4)$$

Equations (3) and (4) correspond to the coupled differential-integral eigenmode equations and are generally difficult to solve analytically. Significant progress, however, can be made by noting that the basic mode structures contain two spatial regions; that is the long-scale [$\theta \sim O(\omega_A/\omega) \sim \epsilon^{-1}$] inertial and the short-scale [$\theta \sim O(1)$] ideal regions. Analytical dispersion relation can then be derived by asymptotically matching the solutions in the two regions, and the details are given in the next Section.

III. Dispersion Relation

In the inertial region, we have $|\theta| \sim O(\epsilon^{-1}) \gg 1$. It can then be straightforwardly demonstrated that, with the formal $|k_\theta \rho_E| \sim \epsilon^{1/2}$ ordering, energetic-particle nonadiabatic contributions are negligible due to the finite size orbit effects. Core particle contributions, meanwhile, remain essentially the same as the negligible orbit-size case^{12,13} and, hence, will be omitted in the present analysis. Equation (4), therefore, reduces to

$$\left(\frac{d^2}{d\theta^2} + \Omega^2 \right) \delta\psi_0(\theta) = 0. \quad (5)$$

The solution is

$$\delta\psi_0(\theta) = e^{i\Omega|\theta|}, \quad (6)$$

with the $\text{Im}\Omega > 0$ causality constraint.

In the ideal region, meanwhile, we have $|\theta| \sim O(1)$, and Eq. (4) reduces to

$$\left[\frac{d^2}{d\theta^2} + \frac{\alpha \cos \theta}{f} - \frac{(\alpha \cos \theta - s)^2}{f^2} \right] \delta\psi_I - f^{-1/2} g\eta_E < \Omega_d J_0 \delta G >_E = 0. \quad (7)$$

From Eq. (7), we can construct the following quadratic form

$$\delta\psi_I^* \frac{d\psi_I}{d\theta} \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} d\theta \left[- \left| \frac{d\delta\psi_I}{d\theta} \right|^2 + \left(\frac{\alpha \cos \theta}{f} - \frac{(\alpha \cos \theta - s)^2}{f^2} \right) |\delta\psi_I|^2 \right] - \int_{-\infty}^{\infty} d\theta f^{-1/2} g \eta_E \delta\psi_I^* < \Omega_d J_0 \delta G >_E = 0. \quad (8)$$

Asymptotically matching $\delta\psi_I$ and $\delta\psi_0$ gives

$$\delta\psi_I^* \frac{d}{d\theta} \delta\psi_I \Big|_{-\infty}^{\infty} = 2i\Omega. \quad (9)$$

Substituting Eq. (9) into Eq. (8) yields the desired dispersion relation

$$-i\Omega + \delta W_f + \delta W_K = 0; \quad (10)$$

where δW_f is the well-known MHD fluid δW , i.e.,

$$\delta W_f = \frac{1}{2} \int_{-\infty}^{\infty} d\theta \left\{ \left| \frac{d\delta\psi_I}{d\theta} \right|^2 + \left[\frac{(s - \alpha \cos \theta)^2}{f^2} - \frac{\alpha \cos \theta}{f} \right] |\delta\psi_I|^2 \right\}; \quad (11)$$

δW_K is the kinetic δW due to the energetic-particle “compression”, i.e.,

$$\delta W_K = \frac{1}{2} \eta_E \int_{-\infty}^{\infty} d\theta < \delta S^* \delta G >_E, \quad (12)$$

and

$$\delta S = \Omega_d f^{-1/2} g J_0(\lambda_\rho) \delta\psi_I. \quad (13)$$

Note that the causality constraint requires

$$\delta W_f + \delta W_{Kr} < 0 \quad (14)$$

with $\delta W_{Kr} \equiv \text{Re}(\delta W_K)$. It is also interesting to note that the dispersion relation, Eq. (10), has the same form as that of the “fishbone” instability.⁷⁻⁹ We therefore, term it as the generic “fishbone” dispersion relation.

IV. General Stability Analyses

Since we are interested in the energetic particle-wave resonances, we let

$$\delta W_K = \delta W_{Kr} + i\delta W_{Ki}. \quad (15)$$

Furthermore, we let $\omega = \omega_r + i\gamma$ with $|\gamma/\omega_r| \ll 1$ for near marginal stability studies.

From Eq. (10), it is clear that the stability properties depend crucially on Ω . Noting that $\Omega^2 \equiv \omega(\omega - \omega_{*pi})/\omega_A^2$, we can then identify two types of modes⁸ depending on the sign of $\Omega^2[\omega = \omega_r]$. Specifically, for

$$\omega_r < 0 \quad \text{or} \quad \omega_r > \omega_{*pi}, \quad (16)$$

we have Ω being real and, hence, the continuum mode. On the otherhand, for ω_r inside the $(0, \omega_{*pi})$ ion diamagnetic gap, we have Ω being imaginary and, hence, the gap mode. In the following, we shall discuss the stability properties of these two types of modes separately.

For the continuum mode, ω_r is determined by the real part of dispersion relation, Eq.(10); i.e.,

$$\delta W_f + \delta W_{Kr} = -\gamma \frac{\partial \Omega_r}{\partial \omega_r} < 0. \quad (17)$$

The growth rate, meanwhile, is given by the imaginary part of the dispersion relation and can be written as

$$\gamma = (\delta W_{Ki} - \Omega_r) / \left(-\frac{\partial \delta W_{Kr}}{\partial \omega_r} \right). \quad (18)$$

The threshold condition is, thus,

$$\delta W_{Ki} = \Omega_r; \quad (19)$$

i.e., balancing the energetic-particle drive with the continuum damping. For the mode to be unstable, it requires $\delta W_{Ki} > \Omega_r$, and

$$\partial \delta W_{Kr} / \partial \omega_r < 0. \quad (20)$$

Since this mode is induced by the energetic particles, we shall further term it as the energetic particle continuum mode.

For the gap mode, meanwhile, the dispersion relation, Eq. (10), can be rewritten as

$$\Omega' + \delta W_f + \delta W_K = 0; \quad (21)$$

where $\Omega' \equiv -i\Omega = \sqrt{\omega(\omega_{*pi} - \omega)}/\omega_A$. The real part of Eq. (21), which determines ω_r , is

$$\Omega'_r + \delta W_f + \delta W_{Kr} = 0; \quad (22)$$

where the causality constraint implies $\Omega'_r > 0$. The imaginary part of the dispersion relation then gives rise to the growth rate

$$\gamma \simeq \frac{2\delta W_{Ki}(\delta W_f + \delta W_{Kr})}{\omega_{*pi} - 2\omega_r} \omega_A^2. \quad (23)$$

Noting that $\delta W_f + \delta W_{Kr} < 0$ and δW_{Ki} is positive definite, the gap mode can be driven unstable only when

$$\omega_{*pi} > \omega_r > \omega_{*pi}/2. \quad (24)$$

Note that the gap mode is basically a MHD mode, and hence, will be called as the MHD gap mode.

In the rest of this Section, we are going to derive specific expressions of δG and, hence, δW_K for circulating and trapped particles. To do this, we need to know $\delta\phi_I(\delta\psi_I)$. Observing that, with $\delta W_K \sim \epsilon$, the lowest-order $\delta\psi_I$ is sufficient for the present purpose. If we further assume $s \sim \alpha < 1$, it is then reasonable to adopt $\delta\psi_I = 1$.

A. Circulating particle contribution

It is more convenient to analyze the circulating particles via a straightforward transformation from the guiding centers to the drift centers. δW_K , Eq. (12), then becomes

$$\delta W_K = \frac{2\pi e q^2 R_0^2}{c^2} \omega \int_{-\infty}^{\infty} d\theta < \delta S_d^* \delta G_d >; \quad (25)$$

where $\theta \sim O(\theta_k \sim 1/sk_\theta \rho_E) \sim \epsilon^{-1/2} > 1$, $\delta G_d = \delta G \exp(i\theta_d)$, $\theta_d = -\lambda_d \cos \theta$, $\lambda_d = k_\perp \rho_d = k_\perp \Omega_d/\omega_t$, and

$$\delta S_d^* = \Omega_d f^{-1/2} g J_0(\lambda_\rho) \exp(-i\theta_d). \quad (26)$$

δG_d is given by the following drift-center gyrokinetic equation

$$(\omega_t \partial_\theta - i\omega) \delta G_d = i \frac{e}{m} \frac{Q F_0}{\omega} \Omega_d J_0(\lambda_\rho) \sin \theta_0 \exp(i\theta_d). \quad (27)$$

Here, we have explicitly denoted the subsidiary two scales; $\theta_0 \sim 0(1)$ and $\theta_k \sim O(\epsilon^{-1/2})$. Noting that $\partial_\theta \simeq \partial_{\theta_0}$ and

$$\Omega_d J_0 \sin \theta_0 e^{i\theta_d} = \Omega_d J_0 \sum_{n=1}^{\infty} 2(-1)^n i^{n+1} \frac{n J_n(\lambda_d)}{\lambda_d} \sin n\theta_0 \equiv \sum_{n=1}^{\infty} \delta S_{dn} \sin n\theta_0, \quad (28)$$

Eq. (27) can then be readily solved and we obtain

$$\delta G_d = \frac{e}{m} Q F_0 \sum_{n=1}^{\infty} \frac{\delta S_{dn}}{n^2 \omega_t^2 - \omega^2} (\sin n\theta_0 - i \frac{n\omega_t}{\omega} \cos n\theta_0). \quad (29)$$

Equation (29) into Eq. (25), δW_K for the circulating (untrapped) energetic particles then becomes

$$\delta W_{KU} = \frac{\pi e^2 q^2 R_0^2 \omega}{m c^2} \sum_{n=1}^{\infty} \left\langle \frac{Q F_0}{n^2 \omega_t^2 - \omega^2} \int d\theta |\delta S_{dn}|^2 \right\rangle. \quad (30)$$

Here, $|\delta S_{dn}| = 2\Omega_d J_0(\lambda_\rho) n J_n(\lambda_d) / \lambda_d$ and we have dropped the subscript E to simplify the notations.

Recalling that, from Eq. (19), we have threshold $\delta W_{Ki} = \Omega_r \simeq \omega_r / \omega_A$ for the energetic-particle continuum mode and, from Eq. (24), $\omega_r \sim \omega_{\pi pi} < \omega_{tE}$ for the MHD gap mode, we can then conclude that, in general cases, only the $n = 1$ term in Eq. (30) needs to be considered. Adopting Padé's approximant for the Bessel functions; i.e.,

$$J_0(\lambda_\rho) 2J_1(\lambda_d) / \lambda_d \simeq (1 + \Delta^2 k_\perp^2 / k_\theta^2)^{-1}; \quad (31)$$

where $\Delta^2 \equiv k_\theta^2(\rho^2 + \rho_d^2/2)/4$, Eq. (30) then becomes, keeping only the $n = 1$ term,

$$\delta W_{KU} \simeq \frac{\pi^2 e^2 q^2 R_0^2}{2 m c^2 s} \left\langle \frac{Q F_0 \Omega_d^2}{\Delta (1 + \Delta^2)^{3/2}} \frac{\omega}{\omega_t^2 - \omega^2} \right\rangle. \quad (32)$$

We note also that, in deriving Eqs. (29) and (30), we have assumed $\partial_\theta = \partial_{\theta_0}$ and, hence, ignored the θ_k -scale variations of λ_ρ , λ_d and, hence,

δS_{dn} . Keeping the $O(\epsilon^{1/2})\partial_{\theta_k}$ corrections leads to broadening of wave-particle resonance conditions due to finite-size orbits. Specifically, Eq. (32) is modified (see Appendix) to become

$$\delta W_{KU} = \frac{\pi^2 e^2 q^2 R_0^2}{4 mc^2} \left\langle \frac{Q F_0 \Omega_d^2}{\Lambda(1 + \Delta^2)^2} \int_0^\infty dx e^{-x} \left[\frac{1}{\omega_t - \omega + (\Lambda/2)\omega_t x} - \frac{1}{\omega_t + \omega - (\Lambda/2)\omega_t x} \right] \right\rangle; \quad (33)$$

where $\Lambda = s\Delta/(1 + \Delta^2)^{1/2}$. Note that the finite- Λ resonance broadening effects could, in principle, also enhance the instability drive.

B. Trapped Particle Contribution

Similar to the circulating particles, the kinetic contribution of the trapped energetic particles can be more readily derived in terms of banana centers as, omitting the subscript E,

$$\delta W_{KT} = 2\pi\eta B_0 \int d\mathcal{E} \int d\mu \sum_{l=-\infty}^{\infty} \oint \frac{d\theta}{|v_{||}|} \delta S_b^* \delta G_b; \quad (34)$$

where

$$\delta S_b^* \equiv \Omega_d f^{-1/2} g J_0(\lambda_\rho) \exp(-i\theta_b), \quad (35)$$

$\theta_b = \int^\theta (\omega_d - \bar{\omega}_d) d\tau$, $d\tau = qR_0 d\theta/|v_{||}|$, $\bar{A} \equiv \oint A d\tau/\tau_b$, $\tau_b = \oint d\tau$, $k_\perp^2 = 1 + (2\pi l + \theta)^2 s^2$, and δG_b is the solution of gyrokinetic Eq. (3) with $\delta G_b = \delta G e^{i\theta_b}$. The subscript b , thus, denotes the banana centers.

While one can substitute the known solutions of δG or δG_b (see Ref. 17) into Eq. (34) to derive a formal expression of δW_{KT} , we shall try to further simplify the analysis by assuming the energetic particles are deeply trapped between $2\pi l \pm \theta_T$. We then derive, for $|\omega| \sim |\bar{\omega}_{dE}| < |\omega_{bE}|$,

$$\delta W_{KT} \simeq \frac{8\pi^2 e^2 q R_0 B_0}{mc^2} \int d\mathcal{E} \int d\mu Q F_0 \Omega_d^2 \tau_b \left(\frac{1}{\bar{\omega}_d - \omega} \right). \quad (36)$$

In deriving Eq. (36), we have noted that, compared to the $l = 0$ term, $l \neq 0$ terms are negligible due to the f^{-1} factor. Note that only physics of the precessional drift is kept here.⁹

On the otherhand, for $|\omega - \bar{\omega}_{dE}| \sim |\omega_{bE}|$, the $l \neq 0$ terms dominate. In fact, similar to the circulating particles, we have two θ scales; i.e., $\theta_0 \sim O(\theta_T) \lesssim 1$ and $\theta_k \sim O(1/k_\theta \rho_b) \sim \epsilon^{-1/2}$ with $\rho_b = \Omega_d/\omega_b$. Carrying out straightforward algebra, we then have

$$\delta W_{KT} \simeq \frac{8\pi^2 e^2 q R_0 B_0 \theta_T^2}{mc^2} \int d\mathcal{E} \int d\mu Q F_0 \Omega_d^2 \tau_b \times \sum_{l \neq 0} \sum_p \frac{p^2 J_p^2(\lambda_b)}{\lambda_b^2} \frac{\bar{\omega}_d - \omega}{(\bar{\omega}_d - \omega)^2 - p^2 \omega_b^2}. \quad (37)$$

Here, $\lambda_b = \theta_T k_\perp \rho_b$. Again, using Padé's approximant for the Bessel functions and replacing $\sum_{l \neq 0}$ by integration, Eq (37) becomes

$$\delta W_{KT} \simeq \frac{\pi^2 e^2 q R_0 B_0 \theta_T^2}{mc^2 s} \int d\mathcal{E} \int d\mu Q F_0 \Omega_d^2 \tau_b \frac{1}{\Delta_b (1 + \Delta_b^2)^{3/2}} \frac{\bar{\omega}_d - \omega}{(\bar{\omega}_d - \omega)^2 - \omega_b^2}. \quad (38)$$

Here, $\Delta_b^2 = \lambda_b^2/8$. Note also that, in Eq. (38), we have only kept $p = \pm 1$ terms; which, according to discussions similar to the case of circulating particles, are generally the dominant ones.

Within our orderings, $\omega_{*pi} < \omega_{dE}$, ω_{bE} , $\omega_{tE} < \omega_{*E}$, we get $\delta W_{Kr} > 0$ for $\omega \lesssim \omega_{*pi}$ from Eqs. (32), (36), and (38). The causality condition Eq. (14) thus tells us the MHD gap mode can only be excited when the system is ideal MHD unstable ($\delta W_f < 0$). Meanwhile, from Eq. (17) and Eq. (20), we recall that the existence of the unstable energetic-particle continuum mode requires $\partial \delta W_{Kr} / \partial \omega_r < 0$, and $\delta W_{Kr} < 0$ for ideal MHD stable plasmas ($\delta W_f > 0$). By investigating Eqs. (32), (36), and (38), we find that, given proper distribution functions, both conditions can indeed be satisfied for circulating as well as trapped energetic particles. Generally speaking, it implies that the energetic-particle continuum modes have characteristic frequencies $\omega_r \sim \omega_{tE}$ for circulating particles, and $\omega_r \sim \bar{\omega}_{dE}$ or ω_{bE} for trapped particles. Specific results pertinent to a slowing-down beam ion distribution will be presented in the next section.

The growth rate of energetic particle driven instability is proportional to δW_{Ki} ; which is generally positive definite for $\omega_{*E} > \omega_r$. From, Eqs. (32), (36) and (38), δW_{Ki} is proportion to the β_E and its radial gradient. In

previous analyses,^{11,12} $k_\theta \rho_E \lesssim |\Omega| \sim \epsilon$, one has the results with negligible orbit-size effects and $\delta W_{Ki} \propto k_\theta \rho_E$ (due to ω_{*E}). In the present analysis, $|\Omega| \lesssim k_\theta \rho_E \lesssim O(1)$, the finite-size orbit effects are clearly exhibited by the Δ and Δ_b factors in, respectively, Eqs. (32) and (38). Note that these factors plus QF_0 render δW_K weakly dependent on k_θ . Finally, we note that for $k_\theta \rho_E, k_\theta \rho_{bE} > 1$, δW_K s decay asymptotically as $(k_\theta \rho_E)^{-3}$ or $(k_\theta \rho_{bE})^{-3}$. This property can also be readily obtained from the large argument expansions of the Bessel functions.

V. Slowing-Down Beam Ion Distribution Function

In this Section, we shall derive the specific expressions for the growth rates of KBM using the following slowing-down beam ion distribution function:

$$F_{oE} = \frac{B_0 \beta_E}{2^5 \sqrt{2} \pi^2 m \mathcal{E}_m} \frac{\sqrt{1 - \alpha_0 B_0}}{1 - \alpha_0 B_0/2} \mathcal{E}^{-3/2} \delta(\alpha - \alpha_0); \quad (39)$$

where $\alpha = \mu/\mathcal{E}$ is the pitch angle, B_0 is the ambient magnetic flux at magnetic axis, and $\mathcal{E} \leq \mathcal{E}_m$ with $\mathcal{E}_m$ being the maximum beam energy.

A. Circulating Particles

Substituting Eq. (39) into Eq. (32), we find

$$\delta W_{KU_r} \simeq \frac{\pi}{2^{3/2} s} \alpha_E \frac{\omega_r}{\omega_{tm}} \left[1 - \frac{\omega_r}{2\omega_{tm}} \ln \left| \frac{\omega_r + \omega_{tm}}{\omega_r - \omega_{tm}} \right| \right], \quad (40)$$

and, for $|\omega_r| < \omega_{tm}$,

$$\delta W_{KU_i} \simeq \frac{\pi^2}{2^{5/2} s} \alpha_E \frac{\omega_r^2}{\omega_{tm}^2}; \quad (41)$$

where $\alpha_E \equiv q^2 R_0 \beta'_E$, and the subscript “ m ” indicates that the parameters are evaluated at $\mathcal{E} = \mathcal{E}_m$.

(i) For the MHD gap mode, Eqs. (22) and (23) become, denoting $\delta W_r = \delta W_f + \delta W_{Kr}$,

$$\omega_r \simeq \frac{\omega_{*pi} + [\omega_{*pi}^2 - 4\omega_A^2 (\delta W_r^2 - \delta W_{Ki}^2)]^{1/2}}{2}; \quad (42)$$

and

$$\gamma \simeq \alpha_E \frac{\pi^2}{2^{3/2}s} \frac{\omega_r^2 \omega_A^2}{\omega_{tm}^2} \frac{(-\delta W_r)}{\omega_{*pi}}. \quad (43)$$

γ , thus, increases with k_θ and is limited by $k_\theta \rho_m, k_\theta \rho_{dm} \lesssim 1$. Note the causality constraint $\delta W_r < 0$ becomes, approximately,

$$\delta W_f + \alpha_E \frac{\pi}{2^{3/2}s} \frac{\omega_r}{\omega_{tm}} \simeq \delta W_f < 0. \quad (44)$$

Equation (44), thus, reconfirms the observation that the MHD gap mode is unstable only if the plasma is ideal MHD unstable ($\delta W_f < 0$).

(ii) For the energetic-particle continuum mode, instabilities can occur even if ideal MHD is stable ($\delta W_f \geq 0$). Taking $\delta W_f \simeq 0$, we find from Eqs. (17) and (40) that at the instability threshold

$$\omega_r \simeq 0.834 \omega_{tm}. \quad (45)$$

The threshold condition, meanwhile, is given by Eqs. (19) and (41); i.e.,

$$\alpha_{EC} \simeq 0.57 s \omega_{tm}^2 / \omega_A \omega_r. \quad (46)$$

Near the threshold, the growth rate is

$$\frac{\gamma}{\omega_r} \simeq \frac{\pi}{2} \left(\frac{\alpha_E}{\alpha_{EC}} - 1 \right) \left[\ln \left| \frac{\omega_{tm} + \omega_r}{\omega_{tm} - \omega_r} \right| + \frac{\omega_{tm} (2\omega_r^2 - \omega_{tm}^2)}{\omega_r (\omega_{tm}^2 - \omega_r^2)} \right]^{-1}. \quad (47)$$

From Eqs. (45) to (47), we note α_{EC} and γ depends weakly on k_θ . In fact, referring to Eq. (32), it can be shown that γ is broadly peaked about $k_\theta \rho_{dm} \sim \epsilon_E^{1/3}$, where $\epsilon_E \equiv L_{pE}/R_o$ and L_{pE} is the scale length of the energetic-particle pressure.

B. Trapped Particles

We will use again the slowing-down beam ion distribution function, Eq. (39), for the trapped particle calculations and ignore the finite Larmor-radius effects of $J_0^2(\lambda_\rho)$; but keep the finite banana-width (finite λ_b) contributions. Corresponding expressions of δW_{KT} can then be derived using Eqs. (36) and (38) for, respectively, $|\omega| \sim \bar{\omega}_{dE} < \omega_{bE}$ and $|\omega| \sim \omega_{bE} > \bar{\omega}_{dE}$. Note the

case with $\bar{\omega}_{dE} \gtrsim \omega_{bE}$ (i.e., precessional drift-bounce resonance) is effectively suppressed by the finite- λ_b effects.

(i.) For $|\omega| \sim |\bar{\omega}_{dE}| < |\omega_{bE}|$, the mode is driven by the precessional drift resonance. We have

$$\delta W_{KT r} \simeq \frac{\pi}{\sqrt{2}} \alpha_E \left[1 - \frac{\omega_r}{\bar{\omega}_{dm}} \ln \left| \frac{\omega_r}{\bar{\omega}_{dm} - \omega_r} \right| \right], \quad (48)$$

and

$$\delta W_{KT i} \simeq \frac{\pi^2}{\sqrt{2}} \frac{\omega_r}{\bar{\omega}_{dm}} \alpha_E. \quad (49)$$

(i.a) The MHD gap mode can be excited when

$$\delta W_r \equiv \delta W_f + \delta W_{KT r} \simeq \delta W_{fC} < 0. \quad (50)$$

Here, δW_{fC} denotes ideal MHD δW with only the core component contribution. While ω_r is given by Eq. (42), the corresponding growth rate is

$$\frac{\gamma}{\omega_r} \simeq \alpha_E \sqrt{2} \pi^2 \frac{\omega_A^2}{\bar{\omega}_{dm}} \frac{(-\delta W_r)}{\omega_{*pi}}. \quad (51)$$

γ , thus, scales inversely with ω_{*pi} or k_θ . Equation (51), however, assumes $\omega_{*pi} \gtrsim 2|\delta W_r|/\omega_A$; i.e., the purely ideal ballooning mode is kinetically stabilized by ω_{*pi} .

(i.b) As to the energetic-particle continuum mode, the instability, again, can occur even if $\delta W_f \geq 0$. Taking $\delta W_f \simeq 0$, we have

$$\omega_r \simeq 0.78 \bar{\omega}_{dm}. \quad (52)$$

The corresponding threshold value of α_E is then given by

$$\alpha_{EC} \simeq 0.14 \bar{\omega}_{dm} / \omega_A. \quad (53)$$

Meanwhile, near the threshold, the growth rate is

$$\frac{\gamma}{\omega_r} \simeq \pi \left(\frac{\alpha_E}{\alpha_{EC}} - 1 \right) \left[\ln \left| \frac{\omega_r}{\bar{\omega}_{dm} - \omega_r} \right| + \frac{\bar{\omega}_{dm}}{\bar{\omega}_{dm} - \omega_r} \right]^{-1}. \quad (54)$$

Thus, both α_{EC} and γ scale with k_θ for $k_\theta \rho_{bm} \lesssim 1$.

(ii) For $|\omega| \sim |\omega_{bE}| > |\bar{\omega}_{dE}|$, the mode is driven by the bounce resonance. We have

$$\delta W_{KTr} \simeq \frac{\pi}{2} \frac{\theta_T}{s} \alpha_E \frac{\omega_r}{\omega_{bm}} \left[1 - \frac{\omega_r}{2\omega_{bm}} \ln \left| \frac{\omega_{bm} + \omega_r}{\omega_{bm} - \omega_r} \right| \right], \quad (55)$$

and, for $|\omega_r| < \omega_{bm}$,

$$\delta W_{KTr} \simeq \frac{\pi^2 \theta_T}{4s} \frac{\omega_r^2}{\omega_{bm}^2} \alpha_E. \quad (56)$$

We note that Eqs. (55) and (56) have the same functional dependencies as those of Eqs. (40) and (41) for the circulating particles, and, therefore, the stability properties can also be expected to be similar.

(ii.a) The MHD gap mode, again, can be excited when

$$\delta W_r \simeq \delta W_f + \frac{\pi}{2} \frac{\theta_T}{s} \alpha_E \frac{\omega_{*pi}}{\omega_{bm}} \simeq \delta W_f < 0; \quad (57)$$

i.e., only when $\delta W_f < 0$. The corresponding growth rate is then

$$\gamma \simeq \alpha_E \frac{\pi^2}{2} \frac{\theta_T}{s} \left(\frac{\omega_r \omega_A}{\omega_{bm}} \right)^2 \left(-\frac{\delta W_r}{\omega_{*pi}} \right). \quad (58)$$

(ii.b) Similar to the case of circulating particles, the energetic-particle continuum mode can be destabilized when α_E exceeds the threshold value even if the plasma is ideal MHD stable ($\delta W_f \geq 0$). In this case, we have for $\delta W_f \simeq 0$,

$$\omega_r \simeq 0.834 \omega_{bm}. \quad (59)$$

Meanwhile,

$$\alpha_{EC} \simeq 0.4 \frac{s}{\theta_T} \frac{\omega_{bm}^2}{\omega_r \omega_A}, \quad (60)$$

and, near the threshold,

$$\frac{\gamma}{\omega_r} \simeq \frac{\pi}{2} \left(\frac{\alpha_E}{\alpha_{EC}} - 1 \right) \left[\ln \left| \frac{\omega_{bm} + \omega_r}{\omega_{bm} - \omega_r} \right| + \frac{\omega_{bm}(2\omega_r^2 - \omega_{bm}^2)}{\omega_r(\omega_{bm}^2 - \omega_r^2)} \right]^{-1}. \quad (61)$$

Note that, similar to the case of circulating particles, the growth rate of the energetic-particle continuum mode is broadly peaked about $k_\theta \rho_{bm} \sim \epsilon_E^{1/3}$ and depends weakly on k_θ .

It is instructive to assess the relative importance between the precessional drift and bounce resonances. For the MHD gap mode, one can readily see from Eqs. (51) and (58) that the precessional drift resonance, Eq. (51), dominates. As to the energetic particles continuum mode, Eqs. (53) and (60) clearly indicate that the precessional drift resonance has a lower threshold for $k_\theta \rho_{bm} \lesssim 1$. For high α_E values, such that continuum modes corresponding to both resonances are excited with comparable growth rates, the precessional-drift mode tends to have higher k_θ , $k_\theta \rho_{bm} \sim 1$; while the bounce mode will have $k_\theta \rho_{bm} \sim \epsilon_E^{1/3}$.

VI. Summary and Discussions

In the the present work, we have carried out a matched asymptotic analysis and derived an analytic dispersion relation for the kinetic ballooning modes (KBM) resonantly excited by energetic ions/alpha particles in tokamaks. Our analysis contains effects due to finite-size orbits of both circulating and trapped energetic particles; such as Larmor radii, magnetic drift orbits and banana widths, which render energetic particle contributions effective only in the ideal region instead of the extended inertial region as in the case of negligible orbit sizes. The dispersion relation, furthermore, include various wave-particle resonances and is of the generic “fishbone” form.

We have analyzed this KBM dispersion relation and found that it exhibits two types of unstable modes. One is the MHD gap mode with $\omega_{*pi}/2 \lesssim \omega_r \lesssim \omega_{*pi}$ which exists only when the tokamak plasma is ideal MHD unstable, i.e., $\delta W_f < 0$. The other is the energetic-particle continuum mode with $\omega_r > \omega_{*pi}$; that is, ω_r lies inside the continuum. Typically, ω_r corresponds to the characteristic energetic particle frequencies such as $\bar{\omega}_{dE}$, ω_{bE} or ω_{tE} . The energetic-particle continuum mode can be excited even if $\delta W_f > 0$; but there is a threshold in α_E due to the resonant continuum damping.

To illustrate these properties more explicitly, we have further analyzed the dispersion relation using a slowing-down beam ion distribution function and derive specific expressions of ω_r , γ , and threshold α_E (if applicable) for both types of modes as well as circulating and trapped particles.

Several approximations have been made in this paper in order to simplify the analyses and, thereby, illuminate the physics. We expect straightforward extensions of the present work to more realistic tokamka equilibria and/or

to include additional effects such as core-component damping, global continuum damping, etc. The qualitative features of the instability properties are, however, expected to be similar to those obtained here.

Appendix A. Derivation of Equation (33)

Note that Eq. (31) can be written as

$$J_0(\lambda_\rho)2J_1(\lambda_d)/\lambda_d \simeq (1 + \Delta^2 k_\perp^2/k_\theta^2)^{-1} = (1 + \Delta^2)^{-1}(1 + \Lambda^2 \theta^2)^{-1}; \quad (\text{A1})$$

where $\Delta^2 \equiv k_\theta^2(\rho^2 + \rho_d^2/2)/4$ and $\Lambda^2 \equiv s^2 \Delta^2/(1 + \Delta^2)$. Fourier transforming Eq. (27) to z -space by $\int_{-\infty}^{\infty} e^{-iz\eta} d\eta$, we have

$$i(1 - \Lambda^2 \partial_z^2)(\omega_t z - \omega) \delta \hat{G}_d = \pi \frac{e}{m} \frac{Q F_0}{(1 + \Delta^2)} \frac{\Omega_d}{\omega} [\delta(z-1) - \delta(z+1)], \quad (\text{A2})$$

and the solution is

$$\delta \hat{G} \simeq \frac{i\pi}{2(\omega_t z - \omega)} \frac{e}{m} \frac{Q F_0 \Omega_d}{\omega(1 + \Delta^2)} \frac{1}{\Lambda} [e^{-|z+1|/\Lambda} - e^{-|z-1|/\Lambda}]. \quad (\text{A3})$$

Similarly, we have, from Eq. (26),

$$\delta \hat{S}_d^* \simeq -\frac{i\pi \Omega_d}{2(1 + \Delta^2)\Lambda} [e^{-|z+1|/\Lambda} - e^{-|z-1|/\Lambda}]. \quad (\text{A4})$$

Substituting Eqs. (A3) and Eqs. (A4) into Eq. (25) along with Parseval's theorem, we find

$$\delta W_{Ku} = \frac{\pi e^2 i^2 R_0^2}{4 m c^2} \left\langle \frac{Q F_0 \Omega_d^2}{(1 + \Delta^2)^2 \Lambda^2} \int_{-\infty}^{\infty} dz \frac{1}{\omega_t z - \omega} [e^{-|z-1|/\Lambda} - e^{-|z+1|/\Lambda}]^2 \right\rangle. \quad (\text{A5})$$

In deriving Eq. (A5), we have noted that $|\Lambda| \sim \epsilon^{1/2} < 1$, and thus, the integrand is peaked at $z = \pm 1$. Letting $x = 2|z \pm 1|/\Lambda$ in Eq. (A5), we then derive Eq. (33).

Acknowledgements

The authors would like to acknowledge useful discussion with Dr. G.Y. Fu and Prof. T.H. Stix. One of the authors (S.T. Tsai) is grateful for the hospitality of the Princeton Plasma Physics Laboratory during the time this work was performed. Work supported by U.S. Department of Energy Contract No. DE-AC02-76-CHO3073 and NSFC.

References

- ¹K. McGuire, *et al.* Phys. Rev. Lett. **50**, 89 (1983).
- ²W.W. Heidbrink *et al.* Phys. Rev. Lett. **57**, 835 (1986).
- ³K.-L. Wong *et al.* Phys. Rev. Lett. **66**, 1874 (1991).
- ⁴M.N. Rosenbluth, S.T. Tsai, J.W. Van Dam and M.G. Engquist, Phys. Rev. Lett. **51**, 1969 (1983).
- ⁵R.J. Hastie, Y.P. Chen, F.J. Ke, S.T. Tsai, and L. Chen, Chinese Phys. Lett. **4**, 561 (1987).
- ⁶L.J. Zheng and S.T. Tsai, Phys. Fluids B **4**, 1416 and 3329 (1992).
- ⁷L. Chen, R.B. White, and M.N. Rosenbluth, Phys. Rev. Lett. **52**, 1122 (1984).
- ⁸R.B. White, L. Chen, F. Romanelli, and R. Hay, Phys. Fluids **28**, 278 (1985).
- ⁹J. Weiland and L. Chen, Phys. Fluids **28**, 1399 (1985).
- ¹⁰H. Biglari and L. Chen, Phys. Fluids **29**, 1760 (1986).
- ¹¹B. Coppi and F. Porcelli, Phys. Rev. Lett. **57**, 2277 (1986).
- ¹²L. Chen, in **Theory of Fusion Plasmas**, edited by J. Vaclavik, F. Troyon and E. Sindoni (Assoc. EURATOM, Bologna, 1988), p. 327.
- ¹³H. Biglari and L. Chen, Phys. Rev. Lett. **67**, 3681 (1991).
- ¹⁴H.L. Berk, B.N. Breizman, and H. Ye, Phys. Lett. A, **162**, 475 (1992).
- ¹⁵G.Y. Fu and C.Z. Cheng, Princeton Plasma Physics Laboratory Report - 2852 (1992).
- ¹⁶J.W. Connor, R.J. Hastie and J.B. Taylor, Phys. Rev. Lett. **40**, 396 (1978).
- ¹⁷L. Chen and S.T. Tsai, Plasma Phys. **25**, 349 (1983).

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Prof. I. Kawakami, Hiroshima Univ., JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Research Info. Ctr., National Inst. for Fusion Science, JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. S. Tanaka, Kyoto Univ., JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	Library, Kyoto Univ., JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	S. Mori, Technical Advisor, JAERI, JAPAN
Dr. P.H. Sekanaka, Instituto Fisica, BRAZIL	Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL	J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
Documents Office, Atomic Energy of Canada Ltd., CANADA	D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Prof. T.W. Johnston, INRS-Energie, CANADA	Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Dr. C.R. James., Univ. of Alberta, CANADA	Dr. O. Petrus, AL I CUZA Univ., ROMANIA
Dr. P. Lukác, Komenského Univerzita, CZECHO-SLOVAKIA	Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
The Librarian, Culham Laboratory, ENGLAND	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS	Dr. L. Stanflo, Univ. of UMEA, SWEDEN
P. Mähönen, Univ. of Helsinki, FINLAND	Library, Royal Inst. of Technology, SWEDEN
Prof. M.N. Bussac, Ecole Polytechnique., FRANCE	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
Prof. E. Economou, Univ. of Crete, GREECE	Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
Ms. C. Pini, Univ. of Ioannina, GREECE	Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
Dr. T. Mui, Academy Bibliographic Ser., HONG KONG	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA	Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Librarian, International Center for Theo Physics, ITALY	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Prof. G. Rostangri, Istituto Gas Ionizzati Del Cnr, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN	Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

**DATE
FILMED**

7 / 30 / 93

END