MASTER COPY

MASTER

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

TECHNICAL DATA RECORD

NAA SR
TDR NO. 11127

APPROVALS

PAGE 1 OF 41

DATE 6-9-65

GO NO 7611

S. A NO 11313

SECURITY CLASSIFICATION

(CHECK ONE BOX ONLY)

UNCL. ☒ AEC ☐ DOD
CONF. ☐
SECRET ☐

(CHECK ONE BOX ONLY)

RESTRICTED DATA ☐
DEFENSE INFO. ☐

AUTHORIZED CLASSIFIER SIGNATURE

DATE

AUTHOR

DEPT. & GROUP NO.

H. E. Johnson

727 - 75

TITLE THERMO-PHYSICS TECHNICAL NOTE NO. 43:
SNAP FUEL PARTICLE DISPERSION

PROGRAM

SUBACCOUNT TITLE

Aerospace Safety

Development of Test Procedures

DISTRIBUTION

K. Adler 727
R. Balent 720
K. Buttrey 727
D. Cockeram 720
P. Cohn 791
R. Cranor 727
A. Crawford 735
R. Cummings 727

STATEMENT OF PROBLEM

The purpose of this study is to survey the available information on particle dispersion in the atmosphere to provide the foundation necessary to establish meaningful experiments and procedures for investigating the phenomena in facilities such as the hyperthermal wind tunnel.

ABSTRACT

An evaluation of particle dispersion from natural and artificial sources has been made relating this information to circulation processes in the mesosphere, stratosphere, and troposphere. The data on natural particle and nuclear debris dispersion have been interpreted in terms of their significance in regard to SNAP fuel reentry dispersion and extended by various fall out time and dispersion calculations. Particles of 0.01 to about 1 micron radius should be very widely dispersed and of no potential hazard. Particles above 1 micron size up to about 100 micron radius are in a transition zone between the nonhazardous and hazardous fractions. Particles over 100 microns radius, which some experimental and theoretical evidence indicate will be formed up to sizes over 1000 microns, will fall very rapidly to the ground creating potentially hazardous conditions unless further ablation or explosive disintegration processes reduce these particles to the nonhazardous size range. Particle disintegration mechanisms as understood from various experiments and theoretical considerations have also been reviewed. Further experimentation to more reliably determine particle size distribution as a function of size and to better understand disintegration mechanisms are recommended to more accurately evaluate potential hazards and to develop methods to eliminate the larger sized hazardous fraction of particles.

This report was prepared as an account of work sponsored by the United States Government. Neither the U. S. Government, nor any of its employees nor any of its contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 2 OF 41

Vertical Regions of the Atmosphere and Their Circulation

The regions of the atmosphere according to their vertical temperature distribution are the troposphere, stratosphere, mesosphere, and thermosphere. In addition to the regions based on temperature, there are two regions based on composition which are known as the homosphere and heterosphere. The lower region, the homosphere, is of relatively even composition with the exception of water vapor, ozone, and a few other trace molecules. Turbulence plays a dominant role in stirring the atmosphere below about 105km to maintain a more or less constant composition even though these regions vary in thermal stability.⁽¹⁾⁽²⁾ Above about 105 km a region in diffusive equilibrium prevails because of the absence of convection and turbulence which are the dominant processes below 105km. The upper region, the heterosphere, contains heavier elements in the lower layers whereas lighter elements predominate at higher levels due to gravitational separation. The relative stability of this region is due to the large increase in temperature with altitude. (See Figure 1)

The Troposphere and Tropopause

The troposphere is generally much more unstable than the stratosphere because the tropospheric lapse rate or decrease of temperature with elevation is about $6.5^{\circ}\text{K}/\text{km}$ whereas the stratosphere is either isothermal or shows an inversion or increase of temperature with altitude. The troposphere is well mixed because of the processes of convection, large scale horizontal and vertical motions, and precipitation. Mixing within a hemisphere, however, is much faster than mixing between hemispheres requiring months in the first case to several years in the latter.⁽¹⁾ Contaminant life is dependent on the nature and size of the atmospheric constituents. Stable gases such as CO_2 and CH_4 have lifetimes of about two to four years whereas submicron particles from nuclear explosions can be washed out by the action of clouds and precipitation in a matter of from two to four weeks. Table 1 which is a summary of mixing processes and times in the atmosphere indicates 1-2 year storage times for gases and 2-4 weeks for aerosols. Submicron particles will show variations from listed storage times because of changing weather processes and to slower exchange rates above about 6 km since rain layers are generally found below

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 3 OF 41

that level. Significant seasonal variations in deposition rates of nuclear bomb debris appear as a spring maximum and winter minimum.

The upper boundary of the troposphere, the tropopause, is a thermal transition zone between the troposphere, which is unstable in regard to the vertical circulation of air, and the stratosphere, where the temperature is constant or increasing so vertical circulation is relatively inhibited. Exchange across the tropopause occurs mainly at middle and high latitudes. The tropopause is not continuous around the globe but evidence exists that three tropopauses located in tropical, middle, and polar latitudes, as shown in Figure 2, are present with breaks between them.⁽³⁾ A possible mechanism to account for air mass exchange across the tropopause from the stratosphere to the troposphere and reverse was proposed by Staley who suggested that an annual variation in tropopause height results in a spring incorporation of stratosphere air into the troposphere and an autumn incorporation of troposphere air into the stratosphere.⁽⁴⁾ Thus in spring, an upward movement of the tropopause in middle and high latitudes results in the incorporation of stratosphere air into the troposphere, and likewise in fall, a downward movement incorporates troposphere air into the stratosphere. The most rapid exchange between air of the lower stratosphere and the troposphere occurs in the polar front region.⁽¹⁾

Stratosphere

The stratosphere which is a region of constant and increasing temperature is comparatively stable against convection with a lower rate of vertical mixing than in the troposphere. That the stratosphere can be dynamic rather than static, however, is evidenced by the presence of jet streams which circulate zonally from east to west in a closed loop around the pole at speeds the order of 100 km/hr. The meandering of this loop in meridional directions and the general transport processes in the stratosphere do not result, however, in the same degree of vertical and horizontal mixing of air as in the troposphere.⁽⁵⁾

The downward flow of lower stratospheric air under 20 km which results in intermixing with tropospheric air at mid and high latitudes in winter and spring can be interpreted in various ways. This down flow may be considered as due to a seasonal change in the rate of horizontal and vertical mixing or due to large scale meridional movement of stratospheric air poleward and downward, or as shifts in tropopause location. The upper

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 4 OF 41

stratosphere above 20 km is a more stable air mass than the lower stratosphere. Evidence in regard to circulation in the upper stratosphere is scant. The exchange processes may, however, differ from the lower stratosphere.⁽⁵⁾

According to Kellogg's summary of residence times given in Table 1, the lower stratosphere storage time is 1 to 2 years as compared to an upper stratosphere storage time of about 5-10 years. Movement of high stratosphere material to the low stratosphere is thought to occur during the winter months at high latitudes. This transfer occurs during the so-called impulsive warmings which may be the main way that aerosol material can move from high to low stratosphere levels. Fairly rapid horizontal spread can occur between about 20-30 km levels over a period of weeks to a few months to result in good mixing at high latitudes. Mixing in the equatorial zone at latitudes less than about 30 degrees appears to slow the horizontal exchange of air between hemispheres by acting as a buffer. This buffer zone is apparently the result of comparatively steady, zonal flow undisturbed by high level cyclonic storms which affect the mid latitude lower stratosphere regions. The low stratosphere buffer zone will cause a lower rate of mixing than at higher latitudes. Movement of material into the troposphere and hence to the ground comes about following the replenishment of the lower stratosphere in the winter and its incorporation in the troposphere during the spring peak.⁽⁵⁾

Mesosphere and Lower Thermosphere

Above 60 km a change in the vertical temperature lapse rate occurs so the temperature profile becomes similar to the troposphere. Vertical and horizontal mixing in this region is greater than in the stratosphere. This region is less thermally stable than the upper stratosphere. That the mesosphere is a region of strong wind shears as well as turbulence is evidenced by rocket and meteor trails. The region of strong mixing extends up into the lower thermosphere or up to about 105 km in spite of the more stable lapse rate occurring above the mesopause at about 80 km. Superimposed on the small scale turbulent motion is a large scale meridional circulation which results in additional mixing in this region. A possible reversal in wind direction every 3 or 4 km above 60 km which results in an effective wave length of from 6 to 8 km has also been reported.⁽¹⁾ This

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 5 OF 41

wave length appears to increase with height. Near the bottom of the mesosphere, at 55 km, the temperature approaches high values and stable lapse rate conditions. Under normal conditions there should be very little vertical motion at this level; however, large temperature fluctuations during periods of instability at high latitudes cause significant mixing during winter.⁽⁶⁾

The mesosphere mixing time from Kellogg's summary is much shorter than in the stratosphere. The estimated period of 1 to 10 weeks for mesosphere mixing as compared to 5 to 10 years for stratosphere mixing is indicative of much higher instability and wind shear in the mesosphere than in the stratosphere. The transport and distribution of materials in the upper atmosphere was discussed by Kalkstein who analyzed the results of the Rh¹⁰² tracer injected about 43 kilometers above Johnston Island at 16° north latitude on August 11, 1958 during the Orange shot of the Hardtack series of nuclear weapons tests.⁽⁶⁾ The bulk of the material rose above 100 km. His analysis of the bomb debris movement by tracer studies is as follows: (1) Stabilization of the debris occurred above 100 km followed by fairly horizontal diffusion in a matter of weeks. (2) The debris diffused down to about 80 km. (3) The material settled in a fairly uniform latitudinal direction down to about 55 km where it probably became concentrated. The debris is subject to significant vertical and horizontal mixing throughout this region with mixing times of weeks or months which are very short compared to stratosphere mixing times of from 5 to 10 years. (4) The material then moved down during impulsive warmings from the high stratosphere to low stratosphere in high latitudes and hence to the ground as previously discussed. The schematic drawing of Figure 3 as taken from Kalkstein portrays the sequence of events.

Particle Distribution in the Atmosphere

Natural Particle Distribution in the Upper Atmosphere

The upper atmosphere contains a natural distribution of particles from varied sources. According to Junge, et al.,⁽⁷⁾ the three major stratosphere aerosol components can be classified into particles of the following radii: (1) smaller than 0.1 μ , (2) between 0.1 μ and 1.0 μ (3) larger than 1.0 μ . Particles less than 0.1 μ radius are mainly of

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 6 OF 41

tropospheric origin with a lower limit of 0.01μ because coagulation times are shorter than residence times for particles of radius under 0.01μ . Half lives for particles in process of coagulation, given in Table 2, indicate that particles of less than 0.01μ located above 20 km will disappear very rapidly.

The second population range between 0.1μ and 1.0μ , which show a concentration distribution varying inversely with the square of the radius, reach a concentration maximum at about 20 km. This population consists of particles containing hygroscopic sulfur which are formed in the layer where they appear. The third population particles of radius more than 1.0μ are present in low concentrations; however, large concentration fluctuations occur in this particle range. These particles are of probable extra-terrestrial origin. Fall speed considerations make it likely that the majority of particles over one micron radius must have an outer space origin. Ablation temperatures will not be reached by incoming meteoritic particles of radius under one micron even at maximum entry velocities. Particles of radius over 100μ will ablate at all but the lowest velocities.⁽⁸⁾ The range of possible meteorite entry velocities is considered to extend from 11 to 72 km/sec.⁽⁹⁾

Nuclear Bomb Particle Distribution in the Upper Atmosphere

According to Junge, et al.⁽⁷⁾ particles injected into the stratosphere by atomic explosions vary in radius from about 0.02 to 5μ . The largest of these particles have been observed to fall from 30 km to the ground within two months whereas the smallest particles remain for years. Very small particles under several hundredths micron radius are expected to attach to natural occurring particles. Particles of more than a few hundredths micron radius will tend to remain as a more or less separate population intermixed with the natural particles. According to Friend,⁽¹⁰⁾ small particles of nuclear debris from high altitudes may coalesce with the larger sulfate particles at lower altitudes and then become distributed with the natural particles.

The distribution of aerosols in the lower stratosphere is controlled to a considerable degree, by exchange processes between the stratosphere and troposphere.⁽⁷⁾ These processes affect the distribution of particles

ranging from 0.01 to 10⁴ and larger. The stratosphere to troposphere exchange mechanisms, according to Junge, et al., include vertical mixing by turbulence, diffusion, and large scale exchange related to troposphere pressure systems, horizontal and meridional mixing across the tropopause, and gravitational settling in accordance with Stokes Law.

The behavior of submicron size stratosphere debris indicates that circulation and mixing rather than settlement control fallout processes.⁽¹¹⁾ Studies of world wide fallout or short lived isotopes indicate that stratosphere storage times decrease with increasing latitude but increase with increasing altitude.⁽¹²⁾ A stratosphere reserve results after the heavier particulates have been removed by **gravitational settling**.⁽⁵⁾ Land surface bursts distribute their products locally if the particle diameters are over 50 microns diameter. Smaller particles result in world wide fallout.⁽¹³⁾

Expected SNAP Reactor Fuel Particle Dispersion

Mechanisms of Particle Formation

Opik⁽¹⁴⁾ classifies meteoroid ablation into the following types: (1) vaporization, (2) fusion and spraying of the liquid, (3) sputtering, (4) fragmentation, (5) a combination of these processes. Vaporization is the final stage of the other types of ablation. Fusion is melting into one drop, whereas spraying is the formation of many drops. Sputtering or loss of single atoms from the surface is only of importance for micro-meteors of high velocity. Fragmentation of dustball meteorites forms many small probably solid fragments. Combinations of various processes may also occur such as the final vaporization of particles formed by fusion or spraying. The size of the meteoroid, its velocity, heat conductivity, melt viscosity, presence of an air cap, and rotation of the meteoroid will determine the type of ablation. Meteoroid ablation processes do not directly apply to SNAP reactor ablation upon reentry because the entry angle and speed of entry of SNAP systems are much less than for meteoroids.

Littman and Levy-Pascal⁽¹⁵⁾ suggest from their experimental work on the combustion behavior of zirconium-**uranium-hydride** that disintegration of fuel rods includes some or all of the following steps: (1) gradual heating which strips the cladding and causes loss of hydrogen,

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 8 OF 41

(2) oxidation from aerodynamic heating which forms an oxide layer and a molten core, (3) release of hydrogen or fission products or mechanical failure which causes ejection of molten material, (4) complete explosive disintegration of molten material. They conclude that conditions for complete dispersal of fuel exist but these conditions are poorly understood.

A graphic portrayal of possible ways that particles are formed is given in Figure 4. Observed combustion products taken from experiments reported by AVCO⁽¹⁵⁾ and illustrated in Figure 4 include solid particles and spheres with an oxide shell, hollow particles with an outer oxide shell, and small droplets released from the oxide by expansion of molten or boiling metal. Solid and hollow oxide spheres may be preferentially formed because initial reentry conditions will favor slow oxidation with later burning processes limited by diffusion of oxygen through the oxide shell. Some conclusions made from the AVCO study are (1) solid oxide particles up to 700 μ will be found along the reentry trail, and (2) oxide particles with diameters of several millimeters, which are the approximate size of stable molten droplets, will be formed.

Size of Particles Formed During Reentry

Beal, et al.⁽¹⁷⁾ in their analysis of ablation processes of metals mention that ablation by melting, if it occurs, is much more effective than ablation by vaporization because the heat required for melting is generally less than that required for vaporization. During melting ablation the run off liquid from the primary body will form secondary bodies or drops. As the molten material is subject to shearing and pressure forces from high velocity air, the drops formed can be further reduced in size unless they are smaller than the critical size which depends on liquid properties, velocity, and air density. Notable is the fact that the smallest droplet size occurs if the liquid melt is dispersed at low altitude and high velocity. Dispersion at 200 kft (61 km) results in stable droplet sizes from 1 to 10 millimeters. A stable droplet formed at a specific velocity is subject to deceleration so it is stable to breakup during the rest of its trajectory.

In a later report, Beal, et al.,⁽¹⁸⁾ state that ablation could best be performed at about 30,000 feet (9.1 km) resulting in stable droplet sizes

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 9 OF 41

of less than one micron. At higher altitudes where the stable drop size increases a mechanism such as drop explosion will be required to obtain fine disintegration. Observations of zirconium ablation tests showed that non-spherical zirconium oxide particles of sizes mostly under 10 μ were formed. Particle size distributions were close to those predicted from test conditions.

Other studies by Beal, et al.⁽¹⁹⁾ are concerned with stability of liquid zirconium drops as they move at high velocity through air. Experiments were run to determine whether oxidation of the surface of the droplet would interfere with the process of unstable, oversize drop reduction to stable droplet sizes. Tests performed by quickly enveloping a molten zirconium drop in air observed by a motion picture camera indicated that oxidation does not prevent breakup of unstable drop sizes to smaller droplets as predicted from the Weber number. The Weber number is $\frac{r \rho u^2}{\sigma}$ where r is the radius, ρ is the air density, u is the velocity, and σ is the surface tension. Breakup was expected when the Weber number exceeded a value of about 6.

Beal's theoretical data indicate that reentry of fuel rods at about 200 kft at a velocity of about 17 kft/sec will result in molten zirconium droplets of about 2 mm diameter (1000 μ radius) as can be seen in Figure 5. These droplets will react with atmospheric gases during reentry. Levy-Pascal's study⁽²⁰⁾ as well as others of the reaction of liquid Zr-10 w/o U droplets provide further understanding of what happens when these droplets or the fuel react with the atmosphere. Particles of 50 μ to 200 μ radius were found to undergo partial or complete oxidation dependent on experimental conditions. When oxygen was present particle disintegration to submicron size ranges occurred. Explosions were found to occur within 0.1 sec after expulsion in an oxygen containing atmosphere under a total pressure of 0.4 to 0.95 atm. This study indicates that explosive disintegration of large particles (for example, the 2 mm diameter molten droplets) may be of some significance in reducing particle size. Hollow oxide spheres, which will effectively fall as smaller particles, were observed in this experiment. Molten droplets of zirconium alloy should be able to ignite and burn upon entering the earth's atmosphere up to about 275 kft (84 km) according to the AVCO

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 10 OF 41

study.⁽¹⁶⁾ However, more precise experimental work at reentry pressures and oxygen compositions are needed to reliably predict oxidation of molten droplets, and fuel or fuel particles.

As illustrated in Figure 4, taken from the AVCO study, particles of various types can be produced by reentry ablation. Reaction of oxygen with molten particles can produce both solid and hollow oxide spheres and small droplets which result from the rupture of the oxide shell by molten or boiling metal. If the reentering stable droplets are 1000 μ in radius solid particles of the same size are possible. Conclusions from the AVCO experiments indicate that solid oxide spheres of 300 μ and hollow spheres up to 700 μ will also be formed. They indicated also that rupture of the oxide shell by boiling liquid would result in residual particles of the order of 5 μ . Boiling may be difficult to attain, however, because of radiant heat losses. Expansion of the molten metal inner core may also rupture the oxide shell although no evidence was found to establish this mechanism.

The study of the reaction of liquid Zr 10 w/o U droplets in flowing atmospheric gases at various partial pressures by Levy-Pascal, et al⁽²⁰⁾ showed that the presence of oxygen is required to cause final disintegration into submicron particles. The molten droplets which leave the surface of the melting fuel elements will react with oxygen and nitrogen in the atmosphere forming oxides and nitrides. Explosive reactions occur only at total pressures below 1 atm with the number of exploding particles increasing as the pressure decreases. Some of the resultant particles were in the submicron range. A definite trend toward size reduction was observed. Table 3 shows typical size range data for collected particles. Explosions were observed to follow particle swelling indicative of a pressure differential between the inside and outside of the particle. Hollow expanded spheres as well as solid spheres were collected after the runs. Initiation times for explosion were short, so that a particle leaving the fuel rod would explode in the rod's vicinity. These tests, however, were somewhat unrealistic in the sense that the material was heated very quickly whereas actual shallow angle reentry aerodynamic heating will heat the material slowly with relatively small temperature gradients through the fuel rod. Ablation during reentry could then occur by another mechanism.⁽²¹⁾

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 11 OF 41

Distribution of particles formed by the ablation of zirconium hydride 10 w/o uranium fuel samples with and without additives as a function of particle size reported by Foster⁽²²⁾ is shown in Figures 6 and 7. The additives were used in an attempt to change the properties of the fuel to enhance reentry burnup. Enhanced burnup is required either directly from the fuel element or from the droplets produced from the fuel rod in order to obtain what the author states is the safe dispersal requirement of 1μ particle size or less. The plot of particle size vs particle number shows a rapid decrease with increasing particle size whereas the plot of particle size vs total volume of particles shows a minimum total particle volume at about 25 microns size with maximums below 5 microns and above 1000μ . They conclude that different ablation processes caused the observed diameter groupings. The smaller particles were produced at the specimen surface by physicochemical processes but the larger products were caused by mechanical processes. The larger products were subject to further ablation. Ablation characteristics can be changed by the use of additives, however, alloying additions made directly to the melt appear to reduce the ablation tendency.

Explosive fracturing of molten particles with oxide shells may result in the volume distribution for various particle sizes shown in Figure 7. As previously mentioned Foster accounts for the double maximums by assuming that the larger particles probably result from mechanical processes whereas the smaller particles result from some physicochemical processes. Maximums for particles of diameter over 1000μ and particles under 5μ suggest, however, particle breakup from large into small pieces. For reentry at 200 kft (61 km) at velocities of about 17 kft/sec, expected particle sizes may range from 1000μ radius for oxide particles which retain their stable molten droplet sizes down to submicron sizes resulting from particle disintegration.

Fall-Out Time and Dispersion Calculations

Terminal Velocity - Fall-Out Time Estimates

A spherical particle falling in a quiescent atmosphere is subject to a gravitational force of $\frac{4}{3}\pi r^3 g (\rho_s - \rho_a)$ in the direction of motion. $\frac{4}{3}\pi r^3$ is the volume of a sphere of radius r . ρ_s and ρ_a are the

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 12 OF 41

densities of the particle and atmosphere, respectively, and g is the acceleration of gravity. The atmospheric drag force for Stokes flow is $-6\pi\mu rV$ for Reynolds numbers less than 4 where μ is the viscosity, r is the particle radius, and V is the velocity. Reynolds number is equal to $2\rho_a V r/\mu$. With no slipping of the flow about the sphere, a terminal velocity V_T is reached where both forces are equal as given by the equation:

$$V_T = \frac{2gr^2(\rho_s - \rho_a)}{9\mu} \quad (1)$$

Modifications of the atmospheric drag force to account for the occurrence of slip flow for particle radii ranging from 1 to 100 microns for various atmospheric mean free path, L , to r ratios are as follows:⁽²³⁾

$$L < r$$
$$F = -6\pi\mu rV (1 - L/r) \quad (2)$$

$$L \sim r$$
$$F = - \frac{6\pi\mu rV}{1 + L/r (A + B \exp \left\{ \frac{-cr}{L} \right\})} \quad (3)$$

$$L > r$$
$$F = - \frac{6\pi\mu r^2 V}{(A + B)L} \quad (4)$$

Values of V_T are obtained by equating equations (2), (3), and (4) with the gravitational force.

Figures 8 and 9 show respectively terminal velocities and time of fallout for altitudes up to 70 km for particle densities of 6 gm/cm³. These data are extrapolated to particle radii $< 1\mu$ and $> 100\mu$. Terminal velocities were calculated for the appropriate L/r ratio using data from the 1962 Standard Atmosphere and constant values where $A = 1.257$, $B = 0.400$, and $c = 1.10$. Terminal velocities were calculated for each 10 km altitude interval with the time for each particle to fall through each 10 km interval determined from average velocity values. Total time to fall from any given altitude to the ground were obtained by summing times over the 10 km intervals.

The Naval Radiological Defense Laboratory⁽²⁴⁾ developed a simplified formula for terminal velocities of irregular particles falling through a model atmosphere defined by the Air Research and Development Command.

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 13 OF 41

The particles are not assumed spherical as before, but are cylinders of diameter d and length $2d$. Values of d range from 20 to 10,000 microns. As before, approximations were made by summations of fallout times through 10 km intervals using terminal velocities averaged over each 10 km interval. Data from the U. S. Standard Atmosphere 1962 were used in place of the ARDC model atmosphere for purposes of this estimate. Figures 10 and 11 show terminal velocities and fall out times for altitudes up to 60 km calculated by this method. The NRDL formula is as follows:

$$V_h = \left[\frac{1.325}{\rho_0} b \eta \log_{10}^3 (bd + 1.163) \right] F_v \quad (5)$$

where:

V_h = particle terminal velocity (cm-sec⁻¹)

η = air coefficient of viscosity (g cm⁻¹ sec⁻¹)

$$b = \left[\frac{2 g \rho_0 (\rho - \rho_0)}{\eta^2} \right]^{1/3}$$

ρ = particle density (g cm⁻³)

ρ_0 = air density (g cm⁻³)

g = acceleration of gravity (cm sec⁻²)

d = particle size (cm) defined as a cylinder of diameter d and length $2d$.

$$F_v = 1 + \frac{L}{d} \left[2.514 + 0.300 \exp (-0.55 \frac{d}{L}) \right]$$

L = mean free path of air molecules (cm)

Particles of ellipsoidal shape will, for the most part, fall at velocities less than corresponding spheres of the same volume. The ratios of fall velocities for particles of more or less ellipsoidal shape to the fall velocities of spheres of the same volume are given in Figure 12 taken from Facy⁽²⁵⁾ as adapted from computations by Oberbeck and Lamb. This diagram shows that fall velocities for particles of various ellipsoidal shapes **may equal** but do not significantly exceed the fall velocities of spherical particles of equal volume. Likewise, according to Facy, the fall velocities of irregularly shaped particles are smaller than the fall velocities of spheres of the same volume.

A cylinder of diameter, d , and length $2d$, of the type illustrated in Figures 10 and 11, if assumed to fall like an ellipsoid of semimajor axis twice the semiminor axes will fall at about 0.9 times the fall velocity

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 14 OF 41

of a sphere of equal volume. Quasi-spherical particles of rough, irregular surfaces will fall less rapidly than smooth spheres of equivalent mass and density.⁽²⁶⁾

Particle Dispersion Estimates

Sowle's Hazard Estimate for Reentering Reactors

Sowle's results are illustrated in the plot of Figure 13 taken from his report.⁽²⁷⁾ The fraction of original constituent injected into the upper atmosphere at 30 km (92,000 ft) which is deposited per unit area, f , intersects the NBS acceptable level at about the 60 μ radius level. The maximum hazard which the author describes for particles over 30 to 60 μ is for the case where the particle trajectory is parallel to the mean wind. He also reports that the hazard probability for particles of less than 100 μ radius is less than 10%.

Sutton Turbulent Diffusion Analysis (Beryllium reentry study program)

Field⁽²⁸⁾ applied Sutton's theories which attempt to combine the elements of diffusion theory, turbulence, and eddy diffusion to the problems of beryllium reentry into the atmosphere. Particles of beryllium under 5 μ diameter were considered to behave similarly to particles of molecular diffusion size. An average wind velocity of 5 meters/second is assumed. Derived formulas are as follows:

$$d_{x_{\max}} = \left(\frac{2h^2}{3c^2} \right)^{\frac{1}{2-n}}$$

$$x_{\max} = \frac{Q}{(2/3 e \pi)^{3/2} h^3}$$

where:

$d_{x_{\max}}$ = distance of maximum concentration downwind from source (meters)

h = source height (meters)

c = diffusion coefficient

n = stability parameter where $0 \leq n \leq 1$

$x_{\max}$ = maximum concentration (gm/m^3)

Q = total quantity of material

Assuming that these equations can be applied to SNAP fuel rods for submicron particle sizes, values for distance of maximum concentration, $d_{x_{\max}}$ and maximum concentration, $x_{\max}$, are obtained as follows:

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 15 OF 41

$$d_{x_{\max}} = \left[\frac{2x(6 \times 10^4)^2}{3x(0.20)^2} \right] \frac{1}{(2-0.25)} = 1390 \text{ Kms}$$

$$\begin{aligned} x_{\max} &= \frac{5.18 \times 10^4}{(2/3 \times 2.7183 \times 3.1416)^{3/2} (6 \times 10^4)^3} = 1.8 \times 10^{-11} \text{ gm/m}^3 \\ &= 716 \text{ particles/1000 m}^3 \text{ of } 1 \mu \text{ radius (spherical)} \\ &= 5.5 \times 10^{-7} \text{ gm/km}^3 \text{ per gram of source} \end{aligned}$$

where:

$$h = 60,000 \text{ meters}$$

$$n = 0.25 \text{ (stable atmosphere)}$$

$$c = 0.20 \text{ (stable atmosphere)}$$

$$q = 5.18 \times 10^4 \text{ gm}$$

A Quiescent Atmosphere Model of Gravitational Settling and Dispersion

This model was developed by the Geophysics Corporation of America to determine given particle concentration surfaces resulting at 30 km altitude from a fall from 70 km of particles of various sizes ranging from 2 to 20 microns in diameter. (29) Surfaces of equal concentration are represented as ellipsoids. Maximum particle concentrations occur at the cloud center of gravity located at the center of the ellipsoid. Cloud dispersion effects are concentrated in the value assigned to the eddy diffusion parameter, K_e . The eddy diffusion coefficients used in this model are as follows: longitudinal coefficient, K_x of $10^{10} \text{ cm}^2/\text{sec}$, a latitudinal coefficient, K_y , of $10^8 \text{ cm}^2/\text{sec}$, and a vertical coefficient, K_z of $10^4 \text{ cm}^2/\text{sec}$. The relative dimensions of the ellipsoid concentration surface, x , y , and z as derived from equations (1) and (2) which assume Gaussian particle distribution are given in equations (3), (4), and (5).

$$n(x, y, z, t) = \frac{N}{(2\pi)^{3/2} \sigma_x \sigma_y \sigma_z} \exp \left[-\frac{x^2}{2\sigma_x^2} - \frac{y^2}{2\sigma_y^2} - \frac{z^2}{2\sigma_z^2} \right] \quad (1)$$

$$\sigma^2 = 2 K_e t \quad (2)$$

$$x = \sqrt{2\sigma_x^2 \left\{ \ln \frac{N}{n} - \ln(2\pi)^{3/2} - \ln(\sigma_x \sigma_y \sigma_z) \right\}} \quad (3)$$

$$y = \sqrt{2\sigma_y^2 \left\{ \ln \frac{N}{n} - \ln(2\pi)^{3/2} - \ln(\sigma_x \sigma_y \sigma_z) \right\}} \quad (4)$$

$$z = \sqrt{2\sigma_z^2 \left\{ \ln \frac{N}{n} - \ln(2\pi)^{3/2} - \ln(\sigma_x \sigma_y \sigma_z) \right\}} \quad (5)$$

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 16 OF 41

where:

n = concentration of particles at any point in Cartesian coordinates, x, y, z, t (particle/cm³)

$\sigma_x, \sigma_y, \sigma_z$ = standard deviations of the particle distribution along each of the three coordinates (cm)

N = total number of particles initially released

σ^2 = relative dispersion (cm²)

K_e = eddy diffusion parameter (cm²/sec)

Maximum particle concentrations and constant concentration ellipsoid surface dimensions for SNAP fuel particles calculated by these equations are given in Table 4. The total number of particles, N , of 6.0 gm/cm³ density initially released is determined by dividing the total mass of the fuel rods, 5.18×10^4 gms, by the mass of each particle using various spherical and cylindrical particle sizes. Values of $\sigma_x, \sigma_y, \sigma_z$ were determined from their respective eddy coefficients K_x, K_y, K_z and the time of fallout for the particle under consideration using equation (2). Values of x, y , and z were solved for ellipsoid concentration surfaces of 1 particle per 10^3 meter³, 10^4 meter³, and 10^5 meter³. Maximum concentrations of n were solved by determining the value of n where $x=y=z=0$. Calculations are done for all distances from 60 to 30 km and from 60 to 0 km assuming stratosphere conditions extend down to ground level. Hypothetical concentration surfaces at ground level provide an upper level of values since actual dispersion in the troposphere will generally exceed similar effects in the upper atmosphere.

Accumulated Deposit Densities and Area Coverage by Hage

Table 5 taken from Hage⁽²⁶⁾ shows accumulated deposition areas and densities at the earth's surface from fallout from 30km for smooth spheres of 5.0 gm/cm³ density and diameters of 11 μ and 200 μ . Surface deposition is shown for summer and winter in equatorial and polar latitudes for several variable percentages of the original mass. Deposition densities vary from 1.7×10^{-2} to 7.3×10^{-9} gm/km² per gram of source material for 20, 50, and 80 per cent of the source mass. Their calculations included the following mechanisms: (1) lateral eddy diffusion, (2) variations in fall speed due to altitude, (3) variations in fall speed from variable particle sizes and (4) variations from wind transport

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NAA SR TDR 11127

NO. _____

DATE 6-9-65

PAGE 17 OF 41

profile vectors in altitude, latitude, and season.

Circulation and Distribution of SNAP Fuel Particles

Submicron Particles

Submicron particles injected into the mesosphere between about 60 and 65 km should behave in somewhat the same fashion as the nuclear bomb tracers outlined in Kalkstein's study. Particles of 0.01 micron radius have a fall velocity of about 0.3 cm/sec at 60 km according to Figure 8 which means that about three weeks will be required for particles of that size to reach the normally stable high temperature layer at about 55 km. Since mixing time in the mesosphere from Table 1 is 1 to 10 weeks, the particles of this size or smaller may be reasonably well distributed by the time the stable layer is reached. In reality, however, the stable layer at times extends to 60 km so the actual mixing time is probably longer. Due to 5 to 10 years for both the mixing time and residence time in the stratosphere, this material should be widely distributed by the time the lower stratosphere is reached. The normal circulation pattern can be expected to change during the winter instability period which will cause a fraction of the material to move from the high to low stratosphere. We can thus expect that a well mixed reservoir in the upper stratosphere should gradually be transported to the lower stratosphere. Kalkstein⁽⁶⁾ assumes that every year 1/10 of the upper stratosphere is exchanged with the lower stratosphere. Long residence in the stratosphere is further vindicated for particles of 0.01 microns radius because a period of about 10 years is required for the particles to fall from 60 to 20 km.

Particles of radius of about one micron should fall from the upper to lower stratosphere (60-20 km) in about a month's time assuming terminal fall velocities of Figure 8 are valid for this case. These particles will drop at increasingly slower terminal velocities as they reach lower altitudes. Prior to dropping to an altitude of 30 km, the fall speed of these particles will be less than 1 cm/sec. Since average horizontal wind velocities exceed 1 cm/sec terminal velocities by a factor of roughly about 1000 in the lower stratosphere,⁽²⁷⁾ particles of 1 micron radius should not only be widely dispersed but their motions should be controlled by lower stratosphere circulation processes as for very small submicron

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 18 OF 41

particles. Particles of 0.1 micron radius which theoretically take about a year to fall from 60 km to 20 km will reach terminal velocities of 1 cm/sec at about 50 km. Particles of radius less than 1 micron should become dispersed with the naturally occurring lower stratosphere aerosols which have been found in sizes ranging between 0.01 micron and 1 micron. Turbulence and convection in the troposphere will also disperse the particles of 1 micron radius or less as indicated by the fact that one micron particles will theoretically require about 4 months (Figure 9) to fall to the ground from 10 km whereas estimated mixing times within this region are a few weeks. (Table 1). Actual storage times in the troposphere are from about two to four weeks because variable weather processes such as precipitation may remove this material more rapidly than indicated by settling velocity data.

Removal of the submicron particles reservoir from the low stratosphere to troposphere should occur primarily at middle and high latitudes during the spring. Whether this material can move from one hemisphere to another apparently depends largely on cross-hemispheric mixing before the material passes into the lower stratosphere. There is a barrier in the lower stratosphere which acts to limit exchange between hemispheres. Since the circulation in the stratosphere is primarily zonal rather than meridional and the upper stratosphere is normally a comparatively stable layer, inter-hemispheric transport of the small particles (≈ 0.01 microns) may be much greater than for larger particles of the order of 1 micron due to the large difference in settling velocity.

Particles Over 1 Micron Radius

The theoretical study of the release of zirconium-uranium-hydride fuel at 30 km altitude by Sowle⁽²⁷⁾ shows that no radiological hazard exists if the mean particle radius is less than 30 to 60 microns with a less than 10% hazard probability if the average radius is less than 100 microns. The mechanisms which they considered were significant in spreading a line source of particles over a large area before reaching the ground were the following: (1) turbulent eddy diffusion, (2) the length of burn path, and (3) the variable terminal velocities of particles of different sizes. Experience concerning debris injected by nuclear explosion at high altitudes indicates that atmospheric circulation patterns will widely

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 19 OF 41

disperse SNAP fuel burnup particles in the submicron size range. Sowle's study indicates that SNAP fuel particle dispersion is sufficient to prevent radiological hazard not only from submicron particles but for sizes up to 30 to 60 microns radius. (Figure 13)

Experience from land surface nuclear bombs indicate that particles of less than 25 micron radius will be dispersed on a worldwide basis by tropospheric processes.⁽¹³⁾ Consequently, particles dispersed at 60-65 km by burnup of SNAP fuel rods should not reach hazardous concentrations providing the particles that reach the troposphere are less than 25-60 microns radius. Efforts should be aimed at ways to ensure the breakup of fuel material so that ultimate ablation of the fuel rods and large fuel particles will result in products that reach the troposphere of radius preferably less than about 25 microns.

Although particle distribution models which present a highly probable situation assuming mean conditions are useful in predicting SNAP fuel particle dispersal hazards, these models necessarily neglect the tremendous variability of atmospheric processes which under favorable circumstances might disperse even relatively large particles much more widely than expected from the model. Reentry altitude, latitude, and season of entry may thus be of great importance in determining whether or not a hazard will exist for particles particularly in the marginal radius size range from about 25 to 100 μ . To illustrate the extent of wind variability, Figure 14, which was taken from a recent report by Kantor and Cole,⁽³⁰⁾ gives mean January and July zonal winds from the equator to 75°N to a height of 120 km. (See also Table 6). Although January and July do not represent extremes of all heights and latitudes, Figure 14 data are typical of wind regimes over the northern hemisphere. Day to day variability of winds in January may at some latitudes (30° and 60°) and altitudes (30-40 km) may even result in stronger easterlies than in July. Transfer of air from the high stratosphere to the low stratosphere during the winter months followed by the incorporation of this air in the spring months at mid to high latitudes, may act to enhance fallout of particles entering the upper atmosphere during these months. Entry at other seasons of the year may have the opposite effect of delaying fallout time and of increasing particle dispersion.

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TR 11127
DATE 6-9-65
PAGE 20 OF 41

The principal hazard is for particles of radius over about 100μ . Particles of this size can be expected to settle out of the atmosphere from 60 km in a matter of hours as indicated by the plot of Figure 9. That a significant radiological hazard exists for particles of radius above 100μ is implied by Sowle's calculations. Sowle, however, assumes a number distribution within 10% of a given particle radius. Foster's work, however, indicates that actual distributions show a double peak when total volume is plotted against particle size even though the number distribution decreases by a factor of about 1000 in going from particle sizes of 25μ to 1000μ . This particle size total volume data would thus suggest that there are two distributions with a nonhazardous fraction of radius under 5μ resulting from particle disintegration, and a potentially hazardous fraction resulting from larger particles which do not further ablate or explosively disintegrate. If a significant fraction of the original fuel rod volume is retained by the larger fraction, of some 10 to 100 times the radius of the smaller particles, then a possible fall-out hazard could occur due to the rapid settling velocity and slower dispersion rate of these particles.

Particle concentrations calculated by the Geophysics Corporation of America method with results given in Table 4 indicate that stratosphere processes will disperse particle concentrations of spherical particles up to 10μ radius to comparatively low values between 60 and 30 km. Extrapolation of these data down to zero elevation is also done by assuming diffusion coefficients in the stratosphere apply also to the troposphere in order to provide an upper limit to concentrations at ground level. Mixing times in the troposphere are considerably less than in the stratosphere. (Table 1). Sowle reports that the concentration of particles less than 30-60 microns from reentering U-Zr-H fuel is insufficient to cause a radiological hazard. The radiological hazard is less than 10% if the average particle radius is less than 100μ . He assumes that the reentry burnup altitude is 30 km. Application of Sutton's calculations for submicron particles as used by Fields, and Hage's data given in Table 5 outlined previously show also comparatively low fallout densities for particles near or slightly above the submicron size range. Of the two probable fractions formed by the ablation of SNAP fuel rods on

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 21 OF 41

reentry, the larger fraction is potentially much more hazardous. However since fewer particles of large size can be formed from a given amount of source material, even the larger particles will not necessarily produce a hazardous concentration at ground level. In addition, the larger fraction will, to some extent, be partially controlled or influenced by circulation processes. Consequently, the problem to be answered before realistic calculations of fallout hazard can be made is the problem of size and volume distribution of the particles formed by reentry of SNAP fuel rods in the upper atmosphere.

Problem Areas and Methods of Solution

The principal problem is to determine the final dispersion concentration at ground level of the potentially hazardous fraction of reentering SNAP fuel particles. The hazardous fraction is predominantly of particles greater than 100μ radius with a peak in volume concentration according to Foster's data at $>500\mu$ radius. Particles which are not spherical but of ellipsoidal, cylindrical, and irregular shapes will fall predominately at slower velocities and will become even more widely dispersed than spherical particles. The first problem encountered is to establish the number and volume distribution by size of particles formed by the ablation of SNAP fuel rods at expected reentry elevations. Ablation tests simulating actual reentry pressures and atmospheric compositions at the chosen altitude should be done to better establish the particle size distribution initially formed. Secondary ablation of the primary larger particles is another process which should be more thoroughly investigated because this process may act to significantly reduce the relatively hazardous large particle fraction to a less hazardous small particle fraction. In addition to secondary ablation of primary particles, explosive disintegration may be even a more significant factor in reducing large particle sizes to nonhazardous levels. Experimental work to determine particle size distribution at reentry altitudes as well as secondary ablation and explosive disintegration of large particles into small particles occurring down to tropopause level are needed to establish data for improved hazard calculations.

Particle size distribution must be obtained in terms of particle volume as well as particle numbers. The shape and volume number

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 22 OF 41

distribution of particles, resultant from the initial stages of rod ablation during reentry, should first be more reliably determined followed by analysis of the final distribution of particles entering the tropopause, resulting from particle explosion and further ablation of larger particles. The expected final predominant distribution will consist of large particles possibly of solid oxide composition of the order of 500 μ radius and many smaller particles in the submicron range. This distribution needs to be determined for each probable reentry altitude, trajectory since the extent that potentially hazardous large particles are fragmented into relatively safe submicron particles depends partially on the total pressure and atmospheric composition at fuel burnup altitude. Experiments by Levy-Pascal⁽²⁰⁾ show that total pressure and oxygen content are particularly important in determining the amount of particle disintegration that occurs. As previously mentioned, the importance of secondary reentry processes in influencing final size distribution needs to be properly evaluated since further ablation and explosive disintegration of primary ablation products may be of critical importance in reducing particle sizes to the submicron range.

The secondary aim of an experimental program should be to try to discover ways to improve fuel particle disintegration processes. Variables which affect particle size and shape formation are the following: stable droplet size decrease with decreasing altitude, explosive disintegration of particles with an outer oxide shell by melting expansion or boiling of the interior liquid metal will fracture large into small particles, total and partial pressures of atmosphere components will affect the extent of this disintegration, and further ablation of the relatively large fragments as they fall will further reduce particle sizes.

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 23 OF 41

REFERENCES

- (1) W. W. Kellogg, "Pollution of the Upper Atmosphere by Rockets," Space Science Reviews, 3, 275-316, 1964
- (2) J. C. G. Walker, "The Upper Atmosphere," Space/Aeronautics, 42, 4, October 1964
- (3) P. A. Sheppard, "Atmospheric Tracers and the Study of the General Circulation of the Atmosphere," Reports on Progress in Physics, Vol 26, 1963
- (4) D. O. Staley, "Stratospheric Mass Variation and Fission Product Radioactivity in Surface Air," Journal of the Atmospheric Sciences, Vol. 20, 1963
- (5) R. Bjornerstedt and Kay Edvarson, "Physics, Chemistry, and Meteorology of Fallout," Annual Review of Nuclear Science, Vol. 13 1963
- (6) M. I. Kalkstein, "Rhodium-102 High-Altitude Tracer Experiment," Science, Vol. 137, No. 3531, August 31, 1962
- (7) C. E. Junge, C. W. Chagnon, and J. E. Manson, "Stratospheric Aerosols," Journal of Meteorology, Vol. 18, February 1961
- (8) N. P. Carleton, "The Relation of Recent Atmospheric Dust Measurements of Volz and Goody to the Problem of Meteoritic Influx," Journal of the Atmospheric Sciences, Vol. 19, September 1962
- (9) C. J. Maiden, "Meteoroid Impact," Space Exploration, LeGalley and McKee Editors, 1964.
- (10) J. P. Friend, "The Behavior of Nuclear Debris from the Upper Atmosphere Rocket Detonations as Determined in the High-Altitude Sampling Program," SCR-420, Proceedings of the Upper Atmosphere Sampling Symposium, Part 1, Sandia Lab, Albuquerque, N.M., April 11-13, 1961
- (11) E. A. Martell, "Evidence for High Stratosphere Holdup of Nuclear Bomb Debris," SCR-420, Proceedings of the Upper Atmosphere Sampling Symposium, Part 1, Sandia Lab, Albuquerque, N.M., April 11-13, 1961
- (12) E. A. Martell and P. J. Drevinsky, "Atmospheric Transport of Artificial Radioactivity," Science, Vol. 132, No. 3439, November 25, 1960
- (13) A. D. Anderson, "The NRDL Dynamic Model for Fallout from Land Surface Nuclear Bursts," USNRDL-TR-410, 1960
- (14) E. J. Opik, "Physics of Meteor Flight in the Atmosphere," Interscience Publishers, Inc., New York, 1958

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 24 OF 41

References (Continued)

- (15) F. E. Littman and A. E. Levy-Pascal, "Combustion and Disintegration of Zirconium Hydride-Uranium Fuel Rods, "SC-DC-3553, Proceedings Aerospace Nuclear Safety, ANS-National Topical Meeting, Albuquerque N.M., October 1-4, 1963
- (16) Research and Advanced Development Division, AVCO Corp., "Investigation of Reentry Destruction of Nuclear Auxiliary Power Plant," AFSWC-TR-61-69, October 1961
- (17) J. L. Beal, W. Smith, F. Haas, R. Pleuthner, "Preliminary Analysis Dispersal of Reactor Core Materials by Reentry Processes," TID-17172 July 1962
- (18) J. L. Beal, W. Brown, R. Pleuthner, "Dispersal of Reactor Core Materials by Reentry Processes," Quarterly Report, November 1963
- (19) J. Beal, W. Brown, F. Vassallo, "Dispersal of Reactor Core Materials by Reentry Processes," SC-DC-64-1494X, Quarterly Report, July 1964
- (20) A. Levy-Pascal, "Particle Disintegration Study for Reentering Nuclear Auxiliary Power Systems," Final Report, AFSWC-TDR-63-51 June 1963
- (21) J. Stockham and R. Snow, "Ablation Product Coalescence," IITRI C6007-12, February 15, 1963 to February 14, 1964, March 13, 1964
- (22) E. L. Foster, Jr., "Reentry Burnup Enhancement of Hydrided Zirconium Uranium Alloy Fuel," SC-DC-3553, Proceedings Aerospace Nuclear Safety, ANS-National Topical Meeting, October 1-4, 1963
- (23) Geophysics Corporation of America, "Circulation Processes and Mixing Mechanisms in the Upper Atmosphere Above 30 km," SC-DC-64-326, Quarterly Progress Report, September 15, 1963 to December 15, 1963
- (24) D. Clark, Jr and W. Cobbin, "Some Relationships Among Particle Size, Mass Level, and Radiation Intensity of Fallout from a Land Surface Nuclear Detonation," USNRDL-TR-639, March 21, 1963
- (25) L. Facy, "Radioactive Precipitations and Fallout," Nuclear Radiation in Geophysics, H. Israel and A. Krebs, Editors, 1962
- (26) K. Hage, "Particle Fallout and Dispersion Below 30 km in the Atmosphere," SC-DC-64-1463, Final Report, Sandia Corp., Albuquerque, N.M., August 1964
- (27) D. Sowle, "The Hazard Associated with the Dispersal of Small Particles from Reentering Reactors as a Function of Particle Size," AFSWC-TDR-62-81, August 1962
- (28) A. Field, Jr., "Study of Beryllium Burnup and Dispersion for Reentering Nuclear Auxiliary Power Systems," AFSWC-TDR-62-119, Dec. 1962

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 25 OF 41

References (Continued)

- (29) Geophysics Corporation of America, "Circulation Processes and Mixing Mechanisms in the Upper Atmosphere Above 30 km," SC-DC-64-764, Quarterly Progress Report No. 2, December 15, 1963 through March 15, 1964.
- (30) A. J. Kantor, A. E. Cole, "Zonal and Meridional Winds to 120 Kilometers," Journal of Geophysical Research, Vol. 69, No. 24, December 15, 1964

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127

DATE 6-9-65

PAGE 26 OF 41

TABLE 1

Summary of Mixing Processes, Rates, (1)
and Storage Times in the Upper Atmosphere

Thermo- sphere	Molecular diffusion	~ 1 week	5-10 years
Meso- sphere	Turbulent diffusion	1-10 weeks	5-10 years
Strato- sphere	Turbulent diffusion and large scale cir- culations	5-10 years	5-10 years in upper stratosphere 1-2 years in lower stratosphere
Tropo- sphere	Turbulence and convection	Few weeks	1-2 years for gases 2-4 weeks for aerosols
Probable mixing mechanism		Estimated mixing time within region	Estimated storage time (Time for exchange with tropo- sphere or loss to ground and oceans)

TABLE 2

Half-Lives in Days for Particle Coagulation (7)

Altitude (km)	Radii in Microns			
	0.0025	0.005	0.01	0.02
27.5	1.6	6.4	24	90
20.0	5.1	20	80	280
12.5	17	61	210	740

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. 'NAA SR TDR 11127DATE 6-9-65PAGE 27 OF 41

TABLE 3

Typical Size Range of Collected Particles⁽²⁰⁾

Atmosphere	Argon			Argon-Oxygen Mixture**			Argon-Nitrogen Mixture		
Original Particle Size (microns)	45-53	88-105	177-210	44-53	88-105	177-210	44-53	88-105	177-210
Final Spherical Combustion Product Size (Microns)	12-100	25-105	25-200	12-75*	12-75	50-200	25-75	50-150	50-200

* In some cases, no spheres were found. All particles disintegrated into fine dust after passing through plasma flame.

** Large hollow expanded particles not included.

TABLE 4

Particle Dispersion Concentrations*

Particle Shape	Size (μ)**	Fall Distance (km)	Maximum Concentration particles/1000m ³	Ellipsoid Dimensions for concentration Surface = 1 particle/ 1000m ³		
				x(km)	y(km)	z(km)
sphere	1	60-30	654	4520	452	4.5
sphere	1	60-0	5.2	11460	1146	11.5
sphere	10	60-30	30.5	914	91.4	.9
sphere	10	60-0	2.1	1038	103.8	1.0
cylinder	20	60-30	44.5	589	58.9	.6
cylinder	20	60-0	1.8	681	68.1	.7

* For ellipsoid of axes x, y, z. Maximum concentration assumed at $x = y = z = 0$.

**For sphere = radius; for cylinder, length = 2 x Diameter.

ATOMICS INTERNATIONAL
A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127
DATE 6-9-65
PAGE 28 OF 41

Table 5

Computed Area Coverages and Average Accumulated Deposit Densities at the Earth's Surface Due to the Fallout and Lateral Eddy Diffusion of Smooth Spheres (Density 5.0 gm cm^{-3}) From A Point Source at 30 km. (26)

(See Supplemental Table 6 for assumed wind profiles used in calculations)

Wind Profile Number (Table 6)	Latitude Belt and Season	Mass-Median Diameter (Microns)	Range of Fall Times in Days (80% of Total Mass)	Percentage of Total Mass					
				20		50		80	
				Area Covered (km^2)	Ave. Density ($\text{gm gm}^{-1} \text{km}^{-2}$)	Area Covered (km^2)	Ave. Density ($\text{gm gm}^{-1} \text{km}^{-2}$)	Area Covered (km^2)	Ave. Density ($\text{gm gm}^{-1} \text{km}^{-2}$)
(1)	Equatorial summer	200	0.046 to 0.22	1.9×10^2	1.1×10^{-3}	8.8×10^2	5.7×10^{-4}	4.3×10^2	1.9×10^{-4}
(2)	Equatorial winter			1.2×10^1	1.7×10^{-2}	5.5×10^1	9.1×10^{-3}	2.1×10^2	3.8×10^{-3}
(3)	Polar summer			1.9×10^2	1.1×10^{-3}	1.0×10^{-3}	5.0×10^{-4}	4.4×10^3	1.8×10^{-4}
(4)	Polar winter			4.4×10^2	4.5×10^{-4}	1.7×10^3	2.9×10^{-4}	8.0×10^3	1.0×10^{-4}
(1)	Equatorial summer	11	6.3 to 49	2.7×10^6	7.4×10^{-8}	1.5×10^7	3.3×10^{-8}	6.6×10^7	1.2×10^{-8}
(2)	Equatorial winter			3.7×10^5	5.4×10^{-7}	1.6×10^6	3.1×10^{-7}	6.7×10^6	1.2×10^{-7}
(3)	Polar summer			5.3×10^6	3.8×10^{-8}	2.3×10^7	2.2×10^{-8}	1.1×10^8	7.3×10^{-9}
(4)	Polar winter			8.4×10^6	2.4×10^{-8}	4.3×10^7	1.2×10^{-7}	----	----

ATOMICS INTERNATIONAL

A Division of North American Aviation, Inc.

NO. NAA SR TDR 11127DATE 6-9-65PAGE 30 OF 41

TABLE 6

Selected Wind Profiles Used in Fallout Computations⁽²⁶⁾

(See Table 5)

Altitude (km)	San Andres Island				Sault St. Marie			
	(1)		(2)		(3)		(4)	
	12GCT 16 Aug'62		00GCT 19 Feb'62		12GCT 17 Aug'62		12GCT 21 Feb'62	
	deg	msec ⁻¹	deg	msec ⁻¹	deg	msec ⁻¹	deg	msec ⁻¹
1	090	17	084	12	287	09	032	02
2	090	13	076	10	301	15	296	08
3	070	07	085	07	304	17	281	22
4	060	07	065	06	300	22	282	26
5	072	14	140	02	297	28	280	32
6	080	12	120	02	292	32	272	30
7	095	12	118	04	292	35	265	35
8	094	10	117	06	290	33	266	39
9	100	06	205	06	292	35	261	40
10	110	05	292	12	293	36	261	44
11	100	04	250	15	297	34	263	43
12	240	04	257	17	294	29	267	40
13	160	04	258	23	294	27	268	42
14	150	05	285	23	293	22	272	36
15	210	03	303	16	294	19	270	34
16	150	08	285	07	288	13	273	27
17	125	13	245	04	284	10	265	30
18	128	13	200	01	287	09	267	29
19	115	13	064	02	272	06	273	26
20	115	20	084	02	281	04	280	25
21	135	18	092	01	109	01	288	23
22	115	16	245	05	112	01	292	19
23	118	20	243	02	112	01	282	21
24	118	30	114	01	111	03	284	24
25	118	34	096	04	105	05	300	23
26	115	34	095	05	096	07	300	27
27	115	26	093	08	070	09	313	19
28	118	36	081	14	073	08	318	20
29	118	37	076	15	096	07	302	25
30	118	37	085	17	098	07	295	26

6-9-65

Page 31 Of 41

FIG. 1 THE TEMPERATURE DISTRIBUTION AND NOMENCLATURE OF THE UPPER ATMOSPHERE (1)

FIG. 2 MERIDIONAL SECTION REPRESENTATIVE OF NORTH AMERICA AREA SHOWING TROPOPAUSE AND ISOTHERMS (THIN LINES). (ADAPTED FROM REFERENCE 3, FIG. 6)

FIG. 3 (a) VARIATION OF TEMPERATURE WITH ALTITUDE (b) REPRESENTATION OF POSSIBLE HISTORY OF R_h^{102} TRANSPORT, ⁽⁶⁾

FIG. 4 COMBUSTION OF NON VOLATILE METAL PARTICLES IN A HOT OXIDIZING ATMOSPHERE. (16)

FIG. 5 MAXIMUM STABLE DROPLET SIZE FROM AN ABLATING BISMUTH MELT DURING ENTRY IN THE EARTH'S ATMOSPHERE.

FIG. 6 DISTRIBUTION OF THE NUMBER OF PARTICLES OBTAINED DURING THE ABLATION OF THREE FUEL SAMPLES AS A FUNCTION OF PARTICLE SIZE.

FIG. 7 VOLUME DISTRIBUTION OF PARTICLES OBTAINED DURING THE ABLATION OF THREE FUEL SAMPLES.

Classification
Unclassified
Date
6/19/67
Author: Charles S. Jones

FIG. 9 TIME REQUIRED FOR SPHERICAL PARTICLES OF A GIVEN RADIUS TO FALL FROM A GIVEN INITIAL ALTITUDE TO SEA LEVEL. (DENSITY OF PARTICLES = 6.0 gm/cm^3)

FIG. 10 TERMINAL VELOCITY AS A FUNCTION OF ALTITUDE FOR A CYLINDER OF DIAMETER, d , AND LENGTH, $2d$. (DENSITY OF PARTICLES = 6.0 g/cm^3)

 Classified
 Declassify on: OADR
 Date: 6/6/85
 Author: [Signature]
 Authorized Executive Signature

NAA SR TDR 11127
 6-9-65
 Page 37 of 41

PARTICLE DIAMETER IN MICRONS (DIAMETER, d , LENGTH = $2d$)

FIG. 11 TIME REQUIRED FOR CYLINDRICAL PARTICLES OF DIAMETER, d , AND LENGTH, $2d$, TO FALL FROM A GIVEN INITIAL ALTITUDE TO SEA LEVEL. (DENSITY OF PARTICLES = 6.0 gm/cm^3)

6-9-65

Page 39 Of 41

<i>Unclassified</i> Classification RD/DI Group		
<i>St. Fields</i> Authorized Classifier Signature		Date: 6/10/65

FIG. 12 RATIO OF TERMINAL VELOCITY OF ELLIPSOID
TO TERMINAL VELOCITY OF SPHERE OF SAME
VOLUME.

 FIG. 13 FRACTIONAL MASS PER UNIT AREA FOR $Li-Zr-H$.⁽²⁷⁾

PREPARED BY:	ATOMICS INTERNATIONAL A DIVISION OF NORTH AMERICAN AVIATION, INC.	PAGE NO. 41 OF 41
CHECKED BY:		NAA SR TDR REPORT NO. 11127
DATE: 6-9-65		MODEL NO.

FIG. 14 MEAN JANUARY AND JULY ZONAL WINDS (m/sec) TO 120 Km
BETWEEN EQUATOR AND 75°N. (30) (DASHED LINES - DATA SCANT OR INDEFINITE)

Classification	RD/DI	Group
<i>Unclassified</i>		
<i>J. R. Fields</i>		Date: 6/10/65
Authorized Classifier Signature		