HEALTH AND SAFETY RESEARCH DIVISION

Nuclear and Chemical Waste Programs
(Activity No. AH 10 05 00 0; ONLWCO1)

RADIOLOGICAL SURVEY OF LATTY AVENUE
IN THE VICINITY OF THE FORMER COTTER SITE,
HAZELWOOD/BERKELEY, MISSOURI (LM001)

W. D. Cottrell and R. F. Carrier*

*Biology Division

Date of Issue — May 1987

B. A. Berven — Environmental Assessments Program Manager
W. D. Cottrell — FUSRAP Project Director

Work performed as part of the
ENVIRONMENTAL ASSESSMENTS PROGRAM

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
operated by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under Contract No. DE-AC05-84OR21400

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

MASTER

eb
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

CONTENTS

FIGURES	v
TABLES	vii
ACKNOWLEDGMENTS	ix
ABSTRACT	xi
INTRODUCTION	1
SURVEY METHODS	2
SURVEY RESULTS	2
GAMMA EXPOSURE RATE MEASUREMENTS	2
SURFACE SOIL SAMPLES	3
SUBSURFACE SOIL SAMPLES AND GAMMA LOGGING OF AUGER HOLES	3
SUBSURFACE WATER SAMPLES	4
SIGNIFICANCE OF FINDINGS	4
REFERENCES	6
APPENDIX—GAMMA PROFILE GRAPHS OF AUGER HOLES AT PROPERTY LM001	77

FIGURES

1	Diagram showing location of the Latty Avenue survey (LM001)	7
2	Diagram showing grid system used at Latty Avenue (LM001)	9
3	Locations of surface soil samples taken at Latty Avenue (LM001) between 0+00, BL and 2+00, BL	11
4	Locations of surface soil samples taken at Latty Avenue (LM001) between 2+00, BL and 4+50, BL	12
5	Locations of surface soil and water samples taken at Latty Avenue (LM001) between 4+50, BL and 7+00, BL	13
6	Locations of surface soil and water samples taken at Latty Avenue (LM001) between 7+00, BL and 9+00, BL	14
7	Locations of surface soil samples taken at Latty Avenue (LM001) between 9+00, BL and 11+50, BL	15
8	Locations of surface soil samples taken at Latty Avenue (LM001) between 11+50, BL and 14+00, BL	16
9	Locations of surface soil samples taken at Latty Avenue (LM001) between 14+00, BL and 16+50, BL	17
10	Locations of surface soil samples taken at Latty Avenue (LM001) between 16+50, BL and 19+00, BL	18
11	Locations of surface soil samples taken at Latty Avenue (LM001) between 19+00, BL and 21+50, BL	19
12	Locations of surface soil samples taken at Latty Avenue (LM001) between 21+50, BL and 24+00, BL	20
13	Locations of surface soil samples taken at Latty Avenue (LM001) between 24+00, BL and 26+50, BL	21
14	Locations of surface soil samples taken at Latty Avenue (LM001) between 26+50, BL and 29+00, BL	22
15	Locations of surface soil samples taken at Latty Avenue (LM001) between 29+00, BL and 31+50, BL	23

16	Locations of auger holes at Latty Avenue (LM001) between 0+00, BL and 2+00, BL	24
17	Locations of auger holes at Latty Avenue (LM001) between 2+00, BL and 4+50, BL	25
18	Locations of auger holes at Latty Avenue (LM001) between 4+50, BL and 7+00, BL	26
19	Locations of auger holes at Latty Avenue (LM001) between 7+00, BL and 9+00, BL	27
20	Locations of auger holes at Latty Avenue (LM001) between 9+00, BL and 11+50, BL	28
21	Locations of auger holes at Latty Avenue (LM001) between 11+50, BL and 14+00, BL	29
22	Locations of auger holes at Latty Avenue (LM001) between 14+00, BL and 16+50, BL	30
23	Locations of auger holes at Latty Avenue (LM001) between 16+50, BL and 19+00, BL	31
24	Locations of auger holes at Latty Avenue (LM001) between 19+00, BL and 21+50, BL	32
25	Locations of auger holes at Latty Avenue (LM001) between 21+50, BL and 24+00, BL	33
26	Locations of auger holes at Latty Avenue (LM001) between 24+00, BL and 26+50, BL	34
27	Locations of auger holes at Latty Avenue (LM001) between 26+50, BL and 29+00, BL	35
28	Locations of auger holes at Latty Avenue (LM001) between 29+00, BL and 31+50, BL	36
29	Locations of auger holes at Latty Avenue (LM001) between 31+50, BL and 34+00, BL	37
30	Locations of auger holes at Latty Avenue (LM001) between 34+00, BL and 36+50, BL	38
31	Locations of auger holes at Latty Avenue (LM001) between 36+50, BL and 38+50, BL	39

TABLES

1	Applicable guidelines for protection against radiation	40
2	Background radiation levels in the St. Louis area	41
3	Gamma exposure rate measurements at Latty Avenue (LM001)	42
4	Results of analyses of soil samples taken at Latty Avenue (LM001)	51
5	Extent of subsurface contamination on the Latty Avenue site (LM001) as indicated by scintillation probe loggings and sample analyses	69
6	Concentration of selected radionuclides in water samples taken from auger holes at the Latty Avenue site (LM001)	74
7	Summary of measurements and sample results at Latty Avenue (LM001)	75

ACKNOWLEDGMENTS

Research for this project was sponsored by the Division of Facility and Site Decommissioning Projects, U.S. Department of Energy, under Contract No. DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc. The authors wish to acknowledge the support of J. E. Baublitz, Deputy Director, Office of Remedial Action and Waste Technology; J. J. Fiore, Director, Division of Facility and Site Decommissioning Projects; and members of their staff. The authors also appreciate the contributions of A. C. Butler, R. W. Doane, B. S. Ellis, J. L. Marley, D. A. Roberts, J. A. Roberts, W. H. Shimpagh, T. R. Stewart, and W. Winton of the RASA group, and C. Smith and R. Smith of Don Stone Associates for participation in collection, analyses, and reporting of data for this survey.

ABSTRACT

A radiological survey was conducted over a proposed construction corridor along Latty Avenue, Hazelwood/Berkeley, Missouri, in the vicinity of the former Cotter site at 9200 Latty Avenue. The survey included gamma exposure rates at the ground surface and at 1 m above the surface throughout the site, sampling of surface soil, sampling of subsurface soil from auger holes, gamma logging of auger holes, and sampling of subsurface water. The results of the survey demonstrated some degree of radioactive contamination in all areas of the construction corridor, extending north and south in some regions onto adjacent private properties. Redistribution of the contamination by flooding, surface runoff, and road and utility line activities was evident. The pattern of contamination ranged from widespread to isolated spots and was found to occur from near the surface to depths of ~1.8 m. The most highly contaminated region was noted on both sides of Latty Avenue adjacent to the former Cotter site. Concentrations of ^{230}Th in soil from that region were as high as 16,000 pCi/g.

INTRODUCTION

Process waste and residues associated with the production and refining of uranium materials were generated by the Mallinckrodt Chemical Works, St. Louis, Missouri, during the period 1942 through the 1950s. According to a Nuclear Regulatory Commission (NRC) report,¹ residues which had been stored at the St. Louis Airport property were moved by the Continental Mining and Milling Company of Chicago, Illinois, to the Latty Avenue Storage Site in Hazelwood, Missouri, in early 1966. The Latty Avenue storage site is located in a low-lying industrialized area in Hazelwood/Berkeley, Missouri, approximately 12 miles northwest of St. Louis (Fig. 1). The Commercial Discount Corporation, Chicago, Illinois, purchased the residues in January 1967. Much of the material was dried and shipped to the Cotter Corporation facilities in Canon City, Colorado. These materials included ore residues and uranium-, thorium-, and radium-bearing process wastes. Materials remaining at the Latty Avenue Storage Site were sold to the Cotter Corporation in December 1969. Records indicate that residues remaining on the site at that time included 74,000 tons of Belgian Congo pitchblende raffinate containing 113 tons of uranium, 32,500 tons of Colorado raffinate containing 48 tons of uranium, and 8700 tons of leached barium sulfate containing about 7 tons of uranium. During August through November 1970, Cotter Corporation dried some of the remaining residues and shipped them to a Cotter mill in Canon City, Colorado. An estimated 10,000 tons of Colorado raffinate and 8700 tons of leached barium sulfate remained at the Latty Avenue site in December 1970.

In April 1974, an NRC inspector was informed that the remaining Colorado raffinate had been shipped to Canon City during the prior year (1973) without drying, and that the leached barium sulfate had been transported to a landfill area in St. Louis County. Twelve to eighteen inches of top soil were reported to have been stripped from the Latty Avenue site surface and supposedly were removed from the site with the leached barium sulfate. However, analysis of soil samples taken during an NRC investigation of the site in 1976 indicated the presence of uranium- and thorium-bearing residues.

In 1977, a survey was conducted by Oak Ridge National Laboratory (ORNL) to characterize the radiological condition of the site. It was determined that alpha and beta-gamma contamination levels on building surfaces exceeded the NRC guidelines for release of decontaminated property for unrestricted use.² Numerous areas having gamma exposure rates of 300–500 $\mu\text{R}/\text{h}$ were clearly above criteria recommended by the International Commission on Radiological Protection (ICRP).³ It was estimated that the top three inches of soil on much of the area contained an average of 140 pCi/g of ^{226}Ra and probably higher average concentrations of ^{238}U , ^{230}Th , and ^{227}Ac .

At the request of the U.S. Department of Energy (DOE), a preliminary survey of properties adjacent to and in the vicinity of the former Cotter site at 9200 Latty Avenue was made in September 1983 to determine if contamination was present on any vicinity properties and to evaluate the scope of work required to perform radiological assessment surveys of these properties.⁴ Elevated gamma radiation levels were observed on all properties adjacent to 9200 Latty Avenue, especially on those properties to the north, east, and south.

A radiological survey was performed by ORNL during the period January through April, 1984, over a section of Latty Avenue in the vicinity of the former Cotter site (the Jarboe Realty and Investment Co.). The area surveyed was a proposed construction corridor along Latty Avenue and included the asphalt-covered street and shoulders and ditches on both sides of the street. The width of the corridor varied from 70 to 295 ft and extended from Coldwater Creek eastward along Latty Avenue to its intersection with interstate highway 170, a distance of 3850 ft. The results of surveys of the fence line surrounding the former Cotter site and of other private properties in the vicinity of that site are provided in separate reports.^{5,6}

SURVEY METHODS

The survey included the following measurements: (1) gamma radiation exposure rates at the surface and at 1 m from the surface throughout the site; (2) concentrations of ^{226}Ra , ^{230}Th , and ^{238}U in surface and subsurface soil on the site; (3) concentrations of ^{226}Ra , ^{230}Th , ^{238}U , and ^{210}Pb in subsurface water on the site; and (4) gamma radiation levels at various depths in auger holes drilled on site. The radiological survey followed a general plan developed at ORNL for vicinity properties in Hazelwood and Berkeley, Missouri.⁷

The site was divided into 50 ft $\times$ 50 ft "survey blocks" by the rectangular grid system shown in Fig. 2. A comprehensive description of survey methods and instrumentation is given in a separate report.⁸

SURVEY RESULTS

Applicable federal guidelines for radiation exposure to the general public are summarized in Table 1. Normal radiation background levels and radionuclide concentrations in soil in the Hazelwood/St. Louis area are presented in Table 2. These data are provided for comparison with the survey results presented in this section.

All direct measurements presented in this report are gross readings: background radiation levels have not been subtracted. Similarly, background concentrations have not been subtracted from radionuclide concentrations measured in environmental samples.

GAMMA EXPOSURE RATE MEASUREMENTS

Results of grid point measurements of gamma exposure rates at the surface and at 1 m above the surface and the range of gamma radiation levels detected during the gamma scan of grid blocks are presented in Table 3. Grid point and grid block locations are shown in Fig. 2. The maximum gamma exposure rate measured on this site during the grid block scan, 720 $\mu\text{R}/\text{h}$, exceeds background by a factor of about 60. The gamma exposure rates measured at grid points at the ground surface ranged from 4 to 490 $\mu\text{R}/\text{h}$ and averaged 17. Grid point gamma exposure rates at 1 m above the ground surface ranged from 4 to 380 $\mu\text{R}/\text{h}$ with a mean of about 16.

SURFACE SOIL SAMPLES

Surface soil samples were collected from systematic locations to provide representative sampling of the site. Samples were normally collected from a depth of 0–15 cm. However, in some cases, for further definition of contamination, sampling was extended to depths of 75 cm. In addition to systematic sampling, samples were collected from selected locations showing elevated gamma radiation levels and are designated as “biased” samples. Sample locations are shown in Figs. 3–15, with results of sample analyses provided in Table 4.

Concentrations of ^{226}Ra in systematic surface soil samples ranged from 1.0 to 12 pCi/g and averaged 2.7 pCi/g; ^{238}U concentrations ranged from 0.99 to 15 pCi/g and averaged 2.1 pCi/g. Mean concentrations of ^{226}Ra and ^{238}U exceeded background levels by factors of about 3 and 2, respectively. Concentrations of ^{230}Th in surface soil ranged from 1.6 to 630 pCi/g with a mean of 54 pCi/g. The maximum concentration of ^{230}Th exceeded the DOE soil guideline by a factor of ~ 120 while the mean concentration exceeded the guideline value by about 10.

The three biased samples were found to contain concentrations of ^{230}Th ranging from 350 to 810 pCi/g. Concentrations of ^{226}Ra and ^{238}U ranged from 8.3 to 12 pCi/g, and 5.4 to 8.7 pCi/g, respectively.

SUBSURFACE SOIL SAMPLES AND GAMMA LOGGING OF AUGER HOLES

Holes were augered at locations shown in Figs. 16–31. One or more soil samples were collected from each hole. The sampling depths chosen usually included the region of maximum gamma radiation level as indicated by the gamma log. At selected locations, Shelby tubes and/or split-spoon samplers were used to collect subsurface samples at known depths. Concentrations of ^{226}Ra , ^{230}Th , and ^{238}U in subsurface soil are given in Table 4.

The maximum concentration of each radionuclide was found in sample 148 taken from a depth of 0 to 15 cm. Those values are 660, 22,000, and 790 pCi/g for ^{226}Ra , ^{230}Th , and ^{238}U , respectively.

Each of the auger holes and core holes were “logged” using a gamma scintillation detector. A plastic pipe (PVC Schedule 40) with a 4-in inside diameter was placed in the hole and a NaI scintillation probe was lowered inside the pipe. The probe was encased in a lead shield with a horizontal row of collimating slits on the side. This collimation allows measurement of gamma radiation intensities resulting from contamination within small fractions of hole depth. Measurements were usually made at 15- or 30-cm intervals. Logging of the core holes was done to determine the extent of subsurface contamination at each location. Profiles of gamma intensity as a function of depth are presented in the Appendix.

The logging technique used here is not radionuclide specific. However, the logging data, in conjunction with the soil analyses data, may be used to estimate the region of contamination in the auger holes. It appears from a comparison of these data that a reading of 1000 cpm or greater using the shielded scintillator indicates the presence of elevated

concentrations of ^{226}Ra and/or ^{238}U . Consequently, soil giving rise to 1000 cpm or greater on the scintillator or containing radionuclide concentrations above criteria (Table 1) as determined from soil analysis was considered as contaminated soil. Using these criteria, an estimate of the region of contamination in each hole was made. These data are listed in Table 5. A word of caution should be added in the interpretation of the data presented in Table 5. Only those holes yielding a positive indication of contamination, using the previously stated criteria, are listed as having a region of contamination; an entry of "none" opposite a location means only that no positive indication of contamination was observed using the "logging" and sampling procedures employed on this site. Because of the difficulty of detecting ^{230}Th with this in-situ hole logging device, concentrations of ^{230}Th in the absence of ^{226}Ra considerably in excess of 5 pCi/g may have gone undetected.

SUBSURFACE WATER SAMPLES

Water samples were collected from auger holes and analyzed for ^{226}Ra , ^{238}U , ^{230}Th , and ^{210}Pb . Locations of the holes from which the samples were taken are shown in Figs. 5 and 6 with the results of laboratory analyses provided in Table 6. Concentrations of ^{226}Ra , ^{238}U , ^{230}Th , and ^{210}Pb were well below the concentration guides for water for unrestricted use (10 CFR 20).

SIGNIFICANCE OF FINDINGS

A summary of survey measurements and results is provided in Table 7. Radioactive contamination was found to some extent in all areas of the construction corridor. The corridor may be further characterized by areas as follows.

1. 0+00 to 4+50 (Figs. 3 and 4): This area lies in a flood plain. The contamination is fairly uniformly distributed over the surface and generally extends to a depth of about 30 cm. In a few isolated areas where the soil has been disturbed, the materials may extend to depths of 120 cm. Soil mixed with the radioactive materials was apparently eroded from the former processing site and carried to this area by flood waters. Concentrations of ^{230}Th in this area ranged from about 5 pCi/g to 280 pCi/g.
2. 4+50 to 9+00 (Figs. 5 and 6): This section of the corridor encompasses drainage ditches, culverts, road shoulders, and asphalt roadways and extends into private property on both the north and south sides of Latty Avenue. The drainage ditches, culverts, and road shoulders are highly contaminated. The private property is contaminated to some extent. The contamination extends to depths greater than 1.8 m along the road shoulders on the north side of Latty Avenue (drill holes 119 and 212, Table 6), extends under the asphalt of the roadway at least 1.8 m from the north edge of the pavement (drill holes 116 and 118, Table 6), and may in some places underlie the complete width of the asphalt surface. Concentrations of ^{230}Th in soil samples from this area were as high as 22,000 pCi/g. The north end of the former Cotter site lies along the south edge of Latty Avenue and a strip of the former site, ~200 ft deep, was included in the construction corridor. Radioactive contamination on this part of the site extends to depths as great as 1.8 m.

3. 9+00 to 30+00 (Figs. 7 to 15): The contamination along this section of Latty Avenue is spotty and, except for areas that have been disturbed by construction, lies near the surface (<0.5 m). In areas that have undergone construction (e.g., laying of pipe lines or rebuilding of roadway), the radioactively contaminated soil may extend to depths of 1.5 m or greater (Table 6). The pattern of contamination in this area suggests that radioactive residues may have spilled from trucks traveling this route to and from the site and that the contamination may have been further redistributed by surface runoff, road repair, and the laying and/or repair of underground utility lines.
4. 30+00 to 38+50 (Figs. 28 to 31): This section of Latty Avenue has been replaced with a new concrete roadway that appears to be elevated above the original grade of the old road. Sampling along this section was confined to areas not covered by concrete. No attempt was made to sample the fill underneath the concrete roadway. Radioactive contamination found along this section of Latty Avenue appears to be confined to isolated spots and concentrations in general are much lower than those measured in other areas of the corridor.
5. Asphalt Road Surface: Much of the asphalt road surface along Latty Avenue has become contaminated with radioactive materials. Nineteen of 47 asphalt samples analyzed were found to contain concentrations of ^{230}Th above the DOE guideline of 5 pCi/g. The concentrations of ^{230}Th in asphalt ranged from <1 to 600 pCi/g.

REFERENCES

1. Nuclear Regulatory Commission I&E Investigation Report No. 76-01, August 20, 1976.
2. Nuclear Regulatory Commission (NRC), *Standards for Protection Against Radiation*, 10 CFR 20.105 (November 17, 1960).
3. International Commission on Radiological Protection, *Recommendations of the International Commission on Radiological Protection*, ICRP Publication 26, Pergamon Press, Elmsford, New York (1977).
4. W. D. Cottrell to B. A. Berven, Report on Trip to the Latty Avenue Site, Hazelwood, Missouri (September 27, 1983).
5. R. F. Carrier and W. D. Cottrell, *Radiological Survey of the Perimeter Fence Line of the Former Cotter Site, Hazelwood, Missouri (LM002)*, Oak Ridge National Laboratory, ORNL/TM-10007 (December 1986).
6. W. D. Cottrell, R. F. Carrier, and C. A. Johnson, *Radiological Survey of Properties in the Vicinity of the Former Cotter Site, Hazelwood/Berkeley, Missouri (LM003)*, Oak Ridge National Laboratory, ORNL/TM-10008 (in press).
7. *Radiological Survey Plan for Vicinity Properties in Hazelwood and Berkeley, Missouri*, letter from B. A. Berven, ORNL/RASA, to A. J. Whitman, DOE/HQ (October 1983).
8. T. E. Myrick, B. A. Berven, W. D. Cottrell, W. A. Goldsmith, and F. F. Haywood, *Procedures Manual for the Remedial Action Survey and Certification Activities (RASCA) Program*, Oak Ridge National Laboratory, ORNL/TM-8600 (September 1982).
9. T. E. Myrick, B. A. Berven, and F. F. Haywood, *State Background Radiation Levels: Results of Measurements Taken during 1975-1979*, ORNL/TM-7343 (November 1981).

7/8

ORNL-DWG 87-11427

Fig. 1. Diagram showing general location of the Latty Avenue survey (LM001).

Fig. 2. Diagram showing grid system used at Latty Avenue (LM001).

Fig. 3. Locations of surface soil samples taken at Latty Avenue (LM001) between 0+00, BL and 2+00, BL.

Fig. 4. Locations of surface soil samples taken at Latty Avenue (LM001) between 2+00, BL and 4+50, BL.

Fig. 5. Locations of surface soil and water samples taken at Latty Avenue (LM001) between 4+50, BL and 7+00, BL.

Fig. 6. Locations of surface soil and water samples taken at Latty Avenue (LM001) between 7+00, BL and 9+00, BL.

Fig. 7. Locations of surface soil samples taken at Latty Avenue (LM001) between 9+00, BL and 11+50, BL.

ORNL-DWG 87-11434

Fig. 8. Locations of surface soil samples taken at Latty Avenue (LM001) between 11+50, BL and 14+00, BL.

Fig. 9. Locations of surface soil samples taken at Latty Avenue (LM001) between 14+00, BL and 16+50, BL.

Fig. 10. Locations of surface soil samples taken at Latty Avenue (LM001) between 16+50, BL and 19+00, BL.

ORNL-DWG 87-11437

Fig. 11. Locations of surface soil samples taken at Latty Avenue (LM001) between 19+00, BL and 21+50, BL.

Fig. 12. Locations of surface soil samples taken at Latty Avenue (LM001) between 21+50, BL and 24+00, BL.

ORNL-DWG 87-11439

Fig. 13. Locations of surface soil samples taken at Latty Avenue (LM001) between 24+00, BL and 26+50, BL.

Fig. 14. Locations of surface soil samples taken at Latty Avenue (LM001) between 26+50, BL and 29+00, BL.

Fig. 15. Locations of surface soil samples taken at Latty Avenue (LM001) between 29+00, BL and 31+50, BL.

Fig. 16. Locations of auger holes at Latty Avenue (LM001) between 0+00, BL and 2+00, BL.

Fig. 17. Locations of auger holes at Latty Avenue (LM001) between 2+00, BL and 4+50, BL.

Fig. 18. Locations of auger holes at Latty Avenue (LM001) between 4+50, BL and 7+00, BL.

Fig. 19. Locations of auger holes at Latty Avenue (LM001) between 7+00, BL and 9+00, BL.

Fig. 20. Locations of auger holes at Latty Avenue (LM001) between 9+00, BL and 11+50, BL.

Fig. 21. Locations of auger holes at Latty Avenue (LM001) between 11+50, BL and 14+00, BL.

Fig. 22. Locations of auger holes at Latty Avenue (LM001) between 14+00, BL and 16+50, BL.

Fig. 23. Locations of auger holes at Latty Avenue (LM001) between 16+50, BL and 19+00, BL.

Fig. 24. Locations of auger holes at Latty Avenue (LM001) between 19+00, BL and 21+50, BL.

Fig. 25. Locations of auger holes at Latty Avenue (LM001) between 21+50, BL and 24+00, BL.

ORNL-DWG 87-11452

Fig. 26. Locations of auger holes at Latty Avenue (LM001) between 24+00, BL and 26+50, BL.

Fig. 27. Locations of auger holes at Latty Avenue (LM001) between 26+50, BL and 29+00, BL.

Fig. 28. Locations of auger holes at Latty Avenue (LM001) between 29+00, BL and 31+50, BL.

Fig. 29. Locations of auger holes at Latty Avenue (LM001) between 31+50, BL and 34+00, BL.

ORNL-DWG 87-11456

Fig. 30. Locations of auger holes at Latty Avenue (LM001) between 34+00, BL and 36+50, BL.

Fig. 31. Locations of auger holes at Latty Avenue (LM001) between 36+50, BL and 38+50, BL.

Table 1. Applicable guidelines for protection against radiation

Mode of exposure	Exposure conditions	Guideline value	Guideline source
Radionuclide concentrations in soil	Maximum permissible concentration of the following radionuclides in soil above background levels averaged over 100 m ² area ²³² Th ²³⁰ Th ²²⁸ Ra ²²⁶ Ra	5 pCi/g averaged over the first 15 cm of soil below the surface; 15 pCi/g when averaged over 15-cm-thick soil layers, more than 15 cm below the surface	U.S. Department of Energy Guidelines for Residual Radioactivity at Formerly Utilized Sites Remedial Action Program and Remote Surplus Facilities Management
Radionuclides in water	Maximum permissible concentration of the following radionuclides in water for unrestricted use ²³⁸ U ²³⁰ Th ²²⁶ Ra ²¹⁰ Pb	4×10^4 pCi/L 2×10^3 pCi/L 30 pCi/L 100 pCi/L	NRC 10 CFR 20.106 Appendix B, Table II
	Maximum contaminant level combined ²²⁶ Ra and ²²⁸ Ra in drinking water	5 pCi/L	EPA-National Primary Drinking Water Regulations (40 CFR 141)

Table 2. Background radiation levels in the St. Louis area

Type of radiation measurement or sample	Radiation level or radionuclide concentration	
	Range	Average
Gamma exposure rate at 1 m above floor or ground surface ($\mu\text{R/h}$)	5–8	6
Concentration of radionuclides in soil (pCi/g)		
^{238}U	0.33–1.2	0.91
^{226}Ra	0.31–1.3	0.96
$^{230}\text{Th}^a$	0.33–1.2	0.91

^a ^{230}Th assumed to be in equilibrium with ^{238}U .

**Table 3. Gamma exposure rate measurements at
Latty Avenue (LM001)**

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
0+00, BL	7	8	7-10
0+50, BL	9	10	9-11
1+00, BL	10	11	9-12
1+50, BL	10	11	9-12
2+00, BL	12	11	11-21
2+50, BL	11	11	9-15
3+00, BL	12	12	9-16
3+50, BL	14	14	13-17
4+00, BL	13	12	11-26
4+50, BL	23	21	11-160
5+00, BL	20	20	15-160
5+40, BL	28	21	21-39
5+50, BL	26	21	21-39
6+00, BL	32	21	21-26
6+50, BL	34	26	17-26
7+00, BL	28	23	14-23
7+50, BL	21	21	13-21
8+00, BL	21	21	11-21
8+50, BL	21	21	6-30
9+00, BL	16	15	11-55
9+50, BL	13	13	11-46
10+00, BL	13	13	9-29
10+50, BL	11	10	9-24
11+00, BL	7	7	6-26
11+50, BL	7	7	6-24
12+00, BL	7	7	6-10
12+50, BL	7	7	6-20
13+00, BL	6	6	5-11
13+50, BL	6	6	5-11
14+00, BL	6	6	05-13
14+50, BL	7	6	06-49
15+00, BL	6	6	05-37
15+50, BL	8	7	05-33
16+00, BL	7	7	06-33
16+50, BL	9	8	06-37
17+00, BL	8	7	05-20
17+50, BL	7	7	06-28

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
18+00, BL	7	6	5-21
18+50, BL	6	6	5-28
19+00, BL	7	6	5-23
19+50, BL	7	7	6-28
20+00, BL	7	7	5-39
20+50, BL	6	7	5-32
21+00, BL	6	6	5-21
21+50, BL	6	6	5-26
22+00, BL	5	5	5-12
22+50, BL	6	9	5-10
23+00, BL	5	7	5-10
23+50, BL	6	6	5-17
24+00, BL	6	6	5-17
24+50, BL	6	6	5-13
25+00, BL	7	6	4-10
25+50, BL	6	6	5-10
26+00, BL	6	5	4-10
26+50, BL	5	5	5-10
27+00, BL	5	5	4-9
27+50, BL	6	6	5-15
28+00, BL	5	6	4-9
28+50, BL	5	6	5-12
29+00, BL	5	6	4-24
29+50, BL	11	19	4-21
30+00, BL	5	5	4-21
30+50, BL	5	5	5-10
31+00, BL	5	5	5-10
31+50, BL	5	6	5-10
32+00, BL	6	5	5-10
32+50, BL	6	6	5-10
33+00, BL	5	5	5-10
33+50, BL	5	6	5-10
34+00, BL	6	5	5-10
34+50, BL	5	6	5-7
35+00, BL	5	6	5-7
35+50, BL	6	5	5-7
36+00, BL	6	6	5-7
36+50, BL	5	6	5-7
37+00, BL	5	6	5-7

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
37+50, BL	5	6	5-7
38+00, BL	5	6	5-7
38+50, BL	6	8	—
0+00, 35R	d	d	7-10
0+50, 35R	10	10	9-11
1+00, 35R	12	11	9-12
1+50, 35R	10	11	9-12
2+00, 35R	12	12	11-21
2+50, 35R	14	15	11-21
3+00, 35R	12	12	—
3+00, 40R	12	12	12-15
3+50, 40R	12	12	12-20
4+00, 40R	16	17	15-59
4+50, 40R	21	19	19-160
5+00, 40R	26	23	21-660
5+50, 40R	79	57	—
5+50, 50R	79	92	92-280
6+00, 50R	180	180	26-720
6+50, 50R	59	46	26-330
7+00, 50R	240	260	13-260
7+50, 50R	79	87	12-130
8+00, 50R	26	24	19-29
8+50, 50R	24	24	21-39
8+80, 40R	26	26	17-26
9+00, 40R	26	26	13-26
9+50, 40R	23	23	13-33
10+00, 40R	16	15	13-59
10+50, 40R	14	14	9-39
11+00, 40R	11	11	9-26
11+50, 40R	9	9	6-9
12+00, 40R	9	8	6-26
12+50, 40R	8	7	6-21
13+00, 40R	6	6	6-7
13+50, 40R	6	5	6-26
14+00, 40R	6	6	6-53
14+24, 50R	9	8	—
14+50, 50R	7	9	5-12
15+00, 50R	7	6	6-99
15+50, 50R	6	8	6-53

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
16+00, 50R	7	8	6-53
16+50, 50R	7	7	6-55
17+00, 50R	9	9	6-49
17+50, 50R	9	9	6-50
18+00, 50R	8	7	6-20
18+50, 50R	9	9	6-22
19+00, 50R	9	9	6-29
19+50, 50R	10	9	6-32
20+00, 50R	9	9	6-33
20+50, 50R	9	9	6-22
21+00, 50R	10	9	6-29
21+50, 50R	9	10	6-17
22+00, 50R	4	4	4-12
22+50, 50R	7	7	4-10
23+00, 50R	5	5	5-19
23+50, 50R	9	8	5-11
24+00, 50R	7	9	5-23
24+50, 50R	9	9	6-21
25+00, 50R	7	8	5-26
25+50, 50R	7	7	5-26
26+00, 50R	7	6	4-26
26+50, 30R	5	4	-
26+50, 50R	4	6	4-32
27+00, 30R	4	5	4-12
27+50, 30R	4	5	4-21
28+00, 30R	4	4	4-26
28+50, 30R	5	5	4-21
29+00, 30R	5	5	8-29
29+50, 30R	8	9	7-29
30+00, 30R	6	6	6-26
30+50, 30R	9	9	-
30+50, 20R	9	9	7-10
31+00, 20R	7	7	7-10
31+50, 20R	7	8	7-10
32+00, 20R	7	7	7-86
32+50, 20R	7	8	7-10
33+00, 20R	7	7	7-10
33+50, 20R	8	7	7-10
34+00, 20R	7	8	7-10

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
34+50, 20R	7	6	6-10
35+00, 20R	7	8	7-10
35+50, 20R	7	7	7-10
36+00, 20R	8	7	7-10
36+50, 20R	6	7	7-10
37+00, 20R	9	9	7-10
37+50, 20R	8	8	7-10
38+00, 20R	7	8	7-10
38+50, 20R	6	7	—
18+00, 65R	7	6	7-9
18+50, 65R	8	8	7-9
19+00, 65R	7	7	7-10
19+50, 65R	10	10	8-11
20+00, 65R	9	9	8-11
20+50, 65R	6	7	6-10
21+00, 65R	10	10	6-10
21+50, 65R	6	6	4-9
22+00, 65R	5	4	4-9
22+50, 65R	6	6	4-9
23+00, 65R	5	5	6-10
23+50, 65R	7	9	7-9
24+00, 65R	8	9	7-9
24+50, 65R	8	8	7-9
25+00, 65R	7	7	7-9
25+50, 65R	6	5	6-9
26+00, 65R	7	7	6-9
26+50, 65R	4	4	—
5+50, 75R	72	68	39-720
6+00, 75R	160	160	39-210
6+50, 75R	42	39	39-460
7+00, 75R	53	37	39-340
7+20, 75R	130	120	—
7+20, 100R	33	26	39-280
7+50, 100R	45	42	24-290
8+00, 100R	92	63	20-110
8+50, 100R	99	79	20-260
7+20, 150R	26	22	33-79
7+50, 150R	32	32	39-400
8+00, 150R	87	59	79-530

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
8+50, 150R	380	490	79-530
7+20, 200R	58	55	33-79
7+50, 200R	76	84	39-200
8+00, 200R	66	66	66-560
8+50, 200R	86	130	99-600
7+20, 210R	42	46	46-99
7+50, 210R	59	55	53-105
8+00, 210R	68	64	53-132
8+50, 210R	66	55	66-329
0+00, 35L	e	e	-
0+50, 35L	11	11	-
1+00, 35L	9	9	-
1+50, 35L	10	10	-
2+00, 35L	11	11	-
2+50, 35L	14	14	-
2+90, 35L	10	10	-
2+90, 40L	10	10	-
3+00, 40L	10	9	-
3+50, 40L	13	14	-
4+00, 40L	16	17	-
4+50, 40L	16	12	-
5+00, 40L	28	26	-
5+50, 40L	26	26	-
5+50, 50L	21	21	19-23
6+00, 50L	21	23	17-28
6+50, 50L	30	28	11-28
7+00, 50L	15	14	14-17
7+50, 50L	15	15	13-16
8+00, 50L	9	9	9-14
8+50, 50L	10	9	9-13
9+00, 50L	9	9	9-11
9+50, 50L	9	9	7-12
10+00, 50L	10	10	9-12
10+50, 50L	9	7	7-10
11+00, 50L	8	8	6-9
11+50, 50L	7	7	6-9
12+50, 50L	7	7	6-10
12+00, 50L	7	8	6-11

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
13+00, 50L	7	6	6–11
13+50, 50L	4	4	5–7
14+00, 50L	5	6	5–7
14+50, 50L	6	6	5–7
15+00, 50L	5	7	5–10
15+50, 50L	8	8	6–10
16+00, 50L	8	9	6–10
16+50, 50L	7	8	6–10
17+00, 50L	9	9	6–10
17+50, 50L	9	9	6–10
18+00, 50L	9	8	6–10
18+50, 50L	9	8	6–10
19+00, 50L	8	8	6–10
19+50, 50L	7	8	6–10
20+00, 50L	9	8	6–10
20+50, 50L	8	8	6–10
21+00, 50L	9	9	6–10
21+50, 50L	8	9	6–10
22+00, 50L	9	8	6–10
22+50, 50L	9	7	6–10
23+00, 50L	9	9	6–10
23+50, 50L	8	9	6–10
24+00, 50L	8	9	6–10
24+50, 50L	8	8	6–10
25+00, 50L	7	8	6–10
25+50, 50L	9	8	6–10
26+00, 50L	7	7	6–10
26+50, 50L	8	8	6–10
27+00, 50L	7	9	6–10
27+50, 50L	7	8	6–10
28+00, 50L	8	8	6–10
28+50, 50L	7	7	6–10
29+00, 50L	7	7	6–10
29+50, 50L	5	5	–
30+00, 55L	9	9	–
30+50, 55L	7	7	–
31+00, 55L	7	7	–
31+50, 55L	6	6	–
32+00, 55L	7	6	–

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
32+50, 55L	7	7	—
33+00, 55L	7	7	—
33+50, 55L	7	8	—
34+00, 55L	6	7	—
34+50, 55L	7	7	—
35+00, 55L	7	7	—
35+50, 55L	6	7	—
36+00, 55L	7	7	—
36+50, 55L	6	8	—
37+00, 55L	7	7	—
37+50, 55L	6	7	—
38+00, 55L	6	7	—
38+50, 55L	9	7	—
5+40, 85L	16	7	—
5+50, 85L	16	16	—
6+00, 85L	17	16	—
6+50, 85L	17	15	—
7+00, 85L	15	13	—
7+50, 85L	11	8	—
8+00, 85L	12	9	—
8+50, 85L	9	6	—
9+00, 85L	7	5	—
9+50, 85L	6	6	—
10+00, 85L	9	10	—
10+50, 85L	5	5	—
11+00, 85L	7	7	—
11+50, 85L	7	7	—
12+00, 85L	7	7	—
12+50, 85L	9	8	—
13+00, 85L	7	7	—
13+50, 85L	4	4	—
14+00, 85L	5	5	—
14+50, 85L	5	5	—
15+00, 85L	4	5	—
15+50, 85L	8	8	—
16+00, 85L	7	7	—
16+50, 85L	7	8	—
17+00, 85L	9	8	—
17+50, 85L	9	9	—

Table 3 (continued)

Grid location ^a	Grid point measurements ^b ($\mu\text{R/h}$)		Range of gamma exposure rates during scan of grid block ^c ($\mu\text{R/h}$)
	Gamma exposure rate at 1 m	Gamma exposure rate at the surface	
18+00, 85L	9	9	—
18+50, 85L	9	9	—
19+00, 85L	7	8	—
19+50, 85L	7	8	—
20+00, 85L	9	8	—
20+50, 85L	9	9	—
21+00, 85L	8	8	—
21+50, 85L	7	8	—
22+00, 85L	7	7	—
22+50, 85L	9	7	—
23+00, 85L	6	7	—
23+50, 85L	8	9	—
24+00, 85L	9	9	—
24+50, 85L	8	8	—
25+00, 85L	8	8	—
25+50, 85L	9	8	—
26+00, 85L	7	7	—
26+50, 85L	9	9	—
27+00, 85L	8	8	—
27+50, 85L	8	8	—
28+00, 85L	8	8	—
28+50, 85L	7	8	—
29+00, 85L	7	7	—
29+50, 85L	5	6	—

^aGrid location shown on Fig. 1.

^bGrid point measurements are discrete measurements at grid point.

^cGrid block measurements are obtained by a gamma scan of the entire block.

^dInaccessible.

Table 4. Results of analyses of soil samples taken at Latty Avenue (LM001)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
Systematic samples ^e					
S1	3+65, 12R	0-15	1.9 ± 0.7	51	1.9
S2A	3+70, 16L	0-15	2.7 ± 0.3	68	2.4
B		15-30	1.8 ± 0.04	28	1.6
S3A	3+75, 32L	0-15	2.5 ± 0.4	72	2.5
B		15-30	1.2 ± 0.3	12	1.2
C		30-45	1.3 ± 0.1	<5.4	1.1
D		45-60	1.1 ± 0.2	<4.5	1.0
S4A	4+15, 36L	0-15	2.0 ± 0.4	47	1.9
B		15-30	1.6 ± 0.2	22	1.6
S5	3+15, 32L	0-15	2.2 ± 0.3	54	2.2
S6	2+88, 28L	0-15	1.5 ± 0.3	17	1.5
S7A	2+50, 32L	0-15	3.6 ± 0.1	19	3.3
B		15-30	3.5 ± 0.3	15	3.2
C		30-45	3.1 ± 0.3	9.5	3.0
D		50-60	2.9 ± 0.3	7.8	2.8
S8	2+50, 30R	0-15	1.6 ± 0.1	14	1.6
S9A	2+25, 32L	0-15	3.6 ± 0.1	21	3.2
B		15-30	3.4 ± 0.1	9.0	3.2
S10	2+25, 32R	0-15	2.8 ± 0.2	25	2.6
S11A	1+75, 18L	0-15	1.7 ± 0.4	30	1.5
B		15-30	1.6 ± 0.1	19	1.4
C		30-45	1.2 ± 0.08	<3.6	1.0
D		45-60	1.3 ± 0.2	<3.2	1.1
S12A	1+75, 18R	0-15	1.9 ± 0.1	38	1.8
B		15-30	1.4 ± 0.3	7.2	1.3
C		30-45	1.1 ± 0.2	2.9	1.0
D		45-60	1.1 ± 0.2	<3.2	0.99
S13	1+25, 18L	0-10	1.4 ± 0.2	31	1.4
S14A	1+25, 18R	0-15	1.4 ± 0.2	16	1.3
B		15-30	1.5 ± 0.4	20	1.4
S15A	0+75, 18L	0-15	1.9 ± 0.4	30	1.9
B	0+75, 18L	15-30	1.5 ± 0.3	16	1.4
C	0+75, 18L	30-45	1.1 ± 0.2	<5.0	1.0
D	0+75, 18L	45-60	1.2 ± 0.2	<3.2	1.0
S16A	0+75, 18R	0-15	1.7 ± 0.05	23	1.5
B		15-30	1.4 ± 0.1	18	1.4
C		30-45	1.2 ± 0.03	7.3	1.2
D		45-60	1.1 ± 0.2	3.8	1.1

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
S17	Creek bed	0–15	1.2 ± 0.1	4.1	1.0
S18	Creek bed	0–15	1.3 ± 0.1	4.3	1.1
S19	Creek bed	0–15	1.2 ± 0.1	4.4	1.1
S20A	3+80, 35R	0–15	3.2 ± 0.4	98	5.3
B		15–30	2.2 ± 0.4	39	3.8
S21A	4+50, 15L	0–15	1.1 ± 0.2	120	4.0
B		15–30	3.0 ± 0.1	55	3.3
S22A	4+31, BL	0–15	3.1 ± 0.3	110	2.9
B		15–30	5.1 ± 0.3	96	15
C		30–45	1.6 ± 0.3	13	12
D		45–60	1.4 ± 0.08	4.1	8.9
S23	16+50, 50L	0–15	1.3 ± 0.07	11	1.3
S24	21+50, 85L	0–15	1.3 ± 0.3	1.9	1.0
S25A	25+50, 50L	0–15	1.3 ± 0.2	2.3	1.1
B		15–30	1.2 ± 0.2	2.6	1.1
C		30–35	1.2 ± 0.2	2.1	1.1
D		45–50	1.2 ± 0.3	1.6	1.2
S26	27+50, 85L	0–15	1.4 ± 0.3	2.3	1.0
S27	29+00, 85L	0–15	1.2 ± 0.2	7.4	1.1
S28	28+50, 12R	0–8	2.3 ± 0.4	82	1.4
S29	24+50, 36R	0–8	1.3 ± 0.2	20	1.0
S30	20+50, 42R	0–15	1.2 ± 0.1	6.3	1.0
S31	15+50, 40R	0–15	1.3 ± 0.2	17	1.1
S32	10+00, 29R	0–15	1.3 ± 0.1	11	1.2
S33	7+50, 28R	0–15	1.5 ± 0.04	27	1.4
S34	6+39, 22R	0–15	2.2 ± 0.1	62	4.0
S35	5+25, 13R	0–15	7.1 ± 0.4	410	5.4
S36A	5+50, 47L	0–15	4.7 ± 0.4	230	3.0
B		45–60	1.6 ± 0.3	29	1.5
S37A	6+00, 50L	0–15	3.6 ± 0.2	170	2.1
B		45–60	1.2 ± 0.1	6.7	1.3
S38A	3+80, 35R	0–15	2.7 ± 0.07	77	2.5
B		15–30	1.7 ± 0.1	20	1.6
C		50–60	1.6 ± 0.2	81	1.2
S39A	4+50, 15L	0–15	4.6 ± 0.2	200	4.0
B		15–30	6.4 ± 0.6	280	6.1
C		30–45	4.0 ± 0.6	140	4.3
D		45–60	2.4 ± 0.03	68	3.3
E		60–75	2.1 ± 0.1	54	3.1

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
S40A	30+55, 60R	0-15	9.3±0.3	430	6.7
B		15-30	10 ±0.3	410	13
C		60-75	1.5±0.09	18	1.5
S41A	30+60, 112R	0-5	12 ±0.6	38	11
B		15-30	2.1±0.1	630	3.6
C		55-60	1.4±0.3	6.3	1.5
S42A	29+19, 08R	0-30	4.4±0.4	240	4.3
B		30-45	1.5±0.3	8.6	1.5
C		45-55	1.5±0.1	7.2	1.2
S43A	28+75, 10R	0-30	5.8±0.8	54	2.6
B		30-45	2.0±0.2	5.4	1.5
C		45-55	1.3±0.2	17	1.2
S44	27+15, 12R	0-15	1.0±0.2	23	1.0
S45	27+85, 11R	0-15	1.2±0.2	11	1.1
S46A	26+69, 10R	0-15	3.5±0.1	140	2.7
B		15-30	1.8±0.06	29	2.0
S47A	26+27, 12R	0-15	2.1±0.3	50	1.0
B		15-30	1.8±0.1	32	1.6
S48	13+67, 25R	0-15	4.5±0.3	5.4	1.2
S49	13+15, 25R	0-15	1.6±0.08	6.3	1.0
S50	11+55, 26R	0-15	2.7±0.1	35	1.6
S51	11+98, 26R	0-15	5.2±0.4	13	1.3
S52	9+06, 24R	0-15	1.9±0.2	20	1.8
S53	9+56, 24R	0-15	2.1±0.08	8.6	1.2
S54	14+38, 41R	0-15	1.3±0.1	<3.2	0.99
S55	15+12, 38R	0-15	4.9±0.2	<3.6	1.3
S56	30+65, 25R	0-15	1.8±0.2	28	1.4
S57	30+39, 28L	0-15	1.3±0.4	17	1.2
S58	29+60, 95L	0-15	2.9±0.4	<5.0	1.1
S59	29+38, 79L	0-15	3.2±0.2	<3.2	1.1
S60	23+70, 22L	0-15	3.8±0.2	<1.8	1.1
S61	23+38, 30R	0-15	1.7±0.2	<3.6	1.0
S62	23+27, 38R	0-15	2.2±0.1	11	1.2
S63	21+56, 44R	0-15	2.9±0.3	<2.7	1.1
<i>Biased samples^f</i>					
B1	30+60, 60R	0-15	8.7±0.9	500	8.7
B2A	13+60, 23R	0-5	8.3±1.0	350	5.4
B	13+60, 23R	5-10	12 ±0.3	810	8.6

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
<i>Auger samples^g</i>					
1	38+64, 05L	0–15	1.2 ±0.07	1.1	0.99
2	37+50, 19L	0–15	1.3 ±0.1	9.0	1.1
3	36+50, 17L	0–15	1.5 ±0.2	17	1.2
4	37+50, 16L	0–15	1.2 ±0.07	3.8	1.0
5	34+50, 15L	0–15	1.3 ±0.03	6.3	1.1
6	33+50, 17L	0–15	1.2 ±0.2	1.8	1.0
7	32+50, 18L	0–15	1.3 ±0.05	3.0	1.1
8	31+50, 17L	0–15	1.3 ±0.1	0.90	0.98
9	30+50, 19L	0–15	1.4 ±0.3	14	1.3
10A	28+50, 11L	0–15	0.81±0.03	1.7	0.39
B		15–30	1.4 ±0.1	0.59	1.1
11A	27+50, 08L	0–15	0.55±0.1	2.7	0.60
B		15–30	1.5 ±0.4	0.54	1.2
12A	26+50, 08L	0–15	0.66±0.1	0.54	0.71
B		15–30	1.2 ±0.1	6.3	1.1
13A	25+50, 07L	0–15	1.6 ±0.4	7.7	0.72
B		15–30	1.1 ±0.1	1.4	1.0
14A	24+50, 06L	0–15	0.94±0.06	8.1	0.96
B		15–30	1.3 ±0.08	1.2	1.1
15A	23+50, 05L	0–15	0.77±0.1	22	0.86
B		15–30	1.3 ±0.08	<0.90	1.2
16A	22+50, 04L	0–15	0.87±0.09	4.6	0.82
B		15–30	0.99±0.1	4.5	1.1
17A	21+50, 03L	0–15	0.83±0.08	2.5	0.91
B		15–30	0.97±0.2	1.2	1.1
18A	20+50, 03L	0–15	0.75±0.1	2.8	0.75
B		15–30	1.0 ±0.1	1.3	1.2
19A	19+50, 01L	0–15	0.87±0.1	1.5	0.97
B		15–30	1.6 ±0.2	1.9	1.5
20A	18+50, 01L	0–15	0.86±0.08	1.6	0.96
B		15–30	1.1 ±1	1.2	1.2
21A	17+47, BL	0–15	0.94±0.1	8.2	1.0
B		15–30	1.1 ±0.3	1.2	1.1
22A	16+49, BL	0–15	0.72±0.07	1.5	0.77
B		15–30	1.2 ±0.1	1.2	1.1
23A	15+48, 01L	0–15	0.76±0.06	3.7	0.84
B		15–30	1.5 ±0.4	1.3	1.4
24A	14+48, BL	0–15	0.83±0.1	8.8	0.90
B		15–30	1.2 ±0.2	1.8	1.2
C		30–60	1.1 ±0.1	1.3	1.1

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
25A	13+48, 01L	0-15	0.72±0.2	1.2	0.69
B		15-30	1.0 ±0.1	1.2	1.1
26A	12+55, BL	0-15	0.58±0.07	1.0	0.60
B		15-30	1.4 ±0.2	1.7	1.4
27A	11+47, BL	0-15	0.62±0.05	2.0	0.67
B		15-30	1.2 ±0.2	1.5	1.2
28A	10+50, 03L	0-15	0.60±0.07	2.7	0.64
B		15-30	1.2 ±0.1	1.2	1.2
29A	9+56, 03L	0-15	1.6 ±0.3	24	1.3
B		15-30	1.1 ±0.1	1.1	1.0
30A	8+47, 05L	0-15	0.75±0.08	3.3	0.80
B		15-30	1.1 ±0.1	0.72	1.0
31A	7+50, 02L	0-15	0.69±0.2	3.7	0.65
B		15-30	1.2 ±0.09	1.4	1.1
32A	6+50, 05L	0-15	0.56±0.2	6.3	0.59
B		15-30	1.0 ±0.09	1.6	1.0
33A	5+50, 07L	0-15	0.32±0.1	1.4	0.43
B		15-30	0.98±0.09	1.2	1.0
34A	11+00, 17R	0-15	2.7 ±0.1	93	2.7
B		15-30	1.3 ±0.3	9.4	1.6
35A	12+18, 17R	0-15	2.1 ±0.2	94	1.7
B		15-30	1.5 ±0.2	7.4	2.2
C		30-45	1.5 ±0.3	25	1.4
36A	12+92, 16R	0-15	1.7 ±0.1	16	1.8
B		15-30	1.2 ±0.09	24	0.96
37A	14+50, 22R	0-15	2.9 ±0.05	130	3.0
B		15-30	1.2 ±0.3	5.6	1.3
38A	15+35, 25R	0-15	2.3 ±0.2	60	2.1
B		15-30	1.1 ±0.3	0.77	1.2
39A	15+38, 13R	0-15	12 ±0.8	500	10
B		15-30	1.7 ±0.4	24	3.3
C		30-45	1.3 ±0.1	2.1	2.7
D		45-60	1.3 ±0.2	1.2	1.3
E		90-105	1.5 ±0.2	1.4	1.0
40A	16+00, 24R	0-15	1.9 ±0.08	37	1.8
B		15-30	2.0 ±0.2	1.2	2.2
41A	16+00, 15R	0-15	5.9 ±0.3	260	6.3
B		15-30	1.2 ±0.3	4.5	1.8
42A	17+00, 23R	0-15	1.9 ±0.1	37	1.7
B		15-30	0.98±0.2	3.2	1.1

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
43A	17+03, 15R	0–15	2.4 ±0.1	41	6.9
B		15–30	1.2 ±0.2	4.5	3.0
C		30–45	1.1 ±0.1	0.77	1.1
D		45–60	1.1 ±0.2	0.90	1.1
44A	18+01, 23R	0–15	1.6 ±0.3	31	1.6
B		15–30	1.2 ±0.09	6.8	1.3
45A	18+01, 15R	0–15	2.6 ±0.5	54	2.3
B		15–30	1.3 ±0.2	8.1	1.5
46A	19+00, 23R	0–15	1.7 ±0.4	23	1.5
B		15–30	1.3 ±0.1	7.2	1.3
47A	19+00, 13R	0–15	19 ±1	1,400	18
B		15–30	1.3 ±0.1	4.5	5.3
C		30–45	1.2 ±0.2	6.8	5.2
48A	20+23, 23R	0–15	2.1 ±0.2	54	1.9
B		15–30	1.6 ±0.2	8.2	1.3
49A	20+23, 16R	0–15	3.6 ±0.2	140	3.2
B		15–30	1.2 ±0.3	6.8	1.8
50A	20+23, 13R	0–15	15 ±1	1,100	16
B		15–30	1.3 ±0.2	4.5	6.3
C		30–45	1.1 ±0.1	3.6	3.4
51A	20+23, 11R	0–15	7.6 ±0.2	620	8.2
B		15–30	1.4 ±0.5	18	2.0
C		30–45	1.3 ±0.3	3.8	1.4
52A	21+00, 25R	0–15	1.7 ±0.1	36	1.8
B		15–30	1.8 ±0.1	30	1.7
53A	21+00, 13R	0–15	10 ±3	550	8.4
B		15–30	1.2 ±0.2	5.0	3.7
C		30–45	1.3 ±0.3	1.4	1.7
54A	22+50, 25R	0–15	1.6 ±0.1	14	1.3
55A	22+50, 12R	0–15	3.3 ±0.08	110	2.9
B		15–30	4.9 ±0.7	210	4.6
C		30–45	1.2 ±0.2	63	2.9
56A	24+00, 08R	0–15	1.7 ±0.1	50	1.9
B		15–30	7.1 ±0.5	660	9.8
C		30–45	5.7 ±1	270	5.8
D		45–60	1.6 ±0.3	19	1.5
57A	25+00, 13R	0–15	0.79 ±0.3	18	1.3
B		45–60	4.3 ±0.6	190	2.5
58A	25+00, 11R	0–15	6.5 ±0.3	360	4.6
B		15–30	1.7 ±0.1	45	1.2

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
59A	25+00, 07R	0–15	5.0 ±0.6	210	4.0
B		15–30	1.3 ±0.1	6.0	1.2
60	26+00, 13R	0–15	3.9 ±0.1	120	3.8
61	27+13, 07R	0–15	1.9 ±0.5	67	2.2
62A	28+00, 06R	0–15	4.0 ±0.3	150	3.1
B		15–30	2.5 ±0.4	69	3.7
63A	29+21, 03R	0–15	7.7 ±0.6	380	4.0
B		15–30	7.9 ±0.8	300	8.3
C		60–75	1.5 ±0.3	9.0	2.7
64A	30+35, 30R	0–15	5.8 ±0.4	390	2.6
B		15–30	1.4 ±0.1	9.5	2.0
65	31+00, 15R	0–15	2.0 ±0.2	53	1.9
66A	32+25, 15R	0–15	28 ±1	1,600	33
B		45–60	1.2 ±0.2	4.9	1.0
67	32+96, 15R	0–15	1.4 ±0.1	14	1.3
68	34+00, 15R	0–15	1.6 ±0.4	14	1.2
69	35+00, 15R	0–15	1.7 ±0.05	13	1.4
70	36+00, 14R	0–15	1.4 ±0.2	12	1.2
71	37+00, 14R	0–15	1.6 ±0.2	25	1.4
72	38+00, 15R	0–15	1.3 ±0.08	5.4	1.1
73A	29+37, 32L	0–15	0.49±0.09	1.1	0.52
B		15–30	29 ±0.7	2,000	25
C		30–45	1.4 ±0.2	5.4	1.3
D		45–60	1.1 ±0.2	6.8	1.2
E		60–75	1.2 ±0.2	3.4	0.38
74A	29+35, 38L	0–15	1.9 ±0.9	23	1.5
B		15–30	1.4 ±0.2	5.4	1.2
75	28+00, 19L	0–15	1.2 ±	8.1	1.1
76	27+00, 19L	0–15	1.3 ±0.09	3.6	1.2
77A	26+00, 21L	0–15	2.1 ±0.3	54	1.4
B		15–30	2.2 ±0.1	54	1.5
C		30–45	2.2 ±0.05	43	1.5
78	25+00, 18L	0–15	1.4 ±0.08	12	1.3
79A	24+00, 16L	0–15	5.4 ±0.5	260	7.2
B		45–60	1.3 ±0.3	9.5	1.8
80	23+00, 16L	0–15	1.8 ±0.1	23	1.8
81A	21+98, 18L	0–15	8.7 ±0.3	230	4.4
B		15–30	6.2 ±0.3	330	8.0
C		60–75	1.4 ±0.03	3.6	3.5

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
82A	21+98, 20L	0-15	2.6 ±0.4	190	3.6
B		45-60	1.3 ±0.2	1.8	1.4
83A	20+93, 15L	0-15	24 ±0.8	1,200	19
B		30-45	1.6 ±0.4	28	5.2
C		60-75	1.4 ±0.3	1.7	3.4
84	20+00, 12L	0-15	1.4 ±0.2	6.8	1.5
85A	19+00, 12L	0-15	17 ±0.7	1,000	10
B		75-90	1.6 ±0.3	19	2.8
86A	18+00, 13L	0-15	4.4 ±0.3	150	7.3
B		90-105	1.5 ±0.2	4.5	1.2
87	18+00, 16L	0-15	2.5 ±0.2	230	2.0
88A	17+00, 14L	0-15	6.3 ±0.3	290	6.6
B		60-75	1.4 ±0.1	8.6	1.7
89A	16+00, 13L	0-15	43 ±0.9	1,600	41
B		75-90	1.5 ±0.07	8.6	1.4
90A	16+00, 16L	15-30	1.5 ±0.09	5.9	1.7
B		60-75	1.5 ±0.3	2.3	1.0
91	32+24, 20R	0-15	1.7 ±0.3	22	1.6
92A	29+40, 26L	0-15	0.56±0.05	34	0.62
B		15-30	0.75±0.1	5.0	0.74
93A	29+21, 05L	0-15	1.0 ±0.06	7.5	1.1
B		15-30	1.1 ±0.2	2.8	1.1
94	29+21, 09R	0-15	3.7 ±0.6	180	2.8
95A	24+50, 02R	0-15	1.2 ±0.2	3.1	1.4
B		15-30	1.1 ±0.07	2.6	1.1
96A	24+50, 10R	0-15	3.9 ±0.3	140	2.4
B		15-30	5.2 ±0.3	160	4.9
C		75-90	1.3 ±0.3	2.2	1.6
97	24+50, 15R	0-15	6.9 ±0.1	250	3.9
98A	24+00, 01R	0-15	0.93±0.2	25	1.1
B		15-30	1.5 ±0.07	1.6	1.4
99	24+00, 14R	0-15	4.6 ±0.3	140	2.6
100A	15+99, 10L	0-15	1.4 ±0.3	31	1.1
B		15-30	1.2 ±0.2	3.0	1.1
101A	14+94, 14L	0-15	3.8 ±0.4	140	4.6
B		15-30	1.4 ±0.09	2.7	2.0
C		45-60	1.5 ±0.3	1.4	1.1
102A	13+80, 20L	0-15	1.1 ±0.1	24	1.1
B		15-30	0.99±0.1	4.5	1.1

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
103A	12+93, 19L	0–15	3.9 ± 0.3	160	3.1
B		15–30	1.2 ± 0.3	13	1.2
C		60–75	1.1 ± 0.2	0.68	1.2
104A	11+65, 15L	0–15	0.77 ± 0.2	13	0.67
B		15–30	1.2 ± 0.08	10	1.2
105A	10+98, 16L	0–15	5.0 ± 0.4	240	4.2
B		15–30	1.3 ± 0.08	5.0	1.4
106A	10+98, 21L	0–15	1.7 ± 0.07	40	1.7
B		15–30	1.3 ± 0.1	6.8	1.3
107A	10+98, 12L	0–15	0.42 ± 0.05	1.5	0.43
B		15–30	1.1 ± 0.2	1.5	0.96
C		60–75	1.2 ± 0.3	0.81	1.1
108A	9+86, 15L	0–15	0.76 ± 0.2	42	0.64
B		15–30	1.9 ± 0.2	22	4.7
C		30–45	1.5 ± 0.1	3.2	1.6
109A	9+86, 20L	0–15	2.6 ± 0.4	21	2.7
B		15–30	2.1 ± 0.2	38	2.0
110A	9+86, 11L	0–15	0.96 ± 0.2	3.3	0.73
B		15–30	1.2 ± 0.2	2.9	1.1
111A	9+00, 11L	0–15	5.2 ± 0.2	250	6.2
B		15–30	1.3 ± 0.2	7.2	3.7
112A	No sample				
B	No sample				
C	8+00, 19L	30–40	1.1 ± 0.2	1.5	1.0
113A	No sample				
B	No sample				
114A	8+00, 13L	0–15	0.79 ± 0.09	4.5	0.81
B		15–30	1.9 ± 0.3	15	1.4
115A	7+00, 19L	0–15	0.62 ± 0.03	4.2	0.60
B		15–30	0.84 ± 0.1	2.1	0.87
116A	6+02, 17L	0–15	13 ± 0.59	600	10
B		15–30	8.5 ± 0.5	410	7.3
C		30–45	5.7 ± 0.3	330	5.0
D		45–60	7.1 ± 0.5	450	5.9
E		60–75	2.0 ± 0.3	63	2.0
117A	6+02, 28L	0–15	11 ± 1	500	2.6
B		15–30	7.2 ± 0.2	330	8.2
C		45–60	4.5 ± 0.4	200	5.9
D		60–75	7.9 ± 0.5	400	8.4

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
118A	6+02, 13L	0-15	0.41 ± 0.08	3.7	2.7
B		15-30	2.6 ± 0.3	98	0.41
119A	5+50, 20L	0-15	19 (+-2)	900	20
B		15-30	4.4 ± 0.2	120	5.3
C		30-45	2.2 ± 0.1	84	2.8
D		45-60	1.4 ± 0.3	20	1.4
E		60-75	4.5 ± 0.6	340	3.3
120A	5+56, 09R	0-15	3.3 ± 0.2	200	4.2
B		15-30	16 ± 1	800	16
C		30-45	56 ± 4	3,000	54
D		45-60	14 ± 0.7	570	11
E		60-75	11 ± 0.6	610	8.2
121A	6+00, 10R	0-15	37 ± 3	2,000	42
B		15-30	5.1 ± 0.3	250	6.3
C		45-60	1.7 ± 0.07	37	2.1
122A	7+00, 08R	0-15	7.2 ± 0.6	370	4.9
B		45-60	1.4 ± 0.4	20	1.9
123A	7+00, 16R	0-15	13 ± 0.7	790	11
B		45-60	1.3 ± 0.1	18	1.8
124A	7+00, 19R	0-15	14 ± 0.5	670	10
B		15-30	3.3 ± 0.08	120	4.7
C		60-75	1.5 ± 0.09	45	1.7
125A	7+00, 05R	0-15	1.4 ± 0.2	51	1.5
B		15-30	41 ± 2	2,400	41
C		30-45	1.4 ± 0.2	40	1.2
126A	7+00, 01R	0-15	0.94 ± 0.08	14	0.98
B		15-30	1.3 ± 0.2	32	1.7
127A	8+00, 11R	0-15	6.2 ± 0.4	330	6.9
B		15-30	6.8 ± 0.51	120	8.5
C		60-75	1.7 ± 0.3	22	2.0
128A	8+00, 20R	0-15	2.7 ± 0.1	66	2.3
B		30-45	1.2 ± 1	6.7	1.1
C		75-90	1.2 ± 0.1	3.6	1.2
129A	8+00, 06R	0-15	0.89 ± 0.3	6.8	0.96
B		15-30	4.6 ± 0.3	220	5.3
130A	9+00, 13R	0-15	5.6 ± 0.4	260	4.4
B		15-30	2.7 ± 0.1	93	3.3
C		60-75	1.2 ± 0.2	3.2	1.4
131A	9+85, 15R	0-15	3.0 ± 0.1	100	2.6
B		15-30	1.4 ± 0.2	4.9	2.0

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
132A	4+80, 29R	0–15	1.4 ±0.09	51	1.0
B		15–30	0.76±0.1	3.0	0.76
133A	5+04, 40R	0–15	0.67±0.07	5.1	0.90
B		15–30	0.99±0.08	0.50	0.91
134A	3+50, 38L	0–15	1.6 ±0.2	15	1.3
B		15–30	1.2 ±0.2	1.3	1.0
C		60–75	1.4 ±0.2	6.4	1.0
135A	3+30, BL	0–15	1.5 ±0.2	18	1.4
B		15–30	1.2 ±0.2	1.3	1.1
136A	3+15, 29R	0–15	1.4 ±0.2	14	1.3
B		15–30	1.2 ±0.2	4.0	1.1
C		60–75	1.5 ±0.6	2.9	1.1
137A	3+55, 30R	0–15	1.0 ±0.1	6.7	0.99
B		15–30	1.0 ±0.3	5.2	1.1
C		45–60	1.1 ±0.2	1.1	1.1
138A	3+58, 35R	0–15	1.4 ±0.2	2.4	1.1
B		15–30	1.2 ±0.07	4.3	1.0
139A	3+94, 04L	0–15	1.0 ±0.2	6.5	0.91
B		15–30	1.1 ±0.3	1.5	0.92
C		60–75	1.2 ±0.1	0.70	1.1
140A	4+28, 32L	0–15	1.1 ±0.2	5.8	0.99
B		15–30	1.0 ±0.06	9.9	0.99
C		105–120	1.2 ±0.1	8.5	1.0
141A	4+43, 16R	0–15	1.3 ±0.08	3.2	1.1
B		15–30	1.3 ±0.06	5.0	1.2
C		45–60	1.2 ±0.2	2.2	1.1
142A	4+25, 30R	0–15	1.6 ±0.3	19	1.5
B		15–30	1.3 ±0.1	6.3	1.2
C		45–60	1.2 ±0.2	0.99	1.1
143A	5+50, 64L	0–15	3.2 ±0.2	120	2.1
B		15–30	1.2 ±0.4	9.0	1.3
C		45–60	1.1 ±0.2	1.4	1.0
144A	6+00, 73L	0–15	1.7 ±0.2	34	1.4
B		15–30	1.2 ±0.1	8.6	1.4
C		45–60	1.2 ±0.1	1.7	1.0
145A	7+00, 30L	0–15	1.9 ±0.1	17	1.9
B		15–30	1.5 ±0.1	9.5	1.8
C		45–60	1.2 ±0.1	1.6	0.96
146A	8+77, 69R	0–15	1.6 ±0.3	14	1.2
B		60–75	1.1 ±0.2	1.2	0.90

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
147A	8+80, 135R	0–15	69 ± 3	6,100	80
B		75–90	1.0 ± 0.3	0.99	1.1
148A	8+75, 200R	0–15	660 ± 30	22,000	790
B		90–105	1.5 ± 0.3	23	2.6
149A	8+20, 210R	45–60	1.8 ± 0.4	14	1.2
B		95–100	2.3 ± 0.4	32	1.6
150A	8+24, 150R	0–15	35 ± 2	2,000	48
B		60–75	2.7 ± 0.1	53	3.3
151A	8+15, 100R	0–15	16 ± 0.6	1,000	24
B		45–60	1.3 ± 0.1	23	5.8
152A	8+13, 39R	0–15	3.3 ± 0.1	190	3.5
B		15–30	1.8 ± 0.3	50	2.8
153A	7+64, 52R	0–15	11 ± 0.8	550	13
B		45–60	1.1 ± 0.2	5.0	1.8
154A	7+70, 111R	90–105	1.2 ± 0.2	6.8	2.7
155A	7+75, 210R	60–75	1.6 ± 0.2	11	1.3
B		90–95	2.8 ± 0.5	57	2.4
156A	7+25, 210R	60–90	1.9 ± 0.2	25	1.7
B		90–95	1.8 ± 0.2	30	1.6
157A	7+36, 94R	75–90	1.2 ± 0.3	14	1.8
158A	7+20, 40R	0–15	200 ± 7	10,000	240
B		60–75	2.2 ± 0.07	67	3.6
159A	6+50, 50R	75–90	1.4 ± 0.3	65	3.0
160A	5+88, 41R	0–15	300 ± 10	16,000	240
B		90–95	1.4 ± 0.3	12	2.9
161A	2+88, BL	45–60	1.2 ± 0.2	1.1	1.1
B		120–125	1.2 ± 0.05	2.7	1.1
162A	2+68, 10L	0–15	2.0 ± 0.2	32	2.0
B		60–75	1.2 ± 0.1	1.7	1.1
C		120–125	1.2 ± 0.2	3.6	1.2
163A	2+42, BL	30–45	1.3 ± 0.1	3.8	1.3
B		120–125	1.2 ± 0.3	5.0	1.4
164A	(Hole designation not used)		—	—	—
165A	2+29, BL	30–45	1.4 ± 0.3	220	1.4
B		120–125	1.2 ± 0.3	0.86	1.2
166A	2+00, 33L	0–15	1.7 ± 0.09	21	1.5
B		105–110	1.1 ± 0.2	1.3	0.98
167A	2+00, 33R	0–15	1.5 ± 0.1	15	1.4
B		105–110	1.2 ± 0.3	20	0.95
168A	1+49, BL	0–15	1.8 ± 0.09	32	1.7
B		105–110	1.0 ± 0.2	1.3	0.90

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
169A	1+00, 33R	0–15	1.7 ± 0.1	21	1.6
B		120–125	1.1 ± 0.1	27	1.0
170A	1+00, 30L	0–15	1.6 ± 0.3	15	1.5
B		105–110	1.2 ± 0.09	6.7	0.98
171A	0+40, BL	0–15	1.1 ± 0.1	5.8	0.95
B		120–125	1.0 ± 0.06	3.1	0.93
172A	5+50, 38L	0–15	2.6 ± 0.2	86	2.2
B		45–60	1.1 ± 0.2	7.0	1.3
C		115–120	1.1 ± 0.1	5.2	0.95
173A	6+00, 38L	0–15	2.3 ± 0.1	49	2.1
B		45–60	1.0 ± 0.3	9.0	1.2
C		60–75	1.4 ± 0.2	18	1.2
174A	20+50, 02R	30–60	1.2 ± 0.07	<2.7	1.1
B		60–90	1.2 ± 0.2	<3.2	1.1
C		90–120	1.5 ± 1	<2.7	1.2
175A	25+50, 04L	30–60	1.1 ± 0.1	<1.1	1.0
B		60–90	1.2 ± 0.2	<3.2	1.0
C		105–120	1.2 ± 0.2	<2.7	0.97
176A	37+02, 14R	15–30	1.3 ± 0.1	<5.4	1.0
B		30–60	1.4 ± 0.2	<2.7	1.0
C		60–90	1.3 ± 0.09	<3.2	1.0
D		90–120	1.2 ± 0.05	<2.7	1.0
177A	30+93, 15R	15–30	1.2 ± 0.1	<5.4	1.3
B		30–60	1.2 ± 0.08	<4.0	0.97
C		60–90	1.1 ± 0.09	<1.8	1.0
D		90–120	1.2 ± 0.06	<3.2	0.96
178A	28+02, 06R	30–60	1.7 ± 0.2	28	1.6
B		60–90	1.3 ± 0.1	<3.6	1.0
C		90–120	1.1 ± 0.2	<4.5	1.1
179A	26+02, 21L	45–60	1.5 ± 0.08	<1.4	1.1
B		60–90	1.5 ± 0.06	<1.8	1.0
C		90–120	1.6 ± 0.2	<4.5	1.1
180A	29+21, 12R	15–30	1.9 ± 0.2	4.0	1.9
B		30–60	1.5 ± 0.2	2.7	1.1
180C		60–90	1.3 ± 0.2	2.9	0.97
D		90–120	1.4 ± 0.2	28	1.0
181A	27+10, 07R	15–30	1.2 ± 0.2	3.7	1.8
B		30–60	1.3 ± 0.1	1.8	1.1
C		60–90	1.5 ± 0.1	2.0	0.97
D		90–120	1.2 ± 0.2	2.1	0.93

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
182A	25+97, 07R	15–30	1.6 ± 0.25	18	2.3
B		30–60	1.3 ± 0.2	3.4	1.1
C		60–90	1.4 ± 0.06	2.5	0.94
D		90–120	1.6 ± 0.05	2.6	0.95
183A	25+04, 13R	60–90	1.4 ± 0.08	2.4	1.7
B		90–120	1.6 ± 0.09	2.1	1.4
184A	24+47, 15R	15–30	1.4 ± 0.2	11	1.4
B		30–60	1.6 ± 0.2	19	1.4
C		60–90	1.2 ± 0.1	2.1	1.1
D		90–120	1.2 ± 0.09	2.6	0.97
185A	23+98, 14R	15–30	1.2 ± 0.2	6.8	1.1
B		30–60	1.5 ± 0.2	15	1.5
C		60–90	1.2 ± 0.08	2.9	0.95
D		90–120	1.3 ± 0.16	2.4	0.95
186A	22+25, 12R	45–60	1.2 ± 0.2	2.1	1.1
B		60–90	1.2 ± 0.1	2.5	1.0
C		90–120	1.4 ± 0.04	3.7	1.0
187A	20+98, 25R	30–60	1.3 ± 0.2	6.3	1.3
B		60–90	1.3 ± 0.2	2.5	1.2
C		90–120	1.4 ± 0.2	2.4	1.2
188A	18+94, 12L	30–60	1.4 ± 0.2	6.2	4.8
B		60–90	1.3 ± 0.2	3.1	3.0
C		90–120	1.4 ± 0.2	3.2	1.4
189A	17+96, 13L	15–30	9.2 ± 0.8	360	6.3
B		30–60	1.2 ± 0.3	4.4	5.9
C		60–90	1.3 ± 0.3	1.7	2.9
190A	12+24, 17R	15–30	1.2 ± 0.2	3.8	1.5
B		30–60	1.5 ± 0.1	1.6	1.3
C		60–90	1.4 ± 0.2	1.9	1.1
191A	12+60, 32L	15–30	1.1 ± 0.2	1.7	1.1
B		30–60	1.2 ± 0.2	2.2	1.1
C		60–90	1.4 ± 0.05	2.1	1.2
D		90–120	1.4 ± 0.2	2.0	1.2
192A	7+96, 19L	0–30	14 ± 0.3	870	10
B		30–60	1.1 ± 0.07	2.7	1.0
C		60–90	1.0 ± 0.3	<1.5	1.0
D		90–120	1.2 ± 0.1	2.1	1.0
193A	30+65, 19R	0–30	1.3 ± 0.1	7.6	1.2
B		30–60	1.1 ± 0.3	0.90	0.99
C		60–90	1.1 ± 0.06	8.1	0.97
D		90–120	1.3 ± 0.08	9.8	0.91

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
194A	30+39, 19L	0–30	1.2 ± 0.2	0.80	1.0
B		30–60	1.1 ± 0.09	0.70	1.0
C		60–90	1.0 ± 0.09	1.0	1.0
D		90–120	1.2 ± 0.2	0.50	1.0
195A	29+09, 10R	0–30	2.7 ± 0.1	53	1.6
B		30–60	2.4 ± 0.3	53	1.8
196A	27+18, 14R	0–30	2.0 ± 0.09	23	1.4
B		30–60	2.4 ± 0.1	29	2.5
197A	26+32, 13R	0–30	2.4 ± 0.2	40	2.8
B		30–60	2.0 ± 0.07	27	2.0
C		60–90	1.3 ± 0.2	2.1	1.1
198A	23+66, 17L	0–30	11 ± 0.2	400	6.2
B		30–60	3.4 ± 0.2	59	3.6
C		60–90	0.91 ± 0.1	3.9	1.1
D		90–120	1.1 ± 0.1	7.4	1.1
199A	23+09, 37R	0–30	1.3 ± 0.2	2.0	1.0
B		30–60	1.2 ± 0.08	1.3	0.94
C		60–90	1.2 ± 0.4	1.3	0.93
D		90–120	1.2 ± 0.1	1.4	0.88
E		120–150	1.5 ± 0.07	2.7	1.1
F		150–185	1.3 ± 0.2	4.1	1.0
G		185–215	1.1 ± 0.2	<3.2	0.92
H		215–245	0.98 ± 0.1	<3.2	0.85
200A	22+40, 45R	0–30	1.2 ± 0.2	4.4	1.0
B		30–60	1.3 ± 0.2	7.9	1.2
C		60–90	1.2 ± 0.3	2.0	1.0
D		90–120	1.1 ± 0.2	2.1	1.0
E		120–150	1.1 ± 0.2	<3.2	1.0
F		150–185	1.2 ± 0.2	<5.4	1.1
G		185–215	1.3 ± 0.09	1.3	1.0
H		215–245	1.1 ± 0.1	0.70	0.91
201A	21+81, 38R	0–30	1.4 ± 0.4	4.6	1.3
B		30–60	1.2 ± 0.2	1.4	1.0
C		60–90	1.1 ± 0.04	1.8	1.0
D		90–120	1.3 ± 0.2	0.80	1.0
E		120–150	1.2 ± 0.2	1.5	1.0
F		150–185	1.2 ± 0.3	1.0	1.0
G		185–225	1.1 ± 0.1	0.70	0.95

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
202A	13+59, 27R	0-30	0.83±0.08	3.3	1.1
B		30-60	1.0 ±0.1	1.3	1.1
C		60-90	1.0 ±0.08	1.3	1.0
D		90-120	0.98±0.07	1.8	1.0
E		165-185	0.98±0.3	1.9	0.93
203A	13+30, 28R	5-30	0.96±0.09	0.80	1.0
B		30-60	0.87±0.1	0.70	0.94
C		60-90	0.93±0.1	0.90	0.91
D		90-120	1.0 ±0.2	2.2	0.98
E		120-165	1.2 ±0.3	1.3	0.93
F		165-185	1.0 ±0.1	0.80	0.95
204A	11+77, 24R	5-30	0.95±0.05	0.60	0.93
B		30-60	0.92±0.08	0.60	0.93
C		60-90	1.1 ±0.09	0.70	1.1
D		90-120	1.3 ±0.1	1.0	1.3
E		120-145	1.2 ±0.2	0.80	1.1
F		145-165	1.4 ±0.2	1.3	1.1
205A	9+48, 22R	5-30	0.97±0.1	<2.3	1.0
B		30-60	0.98±0.09	<1.8	1.1
C		60-90	1.1 ±0.05	<3.6	0.99
D		90-120	1.2 ±0.4	<3.2	1.1
E		120-145	1.3 ±0.1	<2.3	0.97
F		145-165	1.4 ±0.3	<3.6	0.97
206A	9+20, 28R	5-30	1.1 ±0.3	<3.6	1.1
B		30-60	0.96±0.3	<3.2	1.0
C		60-90	1.1 ±0.05	9.0	1.1
D		90-120	1.1 ±0.09	<4.5	0.97
E		120-145	1.1 ±0.2	<3.6	1.0
F		145-165	1.1 ±0.2	<2.3	0.92
207A	9+22, 16R	5-30	0.97±0.08	<0.90	0.99
B		30-60	1.1 ±0.1	<2.3	1.1
C		60-90	1.1 ±0.2	<2.3	0.94
D		90-120	1.0 ±0.1	<3.2	1.0
208A	9+88, 20L	30-60	1.2 ±0.08	<1.5	1.6
B		60-90	1.4 ±0.09	2.1	0.96
C		90-120	1.5 ±0.3	1.8	0.96
209A	7+02, 16R	60-90	1.1 ±0.1	4.1	1.7
B		90-120	1.3 ±0.2	2.4	1.5
C		120-150	1.3 ±0.3	10	1.8
D		150-185	1.3 ±1	6.6	1.4

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
210A	6+00, 28L	60–90	1.4 ± 0.2	11	1.4
B		90–120	1.3 ± 0.2	<2.3	0.94
211A	5+58, 09R	60–90	5.4 ± 0.08	230	4.5
B		90–120	1.2 ± 0.1	14	1.2
C		120–150	1.1 ± 0.2	2.8	1.3
D		150–165	1.5 ± 0.2	33	1.3
212A	5+52, 20L	60–90	10 ± 0.5	450	11
B		90–120	1.1 ± 0.3	<2.7	1.1
C		120–150	1.2 ± 0.07	2.4	0.96
D		150–185	4.7 ± 0.2	200	5.6
E		185–215	1.3 ± 0.1	3.1	0.96
F		215–245	1.3 ± 0.3	11	1.2
213A	14+48, 35R	0–30	1.8 ± 0.2	44	1.8
B		30–60	0.77 ± 0.1	<5.0	0.77
C		60–90	0.85 ± 0.2	<3.6	0.79
D		90–120	0.98 ± 0.6	<5.4	0.80
E		120–145	1.5 ± 0.1	<5.0	1.0
F		145–165	1.2 ± 0.06	<3.2	0.92
214A	5+50, 50L	60–90	1.2 ± 0.2	<0.90	1.1
B		90–120	1.1 ± 0.3	<3.2	1.1
C		120–150	1.2 ± 0.3	3.9	0.89
215A	4+50, 106L	0–30	1.5 ± 0.1	19	1.5
B		30–60	1.1 ± 0.2	<2.7	1.0
C		60–90	1.1 ± 0.3	<3.6	0.99
D		90–120	1.2 ± 0.2	<3.2	0.92
E		120–145	1.4 ± 0.4	<3.2	0.81
F		145–165	1.3 ± 0.2	<3.6	0.76
216A	7+90, 210R	0–30	55 ± 3	2,700	2.4
B		30–60	1.4 ± 0.1	<4.1	0.80
C		60–75	1.3 ± 0.4	<5.4	1.0
D		75–90	1.5 ± 0.2	<5.4	1.0
217A	7+39, 96R	0–30	4.5 ± 0.4	150	5.5
B		30–60	28 ± 2	1,300	24
C		60–75	1.4 ± 0.1	15	3.6
D		60–90	2.1 ± 0.3	86	3.7
E		90–120	1.9 ± 0.2	<4.1	1.8
218A	7+10, 50R	0–30	16 ± 1	520	19
B		30–45	2.6 ± 0.2	49	3.9
C		60–90	1.3 ± 0.08	13	2.3
D		90–120	1.4 ± 0.2	16	1.8

Table 4 (continued)

Sample/ hole number	Location ^a	Depth (cm)	Radionuclide concentration (pCi/g)		
			²²⁶ Ra ^b	²³⁰ Th ^c	²³⁸ U ^d
219A	5+89, 50R	60–90	2.1 ± 0.3	<5.4	6.0
B		90–120	1.7 ± 0.1	<2.3	2.9
220A	8+11, 43R	0–30	3.6 ± 0.2	110	3.1
B		30–60	1.2 ± 0.06	4.7	1.4
C		60–75	1.1 ± 0.2	8.0	1.0
D		75–95	1.1 ± 0.2	<2.3	0.99
E		95–115	1.3 ± 0.4	<3.6	0.89
221A		0–30	82 ± 3	3,700	93
B	8+25, 150R	30–60	2.6 ± 0.4	35	7.0
C		70–105	1.8 ± 0.07	14	3.1
D		105–135	1.4 ± 0.1	17	2.0

^aLocations of soil samples are shown on Figs. 3–31.

^bIndicated counting error is at the 95% confidence level ($\pm 2\sigma$).

^cThe error of the reported radionuclide concentration is $\pm 5\%$ (95% confidence level).

^dTotal analytical error of measurement results is $\pm 3\%$ (95% confidence level).

^eSystematic samples are taken at grid locations irrespective of gamma exposure.

^fBiased samples were taken from areas shown to have elevated gamma exposure rates.

^gAuger samples are those taken from holes drilled to further define the depth and extent of contamination. Holes may be drilled in either contaminated or uncontaminated regions.

Table 5. Extent of subsurface contamination on the Latty Avenue site (LM001) as indicated by scintillation probe loggings and sample analyses

Hole	Location ^a	Depth of hole (m)	Estimated vertical extent of contaminated soil ^b (cm)	Region of maximum contamination (cm)	Radionuclide concentration in region of maximum contamination (pCi/g)		
					²²⁶ Ra	²³⁰ Th ^c	²³⁸ U ^d
1	38+64, 05L	2.7	None				
2	37+50, 19L	2.7	0-15	0-15	1.3 ±0.1	9.0	1.1
3	36+50, 17L	2.6	0-15	0-15	1.5 ±0.2	17	1.2
4	35+50, 16L	2.4	None				
5	34+50, 15L	2.6	0-15	0-15	1.3 ±0.05	6.3	1.1
6	33+50, 17L	2.6	None				
7	32+50, 18L	2.6	None				
8	31+50, 17L	2.7	None				
9	30+50, 19L	2.6	0-15	0-15	1.4 ±0.3	14	1.3
10	28+50, 11L	1.7	None				
11	27+50, 08L	1.7	None				
12	26+50, 08L	1.7	None				
13	25+50, 07L	1.7	0-15	0-15	1.6 ±0.4	7.7	0.72
14	24+50, 06L	1.7	0-15	0-15	0.94±0.06	8.1	0.96
15	23+50, 05L	1.7	0-15	0-15	0.77±0.1	22	0.86
16	22+50, 04L	1.7	None				
17	21+50, 03L	1.7	None				
18	20+50, 03L	1.7	None				
19	19+50, 01L	1.7	None				
20	18+50, 01L	1.7	None				
21	17+47, BL	1.2	0-15	0-15	0.94±0.1	8.2	1.0
22	16+49, BL	1.7	None				
23	15+48, 01L	1.7	None				
24	14+48, BL	2.6	0-15	0-15	0.83±0.1	8.8	0.90
25	13+48, 01L	1.7	None				
26	12+55, BL	1.7	None				
27	11+47, BL	1.1	None				
28	10+50, 03L	1.7	None				
29	9+56, 03L	1.7	0-15	0-15	1.6 ±0.3	24	1.3
30	8+47, 05L	1.7	None				
31	7+50, 02L	1.7	None				
32	6+50, 05L	1.7	0-15	0-15	0.56±0.2	6.3	0.59
33	5+50, 05L	1.7	None				
34	11+00, 17R	1.1	0-15	0-15	2.7 ±0.1	93	2.7
35	12+18, 17R	1.1	0-60	0-15	2.1 ±0.2	94	1.7
36	12+92, 16R	1.1	0-60	15-30	1.2 ±0.09	24	0.96
37	14+50, 22R	1.1	0-15	0-15	2.9 ±0.05	130	3.0
38	15+35, 25R	1.2	0-15	0-15	2.3 ±0.3	60	2.1
39	15+38, 13R	1.2	0-30	0-15	12 ±0.8	500	10
40	16+00, 24R	1.2	0-15	0-15	1.9 ±0.08	37	1.8
41	16+00, 15R	1.2	0-15	0-15	5.9 ±0.3	260	6.3
42	17+00, 23R	1.2	0-15	0-15	1.9 ±0.1	37	1.7
43	17+03, 15R	1.1	0-15	0-15	2.4 ±0.1	41	6.9
44	18+01, 23R	1.2	0-15	0-15	1.6 ±0.3	31	1.6
45	18+01, 15R	1.2	0-15	0-15	2.6 ±0.5	54	2.3

Table 5 (continued)

Hole	Location ^a	Depth of hole (m)	Estimated vertical extent of contaminated soil ^b (cm)	Region of maximum contamination (cm)	Radionuclide concentration in region of maximum contamination (pCi/g)		
					²²⁶ Ra	²³⁰ Th ^c	²³⁸ U ^d
46	19+00, 23R	1.2	0-15	0-15	1.7±0.4	23	1.5
47	19+00, 13R	1.2	0-15	0-15	19 ±1.0	1400	18
48	20+23, 23R	1.2	0-15	0-15	2.1±0.2	54	1.9
49	20+23, 16R	1.1	0-15	0-15	3.6±0.2	140	3.2
50	20+23, 13R	1.2	0-15	0-15	15 ±1.0	1100	16
51	20+23, 11R	0.8	0-30	0-15	7.6±0.2	620	8.2
52	21+00, 25R	1.2	0-30	0-15	1.7±0.1	36	1.8
53	21+00, 13R	1.1	0-15	0-15	10 ±3.0	550	8.4
54	22+50, 25R	1.2	0-15	0-15	1.6±0.1	14	1.3
55	22+50, 12R	1.2	0-45	15-30	4.9±0.7	210	4.6
56	24+00, 08R	1.1	0-75	15-30	7.1±0.5	660	9.8
57	25+00, 13R	1.2	0-90	45-60	4.3±0.6	190	2.5
58	25+00, 11R	1.1	0-60	0-15	6.5±0.3	360	4.6
59	25+00, 07R	1.1	0-15	0-15	5.0±0.6	210	4.0
60	26+00, 07R	0.9	0-90	0-15	3.9±0.1	120	3.8
61	27+13, 07R	1.5	0-75	0-15	1.9±0.5	67	2.2
62	28+00, 06R	1.1	0-60	0-15	4.0±0.3	150	3.1
63	29+21, 03R	1.1	0-60	0-15	7.7±0.6	380	4.0
64	30+35, 30R	0.6	0-15	0-15	5.8±0.4	390	2.6
65	31+00, 15R	1.2	0-15	0-15	2.0±0.2	53	1.9
66	32+25, 15R	1.2	0-45	0-15	28 ±1.0	1600	33
67	32+96, 15R	1.1	0-15	0-15	1.4±0.1	14	1.3
68	34+00, 15R	1.2	0-30	0-15	1.6±0.4	14	1.2
69	35+00, 15R	1.2	0-15	0-15	1.7±0.5	13	1.4
70	36+00, 14R	1.2	0-15	0-15	1.4±0.2	12	1.2
71	37+00, 14R	1.2	0-15	0-15	1.6±0.2	25	1.4
72	38+00, 15R	1.2	0-15	0-15	1.3±0.08	5.4	1.1
73	29+37, 32L	1.8	15-30	15-30	29 ±0.7	2000	25
74	29+35, 38L	1.2	0-15	0-15	1.9±0.9	23	1.5
75	28+00, 19L	1.2	0-15	0-15	1.2±0.2	8.1	1.1
76	27+00, 19L	1.1	None				
77	26+00, 21L	1.1	0-45	0-30	2.2±0.10	54	1.5
78	25+00, 18L	1.2	0-30	0-15	1.4±0.08	12	1.3
79	24+00, 16L	0.9	0-45	0-15	5.4±0.5	260	7.2
80	23+00, 16L	1.2	0-30	0-15	1.8±0.1	23	1.8
81	21+98, 18L	1.1	0-60	15-30	6.2±0.3	330	8.0
82	21+98, 20L	1.1	0-45	0-15	2.6±0.4	190	3.6
83	20+93, 15L	1.1	0-60	0-15	24 ±0.8	1,200	19
84	20+00, 12L	1.1	0-30	0-15	1.4±0.2	6.8	1.5
85	19+00, 12L	1.1	0-105	0-15	17 ±0.7	1,000	10
86	18+00, 13L	1.1	0-90	15-30	4.4±0.3	150	7.3
87	18+00, 16L	1.1	0-60	0-15	2.5±0.2	230	2.0
88	17+00, 14L	1.1	0-60	0-15	6.3±0.3	290	6.6
89	16+00, 13L	0.9	0-75	0-15	43 ±0.9	1,600	41
90	16+00, 16L	1.2	0-15	15-30	1.5±0.07	8.6	1.4
91	32+24, 20R	1.5	0-45	0-15	1.7±0.3	22	1.6

Table 5 (continued)

Hole	Location ^a	Depth of hole (m)	Estimated vertical extent of contaminated soil ^b (cm)	Region of maximum contamination (cm)	Radionuclide concentration in region of maximum contamination (pCi/g)		
					²²⁶ Ra	²³⁰ Th ^c	²³⁸ U ^d
92	29+40, 26L	1.2	0-15	0-15	0.56 ± 0.05	34	0.62
93	29+21, 05L	1.2	0-15	0-15	1.0 ± 0.06	7.5	1.1
94	29+21, 09R	0.9	0-60	0-15	3.7 ± 0.6	180	2.8
95	24+50, 02R	1.1	None				
96	24+50, 10R	1.2	0-75	30-60	Not sampled at point of max cont.		
97	24+50, 15R	1.2	0-30	0-15	6.9 ± 0.1	250	3.9
98	24+00, 01R	1.2	0-15	0-15	0.93 ± 0.2	25	1.1
99	24+00, 14R	1.2	0-75	0-15	4.6 ± 0.3	140	2.6
100	15+99, 10L	1.1	0-15	0-15	1.4 ± 0.3	31	1.1
101	14+94, 14L	1.1	0-15	0-15	3.8 ± 0.4	140	4.6
102	13+80, 20L	1.2	0-15	0-15	1.1 ± 0.1	24	1.1
103	12+93, 19L	1.2	0-15	0-15	3.9 ± 0.3	160	3.1
104	11+67, 15L	1.1	0-15	0-15	0.77 ± 0.2	13	0.67
105	10+98, 16L	1.1	0-15	0-15	5.0 ± 0.4	240	4.2
106	10+98, 21L	1.1	0-15	0-15	1.7 ± 0.07	40	1.7
107	10+98, 12L	1.1	None				
108	9+86, 15L	0.9	0-30	0-15	0.76 ± 0.2	42	0.64
109	9+86, 20L	1.1	0-30	15-30	2.1 ± 0.2	38	2.0
110	9+86, 11L	1.1	None				
111	9+00, 11L	0.9	0-15	0-15	5.2 ± 0.2	250	6.2
112	8+00, 19L	1.2	0-30	15-30	Not sampled		
113	8+00, 29L	1.2	0-30	15-30	Not sampled		
114	8+00, 13L	1.2	0-30	15-30	1.9 ± 0.3	15	1.4
115	7+00, 19L	1.2	None				
116	6+02, 17L	2.7	0-75	0-15	13 ± 0.6	600	10
117	6+02, 28L	1.2	0-90	0-15	11 ± 1.0	500	2.6
118	6+02, 13L	1.8	0-75	0-15	2.6 ± 0.3	98	0.41
119	5+50, 20L	1.8	0-180	0-15	19 ± 2.0	900	20
120	5+56, 09R	1.8	0-105	30-45	56 ± 4.0	3,000	54
121	6+00, 10R	1.8	0-90	0-15	37 ± 3.0	2,000	42
122	7+00, 08R	1.8	0-90	0-15	7.2 ± 0.6	370	4.9
123	7+00, 16R	1.8	0-90	0-15	13 ± 0.7	790	11
124	7+00, 19R	1.8	0-90	0-15	14 ± 0.5	670	10
125	7+00, 05R	1.8	0-90	15-30	41 ± 2.0	2,400	41
126	7+00, 01R	1.8	0-90	15-30	1.3 ± 0.2	32	1.7
127	8+00, 11R	1.5	0-90	0-15	6.2 ± 0.4	330	6.9
128	8+00, 20R	1.8	0-30	0-15	2.7 ± 0.1	66	2.3
129	8+00, 06R	1.8	0-90	15-30	4.6 ± 0.3	220	5.3
130	9+00, 13R	1.8	0-60	0-15	5.6 ± 0.4	260	4.4
131	9+85, 15R	1.5	0-15	0-15	3.0 ± 0.1	100	2.6
132	4+80, 29R	1.8	0-15	0-15	1.4 ± 0.09	51	1.0
133	5+04, 40R	1.8	0-15	0-15	0.67 ± 0.07	5.1	0.90
134	3+50, 38L	1.8	0-15	0-15	1.6 ± 0.2	15	1.3
135	3+30, BL	1.2	0-15	0-15	1.5 ± 0.2	18	1.4
136	3+15, 29R	1.2	0-15	0-15	1.4 ± 0.2	14	1.3
137	3+55, 30R	1.1	0-75	0-15	1.0 ± 0.1	6.7	0.99

Table 5 (continued)

Hole	Location ^a	Depth of hole (m)	Estimated vertical extent of contaminated soil ^b (cm)	Region of maximum contamination (cm)	Radionuclide concentration in region of maximum contamination (pCi/g)		
					²²⁶ Ra	²³⁰ Th ^c	²³⁸ U ^d
138	3+58, 35R	1.8	105–180		Not sampled		
139	3+94, 04L	1.8	0–15	0–15	1.0±0.2	6.5	0.91
140	4+28, 32L	3.0	0–15	0–15	1.1±0.2	5.8	0.99
141	4+43, 16R	1.8	None				
142	4+25, 30R	1.8	0–15	0–15	1.6±0.3	19	1.5
143	5+50, 64L	1.8	0–15	0–15	3.2±0.2	120	2.1
144	6+00, 73L	1.2	0–15	0–15	1.7±0.2	34	1.4
145	7+00, 30L	1.8	0–15	0–15	1.9±0.1	17	1.9
146	8+77, 69R	1.8	0–15	0–15	1.6±0.3	14	1.2
147	8+80, 135R	1.8	0–75	0–15	69 ± 3.0	6,100	80
148	8+75, 200R	1.8	0–105	0–15	660 ± 30	22,000	790
149	8+20, 210R	0.9	0–105	90–105	2.3±0.4	32	1.6
150	8+24, 150R	1.8	0–90	0–15	35 ± 2.0	2,000	48
151	8+15, 100R	1.8	0–105	0–15	16 ± 0.6	1,000	24
152	8+13, 39R	1.8	0–90	0–15	3.3±0.1	190	3.5
153	7+64, 52R	1.8	0–75	0–15	11 ± 0.8	550	13
154	7+70, 111R	1.8	0–90	30–45	Not sampled		
155	7+75, 210R	0.9	0–120	0–30	Not sampled		
156	7+25, 210R	0.9	0–105	15–30	Not sampled		
157	7+36, 94R	1.8	0–90	30–45	Not sampled		
158	7+20, 40R	1.8	0–90	0–15	200 ± 7	10,000	240
159	6+50, 50R	1.7	0–105	15–30	Not sampled		
160	5+88, 41R	0.9	0–90	0–15	300 ± 10	16,000	240
161	2+88, BL	1.2	None				
162	2+68, 10L	1.2	0–30	0–15	2 ± 0.2	32	2.0
163	2+42, BL	1.2	None				
165	2+29, BL	1.2	0–120	30–45	1.4±0.3	220	1.4
166	2+00, 33L	1.1	0–30	0–15	1.7±0.09	21	1.5
167	2+00, 33R	1.1	0–30	90–105	1.2±0.03	20	0.95
168	1+49, BL	1.1	0–30	0–15	1.8±0.09	32	1.7
169	1+00, 33R	1.2	0–30	120–135	1.1±0.1	27	1.0
170	1+00, 30L	1.1	0–30	0–15	1.6±0.3	15	1.5
171	0+40, BL	1.2	0–15	0–15	1.1±0.1	5.8	0.95
172	5+50, 38L	0.9	0–45	15–30	Not sampled		
173	6+00, 38L	0.9	0–75	0–15	2.3±0.1	49	2.1
174	20+50, 02R	1.2	None				
175	25+50, 04L	1.2	0–15	0–15	Not sampled (see hole No. 13)		
176	37+02, 14R	1.2	0–15	0–15	Not sampled (see hole No. 71)		
177	30+98, 15R	1.2	0–15	0–15	Not sampled (see hole No. 65)		
178	28+02, 06R	1.2	0–60	0–15	Not sampled (see hole No. 62)		
179	26+02, 21L	1.2	0–45	0–15	Not sampled (see hole No. 77)		
180	29+21, 12R	1.2	0–120	90–120	1.4±0.2	28	1.0
181	27+10, 07R	1.2	0–15	0–15	Not sampled (see hole No. 61)		
182	25+97, 07R	1.2	0–30	0–15	Not sampled (see hole No. 60)		
183	25+04, 23R	1.2	0–60	45–60	Not sampled (see hole No. 57)		
184	24+47, 15R	1.2	0–15	0–15	Not sampled (see hole No. 97)		

Table 5 (continued)

Hole	Location ^a	Depth of hole (m)	Estimated vertical extent of contaminated soil ^b (cm)	Region of maximum contamination (cm)	Radionuclide concentration in region of maximum contamination (pCi/g)		
					²²⁶ Ra	²³⁰ Th ^c	²³⁸ U ^d
185	23+98, 14R	1.2	0-60	0-15	Not sampled (see hole No. 99)		
186	22+52, 12R	1.2	0-45	15-30	Not sampled (see hole No. 55)		
187	20+98, 25R	1.2	0-30	15-30	Not sampled (see hole No. 52)		
188	18+94, 12L	1.2	0-90	0-15	Not sampled (see hole No. 85)		
189	17+96, 13L	0.9	0-30	15-30	9.2 ± 0.8	360	6.3
190	12+24, 17R	1.2	0-15	0-15	Not sampled (see hole No. 35)		
191	12+60, 32L	1.2	None				
192	7+96, 19L	1.2	0-30	0-30	14 ± 0.3	870	10
193	30+56, 19R	1.2	0-30	0-30	1.3 ± 0.1	7.6	1.2
194	30+39, 19L	1.2	None				
195	29+09, 10R	0.6	0-120	0-60	2.7 ± 0.1	53	1.6
196	27+18, 14R	0.6	0-90	30-60	2.4 ± 0.1	29	2.5
197	26+32, 13R	0.9	0-60	0-30	2.4 ± 0.2	40	2.8
198	23+66, 17L	1.2	0-60	0-30	11 ± 0.2	400	6.2
199	23+09, 37R	2.4	None				
200	22+40, 45R	2.4	None				
201	21+81, 38R	2.3	None				
202	13+59, 27R	1.8	None				
203	13+30, 28R	1.8	None				
204	11+77, 24R	1.7	None				
205	9+48, 22R	1.7	None				
206	9+20, 28R	1.7	None				
207	9+22, 16R	1.2	None				
208	9+88, 20L	1.2	0-30	15-30	Not sampled (see hole No. 109)		
209	7+02, 16R	1.8	0-75	0-15	Not sampled (see holes No. 122, 123, 124)		
210	6+00, 28L	1.2	0-75	0-15	Not sampled (see hole No. 117)		
211	5+58, 09R	1.7	0-165	30-45	Not sampled (see hole No. 120)		
212	5+52, 20L	2.4	0-180	60-90	10 ± 0.5	450	11
213	14+48, 35R	1.7	0-30	0-30	1.8 ± 0.2	44	1.8
214	5+50, 50L	1.5	0-15	0-15	Not sampled (see holes No. 143, 172)		
215	4+50, 106L	1.7	0-15	0-15	1.5 ± 0.1	19	1.5
216	7+90, 210R	0.9	0-30	0-30	55 ± 3.0	2,700	2.4
217	7+39, 96R	1.2	0-90	30-60	28 ± 2.0	1,300	24
218	7+10, 50R	1.4	0-120	0-30	16 ± 1.0	520	19
219	5+89, 50R	1.2	0-60	0-15	Not sampled (see hole No. 160)		
220	8+11, 43R	1.2	0-30	0-30	3.6 ± 0.2	110	3.1
221	8+25, 150R	1.4	0-135	0-15	82 ± 0.3	3,700	93

^aSee Figs. 16-31 for location.

^bContaminated soil is defined as soil having concentrations of ²²⁶Ra or ²³⁰Th equal to or greater than the criteria given in Table 1; or giving rise to 1,000 cpm or more on the shielded scintillator.

^cThe error of the reported radionuclide concentration is less than ± 5% (95% confidence level).

^dThe total analytical error is less than ± 3% (95% confidence level).

Table 6. Concentration of selected radionuclides in water samples taken from auger holes at the Latty Avenue site (LM001)

Sample number	Grid location ^a	Concentration of radionuclide ^b (pCi/L)			
		²²⁶ Ra	²³⁸ U	²³⁰ Th	²¹⁰ Pb
30	8+47, 05L	<0.27	<6.7	<0.81	0.27 ± 2.2
31	7+50, 02L	<0.27	<6.7	<0.81	0.27 ± 2.2
32	6+50, 05L	<0.27	<6.7	<1.1	2.2 ± 2.4
33	5+50, 07L	<0.81	<6.7	<0.81	0.27 ± 2.2
151	8+15, 100R	<8.1	30	1.1 ± 0.14	<1.9
152	8+13, 39R	1.9 ± 3.5	3.7 ± 1.3	2.0 ± 0.16	0.27 ± 2.2
157	7+36, 95R	<0.54	53	<1.6	0.27 ± 2.2
159	6+50, 50R	<0.54	63	0.86 ± 0.11	2.7 ± 2.7

^aSee Figs. 16–31 for locations of auger holes.

^bIndicated errors associated with concentrations are 2 σ (95% confidence level).

**Table 7. Summary of measurements and sample results
at Latty Avenue (LM001)**

Measurement or sample type	Number of measurement/ samples	Range	Mean
Gamma exposure rate at 1 m ($\mu\text{R/h}$) ^a	336	4–380	16
Gamma exposure rate at surface ($\mu\text{R/h}$) ^a	336	4–490	17
Scan, gamma exposure rate near surface ($\mu\text{R/h}$) ^b	–	4–720	–
Concentration of ²²⁶ Ra in surface soil (pCi/g), systematic locations ^c	111	1.0–12	2.7
Concentration of ²³⁰ Th in surface soil (pCi/g), systematic locations ^c	111	1.6–630	54
Concentration of ²³⁸ U in surface soil (pCi/g), systematic locations ^c	111	0.99–15	2.1
Concentration of ²²⁶ Ra in surface soil (pCi/g), biased locations ^d	3	8.3–12	9.7
Concentration of ²³⁰ Th in surface soil (pCi/g), biased locations ^d	3	350–810	550
Concentration of ²³⁸ U in surface soil (pCi/g), biased locations ^d	3	5.4–8.7	7.6
Maximum ²²⁶ Ra concentration in soil from auger holes, (pCi/g) ^e	221	0.56–660	–
Maximum ²³⁰ Th concentration in soil from auger holes, (pCi/g) ^e	221	0.90–22,000	–
Maximum ²³⁸ U concentration in soil from auger holes, (pCi/g) ^e	221	0.41–790	–
Concentration of ²²⁶ Ra in water from auger holes, (pCi/L) ^f	8	<0.27–<8.1	1.6
Concentration of ²³⁰ Th in water from auger holes, (pCi/L) ^f	8	<0.81–2.0	1.1
Concentration of ²³⁸ U in water from auger holes, (pCi/L) ^f	8	3.7–63	22
Concentration of ²¹⁰ Pb in water from auger holes, (pCi/L) ^f	8	<0.27–2.7	1.0

^aAt grid points.

^bScan of entire property.

^cSystematic samples, Table 4.

^dBiased samples, Table 4.

^eAuger hole samples, Table 4.

^fWater samples, Table 6.

APPENDIX

GAMMA PROFILE GRAPHS OF AUGER HOLES AT PROPERTY LM001

ORNL-DWG 87-11458

Fig. A.1. Gamma profile of auger hole 1.

ORNL-DWG 87-11459

Fig. A.2. Gamma profile of auger hole 2.

ORNL-DWG 87-11460

Fig. A.3. Gamma profile of auger hole 3.

ORNL-DWG 87-11461

Fig. A.4. Gamma profile of auger hole 4.

Fig. A.5. Gamma profile of auger hole 5.

ORNL-DWG 87-11463

Fig. A.6. Gamma profile of auger hole 6.

ORNL-DWG 87-11464

Fig. A.7. Gamma profile of auger hole 7.

ORNL-DWG 87-11465

Fig. A.8. Gamma profile of auger hole 8.

ORNL-DWG 87-11466

Fig. A.9. Gamma profile of auger hole 9.

Fig. A.10. Gamma profile of auger hole 10.

ORNL-DWG 87-11468

Fig. A.11. Gamma profile of auger hole 11.

ORNL-DWG 87-11469

Fig. A.12. Gamma profile of auger hole 12.

ORNL-DWG 87-11470

Fig. A.13. Gamma profile of auger hole 13.

ORNL-DWG 87-11471

Fig. A.14. Gamma profile of auger hole 14.

ORNL-DWG 87-11472

Fig. A.15. Gamma profile of auger hole 15.

ORNL-DWG 87-11473

Fig. A.16. Gamma profile of auger hole 16.

ORNL-DWG 87-11474

Fig. A.17. Gamma profile of auger hole 17.

ORNL-DWG 87-11475

Fig. A.18. Gamma profile of auger hole 18.

ORNL-DWG 87-11476

Fig. A.19. Gamma profile of auger hole 19.

Fig. A.20. Gamma profile of auger hole 20.

ORNL-DWG 87-11478

Fig. A.21. Gamma profile of auger hole 21.

Fig. A.22. Gamma profile of auger hole 22.

ORNL-DWG 87-11480

Fig. A.23. Gamma profile of auger hole 23.

ORNL-DWG 87-11481

Fig. A.24. Gamma profile of auger hole 24.

ORNL-DWG 87-11482

Fig. A.25. Gamma profile of auger hole 25.

ORNL-DWG 87-11483

Fig. A.26. Gamma profile of auger hole 26.

ORNL-DWG 87-11484

Fig. A.27. Gamma profile of auger hole 27.

ORNL-DWG 87-11485

Fig. A.28. Gamma profile of auger hole 28.

ORNL-DWG 87-11486

Fig. A.29. Gamma profile of auger hole 29.

ORNL-DWG 87-11487

Fig. A.30. Gamma profile of auger hole 30.

ORNL-DWG 87-11488

Fig. A.31. Gamma profile of auger hole 31.

ORNL-DWG 87-11489

Fig. A.32. Gamma profile of auger hole 32.

ORNL-DWG 87-11490

Fig. A.33. Gamma profile of auger hole 33.

ORNL-DWG 87-11491

Fig. A.34. Gamma profile of auger hole 34.

ORNL-DWG 87-11492

Fig. A.35. Gamma profile of auger hole 35.

ORNL-DWG 87-11493

Fig. A.36. Gamma profile of auger hole 36.

ORNL-DWG 87-11494

Fig. A.37. Gamma profile of auger hole 37.

ORNL-DWG 87-11495

Fig. A.38. Gamma profile of auger hole 38.

ORNL-DWG 87-11496

Fig. A.39. Gamma profile of auger hole 39.

ORNL-DWG 87-11497

Fig. A.40. Gamma profile of auger hole 40.

ORNL-DWG 87-11498

Fig. A.41. Gamma profile of auger hole 41.

ORNL-DWG 87-11499

Fig. A.42. Gamma profile of auger hole 42.

ORNL-DWG 87-11500

Fig. A.43. Gamma profile of auger hole 43.

ORNL-DWG 87-11501

Fig. A.44. Gamma profile of auger hole 44.

ORNL-DWG 87-11502

Fig. A.45. Gamma profile of auger hole 45.

ORNL-DWG 87-11503

Fig. A.46. Gamma profile of auger hole 46.

ORNL-DWG 87-11504

Fig. A.47. Gamma profile of auger hole 47.

ORNL-DWG 87-11505

Fig. A.48. Gamma profile of auger hole 48.

ORNL-DWG 87-11506

Fig. A.49. Gamma profile of auger hole 49.

Fig. A.50. Gamma profile of auger hole 50.

ORNL-DWG 87-11508

Fig. A.51. Gamma profile of auger hole 51.

ORNL-DWG 87-11509

Fig. A.52. Gamma profile of auger hole 52.

ORNL-DWG 87-11510

Fig. A.53. Gamma profile of auger hole 53.

ORNL-DWG 87-11511

Fig. A.54. Gamma profile of auger hole 54.

ORNL-DWG 87-11512

Fig. A.55. Gamma profile of auger hole 55.

ORNL-DWG 87-11513

Fig. A.56. Gamma profile of auger hole 56.

ORNL-DWG 87-11514

Fig. A.57. Gamma profile of auger hole 57.

Fig. A.58. Gamma profile of auger hole 58.

Fig. A.59. Gamma profile of auger hole 59.

ORNL-DWG 87-11517

Fig. A.60. Gamma profile of auger hole 60.

ORNL-DWG 87-11518

Fig. A.61. Gamma profile of auger hole 61.

Fig. A.62. Gamma profile of auger hole 62.

Fig. A.63. Gamma profile of auger hole 63.

ORNL-DWG 87-11521

Fig. A.64. Gamma profile of auger hole 64.

ORNL-DWG 87-11522

Fig. A.65. Gamma profile of auger hole 65.

Fig. A.66. Gamma profile of auger hole 66.

ORNL-DWG 87-11524

Fig. A.67. Gamma profile of auger hole 67.

Fig. A.68. Gamma profile of auger hole 68.

Fig. A.69. Gamma profile of auger hole 69.

ORNL-DWG 87-11527

Fig. A.70. Gamma profile of auger hole 70.

ORNL-DWG 87-11528

Fig. A.71. Gamma profile of auger hole 71.

ORNL-DWG 87-11529

Fig. A.72. Gamma profile of auger hole 72.

ORNL-DWG 87-11530

Fig. A.73. Gamma profile of auger hole 73.

ORNL-DWG 87-11531

Fig. A.74. Gamma profile of auger hole 74.

ORNL-DWG 87-11532

Fig. A.75. Gamma profile of auger hole 75.

ORNL-DWG 87-11533

Fig. A.76. Gamma profile of auger hole 76.

ORNL-DWG 87-11534

Fig. A.77. Gamma profile of auger hole 77.

Fig. A.78. Gamma profile of auger hole 78.

ORNL-DWG 87-11536

Fig. A.79. Gamma profile of auger hole 79.

ORNL-DWG 87-11537

Fig. A.80. Gamma profile of auger hole 80.

ORNL-DWG 87-11538

Fig. A.81. Gamma profile of auger hole 81.

Fig. A.82. Gamma profile of auger hole 82.

ORNL-DWG 87-11540

Fig. A.83. Gamma profile of auger hole 83.

ORNL-DWG 87-11541

Fig. A.84. Gamma profile of auger hole 84.

ORNL-DWG 87-11542

Fig. A.85. Gamma profile of auger hole 85.

Fig. A.86. Gamma profile of auger hole 86.

ORNL-DWG 87-11544

Fig. A.87. Gamma profile of auger hole 87.

ORNL-DWG 87-11545

Fig. A.88. Gamma profile of auger hole 88.

ORNL-DWG 87-11546

Fig. A.89. Gamma profile of auger hole 89.

ORNL-DWG 87-11547

Fig. A.90. Gamma profile of auger hole 90.

ORNL-DWG 87-11548

Fig. A.91. Gamma profile of auger hole 91.

Fig. A.92. Gamma profile of auger hole 92.

ORNL-DWG 87-11550

Fig. A.93. Gamma profile of auger hole 93.

ORNL-DWG 87-11551

Fig. A.94. Gamma profile of auger hole 94.

ORNL-DWG 87-11552

Fig. A.95. Gamma profile of auger hole 95.

Fig. A.96. Gamma profile of auger hole 96.

ORNL-DWG 87-11554

Fig. A.97. Gamma profile of auger hole 97.

Fig. A.98. Gamma profile of auger hole 98.

Fig. A.99. Gamma profile of auger hole 99.

ORNL-DWG 87-11557

Fig. A.100. Gamma profile of auger hole 100.

ORNL-DWG 87-11558

Fig. A.101. Gamma profile of auger hole 101.

Fig. A.102. Gamma profile of auger hole 102.

ORNL-DWG 87-11560

Fig. A.103. Gamma profile of auger hole 103.

ORNL-DWG 87-11561

Fig. A.104. Gamma profile of auger hole 104.

ORNL-DWG 87-11562

Fig. A.105. Gamma profile of auger hole 105.

ORNL-DWG 87-11563

Fig. A.106. Gamma profile of auger hole 106.

ORNL-DWG 87-11564

Fig. A.107. Gamma profile of auger hole 107.

ORNL-DWG 87-11565

Fig. A.108. Gamma profile of auger hole 108.

ORNL-DWG 87-11566

Fig. A.109. Gamma profile of auger hole 109.

ORNL-DWG 87-11567

Fig. A.110. Gamma profile of auger hole 110.

ORNL-DWG 87-11568

Fig. A.111. Gamma profile of auger hole 111.

Fig. A.112. Gamma profile of auger hole 112.

ORNL-DWG 87-11570

Fig. A.113. Gamma profile of auger hole 113.

ORNL-DWG 87-11571

Fig. A.114. Gamma profile of auger hole 114.

ORNL-DWG 87-11572

Fig. A.115. Gamma profile of auger hole 115.

ORNL-DWG 87-11573

Fig. A.116. Gamma profile of auger hole 116.

ORNL-DWG 87-11574

Fig. A.117. Gamma profile of auger hole 117.

ORNL-DWG 87-11575

Fig. A.118. Gamma profile of auger hole 118.

Fig. A.119. Gamma profile of auger hole 119.

ORNL-DWG 87-11577

Fig. A.120. Gamma profile of auger hole 120.

Fig. A.121. Gamma profile of auger hole 121.

Fig. A.122. Gamma profile of auger hole 122.

Fig. A.123. Gamma profile of auger hole 123.

Fig. A.124. Gamma profile of auger hole 124.

Fig. A.125. Gamma profile of auger hole 125.

ORNL-DWG 87-11583

Fig. A.126. Gamma profile of auger hole 126.

Fig. A.127. Gamma profile of auger hole 127.

Fig. A.128. Gamma profile of auger hole 128.

Fig. A.129. Gamma profile of auger hole 129.

Fig. A.130. Gamma profile of auger hole 130.

ORNL-DWG 87-11588

Fig. A.131. Gamma profile of auger hole 131.

Fig. A.132. Gamma profile of auger hole 132.

Fig. A.133. Gamma profile of auger hole 133.

Fig. A.134. Gamma profile of auger hole 134.

ORNL-DWG 87-11592

Fig. A.135. Gamma profile of auger hole 135.

Fig. A.136. Gamma profile of auger hole 136.

ORNL-DWG 87-11594

Fig. A.137. Gamma profile of auger hole 137.

Fig. A.138. Gamma profile of auger hole 138.

Fig. A.139. Gamma profile of auger hole 139.

ORNL-DWG 87-11597

Fig. A.140. Gamma profile of auger hole 140.

ORNL-DWG 87-11598

Fig. A.141. Gamma profile of auger hole 141.

Fig. A.142. Gamma profile of auger hole 142.

ORNL-DWG 87-11600

Fig. A.143. Gamma profile of auger hole 143.

ORNL-DWG 87-11601

Fig. A.144. Gamma profile of auger hole 144.

ORNL-DWG 87-11602

Fig. A.145. Gamma profile of auger hole 145.

Fig. A.146. Gamma profile of auger hole 146.

Fig. A.147. Gamma profile of auger hole 147.

Fig. A.148. Gamma profile of auger hole 148.

ORNL-DWG 87-11606

Fig. A.149. Gamma profile of auger hole 149.

Fig. A.150. Gamma profile of auger hole 150.

ORNL-DWG 87-11608

Fig. A.151. Gamma profile of auger hole 151.

Fig. A.152. Gamma profile of auger hole 152.

ORNL-DWG 87-11610

Fig. A.153. Gamma profile of auger hole 153.

Fig. A.154. Gamma profile of auger hole 154.

ORNL-DWG 87-11612

Fig. A.155. Gamma profile of auger hole 155.

Fig. A.156. Gamma profile of auger hole 156.

ORNL-DWG 87-11614

Fig. A.157. Gamma profile of auger hole 157.

Fig. A.158. Gamma profile of auger hole 158.

Fig. A.159. Gamma profile of auger hole 159.

Fig. A.160. Gamma profile of auger hole 160.

Fig. A.161. Gamma profile of auger hole 161.

ORNL-DWG 87-11619

Fig. A.162. Gamma profile of auger hole 162.

Fig. A.163. Gamma profile of auger hole 163.

Fig. A.164. Gamma profile of auger hole 165.

ORNL-DWG 87-11622

Fig. A.165. Gamma profile of auger hole 166.

Fig. A.166. Gamma profile of auger hole 167.

ORNL-DWG 87-11624

Fig. A.167. Gamma profile of auger hole 168.

Fig. A.168. Gamma profile of auger hole 169.

Fig. A.169. Gamma profile of auger hole 170.

Fig. A.170. Gamma profile of auger hole 171.

ORNL-DWG 87-11628

Fig. A.171. Gamma profile of auger hole 172.

ORNL-DWG 87-11629

Fig. A.172. Gamma profile of auger hole 173.

251 / 252

ORNL-DWG 87-11630

Fig. A.173. Gamma profile of auger hole 215.

INTERNAL DISTRIBUTION

1. B. A. Berven	30. C. R. Richmond
2-6. R. F. Carrier	31. T. H. Row
7. R. O. Chester	32. P. J. Walsh
8-17. W. D. Cottrell	33. J. K. Williams
18. J. T. Ensminger	34. J. P. Witherspoon
19. R. D. Foley	35. D. A. Witt
20. R. A. Griesemer	36. M. G. Yalcintas
21. A. R. Hawthorne	37. Biology Library
22. C. A. Johnson	38-39. Central Research Library
23. S. V. Kaye	40. ORNL Technical Library, Y-12
24. G. D. Kerr	41. IR&A Publications Office
25. C. A. Little	42-43. Information Analysis Library
26. J. L. Marley	44-45. Laboratory Records
27. C. J. Owen	46. Laboratory Records - RC
28. P. T. Owen	47. ORNL Patent Office
29. H. A. Pfuderer	

EXTERNAL DISTRIBUTION

- 48-52. S. W. Ahrends, U.S. Department of Energy, Technical Services Division, Oak Ridge Operations Office, Oak Ridge, TN 37831
53. J. D. Berger, Oak Ridge Associated Universities, P.O. Box 117, Oak Ridge, TN 37831
- 54-56. J. J. Fiore, U.S. Department of Energy, Director, Division of Facilities and Site Decommissioning Projects, Office of Nuclear Energy, 19901 Germantown Road, Germantown, MD 20874
- 57-60. C. E. Miller, U.S. Department of Energy, Director, Surplus Facilities Management, Richland Operations Office, P.O. Box 550, Richland, WA 99352
- 61-62. J. F. Nemec, Bechtel National, Inc., 800 Oak Ridge Turnpike, Oak Ridge, TN 37831
63. G. P. Turi, U.S. Department of Energy, Division of Facilities and Site Decommissioning Projects, Office of Nuclear Energy, 19901 Germantown Road, Germantown, MD 20874
64. J. W. Wagoner II, U.S. Department of Energy, Division of Facilities and Site Decommissioning Projects, Office of Nuclear Energy, 19901 Germantown Road, Germantown, MD 20874
65. A. Wallo, Aerospace Corporation, Environmental and Conservation Directorate, 955 L'Enfant Plaza, SW, Suite 4000, Washington, DC 20024
66. Office of Assistant Manager, Energy Research and Development, Oak Ridge Operations Office, Oak Ridge, TN 37831
- 67-96. Office of Scientific and Technical Information, U.S. Department of Energy, Oak Ridge, TN 37831