DES

The mass splittings reported in this paper indicate that the Σ_c is the first multiplet for which the intrinsic light quark mass difference is not the domi

013330

1 OF 2

PURDUE HIGH ENERGY PARTICLE PHYSICS

Progress and Research Proposal

11/1/93 - 10/31/94

(3 volumes, Budgets, Progress, Proposal)

Volume 2/3 - PROGRESS

Principal Investigators

Experimental

**Task A: J. P. Finley, J. A. Gaidos, F. J. Loeffler,
R. L. McIlwain, D. H. Miller, T. R. Palfrey,
E. I. Shibata, I. P. Shipsey**

Task D: D. D. Carmony, L. J. Gutay

Task E: D. S. Koltick, R. L. McIlwain, E. I. Shibata

Task G: V. E. Barnes, D. Bortoletto, A. F. Garfinkel

Theoretical

**Task B: R. H. Capps, T. E. Clark, T. K. Kuo,
S. T. Love**

Task F: E. Fischbach

**PURDUE UNIVERSITY
West Lafayette, Indiana 47907**

AN EXPERIMENTAL HIGH ENERGY PHYSICS PROGRAM

within the

DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY
West Lafayette, Indiana 47907

Supported by

U.S. DEPARTMENT OF ENERGY

Through

Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana

PROGRESS

for

11/1/93 - 10/31/94

Current Grant Number: DE-FG02-19ER40681 Task A

Principal Investigators: J. Finley, J. A. Gaidos, F. J. Loeffler, R. L. McIlwain,
D. H. Miller*, T. R. Palfrey, E. I. Shibata, I. P. Shipsey

April 1993

*Spokesperson

MASTER

REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

1. Recent CLEO Publications	03
2. Recent Particle Astrophysics Publications	05
3. Review of CLEO 1992-1993	06
4. R& D on Gas Microstrip Detectors	13
5. Review of Particle Astrophysics	16

1. CLEO PUBLICATIONS

1. Inclusive and Exclusive Decays of B Mesons to Final States Including Charm and Charmonium Mesons. D. Bortoletto et al., Physical Review D **45**, 21 (1992)
2. Measurement of Baryon Production in B-Meson Decay. G. Crawford et al., Physical Review D **45**, 752 (1992)
3. D_s^+ decays to $\eta\pi^+$ and $\eta'\pi^+$. P. Avery et al., Physical Review Letters **68**, 1275 (1992)
4. D_s^+ decays to $\eta\rho^+$, $\eta'\rho^+$, and $\phi\rho^+$. P. Avery et al., Physical Review Letters **68**, 1279 (1992)
5. Measurements of Semileptonic Branching Fractions of B Mesons at the $\Upsilon(4S)$ Resonance. S. Henderson et al., Physical Review D **45**, 2212 (1992)
6. Two-body D_s^+ decays to $\eta\pi^+$, $\eta'\pi^+$, $\eta\rho^+$, $\eta'\rho^+$, and $\phi\rho^+$. M. Daoudi et al., Physical Review D **45**, 3965 (1992)
7. Electronic Branching Ratio of the τ lepton. R. Ammar et al., Physical Review D **45**, 3976 (1992)
8. Observation of the Decay $\Xi_c^0 \rightarrow \Omega^- K^+$. S. Henderson et al., Physics Letters B **283**, 16 (1992)
9. The CLEO II detector. Y. Kubota et al., Nuclear Instruments and Methods A **320**, 66 (1992)
10. A Measurement of the tau lepton lifetime. M. Battle et al., Physics Letters B **291**, 488 (1992)
11. Measurement of the $D^*(2010)$ Branching Fractions. F. Butler et al., Physical Review Letters **69**, 2041 (1992)
12. Isospin Mass Splittings from Precision Measurements of $D^* - D$ Mass Differences. D. Bortoletto et al., Physical Review Letters **69**, 2046 (1992)
13. Shape Studies of Quark Jets vs. Gluon Jets at $\sqrt{s} = 10$ GeV. M. S. Alam et al., Physical Review D **46**, 4822 (1992)
14. Exclusive $\chi(2P)$ Production in $\Upsilon(3S)$ Decay. G. Crawford et al., Physics Letters B **294**, 139 (1992)
15. Measurement of τ Decays Involving η Mesons. M. Artuso et al., Physical Review Letters **69**, 3278 (1992)
16. Measurement of the Tau Lepton Electronic Branching Fraction. D. S. Akerib et al., Physical Review Letters **69**, 3610 (1992)
17. Search for $\tau^- \rightarrow \gamma\mu^-$: A Test of Lepton Number Conservation. A. Bean et al., Physical Review Letters **70**, 138 (1993)
18. Tau Decays with One Charged Particle plus Multiple π^0 's. M. Procaro et al., Physical Review Letters **70**, 1207 (1993)
19. Lepton Asymmetry Measurements in $\bar{B} \rightarrow D^* l \nu_l$ and Implications for V-A and the Form Factors. S. Sanghera et al., Physical Review D **47**, 791 (1993)

20. Study of D^0 decays into K^0 and K^{*0} . M. Procaro et al. CLNS 92/1167, CLEO92-10(submitted to Physical Review D)
21. Search for Exclusive $b \rightarrow u$ Semileptonic Decays of B Mesons. A. Bean et al. CLNS 92/1174, CLEO 92-12(submitted to Physical Review Letters)
22. Production and Decay of the $D_{s1}(2536)^+$. J.P. Alexander et al. CLNS 93/1191, CLEO 93-1(submitted to Physics Letters B)
23. A Measurement of the Tau Lepton Mass. R. Ballest et al. CLNS 93/1194, CLEO 93-2(submitted to Physical Review Letters)
24. A Limit on the Tau Neutrino Mass. D. Cinabro et al. CLNS 93/1195, CLEO 93-3(submitted to Physical Review Letters)

2. PARTICLE ASTROPHYSICS PUBLICATIONS

1. Detection of TeV Photons from the Active Galaxy Markarian 421, Punch et al., *Nature*, Vol. 358, (1992) 477
2. Search for TeV Gamma Ray Emission from AGN's Using the Whipple Imaging Telescope, Fennell et al., *Proceedings of the Compton Observatory Symposium*, Oct. 1992, American Institute of Physics (to be published).
3. GRANITE-A Stereoscopic Imaging Cherenkov Telescope System, Schubnell et al., *Proceedings of the Compton Observatory Symposium*, Oct. 1992, American Institute of Physics (to be published).
4. Search for TeV Gamma Rays from Geminga, Fegan, et. al., *Proceedings of the Compton Observatory Symposium*, Oct. 1992, American Institute of Physics (to be published); expanded paper to be published in *Astrophysical Journal* (1993).
5. A Method for Searching the Whipple Observatory Gamma Ray Data Base for Evidence of GRB's, Chantell et al., *Proceedings of the Compton Observatory Symposium*, Oct. 1992, American Institute of Physics (to be published).
6. The Very High Energy Photon Spectrum of Markarian 421, Mohanty et al., (submitted to *Astrophysical Journal Letters*).
7. Observations of the Crab Pulsar with the ASGAT Atmospheric Cherenkov Telescope, Vancanti et al., *Taos Conference on Pulsars 1992*, (to be published).
8. Observations of the Crab Pulsar with the ASGAT Atmospheric Cherenkov Telescope. Goret et al., *Astronomy and Astrophysics*, (May) 1993.
9. Mirrors for Gamma Ray Astronomy, Loeffler, *Nuclear Instruments and Methods*, (accepted for publication) 1993.

3. REVIEW OF CLEO, 1992-1993

Currently the CLEO experiment is one of the forefront experiments for the study of heavy flavor physics. The exceptional performance of the accelerator CESR, together with the stable and high performance operation of CLEO-II, is allowing some of the most precise and sensitive studies of the b quark, charm quark and the tau lepton, as well as low-multiplicity physics such as the physics of two-photon collisions.

3.1. REVIEW OF 1992

For calendar year 1992 a total of 1.4 fb^{-1} was accumulated. On our best days, we collected in excess of 12 pb^{-1} . CLEO-II now has a total of 1.5 fb^{-1} on the $\Upsilon(4S)$, corresponding to 3 million B meson decays. The luminosity collected each calendar year by CLEO is shown in see Fig. 1.

Figure 1: CLEO-II luminosity by calendar year

3.2. PHYSICS IN 1992-93

Some of the highlights have been:

- Measurements of the rate for b-quark decays into u-quark indicating a value of $|V_{ub}|$ significantly lower than indicated by previous experimental results.
- Highest precision measurement to date of $B\bar{B}$ mixing, using two complementary techniques.
- The definitive measurements of tau-decays to one-prong plus multiple π^0 's ($\tau^- \rightarrow \nu_\tau \pi^- n \pi^0$).
- The world's single best measurement of the electronic branching ratio of the tau-lepton (B_e), as well as the best limit on the rare decay of the tau to the $\mu\gamma$ final state.
- The world's first observation of the Cabibbo-suppressed semileptonic charm decay: $D^- \rightarrow \pi^0 \ell^- \nu_\ell$.
- The first measurement of the masses of the three Σ_c states in the same experiment.
- The first simultaneous measurement of all the direct cascade transitions from the $\Upsilon(3S)$ to the $\Upsilon(2S)$ and $\Upsilon(1S)$: the $\gamma\gamma$, $\pi^0\pi^0$, and $\pi^+\pi^-$ transitions.

3.3. DATA SET AND RUNNING

1992 has been a year of unequalled productivity since the inception of the CLEO experiment in the early 1980's. We have had a very high duty cycle for data collection while the storage ring (CESR) has been delivering luminosity, and we have kept the calibration and operation of the detector at the highest levels of performance achieved to date. The fact that the experiment as a whole continues to grow at the rate of adding 1-2 new institutions per year attests to the high visibility of the experiment and its attractiveness as a source of interesting physics.

The total CLEO-II data set processed and available for analysis is presented in Table I.

YEAR	Center-of-mass energy (GeV)	$\mathcal{L}$ (/pb)
1989	10.35 ($\Upsilon(3S)$)	132
1990	10.55-10.62 ($\bar{B}B^*$ scan)	130
1990	9.46 ($\Upsilon(1S)$)	15
1990-91	10.55-10.59 ($\Upsilon(4S)$ +continuum)	672
1991-92	10.55-10.59 ($\Upsilon(4S)$ +continuum)	680
1992	10.55-10.59 ($\Upsilon(4S)$ +continuum)	318
1992	9.46 ($\Upsilon(1S)$)	48
1992-1993	10.55-10.59 ($\Upsilon(4S)$ +continuum)	613
1989-1993, total	9.46-10.62	2608

Table I: Summary of CLEO-II dataset

3.4. SPECIFIC PURDUE ACTIVITIES

Since July 1, 1992, David H. Miller has been the spokesperson for the CLEO collaboration and has been deeply involved in the day to day running, the next phase upgrade of the silicon vertex detector, and the longer term high luminosity running.

3.5. DETAILED ANALYSIS OF BOUND STATE UPSILON DECAYS

David N. Brown completed his doctoral thesis entitled "Inclusive spectra in the Upsilon energy region and a study of quark-jet versus gluon-jet fragmentation" in November 1992 under the supervision of Professor Shibata. He measured the various particle ($\pi^\pm$, $K^\pm$, p , $\bar{p}$, K_s^* , π^* , Λ^* , ϕ^* , D^* , ρ^* , K^{*0} , and $e^\pm$) yields as a function of momentum from $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(3S)$, $\Upsilon(4S)$, the nearby continuum, and χ'_b states. These measurements of so many different particle species required the use of several particle identification methods. As part of his service work for CLEO David worked on charged particle identification by specific ionization measurements and combined that information with that from the time of flight system. David was also the caretaker of the fitting/plotting program MN.FIT, which is

used by the whole collaboration and was used to fit his various spectra. The decays of χ'_b with $J = 1$ were found to exhibit properties most similar to those observed in continuum $q\bar{q}$ decays, while the properties of χ'_b with $J = 0$ and 2 are most like the ggg decays of the $\Upsilon(1S)$ state. Dr. Brown is now a research associate in experimental high energy physics at the University of Pittsburgh.

Jay M. Dominick completed his doctoral thesis entitled "Hadronic transitions between bottomonium states" in March 1993 under the supervision of Professor Shibata. The branching ratios for $\Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(1S)$, $\Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(2S)$, $\Upsilon(2S) \rightarrow \pi^+\pi^-\Upsilon(1S)$ were measured and the first indication of $\pi^+\pi^-$ transitions between χ states was observed using the decay chain $\Upsilon(3S) \rightarrow \gamma_1\chi_{b1}(2P)$, $\chi_{b1}(2P) \rightarrow \pi^+\pi^-\chi_{b1}(1P)$, $\chi_{b1}(1P) \rightarrow \gamma_2\Upsilon(1S)$. A search for the h_b through the decay $\Upsilon(3S) \rightarrow \pi^+\pi^-h_b(9900)$ yielded an upper limit of $L0.18\gamma(3S) \rightarrow \pi^+\pi^-h_b$ contributions to the tracking of charged particles with momenta less than ~ 100 MeV/c; such momenta occurred frequently in the decays he studied. Dr. Dominick is now a research associate in experimental high energy physics at Southern Methodist University.

3.6. EXCLUSIVE B DECAYS

David Miller and his student, Michael Modesitt, are involved in the analysis of exclusive hadronic B decays. In particular, we are analysing exclusive decays of the type $D^{(*)}D_s^{(*)}$ where the virtual W has decayed to a $D_s^{(*)}$. This will allow a detailed determination of the decay constant of the D_s . This is part of the more general physics goals detailed below.

Studies of the factorizability of the $b \rightarrow c$ vertex The "factorization" hypothesis has been widely used in theoretical predictions for hadronic decays of heavy flavored hadrons. It implies that the hadronization of the virtual W is independent of the hadronization at the lower vertex. This notion leads immediately to relationships between decays: $B^0 \rightarrow D^*\ell\nu$ and $B \rightarrow D^*\rho^+$, for example, which depend only on the W-mass (measurable in both cases) and polarization (also measurable). In the latter case, one needs also to consider the probability of the W hadronizing into a ρ^+ (which, in turn, depends on the ρ^+ decay constant f_{ρ^+}). By comparing the rates and polarizations at a fixed W-mass for the leptonic vs. hadronic B-decays, we can therefore both test the factorization hypothesis as well as extract information on hadron decay constants.

Our studies of $\frac{d\Gamma}{dq^2} \left(\frac{D^*\rho}{D^*\ell\nu} \right) |_{M_\rho=M_{\ell\nu}}$ indicate that the factorization hypothesis seems to be valid at the 10% level for B-decay. We are now evaluating this hypothesis for the higher q^2 decay of, e.g., $B \rightarrow D^*D_s^*$, compared with the semileptonic decay at $M_{\ell\nu} = M_{D_s^*}$.

Studies of internal vs. external W-emission In charm decays, we observe decays such as $D^0 \rightarrow K^0\pi^0$ with rates which are roughly half that of $D^0 \rightarrow K^-\pi^+$. Phenomenologically, the difference between the two can be understood as the difference between so-called "internal" ($D^0 \rightarrow K^0\pi^0$) W-emission vs. "external" ($D^0 \rightarrow K^-\pi^+$) W-emission. The former case is also referred to as the "color-suppressed" decay, as one needs to have the quarks which form the π^0 bind in a color singlet. Naively, random color matching argues that this should happen only 1/3 of the time (hence, a factor of 9 suppression in rate, with an additional factor of

two due to the π^0 wave function). We note here that such states as $D^0\pi^0$, $D^0\eta^0$, and $D^0\eta'$ all require high precision calorimetry for optimal acceptance.

Although such color-suppressed decays of the B-meson for which the W- produces a $c\bar{s}$ ($B \rightarrow \psi K_s^0$, e.g.) have been studied for at least five years, color-suppressed decays of the B-meson for which the W- hadronizes as a $u\bar{d}$ ($B \rightarrow D^0\pi^0$ e.g.) have not yet been observed. Upper limits have been set which indicate suppression factors on the order of 10 relative to the non-color suppressed $B \rightarrow D^0\pi^-$ case, unlike the charm case. Plausible arguments have been offered which explain this situation as a result of the higher likelihood of a charm quark binding to an anti-charm to form a ψ than a charm quark binding to a light quark, although a coherent picture has not yet emerged.

Measurement of the Kobayashi-Maskawa matrix elements $|V_{cb}|$ and $|V_{ub}|$ The study of inclusive and exclusive semi-leptonic decays will provide the most precise determination of the CKM matrix elements $|V_{ub}|$ and $|V_{cb}|$. In addition, these parameters can be extracted from a variety of hadronic B-decay modes. Typically, the technique here is to determine the branching ratios in a variety of hadronic B-decay modes. Using the B-lifetime, now being measured with increasing precision at LEP, one can determine a partial width for each mode, which, when used with a particular model specifying form factors, etc., allows one to extract the fundamental coupling of the b-quark to the c-quark or the u-quark for exclusive modes. The KM-matrix element $|V_{ub}|$ can similarly be derived from observation of decays $D_s X$, where the D_s is presumed to couple directly to the W and X is some charmless particle, by combining the measurement of the branching ratio with the B-lifetime. ARGUS performed this exercise recently and determined a rather large value of $|V_{ub}|$, after searching in final states such as $D_s n\pi$. We have also searched in these modes, and, to date, have no signal at a level of sensitivity roughly twice the ARGUS value.

HQET (Heavy Quark Effective Theory) tells us that the KM-matrix element $|V_{cb}|$ can be extracted at maximum W-mass in the decay D^*W . If the factorization hypothesis holds, and the D^* spectrum is, in fact, relatively insensitive to the decay of the W, then one can also extract a value of $|V_{cb}|$ from the inclusive D^* spectrum in B-decay. If not, then we will be able to at least determine the limitations of the factorization hypothesis. This analysis is being actively pursued, and will be reported on at the APS meeting in Washington this spring.

3.7. INCLUSIVE PARTICLE YIELDS IN B-DECAY

Measurements of the inclusive production rates of baryons (protons, Λ_c 's, Σ_c 's, e.g.) and mesons (D , D^* , D^{**} , D_s , ϕ , e.g) are instrumental in helping us discriminate between models of B-decay. Within the last year, we have reported at the summer ICHEP conference the first observation of Σ_c production in B-decay. Asymmetries between Σ_c^{++} and Σ_c^0 production rates can give us vital information in determining the rates for external-W and internal decay processes.

3.8. CHARM PHYSICS AT CLEO

The charm physics program at CLEO-II has evolved along three parallel streams, concentrating on: hadronic decays of charmed mesons, hadronic decays of charmed baryons, and semileptonic decays of charmed particles. There are Purdue people actively working in the latter two areas.

Shipsey and his student, Miao, have begun a systematic study of charmed baryon semileptonic decay. Theoretically charmed baryon semileptonic decays are simpler than charmed meson semileptonic decays, because the latter can produce two ground states, K, K^* while the former can only produce a Λ . Charmed baryon semileptonic decay is therefore an ideal place to test HQET.

At a pragmatic level, in order to understand the decay mechanisms of the Λ_c , a reliable normalization is necessary. The decay most often used to normalize is the accessible $\Lambda_c \rightarrow pK\pi$. However, there is no reliable prediction for the rate of this decay and the experimental measurements differ by a factor of two.

$\Lambda_c \rightarrow X\Lambda\ell\nu$ offers the possibility of absolute normalization of all Λ_c decay modes because the inclusive leptonic decay of Λ_c can be calculated from the charm semi-leptonic decay width and the Λ_c lifetime to be $Br(\Lambda_c^+ \rightarrow \ell^+ x) = (3.4 \pm 0.4)\%$.

Using the Purdue computers, we look for $\Lambda\ell$ pairs of the appropriate sign and mass. A yield of 400 events on a small background see [Fig. 2]. corresponds to a cross-section of $4.72 \pm 0.26^{+0.52}_{-0.61} pb$. When combined with the measured cross-section for $\Lambda_c \rightarrow pK\pi$, we obtain the absolute normalization $Br(\Lambda_c \rightarrow pK\pi) = f(6.7 \pm 0.4 \pm 1.3)\%$, where f is the fraction of semileptonic decays accounted for. From charmed meson studies, $f = 0.6 \pm 0.2$.

Polarization of the Λ in $\Lambda_c \rightarrow \Lambda\ell\nu$ arises from non-trivial population of final states by the different helicity amplitudes. Solving the quadratic ambiguity, we observe, for the first time, a striking decay asymmetry of the Λ in this decay, shown in Fig. 3. Parity violation is near maximal with the sign arising from the V-A nature of the hadronic current.

Dr. Peining Wang began a study of the isospin mass splitting of the Σ_c while still a student. Several experiments have reported the production of Σ_c^{++} and Σ_c^0 . To date, only a single Σ_c^+ event consistent with the decay chain $\Sigma_c^+ \rightarrow \Lambda_c^+\pi^0$, $\Lambda_c^+ \rightarrow pK^-\pi^+$, has been reported from the BEBC bubble chamber experiment in 1980. In the past year, we have observed for the first time the Σ_c^+ baryon with statistical significance. Also we were able to make the first systematic measurement for the mass splittings between the Σ_c isospin states.

Many theoretical calculations which take into account both the effect of the intrinsic mass difference between the up and down quarks in the so-called heavy-light baryon system and the effect of electromagnetic interactions which consists of electrostatic Coulomb interactions between the quarks and spin-spin interactions, predict Σ_c isospin mass splittings ranging from $+18.0 MeV/c^2$ to $-6.5 MeV/c^2$. Several calculations also include the contributions of QCD effects. Different dynamical models of hadrons with a heavy quark can be tested from the measurement of the mass differences between hadrons that are members of the same isospin multiplet. Sources that play essential roles in determining the mass splittings between isospin multiplet states are the difference between the up quark and down quark masses and the electromagnetic interactions, which consists of electrostatic Coulomb interactions and

spin-spin interactions, between quarks. It has been suggested that the chromoelectric and chromomagnetic effects, such as the hyperfine mass splittings from the spin-spin interaction, may be important.

Here we present results on the observation of the Σ_c^+ baryon in nonresonant e^+e^- annihilation around $\sqrt{s} = 10.5$ GeV through the decay to $\Lambda_c^+\pi^0$. The data sample was collected with the CLEO II detector operating at the Cornell Electron Storage Ring (CESR) between 1990 and 1992. It consists of 1.48 fb^{-1} from the $\Upsilon(4S)$ resonance and from center-of-mass energies just below and above the $\Upsilon(4S)$ resonance.

We search for the Σ_c^+ charmed baryons in the decay mode $\Lambda_c^+\pi^0$, which is expected to dominate Σ_c^+ decays. We first search for the Λ_c^+ charmed baryons in the decay modes $\Lambda_c^+ \rightarrow pK^-\pi^+$, $\Lambda_c^+ \rightarrow \Lambda^0\pi^+\pi^-\pi^+$, $\Lambda_c^+ \rightarrow p\bar{K}^0$ and $\Lambda_c^+ \rightarrow \Lambda^0\pi^+$, where $\Lambda \rightarrow p\pi^-$, $K_s^0 \rightarrow \pi^+\pi^-$. To reduce combinatorial background, only combinations with $x_p > 0.5$ are used, where $x_p = p/p_{\text{max}}$, $p_{\text{max}} = \sqrt{E_{\text{beam}}^2 - m_{\Lambda_c^+}^2}$ and p is the momentum of the Λ_c^+ . Λ_c^+ baryons produced in B meson decays are kinematically limited to $x_p < 0.5$. Thus, when we form $\Lambda_c^+\pi^0$ combinations to search for Σ_c^+ , we exclude contributions from Σ_c^+ baryons from B meson decays. To reduce the background due to fake Λ_c^+ candidates, particle identification is used for all four decay modes of the Λ_c^+ . A combined weight for each hadronic particle type (π , K , p), is formed using information from both the dE/dx and (when available) TOF measurements, such that the sum of the three weights is normalized to one.

For the modes involving a K_s^0 (or a Λ^0), a secondary vertex is first reconstructed from a pair of oppositely charged tracks which intersect at a radial distance of more than 1 mm from the primary vertex. The pair is identified as a K_s^0 (Λ^0) candidate if the invariant mass of the pair when interpreted as $\pi^+\pi^-$ ($p\pi^-$) is within 3 standard deviations (3σ) of the measured K_s^0 (Λ^0) mass. The χ^2 , which measures how well the net momentum of the reconstructed K_s^0 (Λ^0) is extrapolated back to the beam position, is required to be less than 3.0.

A photon candidate is constructed from a shower cluster distributed over many CsI crystals. The photon candidate is excluded if it matches the projection of charged tracks into the calorimeter or has less than 50 MeV of energy deposited in the CsI crystals. Only isolated showers, which have no large energy deposition in nearby crystals, are used. The π^0 is reconstructed from a pair of photon candidates, each of which has $|\cos\theta| < 0.71$, where θ is the angle of the photon direction with respect to the beam direction. This corresponds to the highest resolution region of our calorimeter. The neutral pions are then selected by requiring that the $\gamma\gamma$ invariant mass be within 2.5σ of the known π^0 mass. A kinematic fit to the known π^0 mass is applied to the π^0 candidates.

From 2402 ± 103 Λ_c^+ candidate events, we obtain 111 ± 16 Σ_c^+ events centered at $(168.5 \pm 0.4 \pm 0.3) \text{ MeV}/c^2$. The first error is statistical and the second is systematic. We also find 126 ± 15 Σ_c^{++} events and 124 ± 17 Σ_c^0 events.

The mass difference $M(\Sigma_c^+) - M(\Lambda_c^+)$ is measured to be $(168.5 \pm 0.4 \pm 0.2) \text{ MeV}/c^2$. We also measure the mass differences $M(\Sigma_c^{++}) - M(\Lambda_c^+)$ and $M(\Sigma_c^0) - M(\Lambda_c^+)$ to be $(168.2 \pm 0.3 \pm 0.2) \text{ MeV}/c^2$ and $(167.1 \pm 0.3 \pm 0.2) \text{ MeV}/c^2$, respectively. The mass differences $M(\Sigma_c^{++}) - M(\Sigma_c^0)$ and $M(\Sigma_c^+) - M(\Sigma_c^0)$ are determined to be $(1.1 \pm 0.4 \pm 0.1) \text{ MeV}/c^2$ and $(1.4 \pm 0.5 \pm 0.3) \text{ MeV}/c^2$, respectively.

The mass splittings reported in this paper indicate that the Σ_c is the first baryon isospin multiplet for which the intrinsic light quark mass difference is not the dominant source of isospin mass splitting.

4. GAS MICROSTRIP DETECTORS

Gas Microstrip Detectors (GMD) offer an exciting alternative for high precision, low mass tracking at both the SSC and B Factories. GMD prototype chambers have already achieved 30 μm spatial resolution, a detector memory comparable to the SSC crossing time and MRad radiation hardness. A GMD has been operated successfully in experiment NA12 at CERN.

A new GMD R&D group lead by Ian Shipsey was formed at Purdue during Fall 1991. The group are proposing a GMD as an Intermediate Angle Tracker for SDC and as high precision tracking and triggering layers within the new drift chamber of the upgraded CLEO II. More detail can be found in the proposal.

During 1992 the Purdue group was the first in the U.S. to build a GMD entirely in-house from mask design and mask production through lithography and chemical processing to working detector. Results have been presented at SDC meetings, the Cornell CP Workshop and the Argonne Undergraduate Student Symposium. This work has been recognised by a National Science Foundation Young Investigator (NYI) Award to Ian Shipsey in 1992, and an SSC National Fellowship in 1993.

4.1. Mask Manufacture

The conventional method of manufacture of a GMD starts with the design of a mask. Using BRUTUS, a Purdue written software graphical interface package, we encode a pattern. One of our first designs is shown in Fig. 4. Twenty anodes and twenty cathodes, 10 μm and 60 μm wide respectively, with 240 μm pitch are OR'ed to two readout pads one for anodes the other for cathodes. This is done to minimise the number of electronic channels required. The active area of the GMD is 6 cm x 0.6 cm.

The pattern is compiled on the computer of an E-beam microfabricator in the School of Electrical Engineering at Purdue. E-beam lithography is the most accurate technique available for mask manufacture but is limited in the size of the mask that can be produced by the need for vacuum and commercially by the large market for small masks.

A Hoya Corporation glass blank metalized with chrome and subsequently coated in PMMA, an electron sensitive photo-resist, is used to make the mask. The computer steers the e-beam across the blank and combines this with a travelling stage to cover the mask area, typically 6" x 6". Where the e-beam strikes the PMMA, a chemical reaction is initiated making the PMMA preferentially soluble. Five separate detectors are drawn by the e-beam on each mask.

The PMMA is then developed (dissolved) to reveal the chromium layer underneath, which is in turn etched (dissolved) in phosphoric acid. Finally the unexposed PMMA is stripped (dissolved) to reveal, the negative image mask shown in Fig. 5. The quality of the Purdue mask is as good as most commercially available and can be produced for less than \$400 including the price of the blank. This should be compared to the commercial price of a similar sized mask which is in the range \$2,000-\$4,000.

4.2. Substrate preparation

Initially we have chosen glass as the substrate material. This is because it is available in the right resistivity range, it is smooth, making it easy to print on, and inexpensive. The two glasses we have used are Tempax, a borosilicate glass made by Schott Glaswerke in Mainz, Germany and Iron-Vanadium "Pestov" glass also made by Schott. These two glasses differ in resistivity by three orders of magnitude and while Tempax has been demonstrated to charge up Iron-Vanadium is not expected to charge up. Fe-Va is a candidate for the SDC Intermediate Angle Tracker.

All of the process steps are carried out by us using the facilities of the Electrical Engineering Solid State Labs. We metalize the glass with aluminum by vacuum thermal vapor deposition. We choose aluminum because it is inexpensive; however gold, being inert, is a better choice. The metal thickness is typically 2,000-3,000 Angstroms. Next, using a spinner we apply a 1 μm thick layer of negative photo-resist, such as Kodak 747R.

4.3. GMD production

The substrate and the mask are aligned and held together in vacuum by a Karl-Suss contact mask aligner to contact print a GMD using UV light. The UV light triggers the photo-chemistry of the resist. Subsequently the print undergoes three chemical steps: development of the PR, etching of the metal layer and stripping of the remaining PR. The result is a positive image detector, an example of which is shown in Fig. 6.

4.4. Performance of a Purdue GMD

The substrate, with five GMD's printed on it, is housed in a gas-tight metal box. The GMD is connected to readout electronics by wire bonding by hand a cable to the readout pads using a heat-cured silver epoxy. HV is applied to the anodes which are readout through an Ortec 142PC charge-sensitive pre-amplifier and pulse shaping amplifier. The detector is irradiated with an Fe^{55} source. The signal from the pre-amplifier and the induced signal on the neighbouring cathodes seen by a second 142PC pre-amplifier is shown in Fig. 7. The spectrum from the OR of twenty anodes is shown in Fig. 8 for a Tempax GMD and for a Fe-Vanadium GMD. Both of these spectra clearly show the 6 KeV Fe^{55} absorption peak and the 3 KeV Argon escape peak. The FWHM is about 20%; the gain is a few hundred. In the figure the tempax spectrum has been recorded with a LeCroy QVT. The Fe-Va spectrum has been recorded by a data acquisition system consisting of a Macintosh Quadra 700 interfaced by GPIB bus to a Kinetics GPIB/CAMAC crate-controller and a LeCroy 2259A peak sensing ADC. Gain and efficiency curves for Tempax and Fe-Va are shown in Fig. 9. Both substrates are found to have similar gain at the same operating voltage.

There are differences between the Tempax and Fe-Va however. Tempax shows a decline in gain at modest fluence with a time constant of minutes. If the HV is removed the detector recovers after a few hours. At a location some distance from the irradiated point the same decline in gain is seen even though there has been no irradiation at the second point. This global effect can be understood as arising from the motion of ions in the bulk of the glass. Fe-Va does not show this effect.

We have exposed the detectors to a flux equivalent to several years at the SSC using a Copper X ray tube. Tempax shows a clear charging phenomena at high gain To our surprise, and that of our SDC colleagues, Fe-Va appears to show the same charge-up behavior. Our results are shown in Fig. 10.

5. PROGRESS REPORT: PARTICLE ASTROPHYSICS

The Purdue particle astrophysics group is involved in projects pursuing the search for and study of astrophysical sources of very high energy photons. High energy charged cosmic rays have a well studied spectrum from a few GeV up to energies of 10^{22} eV and exhibit no anisotropy until near 10^{19} eV. Current models suggest that most cosmic rays of energies up to several Teravolts are accelerated through the interaction with shock waves generated by supernova explosions. Although there are only conjectures for sources and acceleration mechanisms for the higher energy regime, a reasonable picture may be forming from data obtained within the past two years. The essential ingredients are a high velocity magnetic field and method of confinement for an adequate period. Active Galactic Nuclei (AGN), which are presumed to be powered by supermassive black holes whose accretion mechanism generates a wide range of electromagnetic radiation are the greatest sources of concentrated energy on the galactic scale. In addition, certain types of AGN's exhibit powerful relativistic jets which recent results suggest are powerful particle beams.

The number of discrete galactic sources decreases dramatically with energy; a few dozen X-ray binaries, fewer MeV and GeV sources and, as yet, only the Crab Nebula at TeV energies. Extra-Galactic sources are even rarer. During the past two years, the EGRET detector on the CGRO satellite has recorded more than dozen discrete sources of GeV photons, some twenty of which are AGN. Early last year, the Whipple collaboration began observations on the AGN's detected by EGRET that were in the viewing range from Mt. Hopkins. A definitive signal was resolved from Markarian 421, the first TeV source ever reported at cosmic distances. The Mk 421 results are shown in Fig. 11 along with Crab data as a comparative monitor. The raw data rate was 172 events/minute (typical 3 Hz running) with a background subtracted signal rate of 0.3 γ 's /minute, corresponding to 0.9 σ /hour of statistical significance with a signal flux of 30% of that of the Crab, which typically yields 1 event/minute of signal. Mk 421 has a cosmological redshift of $z=0.03$, indicating a distance of 400 million light-years. At these distances, the optical depth due to absorption by infra-red photons is a strong function of z and can be used as a distance indicator. TeV gamma ray sources are rare objects; a few such objects constitutes a set of unique laboratories.

The Purdue group is a member of the Whipple Collaboration, based on Mt. Hopkins in southern Arizona. This collaboration utilizes two large telescopes (see Fig. 12) comprised of several hundred mirrors forming a tessellated pattern on the support structure. These telescopes are designed to record Cherenkov photons created by electromagnetic cascades in the upper atmosphere by incident relativistic cosmic rays. Photon initiated air showers are resolved by selections on the parameter space to which the characteristic elliptical-like images are fitted.

Currently, in addition to maintaining an extensive observing program, the focus of the Whipple collaboration has been to expand the capabilities for source detection by implementing stereoscopic viewing with the addition of a second large imaging telescope (the 11-meter). The 11-meter is about 142 meters from the original 10-meter and has been operational for about a year. The next step involves increasing the size of the 11-meter camera box from 37 to 91 photomultipliers, similar to the 10-meter, and to upgrade the data acquisition capabilities of both telescopes. With twin high resolution camera boxes, data from

coincident showers will allow source resolution to 2 arc-minutes, significantly more accurate than current X-ray and γ -ray instruments.

The first major step of the upgrade, implementing a new data acquisition system, was the responsibility of the Purdue group. This has been accomplished and is now controlling data acquisition on the 11-meter as of March 1993. This new system utilizes six pairs of control modules consisting of a CAMAC Crate Controller and List Processor, three at each telescope, on a common ethernet line. The controllers run as a dedicated task on a VAX computer which harvests the data and stores it on a disk. The List Processor reads off the data at back plane speed ($\sim 1\mu\text{s}/\text{word}$) and stores it in local memory for transfer to the VAX through the crate controller in an asynchronous mode. This system is capable of taking data at 50Hz with negligible dead-time. It is possible to log-in remotely over Internet and modify the details of data-acquisition. With this system, a high-level/low-level triggering scheme will be effected between the telescopes to lower the threshold energy for shower detection. Local data monitoring and data diagnostic checking will be implemented via a PC on the ethernet link; this is also the responsibility of the Purdue group and development is underway.

In addition to continued observing of the Crab Nebula and Markarian 421, we are taking data on newly discovered AGN's, some interesting Wolf-Rayet stars (very hot, massive stars which emit enormous winds) and several gamma ray binary systems. A limit on the TeV flux from Geminga, a nearby binary with a neutron star, which has been detected only at X-ray and γ -ray energies, has been established and has been submitted for publication; our result is in disagreement with DC fluxes reported by other groups.

The collaboration of T. R. Palfrey with the ASGAT (ground-based gamma ray astronomy at Themis in France) is continuing. During the last year he helped complete the analysis of the 1992 Crab data which has been accepted for publication. In addition he has carried out the simulations for the new focal-plane instrumentation of the telescopes which is to be installed in early 1993. He has also studied the usefulness of a projected imaging telescope of low collecting power and good angular resolution placed near the center of the current array. This latter is one piece of a larger plan by Saclay and the Ecole Polytechnique to build one or more large-area, high-resolution imaging Cherenkov telescopes.

In addition to our primary research with the imaging telescopes, we have designed and built an Air Cherenkov Array, an array of eight units, each consisting of four upward-facing photomultiplier tubes, as illustrated in Fig. 13. The photomultipliers in each box are required to form a coincidence within a narrow time window, whereas box coincidences are within a broader gate. This is a unique detector which samples a large solid angle and records Cherenkov showers with an energy threshold of $> 50\text{ TeV}$. Currently this array is collecting data during dark nights at a rate of 0.2 Hz. These triggers will be analysed early this summer to study shower resolution and possible coordination with the 10- and 11- meter telescopes.

In summary, during the past year we have discovered the first AGN in the TeV energy regime and have made excellent progress towards upgrading the telescopes of the Whipple collaboration. With continued interaction between the Whipple and EGRET observatories, a coherent picture of high energy cosmic rays and their acceleration mechanisms may be on the way to being established.

Fig. 2: The Λl invariant mass distribution.

Fig. 3: The angular distribution of protons in the Λ rest frame with respect to the Λ direction in the Λ_c rest frame.

Fig. 4: The anode cathode pattern of a Purdue GMD.

Fig. 5: A photo-montage of a Purdue chrome on glass mask showing the cathode and anode traces joining the readout pads.

Fig. 6: Purdue GMD produced by wet etch.

Fig. 7: A Purdue Tempax GMD. The upper pulse is the anode signal after an Ortec 142PC amplifier. The induced pulse on the neighboring cathode is the lower trace.

22598 II
Monday, November 23, 1992 22:10

Fig. 8: The Fe^{55} spectra obtained with a Purdue Tempax GMD and a Purdue Iron-Vanadium GMD are shown. FWHM is about 20%.

Fig. 9: Gain and efficiency curves for Tempax and Fe-Va.

Dose is $\sim 1.3 \text{ mC/cm}$

Fig. 10: High Flux X-ray exposure of a Tempax and Fe-Va GMD.

Fig. 11 Distributions in orientation angle for Markarian 421 and Crab Nebula data. The orientation angle, α , is the angle between the major axis of the shower image and the line from the source position in the image plane and the centroid of the image; this angle should peak near 0° .

Fig. 12:

the Air Cherenkov Array on Mt. Hopkins. Each of the eight units contains four photomultipliers. By incorporating programmable delay modules, a small area of sky can be selected for tracking. The 10-meter telescope is within this array.

Fig. 13:
Air Cherenkov
Array

Purdue University
Contract DE-AC02-76ER01428 (Task B)

A THEORETICAL HIGH ENERGY PHYSICS PROGRAM

within the

**DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY
West Lafayette, Indiana 47907-1396**

Supported by

U.S. DEPARTMENT OF ENERGY

Through

**Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana**

Current Contract Number: DE-AC02-76ER01428

TASK B

**ANNUAL PROGRESS REPORT
For May 1992 - April 1993**

Investigators: R. H. Capps, T. E. Clark, T.K. Kuo, S.T. Love

Research Associates: B. Haeri, H. Lew

Contract Renewal Proposal

for 1994

List of Personnel

R. H. Capps	(Professor)
T.E. Clark	(Professor)
T.K. Kuo	(Professor)
S.T. Love	(Professor)
B. Haeri	(Research Associate)
H. Lew	(Research Associate)

TABLE OF CONTENTS

I. Task B Overview	3
II. Individual Progress Reports	
R.H. Capps	3
T.E. Clark and S. T. Love	6
T.K. Kuo	9
B. Haeri	10
H. Lew	11
III. 1992 - 1993 Publication List	12

I. Task B Overview

During the previous year, our theory group has conducted investigations into a range of various research areas.

R.H. Capps has pursued his study to affine Kac-Moody algebras obtaining multiplicity formulas for a class of representations. He has also extended his analysis to construct representations of the quantum group $SU(3)_q$.

T.E. Clark and **S.T. Love**, in collaboration with a research associate **B. Haeri**, have studied the non-perturbative mass bounds imposed on scalar and fermion systems due to triviality and vacuum stability constraints. To achieve this, they constructed an exact Wilson renormalization group equation for the coupled scalar-fermion system, established the Gaussian fixed point and numerically integrated the solution in a local approximation.

T.K. Kuo has continued his investigations in resonance neutrino oscillations in matter involving three flavors of neutrinos. He has also explored further consequences of an $SP(6)_L \times U(1)_Y$ model with special emphasis on the properties of a new neutral gauge boson Z' .

During the past year we have conducted an active search for a new assistant professor in high energy theory. There have been several outstanding candidates and we hope to make an offer in the near future. At the conclusion of the current academic year, the research associate appointment of Dr. H. Lew will terminate. Dr. A. Papadopoulos, who has recently completed his Ph.D. at the University of California, Berkeley will join the group as a research associate. The appointment of Dr. B. Haeri has also been extended for the next academic year.

II. Individual Progress Reports

Richard H. Capps

A. Multiplicity Formulas for a Class of Representations of Affine Algebras

A common procedure for studying a finite Lie algebra of rank n is to choose n independent, commuting generators and to diagonalize them in every representation. The weight vector of a state in a representation is a vector in an n -dimensional Euclidian space, with components equal to the components of the commuting generators. These components are quantum numbers of the state. A root vector is a weight in the representation of the generators.

Dynkin defined a set of n integers that characterize each weight. A dominant weight is one such that none of these Dynkin components are negative.

Various reflection symmetries exist that allow the weights to be grouped into classes called Weyl orbits. For example, the six outer weights of the $SU(3)$ octet form an orbit. The multiplicity of a weight is the number of states with the same set of n quantum numbers. For example, the multiplicity of the center state of the $SU(3)$ octet is 2. The multiplicities of all the weights in an orbit must be the same. Since there is exactly one dominant weight in each orbit, the multiplicity structure of an irrep (irreducible representation) is specified by giving the multiplicities of all the dominant weights.

Affine Kac-Moody algebras are a type of infinite dimensional generalization of finite algebras. The properties discussed above apply to these algebras also, with one exception. An affine weight is specified by n Dynkin components, and one additional parameter called a null-depth parameter, which ranges through all the non-negative integers. Thus each dominant set of Dynkin components corresponds to an infinite sequence of dominant weights, corresponding to increasing null-depth. The multiplicities increase without bound along such a sequence.

In a previous paper I defined the width of an irreducible representation (a finite number) to be the number of dominant sets of Dynkin components.^[1] A multiplicity formula for the recursion sequence of all width-one irreps of all affine algebras was found that is simpler and faster to use than any formula found previously in the literature. The formula works equally well for twisted and untwisted affine algebras.

Last spring and summer a graduate student, Michael Lyons, and I finished work on the width-two irreps of the affine algebras. We listed all these irreps and were able to derive recursion formulas for the multiplicities that are simpler and faster than any in the literature. In the twisted cases, we constructed tables of the first 30 or 40 multiplicities, since these tables are not in the literature. The work has been submitted to Communications in Mathematical Physics. (The average length of time between submission and publication in this journal is more than a year.)

B. Representations of Quantum $SU(3)$

Quantum algebras, which are deformations of ordinary Lie algebras, have received a lot of attention in the last few years. The q -deformation of a real number x is denoted by $[x]_q$ and defined by

$$[x]_q = \frac{q^x - q^{-x}}{q - q^{-1}}$$

where q is a complex number. It is easy to show that as $q \rightarrow 1$, $[x]_q \rightarrow x$, so

that the deviation of q from 1 measures the amount of deformation. In the quantum algebras, some of the quantities in the commutation relations are replaced by their q -deformations.

Quantum algebras have played a role in many different physical problems, including deformed molecules and nuclei, anisotropic spin-chain hamiltonians, squeezed light, and knot theory.

The best understood of the quantum algebras is $SU(2)_q$. If the generators of ordinary $SU(2)$ are j_0 and $j_{\pm}$, the corresponding generators of quantum $SU(2)$ are denoted by J_0 and $J_{\pm}$. Curtwright and Zachos have exhibited explicit formulas for the matrices J_0 and $J_{\pm}$ for every irrep (irreducible representation).^[2] These are obtained by replacing some of the factors in the corresponding formulas for the $SU(3)$ generators $j_{\pm}$ by q -deformations.

I have been studying the question of specific representations for the generators of $SU(3)_q$. It is well-known that ordinary $SU(3)$ may be generated by two sets of $SU(2)$ generators, oriented at an angle of 120° in the two-dimensional space of hypercharge and t_0 (z component of isotopic spin). I will call these two sets of generators t -spin and u -spin, following the usage common in particle physics. The two diagonal quantum numbers are t_0 and u_0 .

Every irrep ψ of $SU(3)$ may be specified by a pair of non-negative integers (p,q) , where ψ is the largest irrep obtainable from the direct product of p quark irreps and q antiquark irreps. For simplicity I will assume that $p \geq q$. For some pairs of the quantum numbers (t_0, u_0) in an irrep, the state multiplicity may be greater than 1. The maximum multiplicity in an irrep is q (when $p \geq q$). It is known that when $q = 0$, and no multiplicities are greater than 1, a straightforward generalization of the above mentioned formulas of Curtwright and Zachos leads to specific forms for the generator matrices of $SU(3)_q$.

However, when the multiplicity of a state is greater than 1, the situation is more complicated. This is because a state of fixed total t -spin is a superposition of states of different total u -spins. One needs to specify the transformation matrices between the t -spin and u -spin bases to obtain a solution.

I have made two steps forward on this problem, for quantum $SU(3)$. First, I have developed two simple, but different, procedures for finding all these transformation matrices for any irrep of $SU(3)_q$. Second, I have found a general formula for all the transformation matrices in the cases $q = 1$. The matrix elements are all of the form $([a]_q[b]_q/[c]_q[d]_q)^{1/2}$, where a, b, c and d are integral expressions linear in t_0, u_0 , and p . I am in the process of writing

up these results.

References

1. Richard H. Capps, J. Phys. A.: Math. Gen. **23**, 4727 (1990).
2. T. Curtwright and C. Zachos, Phys. Lett. **234B**, 237 (1990).

T.E. Clark and S.T. Love

Of the various sectors comprising the Standard Model of particle physics, the one which is least constrained experimentally is the scalar sector which is responsible for the electroweak symmetry breaking as well as the fermion mass generation. The scalar self interactions and fermion-scalar Yukawa interactions generating this behavior are characterized by couplings which are not asymptotically free. Consequently, any conclusions drawn from a perturbative analysis of their short distance structure is at best indicative and require confirmation using nonperturbative techniques.

For instance, a study of the one-loop effective potential in the self coupled $\lambda\phi^4$ scalar model produces a Landau singularity which has in turn been used to place a perturbative triviality bound^[1] on the scalar mass. The qualitative nature of these conclusions has been largely substantiated by various nonperturbative studies.^[2-3] Here it is found that a nontrivial self coupled scalar model can only be consistently defined as a cutoff theory; that is, an effective description of the dynamics below some momentum scale Λ . The only consistent $\Lambda \rightarrow \infty$ limit of this model is one characterized by a vanishing renormalized coupling. This free field behavior is a reflection of the Landau singularity already observed in renormalization group improved perturbation theory. In general, the perturbative estimate is inadequate as it is being applied in a range outside the domain of validity of perturbation theory. If however, the scalar mass, m_H , is such that $m_H \ll \Lambda$, then the low energy value of the scalar self coupling is insensitive to its initial value due to the renormalization group trajectory focusing effects of the trivial (Gaussian) infrared fixed point. On the other hand, when discussing upper bounds on the scalar mass, one must consider values of m_H which are not much smaller than Λ itself. For instance for $\Lambda \sim 1$ TeV, the upper bound on m_H using perturbation theory^[1] is found to be $m_H \sim 800$ GeV. Certainly these quantitative results require substantiation in a nonperturbative framework.

In fact it turns out that various such estimates^[2-3] have been obtained yielding a maximum scalar mass of $m_H \sim 650-800$ GeV again for $\Lambda \sim 1$ TeV.

This rough quantitative agreement of these nonperturbative predictions with the perturbative estimate is certainly not a priori guaranteed. In addition to the fact that the scalar self coupling is sufficiently large to render the 1-loop perturbative result suspect, it also follows that such a coupling induces additional irrelevant operators whose effects are not necessarily insignificant for scales not too far removed from Λ . In general, the coefficients of these operators are suppressed by powers of m_H/Λ . While for $m_H \ll \Lambda$ such terms are clearly quite small, this need not be the case as m_H approaches Λ .

Of the various nonperturbative studies which have been performed to study the critical dynamics of these scalar self coupled systems, most have employed lattice Monte Carlo simulations.^[2] A somewhat different approach^[3] to the study of these systems employs the continuous Wilson renormalization group.^[4] In this case, one specifies the model at the ultraviolet (UV) cutoff scale Λ and, by demanding the cutoff independence of the renormalized Green functions, one constructs a functional differential equation for the action at the longer distance scale. This action contains not only the relevant and marginal operators but also the irrelevant operators with the coefficient of each operator fixed in terms of the initial action defined at scale Λ . Moreover as the construction is independent of the strength of the initial coupling, it constitutes a nonperturbative procedure. Using a local action approximation to these exact functional equations, an upper bound for the scalar mass consistent with the other nonperturbative estimates has been extracted.^[3] Recently, the sensitivity of this bound to the choice of action at scale Λ has also been investigated.^[5] The resulting maximum scalar mass was found to be insensitive to the initial action.

In collaboration with one of our research associates, B. Haeri, we extended the Wilson renormalization group approach to the construction of renormalized effective Lagrangians in theories containing fermions in addition to scalars.^[6-7] We established the existence of the Gaussian fixed point so that a consistent interpretation of the model again requires the introduction of an ultraviolet cutoff Λ . For scalar and fermion masses not too far removed from this UV cutoff, some nonperturbative tools must be employed. We use the continuous Wilson renormalization group to construct a functional differential equation for the action functional which can serve as just such a nonperturbative procedure. To obtain this equation, one varies the UV cutoff and demands the resultant action functional reproduces the same physics (i.e. renormalized Green functions) on all momentum scales less than the new (lower) cutoff. That is, the change in cutoff is compensated by changing the coupling constants of a complete set of local operators. This set includes

irrelevant operators in addition to the relevant and marginal operators. Thus the various action functionals satisfying this equation correspond to points on a particular (Wilson) renormalization group flow each producing the same physics on momentum scales less than the UV cutoff used in its definition.

This exact Wilson renormalization group equation is tantamount to an infinite number of coupled consistency equations for the various coupling constants of the complete set of operators. As such, its analysis and extraction of physical results from it requires additional approximations. We employ a local action approximation in terms of which the functional differential equation reduces to a partial differential equation for a generalized effective potential in a model containing one scalar and one fermion. In the limit of small scalar self coupling and Yukawa coupling at the UV cutoff scale, this equation reduces to the familiar 1-loop perturbative renormalization group equations for these couplings. For larger couplings at the UV cutoff scale, however, the equation for the generalized effective potential requires a numerical evaluation. We perform such an analysis under the further assumption that only terms up to bilinear in the fermion fields are retained in the generalized potential. The resulting numerical solutions are then analyzed for various choices of the initial Yukawa and scalar self couplings and for different initial cutoffs. In particular, the role of the induced irrelevant operators in driving the theory toward the trivial (Gaussian) fixed point is emphasized. The exact numerical solutions are also contrasted with the analytic solutions of the one loop perturbative equations extrapolated well outside their domain of validity.

By examining the numerical solutions resulting from different values of the scalar self coupling and Yukawa coupling chosen at various cutoff scales and resulting in nontrivial scalar vacuum expectation values which spontaneously breaks a discrete symmetry forbidding an explicit fermion mass term, we obtain the triviality and vacuum stability bounds on both the scalar and fermion masses. Our results indicate that the scalar mass bound increases in the presence of the Yukawa coupling. The amount of this increase depends on the strength of the initial scalar self coupling; larger self couplings yielding smaller percentage increases. For the range of couplings we considered, the percent increase in the scalar mass bound varied from roughly 15-20% to 6-10%. In addition, using the values of the mass upper bounds as a function of the cutoff, we determine the allowed ranges of scalar and fermion masses for a specific choice of the cutoff.

References

1. R. Dashen and H. Neuberger, Phys. Rev. Lett. **50** (1983) 1897; M.A.B. Beg, C. Panagiotakopoulos and A. Sirlin, Phys. Rev. Lett. **52** (1984) 833.
2. J. Kuti, L. Lin and Y. Shen, Phys. Rev. Lett. **61** (1988) 678; Nucl. Phys. (Proc. Suppl.) **4** (1988) 397; Y. Shen, J. Kuti and S. Meyer, *ibid.* 216; M. Lüscher and P. Weisz, Nucl. Phys. **B318** (1989) 705; G. Bhanot, K. Bitar, U. Heller and H. Neuberger, Nucl. Phys. **B353** (1991) 551.
3. P. Hasenfratz and J. Nager, Z. Phys. **C37** (1988) 477.
4. K.G. Wilson, Phys. Rev. **B4** (1971) 3174; **B4** (1971) 3184; F.J. Wegner and A. Houghton, Phys. Rev. **A8** (1973) 401. For reviews, see W.G. Wilson and J.B. Kogut, Phys. Rept. **12C** (1974) 75; F.J. Wegner, in *Phase Transitions and Critical Phenomena*, ed. by C. Domb and M.S. Green (Academic Press, New York, 1976) Vol. 6, p. 8; S. Weinberg, in *Proceedings of the 1976 International School of Subnuclear Physics, Erice*. See also, J. Polchinski, Nucl. Phys. **B231** (1984) 269.
5. R. Akoury and B. Haeri, *The Cutoff Dependence of the Higgs Mass in an Extension of the Standard Model*, PURD-TH-92-05, UM-TH-92-09.
6. T.E. Clark, B. Haeri and S.T. Love, *Wilson Renormalization Group Analysis of Theories with Scalars and Fermions*, PURD-TH-92-09, to be published in Nucl. Phys. **B**.
7. R. Akhoury, T.E. Clark, B. Haeri and S.T. Love *Cutoff Dependence of the Higgs Mass*, PURD-TH-92-13, to appear in the *Proceedings of DPF'92*, Fermilab (World Scientific).

T. K. Kuo

A. Extensions of the Standard Model

We have been investigating the ramifications of a model proposed earlier. [1] This is a gauge theory which unifies the three generations so that the usual $SU(2)_L \times U(1)_Y$ group is extended to $SP(6)_L \times U(1)_Y$. A characteristic feature of this model is that there are no extra fermions, but it does have quite a few extra gauge bosons. One of the most interesting possibilities

is the existence of a new, light ($\lesssim 1$ TeV), neutral gauge boson Z' . The coupling strengths of Z' are completely fixed by the theory. It does not obey universality and can cause lepton flavor violating decays, such as $Z^0 \rightarrow \tau\mu$. This process was in fact first suggested in this context. We are now investigating the various processes which may be sensitive to the presence of Z' . [2] They include universality violations, asymmetries, and the process $e^+e^- \rightarrow W^+W^-$, all with respect to LEP.

B. Neutrino Oscillations in Matter

Resonance neutrino oscillation (the MSW effect) offers a very attractive solution to the solar neutrino problem. The usual analysis, however, is done with the simplifying assumption of only two flavors of neutrinos. The result suggests that neutrino parameters are constrained to lie in two very small regions. We have studied the problem with three flavors, using a formalism developed earlier. [3]. It is found [4] that the allowed parameter regions are distorted and enlarged considerably when compared to the results of the two flavor analysis. Thus, a lot more experimental inputs are needed before we can pin down the neutrino parameters. In particular, the design of new experiments should not be based on the two flavor analysis, which may well be off the mark.

References

1. T. K. Kuo and N. Nakagawa, P. R. **D31**, 1161 (1985); A. Bagnoid, T. K. Kuo, and N. Nakagawa, Int. J. of Mod. Phys. **2**, 1351 (1987).
2. G. Park and T. K. Kuo, P. R. **D45**, 1720 (1992); Zeit. f. Phys. C., to appear (1993); T. K. Kuo, G. Park and M. Zralek, preprint (PURD-TH-93-1), (1993).
3. T. K. Kuo and J. Pantaleone, Rev. Mod. Phys. **61**, 937 (1989).
4. D. Harley, J. Pantaleone and T. K. Kuo, Phys. Rev. **D**, to appear, (1993); preprint (IUHET-246), (1993).

B. Haeri

During 1992 my research was partly aimed at continuing development of Wilson renormalization group methods. In collaboration with R. Akhoury I calculated the absolute upper bound of the Higgs mass in the context of a scalar field theory in which the Higgs particle is coupled to a $SU(2) \times U(1)$

scalar singlet. We used perturbation theory to perform the calculation of the Higgs mass, and thus give a comparison with the Higgs mass upper bound found previously using non-perturbative solutions of the Wilson renormalization group equations. The main advantage of the Wilson renormalization group is the possibility of predicting an absolute upper bound surpassing 1 TeV, a region where perturbation theory breaks down. We find though that the absolute upper bound on the Higgs mass remains below 1 TeV (about 800 GeV), when considering a simple extension of the standard model involving the addition of a scalar singlet coupled to the Higgs.

I also worked on developing a model to explain why the top mass is potentially close in value to the electroweak scale. The model introduces a new interaction for some members of the third generation of quarks and leptons, strong enough to form massless condensates involving the top quark. These massless boundstates are the Nambu-Goldstone particles necessary to dynamically break electroweak symmetry. The new interactions dynamically generate a top quark mass on the order of the electroweak scale in the process of forming the Nambu-Goldstone particles.

H. Lew

In the past year my research involved neutrino physics, the problem of electric charge quantization, further investigation of various aspects of quark-lepton symmetric models and a study of the interplay between particle physics models and the standard hot big bang model of cosmology. The specifics of this research are outlined in the following:

(i) *Neutrino physics* [1]

It is possible that the tau lepton may have a longer lifetime than that predicted by the standard model. The simplest extension to the standard model incorporating a longer tau lifetime involves the addition of a gauge singlet Weyl fermion. We considered the most general model of this kind and evaluated the experimental constraints on the various parameters of this model. It was shown that the model can be made consistent with the standard cosmology model for a range of parameters. We then examined possible signatures of the model in certain experiments searching for heavy neutral leptons.

(ii) *Electric charge quantization* [2]

Experimentally it has been known for a long time that the electric charges of the observed particles appear to be quantized. An approach to understanding electric charge quantization that can be used for gauge theories with explicit $U(1)$ factors – such as the standard model and its variants – was

reviewed and discussed. This approach uses the allowed invariances of the Lagrangian and their associated anomaly cancellation equations. We demonstrated that charge may be de-quantized in the three-generation standard model with massless neutrinos, because differences in family-lepton-numbers are anomaly-free. We also reviewed the relevant experimental limits. This approach to charge quantization suggests that the minimal standard model should be extended so that family-lepton-number differences are explicitly broken. Some candidate extensions (e.g. the minimal standard model augmented by Majorana right-handed neutrinos) were examined.

(iii) *Quark-lepton symmetry and the cosmological domain wall problem* [3]

Quarks and leptons may be related to each other through a spontaneously broken discrete symmetry. Models with acceptable and interesting collider phenomenology have been constructed which incorporate this idea. However, the standard Hot Big Bang model of cosmology is generally considered to eschew spontaneously broken discrete symmetries because they often lead to the formation of unacceptably massive domain walls. We showed that there are a number of plausible quark-lepton symmetric models which do *not* produce cosmologically troublesome domain walls. We also considered some interesting questions concerning anomalous discrete symmetries.

(iv) *Shadow matter* [4]

The cosmological and astrophysical implications of a shadow matter model which could also have interesting experimental consequences were examined. The model has identical microphysics for both the ordinary and shadow worlds but requires a macroscopic asymmetry from nucleosynthesis constraints. It was proposed that this macroscopic asymmetry can be generated at the quark-hadron phase transition.

(v) *Baryogenesis and domain walls* [5]

It has been recently speculated that global symmetries are broken by gravity. We proposed a scenario for the generation of the baryon asymmetry in the early Universe in which the domain walls predicted by theories with discrete symmetries become unstable due to these Planck scale effects. The relative motion of the decaying walls can provide a mechanism for the departure from thermal equilibrium necessary to have a nonvanishing baryon asymmetry. In particular, we examined this idea within the frameworks of the Left-Right and Quark-Lepton symmetric models.

References

1. R. Foot and H. Lew, *Mod. Phys. Lett. A* **7**, 2921 (1992); *Phys. Rev. D* **47**, 2011 (1993).
2. R. Foot, H. Lew and R.R. Volkas, *J. Phys. G* **19**, 361 (1993).
3. H. Lew and R. R. Volkas, *Phys. Rev. D* **47**, 1356 (1993).
4. H. Lew, "A Shadow Matter Universe", Purdue University Report No. PURD-TH-93-04.
5. H. Lew and A. Riotto, "Baryogenesis, Domain Walls and the Role of Gravity", Purdue University Report No. PURD-TH-93-06.

III. Publications (May 1992 - April 1993)

1. W.A. Bardeen and S.T. Love, *Effective Lagrangian for Scale Invariant Quantum Electrodynamics*, *Phys. Rev. D* **45** (1992) 4672-4680.
2. M. Carena and C.E.M. Wagner, *Bound State Mass in Quenched QED*, *Phys. Lett. B* **285** (199) 277-284.
3. R. Foot and H. Lew, *Implications of a Model with a Longer Tau Lifetime*, *Mod. Phys. Lett. A* **7** (1992) 2921-2930.
4. R. Foot and H. Lew *Possible Connection Between a Longer Tau Lifetime and the Tau Neutrino Mass*, *Phys. Rev. D* **47** (1993) 2011-2103.
5. R. Foot, H. Lew and R.R. Volkas, *Glueball Mediated Decay of Massive Neutrios in Quark-Lepton Symmetric Models*, *Int. J. Mod. Phys. A* **8** (1993) 983-1003.
6. R. Foot, H. Lew and R.R. Volkas, *Electric Charge Quantization*, *J. Phys. G* **19** (1993) 361-372.
7. H. Lew and R.R. Volkas, *Discrete Quark-Lepton Symmetry Need Not Pose a Cosmological Domain Wall Problem*, *Phys. Rev. D* **47** (1993) 1356.

Preprints (submitted for publication)

1. D. Harley, T.K. Kuo and J. Pantaleone, *Solar Neutrinos with Three Flavor Mixings*, PURD-TH-92-08, to be published in Phys. Rev. D.
2. T.E. Clark, B. Haeri and S.T. Love, *Wilson Renormalization Group Analysis of Theories with Scalars and Fermions*, PURD-TH-92-10, to be published in Nucl. Phys. B.
3. T.K. Kuo and G. Park, *Test for Heavier Gauge Bosons in the $Sp(6)_L \times U(1)_Y$ Model*, PURD-TH-92-15, to be published in Zeit. f. Phys. C.
4. H. Lew, *A Shadow Matter Universe*, PURD-TH-93-04.
5. H. Lew and A. Riotto, *Baryogenesis, Domain Walls and the Role of Gravity*, PURD-TH-93-06.
6. D. Harley, T.K. Kuo and J. Pantaleone, *The Solar Neutrino Problem: Implication of Three Flavor Mixings*.
7. R.H. Capps, *Weight Systems of Affine Kac-Moody Algebras and Coxeter Diagram Symmetries*.
8. R.H. Capps and M.A. Lyons *Multiplicity Formula for a Class of Representations of Affine-Kac-Moody Algebras*.

Conference Proceedings (to be published)

1. R. Akhoury, T.E. Clark, B. Haeri and S.T. Love, *Cutoff Dependence of the Higgs Mass*, PURD-TH-92-13, to appear in the *Proceedings of DPF'92*, Fermilab 1992.
2. R. Foot, H. Lew and R. Volkas, *Quark-Lepton Symmetry*, PURD-TH-92-14, to appear in the *Proceedings of DPF'92*, Fermilab 1992.
3. T.K. Kuo, *Neutrino Oscillations in Matter*, to appear in the *Proceedings of the Conference on Realistic Nuclear Structures*.

AN EXPERIMENTAL HIGH ENERGY PHYSICS PROGRAM

within the

DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY

West Lafayette, Indiana 47907

Supported by

U.S. DEPARTMENT OF ENERGY

Through

Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana

Current Grant Number: DE-FG02-91ER40681

TASK D

PROGRESS REPORT
for the period ending
May, 1993

Principal Investigators: D. D. Carmony and L. J. Gutay.

Spokesperson: L. J. Gutay (317) 494-5047 FTS and Commercial

PROGRESS REPORT
for the period ending
May, 1993

TABLE OF CONTENTS

	page
I. The L3 Detector to Study e^+e^- Collisions at CERN.....	3
II. The C0 Experiment to Study $\bar{p}$ -p Collisions at FNAL.....	15
III. Accelerator Physics at Fermilab.....	16
IV. Development Work for the SDC Detector at SSC.....	17
V. Publications since 1985.....	21

I. The L3 Detector to Study e^+e^- Collisions at CERN

1.1 Muon Chamber Resolution Improvement

The L3 experiment is taking data at $\sqrt{s} = 100$ GeV/c at the present in the LEP1 tunnel. According to CERN plans, in 1994 there will be a changeover to LEP2 with $\sqrt{s} = 200$ GeV/c. In order to retain the presently attainable momentum resolution of 1.5% without increasing the magnetic field, at twice the center of mass energy, the chamber resolution must be improved and have to be reduced.

In order to obtain a compact, modular, high resolution chamber with an accurate wire positioning, a design was chosen where the sense and field wires alternate with each other with a 4.5mm separation. Due to the closeness (4.5 mm) of the neighboring wires, they are coupled inductively and signals on sense wires distorted.

To correct for induction effects, we measured the time dependence of the pulse shapes for all 16 sense wires for a given track, by a 100 Mh Flash ADC readout system. The coordinates were fitted to a track segment and residual errors determined. The residual distributions were then compared before and after the correction was made.

Experimental Setup

The chambers of Octant-17 and the triggering scintillators were built by the L3 collaboration. The readout electronics and triggering system were built by the Purdue group.

A cosmic ray trigger came from the coincidence of scintillator arrays located above and below the octant. To eliminate soft muons two layers of lead bricks were interposed between the chamber and the lower set of scintillators. This arrangement is shown in Fig. I-1.

The pulses on the wire were readout and stored by our FADC system. It digitized the pulse every 10 ns. The digitized and stored pulse was read by our VME system. It preanalyzed the data and wrote it on magnetic tape.

Results:

The resolution studies were initiated during 1991. The software for multi-microprocessor VME data acquisition has been implemented and the Flash ADC readout system designed. During December 1991 the 17-th octant has been instrumented, the readout system installed and the first run completed.

We analysed the data at Purdue during first months of 1992. The most important conclusions of that analysis are summarized below:

1. the between-wires induction and cross talk contribute substantially to overall chamber resolution. The correction for these effects is possible. It requires detailed studies of pulse shape dependence on pulse height and drift distances.
2. the algorithm, which associates a drift distance with the earliest charge arrival time, produces the best resolution.
3. the multiple scattering of cosmic particles dominates the resolution. A method of eliminating low energy spectrum must be developed.
4. due to the fast rise of the sense wire pulse the 10 ns measurement repetition time of our Flash ADCs is not adequate for detailed studies.

For the next run, conducted during May, 92, a number of improvements have been made. We installed a pulse splitter on selected cells: the same signal was fed into two FADC units with relative delay of 5 ns. By doing so the effective FADC frequency has been increased to 200 MHz. The lead absorber

placed between the chambers and the lower trigger scintillator, in conjunction with the time-of-flight measurement, allowed us to eliminate muons with energy below ≈ 600 MeV.

In the data analysis we concentrated on investigating the induction and cross-talk effects and their influence on the spatial resolution of the chamber.

A correction algorithm has been developed. When it was applied to FADC data, it led to substantial resolution improvement. We have also shown, that our method can successfully be applied to L3 experimental data in spite of the fact, that in the latter case the pulse shape information is not available.

Fig. I-2a shows signals on 5 wires randomly selected from the middle of a cell. A track, fitted to all 16 hits, and residual errors are also shown. A small depression before each hit, seen as a lowering of the base line, is due to induced signals of opposite polarity. The effect of these induced signals is to delay the time at which signals cross a threshold.

The correction algorithm is based on the assumption, suggested by modeling calculations, that the avalanche on a given sense wire generates induced signals on all other wires in proportion to its own amplitude. In a first approximation the proportionality parameters depend only on a relative distance between the wires.

The observed amplitude D_i on the i -th sense wire can be corrected by subtracting from the genuine amplitude A_i , solely due to the avalanche on the wire, all induction and cross-talk signals generated on the wire by avalanche signals on all other sense wires.

On the i -th sense wire the sum of induced signals from remaining sense wires is equal to

$$D_{\text{sense}}^i = \sum_{k=1}^n a_{k,i} * A_k$$

where $a_{k,i} * A_k$ is a proportion of the signal A_i on wire k induced on the i -th wire, and $a_{i,i} = 0$.

The signals induced on field wires, which are connected together, are grounded through large capacitors. Because of capacitive coupling to ground some amount of cross-talk can not be avoided: the potential of the field wires changes by a fraction of the sum of the signals induced by all sense wires on all field wires

$$D_{\text{field}} = \sum_{i=1}^n \sum_{k=1}^{n+1} b_{k,i} * A_i$$

where k and i enumerate field and sense wires - respectively, and b are sense-to-field induction parameters.

So the observed amplitude D_i is equal to

$$D_i = A_i - D_{\text{sense}}^i - c * D_{\text{field}}$$

where c is a cross-talk factor.

The problem of correcting n amplitudes recorded on n wires is equivalent to solving a linear equation

$$\vec{D} = \alpha * \vec{A}$$

provided, the elements of the correction matrix α (expressed by a , b and c parameters) are known.

The procedure is repeated for each FADC time channel (up to 256).

In order to reduce the number of parameters we took into account the contributions from first- and second-neighbor wires only. Within this assumption we were able to determine the unknown parameters from the data

themselves by a fitting procedure. In the fit, the baseline preceding a hit on each wire by 20 ns (≈ 1 mm) is required to be equal to 0 within baseline-noise accuracy.

The effect of the correction on individual FADC time spectra is illustrated on Fig. I-2b. By comparing with Fig. I-2a one can notice, that a dip before the hits has been eliminated and the signal amplitudes have increased.

The overall resolution is shown on Fig. I-3, where the results for uncorrected and corrected data are compared.

Fig. I-4 shows a systematic error in an average track position as determined by comparing the corrected and uncorrected data.

1.2 Improving Muon Acceptance and Momentum Resolution in the L3 Muon Tracker

In the L3 muon chambers the sense, field, guard and mesh wires of the unit cell occupy a common volume. The breaking or loosening of a wire may require lower voltages of several wires in the cell which contains the bad wire. In order to maintain good muon momentum resolution, bad cells and neighboring cells with reduced high voltages have to be left out of the muon momentum analysis. The left out cells and thus rejected tracks, however, became substantial in number with time. Hence, it had become imperative to change the approach to bad cells. Thus, the Purdue group initiated a study of the first order variation of the electric field in a drift cell as a function of wire voltages both inside the cell and neighboring cell.

It was assumed that the decrease in momentum resolution in the bad cell and in its neighboring cell results from the deviations in the electric field from its nominal value and changes in gain at the sense wire. To calculate

the actual field value, it was assumed that the field variations are linear to the voltage variations.

$$E = E^0 - E^0 [\sum a_i (V_i^0 - V_i)]$$

Using the program WIRCHA, 276 combinations of V_i values were used to calculate corresponding E values. All 276 combinations were used in a least squares fit to determine the a_i 's in the above expression. By using this analytic expression on cases with small gain changes, we were able to recover to cases where $|V_i^0 - V_i| \ll V_i^0$. This resulted in a 30% increase of the accepted muon sample with good resolution.

1.3 Ongoing Physics Analysis by the Purdue Group at L3

After the 1992 run of LEP, over 1 million Z^0 decays were recorded by the L3 detector the Purdue group has been pursuing topics which exploit this large sample of Z^0 's for interesting physics. Our analysis has also concentrated on physics topics which rely upon muon identification and so capitalizes on our expertise with the L3 muon chambers.

Using the inclusive lepton sample of Z^0 decays we have published in the past year a paper on the forward-backward asymmetry of $Z^0 \rightarrow b\bar{b}$ events, $A_{b\bar{b}}$, and a measurement of mixing for both B_d and B_s mesons. We have measured $A_{b\bar{b}}$ to be $0.086 \pm 0.015_{\text{stat}}$. From this we have been able to derive a value for $\sin^2 \bar{\theta}_W$ after taking QED and QCD corrections into account of 0.2336 ± 0.0029 , a precision of 1% which is competitive with $\sin^2 \bar{\theta}_W$ derived from purely leptonic decays of the Z^0 .

Since LEP has copious production of B_s mesons our measurement of mixing is a measurement of B_d and B_s mixing combined. This allows us to determine a direct measurement of B_s mixing by combining our result with a value for B_d mixing determined at machines operating below B_s production threshold. Current values of B_d mixing favor, in the Standard Model, maximal mixing for

the B_s meson, hence, deviations from this would be indicative of new physics. We have published on data from our 1990 and 1991 running periods a measured value of $\chi_B = .121 \pm .017(\text{stat.}) \pm .006(\text{sys.})$. We can infer a value for B_s mixing by using the CLEO/ARGUS value for B_d of $\chi_d = .16$ giving with our measurement $\chi_s = .46 \pm .21$, a value consistent with maximal mixing, i.e., $\chi_s = 0.5$.

Recently, we made a search for events containing the as yet undiscovered B_c meson, a bound state of the heavy b and c quarks whose production would be energetically accessible at LEP. It has been estimated that in a sample of 10^6 Z's from 10 to 100 B_c mesons might be produced. An event with a decaying B_c would have a dramatic signature as a predominately 3-jet event with few decay products. We looked at events with three highly energetic leptons in the same hemisphere, in particular decays of the B_c meson which would have a lepton from semi-leptonically decaying b quark and two leptons from decay of J/ψ . Some interesting events were found, however, there are also several background processes to contend with. We plan to continue to look further at this for the upcoming LEP run.

The Purdue group is also involved in searching for lepton flavor violating decays of the Z^0 . Such decays are forbidden in the Standard Model, however, theoretical extensions provide for branching ratios for $Z^0 \rightarrow \mu\tau$ as high as 10^{-4} and would be within reach with our statistics. The high momentum resolution of L3 is key to removing background coming from decays of $Z^0 \rightarrow \tau^+\tau^-$. Our preliminary result gives a 95% CL limit of 2.9×10^{-5} for the process of $Z^0 \rightarrow \mu\tau$.

We have continued to work on $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$ events looking at the parameters of cross section and forward-backward asymmetry. Both of these parameters have interest as an input to an indirect determination of the top mass. For the 1992 running period the preliminary value of $\sigma_{\mu+\mu-}$ is

1.46 nbarn $\pm .01$ stat $\pm .01$ sys. The work on the muon chamber high voltage and dead cell map that our group provides is proving to be an increasingly important part of this analysis. The low systematic error on the above cross section measurement has been obtained by accurately identifying and correcting or removing sub-nominally operating drift cells. Over last years running period the number of disconnected channels has risen over 2-fold decreasing the acceptance of the muon chambers by 7.5%.

We studied the feasibility of improving the 2.5% muon momentum resolution by analytically removing cross-talk effects in the reconstructed muon track hits. It was found that for 45 GeV muons the single wire resolution could be improved by upwards of 20% which translates into a 10% improvement on momentum resolution for the entire track.

1.4 Silicon Microvertex Detector for L3

In order to obtain an accurate interaction vertex and muon tracking a Silicon Microvertex Detector (SMD) was developed. The alignment of the SMD was done with lasers. It has been installed and T. Adam finished the laser alignment.

GOOD event Example: no combining

Fig. I-1

Ev.-15100141 : Fit without cross-talk correction

Fig. I-2a

Ev.-15100141 : Fit after cross-talk correction

Fig. I-2b

Fig. I-3

Fig. I-4

II. The C0 Experiment to Study $\bar{p}$ -p Collisions at FNAL

It was suggested by C.N. Yang that multiparticle production in e^+e^- and $\bar{p}$ -p collisions are related to each other. He predicted that the width of the backward-forward distribution as function of multiplicity are similar in $\bar{p}$ -p and e^+e^- collisions. A preliminary comparison of our FNAL and L3 data contradict this prediction.

In our FNAL data, we found that the average transverse momentum $\langle p_t \rangle$ sharply increases with multiplicity for heavy particles. We studied the L3 data and found the same effect.

Finally, there is the great theoretical interest in transverse flow in $\bar{p}$ -p high multiplicity collision. We have shown that there can be a common flow velocity for π , K and $p^\pm$ at Fermilab energies. There is an indication of reaching a new threshold above $\sqrt{s} = 1$ TeV.

III. Accelerator Physics at Fermilab

To enhance Fermilab's research potential, FNAL has embarked on a \$240 million upgrade program. This program hopes to achieve three major goals:

1. Eliminate the interference between the collider detectors and the main ring.
2. Possibility of simultaneous operation of collider and limited fixed target program.
3. To increase the collider luminosity.

The major obstacle in intensity improvement is the transfer of the Linac beam into the Main Injector.

Strong coupled-bunch mode oscillation and bunch length increase during acceleration have been observed since the construction of the Fermilab Booster synchrotron. These effects lead to emittance growth and beam losses. These instabilities increase with intensity thus they are a limiting factor to obtain higher luminosity. To eliminate this quarter of a century old problem was Kathy Harkay's thesis project.

We had a hunch that the wake field of a bunch passing through a resonant cavity generate a parasitic resonance mode which adversely affects the motion of subsequent bunches. Kathy and Colestock of Fermilab developed an ingenious opto-electronics technique to excite cavity and found some parasitic resonances. After intuitive arguments she picked out some of the unwanted modes, built a damper and tested it in the beam on a few cavities. Finally, the dampers were installed in all seventeen cavities. The result was a smashing success. The bunch oscillation was removed and the bunch with increase was slowed down. This is a remarkable result with world wide benefits.

IV. Development Work for the SDC Detector at SSC

Task D is member of the SDC muon group. Earlier, we had worked on crimping stability and determination of elastic limits of wires used in the muon drift tubes. Our recent work has centered on testing the performance of drift tube and anode wire tensioning device.

Historically, the wire tensions of drift chambers have been measured using external magnetic field and/or mechanical method of excitation. However, in some experimental situations the conducting tube that holds the tensioned anode wire is inside a large solid structure, and these methods can not be used. Such is the case of the muon tracker chamber system planned for the SDC detector at SSC. In the SDC detector the unit drift cell is a cylindrical aluminum tube with two strips of field shaping electrodes, which are parallel with each other and insulated from the aluminum tube. The anode wire is located along the cylindrical symmetry axis. Thus two halves of the tube are mirror images of each other. The plane of symmetry is parallel to the field shaping electrodes and runs through the axis of the cylinder. The tubes are glued into towers of large rigid structures. The tubes have to provide 20 μm track location accuracy for the 10 year lifetime of the detector. This requires that wire tensions be monitored continuously in situ with 1% accuracy. The ten meter drift tube length makes it imperative to maintain a large tension of one kilogram on the anode wire to minimize the magnitude of the sagitta, induced by gravity. Thus the development of a tension monitoring device integrated into the high voltage system, which can be operated concurrently with data taking, is an essential design feature of the system.

Tension measurement requires that the tensioned wire be forced into mechanical oscillation whose frequency and amplitude we can vary to find its

resonance frequency. Since the drift tubes are rigidly attached to each other and thus they cannot move, and the assembled tubes cannot be immersed in magnetic field, the only method left to make the anode wire to oscillate is the electrostatic force. In an operating SDC drift tube the shaping electrodes are held at a common potential by a hardwired endcap on the H.V. end of the tube. The fact that the anode wire is strung along the symmetry axis of the tube-shaping electrode system makes it difficult to create a net electrostatic force on it and thus to induce mechanical oscillation. Due to the cylindrical and mirror symmetry of the configuration, in the absence of anode wire sagging due to gravity, no net force acts on the anode wire even in the presence of a potential difference between the shaping electrodes and the anode wire. The sagging of the anode wire breaks one or both of the symmetries and a small force between the anode wire and shaping electrodes develops. The small oscillation amplitude of the wire even at resonance makes it difficult to determine the resonance frequency. These problems demand a highly sensitive instrument which can induce wire oscillation and detect a very small capacitance change of the anode wire ($\delta C_w \approx 1-10\text{fF}$) due to oscillation in the presence of large tube-electrode-wire capacitance ($C_w \approx 100\text{ pF}$). We have developed a method and constructed such a measuring device. To measure the minute variation of capacitance of the anode wire tube-shaping electrode system (C_w) as a function of the anode wire driving frequency, thus the oscillation amplitude, we used the properties of parallel LC resonance circuits. As is well known² and shown in Fig. IV-1, the impedance of this circuit has a maximum at $\omega_0^2 = 1/LC$ as a function of angular frequency ω . If the frequency of the HF generator is set at ω_1 which is slightly different from ω_0 , then the operating point is set on one of the sharply varying wings of the resonance curve. Using this operating point, small changes in frequency cause large changes in impedance. The same is

true if at the fixed ω_1 operating point the capacitance of the resonance circuit (via the C_w of the oscillating anode wire) is varied. This attribute of the slightly off tuned resonance circuit is well suited to detect the changes in C_w of the oscillating anode wire. In our case, $C_w \approx 100$ pF while the variation of C_w (δC_w) due to the variation is $\delta C_w \approx 10$ fF. Thus the variation in capacitance is from four to five order of magnitude smaller than magnitude of the varying capacitance. With the help of an off tuned resonance circuit this sensitivity can be reached without any difficulty. The resonance circuit in Fig. IV-2 consists of the inductance L , capacitances C_2 and C_w . Capacitance C_1 and the impedance of the resonance circuit form a voltage divider chain, fed by the HF oscillator which provides a few MHz of 1V peak to peak signal. Due to the oscillating anode wire, the capacitance C_w and thus the impedance of the resonance circuit varies periodically. As a result a modulated HF signal appears on the voltage divider. The amplitude modulation of this signal is proportional to the amplitude of the mechanical oscillation of the wire. The demodulation is done by a diode and the $R_2 C_3$ filter. The emerging signal is amplified. The amplifier output can be viewed on a scope or measured by an ADC system.

Fig IV - L

Fig IV - 2

V. Publications since 1985

1. Report of Working Group on Detectors for Hadrons and Event Parameters at Colliders, W. Carithers, D.D. Greiner, W. Meyers, L.S. Schroeder, H. Steiner, T.J.M. Symons, G. Young, D. Beavis, G. van Dalen, D.D. Carmony, A.S. Goldhaber, G.D. Westfall, and H. Wieman, Proceedings of the Work-shop on Detectors for Relativistic Nuclear Collisions, pp 75-83, Lawrence Berkeley Lab LBL-18225 (1984).
2. The Description of Fragmentation as a Critical Phenomenon in High Energy P-Nucleus Collisions, A.S. Hirsch, A. Bujak, J.E. Finn, L.J. Gutay, R.W. Minich, N.T. Porile, R.P. Scharenberg, B.C. Stringfellow, and F. Turkot, Nucl. Phys. A418, 267c (1984).
3. Nuclear Charge Dispersion in the Fragmentation Mass Region and the Thermal Liquid Drop Model, N.T. Porile, A. Bujak, J.E. Finn, L.J. Gutay, A.S. Hirsch, R.W. Minich, G. Paderewski, R.P. Scharenberg, B.C. Stringfellow, and F. Turkot, Phys. Lett. 156B 177 (1985).
4. Mass Yield Distribution for the Interaction of Silver with 300 GeV Protons, A. Bujak, J.E. Finn, L.J. Gutay, A.S. Hirsch, R.W. Minich, G. Paderewski, N.T. Porile, R.P. Scharenberg, B.C. Stringfellow, and F. Turkot, Phys. Rev. C 32 620 (1985).
5. Accelerator Internal Target Experiments Using a Supersonic Gas Jet, B.C. Stringfellow, A.T. Bujak, Y.H. Chung, J.E. Finn, L.J. Gutay, A.S. Hirsch, M. Mahi, G.L. Paderewski, N.T. Porile, T.C. Sangster, R.P. Scharenberg, and F. Turkot, Nuclear Instruments and Methods in Phys. Rev. A251, 242-253, (1986).
6. Fragment Emission in Proton Xenon Interactions at Near Threshold Energies. N.T. Porile, A. Bujak, D.D. Carmony, Y.H. Chung, L.J. Gutay, A.S. Hirsch, M. Mahi, G.L. Paderewski, T.C. Sangster, R.P. Scharenberg, B.C. Stringfellow Proceedings, Nuclear Dynamics 113-115. (Copper Mountain, 1986).
7. FNAL E-735, A Search for A Deconfined Quark Gluon Phase of Strongly Interacting Matter in $\bar{p}$ -p Interactions at $\sqrt{s} = 2$ TeV, L.R. Fortney, A.T. Goshaw, S. Oh, W.J. Robertson, W.D. Walker, V.H. Areti, C. Hojvat, D. Reeves, F. Turkot, E.W. Anderson, C. Lindsey, M. Duong-Van, J.M. Bishop, N.N. Biswas, J. LoSecco, P. Kenney, J. Piekarz, A. Bujak, D.D. Carmony, R.J. De Bonte, L.J. Gutay, A.S. Hirsch, D.S. Koltick, N. Morgan, N.T. Porile, R.P. Scharenberg, B.C. Stringfellow, A. Erwin, C. Findeisen, K. Nelson, and M. Thompson, Proceedings of XVII International Symposium on Multiparticle dynamics (Seewinkel, Austria, June 1986), World Scientific Press pp 839-842.
8. FNAL Experiment E-735 at the C0 Collider Intersection, L.R. Fortney, A.T. Goshaw, S. Oh, W.J. Robertson, W.D. Walker, V.H. Areti, C. Hojvat, D. Reeves, F. Turkot, E.W. Anderson, C. Lindsey, M. Duong-Van, J.M. Bishop, N.N. Biswas, J. LoSecco, V.P. Kenney, J. Piekarz, A. Bujak, D.D. Carmony, R.J. De Bonte, L.J. Gutay, A.S. Hirsch, D.S. Koltick, N. Morgan, N.T. Porile, R.P. Scharenberg, B.C. Stringfellow, A. Erwin, C. Findeisen, K. Nelson, and M. Thompson, Proceedings of the International conference on antiproton-proton Interactions, Aachen (1986).

9. Light-Fragment Production in Proton-Xenon Interactions between 1 and 19 GeV, T.C. Sangster, A.T. Bujak, D.D. Carmony, Y.H. Chung, L.J. Gutay, A.S. Hirsch, M. Mahi, G.L. Paderewski, N.T. Porile, R.P. Scharenberg, B.C. Stringfellow, Phys. Lett. B 188 29-32 (1987).
10. Fragment Production from 1-GeV to 20-GeV P + Xe Reactions. T.C. Sangster, A. Bujak, D.D. Carmony, Y.H. Chung, L.J. Gutay, A.S. Hirsch, M. Mahi, G.L. Paderewski, N.T. Porile, R.P. Scharenberg, B.C. Stringfellow. Proceedings, High Energy Heavy Ion Study 363-371. (Berkeley, 1987).
11. Energy dependence of Proton Induced Xenon Fragmentation and the Approach to Liquid-gas Criticality in Nuclear Matter, A.S. Hirsch, A.T. Bujak, D.D. Carmony, Y.H. Chung, L.J. Gutay, M. Mahi, G.L. Paderewski, N.T. Porile, T.C. Sangster, R.P. Scharenberg, and B.C. Stringfellow, Phys. Rev. Letters. 60 1936-1939 (1988).
12. Fragment Emission in Proton-Xenon Interactions in the Near-Threshold Regime, N.T. Porile, A.T. Bujak, D.D. Carmony, Y.H. Chung, L.J. Gutay, A.S. Hirsch, M. Mahi, G.L. Paderewski, T.C. Sangster, R.P. Scharenberg, and B.C. Stringfellow, Nuclear Physics A 471 149c-169c (1987).
13. Degradation of Plastic Scintillator Exposed to Tevatron Tunnel Background Radiation, C.S. Lindsey, E.W. Anderson, C. Hojvat, F. Turkot, A. Bujak, Nuclear Instruments and Methods in Physics A254 212-214 (1987).
14. Multiplicity dependence of the transverse-momentum spectrum for centrally produced hadrons in antiproton-proton collisions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P.D. Berry, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, C.S. Lindsey, J.M. Losecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, R.P. Scharenberg, B.C. Stringfellow, S.R. Stampke, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, and D.K. Wesson. Phys. Rev Lett. 60 1622-1625 (1988).
15. Approach to Criticality in the Fragmentation of Xenon, N.T. Porile, A.J. Bujak, D.D. Carmony, Y.H. Chung, L.J. Gutay, A.S. Hirsch, M. Mahi, G.L. Paderewski, T.C. Sangster, R.P. Scharenberg, and B.C. Stringfellow, Phys. Rev. C39, 1914 - 1928 (1989).
16. The Transverse Momentum Spectrum of Charged Particles Produced in $\bar{p}p$ Collisions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, E. W. Anderson, H. Areti, S. Banerjee, P. D. Berry, N. N. Biswas, A. Bujak, D. D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A. T. Goshaw, L. J. Gutay, A. S. Hirsch, C. Hojvat, V. P. Kenney, D. Koltick, C. S. Lindsey, J. M. LoSecco, T. McMahon, A. P. McManus, N. Morgan, K. Nelson, S. H. Oh, J. Piekarz, N. T. Porile, D. Reeves, R. P. Scharenberg, B. C. Stringfellow, S. R. Stampke, M. Thompson, F. Turkot, W. D. Walker, C. H. Wang, and D. K. Wesson, Proceedings of the Rencontres de Moriond, Les Arcs, France (1988).

17. A Study of 1.8 TeV Proton-Antiproton Collisions at the C0 region of the Fermilab Tevatron Collider, T. Alexopoulos, C. Allen, E. W. Anderson, H. Areti, S. Banerjee, P. D. Berry, N. N. Biswas, A. Bujak, D. D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A. T. Goshaw, L. J. Gutay, A. S. Hirsch, C. Hojvat, V. P. Kenney, D. Koltick, C. S. Lindsey, J. M. LoSecco, T. McMahon, A. P. McManus, N. Morgan, K. Nelson, S. H. Oh, J. Piekartz, N. T. Porile, D. Reeves, R. P. Scharenberg, B. C. Stringfellow, S. R. Stampke, M. Thompson, F. Turkot, W. D. Walker, C. H. Wang, and D. K. Wesson, Proceedings of the Division of Particles and Fields Meeting, Storrs, CT (1988).
18. Preliminary Results on $p(t)$ Distributions for TOF Identified Hadrons from Anti-P P Collisions at $\sqrt{s} = 1.8$ -TeV. E-735 Collaboration. FNAL-E735 experiment. V.P. Kenney, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P.D. Beery, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, D. Koltick, C.S. Lindsey, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, K. Piekartz, N.T. Porile, D. Reeves, R.P. Scharenberg, B.C. Stringfellow, S.R. Stampke, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson, 1988. Proceedings, Current Issues in Hadron Physics 193-196. (Les Arcs, 1988).
19. Lambda and Anti-lambda Production from Anti-P P Collisions at 1.8-TeV Center of Mass Energy. E735 collaboration. Fnal-735 experiment. S.H. Oh, S. Banerjee, P.D. Beery, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, Y. Choi, C. Findeisen, A.T. Goshaw, C. Hojvat, C.S. Lindsey, J.M. LoSecco, A.P. McManus, N. Morgan, K. Nelson, J. Piekartz, D. Reeves, S.R. Stampke, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson Proceedings, Hadronic matter in collision 594-606. (Tucson, 1988).
20. Measurement of p_t as a function of N_C at the FNAL $p\bar{p}$ Collider. S.R. Stampke, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P.D. Beery, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, P. Cole, Y. Choi, R. DeBonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, C.S. Lindsey, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekartz, N.T. Porile, D. Reeves, R.P. Scharenberg, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson (Duke U. & Fermilab & Iowa State U. & Notre Dame U. & Purdue U. & Wisconsin U., Madison), 1988. Proceedings, Hadronic matter in collision 1988 67-77. (Tucson, 1988).
21. Recent results from E735 at the Fermilab Tevatron Proton Anti-proton Collider with $\sqrt{s} = 1.8$ -TeV. C.S. Lindsey, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P.D. Beery, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, P. Cole, Y. Choi, R. DeBonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekartz, N.T. Porile, D. Reeves, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson. Nucl. Phys. A498 181C-192C (1989).

22. Λ^0 and $\bar{\Lambda}^0$ Production from Proton-Antiproton Collisions at $\sqrt{s} = 1.8$ TeV, S. Banerjee, P. D. Berry, N. N. Biswas, A. Bujak, D. D. Carmony, T. Carter, Y. Choi, C. Findeisen, A. T. Goshaw, C. Hojvat, C. S. Lindsey, J. M. LoSecco, A. P. McManus, N. Morgan, K. Nelson, S. H. Oh, J. Piekarz, D. Reeves, S. R. Stampke, M. Thompson, F. Turkot, W. D. Walker, C. H. Wang, and D. K. Wesson, Phys. Rev. Lett. 62, 12-15 (1989).
23. Mass Identified Particle Yields in Antiproton-Proton Collisions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, E. W. Anderson, H. Areti, S. Banerjee, P. D. Berry, N. N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A. T. Goshaw, L. J. Gutay, A. S. Hirsch, C. Hojvat, V. P. Kenney, C. S. Lindsey, J. M. LoSecco, T. McMahon, A. P. McManus, N. Morgan, K. Nelson, S. H. Oh, J. Piekarz, N. T. Porile, D. Reeves, R. P. Scharenberg, B. C. Stringfellow, S. R. Stampke, M. Thompson, F. Turkot, W. D. Walker, C. H. Wang, and D. K. Wesson, Phys. Rev. Lett. 64, 991-994 (1990).
24. Recent Results from E-735 at the Fermilab Tevatron Proton-Antiproton Collider at $\sqrt{s} = 1.8$ TeV, C. Allen, A. Bujak, D. D. Carmony, P. Cole, Y. Choi, R. De Bonte, L. J. Gutay, A. S. Hirsch, D. Koltick, T. McMahon, N. K. Morgan, N. T. Porile, R. P. Scharenberg, B. C. Stringfellow et al., Nuclear Physics B12, 267-276 (1990).
25. A Cylindrical Drift Chamber for the Measurement of High Charged Particle Multiplicities in Hadronic Events, (C. Allen, A. Bujak, D. Carmony, P. Cole, Y. Choi, R. De Bonte, J.E. Finn, L.J. Gutay, A.S. Hirsch, D. Koltick, T. McMahon, N.K. Morgan, N.T. Porile, R.P. Scharenberg, and B.C. Stringfellow). Nucl. Instr. and Methods A294 108-116(1990).
26. Search for Quark-Gluon Plasma in $\bar{p}p$ Collisions at $\sqrt{s} = 1.8$ TeV, (with T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, J. Bishop, N.N. Biswas, A. Bujak, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, C.S. Lindsey, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, and D.K. Wesson). Proceedings of the Rio de Janeiro International Workshop on Relativistic aspects of nuclear physics, pp. 3-16 (World Scientific Publishing Co., 1990).
27. Deconfinement Signature, Mass Dependence of the Transverse Flow and Time Evolution in Antiproton-Proton Collisions at $\sqrt{s} = 1.8$ TeV, (with T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. K. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, W.R. Roberston, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson, and H. Zuong). Physica Scripta T32 122-125 (1990).
28. Mass Limits for Excited Electrons and Muons from Z^0 Decay, L3 Collaboration/A. Bujak, L. Gutay, T. McMahon, B. Stringfellow, Phys. Lett. B247, 117-184, September 6, 1990.

29. A Determination of Electroweak Parameters from $Z^0 \rightarrow \mu^+ \mu^- (\gamma)$, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, B. Stringfellow, Phys. Lett. B247, 473-480, September 13, 1990.
30. A Test of QCD Based on 4-Jet Events from Z^0 Decays, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, B. Stringfellow, Phys. Lett. B248, 227-234, September 27, 1990.
31. Determination of α_s from Jet Multiplicities Measured on the Z^0 Resonance, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, B. Stringfellow, Phys. Lett. B248, 464-472, October 4, 1990.
32. A Precision Measurement of the Number of Neutrino Species, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, B. Stringfellow, Phys. Lett. B249, 341-352, October 18, 1990.
33. A Determination of Electroweak Parameters from Z^0 Decays into Charged Leptons, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B250, 183-192, November 1, 1990.
34. Test of QCD in $e^+e^- \rightarrow \gamma\gamma$ at LEP, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow. Phys. Lett. B250, 199-204, November 1, 1990.
35. Search for Excited Taus, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B250, 205-211, November 1, 1990.
36. Search for the Neutral Higgs Bosons of the Minimal Supersymmetric Standard Model from Z^0 Decays, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B251, 311-320, November 15, 1990.
37. A Search for Heavy Charged and Neutral Leptons from Z^0 Decays, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B251, 321-330, November 15, 1990.
38. Search for the Charged Higgs Boson in Z^0 Decay, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B252, 511-517, December 20, 1990.
39. Search for a Low Mass Neutral Higgs Boson in Z^0 Decay, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B252, 518-524, December 20, 1990.
40. Search for Excited Neutrinos from Z^0 Decays, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B252, 525-532, December 20, 1990.
41. A Measurement of $B^0-\bar{B}^0$ Mixing in Z^0 Decays, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B252, 703-712, December 20, 1990.
42. A Measurement of the $Z^0 \rightarrow b\bar{b}$ Forward-Backward Asymmetry, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B252, 713-721, December 27, 1990.

43. A Quark - Gluon Plasma Search in Anti-P P at $\sqrt{s} = 1.8$ TeV. By E735 Collaboration. T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P.D. Beery, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, C.S. Lindsey, L.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson. Nucl.Phys. A525:165-170 (1991).
44. Hyperon Production from Proton-Antiproton Collisions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, H. Areti, S. Banerjee, Bhat, J. Bishop, N.N. Biswas, A. Bujak, D.D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, C.S. Lindsey, C. Loomis, J. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, K. Piekarz, N.T. Porile, D. Reeves, R.P. Scharenberg, R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson, and H. Zuong, Physical Review D46 2773-2786 (1991).
45. Search for the Neutral Higgs Boson, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B257, 450, March 28, 1991.
46. Determination of α_s from Energy-Energy Correlations Measured on the Z^0 Resonance, L3 collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, Phys. Lett. B257, 469, March 28, 1991.
47. Recent Results From E735 — Search For QGP in $\sqrt{s} = 1.8$ TeV pbar-p Collisions, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. K. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, W.R. Roberston, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson, and H. Zuong). Proceedings of the 7th Winter Workshop on Nuclear Dynamics, Key West, January 27 - February 1 1991. (to be published).
48. Measurement of the Inclusive Production of Neutral Pions and Charged Particles on the Z^0 Resonance, L3 Collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, To be published in Physics Letters B.
49. Search for Leptoquarks in Z^0 Decays, L3 Collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, To be published in Physics Letters B.
50. Measurement of $Z^0 \rightarrow b\bar{b}$ Decays and the Semileptonic Branching Ratio $Br(b \rightarrow L + X)$, L3 Collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, To be published in Physics Letters B.
51. Measurement of Electroweak Parameters from Hadronic and Leptonic Decays of the Z^0 , L3 Collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, To be published in Zeitschrift fur Physik G.
52. Search for Narrow High-Mass Resonances in Radiative Decays of the Z^0 , L3 Collaboration/A. Bujak, L. Gutay, T. McMahon, and B. Stringfellow, To be published in Physics Letters B.

53. Correlations in Hadron Production in 1.8 TeV Anti-p p collisions, V.P. Kenney, N.N. Biswas, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P.D. Berry, J. Bishop, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, C.S. Lindsey, J.M. Losecco, T. McMahon, A.P. McManus, N. K. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, W.P. Schareneberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson, and H. Zuong, Proceedings of the Singapore Conference, Vol. 1 459-460 (1990).
54. A Search for the Signature of a Deconfined Quark Gluon Phase of Strongly Interacting Matter in $\bar{p}$ -p Interactions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, E.W. Anderson, V. Balamurali, S. Banerjee, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, N.T. Porile, D. Reeves, W.J. Robertson, R.P. Scharenberg, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and Y. Zhuan, Proceedings of the XXVI International Conference on High Energy Physics, Dallas (1992).
55. Particle Production, Hadron Deconfinement and Thermodynamics in $\bar{p}$ -p Interactions, T. Alexopoulos, C. Allen, E.W. Anderson, V. Balamurali, S. Banerjee, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, N.T. Porile, D. Reeves, W.J. Robertson, R.P. Scharenberg, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and Y. Zhuan, Proceedings of the NATO Advanced Study Institute on Particle Production in Highly Excited Matter. Il Ciocco, Italy (1992).
56. Results from E735 at the Tevatron Proton - Anti-proton Collider with $\sqrt{s} = 1.8$ TeV, C.S. Lindsey, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, S. Banerjee, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, A. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, V.P. Kenney, D. Koltick, C. Loomis, J.M. Losecco, T. McMahon, A.P. McManus, N. K. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, W.R. Roberston, R.P. Schareneberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, D.K. Wesson, and H. Zuong, Quark Matter, Gatlinberg TN, Nucl. Phys. A544: 343-356 (1992).
57. Mass Identified Particle Production in Proton-Antiproton Collisions at $\sqrt{s} = 300, 540, 1000, \text{ and } 1800$ GeV, T. Alexopoulos, C. Allen, E.W. Anderson, V. Balamurali, S. Banerjee, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, N.T. Porile, D. Reeves, W.J. Robertson, R.P. Scharenberg, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and Y. Zhuan, Submitted to Physical Review D .

58. Inclusive Photon Production from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, E.W. Anderson, V. Balamurali, S. Banerjee, P.D. Berry, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, Y. Choi, P. Cole, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, N.T. Porile, D. Reeves, A. Rimai, W.J. Robertson, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and Y. Zhuang, Submitted to Physical Review Letters.
59. Charged Particle Multiplicity Distributions in Proton-Antiproton Collisions at $\sqrt{s} = 0.546$ TeV, 1.0 TeV, and 1.8 TeV, T. Alexopoulos, C. Allen, E.W. Anderson, V. Balamurali, S. Banerjee, P.D. Berry, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarczyk, N.T. Porile, D. Reeves, A. Rimai, W.J. Robertson, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and H. Zuong, to be submitted to Physical Review.
60. ϕ^0 Meson Production from $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, V. Balamurali, S. Banerjee, P.D. Berry, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarczyk, N.T. Porile, D. Reeves, A. Rimai, W.J. Robertson, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and H. Zuong, to be submitted to Physical Review Letters.
61. A Study of Source Size in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV Using Pion Interferometry, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, V. Balamurali, S. Banerjee, P.D. Berry, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarczyk, N.T. Porile, D. Reeves, A. Rimai, W.J. Robertson, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and H. Zuong, to be submitted to Physical Review D.

62. Multiplicity Dependence of Transverse Momentum Spectra of Centrally Produced Hadron in pp Collisions at 0.3, 0.54, 1.0, and 1.8 TeV Center of Mass Energy, T. Alexopoulos, C. Allen, E.W. Anderson, H. Areti, V. Balamurali, S. Banerjee, P.D. Berry, P. Bhat, J. Bishop, N.N. Biswas, A. Bujak, D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. deCarlo, A.R. Erwin, C. Findeisen, A.T. Goshaw, L.J. Gutay, A.S. Hirsch, C. Hojvat, J.R. Jennings, V.P. Kenney, C.S. Lindsey, C. Loomis, J.M. LoSecco, T. McMahon, A.P. McManus, N. Morgan, K. Nelson, S.H. Oh, J. Piekarz, N.T. Porile, D. Reeves, A. Rimai, W.J. Robertson, R.P. Scharenberg, S.R. Stampke, B.C. Stringfellow, M. Thompson, F. Turkot, W.D. Walker, C.H. Wang, J. Warchol, D.K. Wesson, and H. Zuong, to be submitted to Physical Review D.

Purdue University
Grant DE-AC02-76ER01428 Task E

AN EXPERIMENTAL HIGH ENERGY PHYSICS PROGRAM

within the

DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY
West Lafayette, Indiana 47907

Supported by

U.S. DEPARTMENT OF ENERGY

Through

Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana

Current Grant Number: DE-AC02-76ER01428

Task E

PROGRESS REPORT

Principal Investigators: D. S. Koltick, R. L. McIlwain, E. I. Shibata

April 1993

Table of Contents

I.	Personnel	2
II.	Summary	3
III.	TOPAZ.....	3
	A. Performance of the Vertex Chamber	
	B. Study of Electroweak Interactions	
IV.	SSC.....	8
	A. Brookhaven Beam Test	
	B. Supervisor of TNRLC and MOUs for the Fiber Tracking Group	
	C. Construction of Fiber Ribbons	
	D. Test of Fiber Ribbon Mounting Techniques	
V.	D-Zero.....	16
	A. Status of tau physics	
	B. Multi-Jet physics	
VI.	Publications and Presentations.....	20

I. PERSONNEL

The personnel involved in Task E are:

TOPAZ at KEK

D. S. Koltick	Professor
B. L. Howell	Research Associate
I. Y. Levine	Graduate Student

Solenoid Detector Collaboration (SDC) at SSC and D0-Main Injector Upgrade

D. S. Koltick	Professor
R. L. McIlwain	Professor
E. I. Shibata	Professor (SSC only)
C. L. Talmadge	Research Assistant Professor
B. L. Howell	Research Associate
J. C. Cooper	Research Engineer
C. J. Schmitz	Technical Assistant
I. Ricky	Technical Assistant
T. Crone	Laboratory Assistant
B. K. Abbott	Graduate student
S. Keller	Undergraduate student
D. Schleef	Undergraduate student

II. SUMMARY

This past year has seen many accomplishments for this task. The activities of Task E can be divided into three areas: Topaz experiment at TRISTAN in Japan, the Solenoidal Detector Collaboration at the Superconducting Super Collider Laboratory, and the D-Zero experiment at Fermilab. Important progress has been made in each activity.

A. TRISTAN

The efforts at Topaz are now bearing fruit. We had an invited talk in the electroweak session of this year's International Conference on High Energy Physics at Dallas, Texas. At ICHEP we presented the world's most significant measurements of the electroweak asymmetry for the production of muon and tau lepton pairs. At this year's American Physical Society Division of Particles and Fields meeting at Fermilab, we presented a paper using these measurements to place strong mass limits on the existence of Z' bosons. The limits obtained at Topaz are already among the world's best for some extensions of the standard model. Once this year's data that is now being collected at TRISTAN is included into the analysis, very significant information on the possible existence of Z' bosons will be in hand.

We are also building a significant data base with the vertex chamber at TOPAZ that we built. The impact resolution is under study for hadronic events. Relatively clean samples of B-tagged events is the expected outcome of this effort.

B. SSC

The efforts of the group on the SDC experiment have been significant. These include both experimental and administrative type activities. David Koltick is secretary of the Institutional Board and has been responsible for producing minutes for world-wide distribution. But more importantly, Purdue University has been the Center for Scintillating Fiber Tracking for the SDC. This has included organizing the Memorandum of Understanding between the 10 institutions of the fiber tracking group and SSCL for the conduct of research into scintillating fiber tracking at the SSC.

On the experimental side, we have been a major participant in two beam tests. One at Fermilab in the new muon lab and the other at Brookhaven. The Fermilab experiment has been published. The Brookhaven work will be published shortly.

We have also learned to produce very accurate scintillating fiber ribbons up to 4.5 meters in length. In addition, we have worked with Hercules Corporation to produce an extremely accurate composite cylinder to serve as the base for the scintillating fiber coordinate system. Work is just now starting on the characterization of the lots of HISTE IV, VLPCs that have been produced at Rockwell International. This includes about 100,000 VLPC pixels.

C. D-Zero

Work on D-Zero is focused into two areas. Analysis of the first physics data now being taken at Fermilab and work on the upgraded D-Zero charged particle tracking system, specifically work on the scintillating fiber outer tracker. In the data analysis, we have continued our long tradition of work on the tau lepton. We have already isolated a sample of $Z \rightarrow \tau \tau^*$ decay. Studies on QCD multi-jets are also underway.

III. TOPAZ

This past year the TOPAZ experiment has been smoothly taking data. We have accumulated an integrated luminosity of 67.3 pb^{-1} without any essential problems with the detector.

The vertex chamber, newly installed in the last year, has been operated steadily, our group has obtained impact parameter resolutions of $75 \text{ }\mu\text{m}$ for Bhabha events.

Figure III.1 shows the history of the integrated luminosity per day accumulated by the TOPAZ detector from the beginning of the experiment until now. The luminosity has remarkably improved by the commission of the new QC1 and skew magnet systems a half year after the installation of the superconducting Q-magnet (QCS). Even though we received less luminosity than the other groups due to machine asymmetries, we recorded luminosities of $890 \text{ nb}^{-1}/\text{day}$ this year and have collected total integrated luminosity for the experiment of 159.9 pb^{-1} .

Fig. III.1 Integrated luminosity per day from the beginning of the experiment(1987) till now.

We have actively pursued, the analysis of lepton pair production, which has resulted in a published paper, a contribution paper to the 26th international Conference on High Energy Physics at Dallas and to the DPF meeting at Fermilab, and a doctoral thesis this year. In addition to these activities, an energy scan between 58 and 60 GeV was performed last autumn to examine the possible new resonance suggested by the LEP experiments. The total cross sections for $\gamma\gamma$, multi-hadron, e^+e^- and $\mu^+\mu^-$ productions were measured at 8 energy points. The results of the scan are in good agreement with the Standard Model predictions.

A. Performance of the Vertex Chamber

Using an accumulated data set of about 33 pb^{-1} , intensive efforts have been made to calibrate the VTX in order to achieve the best spatial resolution possible. The average local spatial resolution has been improved to approximately $35 \mu\text{m}$ over the maximum drift length of 3 cm. The average global spatial resolution (track residuals) is limited to $40 \mu\text{m}$. The average impact parameter resolution, estimated using miss distance distributions for Bhabha data, has been reduced to $75 \mu\text{m}$ on average. This should be compared with the average B lifetime of approximately $700 \mu\text{m}$ in the lab frame. The miss distance distribution for the sector pair with the best performance is shown in Figure III.2.

The beam position, profile and stability were also studied using Bhabha events. For one data-taking period, the center of the beam interaction point was found to be $x = +340 \pm 290 \mu\text{m}$ and $y = -758 \pm 106 \mu\text{m}$ with respect to the center of the VTX. This indicates a remarkably small spread in both x and y for such a long data taking period. Preliminary results of a study of the stability of the beam are shown in Figure III.3. After a brief interruption in data taking, the center of the beam is seen to have shifted with respect to the VTX origin by approximately $100 \mu\text{m}$.

With the strong constrain on the interaction point resolution given above, and by combined track reconstruction with the outer chambers, we expect to distinguish secondary vertices for short decay particles with high precision. Further work to improve the efficiency and precision of linking of tracks in the VTX to those in the TCH and the TPC is in progress.

Fig. III.2 Miss distance between collinear two tracks of Bhabha events for the study of impact parameter resolution.

Fig. III.3 Beam position and beam profile measured by the VTX.

B. Study of Electroweak Interactions

We have completed an analysis of $e^+e^- \rightarrow \mu^+\mu^-$ and $e^+e^- \rightarrow \tau^+\tau^-$ reactions at TRISTAN based on a data sample of 75 pb^{-1} of integrated luminosity, collected by the TOPAZ detector prior to 1992. This analysis has culminated in the world's most significant measurement of the forward-backward asymmetry for lepton pairs. At TRISTAN energies, the electroweak asymmetry is approaching its maximum value. Figure III.4 shows our result. The importance of the asymmetry measurement is that it provides the most significant verification that electromagnetic and weak forces interfere with one another as predicted by the standard model. TRISTAN is an ideal place to observe photon- Z^0 interference because the e^+e^- interactions occur at energies below the Z^0 mass, where both electromagnetic and weak forces contribute almost equally to the production of fermion pairs.

During the past year, measurement of the lepton pair asymmetry has led to a Ph.D. thesis, a publication in Physics Letters B, and invited talks at the International Conference on High Energy Physics in Dallas, Texas, the November 1992 APS Meeting of the Division of Particles and Fields, in Chicago, Illinois, and the December 1992 TRISTAN Physics Workshop held at KEK National Laboratory, Japan. In these talks, we reported the constraints placed on new physics possibilities outside the standard model by our measurements. For example, we presented improved limits on the compositeness scale for mu and tau leptons, and their coupling to massive extra Z' bosons. These limits are shown in Tables I and II.

Table I: Compositeness scale parameter ($\Lambda^\pm$) limits derived from TOPAZ $e^+e^- \rightarrow \mu^+\mu^-$ and $e^+e^- \rightarrow \tau^+\tau^-$ data, for the following couplings to a contact interaction describing the compositeness: RR (right-right); LL (left-left); VV (vector-vector); AA (axial vector-axial vector).

$\Lambda^+ / \Lambda^- (\text{TeV})$	$e^+e^- \rightarrow \mu^+\mu^-$	$e^+e^- \rightarrow \tau^+\tau^-$	$e^+e^- \rightarrow l^+l^-$
Limit (95% C.L.)			
RR Coupling	>2.1/3.1	>1.8/2.7	>2.2/4.1
LL	>2.1/3.0	>1.7/2.6	>2.2/3.9
VV	>11.2/3.4	>3.5/4.3	>7.1/4.2
AA	>3.1/7.0	>2.7/3.9	>3.2/8.7

Table II: Mass Limits for various Z' bosons determined from an analysis of TRISTAN e^+e^- data in comparison to $p\bar{p}$ data.

$M_{Z'} (\text{GeV})$	Z_1	Z_ψ	Z_x	Z_η	Z_ν
Limit (95% C.L.)					
TOPAZ (this analysis)	>290	>146	>134	>100	>164
TRISTAN (this analysis)	>430	>166	>245	>145	>196
CDF ($p\bar{p}$ data)	>412	>320	>340	>340	---

Figure III.4

The forward-backward asymmetry for lepton pairs (either μ or τ) as a function of energy.

IV. SSC

A. Brookhaven Beam Test

As shown in Figure IV.1 a system of 96 scintillating fibers was tested in the A2 beam line at the Brookhaven National Laboratory Alternating Gradient Synchrotron in June and July 1992 to evaluate the system performance of a scintillating fiber charged particle tracking system. The fibers were arranged in three superlayers, and each of the three superlayers was made of four layers of scintillating fibers at Purdue by gluing double layered ribbons (doublets), also made at Purdue, to each side of a 1.6 cm thick board. Scintillating fibers of 4.3 m length and 830 μm diameter were optically coupled to clear fiber waveguides of the same diameter and 6 m length and read out with HISTE-III (High-rate Scintillating Fiber Tracking Experiment, third generation) visible light photon counters (VLPCs) made by the Rockwell International Science Center. The VLPCs were kept at their operating temperature ($\sim 7^\circ\text{K}$ in three small cryostats).

A total of 96 channels were instrumented for the test. Due to microbonding problems on some of the VLPC channels, 17 of the 96 channels were not active in the test; the HISTE-IV VLPCs have different microbonding locations and there are no microbonding problems in this latest generation. Analog data were amplified in an amplifier-shaper-discriminator (ASD) module to generate analog output as well as binary hit output for channels over threshold. The binary hit data were then provided to a portotype trigger card, which used the parth of hits through the fibers to identify tracks. The analog data were digitized using fastbus analog-to-digital converters (ADCs) and were read into a 486DX microcomputer. Figure IV.2 shows some typical events which demonstrate that the digital triggering works.

To find the spatial resolution of a doublet layer of scintillating fibers, a straight line was drawn through a pair of hit fibers in the first and and sixth ribbon doublets, and the hit fibers within $1/2$ fiber diameter of this straight line were determined. In Figure IV.3 the spatial distribution of hits relative to the straight line described above are shown for various doublet layers of scintillating fibers. These histograms show narrow peaks, which are characteristic of the digital tracking nature of a scintillating fiber or silicon strip tracking system. The position assigned to a single hit in a doublet is the center position of the hit fiber, and for double hits in a doublet (a hit in one layer plus an adjacent hit in the second layer of the doublet) the position is halfway between the centers of the two hit fibers. The root mean square deviations are 207, 178, 200, and 208 μm for doublet layers 2, 3, 4, and 5 respectively and contain uncertainties from the determination of the track coordinates from the first and sixth doublets as well as that from the doublet layer itself. For the outer doublet layers 2 and 5 the doublet resolution is simply the root mean square deviation divided by $\sqrt{2}$, or 146 μm , assuming all doublets to have the same resolution. For the inner doublet layers 3 and 4 the doublet resolution are the root mean square deviations divided by $\sqrt{3/2}$, which give 145 μm for doublet layer 3 and 163 μm for doublet layer 4. Thus, the average spatial resolution is about 140 μm , which is within 20% of the expectation of $\frac{830}{2\sqrt{12}} \mu\text{m} = 120 \mu\text{m}$ for a scintillating fiber doublet. This measured resolution includes cases in which there is only one hit in a fiber doublet and the assigned position for the hit is the center of the hit fiber. Of course, the resolution is better in cases in which two fibers in a doublet have hits, as will be the case for the SDC system with its higher photostatistics.

Schematic diagram of the three superlayer scintillating fiber tracking system in a 5 GeV/c π^- test beam at BNL.

Figure IV.1

Event Number - 2519

Event Number - 17995

ADC

Trigger

Both

Typical event displays showing hits according to the ADC system, the digital trigger system, or both systems. These events have been identified as straight beam tracks by the prototype digital trigger.

Figure IV.2

Spatial distribution of hits in the four inner doublet layers. The root mean square deviations are 207, 178, 200, and 208 μm for doublet layers 2, 3, 4, and 5 respectively. These deviations give fiber doublet resolutions of 146, 145, 163, and 146 μm for these doublet layers.

Figure IV.3

A Monte Carlo simulation of the fiber tracker was developed by Purdue in order to understand the average number of photoelectrons produced by a minimum ionizing particle. The fiber diameter, the cladding thickness and the spacing between fibers were simulated in the Monte Carlo. Errors in placing the fibers on the boards were determined by finding tracks in the data. In the tracking algorithms, the placement of the fibers on the board was allowed to vary. The distribution between the tracks and the hit fibers was plotted as a function of offsets. The offsets that produced a distribution centered at zero were then used in the Monte Carlo to simulate the actual test geometry. Hits were not allowed in the seventeen known dead channels. Noise was determined in the data by finding the number of fibers that fired but were not associated with a track. Noise in the Monte Carlo was simulated by assigning a small probability for a fiber to randomly fire. This probability was adjusted until the fibers not associated with a track in the Monte Carlo and the actual data agreed. The noise occupancy of a VPLC channel from this adjustment is about 3%, which is in agreement with the examination of extra hits not associated with tracks in the tracking analysis.

The input to the Monte Carlo was the number of photoelectrons produced by a particle as it traversed the center of the fiber. The number of photoelectrons produced by a track was then determined by scaling the Monte Carlo input according to how far the track was from the center of the fiber.

Using Poisson statistics, the probability for a fiber to fire was determined. Tracks were defined by the same method as in the track finding algorithm and the number of fibers that participated in a track was found. In the Monte Carlo the number of fibers that participated in a track was then plotted as a function of photoelectrons. The number of photoelectrons produced by a fiber was then calculated by comparing the number of fibers on a track in the data to the number of fibers on a track in the Monte Carlo. The photoelectron value that gave the lowest χ^2 was then multiplied by $\pi/4$ to give the average number of photoelectrons produced by a fiber. These averages are plotted in Figure IV.4 as a function of the distance from the joint between the clear fiber and scintillating fiber. An exponential fit to these data gives a light attenuation length of 8.2 m for our system of 4.3 m long scintillating fibers and 6 m long clear waveguides, both of 830 μm diameter. Separate tests, including our own, have shown that attenuation lengths are strongly dependent on method of light generation (e.g., traversal of a minimum ionizing particle or radioactive or ultraviolet source) and lengths of clear waveguides, so it is important to note that our BNL system has lengths and excitation that compare closely to what is proposed for SDC. The exponential fit gives 1.2 ± 0.1 photoelectrons at a distance of 4.3 m from the joint between the clear fiber and scintillating fiber; this distance corresponds to the worst case photoelectron yield at $\eta = 0$ in SDC. The average over the pseudorapidity region of interest in SDC ($0 < |\eta| < 2.3$) is about 5 photoelectrons.

Since the BNL beam test, we have been able to increase the number of photoelectrons in several straightforward steps:

- A factor of 1.7 increase in the number of photons has been realized by using a double-clad scintillating fiber instead of the single-clad fibers used at Brookhaven.
- The diameter of the fiber will be increased from 835 μm to 925 μm .
- The quantum efficiency of later version VLPCs is higher. At BNL VLPCs with a

quantum efficiency of 55% were used; some of the pixels in the HISTE-IV VLPCs have 85% quantum efficiency.

- Better optical connectors increase the transmission through a scintillating fiber/clear waveguide joint to as high as 95%. We believe that some of our joints at BNL were as low as 55%.
- Improvements in scintillation dye and concentration optimization will give a gain of about 20%.

With these gains we are now expecting an average of 22 photoelectrons, averaged over $0 < |\eta| < 2.3$, and 5.4 photoelectrons for the worst case photoelectron yield at $\eta = 0$. Thus, the worries about having a sufficient number of photoelectrons for high individual fiber efficiency have dissipated.

Average number of photoelectrons as a function of the distance from the joint between the scintillating fiber and clear waveguide. The solid line is the exponential fit to the data. The fit gives 1.2 ± 0.1 photoelectrons at the farthest end of the fiber, corresponding to the worst case photoelectron yield at $\eta = 0$ in SDC.

Figure IV.4

B. Supervisor of TNRLC and MOUs for the Fiber Tracking Group.

The Purdue group has played a supervisory role in the SDC scintillating fiber tracking group. This has included the management of the TNRLC funding for the group and the development of the Memorandum of Understanding between SSCL and the individual institutions forming the SDC fiber tracking group.

A major result of this role has been in setting the direction for the development of VLPCs and the development of VLPC cassettes at Rockwell International. This past year we worked out the details of a multi-task contract between SSCL and Rockwell. The goals of this contract include the development of high quantum efficiency VLPCs approaching ~80%, the development of a high density cryogenic cassette system, and finally a pilot run of VLPCs to test the manufacturability of such a system. We have completed the first task of this contract with the production of ~75,000 HISTE IV, VLPCs pixels, and have just recently started work on the next 4 tasks of the contract. This means that about half of the original contract is now in effect.

We have started the comparison of HISTE IV devices with the original Solid State Photo Multipliers, SSPMs. This will be a continuing effort throughout the rest of this year.

C. Construction of Fiber Ribbons

In order to identify the parameters of good ribbon construction and in order to develop a safe and viable handling and measuring techniques, we have constructed a number of scintillating fiber multi-layer ribbons and verified their accuracy and uniformity using a Coordinate Measuring Machine (CMM). This work has been performed in conjunction with FEPCO Coordinate Measuring Services, Noblesville, Indiana. CMM measurement has proved to be an essential tool in providing feedback in the construction of accurate fiber ribbons and in identifying important ribbon characteristics.

Purdue's ribbon production facility became operational in May of 1992. Since that time, at least 79 doublet ribbons have been made, many containing 255 channels and some as long as 4.3 m. One full-time staff technician, Tracy Crone, is employed extensively for the task of ribbon making. Ribbons are produced in a clean environment with filtered lighting (no wavelengths below 560 nm), and where humidity and temperature are monitored.

The construction technique relies on a precisely machined jig plate, to which fibers are clamped while an adhesive is applied. We use 7075 aluminum for the plate, a hard alloy suited to precise machining. V-shaped grooves are cut along the length of the plate to align the fibers. To make a ribbon, fibers are first placed in the grooves to form the lower layer of a doublet. The upper layer is then added by placing fibers in the recesses between fibers in the lower layer. This insures a complete correlation between the two layers. Foam-padded clamps that exert only a minimum of force are placed every two inches along the ribbon to hold the fibers in the jig. Figure IV.5 illustrates the process. Figure IV.6 shows the verification of the jig plate by CMM measurement. As can be seen our construction techniques have standard deviations, $\sigma < 5$ microns over areas measured in meters. These techniques are truly at the fore front of what is possible in large scale systems.

Ribbons

Figure 1

Figure 2

Figure 3

The basic element of the scintillating fiber tracker is a ribbon doublet formed of two layers of fibers. The Fiber Tracking Group has produced such ribbons for use in beam tests at Fermi and Brookhaven National Laboratories (Fig. 1). Optical measurements have verified that the reference plates for ribbon production have an accuracy of $\sigma \leq 12 \mu\text{m}$ (Fig. 2). The fiber to fiber spacing is accurate to $\sigma \leq 18 \mu\text{m}$ (Fig.3). This includes ribbons of up to 255 fibers, 4.3 meters in length. The ribbons are flexible and form accurately upon a substrate.

D. Test of Fiber Ribbon Mounting Techniques

Unlike wire chamber construction, in which wires are typically stressed so that non-linearity along a wire's length are removed during wire installation, a scintillating fiber tracker requires an entirely different mounting scheme. This occurs because fibers made of polystyrene can not tolerate any appreciable stress. This makes construction with fibers a 3 dimensional problem, whereas construction of wire chambers is a 2 dimensional problem. That is wire chambers depend on the accuracy of a 2 dimensional end plate. A fiber chamber on the other hand depends on the accurate placement in 3 dimensions of the fiber onto a 3 dimensional surface.

We have made significant progress establishing a ribbon mounting procedure that gives us complete control over where fibers and their base, a cylindrical surface, are spatially located. At any time during the process we are able to stop and verify relative positions, so that mistakes are eliminated. We are also able to know exactly where the fiber is located after it is mounted.

We have accomplished this by working inside the volume of a coordinate measuring machine, CCM. This machine is capable of reproducible measurements at the $1.5\mu\text{m}$ level over a volume large enough to accommodate construction of the D-zero upgrade outer tracker. We are working in conjunction with FEPCO services to accomplish these tests. Figure IV.7 shows the Mark I version of our ribbon laying machine used to build up a very accurate 3 dimensional scintillating fiber coordinate system. Figure IV.8 shows the accuracy to which the fibers have been placed along the length of the surface of the cylinder, $\sigma \sim 20\mu\text{m}$. This work is continuing.

V. D-Zero

The D-Zero group will collect $\sim 13\text{pb}^{-1}$ of data during its first data run at the collider. The Purdue group has started to analyze this data set already. There are two physics topics of interest to our group.

1. Task E has had a long history of working on problems associated with understanding the tau lepton. With this experience gained at e^+e^- colliders, we are now attempting to apply some of these same techniques to the analysis of tau lepton physics problems at the tevatron. We have started by isolating a sample of $Z \rightarrow \tau^- \tau^+$ events. Figure V.1 shows such an event. These are very clean events containing an electron and a muon from the decay of a pair of tau leptons. From these measurements, we can check to see if the $Z_{(e^+e^-)}$ created in e^+e^- collision is the same $Z(q\bar{q})$ created in hadronic collisions by comparing the decay widths with those measured at LEP. The tau lepton also serves as an important particle to use in tests of lepton universality because it is the heaviest lepton. These analysis are just in the beginning stages.

2. We are also looking at the thermodynamic properties of the production of multi-jet events. One of the striking differences between e^+e^- machines and hadron machines is the large number of hadronic or QCD events. D-Zero now has ~ 3 M hadronic events written to tape. Looking at the data in a very preliminary manner, we have identified events with as many as 7 and 8 and more jets per event in the central rapidity region. If after careful analysis this interpretation of high jet multiplicity in the events holds true,

LATERAL GROOVE DISPLACEMENT ON JIG PLATE

Figure IV.6 Deviation of grooves cut in a ribbon-making jig plate from absolute straightness as measured by CMM.

Figure IV.8

Max ET= 19.8 GeV
CAEH ET SUM= 163.7 GeV
VTX in Z= -39.7 (cm)

Figure V.1

questions about the underlying production of such large numbers of jets can be asked. For example, are plasmas formed or are these simply very high order radiative events. The number of events with large jet multiplicity is becoming large enough that these questions might be answered by studying thermal dynamic variables. We have just started these studies.

VI. Publications and Presentations

Test Beam Results using Scintillating Fibers Readout by a Multianode Phototube and Visible Light Photon Counters, B. Abbott et al., submitted to Nuclear Instruments and Methods in Physics Research.

Precision Measurement of the Total Cross Section and Charge Asymmetry at TOPAZ for $e^+ e^- \rightarrow \mu^+ \mu^-$ and $e^+ e^- \rightarrow \tau^+ \tau^-$, written for the XXVI International Conference on High Energy Physics, October, 1992.

Cross Sections and Charge Asymmetries for $e^+ e^- \rightarrow \mu^+ \mu^-$ and $e^+ e^- \rightarrow \tau^+ \tau^-$ Reactions at $\sqrt{s} = 52 - 61.4 \text{ GeV}$, B. Howell et al., Phys. Lett. B291, Sept. 17, 1992, 206-212.

Studies of $e^+ e^- \rightarrow \gamma\gamma$ and $\gamma\gamma\gamma$ reactions, K. Shimozawa et al., Phys. Lett. B284 (1992) 144-150.

An Experimental Study of Muon Pair Production in Tagged Two-Photon Interactions, H. Hayashii et al., Phys. Lett. B279, (1992) 422-428.

Experimental Study of b-quark Jets in $e^+ e^-$ annihilation at TRISTAN, K. Nagai et al., Phys. Lett. B278, (1992) 506-510.

Tracking With Scintillating Fibers and Visible Light Photon Counters, M. Atac et al., Nuclear Instruments and Methods in Physics Research A320, (1992) 155-160.

Experimental Study of b-quark Jets in $e^+ e^-$ Annihilation at TRISTAN, K. Nagai et al., Phys. Lett. B278 (1992) 506.

The TOPAZ Vertex Chambers Ceramic Endplate, M. Kobayashi et al., Nuclear Instruments and Methods in Physics Research A312 (1992) 440-450.

Purdue University
Contract DE-AC02-76ER01428 (Task F)

A THEORETICAL HIGH ENERGY PHYSICS PROGRAM

within the

**DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY
West Lafayette, Indiana 47907-1396**

Supported by

U.S. DEPARTMENT OF ENERGY

Through

**Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana**

Current Contract Number: DE-AC02-76ER01428

TASK F

**ANNUAL PROGRESS REPORT
For May 1992 - April 1993**

Investigator: E. Fischbach

**Graduate Students: H. Kloor
D. Krause**

I. TASK F OVERVIEW

Our group is continuing in its program of searching for phenomena which probe elementary particle physics beyond the standard model. This effort has produced a number of new and interesting results this year, along with a major review article on non-Newtonian gravity [*Metrologia* 29, No. 3 (September 1992), pp. 215-260. *Metrologia* is the journal of the International Bureau of Standards in Paris].

Our program is motivated in part by the view that in the period before the SSC comes on line the opportunity exists to explore interesting and fundamental questions in elementary particle physics using low-energy experiments. The unity of physics is such that the results of low-energy experiments bear directly on current and future experiments at the highest energies.

As an example of the connection between high-energy and low-energy phenomena we review the motivation for the ongoing interest in the search for non-Newtonian gravity. We note that there are two natural mass scales that arise in elementary particle physics, the hadron scale characterized by the nucleon mass $m_N \cong 1\text{GeV}$, and the Planck scale $M_{Pl} = (\hbar c/G)^{1/2} \cong 10^{10}\text{GeV}$, where G is gravitational constant. From these constants one can form the mass scale

$$m_o = \frac{m_N^2}{M_{Pl}} \cong 10^{-10} eV,$$

and if m_o corresponds to the mass of a new bosonic field, then the exchange of this field would give rise to a long-range force extending over a distance scale of order 2000 m. Moreover, in a theory of such a long-range bosonic force, the dimensionless ratio $f = m_N/M_{Pl}$ can play the role of the coupling constant of this field (the analog of the electric charge e). Since $f \cong 10^{-19}$ such a coupling would not only have a strength comparable to gravity, but also a range which allows its effects to compete with gravity over macroscopic distances.

It follows from the preceding discussion that searches for deviations from the predictions of Newtonian gravity are simply a low-energy tool from probing elementary particle physics at the highest energy scale, which is the Planck scale. This observation forms a significant part of the theoretical and experimental motivation for the continuing study of possible non-Newtonian gravitational effects.

II. NEW CONSTRAINTS ON LIGHT AXIONS AND MACROSCOPIC-RANGED PSEUDOSCALAR FORCES [Proceedings of the XIIIth Moriond Workshop, Villars, Switzerland, January 31 - February 6 1993, to be published.]

We compare the laboratory constants on the couplings of light pseudoscalars (such as axions) to fermions, to those few light scalars and vectors. We begin by demonstrating why it is that the constraints on pseudoscalars are many orders of magnitude weaker than for vectors and scalars. This then leads to the proposal for a new mechanism which would significantly strengthen the limits that could be obtained via laboratory experiments.

The conventional parity (P)- and time-reversal (T)-conserving coupling of an axion or other light pseudoscalar to fermions is given by

$$\mathcal{L} = -ig_P \psi \gamma_5 \psi \phi \quad (1)$$

where ϕ is the axion and ψ is the fermionic field. The lowest order CP-conserving potential between two fermions arising from axion-exchange is

$$V_P(r) = \frac{g_P^2}{4\pi} \left(\frac{1}{2mr} \right)^2 \left[\frac{3(\vec{\sigma}_1 \cdot \hat{r})(\vec{\sigma}_2 \cdot \hat{r}) - \vec{\sigma}_1 \cdot \vec{\sigma}_2}{r} \right] \quad (2)$$

where m is the fermion mass, $\vec{r} = \vec{r}_1 - \vec{r}_2$ is their separation, and $\vec{\sigma}_{1,2}$ are the fermion spins. The form of this potential can be contrasted with the more familiar scalar-exchange potential V_S ,

$$V_S(r) = -\frac{g_S^2}{4\pi} \frac{1}{r}, \quad (3)$$

where g_S is the scalar coupling constant. A variety of spin-dependent experiments [1,2] set a limit on a long-range force arising from axion-exchange given by [3]

$$\frac{g_P^2}{4\pi} \lesssim 10^{-12}. \quad (4)$$

By contrast, the analogous limit on g_S^2 is [1]

$$\frac{g_S^2}{4\pi} \lesssim 10^{-41} \quad (5)$$

for forces with a range of $\approx 1m$. Since both sets of experiments use similar torsion-balance technology, and in some cases have been carried out by the

same group, the question naturally arises as to why the sensitivities of these experiments differs by a factor of order 10^{29} . The answer lies in the factor $(1/2mr)^2$ in Eq.(3), which arises from the presence of γ_5 in Eq.(1). For the exchange of an axion with Compton wavelength of 1m, this suppression factor is

$$(1/2mr)^2 \cong 1 \times 10^{-26} \quad (6)$$

which accounts for most of the suppression of V_{PS} compared to V_P .

The natural question is then whether this suppression can be circumvented, which would then lead to significantly improved constraints on g_P . To this end we have considered two new mechanisms, including 2-axion exchange and the exchange of axions between bound systems. For the 2-axion exchange mechanism we have evaluated for the first time the potential energy arising from the exchange of axions of arbitrary mass. The general expression is complicated, but in the massless limit it reduces to

$$V_P(2 - axion) = \frac{g_P^4}{64\pi^3 m^2 r^3} = \left(\frac{g_P^2}{4\pi}\right)^2 \left(\frac{1}{2mr}\right)^2 \left(\frac{1}{\pi r}\right) \quad (7)$$

This expression still contains the suppression factor $(1/2mr)^2$, and hence does not represent an improvement over the expression in Eq.(3), despite the elimination of the spin-dependent terms.

However, consideration of *bound system* effects does make a substantial improvement. We have shown that in this case the suppression factor $1/2mr$ is replaced by

$$\frac{\hbar/m_{NC}}{\text{size of bound system}} \quad (8)$$

which for typical systems is much closer to unity. This suggests that we should be able to extract more stringent limits on axion couplings from laboratory experiments, as we discuss in [3]. Detailed calculations to extract these limits are currently under way, as we describe in more detail in the accompanying Proposal.

III. THE EQUIVALENCE PRINCIPLE AND WEAK INTERACTIONS

[Invited talk, Proceedings of the Symposium on Satellite Tests of the Equivalence Principle (STEP), Pisa, Italy, April 6-8 1993, to be published] with: M.P. Haugan and D. Tadić.]

We have reconsidered the possibility that gravity, or another long-range force, may couple anomalously to the weak interactions, and thereby lead to a violation of the Weak Equivalence Principle (universality of free-fall). We show that at the level of sensitivity expected in the STEP experiment being planned by the European Space Agency, detection of an acceleration difference between the test masses could lead to fundamental new constraints on the couplings of various long-range forces to neutrinos, and to higher-order weak interactions. In particular this will allow the first limits to be set on the coupling of gravity to energy arising from the parity-violating weak interaction.

If the weak interactions couple to gravity (or another long-range force) in a manner that violates the Equivalence Principle (EP) then it can be shown that the gravitational mass $m_G^{(i)}$ of test sample i and its inertial mass $m_I^{(i)}$ are related by

$$m_G^{(i)} = m_I^{(i)} + \eta_W E_W^{(i)} / c^2 \quad (1)$$

where $E_W^{(i)}$ is the weak energy, and η_W is the parameter which measures the breakdown of the EP (if $\eta_W \neq 0$). When $\eta_W \neq 0$, this gives rise to an EP-violating acceleration difference between two samples given by

$$\frac{a^{(1)} - a^{(2)}}{a} = \eta_W \left[\frac{E_W^{(1)}}{m_I^{(1)}} - \frac{E_W^{(2)}}{m_I^{(2)}} \right] \quad (2)$$

where a is their average acceleration.

In this paper we calculate for the first time the contributions to E_W which are second-order in the Fermi constant G_F . The purpose of this calculation is two-fold: a) In second order of the weak Hamiltonian H_W this probes the coupling of gravity to the *parity-violating* weak interactions and b) in second order, gravity can also couple to virtual neutrinos, and hence probe for the first time for any anomalous coupling of gravity to neutrinos. Our initial calculations suggest that we could set the following limits on the EP-violating parameters η_μ and η_τ , which describe the coupling of gravity to $\nu_\mu(\bar{\nu}_\mu)$ and $\nu_\tau(\bar{\nu}_\tau)$:

$$\eta_\mu \lesssim 1 \times 10^{-2} \quad (3a)$$

$$\eta_\tau \lesssim 1 \times 10^{-8} \quad (3b)$$

Motivated by these initial results, we are undertaking more detailed calculations, as we describe in the accompanying Proposal.

References

1. See E. Fischbach, et al., Metrologia **29**, 215 (1992).
2. R.C. Ritter, et al., Phys. Rev. **D42**, 977 (1990).
3. D.K. Krause, E. Fischbach, C. Talmadge, D. Sudarsky, and D. Tadić, Proceedings of the XIIIth Moriond Workshop, Villars, Switzerland, January 31 - February 6, 1993, (to be published).

AN EXPERIMENTAL HIGH ENERGY PHYSICS PROGRAM

within the

DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY
West Lafayette, Indiana 47907

Supported by
U.S. DEPARTMENT OF ENERGY

Through
Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana

Current Grant Number: DE-FG02-91ER40681

Task G

PROGRESS REPORT
for the period ending
1 April 1993

Principal Investigators: V. Barnes, D. Bortoletto, A. Garfinkel

Spokesperson: V. E. Barnes (317) 494-5199 FTS & Commercial

Contents

I	PROGRESS REPORT	2
1	Introduction to Progress Report and Task G personnel	2
2	The Collider Detector at Fermilab (CDF)	3
2.1	Introduction	3
2.2	History of Purdue Involvement in CDF Hardware	3
2.3	People Involved	4
2.4	The 1992-83 CDF Run	4
2.5	Preparations for the 1992-93 run 1a by Purdue people.	4
2.6	Activities during the current run by Purdue people.	5
2.7	Preparation for run 1b by Purdue people.	5
2.8	Progress toward run 2 (1995) by Purdue people.	6
2.9	Physics analysis of 1988-89 data by Purdue people.	6
2.10	Physics analysis of 1992-93 data by Purdue people.	7
3	The SDC Detector for pp Collisions at the SSCL (Calorimeters)	8
4	Publications since 1985	9
5	Ph.D. Theses	16

Part I

PROGRESS REPORT

1 Introduction to Progress Report and Task G personnel

This progress report summarizes our research effort for the period from November 1 1992 to April 1993.

The members of task G are Professors Virgil Barnes, Daniela Bortoletto, and Arthur Garfinkel; Project Associates Robert De Bonte (shared) and Alvin Laasanen; Research Associates Qifeng Shen and Mark Shaw; Technicians Ernie Beard and Ian Rickey (both shared); and graduate students Mark Bailey, Todd Huffman (Ph.D. 12/92), Mark Kruse, and James Tonnison.

Task G includes founding members of the CDF and SDC collaborations and is devoted to these two efforts. Since the Purdue group had joint responsibility for the design and fabrication of the CDF central hadron calorimeters it had a strong interest in their use in doing jet related physics. The techniques developed, in collaboration with other CDF members, continue to contribute to many of the CDF physics results from measurements of the inclusive jet spectrum to measurements of the W Boson mass and Pt distribution. We are pleased that our CDF Ph.D. Steve Kuhlman, now at Argonne, continues to serve as co-convenor of the CDF QCD analysis group and that an analysis originated by the Purdue group has led to a March 1993 PRL entitled, "Comparison of Jet Production in $p\bar{p}$ Collisions at $\sqrt{s} = 546$ and 1800 GeV.

In recent times our emphasis has moved from jet physics and QCD to B physics and the top search. This induced us to take an active role in the fabrication, installation, and running of the SVX silicon vertex detector. Mark Bailey, who is a major player in the DAQ system for the SVX is currently doing thesis research on B production through studies of exclusive final states. Postdoc Shaw, who was responsible for quality control and testing of the silicon detectors, is using the SVX to tag B jets associated with top production. Todd Huffman received his Ph.D. in December of 1992 for a thesis measuring B production from a study of the inclusive muon cross section, which is being prepared for submission to PRL. Huffman, along with Jim Tonnison had contributed to the fabrication and testing of the VTX intermediate tracking chamber. Tonnison is doing a thesis on the search for top in the current run's data.

Daniela Bortoletto joined the Purdue faculty, after being a CLEO postdoc, in July of 1992. She became a member of Task G in November 1992 when it split off from Task D. With her considerable expertise and interest in B physics she led Purdue into the SVX II project to build a silicon vertex detector for run 2 in about 1995. A new student Mark Kruse

is extending the DAQ efforts of Bailey et al. for use with the SVX' detector for run 1a to start in October of 1993.

On the hardware front we continue to work closely with the Fermilab CDF staff on a number of projects. For run 2 we are also participating in the design and fabrication of an upgraded end-plug calorimeter based on the use of scintillating tiles with wave-length shifting fiber readout. As members of the SDC collaboration for the SSC, we are engaged in the design and construction of the central calorimeters. These calorimeters are rather similar to those being used in the CDF end-plug upgrade and hence there is a considerable degree of synergism between these two efforts. In both efforts we are emphasizing the use of radioactive sources in calibration procedures. Qifeng Shen, who is involved with these calorimeter efforts is interested in searching for W' massive charged vector bosons in CDF data.

2 The Collider Detector at Fermilab (CDF)

2.1 Introduction

In the past year, CDF has embarked on run 1a which promises to deliver approximately 25 pb^{-1} of integrated luminosity as compared to about 4 pb^{-1} in the previous run in 1988-89. The current run is characterized by the use of the SVX silicon vertex detector and improved muon detection. Physics goals include a search for the top quark, an improved measurement of the W boson mass, as well as a large amount of B physics. The Purdue group is principally involved in the top search and in B physics.

2.2 History of Purdue Involvement in CDF Hardware

Purdue had joint responsibility with the Italian Groups at INFN-Frascati and INFN-Pisa for the fabrication, testing, and calibration of the 50 (thirteen-ton) Central and 50 (seven-ton) End Wall hadron calorimeters. In order to fabricate the steel modules, Purdue HEP set up a special shop in conjunction with the Purdue Central Shop. In it, we made use of a computer controlled thermal machining center which cut steel plate by means of a plasma arc. The system was both highly accurate and cost effective. The finished scintillator was provided by the Frascati-Pisa group. The calorimeters were then assembled and cosmic ray tested at Fermilab. The finished calorimeters were all then calibrated in a Fermilab test beam. We also had responsibility for the fabrication, testing, and calibration of the TDC (time to digital) electronics for these calorimeters. More recently, we have been involved in the electronics for the VTX vertex time projection chamber.

We have also played a major role in the fabrication of the SVX silicon vertex detector by taking the responsibility for the quality control and testing of the silicon detectors. In addition to which, we have shared with Fermilab the responsibility for the SVX data acquisition

system. Both the VTX and SVX chambers are operating successfully in the current run.

2.3 People Involved

Faculty members working on CDF are Professors Barnes, Bortoletto, and Garfinkel. Post-doctoral physicists are Drs. A. Laasanen and R. De Bonte (shared), Project Associates, as well as Research Associates Mark Shaw and Qifeng Shen. Thesis students working on CDF are Mark Bailey, Todd Huffman (Ph.D. 12/92), Mark Kruse, and Jim Tonnison. Drs. Shaw, Shen and the students are based at FNAL.

2.4 The 1992-83 CDF Run

The latest run of the Tevatron started in August 1992. The machine delivered a record peak luminosity of $8.0 \times 10^{30} \text{ cm}^{-2}\text{sec}^{-1}$. Up to April 1993, CDF has accumulated about 15 pb^{-1} of integrated luminosity. The data were primarily taken at a center of mass energy of 1.8 TeV. The experiment was capable of being triggered on transverse energy clusters, electrons, muons, or minimum bias events in addition to specialized triggers. A level III trigger was implemented which permits limited track reconstruction and sophisticated calorimetry cleanup. The number of J/ψ collected per pb^{-1} in the central muon chambers alone is more than a factor of four higher than last run (1988-1989), due to the removal of a CFT track requirement for one muon leg. This change also softens the p_T spectrum of J/ψ to which we are sensitive. This run is scheduled to end in June with projected integrated luminosity of $\approx 25 \text{ pb}^{-1}$.

2.5 Preparations for the 1992-93 run 1a by Purdue people.

Graduate students Todd Huffman and Jim Tonnison, based at Fermilab, have worked with members of the Fermilab staff on the VTX time projection chamber which has replaced the VTPC chamber for this run. Huffman has been concerned with design, production, and testing of readout electronics for the VTX. The readout boards containing new low noise preamplifiers met specifications and are currently fully operational on the detector. Tonnison was mainly concerned with developing tracking code for the VTX.

Postdoc Mark Shaw and student Mark Bailey worked at Fermilab on the new silicon vertex detector (SVX). Shaw took responsibility for the quality control of the silicon detectors as delivered from the manufacturer. This was a vital part of the process of defining acceptance standards and refining acceptable production techniques. The SVX has been functioning extremely well in the run to date. Our present chief concern regards the radiation hardness of the device as the run progresses. Mark Bailey developed diagnostic tests for the SVX data acquisition system in a program called SVX-CONTROL. This program enabled him to automate debugging of the fabricated circuit boards. He and his coworkers

completely tested all elements of the SVX front end electronics using the standard CDF data acquisition program called Run Control. He also played a major role in dealing with cross talk problems between the SVX and VTX. Various improvements were made to shielding and timing patterns which have eliminated the problem. Shaw also spent a fraction of his time working on track reconstruction and analysis software for the SVX.

2.6 Activities during the current run by Purdue people.

While all the Purdue people stand 10 day shifts periodically, the postdocs and students based at Fermilab have taken on special responsibilities to insure the proper functioning of the apparatus and the quality of the data. Mark Shaw has been heavily involved in the maintenance and monitoring of the SVX calibration, including: maintenance of the SVX calibration offline software, daily monitoring of leakage currents, pedestals, and other calibration constants, updating of the CDF database to include up-to-date SVX calibration constants, monitoring of SVX bias supply currents (leakage and that drawn by the guard ring), as well as monitoring the SVX occupancy. Mark Bailey is generally on call to deal with any problems that arise with regard to SVX data acquisition electronics or software. Both he and Shaw have also volunteered to stand "Off-line Shifts" which involves spending a couple of hours a day to monitor the quality of the data. Qifeng Shen has volunteered to be part of shift pool and has been working recently on monitoring the YMON program that checks the performance of much of the hardware, looking for dead or noisy channels etc. Alvin Laasanen continues to have responsibility for the calibration of the hadron calorimeter TDC system.

2.7 Preparation for run 1b by Purdue people.

Purdue is preparing in the following ways for run 1b which is currently scheduled to begin in October of 1993. We are continuing many of the activities we performed for SVX again for SVX'. The SVX' silicon detector system makes use of ac-coupled silicon detectors and a radiation hard SVXH readout chip. Mark Shaw is upgrading SVX calibration offline software for use on a test bench during SVX' construction. He is also adding new calibration routines specific to SVX' as well as having contributed to the quality control system for SVX' detector testing. Mark Kruse is working on the data acquisition system for SVX'. He has modified an SVX standalone data acquisition system for the SVX' which allows calibrations to be done on any test structure, an "ear", "ladder", or up to 6 wedges located in either B0 or IB4 labs. In particular the program SVX-CONTROL replaces the standard data taking program RUN-CONTROL and the SSP and EVENT-BUILDER's hardware functions are assumed by a VAX. This permits measurements of pedestals (whose width are a measure of noise), gains (ie. charge calibrations), thresholds (for latching), and leakage current.

2.8 Progress toward run 2 (1995) by Purdue people.

- End-Plug Calorimeter Upgrade:

Barnes, Laasanen, Shen, and Tonnison are working on the end-plug upgrade project in which the gas based calorimeters will be replaced with scintillating tiles with fiber readout; a technique well suited to use at the SSC, as well as at the higher luminosities and shorter times between bunch crossings planned for future CDF runs. Purdue has been active in the design and Monte Carlo simulation of the optics of these scintillating tiles. Purdue has taken responsibility for the calibration of both the Electromagnetic and the Hadronic compartments of these calorimeters, using a remotely driven moving radioactive sources. We have developed and successfully used this design of source driver in a variety of prototype calorimeters in the past year, and Purdue has built and used such source drivers in all scintillator calorimeter prototypes, both CDF and SSC, in the recent Fermilab test beam, as well as a thin-Lead SDC EM calorimeter prototype tested at BNL in the summer of 1992. The source measurements are important for characterizing the quality and uniformity of construction of the calorimeters, as well as for carrying the results of test beam calibrations through time.

These drivers are also being used in a set of Radiation Hardness-testing small EM calorimeter modules ("Personal Calorimeters") which have been built jointly by Purdue and Fermilab and shipped to Beijing, Saclay, and KEK, for irradiation to high levels in 1 GeV electron beams. The moving sources access every tile in the calorimeter and are our primary way of assessing the progressive degradation of the tiles as a function of dose.

- SVX II Silicon Vertex Detector

Bortoletto, Garfinkel, and Shaw are working on SVX II which will be the 3d generation silicon vertex detector used by CDF. A proposal for SVX II has recently been submitted to Fermilab. The proposed device is characterized by an increased length (3 barrels) as compared to SVX and SVX' (2 barrels). It will be the first use of double sided detectors which provide Z readout in addition to $r - \phi$ measurements. We have contributed to the proposal writing and are now engaged in obtaining and testing sample detectors. A particularly critical aspect of the detectors is their capacitance and the implication for signal to noise ratio. Furthermore, the technology for getting access to all the signal lines on both sides of the detectors needs to be tested and finalized. At the moment all testing is being done at Purdue University School of Electrical Engineering. The availability of testing equipment in the Purdue University Physics Department will allow a more efficient set-up.

2.9 Physics analysis of 1988-89 data by Purdue people.

Analyses using the 1988-89 data in which Purdue played a major roll include a search for stable charged particles, entitled, "Limits on the Production of Massive Stable Charged

Particles", which appeared in Phys.Rev. Rapid Communications in Sept. 1992. We have succeeded in ruling out the existence of unit-charged color triplets with masses between 50 and 139 GeV/c² @95% c.l., with similar limits for particles of charge 2/3 or 4/3. For unit-charged color sextets, octets, and decuplets we exclude @ 95% c.l. masses between 50 GeV/c² and 197, 199, and 255 GeV/c² respectively.

Another analysis involving a test of QCD scaling violation in a comparison of the inclusive jet cross sections at 540 and 1800 GeV entitled, "Comparison of Jet Production in $p\bar{p}$ Collisions at $\sqrt{s} = 546$ and 180 GeV", appeared in PRL 70,1376(1993). Measurements were made of the ratio of the CDF cross sections at 546 and 1800 GeV as a function of the scaling variable $x_t = 2E_t/\sqrt{s}$. Comparisons to next-to-leading order QCD calculations have been made which show deviations at the 1.5 - 2.4 σ level.

A third analysis involving b quark production, as measured by the inclusive muon spectrum, (Todd Huffman's thesis) is being prepared for submission to PRL. It shows that the b quark cross section is significantly above predictions of a next-to-leading order QCD calculation. The results were presented by Huffman at the Chicago DPF meeting.

2.10 Physics analysis of 1992-93 data by Purdue people.

Our interest in the new data largely centers on heavy quark production. Mark Bailey is doing thesis research on B meson production by measuring the exclusive final states $B^- \rightarrow \psi K^-$ and $B^0 \rightarrow \psi K^*$ and charge conjugate modes. Figure 3.1 shows the charged B spectrum with a signal of 60 ± 14 events. Figure 3.2 shows the corresponding differential cross section $d\sigma/dp_T$, which is measured here for the first time. His analysis indicates a discrepancy between the data (solid circles) and the $\mathcal{O}(\alpha_s^3)$ theoretical predictions. (solid line). Figure 3.3 shows the neutral B spectrum with a signal of 24 ± 12 events. Figure 3.4 shows the b quark cross sections derived from these measurements as filled circles as well as those from previous CDF measurements including those from Huffman's thesis ($B \rightarrow \mu X$).

Jim Tonnison is doing thesis research on the top search in the lepton + multi-jet topologies. Mark Shaw has played a major role in the development and testing of b-tagging algorithms using the inclusive electron sample and the SVX silicon vertex detector. He is involved in a top search in the lepton + jets channels which emphasizes the use of the SVX for b tagging. Qifeng Shen is working on a search for heavy W' gauge bosons in the channels $\mu + \nu$ and $e + \nu$. The senior people guide and supervise these activities as well as pursuing related analysis interests.

3 The SDC Detector for pp Collisions at the SSCL (Calorimeters)

The personnel who have worked on this project are Professors Virgil Barnes and Arthur Garfinkel, project associate Alvin Laasanen and graduate students James Tonnison and James Ross, as well as several undergraduate students during summers and during the academic year. We are founding members of the Solenoidal Detector Collaboration (SDC), and Barnes has been appointed a member of the SDC Calorimeter Technical Board. One of our long-range goals is to collaborate in building the calorimeters for the proposed Solenoidal Detector for the SSC. This initiative began in close collaboration with some of our Fermilab CDF colleagues, and is aimed not only at the SSC, but also at the upgrade of the CDF endplug calorimeter. A ray-tracing Monte Carlo program developed by Barnes and Tonnison gave the first indications that excellent light yield and uniformity of response were possible for tiles with small diameter waveshifting fiber readout. This was the inception of the interest in this technology, which has led to its being the technology of choice for the CDF endplug upgrade for 1995 to replace the gas readout calorimetry in the CDF plug and forward regions.

This technology has now also been chosen for all or most of the SDC calorimetry. Test beam studies of prototype EM and Hadron calorimeters at Fermilab in late 1991 gave very encouraging results. Purdue provided calibrations, using moving radioactive sources, of these prototypes, and also of CDF endplug upgrade EM and Hadron calorimeter prototypes which are directly relevant to SDC calorimeter design choices. A thin-lead SDC EM calorimeter prototype was tested at BNL in the summer of 1992 and showed the potential to use radioactive source calibrations in lieu of test beam calibrations of most towers of a calorimeter, at the 2 - 3 % accuracy level. Results obtained for an earlier spaghetti calorimeter during and after the August 1990 Fermilab test beam run were presented at the Fort Worth SSC Conference and the Fermilab International Calorimeter Conference, both in the fall of 1990, and are in the published proceedings of those conferences. Preliminary results of the fall, 1991, calibration of the CDF plug upgrade EM and Hadron calorimeter prototypes at the testbeam were reported at the October, 1991 Capri Calorimeter Conference and are in the published proceedings.

The CDF plug upgrade EM prototype was extensively instrumented with source tubes in every layer of every tower. It achieved good energy resolution: the calorimeter energy response has a σ/E of 1.6% for 100 GeV electrons. The relation between beam energy and calorimeter resolution for a typical tower, is $\sigma/E = 15.2$ /tower to the moving Cs137 calibration source were equalized, using grey-scale masking at the phototube cookie. This technique can compensate for imperfections in production and installation of the scintillating tiles, or can compensate for possible radiation damage of the tiles.

We have had responsibility for the calibration and characterization of every one of the CDF Endplug and SDC Barrel scintillating tile calorimeter prototypes for the Fermilab beam tests. We are also helping with radiation hardness studies taking place at various

foreign laboratories which have $\approx 1 - 2$ GeV intense electron beams. We constructed two years ago twenty small Radiation-Hardness- testing calorimeters ("Personal Calorimeters") in collaboration with Fermilab, some of which were shipped to Beijing, to Saclay and to KEK. Radiation hardness adequate for the SDC barrel calorimeter and the SDC endcap calorimeter has been seen in these modules, with scintillators and wavelength shifting fibers of the type used in the recent testbeam prototypes. The $\approx 40\%$ degradation at EM shower maximum at a dose of ≈ 600 kiloRads (10 SSC years at 10^{31} luminosity) could be corrected by differential grey-scale masking of the fibers from each tile. Even in the absence of masking, the damage profile, measured by the moving source, could be used in software correction of the response of the calorimeter as a function of electron energy. If a second EM calorimeter depth segment is available, software correction can also be made for shower-to-shower depth fluctuations, thereby recovering most of the energy resolution of an undamaged calorimeter.

We have also used a moving source calibration to obtain longitudinal response profiles of the tiles in a prototype module of the CDF Microplug, a calorimeter proposed for installation at very forward angles just outside the CDF beam pipe. The Microplug performed well in the 1991 testbeam. One proposed use of the Microplug is as a radiation damage test bed for various promising new scintillating tiles and readout fibers in realistic radiation fields, where the moving source can monitor the development of the damage profile.

The source driver system is described in the accompanying Proposal document, where ongoing research and development activities are also described. Considerable effort in the past year has gone into calculations and measurements relating to the accuracy achievable with moving radioactive gamma sources. Considerable effort has also gone into design of the source tubes and connectors which are being incorporated into the SDC EM and Hadron calorimeter prototypes which are being fabricated at this time for study later in 1993. Our calculations alerted the collaboration to the fact that locating the source tubes in grooves machined in the hadron steel would lead to excessive sensitivity of the tile response to the source position. We have designed a spring steel strip which lifts the source tube out of the groove and into direct contact with each tile, leading to a much better, and acceptable, level of position sensitivity. As described in the accompanying Proposal, mechanical testing and design upgrades of the source driver and the source wire have also been important efforts.

In summary, we have had a very productive, albeit exhausting, year working on the new scintillating tile technology for calorimetry, and the calibration thereof. We are delighted that the technology is proving so successful and has been chosen for both the SDC and CDF detectors.

4 Publications since 1985

Numbering of publications starts in 1985. For a full list of publications see the 1987 progress report, which goes back to June 1, 1975, at the inception of Task D.

1. Charge Distributions of Hadrons Associated with Hadronic J/ψ Production, H.S. Budd, G.O. Alverson, T.L. Graff, S.R. Hahn, L.E. Holloway, L.J. Koester, U.E. Kruse, W.G. Li, P.T. Lukens, R.D. Sard, P. Schoessow, D.A. Bauer, D.J. Judd, T.B.W. Kirk, T. Lagerlund, S.H. Pordes, R. Raja, L.D. Spires, A.A. Wehmann, J.W. Cooper, V.E. Barnes, C. Davis, R.E. Davis, A.F. Garfinkel, A.T. Laasanen, S. Hossain, R.H. Milburn, and R.K. Thornton, Phys. Rev. D31, 1132 (1985).
2. Design and Fabrication of the CDF Large Angle Hadron Calorimetry, A.F. Garfinkel, V.E. Barnes, K. Chadwick, A. Di Virgilio, S. Kuhlmann, A.T. Laasanen, J.K. Simmons, S. Bartalucci, S. Bertolucci, M. Cordelli, M. Curatolo, B. Exposito, P. Giromini, S. Miozzi, A. Sansoni, F. Bedeschi, G. Bellettini, L. Bosisio, F. Cervelli, R. Del Fabbro, E. Focardi, S. Galeotti, P. Giannetti, M.A. Giorgi, A. Menzione, D. Passuello, L. Ristori, A. Scribano, P. Sestini, A. Stefanini, G. Tonelli, H. Jensen, and H. Kautzky, Proceedings of the Annual DPF Meeting, Santa Fe, NM (1984), p368.
3. Performance of the CDF Central Electromagnetic Calorimetry, P. Schoessow, V. Barnes, S. Bertolucci, W. Burley, K. Chadwick, S. Cihangir, D. Connor, J. Cooper, R. Diebold, J. Elias, A. Garfinkel, S. Hahn, H. Jensen, T. Kamon, H. Kautzky, S. Kobayashi, K. Kondo, A. Laasanen, S. Mikamo, M. Miller, A. Murakami, L. Nodulman, J. Proudfoot, D. Quarrie, A. Sansoni, J. Simmons, N. Solomey, K. Takikawa, D. Underwood, R. Wagner, V. White, A. Wicklund, H. Williams, A. Yamashita, and K. Yasuoka, Proceedings of the Annual DPF Meeting, Santa Fe, NM (Oct. 31 - Nov. 3, 1984), p366.
4. Performance of a Lead Glass Spectrometer at High Energies, A.T. Laasanen, V.E. Barnes, C. Davis, A.F. Garfinkel, J.K. Simmons, J.R. Wilson, A.A. Wehmann, G.O. Alverson, J.N. Bellinger, N.S. Budd, S. Cihangir, T.L. Graff, S.R. Hahn, I. Karlinger, L.J. Koester, W.G. Li, P.T. Lukens, R.D. Sard, M.A. Shupe, D.A. Bauer, J.W. Cooper, S. Hossain, R.H. Milburn, W.P.B. Oliver, and R.K. Thornton, Nucl. Inst. & Methods, A236 307 (1985).
5. Differences Between Proton and π^- Induced Production of the Charmonium χ States, D.A. Bauer, T.L. Graff, P.T. Lukens, J.R. Wilson, G.O. Alverson, G. Ascoli, V.E. Barnes, J.N. Bellinger, S. Cihangir, J.W. Cooper, C. Davis, A.F. Garfinkel, L.E. Holloway, I. Karliner, T.B.W. Kirk, L.J. Koester, A.T. Laasanen, W.G. Li, W.P. Oliver, S.H. Pordes, R.D. Sard, J.K. Simmons, R.K. Thornton, and A.A. Wehmann, Phys. Rev. Lett. 54, 753 (1985).
6. Calibration of the CDF Central Electromagnetic Calorimetry, S. Hahn, V. Barnes, S. Bertolucci, W. Burley, K. Chadwick, S. Cihangir, D. Connor, J. Cooper, R. Diebold, J. Elias, A. Garfinkel, H. Jensen, T. Kamon, H. Kautzky, S. Kobayashi, K. Kondo, A. Laasanen, S. Mikamo, M. Miller, A. Murakami, L. Nodulman, J. Proudfoot, D. Quarrie, A. Sansoni, P. Schoessow, J. Simmons, N. Solomey, K. Takikawa, D. Underwood, R. Wagner, V. White, A. Wicklund, H. Williams, A. Yamashita, and K. Yasuoka, Proceedings of the Annual DPF Meeting, Santa Fe, NM (Oct. 31 - Nov. 3, 1984), p365.

7. Central Jets in the CDF Experiment, Arthur F. Garfinkel, Proceedings of the International Europhysics Conference on High Energy Physics, ed. by O. Botner, Uppsala University (1987).
8. Compositeness and QCD at the SSC, V. Barnes, B. Blumenfeld, R. Cahn, S. Chivukula, S. Ellis, J. Freeman, C. Heusch, J. Huston, K. Kondo, J. Morfin, L. Randall, and D. Soper, 45pp., Proceedings of the Workshop on Experiments, Detectors, and Experimental Areas for the Superconducting Supercollider, July 1987, Berkeley, CA. (World Scientific)
9. Report of the Large Solenoid Detector Group, V. E. Barnes with G. G. Hanson, et al., 46pp., Ibid.
10. Status and Prospects of the Fermilab Collider Detector, TeVatron Collider and Antiproton Source, Virgil E. Barnes for the CDF Collaboration, Proceedings of the Eighth Vanderbilt Conference - Collider Physics: Current Status and Future Prospects; October, 1987 (World Scientific).
11. The CDF Central and Endwall Hadron Calorimeter, V. E. Barnes, A. Di Virgilio, A. F. Garfinkel, S. E. Kuhlmann, and A. T. Laasanen, with authors from Frascati, Pisa and FNAL, Nucl. Inst. & Methods A267, 301-314 (1988).
12. The CDF Front End Electronics: The Rabbit System, S. E. Kuhlmann with authors from FNAL, Nucl. Inst. & Methods A267, 68-81 (1988).
13. Cosmic Ray Test of the CDF Central Calorimeters, A. Di Virgilio with authors from ANL, KEK, Tsukuba, Saga, U. of Penn., FNAL, and U. of Illinois, Nucl. Inst. & Methods A267, 330-350 (1988).
14. Design and Construction of the CDF Central Tracking Chamber, A. Byon with authors from FNAL, and Pisa, Nucl. Inst. & Methods A268, 50-74 (1988).
15. The CDF Vertex Time Projection Chamber System, M. H. Schub with authors from U. of Chicago, FNAL, and Tsukuba, Nucl. Inst. & Methods A268, 75-91 (1988).
16. Transverse Momentum Distributions of Charged Particle Produced in $p\bar{p}$ Interactions at $\sqrt{s} = 630$ and 1800 GeV, V. E. Barnes, A. Byon, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. Schub, et al., Phys. Rev. Lett. 61, 1819-1822 (1988).
17. The CDF Central and Endwall Hadron Calorimeter, V. E. Barnes, A. Di Virgilio, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, et al., Nucl. Instr. & Methods A267, 301-304 (1988).
18. Response Maps of the CDF Central Electromagnetic Calorimeter with Electrons, V. E. Barnes, K. Chadwick, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, J. Simmons, et al., Nucl. Instr. & Methods A267, 315- 329 (1988).

19. The CDF Detector: An Overview, V. E. Barnes, A. Byon, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. Schub, et al., Nucl. Instr. & Methods A271, 387-403 (1988).
20. Inclusive Central Jets at the Tevatron $P\bar{P}$ Collider, Proceedings of Seventh Topical Workshop on Proton Antiproton Colliders Physics, (A. F. Garfinkel), ed. R. Raja and J. Yoh (FNAL, Batavia) 1988.
21. Measurement of the Inclusive Jet Cross Section in $\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. Byon, S. Behrends, A. F. Garfinkel, S. Kuhlmann, A. T. Laasanen, M. Schub, et al., Phys. Rev. Lett. 62, 613- 616 (1989).
22. Measurement of W Boson Production in 1.8 TeV $p\bar{p}$ Collisions, V. E. Barnes, A. Byon, A. F. Garfinkel, A. T. Laasanen, M. H. Schub, et al., Phys. Rev. Lett. 62, 1005-1008 (1989).
23. Recent Results: Collider Detector at Fermilab, V. E. Barnes, Proceedings of the Quark Matter '88 conference, Nucl. Physics A498, 193C-204C (1989).
24. Limits on the Masses of Supersymmetric Particles from 1.8 TeV $p\bar{p}$ Collision, V. E. Barnes, A. Byon, S. Behrends, A. F. Garfinkel, A. T. Laasanen, M. H. Schub, et al., Phys. Rev. Letters. 62, 1825-1828 (1989).
25. Soft Hadron Production at CDF, V. E. Barnes, Proceedings of the XXIVth Rencontres de Moriond (1989).
26. Dijet Angular Distributions from $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. Byon, S. Behrends, A. F. Garfinkel, A. T. Laasanen, M. H. Schub, et al., Phys. Rev. Letters 62, 3020-3023 (1989).
27. Measurement of the Mass and Width of the Z^0 Boson at the Fermilab Tevatron, V. E. Barnes, A. Byon, S. Behrends, A. F. Garfinkel, A. T. Laasanen, M. H. Schub, et al., Phys. Rev. Letters 62, 720-723 (1989).
28. Search for Heavy Charged Particles in 1.8 TeV $P\bar{P}$ Collisions at the Fermilab Collider, (CDF Collaboration w/F. Abe, V. E. Barnes, A. Byon, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, et al.,) Phys. Rev. Letters 63, 1447-1450, (1989).
29. K^* production in $P\bar{P}$ interactions at $\sqrt{s} = 630$ and 1800 GeV, (CDF Collaboration w/F. Abe, V. E. Barnes, A. Byon, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, et al.,) Phys. Rev. D, Rapid Comm., 40, 3791-3794, (1989).
30. Search for the Top Quark in the Reaction $p\bar{p} \rightarrow$ Electron + Jets at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. Letters 64, 142-146 (1990).

31. Search for New Heavy Quarks in Electron-Muon Events at the Fermilab Tevatron Collider, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. Letters 64, 147-151 (1990).
32. Measurement of the Ratio $\sigma(W \rightarrow e\nu)/\sigma(Z \rightarrow ee)$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. Letters 64, 152-156 (1990).
33. Two-Jet Differential Cross Section in $P\bar{P}$ Collisions at $\sqrt{s} = 1.8$ TeV, (CDF Collaboration w/F. Abe, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, et al.,) Phys. Rev. Letters, 64, 157-160, January 1990.
34. Measurement of D^* Production in Jets from $P\bar{P}$ Collisions at $\sqrt{s} = 1.8$ TeV, (CDF Collaboration w/F. Abe, V. E. Barnes, A. Byon, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, et al.,) Phys. Rev. Letters 64, 348-352, January 1990.
35. Search for a Light Higgs Boson at the Fermilab Tevatron Proton-Antiproton Collider, (CDF Collaboration, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, Phys. Rev. D41, 1717-1721, March 1990.
36. Two-Jet Invariant-Mass Distribution at $\sqrt{s} = 1.8$ TeV, (CDF Collaboration, V. E. Barnes, S. Behrends, A. Byon, K. Chadwick, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, Phys. Rev. D41, 1722-1725, March 1990.
37. Pseudorapidity Distributions of Charged Particles Produced in $p\bar{p}$ Interactions at $\sqrt{s} = 630$ and 1800 GeV, V. E. Barnes, A. Byon, K. Chadwick, A. G. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, Phys. Rev. D, Rapid Comm. 41, 2330-2338, April 1990.
38. Jet-Fragmentation Properties in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, S. Behrends, A. Byon, K. Chadwick, A. F. Garfinkel, S. E. Kuhlmann, A. T. Laasanen, M. H. Schub, Phys. Rev. Letters 65, 968-971, August 1990.
39. A Determination of Electroweak Parameters from $Z^0 \rightarrow \mu^+ \mu^- (\gamma)$, L3 collaboration, A. Bujak, L. Gutay, T. McMahon, B. Stringfellow, Phys. Lett. B247, 473-480, September 13, 1990.
40. Calibration of a Scintillating Fiber Calorimeter Prototype Using a Moving Radioactive Source, V. E. Barnes and A. T. Laasanen, Proceedings of the SSC Detector Conference, Ft. Worth, Texas, October 15 - 17, 1990.
41. Measurement of the W-Boson Mass, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, Phys. Rev. Lett. 65, 2243-2246, October 1990.

42. Calibration of Scintillating Calorimeters Using a Moving Radioactive Source, V. E. Barnes and A. T. Laasanen, Proceedings of the International Conference on Calorimetry in High Energy Physics, Fermilab, October 29 - November 1, 1990.
43. Top-Quark Search in the Electron + Jets Channel in Proton-Antiproton Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, Phys. Rev. D43, 664-686, February 1991.
44. Measurement of the W-boson Mass in 1.8 TeV $p\bar{p}$ Collisions, (CDF Collaboration, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. D43, 2070-2093, April 1, 1991.
45. Measurement of the W-Boson PT Distribution in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, et al., Phys. Rev. Lett. 66, 2951-2955, June 1991.
46. Measurement of $\sigma B(W \rightarrow e\nu)$ and $\sigma B(Z^0 \rightarrow e^+e^-)$ in $p\bar{p}$ Collisions at $\sqrt{s}=1800$ GeV, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, et al., M. H. Schub, J. Tonnison, Phys. Rev. D44, 29-52, July 1, 1991.
47. Measurement of QCD Jet Broadening in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. D44, 601-616, August 1, 1991.
48. Measurement of the e^+e^- Invariant-Mass Distribution in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, V. E. Barnes, S. Behrends, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. Lett. 67, 2418-2422, October 1991.
49. Search for $W' \rightarrow e\nu$ and $W' \rightarrow \mu\nu$ in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. Lett. 67, 2609-2613, November 1991.
50. Measurement of $B\bar{B}$ Mixing at the Fermilab Tevatron Collider, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. Lett. 67, 3351-3355, December 1991.
51. Measurement of the W-Boson Mass in 1.8 TeV $p\bar{p}$ Collisions, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, J. Tonnison, et al., Phys. Rev. D43, 2070-2093, April 1991.
52. Determination of $\sin^2 \Theta_{11'}$ from the Forward-Backward Asymmetry in $p\bar{p} \rightarrow Z^0 X \rightarrow e^+e^- X$ Interactions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 67, 1502-1506, September 1991.
53. Calibration of Scintillating Tile Calorimeters using a Moving Radioactive Source, V. E. Barnes, A. T. Laasanen, and J. A. Ross, Proceedings of the Second International Conference on Calorimetry in High Energy Physics, Capri, Italy, 14-18 October, 1991.

54. Measurement of the Z-Boson p_T Distribution in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, M. Bailey, V. E. Barnes, A. Byon, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 67, 2937-2941, November 1991.
55. Inclusive Jet Cross Section in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, M. W. Bailey, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 1104-1108, February 1992.
56. Lepton Asymmetry in W-Boson Decays from $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, M. Bailey, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 1458-1462, March 1992.
57. Search for New Gauge Bosons in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, M. W. Bailey, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 1463-1467, March 1992.
58. A Lower Limit on the Top Quark Mass from Events with Two Leptons in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, M. W. Bailey, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanaen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 447, 1992.
59. Topology of Three-Jet Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, M. W. Baily, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. 45D, 1448, 1992.
60. Measurement of the Ratio $B(W \rightarrow \tau \nu)/B(W \rightarrow e \nu)$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 3398, 1992.
61. Measurement of the B-Meson and b-Quark Cross Sections at $\sqrt{s} = 1.8$ TeV Using the Exclusive Decay $B \rightarrow J/\psi K^\pm$, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 3403, 1992.
62. Progress in the Construction of the CDF Silicon Vertex Detector, M. Bailey, A. F. Garfinkel, N. M. Shaw, et al., To be published in Nuclear Instrument and Methods.
63. Properties of Events with Large Total Transverse Energy Produced in Proton- Antiproton Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, N. M. Shaw, J. Tonnison, et al., Phys. Rev. D45, 2249- 2263, 1992.
64. Measurement of the Isolated Prompt Photon Cross Section in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 68, 2734-2738, 1992.

65. Limit on the Top-Quark Mass from Proton-Antiproton Collisions at $\sqrt{s}=1.8$ TeV, M. W. Bailey, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, N. M. Shaw, J. Tonnison, et al., Phys. Rev. D45, 3921, June 1992.
66. Search for New Particles in CDF, A. Laasanen, Mod. Phys. Lett. A, Vol. 7, No. 29, 2659-2667, June 22, 1992.
67. Measurement of the Production and Muonic Decay Rate of W and Z Bosons in $p\bar{p}$ Collisions at $\sqrt{s}=1.8$ TeV, M. W. Bailey, V. E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett., Vol. 69, No. 1, 28-32, July 6, 1992.
68. Limit on the Rare Decay $W^{\pm} \rightarrow \gamma \pi^{\pm}$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, M. W. Bailey, V.E. Barnes, A. F. Garfinkel, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 69, 2160, 1992.
69. Dijet Angular Distribution in $p\bar{p}$ - Collisions at $\sqrt{s} = 1.8$ TeV, V. Barnes, B. T. Huffman, A. T. Laasanen, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 69, 2896, November 16, 1992.
70. Inclusive $J/\psi, \psi(2S)$, and b-Quark Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV, M. W. Bailey, V. E. Barnes, A. Garfinkel, B. T. Huffman, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. Lett. 69, 3704, December 28, 1992.
71. Limits on the Production of Massive Stable Charged Particles, M. W. Bailey, V. E. Barnes, A. Garfinkel, B. T. Huffman, M. H. Schub, N. M. Shaw, J. Tonnison, et al., Phys. Rev. D46, 1889, September 1, 1992.
72. A Novel Electromagnetic Shower Maximum Position Detector, V. E. Barnes, et al., Nucl. Inst. & Methods A324, 475, January 15, 1993.

5 Ph.D. Theses

1. A Study of π^+p , K^+p , and pp Interactions at 100 GeV/c Using a Hybrid Bubble Chamber-Spark Chamber System, W.M. Morse (1975).
2. Measurement of π^-p Backward Elastic Scattering Between 3.0 GeV/c and 5.5 GeV/c, R.L. Dixon (1975).
3. Study of Single-Pion Production by the Weak Charged-Current, G.M. Radecky (1980).
4. Particle Production in Pi-Minus Proton Interactions at 50 GeV/c, L.J. Dauwe (1980).
5. Study of Charmonium Chi States Produced in Proton and Negative Pion Interactions with a Beryllium Target, J. R. Wilson (1985)

6. (χ -states) Measurements of $\chi(3510)$, $\chi(3555)$ and Neutral Strange Particles in 190 GeV/c π^- Beryllium Interactions, James Simmons (1988).
7. Central Jet Transverse Momentum Distribution and Test of QCD for 1.8 TeV $p\bar{p}$ Collisions, Stephen Kuhlman (CDF), (1988).
8. Central Production of Charged Particles at CDF, AeSook Byon (CDF) (1989).
9. K^* Production in Minimum Bias Events at the FNAL TeVatron Collider, Martin Schub (CDF) (1989).
10. Measuring the Inclusive Central Muon Cross Section in Proton-Antiproton Collisions at 1.8 TeV, B. Todd Huffman (CDF) (1992)

Current Thesis Students:

- Mark Bailey
- Mark Kruse
- Jim Tonnison

Fig. 3-1 Invariant mass distribution ψK^- combinations.

Fig. 3-2 $d\sigma(p\bar{p} \rightarrow B^- X)/dp_T$ distribution. Charged B decays are reconstructed in the decay $B^- \rightarrow \psi K^-$ and $\psi \rightarrow \mu^+ \mu^-$

Fig. 3-3 Invariant mass distribution of the ψK^* combinations

Fig. 3-4 $\sigma(p\bar{p} \rightarrow bX)$ distribution.

AN EXPERIMENTAL HIGH ENERGY PHYSICS PROGRAM

within the

DEPARTMENT OF PHYSICS
PURDUE UNIVERSITY
West Lafayette, Indiana 47907

Supported by
U.S. DEPARTMENT OF ENERGY

Through

Division of Sponsored Programs
Purdue Research Foundation
West Lafayette, Indiana

PLANS FOR THE
CENTRAL HIGH ENERGY COMPUTER FACILITY

for

11/1/93 - 10/31/94

FUNDS REQUESTED: \$92,900

Current Grant Number: DE-FG02-91ER40682 Task S

Contact Persons: L. J. Gutay, D. H. Miller,
(317) 494-5390

April 1993

TASK S
11/1/93 - 10/31/94
Computer

We are requesting funds for continuing upgrades to our central computer facility.

(3) DEC Alpha workstations or equivalent with 80 megabyte of memory and each with 2 gigabyte disk	48,000
(3) Two gigabyte disks for distributed workstation and for data storage on Silicon Graphics	\$7,500
(3) 4mm tape drives	\$4,800
(1) High volume laser printer	\$3,500
(2) Bridges to segment the ethernet loop	\$5,600
(2) Unix workstations Silicon Graphics Indigo or equivalent. Includes trade in of SG4 personal Iris.	<u>\$23,500</u>
	\$92,900

END

DATE
FILMED

7 / 14 / 93