THE EFFECTS OF SOIL MINERAL PHASES
ON THE ABIOTIC DEGRADATION
OF
SELECTED ORGANIC COMPOUNDS

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Annual Progress Report

Dr. Shingara S. Sandu
Claflin College
Orangeburg, SC 29115

June 1992

PREPARED FOR THE
U. S. DEPARTMENT OF ENERGY
UNDER GRANT
DE-FG09-90SR18159

RECEIVED
JUN 21 1993
OSTI

MASTER *ds*

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Tetraphenylboron (TPB) will be used to precipitate radioactive ^{137}Cs from high level nuclear waste water, at the Defense Waste Processing Facility (DWPF) which is operated by the U.S. Department of Energy (US DOE), at the Savannah River Site (SRS), Aiken, SC. The decontaminated wastewater will contain millimolar quantities of TPB that will be processed into salt concretions. The waste processing operations will require about 3×10^5 kg/yr of TPB during its full scale operation. The transportation and use of large amounts of TPB can potentially result in the release of TPB into soil or aquatic environments where its degradation will take place.

Previous studies have shown that TPB degrades abiotically in soils; however, the role of specific mineral surfaces in mediating the reaction kinetics and mechanisms for the abiotic degradation of TPB are not clearly known. Laboratory studies have been undertaken to evaluate the surface facilitated degradation of TPB by aluminum silicate clay minerals-kaolinite and montmorillonite. Preliminary results indicate that the rate of degradation of TPB is much higher in kaolinite-a single layer mineral-than in montmorillonite - a double layer mineral- and Orangeburg loamy soil. The initial products of TPB degradation in both minerals are diphenylboric acid (DPBA) and biphenyl. However, HPLC monitoring of degradation products of TPB in montmorillonite appears to indicate the presence of phenol and monophenyboric acid (MPBA).

Tetraphenylborate (TPB) is used to precipitate radioactive ^{137}Cs from high-level nuclear waste water at the Defense Waste Processing Facility (DWPF) operated by the U.S. Department of Energy (DOE) at the Savannah River Site (SRS), Aiken, SC (1). This process is a part of the procedure for glassification of high-level nuclear wastes in preparation of long-term geological disposal. The decontaminated wastewater contains millimolar quantities of TPB that will be processed into salt concretions. The transportation and use of large amounts of TPB can potentially result in the release of TPB into soil or aquatic environments. Although TPB has been used in a variety of laboratory applications (2, 3, 4), it has not been previously used in industrial scale quantities. Studies (5) have shown that TPB degrades in soils to initially form diphenylboric acid (DPBA) and biphenyl. The degradation of TPB appears to be an

abiotic process as its degradation was observed in sterilized soils collected from several locations in South Carolina (6). DPBA is not the final product of TPB abiotic-degradation as it does not accumulate in the soils, but rather appears to degrade further into other organoborate compounds which subsequently degrade into inorganic boron (6).

The reaction mechanism and the specific soil surface mediating the chemical degradation of TPB in soils are not known. Soils present a heterogeneous surface with a variety of organic and inorganic constituents. The preliminary studies indicate that the degradation of TPB is abiotic and is facilitated by soil mineral surfaces, although the specific mechanisms are not understood. TPB as well as its intermediate product DPBA have been reported to be toxic to microorganisms (7) and plants, dependent on soil or water environments for their survival and growth (6, 8, 9).

Objective

The contribution from various surface sites (Si, Al, Fe, Mn, organic matter) needs to be evaluated in order to be able to predict the rate of transformation in the surface and lower horizons in various soil types. In addition, the intermediate products derived from decomposition of TPB need to be identified and their persistence evaluated. The initial work conducted at SREL (1) provided the basis for design of further investigations on the abiotic degradation of TPB and DPBA. The major objectives of the proposed research are to determine the effects of soil mineral phases on the rates of surface facilitated degradation of tetraphenylborate and diphenylboric acid. In addition, the experiments are currently being conducted to determine the products of the abiotic degradation of diphenylboric acid in soils. Previous work (6) has suggested that the initial reaction involves oxidation of TPB by electron transfer to Lewis acid sites, the present study focuses on the effects of the major redox active solid phases.

MATERIALS AND METHODS

Adsorbate clay minerals, KGa-2 (poorly crystallized kaolite, Warren County Ga) and SWY1-(Na-montmorillonite Crook County Wyoming) were procured, from the Department of Geology, University of Missouri, Columbia, Missouri. Manganite and lepidocrocite were purchased from Ward's Natural Science Establishment, Inc., Rochester, New York. Tetraphenylboron (TPB) diphenylboric acid (DPBA) biphenyl (BP), phenol, and monophenylboric acid (MPBA), were obtained from Aldrich Chemical Company, Inc. Milwaukee, WI.. All chemicals were of AR quality. The surface area, cation exchange capacity and pH of the minerals were determined by use of the recommended methodology (10). The clay minerals were cleaned of carbonates, soluble salts, organic matter, free iron oxides and saturated with Ca and Na (10) prior to their use as substrate in this study.

Half a gram of kaolinite or montmorillonite was transferred to 15 mL. pyrex glass centrifuge tube. One hundred μ L, containing required amounts of TPB was added to 0.5g of clay minerals. The contents of tube were thoroughly mixed. The centrifuge tubes containing clay, treated with TPB, were arranged in a test tube rack and were allowed to incubate in darkness at room temperature ($25^{\circ}\text{C} \pm 2$). The tubes were removed at pre-determined time, and extracted with 4 mL of acetonitrile by shaking manually for fifteen minutes. The extraction mixture was centrifuged for 30 minutes at 10,000 rpm in Fisher Centrifuge TM Centrifuge. The supernatant liquid was passed through a 0.2 μ m nylon syringe filter and analyzed, using high performance liquid chromatography (HPLC). The HPLC analysis was performed using Hewlett -Packard model 1050 liquid chromatograph, with a UV100 variable wavelength detector, autosampler, LC chem station and a VYDAC-C₁₈ reverse phase column. The solvent system employed was water (HPLC grade), adjusted to pH 3.2 with H₃PO₄ and acetonitrile. The elution program was a 20 min. gradient from 95:10 water/acetonitrile to 100% acetonitrile at a flow rate of 1 mL min⁻¹. The UV detector sensitivity was generally 0.02 to 0.04 absorbance units for a full-scale deflection, monitored at 230 nm with an injection volume of 20 μ L. The TPB added to one set of clay minerals was extracted within the first about three minutes which has been recognized as time zero extraction.

RESULTS AND DISCUSSION

The TPB concentrations decreased exponentially with time in kaolinite and montmorillonite clays. About 20 percent of the initial TPB remained in clays at the end of the experimental period. The degradation rate of TPB on kaolinite was much higher than montmorillonite. Over 80% of TPB added to kaolinite disappeared in 8 hours where as it took over 12 days for montmorillonite to achieve that level of TPB degradation. It has been suggested that the initial reaction leading to the degradation of TPB, involves its oxidation at Lewis acid sites. Geske (1959) investigated the electrochemical oxidation of TPB using Pt electrodes and found that the initial products of electrochemical reaction were DPBA and biphenyl. Mills et al (11) suggested the following step wise degradation of TPB.

The initial degradation products (DPBA & Biphenyl) of TPB, identified in this study are consistent with the previous reports. However; HPLC monitoring of TPB degradation in montmorillonite, appears to indicate the presence of phenol and monophenylboric acid (MPBA).

Kaolinite is composed of one layer of silica tetrahedra and a sheet of aluminum octahedra where an aluminum atom is linked to six oxygen atoms (coordination #6) and or hydroxal groups through coordinate covalent bonds. Due to the presence of one silica sheet and one alumina sheet, kaolinite is known as 1:1 mineral. Its structure suggests that metal ions are likely to be exposed to perform the function of Lewis acid. Montmorillonite is composed of two sheets of silica tetrahedra and one sheet of aluminum octahedra which is sandwiched between two sheets of silica tetrahedra thus limiting or inhibiting the role of metal ions as a Lewis acid.

Kaolinite, due to its exposed metal ions which have been hypothesized (11) as potential sites for the redox decomposition of TPB provides an ideal substrate for the surface catalyzed degradation of TPB. Contrary to this montmorillonite does not provide ideal surface conditions for the acceptance of electrons that need to be donated by TPB during its redox reaction. Consequently the degradation of TPB in montmorillonite is slow. This is initial data. The laboratory studies on the fate of TPB degradation in soils is an on going project and more data will become available soon

1. Mills, G. L., D. Schwind and D. C. Adriano, Preconcentration and analysis of tetraphenylboron and diphenylborinic acid in natural waters using C₁₈ reverse-phase liquid chromatography. *Chemosphere*, 17,937-942 (1988).
2. Flaschka H. and A. J. Bernard, Jr. In Reilly ed., *Advances in Analytical Chemistry and Instrumentation*. Interscience, New York, pp. 1-118 (1960).
3. Jalicoeur, C., N. D. and A. Cabarra, *Can. J. Chem.*, 49: 2008-2013 (1971).
4. Amatt, S and S.C. Abrahams, *Acta Cryst.* 11, 449-450 (1958).
5. Adriano, D. C. G. L. Mills, J. L. Afre and R. J. Hart, *Proceedings of the International Conference on Environmental Contamination, United Nations Environment* (1985).
6. Mills, G. L., D. Kaplan, D. Schwind, and D. Adriano, Surface-facilitated chemical degradation of tetraphenylboron in soil. *J. Environ. Qual.* In Press. (1989).
7. Shann, J. B., P. M. Bertsch and S. M. Presnell. *J. Environ. Qual.* (submitted) (1989). (in press) 1990.
8. Burkman, W. G., D. C. Adriano and J. L. Afre. Phytoavailability of boron from organic sources. *J. Environ. Qual.*, 15: 416-421 (1987).
9. Kaplan, D. I., W. G. Burkman and D. C. Adriano. Response of loblolly pine (*Pinus taeda* L.) seedlings to soil-applied organo-borates. 37:73-83 (1988).

10. Klute Arnold. Methods of Soil Analysis Part I Physical and Menorological methods. American Society of Agronomy, Inc., Medison, WI. 53711 (1965).
11. Mills, Gary, Kaplin, D., Schwind, Dona and Adriano Dony, Surface-Facillated Chemical Degradation of Tetraphenalboron in Soil, J. Environ Quality: 19, 135, (1990).
12. Geske, D. H., J. Phys. Chem 63:1062, 1959.

END

**DATE
FILMED**

7 / 9 / 93