Fermi National Accelerator Laboratory

FERMILAB-TM-1916

Coupling in the Tevatron

Norman M. Gelfand

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

December 1994

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Coupling in the Tevatron

Norman M. Gelfand *

Fermi National Accelerator Laboratory

December 20, 1994

Introduction

The performance of the Fermilab Tevatron Collider at the commencement of run Ib was far below expectations. After a frustrating period of several months, a low- β quad downstream ¹ of the interaction point at B0 was found to be rolled. This rolled quadrupole coupled the horizontal and vertical motion of the Tevatron beams. It also made matching the beam from the Main Ring to the Tevatron impossible, resulting in emittance blow up on injection. The net result of the roll was a significant reduction in the Tevatron luminosity. When the roll in the quadrupole was corrected the performance of the Tevatron improved dramatically. This note will discuss the experimental data indicating the presence of coupling and subsequent calculations which show how coupling can affect the luminosity. It is not intended to exhaust a discussion of coupling, which hopefully will be understood well enough to be discussed in a subsequent note.

*This work performed at the Fermi National Accelerator Laboratory, under contract DE-AC02-76CHO3000 with the U.S. Department of Energy

¹Downstream and upstream in this note shall refer to the motion of the protons. Thus B11 is downstream of B0.

Coupling

Introduction

Two sets of measurements made with the Tevatron, the measurements of the change in the closed orbit when the strength of a dipole corrector is changed, and the variation in tune as the strength of a quadrupole is changed, both led to the conclusion that the Tevatron was coupled.

In the first case the evidence was the large change in the closed orbit in the crossed plane, the plane orthogonal to the plane of the dipole whose strength was varied. This is shown in figure 1 where the kick was generated using the correction dipole HE13. The rms size of the slope in the crossed plane is $\approx 40\%$ of the rms slope in the kick plane.

In the second case the evidence was anomalously small values of tune shift measured at locations outside of the interaction region. The small tune shifts imply values of β outside of the interaction region comparable to those expected at the interaction point (figure 2). This is clearly inconsistent with any of the usual models of the Tevatron. The values of β calculated from the measurements could not be understood as due to differences, from their nominal values, in the strengths of the lattice elements in the low- β insertion.

It is possible to understand both of these observations if we introduce coupling into the model we have of the Tevatron lattice.

β Measurements

Introduction

The primary source of information about the lattice in this run comes from a series of measurements, using a method pioneered here at Fermilab by Glenn Goderre. The fundamental idea is to apply to the beam a single, known dipole kick, either horizontal or vertical, at a specific location, and measure the resulting change in the closed orbit, at the locations of all the BPMs. To increase the precision of the resulting data, the measurements, of the changed closed orbit, are repeated with a number of different values for the dipole kicks and the slope of the closed orbit value with respect to the size of the dipole kick is calculated.

The correction dipoles located at HE11, HE13, VE11 and VE14 are used

to produce the kicks and the closed orbits are measured in *both* planes for each dipole kick. These pairs of correctors are chosen because the phase advance between the locations is large ($\Delta\phi_x \approx 69^\circ$) and ($\Delta\phi_y \approx 95^\circ$) and yet is not near a multiple of 180° .

When the measurements are made, using the console program C91, the results appear in files ² containing the slope of the variation in the closed orbit positions at the BPM as the dipole kick is varied.

The procedure just described was developed in order to measure the β function in the Tevatron. The change in the closed orbit at point k , Δu_k due to a dipole kick ($\delta u'_{ui}$) at point i , in the case of an *uncoupled* machine, can be written as:

$$\Delta u_k / \delta u'_{ui} = \sqrt{\beta_{ui} \cdot \beta_{uk}} \cdot \cos(\Psi_{ui} - \Psi_{uk} - \mu_u/2) / (2 \cdot \sin(\mu_u/2)) \quad (1)$$

where:

β_{ui} is the value of β_u at the point i ;

Ψ_{ui} is the phase in the u plane at point i ; and

μ_u is the tune in the u plane.

In the crossed plane, the plane orthogonal to u , there would not be any change in the closed orbit when a kick is applied in the u plane.

As was seen in Figure 1, that is not the case with the data. The slopes in the horizontal plane are comparable to those in the vertical plane when the vertical corrector is varied. Thus the Tevatron is coupled and the coupling is strong.

In a coupled machine the 4×4 transfer matrix, which transforms the initial phase space coordinates of a particle (x, x', y, y'); around the lattice back to the starting point in the lattice, is no longer just two 2×2 matrices along the diagonal, with zero for the off diagonal 2×2 matrices. As a consequence of this the usual representation of the motion of a particle in the lattice in terms of the horizontal and vertical β functions, viz. β_H and β_V , is not appropriate.

Despite the fact that calculating, and using, the normal β functions is not a correct way of describing the behavior of the beam, it can be helpful in trying to understand what is happening.

²I want to thank W. Marsh for providing a program which made it possible to access the data in the C91 files.

Consider a horizontal kick $\Delta x'$ at a location a , e.g. E13, and a lattice element with a skew quadrupole field at a location b . The amplitude of the horizontal closed orbit at b , x_b , given by equation (1) is:

$$x_b = \Delta x'_i \cdot \sqrt{\beta_{xa} \cdot \beta_{xb}} \cdot \cos(\Psi_{xa} - \Psi_{xb} - \mu_x/2)/(2 \cdot \sin(\mu_x/2)) \quad (2)$$

A finite horizontal amplitude at the position of the coupling, b , results in a vertical kick, $\Delta y'$, equal to $x_b \cdot r_b$ where: r_b is the skew quadrupole strength of the coupling element.

This vertical kick will modify the vertical closed orbit whose amplitude, at a point c is again given by equation (1):

$$y_c = r_b \cdot x_b \cdot \sqrt{\beta_{yc} \cdot \beta_{yb}} \cdot \cos(\Psi_{yc} - \Psi_{yb} - \mu_y/2)/(2 \cdot \sin(\mu_y/2)) \quad (3)$$

Combining equation (2) and equation (3) we see that the amplitude in the crossed plane is proportional to:

$$y_c \propto r_b \cdot \sqrt{\beta_{xb} \cdot \beta_{yb}}.$$

The effect in the crossed plane, due to a coupling element of a given magnitude, will be greatest if the element is located at a place in the lattice where the product

$$\sqrt{\beta_x \cdot \beta_y}$$

is large.³

We can compute the lattice functions for the design lattice of the Tevatron and compute $\sqrt{\beta_x \cdot \beta_y}$ for the different points in the lattice. The results, for a section of the lattice, are plotted in figure 3. It is clear that the largest values for $\sqrt{\beta_x \cdot \beta_y}$ occur in the low- β triplet, and in particular in the outside elements of the triplet, the Q2 magnets. It is therefore natural to look for the source of the coupling, observed in the cross plane orbit data taken in the Tevatron, in the quadrupoles of the low- β triplets.

³We see also the value, if not the necessity, of using more than one corrector in each plane. For a given choice of corrector the argument of the cosine function in equation (3) might be close to an odd multiple of $\pi/2$. In this case we would not see any effect in the cross plane. By choosing two correctors, with a large phase advance between them, we are able to see the effect of the coupling in at least one set of data.

These closed orbit data unfortunately do not allow the identification of which element is producing the coupling, or even on which side of the interaction region (IR) the coupling occurs. This can be seen by looking once again at equations (2) and (3) and noticing that the phase advance across the elements of the triplet are small (due to the large value of β in the triplet elements compared to their lengths) and that the phase advance across the low- β insertion is very close to 180° .

We can get the same vertical kick, $\Delta y'$, from different elements in the triplet, by adjusting the, unknown, strength of the coupling element to compensate for the different values of β in the different elements of the triplet. The value y_c will not change because the phase advance $\Psi_{yc} - \Psi_{yb}$ will not change significantly as I change the element in the triplet causing the coupling. Thus the triplet element causing the coupling can not be identified.

Across the IR the phase advance, in both planes, is $\approx 180^\circ$. If the size of the closed orbit change due to the kick at a is δx on the upstream side of the IR it will be $-\delta x \cdot \sqrt{\beta_d/\beta_u}$ at the corresponding point on the downstream side. Adjusting the value of r_b the magnitude of vertical closed orbit at the point c will be the same, regardless of which side of the IR we have the coupling, because the phase advance in the y plane from point b to point c has also changed by 180° .

Measurements of the change in the closed orbit inside the IR can eliminate the ambiguities.

Analysis of the Data

The normal model of the Tevatron, whether in SYNCH, Tevlat, Tevconfig, MAD ..., contains no skew elements ⁴ except for the skew quadrupole and skew sextupole elements located in the Tevatron spool packages, and the high order skew moments in the Tevatron magnets. The high order skew elements are not strong enough to account for the magnitude of the observed closed orbits in the crossed planes. The values in the skew quadrupoles are adjusted to bring the tunes in the two planes together, i.e. to reduce the coupling, and therefore should not be the source of the coupling.

In order to study the source of the coupling we have to introduce into the description of the Tevatron lattice skew elements, whose strengths can

⁴Any of these models can, of course, be modified to include effects, such as rolls, that could produce coupling.

be varied. Zero length skew quadrupoles were inserted into the description of the lattice at the location of each of the quadrupoles in the lattice. These *pseudo* elements do not exist in the real machine but are used to reproduce effects, e.g. the change in the closed orbit in the cross plane, that are seen during Tevatron operation.

This modified lattice has been used with a version of the tracking code Tevlat ⁵ which allows fitting experimental data with MINUIT, varying the parameters of lattice elements in the course of the fit.

We begin with the closed orbit data, both for the kick plane and in the crossed plane, in C91 files # 18,19,20 and 21 taken on 3/15/94. The slopes of the original data have been converted into a closed orbit by assuming a kick of $30\mu r$ at the corrector. The results of fitting all four closed orbits, simultaneously with a single set of parameters are shown in figures 4a,4b,4c,4d. ⁶ Reasonable, though not totally satisfactory, solutions are found assuming skew quadrupole fields at the location of Q2 quadrupoles down stream of the IR at both B0 and D0.

It is possible, but not really meaningful, to convert the fitted values of these skew moments into a roll of the corresponding quadrupole. There is no reason to believe that the quad selected for the skew moment is the actual rolled quad. Nonetheless if we use the fitted skew moments and assume that the computed value is due to a roll in the appropriate quads in the B0 and D0 interaction region, we find that the roll would be of the order of a few mr .

Subsequent to the analysis described above, a survey was performed of the accessible low- β magnets at B0. A roll of $\approx 7mr$ was found in the downstream Q2. With the culprit identified it made sense to redo the fitting with the skew moment represented by a roll in the magnet. In this refitting the strengths of the skew correctors at A0(T:SQA0), B0(T:SQB0), and D0(T:SQD0) were fixed at values given by the currents in the spool elements used when the data were taken. The strength of the skew corrector circuit SQ was left as a parameter in the fitting, as were the strengths of the tuning quadrupoles and the rolls of the downstream Q2 quadrupoles at B0 and D0. The strengths of

⁵The original version of Tevlat was written by A. Russell

⁶The data in figures 4a,5a,6a and 7a are from the kick at HE11; in figures 4b,5b,6b and 7b are from the kick at HE13; in figures 4c,5c,6c and 7c are from the kick at VE13; in figures 4d,5d,6d and 7d are from the kick at VE14. The figures labeled as differences are the differences between the fitted closed orbit and the measured closed orbit.

the low- β quads were not varied and were set at the values given by C49.

The results of the fitting are shown in figure 5a,5b,5c, and 5d. The quality of the fits is similar to the fits done with the pseudo-quads. The fit gives a value of $+7.0mr$ for the roll in the downstream Q2 at B0 and $-1.4mr$ for the downstream Q2 at D0. I would estimate that the uncertainty in these values is $\approx 1mr$. The agreement with the survey result is striking, but should not be overemphasized. We have represented all the skew effects by the roll in a single magnet which is certainly not the case. We get the correct value in this instance because of the large roll in a single magnet. When the skew effects are distributed among several magnets we cannot expect to find the individual rolls using this technique for the reasons given earlier.

The roll of the downstream Q2 at B0 was corrected and the β measurements were repeated, again using C91, and the data was stored in files #201, 202, 203, 204 (the data were taken on 7/22/94). The crossed plane amplitudes have been reduced by approximately a factor of 2 from the earlier measurements. Even after correcting the roll in the upstream Q2 at B0 there is still a considerable amplitude in the crossed plane. Not all the coupling has been removed by unrolling the quad upstream Q2 at B0.

The data have been fit using the same procedures used to fit the earlier data. The results are shown in fig #6a, 6b, 6c and 6d. ⁷ The results show a reduction in the roll of the upstream Q2 at B0 to $\approx 0.3mr$. The fit now indicates a roll of the downstream Q2 at D0 of $\approx -1.5mr$ compared to the $\approx -1.4mr$ found earlier. It is reassuring that the same value for the roll at D0 is found here as in the earlier solution since no change was made to the magnets at D0.

During the long shutdown in Sept. 1994, the accessible triplet quads at B0 and D0 were surveyed and any measured rolls were removed. The β measurements were repeated. The results contained in files #250,251,252, and 253, (the data were taken on 9/30/94) show a further reduction in the slope of the closed orbit in the crossed plane. The data have been fit as before and here again the rolls of the upstream quad at D0 and at B0 were allowed to vary. The results of the fitting are shown in figure # 7a ,7b ,7c, and 7d. The fitted values for the roll in the Q2 at B0($\approx 0.5mr$) and D0($\approx 0.2mr$) are now very small and are consistent with zero.

Though the coupling from the interaction regions is considerably de-

⁷Please note that the vertical scale has been changed.

creased there is still a fair amount of coupling left in the Tevatron. Comparing the recent crossed plane data to the original data we find that the crossed plane amplitudes have been reduced by approximately a factor of 2. The fitting described above assumed that the coupling was in the low- β regions and so does not give a particularly good fit to the crossed plane closed orbit data because the coupling is not where I have assumed it to be. Further study will be needed to identify the source of this coupling.

A summary of the results is contained in table I.

Table I
Summary of Results from Fitting Closed Orbit Data

	Rolled B2	Unroll B2	Unroll all
C91 Files used in fit	18,19, 20,21	201,202, 203,204	251,252, 253,254
Variable	Fitted value	Fitted value	Fitted value
Tune quads ckt 1 (/m)	6.1E-04	6.5E-04	6.5E-04
Tune quads ckt 2 (/m)	-6.5E-04	-6.8E-04	-6.9E-04
Skew quad ckt SQ (/m)	-1.5E-05	-7.5E-05	-7.5E-05
Skew quad ckt SQA0 (/m)	-1.8E-03	-8.31E-04	-5.1E-04
Skew quad ckt SQB0 (/m)	-8.43E-04	1.75E-04	0
Skew quad ckt SQD0 (/m)	-1.6E-04	3.5E-04	0
Rotation, DS Q2 at B0 (mr)	7.0	0.29	0.49
Rotation, DS Q2 at D0 (mr)	-1.4	-1.5	0.20

Tune Shift

Introduction

The traditional method of measuring the value of β at a location in an accelerator is to vary the gradient of a quadrupole at that location and measure the resulting tune shift. In the *absence* of coupling the tune shift is related to the value of β , viz.

$$\delta\nu = (1/4\pi) \cdot \frac{\beta \cdot l \cdot \Delta B'}{[B\rho]} \quad (4)$$

where:

β is the value of the β function where the quadrupole is varied;

l is the length of the quadrupole;

$\Delta B'$ is the change of the gradient in the quadrupole;

$[B\rho]$ is the magnetic rigidity or momentum.

The data that I will analyze were taken on 4/24/94, before the unroll of the downstream Q2 at B0. The current in the different correction quadrupoles at B0 was varied and the tunes measured. Equation 4 was used to compute values of β from the data. The results were seen in figure 2.

In a coupled machine the simple relationship of equation 4 is no longer true. It is however possible to fit the measured tune shift as a function of quadrupole strength ($\Delta B'$ in the above equation) using a model of the Tevatron and MINUIT. The parameters used were the same as the ones used to fit the closed orbit data, viz. the strengths of the tuning quadrupoles, the strength of the skew quadrupole circuit T:SQ, and the rolls in the downstream quad at B0. The value of the roll in the downstream quad at D0 was fixed at $-1.4mr$ from our fit to the closed orbit data. This was done because we have no tune shift data taken by varying the quads near the D0 interaction region.

For the original case, the only case for which we have data, the solution for the roll at B0($+6.5mr$) agrees very well with the values found from fitting the closed orbit data. It is reassuring that two independent methods find the same roll for the quadrupole at B0. That they agree with the survey is even more astonishing.

Summary of Coupling Analysis

Measurements made with closed orbits and tune shifts both indicated that the Tevatron was coupled. A computed model was constructed in which it was assumed that the coupling was due to a roll in the downstream Q2 magnets at B0 and D0. The model was used with MINUIT to find the magnitudes of the rolls. Where both sets of measurements exist they give consistent values for the rolls and which furthermore agree with the survey data taken after the measurements were made.

Analysis of the data taken after surveys of the low- β quadrupoles, and correction of the observed rolls, indicate, as expected, sharply reduced rolls, and hence of the coupling. There is still significant coupling in the Tevatron.

Luminosity

Introduction

The objective for the Tevatron is to produce high luminosity at the interaction regions at B0 and D0. The luminosity is inversely related to the size of the p and $\bar{p}$ beams, when they collide. In an uncoupled accelerator (with no non-linear elements) the size of a beam at a given point in the lattice can be expressed in terms of the emittance (ϵ), which is invariant as one moves from point to point in the lattice, and the function $\beta(z)$, where the coordinate z identifies where in the lattice we are measuring β . The size, σ , of the beam is given, when either the dispersion η or the momentum spread dp/p is zero, by

$$\sigma_{rms} = \sqrt{\beta \cdot \epsilon_{rms}}.$$

The luminosity, $\mathcal{L}$ is then

$$\mathcal{L} \propto \frac{1}{\sqrt{\beta_x \cdot \epsilon_x(rms) \cdot \beta_y \cdot \epsilon_y(rms)}}$$

In the case where we have a coupled machine it is not possible to use the normal horizontal and vertical β functions to calculate the beam size. In order to see the effect of the coupling on the beam size, and hence on the luminosity Tevlat was used to track a distribution of 1000 particles for a thousand turns. The initial particle distribution was generated assuming the design lattice with no coupling. The values σ_x and σ_y for the distribution were chosen so that the particles had a 95% normalized emittance, $\epsilon_n(95\%) = 25\pi mm m r$.

In the tracking for each turn the value of σ for the particle distribution was calculated at D0 and B0 for both planes. The σ s vary from turn to turn since the original distribution was not a stationary distribution. The values of σ were therefore averaged over 100 turns. The resulting average values for σ , which go into the calculation of $\mathcal{L}$ are very constant over the several thousand turns for which we have tracked.

Calculations and Results

The computed values of $\sqrt{\sigma_x \cdot \sigma_y}$ at B0 and D0 are plotted in figure 8 for four configurations of the Tevatron:

- The design lattice with $\beta^* \approx 0.35m$.
- The solution found for data taken with the downstream Q2 at B0 rolled.
- The solution found for data taken after the roll of the downstream Q2 at B0 was corrected.
- The solution found for data taken after the shut down in 9/94.

The calculated radius of the beam, $\propto (\sigma_x \cdot \sigma_y)^{1/2}$, at B0 after the roll of the Q2 was removed, $\approx 46\mu$ (see figure 8) is $\approx 74\%$ of the calculated radius using the fit to data where the Q2 was rolled. This decrease in radius should translate into an $\approx 80\%$ increase in the luminosity.⁸ There is a much smaller change in the beam radius at D0 with the changes in the quad rolls, the unrolled value is $\approx 93\%$ of the value with the rolled Q2.

Figure 9 shows the equivalent beam size at the interaction point computed from the *measured* luminosity and beam intensities for a number of stores before and after the unroll of the downstream Q2 at B0.

The equivalent beam radius is

$$(\sigma_x \cdot \sigma_y)^{1/2} = \frac{N_p \cdot N_{\bar{p}}}{\mathcal{L}}$$

where:

$\mathcal{L}$ is the measured luminosity at the start of the store;

N_p is the measured proton intensity at the start of the store;

and $N_{\bar{p}}$ is the measured pbar intensity at the start of the store.

The effect of the unroll of the Q2 is clearly seen. The size of the beam at the interaction region at B0 clearly decreased when the quad was unrolled. Because of the variation in the beam emittances from one store to the next a simple comparison between the calculation described above and the measurements is not possible. We can however average the size of the beam calculated from the luminosity and the bunch intensities from stores before and after the correction of the roll. When we compare the averages we find that the beam size at the interaction region at B0 decreased by $\approx 63 \pm 13\%$. The beam size at D0 did not change significantly. This is in

⁸The luminosity is $\propto 1/r^2$. I have not calculated the size of beam at other than the nominal interaction point and therefore have not calculated the luminosity by integrating over the bunch length

good agreement with the change predicted by the calculation in Tevatron of the beam size.

Conclusion

The data taken in the β measurements can be used to study coupling in the Tevatron. From the measurements it seems possible to construct a reasonable model of the Tevatron which, together with tracking can be used to calculate the Tevatron luminosity.

Acknowledgments

The data analyzed here were obtained by a number of members of the Accelerator Division as part of an extensive series of measurements performed to better understand and improve the Tevatron. Among those responsible for the data used in this note are G. Annala, S. Assadi, G. Goderre, G. Jackson and W. Marsh. I wish to express my gratitude to them first for taking the data with such care, and then for making it available in a form convenient for analysis.

Slopes— Kick at he13 Data from file—019

Input file—/afs/fnal/files/home/room1/gelfand/tracking/minult/data/slopeplot.plotf

11Nov-1994 9-06-22 Plot number— 1 Plot/symbol

11Nov-1994
9-06-22
Plot number— 1
Plot/symbol
ave = -1.876
σ = 39.75

Slopes— Kick at he13 Data from file—019

low beta 4711. from file 19. at 02-aug-1994 10.44.51—slopefilec019.dat

Slope(mm/mr)—H

Input file—/afs/fnal/files/home/room1/gelfand/tracking/minult/data/slopeplot.plotf

11Nov-1994 9-06-22 Plot number— 2 Plot/symbol

11Nov-1994
9-06-22
Plot number— 2
Plot/symbol
ave = -4.687
σ = 15.05

Slopes— Kick at he13 Data from file—019

low beta 4711. from file 19. at 02-aug-1994 10.44.51—slopefilec019.dat

Slope(mm/mr)—V

Figure 1

Comparison of Computed and Measured Lattice Functions.

Figure-2

Tevatron Latttice with Low- β at B0 and D0

Tevatron Latttice with Low- β at B0 and D0

Figure-3b

Fit With Skew Quads Data From file 018

Input file=/afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plot1

Kick at he11

Input file=/afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plot1

Input file=/afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plot1

Input file=/afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plot1

Figure 4a

Input file—/ofs/1nol/files/home/room1/geifand/tracking/minute/skwsquad.plot

140-01-01
 140-01-02
 140-01-03
 140-01-04
 140-01-05
 140-01-06
 140-01-07
 140-01-08
 140-01-09
 140-01-10
 140-01-11
 140-01-12
 140-01-13
 140-01-14
 140-01-15
 140-01-16
 140-01-17
 140-01-18
 140-01-19
 140-01-20
 140-01-21
 140-01-22
 140-01-23
 140-01-24
 140-01-25
 140-01-26
 140-01-27
 140-01-28
 140-01-29
 140-01-30
 140-01-31
 140-01-32
 140-01-33
 140-01-34
 140-01-35
 140-01-36
 140-01-37
 140-01-38
 140-01-39
 140-01-40
 140-01-41
 140-01-42
 140-01-43
 140-01-44
 140-01-45
 140-01-46
 140-01-47
 140-01-48
 140-01-49
 140-01-50
 140-01-51
 140-01-52
 140-01-53
 140-01-54
 140-01-55
 140-01-56
 140-01-57
 140-01-58
 140-01-59
 140-01-60
 140-01-61
 140-01-62
 140-01-63
 140-01-64
 140-01-65
 140-01-66
 140-01-67
 140-01-68
 140-01-69
 140-01-70
 140-01-71
 140-01-72
 140-01-73
 140-01-74
 140-01-75
 140-01-76
 140-01-77
 140-01-78
 140-01-79
 140-01-80
 140-01-81
 140-01-82
 140-01-83
 140-01-84
 140-01-85
 140-01-86
 140-01-87
 140-01-88
 140-01-89
 140-01-90
 140-01-91
 140-01-92
 140-01-93
 140-01-94
 140-01-95
 140-01-96
 140-01-97
 140-01-98
 140-01-99
 140-02-00

Fit With Skew Quads Data From file 019
Kick at he13

```

11000-1004
14-26-14
Plot number- 5
Plot/symbol
N Reg. Stat      K
Ove = 5.8897E-02
O = 1.248
W Total =
Ove = 6.2107E-03
O = 1.251

```

Input File--/afs/fnc/files/home/room1/gelfand/tracking/minute/skewquad.plott

[illegible]

Fit With Skew Quads Data From file 019--Difference.
Kick at he13

11Nov-1984
14-26-14
Plot number- 6
Plot/symbol
H DPT, 30-40
ave= 1.4730E-02
σ= .2249

```
Input file=/afs/lnsl/files/home/room1/oelfand/tracking/minute/skewquad.plot
```

VPR046
 VPR037
 VPR022
 VPR018
 VPR014
 VPR040
 VPR036
 VPR033
 VPR029
 VPR027
 VPR025
 VPR023
 VPR021
 VPR019
 VPR017
 VPR015
 VPR013
 VPR011
 VPR009
 VPR007
 VPR005
 VPR003
 VPR001

Fit With Skew Quads Data From file 019
Kick at he13

```

11Bar-1894
M-28-14
Plot number - 7
Plot/symbol
V type: 28
Ovs = -1.4720E-02
Q = .4727
V total: 0
Ovs = -5.8058E-03
Q = .4148

```

Input file--/afs/tnl/files/home/room1/gettand/tracking/minuit/sknequad.plotf

VPR046
 VPR047
 VPR048
 VPR049
 VPR050
 VPR051
 VPR052
 VPR053
 VPR054
 VPR055
 VPR056
 VPR057
 VPR058
 VPR059
 VPR060
 VPR061
 VPR062
 VPR063
 VPR064
 VPR065
 VPR066
 VPR067
 VPR068
 VPR069
 VPR070
 VPR071
 VPR072
 VPR073
 VPR074
 VPR075
 VPR076
 VPR077
 VPR078
 VPR079
 VPR080
 VPR081
 VPR082
 VPR083
 VPR084
 VPR085
 VPR086
 VPR087
 VPR088
 VPR089
 VPR090
 VPR091
 VPR092
 VPR093
 VPR094
 VPR095
 VPR096
 VPR097
 VPR098
 VPR099
 VPR100
 VPR101
 VPR102
 VPR103
 VPR104
 VPR105
 VPR106
 VPR107
 VPR108
 VPR109
 VPR110
 VPR111
 VPR112
 VPR113
 VPR114
 VPR115
 VPR116
 VPR117
 VPR118
 VPR119
 VPR120
 VPR121
 VPR122
 VPR123
 VPR124
 VPR125
 VPR126
 VPR127
 VPR128
 VPR129
 VPR130
 VPR131
 VPR132
 VPR133
 VPR134
 VPR135
 VPR136
 VPR137
 VPR138
 VPR139
 VPR140
 VPR141
 VPR142
 VPR143
 VPR144
 VPR145
 VPR146
 VPR147
 VPR148
 VPR149
 VPR150
 VPR151
 VPR152
 VPR153
 VPR154
 VPR155
 VPR156
 VPR157
 VPR158
 VPR159
 VPR160
 VPR161
 VPR162
 VPR163
 VPR164
 VPR165
 VPR166
 VPR167
 VPR168
 VPR169
 VPR170
 VPR171
 VPR172
 VPR173
 VPR174
 VPR175
 VPR176
 VPR177
 VPR178
 VPR179
 VPR180
 VPR181
 VPR182
 VPR183
 VPR184
 VPR185
 VPR186
 VPR187
 VPR188
 VPR189
 VPR190
 VPR191
 VPR192
 VPR193
 VPR194
 VPR195
 VPR196
 VPR197
 VPR198
 VPR199
 VPR200
 VPR201
 VPR202
 VPR203
 VPR204
 VPR205
 VPR206
 VPR207
 VPR208
 VPR209
 VPR210
 VPR211
 VPR212
 VPR213
 VPR214
 VPR215
 VPR216
 VPR217
 VPR218
 VPR219
 VPR220
 VPR221
 VPR222
 VPR223
 VPR224
 VPR225
 VPR226
 VPR227
 VPR228
 VPR229
 VPR230
 VPR231
 VPR232
 VPR233
 VPR234
 VPR235
 VPR236
 VPR237
 VPR238
 VPR239
 VPR240
 VPR241
 VPR242
 VPR243
 VPR244
 VPR245
 VPR246
 VPR247
 VPR248
 VPR249
 VPR250
 VPR251
 VPR252
 VPR253
 VPR254
 VPR255
 VPR256
 VPR257
 VPR258
 VPR259
 VPR260
 VPR261
 VPR262
 VPR263
 VPR264
 VPR265
 VPR266
 VPR267
 VPR268
 VPR269
 VPR270
 VPR271
 VPR272
 VPR273
 VPR274
 VPR275
 VPR276
 VPR277
 VPR278
 VPR279
 VPR280
 VPR281
 VPR282
 VPR283
 VPR284
 VPR285
 VPR286
 VPR287
 VPR288
 VPR289
 VPR290
 VPR291
 VPR292
 VPR293
 VPR294
 VPR295
 VPR296
 VPR297
 VPR298
 VPR299
 VPR300
 VPR301
 VPR302
 VPR303
 VPR304
 VPR305
 VPR306
 VPR307
 VPR308
 VPR309
 VPR310
 VPR311
 VPR312
 VPR313
 VPR314
 VPR315
 VPR316
 VPR317
 VPR318
 VPR319
 VPR320
 VPR321
 VPR322
 VPR323
 VPR324
 VPR325
 VPR326
 VPR327
 VPR328
 VPR329
 VPR330
 VPR331
 VPR332
 VPR333
 VPR334
 VPR335
 VPR336
 VPR337
 VPR338
 VPR339
 VPR340
 VPR341
 VPR342
 VPR343
 VPR344
 VPR345
 VPR346
 VPR347
 VPR348
 VPR349
 VPR350
 VPR351
 VPR352
 VPR353
 VPR354
 VPR355
 VPR356
 VPR357
 VPR358
 VPR359
 VPR360
 VPR361
 VPR362
 VPR363
 VPR364
 VPR365
 VPR366
 VPR367
 VPR368
 VPR369
 VPR370
 VPR371
 VPR372
 VPR373
 VPR374
 VPR375
 VPR376
 VPR377
 VPR378
 VPR379
 VPR380
 VPR381
 VPR382
 VPR383
 VPR384
 VPR385
 VPR386
 VPR387
 VPR388
 VPR389
 VPR390
 VPR391
 VPR392
 VPR393
 VPR394
 VPR395
 VPR396
 VPR397
 VPR398
 VPR399
 VPR400
 VPR401
 VPR402
 VPR403
 VPR404
 VPR405
 VPR406
 VPR407
 VPR408
 VPR409
 VPR410
 VPR411
 VPR412
 VPR413
 VPR414
 VPR415
 VPR416
 VPR417
 VPR418
 VPR419
 VPR420
 VPR421
 VPR422
 VPR423
 VPR424
 VPR425
 VPR426
 VPR427
 VPR428
 VPR429
 VPR430
 VPR431
 VPR432
 VPR433
 VPR434
 VPR435
 VPR436
 VPR437
 VPR438
 VPR439
 VPR440
 VPR441
 VPR442
 VPR443
 VPR444
 VPR445
 VPR446
 VPR447
 VPR448
 VPR449
 VPR450
 VPR451
 VPR452
 VPR453
 VPR454
 VPR455
 VPR456
 VPR457
 VPR458
 VPR459
 VPR460
 VPR461
 VPR462
 VPR463
 VPR464
 VPR465
 VPR466
 VPR467
 VPR468
 VPR469
 VPR470
 VPR471
 VPR472
 VPR473
 VPR474
 VPR475
 VPR476
 VPR477
 VPR478
 VPR479
 VPR480
 VPR481
 VPR482
 VPR483
 VPR484
 VPR485
 VPR486
 VPR487
 VPR488
 VPR489
 VPR490
 VPR491
 VPR492
 VPR493
 VPR494
 VPR495
 VPR496
 VPR497
 VPR498
 VPR499
 VPR500

Fit With Skew Quads Data From file 019-Difference.
Kick at he13

```

17Mar-1984
14-20-14
Plot number- 8
Plot/symbol
V BR. 10-01
ave= -2.2751E-02
s= .7144

```

Figure 4b

Fit With Skew Quads Data From file 020

Input file=/afs/tna1/files/home/room1/gelfand/tracking/minuit/skewquad.plott

11Nov-1994 14-26-14
Plot number=9
Plot/symbol
N par= 4
Chi= 8.0904E-03
d= .4975
V total= 4.7808E-04
d= .3694

Fit With Skew Quads Data From file 020 Kick at ve11

Kick at ve11

Input file=/afs/tna1/files/home/room1/gelfand/tracking/minuit/skewquad.plott

11Nov-1994 14-26-14
Plot number=10
Plot/symbol
N par= 4
Chi= 3.4362E-03
d= .2546

Fit With Skew Quads Data From file 020-Difference. Kick at ve11

Input file=/afs/tna1/files/home/room1/gelfand/tracking/minuit/skewquad.plott

11Nov-1994 14-26-14
Plot number=11
Plot/symbol
N par= 4
Chi= 4.4005E-02
d= 1.262
V total= 3.9748E-02
d= 1.236

Fit With Skew Quads Data From file 020 Kick at ve11

Input file=/afs/tna1/files/home/room1/gelfand/tracking/minuit/skewquad.plott

11Nov-1994 14-26-14
Plot number=12
Plot/symbol
N par= 4
Chi= 1.1518E-03
d= .2180

Fit With Skew Quads Data From file 020-Difference. Kick at ve11

Figure 4c

Fit With Skew Quads Data From file 021

Input file: /afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plotf

11 Nov 1994
14-26-14
Plot number-13
Plot/symbol
H BPM, X
Q10 = 1.1619E-03
Q11 = .2240
H BPM, X
Q10 = -2.5689E-03
Q11 = .2239

Fit With Skew Quads Data From file 021

Kick at ve14

Distance from mk-c0 (m)

Kick at ve14

Input file: /afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plotf

11 Nov 1994
14-26-14
Plot number-14
Plot/symbol
H BPM, X
Q10 = 3.5213E-03
Q11 = .1199

Fit With Skew Quads Data From file 021-Difference.

Kick at ve14

Distance from mk-c0 (m)

Input file: /afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plotf

11 Nov 1994
14-26-14
Plot number-15
Plot/symbol
V BPM, X
Q10 = 5.7423E-02
Q11 = 1.053
V BPM, X
Q10 = 5.2579E-02
Q11 = 1.096

Fit With Skew Quads Data From file 021

Kick at ve14

Distance from mk-c0 (m)

Input file: /afs/lnal/files/home/room1/getfand/tracking/minuit/skewquad.plotf

11 Nov 1994
14-26-14
Plot number-16
Plot/symbol
V BPM, X
Q10 = -1.23747E-03
Q11 = .1447

Fit With Skew Quads Data From file 021-Difference.

Kick at ve14

Distance from mk-c0 (m)

Figure 4d

```
input file= /afs/ind/files/home/room1/gelfand/tracking/minut/relat.platt
```

1970-71
1971-72
1972-73
1973-74
1974-75
1975-76
1976-77
1977-78
1978-79
1979-80
1980-81
1981-82
1982-83
1983-84
1984-85
1985-86
1986-87
1987-88
1988-89
1989-90
1990-91
1991-92
1992-93
1993-94
1994-95
1995-96
1996-97
1997-98
1998-99
1999-00
2000-01
2001-02
2002-03
2003-04
2004-05
2005-06
2006-07
2007-08
2008-09
2009-10
2010-11
2011-12
2012-13
2013-14
2014-15
2015-16
2016-17
2017-18
2018-19
2019-20
2020-21
2021-22
2022-23
2023-24
2024-25
2025-26
2026-27
2027-28
2028-29
2029-30
2030-31
2031-32
2032-33
2033-34
2034-35
2035-36
2036-37
2037-38
2038-39
2039-40
2040-41
2041-42
2042-43
2043-44
2044-45
2045-46
2046-47
2047-48
2048-49
2049-50
2050-51
2051-52
2052-53
2053-54
2054-55
2055-56
2056-57
2057-58
2058-59
2059-60
2060-61
2061-62
2062-63
2063-64
2064-65
2065-66
2066-67
2067-68
2068-69
2069-70
2070-71
2071-72
2072-73
2073-74
2074-75
2075-76
2076-77
2077-78
2078-79
2079-80
2080-81
2081-82
2082-83
2083-84
2084-85
2085-86
2086-87
2087-88
2088-89
2089-90
2090-91
2091-92
2092-93
2093-94
2094-95
2095-96
2096-97
2097-98
2098-99
2099-00
2100-01
2101-02
2102-03
2103-04
2104-05
2105-06
2106-07
2107-08
2108-09
2109-10
2110-11
2111-12
2112-13
2113-14
2114-15
2115-16
2116-17
2117-18
2118-19
2119-20
2120-21
2121-22
2122-23
2123-24
2124-25
2125-26
2126-27
2127-28
2128-29
2129-30
2130-31
2131-32
2132-33
2133-34
2134-35
2135-36
2136-37
2137-38
2138-39
2139-40
2140-41
2141-42
2142-43
2143-44
2144-45
2145-46
2146-47
2147-48
2148-49
2149-50
2150-51
2151-52
2152-53
2153-54
2154-55
2155-56
2156-57
2157-58
2158-59
2159-60
2160-61
2161-62
2162-63
2163-64
2164-65
2165-66
2166-67
2167-68
2168-69
2169-70
2170-71
2171-72
2172-73
2173-74
2174-75
2175-76
2176-77
2177-78
2178-79
2179-80
2180-81
2181-82
2182-83
2183-84
2184-85
2185-86
2186-87
2187-88
2188-89
2189-90
2190-91
2191-92
2192-93
2193-94
2194-95
2195-96
2196-97
2197-98
2198-99
2199-00
2200-01
2201-02
2202-03
2203-04
2204-05
2205-06
2206-07
2207-08
2208-09
2209-10
2210-1

Fit With Rolled Quads— Data from file 018
Kick at he11

Distance from mk-c0 (m)

Input file—/afs/ind/files/home/room1/gelland/tracking/minut/rolled.plott

1079-17
1079-20
1079-23
1079-26
1079-28
1079-30
1079-32
1079-34
1079-36
1079-38
1079-40
1079-42
1079-44
1079-46
1079-48
1079-50
1079-52
1079-54
1079-56
1079-58
1079-60
1079-62
1079-64
1079-66
1079-68
1079-70
1079-72
1079-74
1079-76
1079-78
1079-80
1079-82
1079-84
1079-86
1079-88
1079-90
1079-92
1079-94
1079-96
1079-98
1080-00

File With Rolled Quads- Data from file 018-Difference.
Kick at hell

Distance from mk-c0 (m)

Input file-- /afs/indiana.edu/home/room1/celland/tracking/minut/rolled.platt

[illegible]

Fit With Rolled Quads- Data from file 018
Kick at he11

Distance from mk-c0 (m)

Invert file: /afs/ind/files/home/room1/oatland/tracking/minute/roled.plott

WPR-2.7	WPR-2.8	WPR-2.9	WPR-3.0	WPR-3.1	WPR-3.2	WPR-3.3	WPR-3.4	WPR-3.5	WPR-3.6	WPR-3.7	WPR-3.8	WPR-3.9	WPR-4.0	WPR-4.1	WPR-4.2	WPR-4.3	WPR-4.4	WPR-4.5	WPR-4.6	WPR-4.7	WPR-4.8	WPR-4.9	WPR-5.0	WPR-5.1	WPR-5.2	WPR-5.3	WPR-5.4	WPR-5.5	WPR-5.6	WPR-5.7	WPR-5.8	WPR-5.9	WPR-6.0	WPR-6.1	WPR-6.2	WPR-6.3	WPR-6.4	WPR-6.5	WPR-6.6	WPR-6.7	WPR-6.8	WPR-6.9	WPR-7.0	WPR-7.1	WPR-7.2	WPR-7.3	WPR-7.4	WPR-7.5	WPR-7.6	WPR-7.7	WPR-7.8	WPR-7.9	WPR-8.0	WPR-8.1	WPR-8.2	WPR-8.3	WPR-8.4	WPR-8.5	WPR-8.6	WPR-8.7	WPR-8.8	WPR-8.9	WPR-9.0	WPR-9.1	WPR-9.2	WPR-9.3	WPR-9.4	WPR-9.5	WPR-9.6	WPR-9.7	WPR-9.8	WPR-9.9	WPR-10.0	WPR-10.1	WPR-10.2	WPR-10.3	WPR-10.4	WPR-10.5	WPR-10.6	WPR-10.7	WPR-10.8	WPR-10.9	WPR-11.0	WPR-11.1	WPR-11.2	WPR-11.3	WPR-11.4	WPR-11.5	WPR-11.6	WPR-11.7	WPR-11.8	WPR-11.9	WPR-12.0	WPR-12.1	WPR-12.2	WPR-12.3	WPR-12.4	WPR-12.5	WPR-12.6	WPR-12.7	WPR-12.8	WPR-12.9	WPR-13.0	WPR-13.1	WPR-13.2	WPR-13.3	WPR-13.4	WPR-13.5	WPR-13.6	WPR-13.7	WPR-13.8	WPR-13.9	WPR-14.0	WPR-14.1	WPR-14.2	WPR-14.3	WPR-14.4	WPR-14.5	WPR-14.6	WPR-14.7	WPR-14.8	WPR-14.9	WPR-15.0	WPR-15.1	WPR-15.2	WPR-15.3	WPR-15.4	WPR-15.5	WPR-15.6	WPR-15.7	WPR-15.8	WPR-15.9	WPR-16.0	WPR-16.1	WPR-16.2	WPR-16.3	WPR-16.4	WPR-16.5	WPR-16.6	WPR-16.7	WPR-16.8	WPR-16.9	WPR-17.0	WPR-17.1	WPR-17.2	WPR-17.3	WPR-17.4	WPR-17.5	WPR-17.6	WPR-17.7	WPR-17.8	WPR-17.9	WPR-18.0	WPR-18.1	WPR-18.2	WPR-18.3	WPR-18.4	WPR-18.5	WPR-18.6	WPR-18.7	WPR-18.8	WPR-18.9	WPR-19.0	WPR-19.1	WPR-19.2	WPR-19.3	WPR-19.4	WPR-19.5	WPR-19.6	WPR-19.7	WPR-19.8	WPR-19.9	WPR-20.0	WPR-20.1	WPR-20.2	WPR-20.3	WPR-20.4	WPR-20.5	WPR-20.6	WPR-20.7	WPR-20.8	WPR-20.9	WPR-21.0	WPR-21.1	WPR-21.2	WPR-21.3	WPR-21.4	WPR-21.5	WPR-21.6	WPR-21.7	WPR-21.8	WPR-21.9	WPR-22.0	WPR-22.1	WPR-22.2	WPR-22.3	WPR-22.4	WPR-22.5	WPR-22.6	WPR-22.7	WPR-22.8	WPR-22.9	WPR-23.0	WPR-23.1	WPR-23.2	WPR-23.3	WPR-23.4	WPR-23.5	WPR-23.6	WPR-23.7	WPR-23.8	WPR-23.9	WPR-24.0	WPR-24.1	WPR-24.2	WPR-24.3	WPR-24.4	WPR-24.5	WPR-24.6	WPR-24.7	WPR-24.8	WPR-24.9	WPR-25.0	WPR-25.1	WPR-25.2	WPR-25.3	WPR-25.4	WPR-25.5	WPR-25.6	WPR-25.7	WPR-25.8	WPR-25.9	WPR-26.0	WPR-26.1	WPR-26.2	WPR-26.3	WPR-26.4	WPR-26.5	WPR-26.6	WPR-26.7	WPR-26.8	WPR-26.9	WPR-27.0	WPR-27.1	WPR-27.2	WPR-27.3	WPR-27.4	WPR-27.5	WPR-27.6	WPR-27.7	WPR-27.8	WPR-27.9	WPR-28.0	WPR-28.1	WPR-28.2	WPR-28.3	WPR-28.4	WPR-28.5	WPR-28.6	WPR-28.7	WPR-28.8	WPR-28.9	WPR-29.0	WPR-29.1	WPR-29.2	WPR-29.3	WPR-29.4	WPR-29.5	WPR-29.6	WPR-29.7	WPR-29.8	WPR-29.9	WPR-30.0	WPR-30.1	WPR-30.2	WPR-30.3	WPR-30.4	WPR-30.5	WPR-30.6	WPR-30.7	WPR-30.8	WPR-30.9	WPR-31.0	WPR-31.1	WPR-31.2	WPR-31.3	WPR-31.4	WPR-31.5	WPR-31.6	WPR-31.7	WPR-31.8	WPR-31.9	WPR-32.0	WPR-32.1	WPR-32.2	WPR-32.3	WPR-32.4	WPR-32.5	WPR-32.6	WPR-32.7	WPR-32.8	WPR-32.9	WPR-33.0	WPR-33.1	WPR-33.2	WPR-33.3	WPR-33.4	WPR-33.5	WPR-33.6	WPR-33.7	WPR-33.8	WPR-33.9	WPR-34.0	WPR-34.1	WPR-34.2	WPR-34.3	WPR-34.4	WPR-34.5	WPR-34.6
---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	---------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------	----------

Fit With Rolled Quads- Data from file 018-Difference.
Kick at hell

Distance from mk-c0 (m)

Figure 5a

lnst file- /afs/ind/files/home/room1/galland/tracting/minut/roled.plott

input file- /etc/ind/fin/home/room1/gellond/tracking/mini4/rolled.plott

Input file--/afs/ind/files/home/room1/galland/trackings/minis/rolled.platt

Input file- /ots/inst/files/home/room1/gellond/tracking/minute/rolled.pict

Figure 5b


```
next file= /afs/trng/files/home/room1/galland/tracting/minut/rolled.plott
```

Input file- /afs/ind/Files/home/room1/gottend/tracking/minute/rotated.plott

Input file- /opt/ind/files/home/room1/gelfand/training/minil/rolled-plot1

next file: /afs/ind/files/home/room1/coland/tracking/minut/rolled.pict

Figure 5c

Fit With Rolled Quads- Data from file 021

Input file: /afs/ind/rls/home/room1/gelland/tracking/minut/rolled.plott

11/10/84 09:30:47 0-31-47 Plot number-13 Plot/symbol

Form factor
dipole = 1.1619E-03
dipole = 2240
dipole = -2.3598E-03
dipole = 2.359

Fit With Rolled Quads- Data from file 021 Kick at ve14

Kick at ve14

Input file: /afs/ind/rls/home/room1/gelland/tracking/minut/rolled.plott

11/10/84 09:30:47 0-31-47 Plot number-14 Plot/symbol

Form factor
dipole = 1.1619E-03
dipole = 2240
dipole = -2.3598E-03
dipole = 2.359

Fit With Rolled Quads- Data from file 021-Difference. Kick at ve14

Input file: /afs/ind/rls/home/room1/gelland/tracking/minut/rolled.plott

11/10/84 09:30:47 0-31-47 Plot number-15 Plot/symbol

Form factor
dipole = 5.7423E-02
dipole = 1.053
dipole = 5.2739E-02
dipole = 1.054

Fit With Rolled Quads- Data from file 021 Kick at ve14

Input file: /afs/ind/rls/home/room1/gelland/tracking/minut/rolled.plott

11/10/84 09:30:47 0-31-47 Plot number-16 Plot/symbol

Form factor
dipole = 1.9794E-03
dipole = 1.527

Fit With Rolled Quads- Data from file 021-Difference. Kick at ve14

Figure 5d

```
ls -l /etc/passwd | grep root
```

```
input file=/afs/ins/Files/home/room1/getfend/tracking/minuul/unrolled.p01
```

Figure 6a

[illegible] [illegible]

Figure 6b

```
Input file=/afs/lncl/files/home/room1/galland/tracking/result/unrolled.plott
```

Input file- /afs/fns/fhs/home/room1/gelfand/tracking/minuit/unrolled.plott

Figure 6c

[illegible]

Figure 6d

Input file= /p/s/fns/files/home/room1/oefond/tracking/minid/segmentrel.plott

```
input file--/afs/lns/files/home/room1/gallond/tracking/minut/septunroll.plott
```

Figure 7a

host file=/s/tno/ftse/home/room1/gatend/tracking/minit/septunroll.pltf

1000-8954
11-69-01
Flat number- 5
Flat/symbol
area= — at
dev=- -3.0078E-03
pm 1.150
Wt/dm = —
gm = 0.5205E-03
g= 1.187

```
input file=/afs/frag/frag/home/room1/gelfond/tracking/minut/septunreil.plott
```

5444-1084
 11-07-01
 Plot number - 6
 Plot/symbol
 N GRT. 20-04
 000- -1.4070E-02
 0- .1300

```
input file=/afs/tnsl/files/home/room1/gelland/tracking/minull/septunroll.plott
```

Yeller-9004
11-07-01
Plot number- 7
Plot/symbol
Veg. 2
veg= -5.3045E-03
gr= .1646
Vital: -
temp= -7.243E-04
min. 4.5200E-02

input file- /afs/trn/fber/home/room1/selfand/tracking/minut/septunroll.plots

100m-1004
11-07-01
Plot number - B
Plot/symbol
V BE. 204
ave = -5.000E-01
std = .1540

Figure 7b

After the unrolls in Sept. Data from file 252

Input file: /afs/tnsl/files/home/room1/gelfand/tracking/minut/septunroll.plott

Kick at ve11

Input file: /afs/tnsl/files/home/room1/gelfand/tracking/minut/septunroll.plott

Input file: /afs/tnsl/files/home/room1/gelfand/tracking/minut/septunroll.plott

Input file: /afs/tnsl/files/home/room1/gelfand/tracking/minut/septunroll.plott

Figure 7c

After the unrolls in Sept. Data from file 253

Input file=/afs/tns/tns/home/room1/gatond/tracking/minut/septunroll.plott

Kick at ve14

Input file=/afs/tns/tns/home/room1/gatond/tracking/minut/septunroll.plott

Input file=/afs/tns/tns/home/room1/gatond/tracking/minut/septunroll.plott

Input file=/afs/tns/tns/home/room1/gatond/tracking/minut/septunroll.plott

Figure 7d

$\langle \sigma_x \sigma_y \rangle^{1/2}$ vs turn

22Nov-1994

12-22-24

Multiparticle Tracking. Analysis mode-31 1000 turns.
Input file-/afs/fnol/files/home/room1/gelfand/tracking/plotfiles/sigxsigy.study.plotf

Plot number- 1

Plot/symbol

design
mk-b0 *-*
ave= 4.3405E-02
 σ = 8.2444E-06
mk-d0 *-*
ave= 4.2989E-02
 σ = 9.9299E-06
rolled
mk-b0 *-*
ave= 6.2711E-02
 σ = 1.0631E-04
mk-d0 *-*
ave= 4.6534E-02
 σ = 7.3479E-05
unroll
mk-b0 —
ave= 4.6736E-02
 σ = 1.6035E-04
mk-d0 *-*
ave= 4.3284E-02
 σ = 1.0845E-04
septunroll
mk-b0 +--+
ave= 4.3652E-02
 σ = 2.7398E-05
mk-d0 *-*
ave= 4.3286E-02
 σ = 2.0019E-05

Figure 8

Input file=/afs/fnal/files/home/room1/gelfand/tracking/plotfiles/plotdata.plotf

6Dec-1994
10-33-28
Plot number= 1
Plot/symbol
BO $\langle \sigma_x \sigma_y \rangle^m$
ave= 5.3746E-02
 $\sigma = 9.1626E-03$

Data From PLOTDB
Extracted 11/21-22/94

Input file=/afs/fnal/files/home/room1/gelfand/tracking/plotfiles/plotdata.plotf

6Dec-1994
10-33-28
Plot number= 2
Plot/symbol
DO $\langle \sigma_x \sigma_y \rangle^m$
ave= 5.2154E-02
 $\sigma = 5.1671E-03$

Data From PLOTDB
Extracted 11/21-22/94

Figure 9