MODULATION OF LYMPHOPOIESIS

**COMPREHENSIVE PROGRESS REPORT
FOR THE PERIOD
JANUARY 1, 1991 - JULY 30, 1991**

**CORNELIUS ROSSE, M.D., D.SC.,
PROFESSOR & CHAIRMAN
DEPARTMENT OF BIOLOGICAL STRUCTURE**

**UNIVERSITY OF WASHINGTON
SEATTLE, WASHINGTON 98195**

GRANT DE-FG06-86ER60409

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

LIST OF ABBREVIATIONS

BM	bone marrow
BMC	bone marrow cells
BMSC	bone marrow stromal cells
C	complement
CE maca	CE1460 mammary carcinoma
CFU-N	colony forming unit - neutrophil
CM	conditioned media
DBA	<i>Dolichos bifloris</i> agglutinin
G-CSF	granulocyte - colony stimulating factor
GM-CSF	granulocyte/macrophage colony stimulating factor
H	host
LAK	lymphokine activated killer cell
LTBMC	long term bone marrow cultures
mAb	monoclonal antibody
M-CSF	macrophage - colony stimulating factor
MPA	mycophenolic acid
NH	nonhost
NK	natural killer
SPL	spleen
THY	thymus
W/W	Whitlock Witte

INTRODUCTION

Our research work supported by the Department of Energy has been concerned for a number of years with the understanding of lymphopoiesis. Rather than investigating the noxious effects of various environmental agents on immune function or on specific components of the immune system *per se*, we have taken a much broader approach that has addressed the fundamental question of cell differentiation and cell production in the primary lymphoid organs. An understanding of the mechanisms that regulate the production and differentiation of B cells, natural killer (NK) cells, and T cells is requisite for investigating how the immune system might be affected by any agent, extrinsic or intrinsic, that exerts a more subtle modularity effect on cell systems of host defense than the direct killing of susceptible cells. In our work we have made use of *in vivo* and *in vitro* experimental models in which differentiation of select classes of lymphocytes was profoundly altered. In our search to define the mechanisms that were responsible for such changes, we have contributed significantly to the understanding of the regulation of lymphopoiesis. Approaching lymphopoiesis as a whole has enabled us to draw instructive parallels between the lymphocyte subclasses. These parallels have informed the questions we could formulate and the specific experiments we designed to address them. Our Comprehensive Progress Report documents the effectiveness of this approach.

Our productivity on several interrelated fronts of lymphopoiesis has been possible because of the team of investigators we assembled. The funding limitations that have been imposed on us by DOE's budgetary constraints have forced us, however, during two successive project periods to limit the scope of our approach and to prioritize the specific aims of our program and determine which aims would be best pursued under DOE's sponsorship and which under the sponsorship of various NIH institutes. Thus, recently we have focused primarily on the modulation of B cell production and on the establishment of an *in vitro* system for studying NK cell differentiation. The study of T cell differentiation through analyzing the intrathymic microenvironment and exploring transcriptional regulation of T cell development by specific domains of a retroviral genome, though initially supported by DOE, had to be sustained by other funds. The support DOE has provided for these studies was, however, an important factor in our ability to successfully compete for specifically focused grants. The same evolution occurred with our work on evaluating the role of intrathymic and hematogenous (bone marrow-derived) T cell precursors in thymocytopoiesis, which now continues with NIH support.

As a result of our work and the work of other laboratories there are now available *in vitro* models for studying B cell and NK cell differentiation. These models lend themselves for exploring agents and factors that can modulate lymphopoiesis. Our work sponsored by DOE has substantially contributed to these developments. During the current project period we have demonstrated correspondence between animal models and *in vitro* models of modulated lymphopoiesis. Our finding that G-CSF, a growth factor for neutrophil granulocytes, suppresses lymphopoiesis in long term bone marrow cultures (LTBMC) has important implications both for understanding the regulatory mechanisms of hemopoiesis and for clinical use of recombinant growth factors that are beginning to be widely used for the treatment of a variety of diseases. The relevance of this finding to the interests of the Health Effects Research Division of DOE is enhanced by the fact that suppressed

lymphopoieses in our CE maca animal model must also be ascribed to the secretion of G-CSF by the transplanted tumor. Although there is no direct evidence that this tumor was originally induced by environmental agents, the carcinogenicity of such agents is well documented and is a major justification for the support DOE provides for the type of research our laboratory conducts.

During the present project period we adopted LTBM systems developed by others for the purposes of our specific aims. Also we developed a novel long term culture system for NK cells. The discovery of a new growth factor, O-CSF, specific for osteoclasts and the establishment of a clonal assay system that provides evidence for a new class of hemopoietic progenitor cells, the osteoclast progenitor, are important contributions. Although these findings must be judged primarily on their own merit, they do have implications for lymphopoiesis because of their importance in bone-bone marrow interactions and their possible modulatory effects on lymphopoiesis.

We regard our accomplishments during the past and current project periods as persuasive justification for the support of the studies we propose for the next project period. The establishment of a long term culture system for T lymphopoiesis has defied long standing attempts in many laboratories. Given the important role T cells play in the immune response and in the regulation of other lymphohemopoietic cell lineages through the lymphokines they secrete, the need for an *in vitro* system that lends itself to the analysis of T cell maturation and to the testing of factors that may adversely affect T lymphopoiesis cannot be overemphasized. We believe that we can exploit an advantageous set of circumstances that present an excellent opportunity for initiating a focused experimental program for developing such a system. Our current and past accomplishments are the result of collaboration in a well-coordinated team of investigators, each with his own area of focus and expertise. A member of our team, Dr. Andrew Farr, has accumulated extensive experience and insight into the structure and biology of the thymic environment. An important addition is Dr. Alejandro Aruffo, whose appointment in the Department of Biological Structure is now being processed. Dr. Aruffo's knowledge and expertise in molecular interactions between leukocytes and endothelial cells extends the scope of our work into new domains. The productivity of Dr. Aruffo's collaboration with us is well-documented in the Preliminary Results section of our Research Proposal. By a systematic and selective analysis of molecular interactions between heterogeneous thymic stromal cells and T cell progenitors at different stages of maturation, it will be possible for our program to define the complement of critical cellular interactions on which successive stages of T lymphopoiesis depend. The experiments we propose will lay a rational foundation for the development of a long term culture system for T lymphopoiesis.

The investigation of molecular mechanisms that support lymphopoiesis in the thymus (THY) is a logical extension of the work we have been pursuing under DOE sponsorship. Its relevance to the mission of the Health Effects Research Division of DOE is based on our belief that continues to motivate our work:

1. In order to evaluate the biological modifying effects exerted by various environmental agents, we need a better understanding of the natural defense mechanisms of the host. When these mechanisms become overwhelmed or

compromised through the selective elimination or overproduction of certain cell types involved in host defense, a permissive state is created for the overt manifestation of disease or the uncontrolled growth of host cells altered by the environmental carcinogen.

2. Whether or not neoplastic cells induced by environmental agents establish themselves as cancer may be less dependent on the inducing agent than on the effectiveness of the host mechanisms capable of recognizing transformed cells and controlling their growth.

Accomplishment of the aims of the research we propose will result in a better understanding of molecular interaction that influence the development of T cells that play a critical role in the immune response and in the regulation of hemopoiesis and lymphopoiesis.

COMPREHENSIVE PROGRESS REPORT

I. Personnel Involved January 1, 1991 - July 30, 1991

	<u>Time</u>	<u>Salary</u>
Cornelius Rosse, M.D., D.Sc. Professor and Chairman Principal Investigator	20%	10%
Sylvia B. Pollack, Ph.D. Research Professor	20%	20%
Minako Y. Lee, M.D. Research Associate Professor	40%	40%
Fayth K. Yoshimura, Ph.D. Research Associate Professor	10%	10%
Andrew G. Farr, Ph.D. Associate Professor	10%	0%
Alejandro Aruffo, Ph.D. Research Associate Professor	25%	0%
Jose L. Mejino, M.D. Postdoctoral Fellow	100%	100%
Jody Lottsfeldt Research technician III	20%	20%
Arnita Lawrence Hourly Help	10%	15%
Curt Fox Hourly Help	5%	5%
Lasmy Lim Hourly Help	-	-

SCIENTIFIC STAFF

Description of members of the scientific staff include those individuals who have contributed to the research during the present project period as well as those who will be newly involved during the period for which funds are being requested.

Cornelius Rosse, M.D., D. Sc., Professor and Chairman, Principal Investigator

Was associated as co-investigator with contracts sponsored by the Department of Energy and its predecessors between 1967 and 1979 under the Principal Investigatorship of Dr. N.B. Everett. From 1979 - 1985 he was P.I. on contract: Marrow-Tumor Interactions; from 1986-1988 P.I. on grant: Immunodeficiency Syndromes; and from 1989-1991 P.I. on grant: Modulation of Lymphopoiesis.

Sylvia B. Pollack, Ph.D., Research Professor

Joined the program in 1981 from the Fred Hutchinson Cancer Research Center and brought with her extensive experience with *in vivo* and *in vitro* techniques in tumor immunobiology. Her work focuses on marrow-derived and marrow-dependent natural killer cells. The development of a long term bone marrow culture system that supports NK lymphopoiesis is an important contribution. While our proposed work will benefit from her presence as a member of the Biological Structure Research laboratory, her continued work on NK cells will be supported by her independently funded grants.

Minako Y. Lee, M.D., Research Associate Professor

During the present project period she discovered a new growth factor specific for osteoclasts and developed a clonal assay system for osteoclast progenitors. Her studies with the modulatory effects of G-CSF and CE maca on lymphopoiesis were primarily coordinated by her. She will pursue her interests in bone-bone marrow interaction and the regulation of lymphopoiesis by hemopoietic growth factors supported by independently funded grants, and will remain a member of our team.

Andrew Farr, Ph.D., Associate Professor

Joined the faculty in 1982 from the Basel Institute of Immunology. He has developed a program concerned with intrathymic cell production and cell differentiation with a major focus on thymic stroma and expression on the T cell receptor. Although he did not derive any salary from the DOE grant during the present project period, he has extensively contributed to the accomplishment of our aims. The work we propose for the next grant period is largely based on his accomplishments and he will assume responsibility for a major part of the experimental work we propose.

Alejandro Aruffo, Ph.D., Research Assistant Professor

Following his recruitment in 1990 as scientist to Bristol Meyers Squibb Laboratories in Seattle, he has established highly productive collaborations with Dr. Farr's program and the X-ray crystallographers in the Department of Biological Structure. His affiliate appointment in the Department of Biological Structure is being processed at the time of submitting this proposal. His highly regarded work on molecular interactions responsible for leukocyte binding to endothelium that control lymphocyte recirculation augments and extends the experimental approaches we have been using in our research. Our proposed aims rely on the collaborations he has already established with us.

Fayth K. Yoshimura, Ph.D., Research Associate Professor

Joined our group in 1986 from the Fred Hutchinson Cancer Research Center and brought with her extensive experience in molecular biology, molecular genetics, and virology. Her independently funded program has mainly been concerned with the role of murine retroviral sequences that enhance adjacent cellular genes in the generation of thymic lymphomas. Her collaborations with other members of the group studying lymphocyte development have been germane to new directions in our program.

Jose L. Mejino, M.D., Senior Fellow

Joined the program in 1984 after residency training in internal medicine. He is responsible for studies related to T cell development and cell traffic and collaborates with Drs. Rosse, Farr, and Lee. Dr. Mejino has become our "expert" in flow cytometry.

Marshall P. Welch, M.D., Senior Fellow

Joined the program with extensive training in Dermatology at the National Jewish Center for Immunology and Respiratory Medicine and at the University of Washington. With his knowledge of and background in epithelial biology, he is a valuable resource in our studies of thymic epithelium. He has contributed significantly to the progress we have made in characterizing epithelial cell surface proteins thus far and was directly responsible for the cloning of the gene encoding a novel and functionally important stromal cell surface molecule.

Students

Sherree Friend, Graduate Student

Supported by an NSF fellowship, she has been involved in developing thymic stromal cell lines and the generation of monoclonal antibodies specific for different populations of stromal cells. She will be involved in the generation of anti-integrin antibodies and examining their effects on lymphopoiesis *in vitro*.

Melissa Berry, Graduate Student

Supported by a training grant, she is involved in the cloning of genes encoding unique stromal cell surface molecules. She will be involved in the direct expression cloning of cell interaction molecules.

Noma Ladendorff, Graduate Student

Supported by a training grant, she has been studying the expression of integrins and other cell interaction molecules within the thymus.

Walter Medlin, Medical Student

As part of a summer research training program, he has been studying the role of thymic stromal cells in the clonal elimination of thymocytes in response to superantigens.

Todd Winter, Medical Student

As part of a summer research training program, he has examined the ontogeny of thymic stromal cell development.

Tina Yamigawa, Undergraduate

As a Hughes Scholar, she is doing undergraduate research, supervised by Dr. Farr, on thymocyte binding to thymic stromal cell lines in culture.

Duyen Trang, Undergraduate

Over the past 3 years, she has been doing undergraduate research, supervised by Dr. Farr, on immunohistochemical studies of the thymic environment.

Technical Staff

Joan L. Lottsfeldt, Research Tech II - 20%

Bryan Golden, Research Tech II - 100%

Hourly Help/Student Help

Arnita Lawrence, Curt Fox, Lasmy Lim, M. Bellows

II. PUBLICATIONS

1. Lee, M.Y., Kaushansky, K., Judkins, S.A., Lottsfeldt, J.L., Waheed, A. and Shadduck, R.K. Mechanisms of tumor-induced neutrophilia: Constitutive production of colony-stimulating factors and their synergistic actions. *Blood* 74: 115-122, 1989.
2. Lee, M.Y. and Lottsfeldt, J.L. A hypercalcemia-inducing murine mammary carcinoma elaborates a colony stimulating factor selective for the putative osteoclast progenitor. *Exp. Hematol.* 18:608, 1990.
3. Lee, M.Y., Osborne, W.R.A. and Eyre, D.R. Isolation of a murine osteoclast colony stimulating factor (O-CSF). *Proc. Natl. Acad. Sci. USA* 1991 (Submitted)
4. Lee, M.Y. and Nagata, S. Bone marrow expansion and bone modulation by granulocyte colony stimulating factor. *Blood* 74:269a, 1989
5. Lee, M.Y. and Lottsfeldt, J.L. A murine mammary tumor produces a colony stimulating factor for the putative osteoclast progenitors. *Blood* 76: 151a, 1990
6. Lee, M.Y., Fukunaga, R., Lee, T.J., Lottsfeldt, J.L. and Nagata, S. Bone modulation in sustained hemopoietic stimulation in mice. *Blood*, 1991 (In press)
7. Pollack, S.B., Lee, M.Y., Cohen, E., Lottsfeldt, J.L. and Rosse, C. Modulation of murine natural killer cells by a granulocytosis-inducing tumor. *Cancer Res.* 49: 174-178, 1989.
8. Head, A.A., Lee, M.Y., and Pollack, S.B. Effects of a granulocytosis-inducing murine mammary carcinoma on natural killer cell activity. *Exp. Hematol.* 18:588, 1990.
9. Pollack, S.B., Gage, S. and Tsugi, J. Enrichment of NK cells and their IL-2 responsive precursors from murine bone marrow. *Cell. Immunol.* 1991 (Submitted)
10. Pollack, S.B., Tsugi, J. and Rosse, C. Production and differentiation of NK lineage cells in long term bone marrow cultures in the absence of exogenous IL-2. *Cell. Immunol.* 1991 (Submitted).
11. Pollack, S.B., Tsugi, J. and Rosse, C. Maturation of natural killer cells on bone marrow stromal cells. *Exp. Hematol.* 18:588, 1990.

12. Pollack, S.B., Tsugi, J. and Rosse, C. Natural killer (NK) cells develop in long term bone marrow cultures (LTBMC) in the absence of exogenous growth factors. *FASEB J.* 5:A1458, 1991.
13. Pollack, S.B., Tsugi, J., Linnemeyer, P. and Rosse, C. Production and differentiation of NK cells in murine long term bone marrow cultures. *Nat. Immun. Cell Growth Regul.* 10:158, 1991.
14. Mejino, J.L.V., Jr., Lee, M., Hamilton, B.L. and Rosse, C. The role of hematogenous and intrinsic precursor cells in lymphocyte production in the bone marrow. *Am. J. Anat.* 1991 (In Press)
15. Mejino, J.L.V. Jr. and Rosse, C. Thymocyte lineages derived from hematogenous and intrinsic progenitors. *Exp. Hematol.* 18:587, 1990.
16. Rosse, C. and Mejino, J.L.V. Jr. Thymic regeneration following corticosteroid-induced involution effected by intrathymic T cell progenitors. *Exp. Hematol.* 18:585, 1990.
17. Farr, A.G., Hosier, S., Braddy, S.C., Anderson, S.K., Eisenhardt, D.J., Yan, Z.J. and Robles, C.P. Medullary epithelial cell lines from murine thymus constitutively secrete IL-1 and hematopoietic growth factors and express class II antigens in response to recombinant interferon- γ . *Cell. Immunol.* 119:427-444, 1989.
18. Scofield, K., Yan, Z.J. and Farr, A.G. Characterization of murine thymocyte subpopulations reacting with *Dolichos bifloris* agglutinin. *Cell. Immunol.* 120:114-125, 1989.
19. Farr, A.G., Anderson, S.K. and Kanagawa, O. 1991 Apoptosis of V β 6⁺ thymocytes in the cortical Mls-1^a thymic environment. *Int. Immunol.* (submitted)
20. Farr, A., Nelson, A., and Hosier, S. 1991 Characterization of an antigenic determinant expressed by type I epithelial cells in the murine thymus. *J. Histochem. Cytochem.* (Submitted)
21. Hollon, T. and Yoshimura, F.K. Mapping of functional regions of murine retrovirus long terminal repeat enhancers: enhancer domains interact and are not independent in their contributions to enhancer activity. *J. Virol.* 63: 3353-3361, 1989.
22. Yoshimura, F.K., Tupper, J. and Diem, K. Differential DNA binding of nuclear proteins to a long terminal repeat region of the MCF13 and Akv murine leukemia viruses. *J. Virol.* 63: 4945-4948, 1989.

23. Hollon, T. and Yoshimura, F.K. A non-uniform electrical field electroporation chamber design. *Anal. Biochem.* 182: 419-423, 1989.
24. Hollon, T. and Yoshimura, F.K. Variation in enzymatic transient gene expression assays. *Anal. Biochem.* 182: 411-418, 1989.

III. ACCOMPLISHMENTS

The 24 appendices attached to this Comprehensive Progress Report and renewal application document most of our accomplishments during the 2-1/2 years of the current project period. Much additional work that is still in progress or being prepared for publication is described in this report. We have completed most of the work we proposed in Aims 1,2 and 3 but had to delete Aim 4 because of the budgetary cuts imposed by DOE.

A. ACCOMPLISHMENTS IN AIM 1: ABILITY OF STROMA FROM TUMOR-BEARING MICE TO SUPPORT LYMPHOPOIESIS

The purpose of this Aim was to answer the question: **Can stromal cell cultures from CE maca-bearing mice support B-lymphopoiesis?** A comprehensive set of experiments has been completed that give clear cut answers, but these studies have not yet been written up for publication. The key experimental data on which our conclusions are based are described below:

Long term bone marrow cultures were established under Whitlock-Witte (W/W) conditions from CE maca tumor-bearing mice. Fourteen and twenty-one days after the tumor had been transplanted all the mice used in the experiments had a greatly elevated neutrophil count. Similar cultures were initiated with the bone marrow of normal mice. When the stromal layer was established, the cultures were treated with mycophenolyic acid (MPA) using the protocol determined in our preliminary studies. Such treatments eliminated proliferating cells and stem cells while a functional stromal cell layer was left intact. Such stromal cell cultures were then charged with bone marrow cell (BMC) aliquots from which adherent cells were depleted by passaging through a G-10 column. Culture of such adherent cell-depleted cell fractions by themselves has shown that a stromal cell layer does not become established and LTBM cannot be initiated by these adherent cell-depleted fractions.

The cells used for recharging the stromal cell layer were also depleted of B cells with 14.8 antibody against the B-220 antigen in the presence of C'. At weekly intervals non-adherent cells were harvested from the cultures, the cultures fed, and the total number of cells, as well as the number of 14.8^+ or $s\mu^+$ and $c\mu^+$ cells, were enumerated.

The control for the experiment was seeding normal BMC (depleted of adherent cells and 14.8^+ cells) onto stromal cell cultures established from the BM of normal mice. The ability of stromal cells derived from tumor-bearing mice for supporting B cell production was tested by seeding normal BMC (depleted of adherent cells and 14.8^+ cells) onto stromal cell cultures obtained from the BM of CE maca tumor-bearing mice.

Figure 1 compares the effectiveness of stromal cells obtained from normal BM with those obtained from the marrow of CE maca tumor-bearing mice in initiating and maintaining B lymphopoiesis. When the stromal cell layers were recharged with normal BMC, themselves depleted of adherent cells. Our last progress report presented data for a number of duplicate experiments that corroborated the findings shown in Figure 1. The data

convincingly show that bone marrow stromal cell (BMSC) cultures established 14 and 21 days after tumor transplantation, when *in vivo* lymphopoiesis in the BM has been virtually eliminated, are capable of initiating and maintaining B lymphopoieses *in vitro* as effectively as stromal cell cultures established from the BM of normal mice.

Figure 1

Comparison of stromal cells obtained from normal and tumor-bearing mice to support B lymphopoiesis when recharged with normal bone marrow cells. The top left hand panel shows the 14.8⁺ profile of the normal bone marrow cells (depleted of adherent cells) used for recharging the stromal cell cultures. Six weeks after recharge, as indicated in the top right hand panel, these normal bone marrow cells placed on normal stromal cells maintained production of 14.8⁺ cells. The same is true if the cells are placed on the stroma of tumor-bearing mice as indicated in the lower panel.

Figure 2 gives evidence that explanting BM into the LTBM culture system eliminates the inhibitory effect that is exerted on the B lymphopoiesis *in vivo* in the BM of mice that have been transplanted with CE maca. Cell production was established in all cultures and the number of cells harvested over a six-week period exceeded many times the number of cells used for recharge. Although the inoculum was depleted of 14.8⁺ cells, 14.8⁺ were generated in the cultures as indicated in Figure 2.

It is important to emphasize that although our experiments give a conclusive answer to the question we posed in this Aim, our studies do not rule out that the inhibitory effects CE maca exerts *in vivo* on B lymphopoiesis is mediated via stromal cells. Should that be the case, however, it is clear that explanting stromal cells of tumor-bearing mice removes

them from under the modulatory influence that is exercised *in vivo*; they support the generation of B lymphocytes in LTBM as well as BMSC obtained from normal mice.

Although the data shown in Figure 2 suggest that BMC obtained from tumor bearing mice contain 14.8⁻ B lymphocyte progenitors capable of initiating B lymphopoiesis *in vitro*, we have performed a number of experiments that explored this question.

We used S-17 cells, a cloned BMC line developed by Dr. Kenneth Dorshkind. These cells have been shown to constitutively secrete growth factors and differentiation factors upon which B lymphopoiesis in culture depends. We have gained considerable experience with this cell line as well as other BMSC lines in the course of our studies. Figure 3 shows a representative experiment in which BMC of normal mice and tumor-bearing mice depleted of both adherent and 14.8⁺ cells were plated onto S-17 cells under W/W conditions.

These studies confirmed the conclusions we reached from the data presented in Figure 2. In LTBM, 14.8⁺ and μ ⁺ cells are generated at the same rate and in the same numbers when S-17 stromal cell layers are recharged with normal or tumor-bearing mice BM that has been depleted of B cell precursors and stromal cells. Although other assays can demonstrate a defect in more mature B cell progenitors in tumor-bearing mice (described in the last year's Progress Report), the cell that initiates and maintains B lymphopoiesis is present and unaffected in the BM of mice in which B lymphopoiesis is profoundly suppressed.

Although a number of experiments would be designed to further pursue the role of stromal cells in mediating facilitory or inhibitory effects on B lymphopoiesis, we have chosen to focus on studies that explore the modulatory effects of factors that are secreted by CE maca on *in vitro* lymphopoiesis as defined in our original Aim 2.

Figure 2

Generation of 14.8⁺ cells in W/W long term bone marrow cultures initiated with stromal cells from the bone marrow of CE maca-bearing mice. The top left hand panel shows the 14.8⁺ profile of the bone marrow cell population (depleted of adherent cells) obtained from a 17-day tumor-bearing animal used for recharging the stromal cell cultures. Note that 14.8⁺ cells are absent. The right hand top panel shows the generation of 14.8⁺ cells suggesting that stromal cells derived from tumor-bearing animals can support B lymphopoiesis and also that precursors of B cells capable of initiating B lymphopoiesis in such cultures are present in the bone marrow. The lower panel shows on the left, the initial population of bone marrow cells obtained from normal mice that was placed on stromal cell layers of tumor-bearing mice. Precursors present in the normal bone marrow generated 14.8⁺ cells over a 4-week culture period as shown in the right hand lower panel.

S-17: 24 WELL PLATE exp. 061390
rech. with 1×10^5 adh. and 14.8 depleted BM cells

Figure 3

The generation of 14.8⁺ cells on S17 stromal cell line. Bone marrow from mice transplanted 21 days previously with CE maca (TBM) and bone marrow from normal mice (NBM) were depleted of adherent cells and 14.8⁺ cells. In this experiment, 1×10^5 cells were plated onto S17 stromal cell cultures and the generation of 14.8⁺ cells assessed at 2, 3, and 4 weeks. The experiment indicates that cells capable of initiating B lymphopoiesis supported by S17 stromal cells are present in both tumor-bearing and normal bone marrow.

B. ACCOMPLISHMENTS IN AIM 2: MODULATORY EFFECTS OF FACTORS SECRETED BY CE MACA IN IN VITRO B LYMPHOPOIESIS

We have answered two questions that were posed under this Aim in our original proposal:

1. Does CE maca conditioned medium (CM) interfere with in vitro lymphopoiesis?

In the design of experiments to answer this question we made use of CM of several tumor cell types that we maintain as cell lines or clones. These cells and their products are used for comparing various *in vivo* and *in vitro* biological and experimental parameters pertinent to the regulation of lymphopoiesis. The relevant lines or clones are:

- CE 816 - a parent cell line of CE maca that on transplantation reproduces all biological effects of the *in vivo* passaged tumor
- CESJ - a clone derived from CE 816 extensively characterized (56); constitutively expresses G-CSF and M-CSF but not GM-CSF or other growth factors (Appendix 1; 55)
- BC66 - a clone isolated from Balb/C 66 mammary adenocarcinoma (91); lacking the effects of CE maca; secretes M-CSF but no other growth factors (Appendix 1; 55).
- NIH-3T3 - non-factor producing cell clone
- C127I-MG2 - constitutively produces murine G-CSF (Gift of Dr. Nagata, Osaka Bioscience Institute, Japan).
- C127I - cell line lacking murine G-CSF gene

Two representative experiments are presented in Figure 4 and Figure 5. Following considerable preliminary work for determining the culture media most suitable for growing the tumor cell clones or lines, conditions were developed to use serum free media. In all experiments CM of different concentrations from the various tumor clones was added to W/W type cultures in which normal BMC depleted of adherent cells and 14.8⁺ cells were plated onto S-17 stromal cells. The cultures were maintained up to four weeks and the readout was the generation of 14.8⁺ cells. Figure 4 compares CM from the parent cell line, CE 816, with known *in vivo* effects of suppressed lymphopoiesis with CM from the BC66 clone that lacks *in vivo* lymphopoietic effects. It is evident from Figure 4 that the *in vivo* effect of CE 186 is reproduced *in vitro*. Although fewer than 14.8⁺ cells were generated in the presence of BC66 CM than in control cultures, the most dramatic effects were observed with the cell line that inhibits B lymphopoiesis *in vivo*. In order to distinguish between the possible effects of G-CFS and M-CSF, both produced by CE 816 and its CESJ (SJ-3) clone, and M-CSF, the only growth factor produced by BC66, a series of experiments was performed using CM of the clones grown in serum-free media. The results are shown in Figure 5. The difference noted in Figure 4 is born out also under the experimental

conditions of Figure 5. CM known to contain G-CSF has a pronounced inhibitory effect on the generation of 14.8^+ cells in W/W cultures compared to CM containing only M-CSF. This difference is perceptible at one week in the cultures and persists through the four-week culture period.

14.8 and adherent cell depleted NBM seeded
on S17 stromal cell line

Figure 4

Normal bone marrow depleted of adherent cells and 14.8^+ cells then seeded onto S17 stroma and grown under Whitlock-Witte conditions for 4 weeks. Cultures were supplemented with CE816 or BC66 tumor cell conditioned media in RPMI containing 5% FCS.

Figure 5

Normal bone marrow depleted of adherent cells and 14.8⁺ cells then seeded onto S17 stroma and grown under Whitlock-Witte conditions for 4 weeks. Cultures were supplemented with SF-3 or BC66 tumor cell conditioned HL-1 media (serum free).

The data have raised the intriguing possibility that G-CSF responsible for the marked neutrophil hyperplasia that characterizes the *in vivo* model might also be responsible for the down-regulation of lymphopoiesis. Critical experiments to test this possibility had to depend on the availability of recombinant G-CSF, as CM from tumor cell lines and clones may well contain as yet unidentified factors. Data presented in Appendix 6 give evidence that human recombinant G-CSF is not very effective in inducing neutrophil hyperplasia and bone resorption in comparison with the CE tumor and its clones. We were fortunate that a collaboration could be established with Dr. Nagata of Osaka Bioscience Institute, Japan, who has inserted the murine G-CSF gene into a murine cell line C127I-MG2. He provided the cell line for our experiments that enabled us to compare CM from the cell line constitutively producing recombinant murine G-CSF with the other CMs as well as with the CM from the control cell line (C127I), which lack the murine G-CSF gene.

Figure 6 shows another set of experiments in which the production of 14.8^+ cells is compared in the same type of cultures described above in the presence of a number of CM. The data in Figure 6 were obtained at two weeks of culture. CM from C127I cells gave comparable results to CM of NIH-3T3 cells (not included in the figure). The data reveal a striking correspondence between the effects of CE 816, SJ 4, and C127I-MG2 CM, all of which contain G-CSF. These results are suggestive that G-CSF may be the molecule responsible for the suppression of B lymphopoiesis *in vitro* and as well as *in vivo*.

Experiments are now in progress to confirm these data and exclude the possibility that CM of CE 816 and its clones contains another molecule besides G-CSF that might also suppress lymphopoiesis. The experiments are being performed with rabbit polyclonal antibody raised against murine G-CSF, also obtained from Dr. Nagata.

Appendix 6 presents data on the *in vivo* effects of recombinant G-CSF, GM-CSF and human erythropoietin on neutrophil hyperplasia, activation of osteoclasts, and bone resorption. G-CSF mimics most closely the multiple effects of the *in vivo* transplanted CE maca. The effects of these growth factors on other hemopoietic lineages, particularly lymphocytes, are under investigation. In view of the clinical use of recombinant growth factors, the forthcoming data will be of considerable interest.

Thus the experiments we have refined in this Aim give a clear cut answer to the question we asked: **CE maca and its clones secrete factors that profoundly suppress in vitro B lymphopoiesis; these suppressive effects can be ascribed to G-CSF, constitutively expressed by CE maca and its clones.**

C. SUMMARY OF FINDINGS IN AIM 1 AND AIM 2

We have successfully exploited the W/W type LTBMCM for investigating the mechanisms that are responsible for suppressed lymphopoiesis observed in animals that have been transplanted with CE maca. This mechanism seems to be mediated *in vivo* and *in vitro* by G-CSF. Since both stromal cells and B cell progenitors from the BM of CE maca-bearing mice behave normally under *in vitro* conditions, the most likely explanation of a suppressed lymphopoiesis is the high level of G-CSF secretion by the transplanted tumor. CM derived from the tumor cell line and its clones contain G-CSF, and these media and

not those of other non-G-CSF secreting cell clones suppress B lymphocytopoieses *in vitro*. These effects are comparable to the addition of recombinant murine G-CSF to the culture systems.

CONDITIONED MEDIA EXPS. 031291-032191

Figure 6

Adherent cell and 14.8⁺ cell depleted normal bone marrow seeded on S17 stromal cell line then cultured under Whitlock-Witte conditions.

D. SIGNIFICANCE OF FINDINGS IN AIM 1 AND 2

It may also be concluded from Figures 5 and 6 that M-CSF does not influence lymphocyte production to the same extent as G-CSF. CM of the BC66 tumor did not affect the production of 14.8⁺ cells. These results are consistent with our previous studies (Appendix 1; 55) showing the lack of granulocytic or hypercalcemic effects on the transplanted BC66 tumor cell line in *in vivo* experiments.

Our experiments have important implications for the role of G-CSF in modulating lymphopoiesis. Our consistent observations *in vivo* and *in vitro* give particular significance to our work. The object of the present proposal was to identify an experimental system that can reliably assay factors that down-regulate the production of different classes of lymphocytes. We feel that our work establishes the usefulness of LTBMG grown with stabilized stromal cell lines for analyzing biological response modifiers that can affect B lymphopoiesis. Our studies have also raised promising possibilities for investigating the mechanisms that explain how a growth factor that is stimulatory for one cell lineage, (neutrophils at the present level), may inhibit the differentiation of another lineage, (B lymphocytes at the prior level). Our findings provide the necessary background and rationale for exploring these mechanisms through an independently funded project. We are now in the process of preparing a manuscript that will report our work on the effect of various growth factors, particularly G-CSF on B lymphopoiesis in the BM.

E. OTHER FINDINGS IN THE CE MACA EXPERIMENTAL MODEL

In addition to the accomplishments that directly relate to Aims 1 and 2, a number of important findings have emerged from Dr. Lee's work using the CE maca model. These findings are particularly pertinent in view of the interrelationship that has emerged from our studies between the regulation of different hemopoietic lineages. Two of these incidental findings deserve emphasis:

1. **Transformed bone marrow stromal cells in mice bearing CE maca.** On two occasions we observed extremely vigorous growth of stromal cells in cultures when stromal cells were obtained from CE tumor-bearing mice. In these culture flasks, production of hemopoietic cells or lymphocytes was soon overwhelmed by the vigorous growth of stromal cells. These rapidly growing stromal cells were cloned and were named KLF1 and KLF2 clones. We have never observed metastases from the transplanted CE mammary tumor, but as yet we do not have absolute proof whether or not KLF cells are derived from the transplanted tumor or from bone marrow stromal cells. We tested the tumorigenicity of the cloned cells by inoculating KLF1 and KLF2 subcutaneously into normal (BALB/CS CE) F1 mice. Several weeks later both cell lines formed localized tumors. The most interesting finding was that mice bearing either the KLF1 or KLF2 clone developed the same syndrome as mice transplanted with CE maca tumor: marked neutrophil hyperplasia and hypercalcemia. Figure 7 shows furthermore that the number of 14.8^+ in the spleen of the tumor bearing mice was somewhat reduced in comparison with normal mice; more importantly, 14.8^+ cells were absent in the BM of mice bearing these clones.

These preliminary findings suggest that KLF cells are transformed cells and probably produce the same growth factors as CE maca. In addition to the possibility of these cells being derived from the transplanted tumors, we consider it likely that they arose by stromal cell transformation: MMTV virus insertion near the growth factor genes in stromal cells. Although these cell clones are preserved in our freezer, limitations in our funding do not permit us to explore their experimental potential.

2. While pursuing the CE maca animal model, Dr. Lee described a neutrophil colony-forming assay that detects CFU-N, a progenitor responsive to G-CSF but not M-CSF. Her interest in neutrophil hyperplasia has long been combined with the investigation of bone-bone marrow interactions, particularly through the hyperplasia of osteoclasts. She has developed an assay that detects osteoclast progenitor cells described in Appendices 3,4,5 (Appendix 3; 57). The clones of cells developing from osteoclast progenitor cells exhibit the characteristics of osteoclasts that are capable of resorbing bone. Equally important is her discovery of a colony-stimulating factor specific for osteoclasts (Appendix 3; 57). The factor is produced by the same CESJ clone of CE maca as we have been using for our studies of lymphopoiesis. Although the new osteoclast stimulator is not yet widely recognized, it may have important implications not only for bone metabolism but for the physiology of hemopoiesis and lymphopoiesis. This factor stimulates the proliferation and differentiation of osteoclast progenitors from hemopoietic stem cells. Dr. Lee has successfully isolated the factor and its detailed characterization is in progress. The availability of pure osteoclast stimulator will enable us to dissect the mechanisms involved in osteoclast activation and by expansion, an interesting phenomenon we observed in animals bearing CE maca (58). The relevance of this factor to lymphopoiesis is that lymphocytes are the earliest hemopoietic lineage identifiable in the hemopoietic tissue of CE maca-bearing animals that develops in the medullary cavity of bones of the peripheral skeleton usually occupied by fatty marrow. Osteoclast stimulator is produced by the CESJ tumor but not by the other tumor cell lines that we have analyzed in our studies.

F. ACCOMPLISHMENTS IN AIM 3: IN VITRO SYSTEMS FOR NK LYMPHOPOIESIS

The questions we originally asked in this Aim about NK lymphopoiesis paralleled those we posed for B lymphopoiesis in Aims 1 and 2. The rationale was based on our *in vivo* studies. We have shown (Appendix 7; 70) that within three days of subcutaneous transplantation of CE maca and Bc66, a marked elevation of NK activity occurs in the BM and spleen (SPL). This increase is followed by a decrease in both NK activity and the number of BM NK precursors (determined by limiting dilution assay) in the CE maca but not in the BC66 recipient. Resection of CE maca leads to recovery of NK cell numbers and activity. Because of this similarity in the modulation of NK and B cell lymphopoiesis, we originally proposed to investigate the role of stromal cells in NK lymphopoiesis along similar lines as in Aim 1 and 2 for B cells. The proposed experiments for investigating the modulation of NK lymphopoiesis were predicated on two requirements: 1) the outcome of the studies on B lymphopoiesis, and 2) on the availability of a reliable *in vitro* system in which, without the addition of exogenous factors, NK lymphopoiesis can be maintained by BMSC.

The experiments we described in Aim 1 and 2 revealed that defective stromal cell or B lymphocyte progenitor cell function could not be demonstrated in BM of CE maca tumor-bearing mice when the BM was explanted and tested in LTBM. The *in vivo* effects of CE maca on B lymphopoiesis were duplicated *in vitro* by CM that contained G-CSF.

Before embarking on the studies we originally proposed in Aim 3, we repeated the *in vivo*

Figure 7

Histograms generated from FACS analysis of 14.8⁺ mAb stained spleen cells from normal (a) and KLF2 tumor-bearing mice (b) and bone marrow cells from normal (c) and KLF2 tumor-bearing mice (d). X axis, log fluorescence intensity; Y axis, relative cell number.

transplantation experiments with *in vivo* passaged CE maca tumor cells and the SJ clone for which we have documented evidence with respect to G-CSF production. For comparison, we also transplanted mice with the Bc66 clone that produces M-CSF but not G-CSF (Appendix 8; 40). Assay of NK activity confirmed a biphasic response to CE maca recipients, but no modulation of NK activity was detected either with the SJ or BC66 clones. We interpret the data as suggestive of the insensitivity of NK lymphopoiesis to G-CSF or to the possible secretion of factors other than G-CSF and M-CSF by the uncloned tumor. The biphasic NK cell response in the recipients of the *in vivo* passaged tumor may,

moreover, be due to lymphokines produced by activated lymphocytes that were present in the tumor transplant. In order to test the modulatory effect of various CM on *in vitro* NK lymphopoiesis, as we have done for B lymphopoiesis, it was necessary to refine and stabilize the LTBM system that we have been developing.

The above experiments, as well as numerous reports, have pointed to the importance of lymphokines, particularly IL-2, in the *in vitro* generation of NK cells. It was important therefore that we define a LTBM system in which the role of IL-2 can be controlled. We have investigated the IL-2 response of BM NK cells at different maturational stages and have shown that the least differentiated NK-1.1⁺ precursors responded vigorously to IL-2 and generated high levels of cytotoxicity against YAC-1, whereas the smaller NK-1.1⁺ primary NK cells respond to IL-2 but did not generate as much lytic activity as did their precursors (Appendix 9; 69). Moreover, when CD8⁺ cells were present in BM, lymphokine activated killer (LAK) cell activity was also generated in response to IL-2.

Much of our effort in Aim 3 has been directed toward establishing a LTBM system in which the modulatory influences exerted by biological as well as environmental factors can be reliably assessed. **We have established a unique culture system that for the first time allows the production and differentiation of NK cells from their BM cell progenitors without requiring the addition of exogenous growth factors** (Appendix 10; 75). In agreement with previous reports, we found that myeloid (Dexter) LTBM did not produce lytic NK cells. However when we switched the cultures to lymphoid (W/W) conditions and reseeded ("recharged") the switched cultures with fresh BM cells, non-adherent cells with NK lytic activity and NK-1.1⁺ phenotype were produced within one to two weeks. To eliminate the possibility that LTBM were simply maintaining mature lytic NK and/or IL-2 producing T cells, the recharged bone marrow cell (BMC) population was treated with C' and MAbs to NK-1.1, CD8, and CD4. Such depleted BMC proliferated extensively in lymphoid type BM cultures and also generated cells in two weeks that lysed the NK target YAC-1 but not the LAK target P815. NK-1.1⁺ cells were detected by flow cytometry. When recultured in medium containing 50 U rIL-2 per ml, the non-adherent cells from the recharged long term cultures generated high levels of NK but not LAK lytic activity. These findings have been reported at several meetings (Appendix 11; 73, Appendix 12; 74, Appendix 13; 72) and have been submitted for publication (Appendix 10; 75).

The establishment of LTBM conditions that maintain NK cell precursors and NK cell production without the addition of exogenous growth factors will provide the opportunity for studying in detail the factors that regulate NK cell production. The system will also permit us to undertake the study of biological and environmental factors that perturb NK cell production as we have documented with the mammary adenocarcinoma system. Our studies investigating the effect of CE maca on B lymphopoiesis are one step ahead of those concerned with NK lymphopoiesis. Our plans now include assaying both NK and B cell production in cultures that will be set up to investigate the influence of CE maca CM in comparison with the effects of defined growth factors on NK lymphopoiesis.

G. ACCOMPLISHMENTS IN AIM 4: MODULATION OF T LYMPHOPOIESIS

Because our proposal could not be fully funded by DOE, we had to delete the experiments that were described in Aim 4. Nevertheless, parallel work supported by other grants has yielded information that pertain to Aim 4 and in many respects provide preliminary data for the studies we propose for the next project period. A brief summary of the relevant experiments is presented below.

1. Relationship between myelogenous and intrathymic T cell precursors.

Making use of the differential expression of the Thy-1.1 and Thy-1.2 allele in congenic mice have employed two experimental systems, parabiosis and partbody BM chimeras, to investigate the relative contribution of hematogenous and intrathymic T cell precursors to the maintenance of lymphopoiesis in the postnatal THY. These studies were initiated under DOE sponsorship and are now supported by an NIH grant (Appendices 14,15,16; 61,62,80).

The importance of these studies is that for the first time they point to the importance of intrathymic T cell precursors in maintaining postnatal thymocytopoeisis in the steady state without reliance on hematogenous T cell precursors derived from the BM. These findings are contrary to current concepts of T cell production that envisage a continuous stream of T cell precursors from the BM as a requirement for the maintenance of T cell production within the THY under normal circumstances. Our data document an extensive degree of self-maintenance of intrathymic T cell precursors and suggest that hematogenous precursors that do enter the THY are primarily called upon when cell production within the THY must be re-established following irradiation or grafting. These findings have important implications for the regulation of T cell production and the role of the BM in maintaining it.

During the current project period we have extended these studies in two directions: 1) to examine the phenotype of thymocyte derived from intrathymic and hematogenous precursors, and 2) to investigate the role of hematogenous and intrathymic T cell precursors in the regeneration from cortisone-induced thymic atrophy.

1. We have investigated the expression of different T cell maturation of antigens on host (H) and non-host (NH) type Thy-1⁺ cells by 3 color flow cytometric analysis. Prior to staining the cells with fluorochrome-tagged antibodies for T cell maturation antigens, we have eliminated either the H type or NH type Thy-1⁺ cells by anti-Thy-1.1 or anti-Thy-1.2 antibody plus C'. The data we have reported (Appendix 15; 62) show that both H and NH cells expressed the CD3 antigen, providing evidence for the fact that both type of cells contain T cells in equal proportions. Similarly, a CD4⁻CD8⁻ (double negative) cell fraction exists in both populations as does CD4⁺CD8⁺ (double positive) phenotype. Single positive CD4⁺ and CD8⁺ are likewise represented among H type and NH type cells. Thus, it may be concluded that although NH type cells are few compared to H type cells, there is evidence for a stream of T cell maturation (according to schemes that have been proposed) in both H and NH type cells. It is interesting to note the relative predominance of CD4⁺ cells among the NH type maturational stream.

2. Our earlier studies have shown that irradiating the THY permits the NH type cells to differentiate and contribute to thymocytopoiesis. During the past year, we investigated whether hematogenous cells contribute in a similar manner to the recovery from corticosteroid-induced thymic atrophy. Our findings were reported at a meeting (Appendix 16; 80). Although thymic cellularity was reduced by cortisone treatment from 160×10^6 to 2.5×10^6 cells, a depopulation equivalent to that caused by 1,000 rad, our data conclusively show that the recovery from such profound atrophy is effected almost exclusively by intrathymic cells. Both in the parabionts and partbody BM chimeras, the low proportion of NH type cells remained constant, both at 5 and 15 weeks after administering corticosteroid. The data are, in fact, indistinguishable from the animals that have received saline instead of corticosteroid. The studies assign a much greater importance to intrathymic T cell progenitors in the maintenance of T cell production in the THY than has been hitherto recognized. These findings have an important bearing on the development of experimental model systems that will attempt to duplicate thymocytopoiesis *in vitro*.

2. Lineages of T lymphocytes in the developing thymus

Much of the work of this laboratory sponsored over the years by DOE has been concerned with defining the developmental lineages of lymphocytes and hemopoietic cells (50,68,71,78,79,81). Parallel with investigating the role of hematogenous and intrathymic T cell precursors, Dr. Farr's program has been concerned with defining maturational stages in the murine THY. Much of his work received initial support from DOE.

Using an antibody that reacts with the β chain of the T cell receptor, Dr. Farr has been able to identify, with electronmicroscopic histochemical techniques, thymocytes expressing this receptor at different maturational stages *in situ* with the THY. Some cortical cells exhibited only cytoplasmic staining whereas others in the cortex showed low levels of surface expression as well, almost invariably in regions of the cell membrane in contact with epithelial cell processes. In the medulla, thymocytes expressed high levels of uniform surface labelling with the antibody (27). He and his collaborators have shown that as cells mature they rearrange their receptor genes in an ordered fashion and later express surface protein. In the cortex these cell membrane receptors collect at the surface of contact between the thymocyte bearing them and the epithelial cell where selection for self-MHC recognition and/or tolerance induction may occur. There is evidence that cortical thymocytes may bear low levels of receptors per cell because of this capping and patching property. On the other hand, mature medullary thymocyte and peripheral T cells express high levels of receptor per cell (60).

Using the same ultrastructural immunohistochemical approach, Dr. Farr has analyzed the process of clonal deletion of self-reactive thymocytes within the thymic environment. He has demonstrated that in the case of clonal deletion of thymocytes expressing V β 6 + TCR in response to M1s antigens, the deletion process occurs within the cortical thymic environment. This observation provides an important intellectual framework for considering the cellular interactions mediating this process. (Appendix 19)

Using histochemical and immunocytochemical techniques, Dr. Farr has found a lectin with nominal specificity for alpha-linked N-acetyl-D-galactosamine. *Dolichos bifloris* agglutinin (DBA) was found to preferentially label thymocyte with an immature phenotype from fetal, newborn, and adult mice (26). Through days 14 to 16 of gestation, virtually all thymocytes, bound DBA. This was followed by a dramatic reduction of DBA labelling during the last 4 days of gestation, reaching adult levels of about 2-4% of total thymocytes. At later stages of fetal development, DBA⁺ cells were confined to the subcapsular area of the THY. Affinity purification of thymocyte cell membrane components with insolubilized DBA indicated that virtually all of the lectin binding was mediated by a 120-kDA glycoprotein of the thymocytes cell membrane. The molecule may be useful for positively selecting CD4⁻ CD8⁻ (double negative) thymocyte and possibly other immature populations within the T cell lineage.

DBA labeling was correlated with the expression of the T cell receptor chain in the developing THY at both the light and electron microscopic level using monoclonal antibody (24), which recognizes determinants encoded by the V gene family of the T cell receptor β chain. Cytoplasmic and cell membrane expression of this molecule of the receptor were correlated with maturational stages and topographical location of maturing cell clusters in the embryonic THY. The reciprocal compartmentalization of DBA⁺ and β chain receptor⁺ cells suggested a lineage relationship between subpopulations of thymocyte.

Dr. Farr has also utilized an antibody detecting the d chain of the $\gamma\delta$ TCR to characterize the spatial relationship between these cells and stromal element of the thymus during development. He demonstrated that there was a striking localization of $\gamma\delta$ cells to the vicinity of medullary epithelial cells of the thymus (32). He has also contributed considerable documentation concerning the extrathymic distribution of T cells expressing $\gamma\delta$ TCR and their striking affinity for epithelia in other tissues. Dr. Farr's expertise in morphologic and electron microscopic immunohistochemistry has permitted him to make several unique and important contributions to our understanding of the thymic environment.

3. Characterization and in vitro propagation of thymic epithelial cells

Dr. Farr's program is one of the few that has successfully isolated, characterized, and maintained cell lines of murine thymic epithelium. Several cloned thymic epithelial cell lines have now been generated (28,29). Characterization of the various species of keratin and other cell surface molecules expressed by these epithelial cells made it possible to draw correlations between the phenotypes of epithelial cells in subcapsular, cortical and medullary regions of the THY. (Appendix 17; 30, Appendix 18; 84,). A large panel of monoclonal antibodies has been generated to identify different populations of thymic epithelial cells. These reagents are of the greatest importance in characterizing epithelial cells and cell interactions with differentiating thymocyte. We have obtained evidence that selective clones of thymic stromal cells we believe isolated induce growth factors for hemopoietic cell production and are capable of maintaining long term lymphopoiesis *in vitro*. Our proposed work is based on these studies. They are presented in the section **RESEARCH PROPOSAL** as Preliminary Results and as the rationale for our proposed experiments.

H. SUMMARY OF ACCOMPLISHMENTS

1. Employing co-culture of stromal cells and B lymphocyte progenitors explanted from the BM of CE maca tumor-bearing or normal mice, we have shown in reciprocal experiments with LTBMCM that no defect could be demonstrated either in stromal cell or progenitor cell function that could account for the profound suppression of B lymphopoiesis we have discovered in the BM of tumor-bearing mice. We have also confirmed the normal capacity of B cell progenitors obtained from tumor-bearing mice to initiate and maintain B lymphopoiesis on stromal cell lines that have been selected for the ability to preferentially support B lymphopoiesis.

2. Cell clones derived from CE maca constitutively express G-CSF but not M-CSF, whereas a BC66 adenocarcinoma cell line expresses M-CSF but not G-CSF. CM of CE maca and its clones added to LTBMCM profoundly suppress B lymphopoiesis, whereas B lymphopoiesis is unaffected by CM of the BC66 clone.

3. Recombinant G-CSF added to LTBMCM duplicate the *in vitro* and *in vivo* effects of CE maca and its secreted products. G-CSF is a highly effective negative regulator of B lymphopoiesis.

4. We have developed a new cell line (apparently derived from BMSC), which on transplantation causes neutrophilic hyperplasia and suppression of B lymphopoiesis. CM of the clone also suppressed B lymphopoiesis in LTBMCM.

5. Using lytic activity against NK-sensitive targets and the expression of NK antigens, we have defined maturational stages of NK lymphocytes in the bone marrow.

6. Early NK progenitors are stimulated to a greater extent *in vitro* by IL-2 than more mature NK precursors. Exposure of early NK precursors to IL-2 results in both higher levels of NK lytic activity and an increase in the number of the lytic units.

7. We have developed a long term bone marrow culture system in which NK cell production is initiated by the earliest (phenotypically and functionally unidentifiable) NK progenitors and is maintained in the absence of exogenous growth factors, including those produced by incidental T cells.

8. We have isolated a new growth factor that is constitutively expressed by clones of CE maca and specifically promotes the differentiation and proliferation of osteoclasts.

9. We have developed an *in vitro* clonal assay that identifies an osteoclast progenitor cell that gives rise to colonies of cells capable of resorbing bone.

10. Osteoclast stimulator, when infused into animals, causes hypercalcemia, bone resorption, and hyperplasia of osteoclasts. *In vitro*, the factor supports the development of osteolytic colonies from.

11. We have demonstrated that postnatal lymphopoiesis in the thymus relies extensively on an intrathymic pool of T cell precursors that are independent of the minor population of hematogenous (bone marrow-derived) progenitors.

12. All subclasses of thymocyte phenotypes are generated by both intrathymic and hematogenous T cell precursors.

13. Following selective irradiation of the thymus, intrathymic, and hematogenous precursors participate equally in thymic regeneration, whereas after cortisone-induced severe thymic atrophy, recovery is effected almost exclusively by the intrathymic T cell progenitors.

14. We have generated reagents that recognize subpopulations of thymic stromal cells restricted to the subcapsular, cortical, and medullary areas of the thymus and have established a number of cell clones of thymic epithelial cells that express the characteristics of stromal cells from these different regions.

15. We have correlated the expression of the β,γ and δ chains of the T cell receptor with maturational stages of T cells and have identified the reagents that recognize the earliest stages of T cell differentiation.

16. We have shown that clones of thymic stromal cells secrete growth factors for hemopoietic cell lineages and are capable of supporting lymphopoiesis in long term bone marrow cultures.

END

**DATE
FILMED**

2/11/92

I