DELIVER BY POST

JAN 11

"Approved for public release; distribution is unlimited."

**Three-Year Summary Report of
Biological Monitoring at the Southwest
Ocean Dredged-Material Disposal Site
and Additional Locations Off Grays
Harbor, Washington, 1990 - 1992**

L. D. Antrim
D. K. Shreffler

W. H. Pearson
V. I. Cullinan

Battelle/Marine Sciences Laboratory
Sequim, Washington

December 1992

Prepared for the
U.S. Army Corps of Engineers - Seattle District
under a Related Services Agreement
with the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes **any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.** Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401. FTS 626-8401.

Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

THREE-YEAR SUMMARY REPORT OF BIOLOGICAL
MONITORING AT THE SOUTHWEST OCEAN DREDGED-
MATERIAL DISPOSAL SITE AND
ADDITIONAL LOCATIONS OFF GRAYS HARBOR,
WASHINGTON, 1990 - 1992

L. D. Antrim
D. K. Shreffler
W. H. Pearson
V. I. Cullinan

Battelle/Marine Sciences Laboratory
Sequim, Washington 98382

December 1992

Prepared for
U.S. Army Corps of Engineers - Seattle District
under a Related Services Agreement
with the U.S. Department of Energy
Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Richland, Washington

MASTER

NO DUPLICATION OF THIS DOCUMENT IS UNLIMITED *eb*

EXECUTIVE SUMMARY

The Grays Harbor Navigation Improvement Project was initiated to improve navigation by widening and deepening the federal channel at Grays Harbor. Dredged-material disposal sites were selected after an extensive review process that included inter-agency agreements, biological surveys, other laboratory and field studies, and preparation of environmental impact statements (USACE 1982; 1989). The Southwest Site, located approximately 4 miles outside of Grays Harbor, was designated to receive materials dredged during annual maintenance dredging as well as the initial construction phase of the project.

The Southwest Site was located, and the disposal operations designed, primarily to avoid impacts to Dungeness crab. The Final Environmental Impact Statement Supplement for the project (USACE 1989) incorporated a Site Monitoring Plan in which a tiered approach to disposal site monitoring was recommended (Pearson 1987). Under Tier 1 of the Site Monitoring Plan, Dungeness crab densities are monitored to confirm that large aggregations of newly settled Dungeness crab have not moved onto the Southwest Site. Tier 2 entails an increased sampling effort to determine whether a change in disposal operations is needed.

Four epibenthic surveys using beam trawls were conducted in 1990, 1991, and 1992 at the Southwest Site and North Reference area, where high crab concentrations were found in the spring of 1985. Survey results during these three years prompted no Tier 2 activities. Juvenile Dungeness crab density was higher, in general, at the North Reference area than at the Southwest Site. However, the difference between the two sites in juvenile crab density was not statistically significant ($p < 0.05$) for any survey.

Epibenthic surveys were also conducted at two nearshore sites where construction of sediment berms has been proposed. This work is summarized in an appendix to this report.

ACKNOWLEDGMENTS

This research was funded by the U.S. Army Corps of Engineers (USACE), Seattle District. As the technical representative, Bert Brun (USACE) was supportive and cooperative during all phases of the work. We would like also to thank Alex Sumeri, Lauran Cole, and Eric Nelson (USACE) for their assistance during the surveys and process of report preparation. Dr. David Armstrong, from the University of Washington, provided background and unpublished information. Dr. Ray Bienert, Tom Parkwell, and Bruce Higgins, from Battelle, and Christina Lobo, from Western Washington University, were instrumental in successful completion of the field work. Laura Gully assisted with text processing.

CONTENTS

EXECUTIVE SUMMARY	iii
ACKNOWLEDGMENTS	v
1.0 INTRODUCTION	1
1.1 BACKGROUND	1
1.2 SCOPE AND STRUCTURE OF REPORT	6
2.0 SAMPLING METHODS	7
2.1 TIMING OF SURVEYS	7
2.2 VESSEL AND POSITIONING	7
2.3 EPIBENTHIC TRAWLS	8
2.4 CATCH PROCESSING	9
2.4.1 Dungeness Crab	9
2.4.2 Other Catch Contents	15
2.5 HYDROGRAPHIC PROFILES	15
2.6 VIDEO TOWS	16
2.7 DATA COMPILATION AND STATISTICAL ANALYSES	16
3.0 RESULTS	19
3.1 YOY DUNGENESS CRAB	19
3.1.1 Comparison Between Sites	19
3.1.2 High Density Area	21
3.1.3 Distribution With Depth	21
3.1.4 Annual Variation	22
3.2 1+ AGE-CLASS DUNGENESS CRAB	22
3.3 ADULT DUNGENESS CRAB	22
3.3.1 Density	22
3.3.2 Other Characteristics	23

3.4	NON-CRAB CATCH	23
3.5	HYDROGRAPHIC PROFILES	26
4.0	DISCUSSION	27
4.1	EVALUATION OF SITE MONITORING PLAN	27
4.1.1	YOY Crab Density	27
4.1.2	Choice of Sampling Gear	28
4.2	SUPPLEMENTAL FINDINGS ON DUNGENESS CRAB ECOLOGY	29
4.2.1	General Crab Distribution	29
4.2.2	Association of Juvenile Crab with Debris	29
4.2.3	Effects of El Niño	30
4.2.4	1 ⁺ and Adult Dungeness Crab	30
5.0	CONCLUSIONS	33
6.0	REFERENCES	35
APPENDIX A	1992 PRELIMINARY DATA REPORTS	A.1
APPENDIX B	EPIBENTHIC SURVEYS IN 1990 - 1992 AT TWO PROPOSED BERM SITES NEAR GRAYS HARBOR, WASHINGTON	B.1

FIGURES

1.1	Location of Grays Harbor and the Study Area	2
1.2	The Southwest Site and North Reference Area off Grays Harbor	3
2.1	Location of Individual Trawls at the Southwest Site in June 1985	10
2.2	Location of Individual Trawls at the Southwest Site in June 1990	11
2.3	Location of Individual Trawls at the Southwest Site in June 1991	12
2.4	Location of Individual Trawls at the Southwest Site in April 1992	13
2.5	Location of Individual Trawls at the Southwest Site in June 1992	14

TABLES

2.1	Number of Beam Trawls Completed at the Southwest Site and North Reference Area from Dungeness Crab Surveys by MSL Between 1985 and 1992	9
3.1	Summary of Dungeness Crab Densities (number/ha) from Beam Trawl Surveys at Southwest Site and North Reference Area, June 1985 - June 1992	20
3.2	YOY Crab Density/Ha With Depth at Southwest Site and North Reference Area, June 1985 - June 1992	21
3.3	Summary of Catch Data for Adult Dungeness Crab from Trawls at the Southwest Site, June 1990 - June 1992	24
3.4	Summary of Catch Data for Adult Dungeness Crab from Trawls at the North Reference Area, June 1990 - June 1992	25
3.5	Summary of Hydrographic Profiles From Southwest Site and North Reference Area, June 1985 - June 1992	26

1.0 INTRODUCTION

Grays Harbor is a shallow embayment on the Washington Coast where a federal, deep-draft navigation channel links the Hoquiam/Aberdeen area and Chehalis River to the Pacific Ocean (Figure 1.1). A significant contribution to the local economy has been the export of wood products from Hoquiam and Aberdeen. With the increasing size of deep-draft vessels used to transport wood products for the worldwide market, access to loading piers in Hoquiam/Aberdeen became limited to partially loaded or shallower-draft vessels. Consequently, the Grays Harbor Navigation Improvement Project (USACE 1982) was developed to improve access by widening and deepening the navigation channel at Grays Harbor. The navigation channel has required incremental maintenance dredging for enlargement in 1990, in addition to regular annual dredging to remove accumulated sediments from the channel.

1.1 BACKGROUND

For the Grays Harbor Navigation Improvement Project, the U.S. Army Corps of Engineers, Seattle District (USACE), dredged over 6 million cubic yards (cy) of sediments from the Grays Harbor area for the initial improvements alone (Alex Sumeri, USACE, personal communication). Disposal sites to receive dredged materials from both construction and annual maintenance dredging were selected after an extensive review process that included inter-agency agreements, biological surveys, other laboratory and field studies, and preparation of Final Environmental Impact Statement and Supplement (FEIS and FEISS) (USACE 1982; 1989). One ocean disposal site, the Southwest Site, (Figure 1.2) was located near the Southwest Navigation Lane. During the construction phase of the project, this site received approximately 1.2 million cy of uncontaminated sandy materials dredged from the Outer Bar Reach (Figure 1.2). Subsequent use of the Southwest Site has been for disposal of approximately 0.5 million cy of materials from annual dredging of the Outer Bar Reach.

Prior to designation and use of the Southwest Site, biological surveys determined that the dominant infauna near the Southwest Navigation Lane was a tube-building polychaete Owenia fusiformis and the dominant epifauna was

FIGURE 1.1. Location of Grays Harbor and the Study Area

FIGURE 1.2. The Southwest Site and North Reference Area off Grays Harbor (depth contours are in feet)

Dungeness crab, Cancer magister (Pearson et al. 1987). One survey, in June 1985, found exceptionally high concentrations of young-of-the-year (YOY) Dungeness crab, as well as smaller numbers of adult crabs, especially between

the 30.5- and 36.6-m (100- and 120-ft) contours north of the Southwest Navigation Lane (Pearson et al. 1987). Based on these findings, the disposal site (i.e., Southwest Site) was located farther south than originally planned, to avoid the area of high crab concentrations.

Nearshore areas off Grays Harbor and the estuary itself are important habitat at various stages in the life history of Dungeness crab (Armstrong et al. 1989). Adult females breed with males, carry egg masses, and hatch larvae along the open coast. After hatching, pelagic larval stages are found in coastal waters between December and April. In most years, megalopal larvae, the final larval stage, move nearshore toward shallow waters by April and May. In the Grays Harbor area, settlement to the benthic environment typically occurs in nearshore and estuarine areas between April and June. The abundance of YOY crab in nearshore areas can vary dramatically (i.e., by a factor of 10 or more) from year to year. Settled megalopal larvae rapidly undergo metamorphosis to YOY, or 0⁺-age-class, crab. Crab from the 0⁺ and 1⁺ age classes are considered juveniles, while 2⁺-age-class and older crab are called adults. In the Grays Harbor area, many YOY Dungeness crab move into the estuary for their first summer and winter of life. Estuarine residence results in faster growth for the juvenile crab because of warmer waters, a more protected benthic environment, and greater availability of food. By the end of their second year (1⁺ and older crab), most crab move outside the estuary for the duration of their adult life.

The Southwest Site was located, and the disposal operations designed, to avoid impact to commercially important biological resources in the area (USACE 1989). The site is centered near the Southwest Navigational Lane Buoy designated GH, which is located approximately 4 miles south-southwest of the mouth of Grays Harbor (Figure 1.2). The Southwest Site has the shape of a parallelogram covering approximately 1.7 square miles (1.3 square nautical miles) in area and lies between the 27.4 m (90 ft) and 36.6 m (120 ft) depth contours.

The ocean surveys and inter-agency scoping meetings identified razor clams and Dungeness crab as resources of concern in the area. Juvenile Dungeness crab were found to be more concentrated to the north of the site (Pearson et al. 1987). Razor clam populations were found primarily at depths

less than 27.4 m (90 ft) mean lower low water (MLLW) (USACE 1989). The peak in Dungeness crab larval settlement occurs in nearshore areas during May and June (Armstrong et al. 1989). Whereas weather and sea conditions generally limit dredging operations to the summer months (May through September), dredging operations were scheduled to begin by late June or July. Thus, dredging operations started after most larval settlement was expected to have occurred, to minimize potential impacts to crab before, during, and after settlement.

To confirm that impacts to YOY Dungeness crab were minimized, the FEISS for the navigation improvement project incorporated a Site Monitoring Plan developed by Battelle/Marine Sciences Laboratory^(a) (MSL) (Pearson 1987). The general objective of the Site Monitoring Plan was to indicate the studies necessary for determining whether changes in disposal operations were needed. More specifically, monitoring efforts were to assure concerned parties that the choice of the Southwest Site remained appropriate, and that no new developments occurred to necessitate a change in disposal operations or site location. The site monitoring plan recommended a tiered approach to disposal site monitoring. A tiered approach uses simpler (and relatively cost effective) methods for initial monitoring efforts, or lower tiers, while more complex monitoring techniques occupy higher tiers (Zeller and Wastler 1986).

Under Tier 1 in the Site Monitoring Plan, Dungeness crab densities are monitored before initiation of dredging activities to confirm that the disposal site does not contain an exceptionally high density of YOY Dungeness crab. Tier 2 entails an increased sampling effort to determine whether a change in disposal operations is needed. The trigger for moving to Tier 2 of the evaluation is a YOY crab density within the disposal site that is 100 times higher than the density in the reference area to the north. The trigger was based on the fact that high crab concentrations were found in the spring of 1985 at North Reference area.

(a) The Marine Sciences Laboratory is part of the Pacific Northwest Laboratory, which is operated for the U.S. Department of Energy by Battelle Memorial Institute.

1.2 SCOPE AND STRUCTURE OF REPORT

This report summarizes Tier 1 biological monitoring of the Southwest Site in the years 1990, 1991, and 1992. While the Site Monitoring Plan addressed both physical (i.e., mound height) and biological (i.e., Dungeness crab distribution) characteristics of the Southwest Site, this report summarizes the results of MSL's biological monitoring efforts, which were limited to epibenthic surveys with a beam trawl. Section 2.0 describes field sampling methods. Section 3.0 reviews the results of monitoring efforts from four surveys at the Southwest Site and North Reference area conducted by MSL between 1990 and 1992. In addition, these results are compared with data collected in June 1985 by Pearson et al. (1987). The 1985 survey data were included to provide a broader data base, incorporating findings from the period before initiation of disposal operations at the Southwest Site. Because the two other surveys conducted in 1984 and 1985 by Pearson et al. (1987) occurred in fall months, the results are not directly comparable to the spring surveys. Issues related to the biological surveys conducted under the Site Monitoring Plan are discussed in Section 4.0. Section 5.0 summarizes results of Tier 1 monitoring efforts for crab density.

While surveys in 1990 and 1991 were summarized in final reports to USACE (Higgins and Pearson 1991; Antrim et al. 1992, respectively), two surveys in 1992 were described only in preliminary letter reports to USACE. These letter reports are included here in Appendix A. The results of field sampling at potential berm sites in nearshore areas are discussed in Appendix B. In 1990, work in the nearshore area was limited to two trawls. As the prospects for berm construction advanced in 1991 and 1992, field efforts at the proposed nearshore berm sites were increased to include a greater number of trawls, characterization of surface sediments, and assessment of juvenile razor clam and commercially important fish resources.

2.0 SAMPLING METHODS

As part of Tier 1 monitoring under the Site Monitoring Plan, MSL conducted four epibenthic biological surveys at the Southwest Site and North Reference area offshore from Grays Harbor. Surveys were completed in June of 1990, 1991, and 1992, in advance of each year's annual dredging of the Grays Harbor Bar Channel and disposal of dredged materials at the Southwest Site. Another survey at these offshore sites was completed in April 1992 when the offshore areas were monitored as reference sites for comparison with two nearshore sites. This work was conducted in anticipation of the use of dredged materials to construct sediment berms at the nearshore sites, South Beach and Half Moon Bay. (See Appendix J for a review of this work.)

2.1 TIMING OF SURVEYS

The objective of monitoring biological resources at the Southwest Site is to avoid significant impacts to YOY Dungeness crab. Biological monitoring surveys were timed to occur after the normal peak in crab larval settlement so that an estimate of newly settled YOY crab density could be obtained. Moreover, it was critical that surveys were completed before the start of dredging operations and the use of the disposal site. These concerns were harmonized by conducting epibenthic surveys each year in June and beginning dredging operations at the Outer Bar Reach no earlier than July of each year.

2.2 VESSEL AND POSITIONING

Between 1990 and 1992, two different vessels were used for epibenthic trawls conducted near Grays Harbor by MSL. Surveys in 1990 and 1991 were conducted from the F/V Karelia, a 17-m commercial fishing boat skippered by Vern Heikkila. In 1992, the Ultimate, a 13.7-m charter fishing boat skippered by Phil Westrick was contracted. Both vessels were able to trawl at speeds of 1.5 to 2.5 knots, the optimal speed for beam trawl operations (PSEP 1990).

During all surveys, the vessel position was located using a LORAN system. An automatic range-finder function on the LORAN was used to calculate the distance covered in each trawl. Depths were recorded from the vessel's depth sounder, the accuracy of which was confirmed by the hydrographic probe.

2.3 EPIBENTHIC TRAWLS

Two identical 3-m-plumb-staff beam trawl nets were used for epibenthic surveys between 1990 and 1992. These same nets were also used during previous surveys at the Southwest Site by Pearson et al. (1987) and are the same design used in other Dungeness crab studies in Grays Harbor (Armstrong et al. 1985), Willapa Bay, and Puget Sound (Dinnel et al. 1985). This gear was designed primarily to catch juvenile crab (Gunderson and Ellis 1986; Armstrong et al. 1985). An advantage of beam trawls over other trawl designs is the fixed width for the mouth of the net, which enables better repeatability of net functioning. The beam trawls had an aluminum beam providing a fixed opening 3 m wide, a tickler chain, tom weights, a weighted "leadline" footrope, and a bridle system for towing. The 7.9-m-long (26-ft) nets had a two-seam design with a ribline and were made of 8-mm² (5/16-in.²) knotless nylon mesh. The net liners and cod ends were made of 4-mm² (5/32-in.²) knotless nylon mesh.

The sampling design was the same for all surveys. A minimum of five trawls were completed at each location (Table 2.1). Additional trawls were taken at the discretion of the chief scientist. A particular trawl was considered unacceptable if benthic organisms were absent (e.g., the catch was limited to pelagic organisms), or if the net was twisted or open upon retrieval. Unacceptable trawls were repeated without a specific attempt to return to the precise location of the initial trawl. Trawls were oriented parallel to bottom depth contours unless constrained by wind or currents. Unless indications of net overloading were evident, trawls were conducted for 10 min after the net contacted the bottom with an approximate scope of four to one and a vessel speed of 1.5 to 2.5 knots. These sampling techniques were similar to trawls conducted for the surveys in 1984 and 1985 during preliminary investigations at the ocean disposal sites (Pearson et al. 1987). Duplication of these sampling methods was critical to allow for inter-survey comparisons of Dungeness crab catches. No attempt was made, however, to return to the same trawl stations from one year to the next. The location of individual trawls is shown for the surveys in Figures 2.1 through 2.5 for 1985, 1990, 1991, and April and June 1992, respectively.

TABLE 2.1. Number of Beam Trawls Completed at the Southwest Site and North Reference Area from Dungeness Crab Surveys by MSL Between 1985 and 1992

<u>Survey Date</u>	<u>Southwest Site</u>	<u>North Reference Area</u>
June 1985	5	10
June 1990	6	7
June 1991	5	5
April 1992	5	5
June 1992	5	5

2.4 CATCH PROCESSING

The primary focus of catch processing and Tier 1 monitoring of the Southwest Site was Dungeness crab. The inventory of other catch contents differed between surveys depending on the direction provided by USACE and the goals of each survey.

2.4.1 Dungeness Crab

In 1990, Dungeness crab were sorted into juveniles (carapace width ≤ 50 mm) and adults (carapace width > 50 mm). The total catch of juvenile Dungeness crab was counted and weighed (wet weight). If the number of juvenile Dungeness crab in a trawl exceeded 100, a subsample was weighed and counted, and the total number of juveniles was estimated from the subsample weight and the total juvenile catch weight. Carapace width (CW) measurements were made on a total of 20 juveniles from each trawl. Individual adult Dungeness crab were weighed and counted, and the carapace width, shell hardness, gender, and presence (or absence) of eggs were recorded.

In 1991 and 1992, classification of Dungeness crab into two groups defined by a 50-mm CW proved inadequate because a variety of age classes were found in the trawls. In particular, this classification did not allow for distinction of megalopal larvae and 1⁺ age-class Dungeness crab. Dungeness crab from the 1⁺ age class can range from 30- to 85-mm CW in early summer (Armstrong et al. 1989). Therefore, in 1991 and 1992 Dungeness crab were categorized into four age classes (i.e., megalopal larvae, 0⁺ or YOY, 1⁺,

FIGURE 2.1. Location of Individual Trawls at the Southwest Site in June 1985

FIGURE 2.2. Location of Individual Trawls at the Southwest Site in June 1990

FIGURE 2.3. Location of Individual Trawls at the Southwest Site in June 1991

FIGURE 2.4. Location of Individual Trawls at the Southwest Site in April 1992

FIGURE 2.5. Location of Individual Trawls at the Southwest Site in June 1992

and adults) based on Armstrong et al. (1989). Dungeness crab in the YOY or 0⁺ age class have the shape of adult crab with a CW of 6 to 29 mm. Crab in the 1⁺ age class have a CW of 30 to 85 mm. All crab with CW > 85 mm were classified as adult crab.

2.4.2 Other Catch Contents

General catch characteristics recorded were total weight of the catch, nature and weight of the debris, and total wet weight of organisms from each major epifaunal taxon. The major epifaunal taxa were pooled into the following categories: other crustaceans (e.g., shrimp and hermit crabs), echinoderms (e.g., sand dollars and sea stars), mollusks (e.g., snails and clams), and other biota (e.g., jellyfish). In 1990, fish were identified by species, weighed, and counted. Because fish identification and processing is time consuming, adds significant expense to site monitoring efforts, and is not a key monitoring component of Tier 1 monitoring, fish catches were pooled into two general categories in 1991, flat fish and round fish. In 1992, however, offshore sites were monitored for comparison with nearshore sites where berm construction had been proposed (see Appendix B). To address the issue of unique biological assemblages at the proposed berm sites, fish catches from nearshore and offshore sites in 1992 were examined in greater detail, with commercially important fish identified to species.

Some trawls contained extraordinary volumes of debris that could have caused significant inaccuracies in counts of small crab if an attempt had been made to sort the entire catch. The larger components from these trawls (e.g., flat fish, large round fish, large debris, and adult crab) were removed from the whole catch and counted. The remainder of the catch was spread out on a flat surface and split. The contents of one split fraction was carefully sorted for smaller animals (e.g., juvenile crab, fish, and shrimp). Data from the split fraction were then adjusted with an appropriate multiplier to estimate the full trawl catch.

2.5 HYDROGRAPHIC PROFILES

On each day of field work in 1991 and 1992, hydrographic casts were made using an InterOcean probe to obtain a depth profile of temperature and

conductivity. If time allowed, a hydrographic profile was taken at each site on each day that sampling occurred at that site. Measurements were made at 1-m intervals in the top 5 m of the water column and at 5-m intervals thereafter.

2.7 VIDEO TOWS

For surveys in 1984 and 1985, tows with a sled-mounted video camera proved to be a useful technique for locating and characterizing areas of high crab density (Pearson et al. 1987). Underwater video tows were effective in providing a rapid assessment of the patterns of crab distribution over a large area and in precisely locating abrupt changes in crab density. The Site Monitoring Plan called for video tows and epibenthic trawls. Epibenthic trawls were recommended to complement video tows and to provide the quantitative measure of crab density that video tows could not give.

For Tier 1 monitoring surveys between 1990 and 1992, only the June 1990 survey included video tows (Higgins and Pearson 1991). These video tows were considered unsuccessful because poor visibility (≈ 0.5 m) made identification of Dungeness crab and other marine life difficult. No crabs were observed in 1990 during four lengthy video tows (≈ 1.7 to 3.5 h duration). As a result, this sampling technique was abandoned for subsequent years of monitoring.

2.8 DATA COMPILATION AND STATISTICAL ANALYSES

To normalize catch data from trawls of different lengths, trawl catches were converted to relative density of crab (or other biological categories) as crab per hectare (ha) by dividing the number of crab in each trawl by the area swept. The area swept was calculated from the distance towed and the effective fishing width for the 3-m beam trawl (2.3 m) previously determined for the net design (Armstrong et al. 1985). Because trawl data were not corrected for catch efficiency, all densities mentioned in this report are relative rather than absolute. Catches are reported as mean density (number/ha) for a particular age class of crab followed by the standard deviation (s) in parentheses.

Video tows (Pearson et al. 1987) indicated that YOY Dungeness crab were not randomly spaced but instead were clumped. The patchy distribution of crab contributed to high variability in catches from individual trawls, particularly for YOY Dungeness crab. For crab density estimates by site or by year, the standard deviation was typically as large as the mean. For trawl data, large standard deviations are not unusual because replicate trawls frequently show great variability in total numbers of fish or invertebrates (Clark 1974). Parametric analyses, such as the t-test and analysis of variance, on raw catch data involve specific assumptions about the underlying distribution that cannot necessarily be verified with the small number of trawls at each site (e.g., $n = 5$ or 6). Therefore, crab densities by year or by site were compared using a nonparametric one-way analysis of variance on the ranked data (Daniel 1978). The use of ranked data eliminates the potential bias of extremely large and small trawl catches. Individual age classes of Dungeness crab (i.e., adult or YOY) were analyzed separately. The coefficient of correlation was calculated to evaluate an association between YOY crab and debris.

Statistical analysis of replicate trawls from surveys in 1984 and 1985 indicated that four replicate trawls per site would be sufficient to detect where YOY crab density differs by a factor of 10 (Pearson 1987). This analysis was repeated with 1990 catch data, and a similar conclusion was reached. During surveys in 1991 and 1992, statistical power was estimated in the field using preliminary results of the first four or five trawls to confirm that the sampling effort was sufficient to detect a significant difference between sites. Statistical power, based on the z statistic at $\alpha = 0.05$, was calculated using the observed differences in the mean crab density from the two sites (Snedecor and Cochran 1980).

3.0 RESULTS

The mean density of YOY, 1⁺, and adult Dungeness crab is summarized for each of the five surveys in Table 3.1. Because YOY Dungeness crab are the focus of evaluation criteria in the Site Monitoring Plan, the primary emphasis of this section will be on this age class. Additional trawl and survey information is also reviewed in this section.

3.1 YOY DUNGENESS CRAB

Individual trawls from each survey produced widely varying estimates of YOY crab density at each site. In 1991, for example, YOY crab density at the North Reference area ranged from 352 to 22,316 YOY crab/ha in five trawls (Table 3.1 in Antrim et al. 1992). The variance associated with the mean density estimate at a site [designated by the standard deviation of the sample (s)] was typically equal to or greater than the mean (Table 3.2). As a result, these data require large differences between mean density estimates for definition of statistically significant differences between sites or years.

3.1.1 Comparison Between Sites

The difference in YOY crab density at the Southwest Site and North Reference area never approached the trigger value for moving to Tier 2 of the Site Monitoring Plan. With the exception of April 1992, all surveys showed a higher density of YOY Dungeness crab at the North Reference area than at the Southwest Site. The largest difference was found in 1991 when YOY crab densities differed by a factor of 4.6 between the Southwest Site and North Reference area. In no survey, however, was the difference in YOY crab density significant between the two sites ($p > 0.1$). Higgins and Pearson (1991) reported a significant difference in YOY density between these sites for 1990. Their data analysis, however, included three trawls that were partially or completely outside the Southwest Site (Figure 1.3). In addition, they used a different statistic for data analysis.

TABLE 3.1. Summary of Dungeness Crab Densities (number/ha) from Beam Trawl Surveys at Southwest Site and North Reference Area, June 1985 - June 1992

Date	YOY Crab Density (number/ha)			
	SWODS		NREF	
	Mean	s	Mean	s
June 1985	3,653	5,268	91,217	175,673
June 1990	193	373	609	411
June 1991	1,922	1,862	8,809	9,783
April 1992	20	27	13	20
June 1992	20	23	49	64

Date	1+ Crab Density (number/ha)			
	SWODS		NREF	
	Mean	s	Mean	s
June 1985	0	0	0	0
June 1990	0	0	0	0
June 1991	0	0	1	3
April 1992	4	7	0	0
June 1992	0	0	0	0

Date	Adult Crab Density (number/ha)			
	SWODS		NREF	
	Mean	s	Mean	s
June 1985	15	11	90	165
June 1990	40	35	51	34
June 1991	4	5	15	14
April 1992	10	11	40	42
June 1992	2	5	33	38

TABLE 3.2. Mean YOY Crab Density (number/ha) With Depth at Southwest Site and North Reference Area, June 1985 - June 1992

Date	< 27.4m		27.4 - 36.6m		> 36.6m	
	SWODS	NREF	SWODS	NREF	SWODS	NREF
June 1985	ND ^(a)	285	3,653	130,100	ND	437
June 1990	494	1,046	48	458	ND	399
June 1991	ND	ND	1,922	8,809	ND	ND
April 1992	ND	ND	23	13	ND	ND
June 1992	ND	0	20	61	ND	ND

(a) ND = No data.

3.1.2 High Density Area

No exceptional aggregation of YOY crab was ever found at the Southwest Site. The highest density estimate from an individual trawl at the Southwest Site was 13,000 YOY/ha. In each survey between 1990 and 1992, trawls at the North Reference area traversed the area between the 30.5-m and 36.6-m contours (100-ft and 120-ft) where juvenile crab density was exceptionally high in 1985 (~ 400,000 YOY/ha estimated from one trawl). While no densities of this magnitude were found between 1990 and 1992, two individual trawls in 1991 did find relatively high YOY crab densities, as compared to other trawls taken that year. These trawls produced density estimates of 15,926 and 22,316 YOY Dungeness crab/ha. The YOY densities in this range were found in 30% of the 10 trawls taken by Pearson et al. (1987) in June 1985. Although some trawls with high numbers of YOY crab had large amounts of debris, no correlation was found between catch debris weight and density of YOY crab ($r = 0.095$) with data from all years combined.

3.1.3 Distribution With Depth

The Site Monitoring Plan recommended video tows for rapid assessment of crab distribution over a large area and detection of abrupt changes in crab abundance. Video tows were attempted in 1990 but were unsuccessful because of poor visibility. Subsequent surveys did not include video tows. The YOY Dungeness crab density estimates by depth categories are summarized in Table 3.2. Although trawls were spaced to cover a wide portion of each site, they

were not intended to replicate the effort of video tows. Consequently, data are limited for analysis of crab distribution with depth, and statistical analyses are not possible.

3.1.4 Annual Variation

A significant difference ($p < 0.001$) in YOY crab density was detected between years. For this analysis, data from both offshore sites were combined. The density of YOY Dungeness crab was relatively high offshore in June of 1985 (mean = 62,029) and 1991 (mean = 5,365), and low in both April and June of 1992 (mean = 18 and 34, respectively) (Table 3.1). In both years with a high density of YOY Dungeness crab, mean values were driven by two trawls from the North Reference area with outstandingly high numbers.

3.2 1⁺ AGE-CLASS DUNGENESS CRAB

A total of three 1⁺ age-class Dungeness crab were retrieved from 43 trawls during four surveys between 1990 and 1992. In the June 1985 survey covering the same area, no 1⁺ age-class Dungeness crab were found. Thus, the spring density of 1⁺ age-class Dungeness crab was very low at these offshore sites.

3.3 ADULT DUNGENESS CRAB

3.3.1 Density

Over the 3-year sampling period, the estimated density of adult Dungeness crab at the two offshore sites has been relatively stable. The mean density between 1990 and 1992 ranged from 2 to 15 adult crab/ha at the Southwest Site and from 15 to 90 adult crab/ha at the North Reference area. No trend over time was detected in density of adult crab. No significant difference in adult Dungeness crab was found between the individual surveys ($p > 0.1$). For each survey, the estimated mean density of adult crab was higher at the North Reference area than at the Southwest Site; yet, in no year was the difference between the two sites significant ($p > 0.05$).

3.3.2 Other Characteristics

Adult crab were examined for sex, presence of an egg clutch, shell hardness, size, and weight. These data are summarized in Tables 3.3 and 3.4 for the Southwest Site and North Reference area, respectively. The number of female crab was consistently higher than the number of male crab. In June 1991 and 1992, no males were found out of a total of 47 adult Dungeness crab. The data are, however, too limited to identify a trend in sex ratio of adult crab. The overall male:female ratio was 1:17 for all surveys between 1990 and 1992 combined. From a total of 122 female adult Dungeness crab caught, no crabs were found with egg clutches in surveys at the two offshore sites between 1990 and 1992. A total of 2 soft-shell adults, or 1.3% of all adult crab caught, were found between 1990 and 1992. Although a few larger individuals were found in 1990, the mean CW of adult crab ranged from 10 to 15 cm and was consistent between 1990 and 1992. From a total of 122 adult Dungeness crab, a total of five were marketable crab (i.e., males ≥ 15.2 cm carapace width); four of the five were caught in 1990, and two of the five were from the Southwest Site. No examination for a trend in mean adult crab weight was conducted because of bias introduced into this analysis by missing appendages.

3.4 NON-CRAB CATCH

The species and densities of macroepifauna collected in trawls, excluding Dungeness crab, were similar at the Southwest Site and North Reference area. Catches were dominated by English sole, sand dabs, crangonid shrimp, juvenile Pacific tomcod, and surf smelt. The 3-m beam trawl used for these surveys was designed to sample juvenile crabs and flatfish. It is not recognized, however, as the best gear for sampling larger fish, shrimp, and other epibenthic animals (Mearns and Allen 1978; PSEP 1990). As a result, these data have primary value as a general description of the epibenthic community or for comparison with surveys conducted with the same gear.

TABLE 3.3. Summary of Catch Data for Adult Dungeness Crab from Trawls at the Southwest Site, June 1990 - June 1992

Month-Yr/ Station	Males				Females			
	Total Number	Number Soft Shell	Mean Carapace Width, cm	Mean Individual Weight, g	Total Number	Number Soft Shell	Mean Carapace Width, cm	Mean Individual Weight, g
06-90/203	0	0	ND (a)	ND	0	0	ND	ND
06-90/204	0	0	ND	ND	0	0	ND	ND
06-90/207	1	0	15	492	7	1	12	273
06-90/208	1	0	15	469	10	0	12	183
06-90/209	1	0	17	651	5	0	13	363
06-90/216	0	0	ND	ND	3	0	12	281
06-91/322	0	0	ND	ND	2	0	14	375
06-91/323	0	0	ND	ND	1	0	14	360
06-91/329	0	0	ND	ND	0	0	ND	ND
06-91/342	0	0	ND	ND	0	0	ND	ND
06-91/343	0	0	ND	ND	0	0	ND	ND
04-92/9213	0	0	ND	ND	2	0	12	155
04-92/9214	1	0	14	300	2	0	13	313
04-92/9215	0	0	ND	ND	1	0	15	350
04-92/9216	0	0	ND	ND	1	0	14	300
04-92/9217	0	0	ND	ND	0	0	ND	ND
06-92/502	0	0	ND	ND	0	0	ND	ND
06-92/503	0	0	ND	ND	0	0	ND	ND
06-92/504	0	0	ND	ND	0	0	ND	ND
06-92/505	0	0	ND	ND	1 (b)	0	13	240
06-92/506	0	0	ND	ND	0	0	ND	ND

(a) ND = No data available.

(b) No claws.

TABLE 3.4. Summary of Catch Data for Adult Dungeness Crab from Trawls at the North Reference Area, June 1990 - June 1992

Month-Yr/ Station	Males				Females				
	Total Number	Number Soft Shell	Mean Carapace Width, cm	Mean Individual Weight, g	Total Number	Number Soft Shell	Mean Carapace Width, cm	Mean Individual Weight, g	Number w/Egg Clutch
06-90/201	0	0	ND	ND	8	0	14	532	0
06-90/202	0	0	ND	ND	2	0	13	454	0
06-90/205	1	0	18	780	10	0	16	326	0
06-90/206	1	0	19	740	5	0	14	431	0
06-90/211	0	0	ND	ND	3	0	12	309	0
06-90/212	1	0	14	350	8	0	12	297	0
06-90/213	0	0	ND	ND	1	0	14	428	0
06-91/324	0	0	ND	ND	0	0	ND	ND	0
06-91/325	0	0	ND	ND	5	1	14	362	0
06-91/333	0	0	ND	ND	3	0	13	322	0
06-91/345	0	0	ND	ND	0	0	ND	ND	0
06-91/346	0	0	ND	ND	4	0	13	301	0
04-92/9218	0	0	ND	ND	2	0	12	155	0
04-92/9219	1	0	17	300	2	0	13	313	0
04-92/9220	0	0	ND	ND	1	0	15	350	0
04-92/9221	0	0	ND	ND	1	0	14	300	0
04-92/9222	0	0	ND	ND	0	0	ND	ND	0
06-92/508	0	0	ND	ND	1	0	12	260	0
06-92/509	0	0	ND	ND	0	0	ND	ND	0
06-92/510	0	0	ND	ND	0	0	14	320	0
06-92/511	0	0	ND	ND	19	0	13	310	0
06-92/512	0	0	ND	ND	11	0	12	260	0

(a) One female with only one claw.

(b) One female with no claws, one with only one claw, and one with one claw and one leg missing.

3.5 HYDROGRAPHIC PROFILES

Temperature and salinity data from hydrographic profiles taken in 1991 and 1992 are summarized in Table 3.5. In general, hydrographic profiles taken outside Grays Harbor indicate a well-mixed surface layer approximately 5 m deep and a trend toward decreased temperature and increased salinity, or greater water density, with depth. One important observation is the warmer temperatures measured in June 1992. The average temperatures were approximately 2°C warmer in June 1992 than in June 1991, when the average temperature was 8.9°C ($s = 1.1$) at 15 m and 7.3°C ($s = 0.1$) at 25 m. Surface waters were also warmer in 1992 than in 1991. The total depth of the profiles in 1992 was limited by the length of the cable on the probe. For these profiles, the bottom temperature and salinity values are not representative of the true site conditions.

TABLE 3.5. Summary of Hydrographic Profiles From Southwest Site and North Reference Area, June 1985 - June 1992

<u>Site</u>	<u>Date</u>	<u>Depth, m</u>	<u>Mean Temperature, °C</u>		<u>Mean Salinity, ppt</u>	
			<u>Surface</u>	<u>Bottom</u>	<u>Surface</u>	<u>Bottom</u>
North Reference	06-16-85	34.0	13.7	7.6	30.6	33.3
Southwest	06-20-91	32.0	11.7	7.2	30.9	32.6
North Reference	06-20-91	33.0	12.2	7.2	30.8	32.6
Southwest	06-21-91	37.4	12.7	7.2	29.6	32.6
North Reference	06-21-91	31.5	12.8	7.3	29.5	32.5
North Reference	04-23-92	15.0 ^(a)	12.2	11.8	29.0	31.5
North Reference	06-15-92	25.0 ^(b)	15.0	9.2	28.8	32.8
Southwest	06-16-92	25.0 ^(c)	13.3	9.4	30.3	32.6

(a) Depth of measurement limited by cable length on CTD unit. Total depth was 37.0 m.

(b) Depth of measurement limited by cable length on CTD unit. Total depth was 38.7 m.

(c) Depth of measurement limited by cable length on CTD unit. Total depth was 29.3 m.

4.0 DISCUSSION

4.1 EVALUATION OF SITE MONITORING PLAN

In the Site Monitoring Plan, the objective of Tier 1 monitoring of the Southwest Site is to verify that the area of high crab density has not moved onto the site. Section 4.1.1 provides a discussion of monitoring results. The choice of the beam trawl as the preferred sampling gear has been questioned in public hearings related to Dungeness crab surveys. The rationale for using beam trawls as opposed to other sampling gear is given in Section 4.1.2.

4.1.1 YOY Crab Density

A significant difference in YOY crab density was never found between the two sites. Site monitoring surveys in 1990, 1991, and 1992, therefore, confirmed that the selection and boundary locations of the Southwest Site remain appropriate.

In comparison to the North Reference area, high densities of YOY Dungeness crab were not found at the Southwest Site. From a total of 26 trawls taken at the Southwest Site in spring surveys between 1985 and 1992, the highest density estimate from an individual trawl was 12,890 YOY crab/ha in 1985. Approximately 20% of 32 trawls at the North Reference area produced YOY density estimates greater than 15,000 YOY crab/ha. Therefore, there has been no indication that areas of dense YOY crab aggregation that have been found at the North Reference area exist at the Southwest Site.

Settlement of megalopal larvae and densities of YOY Dungeness crab were exceptionally high in 1985, with density estimates as high as 400,000 YOY/ha (Pearson et al. 1987; Armstrong et al. 1989). None of the surveys between 1990 and 1992 found YOY densities of this magnitude. Estuarine studies confirm no exceptional years since 1985 (Dr. David Armstrong, University of Washington, personal communication). Annual fluctuations in YOY crab abundance may be attributable to natural cycles that occur over an interval of approximately 9 to 10 years (Botsford and Wickham 1978). Nevertheless, surveys between 1990 and 1992 may not have sampled the peak in YOY crab density, either by missing the timing of peak settlement or bypassing

aggregations of YOY crab. While the peak in larval settlement generally occurs in May through mid-June, there is considerable interannual variation in the timing of settlement (Armstrong et al. 1989). In 1992, for example, larval settlement may have been accelerated by oceanographic events such as El Niño (see Section 4.2.3). The nearshore substrate is relatively homogeneous sand and gravel and offers little refuge for juvenile crab. In this environment, aggregations of YOY crab can dissipate quickly as the crab move into more protected estuarine waters or are cropped by predation (Armstrong et al. 1989). The limited coverage provided by a few trawls may have missed aggregations of YOY crab, particularly if aggregations occurred on a fine scale (i.e., YOY crab were not widely dispersed).

4.1.2 Choice of Sampling Gear

In public meetings related to berm construction in nearshore areas, crab pots have been suggested as a more effective sampling technique for Dungeness crab. This criticism could also apply to surveys at or near the Southwest Site.

Crab pots are not an effective gear choice for quantification of juvenile or adult crab densities. Dungeness crab rely on chemoreception to detect chemical food cues (Pearson et al. 1979), and strong reliance by crab on chemical cues is what supports commercial fisheries using baited traps. Although juvenile crab can be retrieved from commercial traps, their catch is incidental and not quantitative without standardization of bait, soak time, and other factors in fishing effort. Further, commercial traps are specifically designed to provide escape routes for juveniles. Also, the zone of attraction from which a baited trap draws crab has been estimated (Pearson and Woodruff 1987) but not confirmed by field studies. To develop trap equipment and techniques for quantitative measures of crab abundance, extensive studies would be necessary.

The beam trawl was designed specifically for quantitative sampling juvenile crab and fish (Gunderson and Ellis 1986). The YOY Dungeness crab are not able to avoid capture by the beam trawl except by burial in the sediment. Another advantage of the beam trawl is the fixed opening size, which provides better replication between trawls and more accurate estimation of area swept.

For site monitoring studies, an additional advantage is consistency with previous surveys.

In general, the efficiency of trawl nets for capture of demersal fish is typically low, i.e., 10% to 50% (Mearns and Allen 1978). Nonetheless, otter trawls are widely used for commercial fisheries. Beam trawls are very similar in design to otter trawls but may not be as effective for capturing larger fish, shrimp, or other highly mobile animals (Mearns and Allen 1978). Highly mobile animals, like adult crab, have several mechanisms to escape capture from a trawl net including burying in the sediment, movement away from the approaching trawl, and crawling out of the net opening. A beam trawl may, therefore, not be the most effective gear for sampling adult Dungeness crab, yet it is acceptable for comparisons of relative density. Beam trawls are, however, valuable for accurate quantification of relative densities of juvenile flatfishes and juvenile crab (Holme and McIntyre 1971). Because YOY crab were the focus of this site monitoring, the beam trawl was the most appropriate and effective gear for surveys.

4.2 SUPPLEMENTAL FINDINGS ON DUNGENESS CRAB ECOLOGY

4.2.1 General Crab Distribution

Epibenthic trawl catches indicated that the distribution of YOY Dungeness crab was patchy at the Southwest Site and North Reference area. Although some integration of high and low YOY crab densities in fine-scale patches would naturally occur over the duration of each trawl, there remained substantial variability in YOY crab density estimates between individual trawls at a given site. Video tows of the ocean bottom by Pearson et al. (1987) provided a visual and qualitative picture of the epibenthic community. These video tows revealed that juvenile crab density changed sharply over short distances, and that juvenile crab appear to be concentrated in aggregations that were most often oriented along depth contours. Presumably aggregations of YOY crab are associated with fine-scale variations in the physical environment.

4.2.2 Association of Juvenile Crab with Debris

Dungeness crab are known to use eelgrass, wood debris, gravel, and shell fragments for protected habitat (Armstrong et al. 1989). With data from the

June 1985 survey, Pearson et al. (1987) found a positive correlation between YOY crab and debris from trawl catches. Off Willapa Bay, Carrasco et al. (1985) found juvenile crab density highest in areas with epibenthic cover of eelgrass debris. Yet, a similar analysis using data from surveys in 1990, 1991, and 1992 found no correlation between debris and YOY crab density. This result may be because survey efforts were not directed toward characterization of physical bottom characteristics.

The study area is subject to strong movements of water driven by wave action as well as tidal, coastal, and wind-driven currents. The general sediment characteristics of the study area are relatively smooth sand with areas of gravel intermixed (Armstrong et al. 1989). In addition to patches of gravel, epibenthic habitat for juvenile crab is available in dispersed areas of debris off Grays Harbor. Debris in trawls from the study area consisted mostly of wood pieces, bryozoans, algae, and eelgrass. Debris items are not likely to provide stable physical habitat for long-term sheltering of juvenile crab in areas of strong water movement. As a result, juvenile crab rapidly move toward more protected estuarine waters.

4.2.3 Effects of El Niño

El Niño is an oceanographic phenomenon that can result in unusually warm surface water. An El Niño of moderate intensity occurred in the winter and spring of 1992. Hydrographic casts in June 1992 confirmed that water temperatures were approximately 2°C warmer than in 1991. Warmer water temperatures could accelerate the development of pelagic megalopal larvae and cause an early onset of settlement. Dr. David Armstrong (University of Washington, personal communication) found that larval settlement in the Grays Harbor estuary in 1992 occurred at three peaks between early April and mid-May and was not of exceptional abundance. Although two surveys in 1992 found very few YOY crab, the surveys may have missed the peak density in the study area. The effects of El Niño on crab larval settlement outside the estuary are unknown.

4.2.4 1⁺ and Adult Dungeness Crab

Juvenile crab from the 1⁺ age class were found at a very low density near the Southwest Site. Because they are not likely to have the necessary

mobility to avoid capture by the trawl net, 1⁺ crab density estimates are considered accurate. This result confirms previous findings that very few 1⁺ age-class crab are found in offshore areas (Armstrong et al. 1989).

No distinct trends in adult crab density or size were identified from this work. The biased sex ratio of approximately 1:17 (male:female) may be indicative of natural segregation by sex or removal of male crabs from the area by the commercial fishery.

5.0 CONCLUSIONS

Tier 1 monitoring efforts at the Southwest Site have confirmed that the selection of the site location and boundaries remains appropriate. In surveys conducted between 1990 and 1992, no dense aggregation of YOY Dungeness crab was located within the Southwest Site. In fact, for no site monitoring survey was the difference in YOY crab density significant between the Southwest Site and the North Reference area.

6.0 REFERENCES

- Antrim, L. D., V. I. Cullinan, and W. H. Pearson. 1992. Dungeness Crab Survey for the Southwest Ocean Disposal Site and Additional Sites off Grays Harbor, Washington, June 1991. PNL-7932, prepared for the Seattle District of the U.S. Army Corps of Engineers by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory, Richland, Washington.
- Armstrong, D. A., D. Gunderson, and J. L. Armstrong. 1985. Juvenile Dungeness Crab Population Dynamics Offshore of the Grays Harbor Estuary, Spring and Summer, 1984. Technical Report to the U.S. Army Corps of Engineers, Seattle District prepared by ARCS, Seattle, Washington.
- Armstrong, D. A., L. Botsford, and G. Jamieson. 1989. Ecology and Population Dynamics of Juvenile Dungeness Crab in Grays Harbor Estuary and Adjacent Nearshore Waters of the Southern Washington Coast. Report to the U.S. Army Corps of Engineers, Seattle District.
- Botsford, L. W. and D. E. Wickham. 1978. "Behavior of Age-Specific, Density-Dependent Models and the Northern Dungeness Crab (Cancer magister) Fishery." J Fish. Res. Board Can. 35:833-943.
- Carrasco, K. R., D. A. Armstrong, D. R. Gunderson, and C. Rogers. 1985. "Abundance and Growth of Cancer magister Young-of-the-Year in the Nearshore Environment." In Proceedings of the Symposium on Dungeness Crab Biology and Management, October 1984. Alaska Sea Grant Report No. 85-3:171-184.
- Clark, S. H. 1974. "A Study of Variation in Trawl Data Collected in Everglades National Park, Florida." Trans. Amer. Fish. Soc. 103(4):777-785.
- Daniel, W. W. 1978. Applied Nonparametric Statistics. Houghton Mifflin Company, Boston.
- Dinnel, P. A., D. A. Armstrong, and C. Dungan. 1985. "Initiation of a Dungeness Crab, Cancer magister, Habitat Study in North Puget Sound," pp. 327-340. In Proceedings of the Symposium on Dungeness Crab Biology and Management, University of Alaska, Alaska Sea Grant Report No. 85-3.
- Gunderson, D. R., and I. E. Ellis. 1986. "Development of a Plumb Staff Beam Trawl for Sampling Demersal Fauna." Fisheries Res. 4:35-41.
- Higgins, B. H., and W. H. Pearson. 1991. Dungeness Crab Survey for the Southwest Ocean Disposal Site off Grays Harbor, Washington, June 1990. PNL-5045, prepared for the Seattle District of the U.S. Army Corps of Engineers by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory, Richland, Washington.
- Holme, N. A., and A. D. McIntyre. 1971. Methods for the Study of Marine Benthos. IBP Handbook 16. Blackwell Scientific Publications, Oxford, England.

Mearns, A. J., and M. J. Allen. 1978. Use of Small Otter Trawls in Coastal Biological Surveys. EPA 600/3-78/083. U.S. Environmental Protection Agency, Office of Research and Development, Corvallis, Oregon.

Pearson, W. H. 1987. Conceptual Plan for the Management and Monitoring of Two Ocean Disposal Sites off Grays Harbor, Washington. PNL-6420, prepared for the Seattle District of the U.S. Army Corps of Engineers by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory, Richland, Washington.

Pearson, W. H. and D. L. Woodruff. 1987. Effects of Dredged Materials on Bait Odor Response in the Dungeness Crab, Cancer magister. PNL-6436, prepared for the Seattle District of the U.S. Army Corps of Engineers by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory, Richland, Washington.

Pearson, W. H., D. L. Woodruff, P. Wilkinson, J. S. Young, H. L. McCartney, and D. C. Klopfer. 1987. Data Report for the 1984/1985 Ocean Surveys to Investigate Potential Ocean Disposal Sites off Grays Harbor, Washington. PNL-6280, prepared for the Seattle District of the U.S. Army Corps of Engineers by Battelle/Marine Sciences Laboratory Pacific Northwest Laboratory, Washington.

Pearson, W. H., P. C. Sugarman, D. L. Woodruff, and B. L. Olla. 1979. Thresholds for Detection and Feeding Behavior in the Dungeness Crab, Cancer magister. J. Exp. Mar. Biol. Ecol. 39:65-78.

PSEP (Puget Sound Estuary Program). 1990. Recommended Guidelines for Sampling Soft-Bottom Demersal Fishes by Beach Seine and Trawl in Puget Sound. Prepared by PTI Environmental Services for the U.S. Environmental Protection Agency.

Snedecor, G. W., and W. G. Cochran. 1980. Statistical Methods, 7th ed. The Iowa State University Press, Ames, Iowa.

USACE (U.S. Army Corps of Engineers). 1982. Grays Harbor Navigation Channel Improvement Project Feasibility Report and Environmental Impact Statement. U.S. Army Corps of Engineers, Seattle, Washington.

USACE (U.S. Army Corps of Engineers). 1989. Final Environmental Impact Statement Supplement, Grays Harbor, Washington, Navigation Improvement Project. U.S. Army Corps of Engineers, Seattle, Washington.

Zeller, R. W., and T. A. Wastler. 1986. "Tiered Ocean Disposal Monitoring Will Minimize Data Requirements," pp. 1004-1009. In Oceans 86 Conf. Rec. IEEE Service Center, Piscataway, New Jersey.

APPENDIX A

1992 PRELIMINARY DATA REPORTS

Pacific Northwest Division
Marine Sciences Laboratory
439 West Sequim Bay Road
Sequim, Washington 98382
(206) 683-4151

May 4, 1992

Mr. Bert Brun
U. S. Army Corps of Engineers
Seattle District
P.O. Box C-3755
Seattle, WA 98124-2255

Dear Bert:

This letter is the Preliminary Cruise Report for the 1992 Grays Harbor Biological Surveys. Tasks 1 and 2 of the 1992 Crab and Razor Clam Surveys were successfully completed between April 21 and 24, 1992. The technical contact at the U. S. Army Corps of Engineers, Seattle District (USACE) was Mr. Bert Brun. The staff from Battelle/Marine Sciences Laboratory (MSL) included Liam Antrim (Project Manager), Dr. Walter Pearson (Technical Advisor), Dave Shreffler (Chief Scientist), and Dr. Ray Bienert, Tom Parkwell, and Christina Lobo (field crew). Favorable weather conditions enabled us to complete the surveys with only one field day lost because of high seas. Phil Westrick, skipper of the M/V Ultimate, provided us with expert assistance and knowledge of the local conditions.

Surveys were conducted at five sites in the area of Westport and the Southwest Ocean Disposal Site (SWODS). Sampling efforts included a total of 18 bottom trawls with a 3-m plumb-staff beam trawl, 11 Van Veen grabs for grain-size analysis, 24 Smith-McIntyre grabs for razor clams, and 3 hydrographic casts for depth profiles of salinity and temperature. The results from each site are discussed separately below. Also included is a table summarizing the trawl data for Dungeness crab.

Southwest Ocean Disposal Site

Sampling at the SWODS was conducted on April 23, 1992. A total of five successful trawls were completed. At this site, as well as all other trawl sites, Dungeness crabs were sorted into four size classes: megalopae (carapace width < 6 mm), 0+ (carapace width 6-29 mm), 1+ (carapace width 30-85 mm), and adults (carapace width > 85 mm). The mean density of all sizes of Dungeness crabs combined was 39 crabs/hectare ($s = 27$; range = 5 to 73/ha). The mean density of megalopae was 5/ha ($s = 5$; range = 0 to 11/ha). The mean density of 0+ was 23/ha ($s = 26$; range = 0 to 57/ha). The mean density of 1+ was 3/ha ($s = 4$; range = 0 to 8/ha). The mean density of adults was 8/ha ($s = 12$; range = 0 to 28/ha). As planned, no surface sediment grabs were taken at this site for either razor clams or grain size analysis.

Mr. Bert Brun
May 5, 1992
Page 2

North Reference Site

The area to the north of the SWODS is considered a reference site for surveys conducted at the SWODS. A total of five successful trawls were completed on April 23, 1992. These trawls were positioned to survey the area where high densities of juvenile Dungeness crabs were located in previous trawl surveys in 1985, 1990, and 1991. The mean total density of Dungeness crabs was 67 crabs/hectare ($s = 42$; range = 28 to 125/ha). The mean density of megalopae was 13/ha ($s = 14$; range = 0 to 33/ha). The mean density of 0+ was 13/ha ($s = 20$; range = 0 to 47/ha). No 1+ were found in any of the five trawls. The mean density of adults was 40/ha ($s = 42$; range = 0 to 94/ha). As planned, no surface sediment grabs were taken at this site for either razor clams or grain size analysis.

South Beach Berm Site

Four successful trawls were conducted at the South Beach Berm Site--one on April 21 and three on April 23. The mean total density of Dungeness crabs was 385 crabs/hectare ($s = 254$; range = 91 to 623/ha). The mean density of megalopae was 14/ha ($s = 6$; range = 7 to 21/ha). The mean density of 0+ was 216/ha ($s = 206$; range = 53 to 514/ha). The mean density of 1+ was 118/ha ($s = 221$; range = 0 to 449/ha). The mean density of adults was 37/ha ($s = 12$; range = 27 to 54/ha). Based on visual inspection, six surface sediment grabs at this site were comprised primarily of clean, fine-grain sand. Razor clams were sampled from four sites along each of three transects ($n=12$ samples) perpendicular to the shore. Samples were taken with the Smith-McIntyre grab at the 30 ft, 35 ft, 40 ft, and 45 ft contours. No razor clams were found. The samples contained primarily fine-grain sand with some pea gravel, sand dollars, crangon shrimp, polychaetes, molluscs, amphipods, and brittle stars.

Half Moon Bay Berm Site

Four successful trawls were conducted at the Half Moon Bay Berm Site on April 21, 1992. The mean total density of Dungeness crabs was 1072 crabs/hectare ($s = 1210$; range = 105 to 2767/ha). This mean total density is mostly because of the high densities of megalopae (657/ha; $s = 673$; range = 57 to 1454/ha) and 0+ (333/ha; $s = 537$; range = 0 to 1135/ha). The mean density of 1+ was 37/ha ($s = 51$; range = 0 to 113/ha). The mean density of adults was 45/ha ($s = 18$; range = 27 to 66/ha). The higher mean total density observed at this site relative to the other three is probably attributable to the presence of more debris in the trawls at Half Moon Bay. Juvenile crabs (megalopae, 0+, 1+) are known to utilize debris for cover. Based on visual inspection, five surface sediment grabs at this site were comprised primarily of clean, fine-grain sand.

Mr. Bert Brun
May 5, 1992
Page 3

Washington Department of Fisheries Razor Clam Site "I"

Razor clams were sampled from four sites along each of three transects (n=12 samples) perpendicular to the shore. Samples were taken with the Smith-McIntyre grab at the 25 ft, 30 ft, 35 ft, and 40 ft contours. The estimated penetration depth of the grab was 3-4 cm. No juvenile or adult razor clams were found. The absence of adult razor clams in the grab samples was expected, because the Smith-McIntyre grab does not typically penetrate the sediment deeply enough to capture adults (pers. comm. Doug Simons, Washington Department of Fisheries). The absence of juvenile razor clams in the grab samples was probably a result of the timing of the April survey. Peak spawning time for razor clams on Washington beaches varies from mid-May through July (Lassuy and Simons 1989). Thus, the April Smith-McIntyre grab samples were probably taken too early in the spring to capture any juvenile razor clams. The samples contained primarily fine-grain sand and sand dollars with some crangon shrimp, polychaetes, and molluscs.

Conclusions

In April 1992, the mean total density of Dungeness crab was 1072/ha (s = 1210) at the Half Moon Bay Berm Site, 385/ha (s = 254) at the South Beach Berm Site, 97/ha at the SWODS (s = 27), and 67/ha (s = 42) at the North Reference Site. Because the crab densities within the two berm sites did not exceed the threshold value (100 times higher than the North Reference Site), Tier 2 activities outlined in the site monitoring plan are not required at this time. However, the final determination of compliance with the site monitoring plan will not be made until the May/June survey is completed.

References

Lassuy, D. R. and D. Simons. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest)--Pacific razor clam. U.S. Fish. Wildl. Serv. Biol. Rep. 82(11.89). U.S. Army Corps of Engineers, TR-EL-82-4. 16 pp.

Please contact me at 681-3659 if you need any additional information before the next scheduled survey in May/June.

Sincerely,

Dave Shreffler
Dave Shreffler
Research Scientist

:DKS

Preliminary results of 1992 Grays Harbor epibenthic surveys. Crab densities are reported as number/hectare not number/haul.

Site	Station	Distance Trawled (nm)	Area Swept (ha)	Densities (No./ha)			
				Megalopae	0+	1+	Adult
Half Moon Bay	GH9201	0.25	0.1066	1454	1135	113	66
	GH9202	0.26	0.1109	144	108	0	54
	GH9203	0.26	0.1109	974	90	18	27
	GH9204	0.29	0.1237	57	0	16	32
				Mean	657.25	333.25	36.75
				Std. Deviation	673.02	536.58	51.47
							44.75
							18.39
							1072.00
							1210.33
South Beach	GI9211	0.35	0.1492	7	114	449	54
	GI9212	0.44	0.1876	11	53	0	27
	GI9224	0.22	0.0938	21	181	21	32
	GI9225	0.26	0.1109	18	514	0	36
				Mean	14.25	215.50	117.50
				Std. Deviation	6.40	205.75	221.22
							37.25
							11.76
							384.50
							253.78
SWODS	GI9213	0.31	0.1321	8	15	8	0
	GH9214	0.25	0.1066	0	0	0	28
	GI9215	0.62	0.2643	11	42	0	4
	GI9216	0.29	0.1236	8	57	0	8
				Mean	14.25	215.50	117.50
				Std. Deviation	6.40	205.75	221.22
							37.25
							11.76
							384.50
							253.78
North Rel.	GI9218	0.44	0.1876	21	16	0	16
	GI9219	0.15	0.0640	0	47	0	78
	GI9220	0.43	0.1834	33	0	0	0
	GI9221	0.67	0.2857	11	4	0	14
				Mean	13.00	13.40	0.00
				Std. Deviation	14.20	19.89	0.00
							40.40
							42.46
							66.60
							41.73

Pacific Northwest Division
Marine Sciences Laboratory
439 West Sequim Bay Road
Sequim, Washington 98382
(206) 683-4151

June 20, 1992

Bert Brun
Army Corps of Engineers
Seattle District
P.O. Box C-3755
Seattle, WA 98124-2255

Dear Bert:

This letter is the Letter Report for the June 1992 Grays Harbor Biological Survey. For this project, the technical contact at the U.S. Corps of Engineers, Seattle District (USACE), was Mr. Bert Brun. The field crew from Battelle/Marine Sciences Laboratory (MSL) were Liam Antrim (Chief Scientist), Dave Shreffler, and Tom Parkwell. Dr. Walter Pearson was the Technical Advisor for the project.

Surveys were conducted from the chartered vessel the M/V Ultimate. The competent seamanship of the skipper, Phil Westrick, helped the survey to be performed efficiently and safely. Favorable weather conditions allowed for field work to be completed in two days, on June 15 and 16, 1992. Bottom trawls were conducted at a total of four sites in the Grays Harbor area: two offshore sites, the Southwest Ocean Disposal site (SWODS) and the reference area to the north of the SWODS (North Reference site); and two nearshore sites where sediment berms have been proposed or built, the South Beach site, and the Half Moon Bay site. As planned, no sediment samples were taken for grain-size analysis or razor clam sampling.

Sampling efforts included a total of 18 successful bottom trawls with a 3-m plumb-staff beam trawl and 3 hydrographic casts for depth profiles of salinity and temperature. Successful trawls were indicated by the presence of benthic organisms in the catch (e.g., sand dollars, snails, hermit crabs, and/or flat fish) and an unfouled net. Detailed sorting of each trawl catch was completed for Dungeness crab and commercially important fish. Other faunas were grouped into the following major epifaunal taxa: other crustaceans, other fish, echinoderms, mollusks, and other biota. Non-faunal debris in each catch was weighed and characterized. Dungeness crab were sorted into four age-classes: megalopal larvae, 0+ age class or young-of-the-year (YOY) (6 to 29 mm carapace width), 1+ age class (30 to 85 mm carapace width), and adults (>85 mm carapace width). Commercially important fish were identified to species.

Mr. Bert Brun
June 20, 1992
Page 2

The results from each site are discussed separately below. Because trawl data were corrected for area swept but not catch efficiency, all densities reported in this letter are relative rather than absolute densities. Also included in this letter report is a discussion of salient points and a table summarizing the trawl data for Dungeness crab. Additional survey data will be provided in subsequent reports to USACE.

Southwest Ocean Disposal Site

Five trawls were completed at the SWODS on June 15, 1992. In one of these trawls, Dungeness crab were not found. From all five trawls at this site, the mean total density of Dungeness crab was 22 crab/hectare (ha) ($s = 21$, range = 0 to 53/ha). YOY crab were found in three of the five trawls, with a mean density of 20 YOY crab/ha ($s = 23$, range = 0 to 53/ha). Adult crab were retrieved on one trawl, resulting in a mean of 2 adult crab/ha ($s = 5$, range = 0 to 10/ha). No megalapae or 1⁺ age-class Dungeness crab were found at the SWODS.

North Reference Site

The area to the north of the SWODS is considered a reference site for surveys conducted at the SWODS. Relatively high densities of YOY Dungeness crab have been found in this area during previous surveys by MSL. Five trawls were completed at the North Reference site on June 15 and 16, 1992. Trawls were positioned to survey the area where high YOY crab densities had been found in previous years. As was the case with the SWODS, only adult and YOY Dungeness crab were found at this site. The mean total density of Dungeness crab was 82 crab/ha ($s = 45$, range = 19 to 144/ha). Three trawls contained YOY crab with a mean density of 49 YOY crab/ha ($s = 64$, range = 0 to 144/ha). Four of the five trawls contained adult Dungeness crab with a mean density of 33 adult crab/ha ($s = 38$, range = 0 to 76/ha).

South Beach Site

Four trawls were completed at the South Beach site on June 16, 1992. No Dungeness crab were found in the trawl conducted at the shallowest depth, approximately 30 feet MLLW. From all four trawls at this site, the mean total density of Dungeness crab was 15 crab/ha ($s = 14$, range = 0 to 31/ha). The mean density of YOY Dungeness crab was 2 YOY crab/ha ($s = 3$, range = 0 to 6/ha). Adult Dungeness crab were captured at a mean density of 8 adult crab/ha ($s = 15$, range = 0 to 31/ha). The mean density of 1⁺ age-class crab was 6 crab/ha ($s = 11$, range = 0 to 23/ha). No megalopae were found in any of the trawls.

Half Moon Bay Site

Four trawls were completed at Half Moon Bay on June 16, 1992. Before and during trawling efforts, a dredge was disposing of dredged materials in Half Moon Bay. The one release of material we witnessed occurred at the eastern end of the bay between the time of our second and third trawls. None of the trawls traversed the area where this one dump occurred. Although all trawls contained Dungeness crab from the YOY (0⁺), 1⁺, and adult age classes, each catch was dominated by 1⁺ age-class crab. No megalopal larvae were found in any of the trawls. The mean total density of Dungeness crab was 1018 crab/ha (s = 46, range = 979 to 1077/ha). YOY crab were found at a mean density of 63 YOY crab/ha (s = 73, range = 16 to 171/ha). The mean density of 1⁺ age-class crab was 876 1⁺ age-class crab/ha (s = 81, range = 785 to 949/ha). Adult Dungeness crab were found at a mean density of 79 adult crab/ha (s = 47, range = 31 to 143/ha). Trawls from this site contained a relatively large proportion of wood debris, the quantity of which decreased with sampling depth.

Summary Notes and Comments

- No megalopal larvae were caught in any of the trawls taken in June 1992. Two possible explanations for this result are that 1) the survey was conducted after the majority of the annual settlement of megalopae had occurred and/or 2) numbers of Dungeness crab larvae were low in 1992.
- At all four sites, the densities of YOY (0⁺ age class) Dungeness crab were low compared to the survey in June 1991. Compared to results from the April 1992 survey, YOY Dungeness crab densities were low at nearshore sites but similar at offshore sites. Because the majority of larval settlement and metamorphosis to YOY crab occurs in May through mid-June (Armstrong et al. 1989), the April 1992 survey may have preceded the peak of settlement and the June 1992 survey may have followed a peak in settlement. A strong settlement could have occurred in offshore and/or nearshore areas between the two surveys. Typically, there is a period of high mortality immediately following larval settlement. Thus, the June 1992 survey may have missed the peak YOY densities. An alternative hypothesis is that larval settlement in 1992 was weak. Armstrong et al. (1989) describes high variability in interannual recruitment of YOY crab.
- In June 1992, the density of 1⁺ Dungeness crab was higher at Half Moon Bay than in both June 1991 and April 1992. According to Armstrong et al. (1987) densities of 1⁺ crab in the Grays Harbor Bar to South Beach areas are highest in June to September, and this peak density results from an inshore migration of 1⁺ crab. Between 1983 and 1985, they found 1⁺ crab densities between 274 and 1175/ha. The results from the June 1992 survey (876 1⁺ crab/ha) lie within this range.

Mr. Bert Brun
June 20, 1992
Page 4

The data in this report should provide USACE and WDF with information to assess the need for any change in dredged-materials disposal activities at the survey sites. If any points requiring clarification arise, please contact me at (206) 681-3620.

References

Armstrong, D. A., T. C. Wainwright, J. Orensanz, P. A. Dinnel, and B. R. Dumbauld. 1987. Model of Dredging Impact on Dungeness Crab in Grays Harbor, Washington. FRI-UW-8702, Fisheries Research Institute, University of Washington, Seattle, Washington. Final Report to U.S. Army Corps of Engineers, Seattle District.

Armstrong, D. A., L. Botsford, and G. Jamieson. 1989. Ecology and Population Dynamics of Juvenile Dungeness Crab in Grays Harbor Estuary and Adjacent Nearshore Waters of the Southern Washington Coast. Report to the U.S. Army Corps of Engineers, Seattle District.

Sincerely,

Liam Antrim
Research Scientist

attachment

TABLE Dungeness Crab Catch Data from Bottom Trawls Conducted for the June 1992 Grays Harbor Biological Survey. (Site 1 = Southwest Ocean Disposal site; Site 2 = North Reference site; Site 3 = South Beach site; Site 4 = Half Moon Bay site)

Site	Area Swept (ha)	Trawl Catch (numbers of crab)					Crab Density Estimates				
		All Dungeness Crabs	Age Class			Total Dungeness Density (#/ha)	Total Megalopae Density (#/ha)	Total 0+ Density (#/ha)	Total 1+ Density (#/ha)	Total Adult Density (#/ha)	
			Megalopae	0+	1+						
1	0.14	5	0	5	0	34	0	34	0	0	
1	0.13	7	0	7	0	53	0	53	0	0	
1	0.09	1	0	1	0	12	0	12	0	0	
1	0.10	1	0	0	1	10	0	0	0	10	
1	0.06	0	0	0	0	0	0	0	0	0	
	Mean (s)					22 21	0 0	20 23	0 0	2 5	
2	0.08	8	0	7	0	98	0	86	0	12	
2	0.11	16	0	16	0	144	0	144	0	0	
2	0.21	4	0	3	0	19	0	14	0	5	
2	0.26	19	0	0	19	74	0	0	0	74	
2	0.16	12	0	0	12	76	0	0	0	76	
	Mean (s)					82 45	0 0	49 64	0 0	33 38	
3	0.17	0	0	0	0	0	0	0	0	0	
3	0.13	3	0	0	3	23	0	0	23	0	
3	0.16	1	0	1	0	6	0	6	0	0	
3	0.16	5	0	0	5	31	0	0	0	31	
	Mean (s)					16 14	0 0	2 3	6 11	8 15	
4	0.13	126	0	2	120	985	0	16	938	31	
4	0.18	185	0	3	170	1033	0	17	949	67	
4	0.09	101	0	16	78	1078	0	171	832	75	
4	0.10	96	0	5	77	979	0	51	785	143	
	Mean (s)					1018 46	0 0	63 73	876 81	79 47	

SUPPLEMENTAL TABLES AND FIGURES

1992 DUNGENESS CRAB SURVEYS FOR THE
SOUTHWEST OCEAN DISPOSAL SITE AND
TWO PROPOSED BERM SITES
OFF GRAYS HARBOR, WASHINGTON

L. D. Antrim
D. K. Shreffler
R. W. Bienert
W. H. Pearson

Battelle/Marine Sciences Laboratory
Sequim, Washington

September 1992

Prepared for the
U.S. Army Corps of Engineers - Seattle District
under a Related Services Agreement
with the U.S. Department of Energy
Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute

TABLES

1	Trawl Catch Data Summary from the April 1992 Epibenthic Survey Near Grays Harbor, Washington	1
2	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Southwest Ocean Disposal Site in April 1992	2
3	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the North Reference Site in April 1992	2
4	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the South Beach Site in April 1992	3
5	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Half Moon Bay Site in April 1992	3
6	Trawl Catch Data Summary from the June 1992 Epibenthic Survey Near Grays Harbor, Washington	4
7	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Southwest Ocean Disposal Site in June 1992	5
8	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the North Reference Site in June 1992	5
9	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the South Beach Site in June 1992	6
10	Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Half Moon Bay Site in April 1992	6
11	Grain Size Analysis of Sediment Grab Samples Collected in April 1992 from the Grays Harbor Area	7
12	Hydrographic Data from Sites Near Grays Harbor, Washington Collected for the Dungeness Crab Survey Between April 21 and 24, 1992	8
13	Hydrographic Data from Sites Near Grays Harbor, Washington, Collected for the Dungeness Crab Survey Between June 15 and 16, 1992	9

FIGURES

1a.	Grain Size Analysis of Sediment Samples Collected from the South Beach Site near Grays Harbor, Washington, in April 1992	10
1b.	Grain Size Analysis of Sediment Samples Collected from the South Beach Site near Grays Harbor, Washington, in April 1992	11
1c.	Grain Size Analysis of Sediment Samples Collected from the Half Moon Bay Site near Grays Harbor, Washington, in April 1992	12
1d.	Grain Size Analysis of Sediment Samples Collected from the Half Moon Bay Site near Grays Harbor, Washington, in April 1992	13

TABLE 1. Trawl Catch Data Summary (Excluding Dungeness Crab Catch) from the April 1992 Epibenthic Survey Near Grays Harbor, Washington (Results are presented in kg/ha.)

Site (a)	Biological Catch Wt.	Debris Wt.	Commercially Important Fish	Other Fish	Crustaceans (b)	Echinoderms	Mollusks	Other Taxa
1	36.33	1.65	28.92	4.88	0.30	0.00	0.91	0.00
1	132.46	5.25	62.01	64.35	1.03	0.00	2.81	0.00
1	27.42	2.12	11.35	5.30	0.23	0.00	0.87	0.00
1	56.87	4.24	30.62	9.71	1.29	0.00	1.05	0.00
1	13.70	1.07	5.90	4.99	0.05	0.00	0.25	0.00
Mean	53.36	2.87	27.76	17.85	0.58	0.00	1.18	0.00
s	46.92	1.79	21.96	26.07	0.55	0.00	0.96	0.00
2	10.96	0.34	3.89	1.44	0.11	0.00	0.27	0.00
2	220.55	7.11	175.46	37.50	2.97	0.00	1.56	0.00
2	14.73	0.33	2.54	5.45	0.41	0.00	0.05	0.00
2	24.17	0.34	11.80	7.07	0.35	0.00	0.07	0.00
2	105.10	1.41	25.92	62.21	1.64	0.00	1.29	0.59
Mean	75.10	1.91	43.93	22.73	1.10	0.00	0.65	0.12
s	90.01	2.95	74.12	26.33	1.20	0.00	0.72	0.26
3	418.36	0.67	0.00	6.57	14.75	408.18 (c)	0.54	0.00
3	355.06	0.00	0.11	0.93	3.73	350.21 (c)	0.10	0.00
3	7.25	0.64	0.37	0.11	0.91	0.00	0.11	0.00
3	13.39	0.00	0.90	0.23	1.22	0.54	0.05	0.00
Mean	198.52	0.33	0.35	1.96	5.15	189.73	0.20	0.00
s	218.85	0.38	0.40	3.09	6.52	220.05	0.23	0.00
4	39.33	73.73	1.20	17.21	12.2	7.50	1.13	0.09
4	2.03	14.97	0.00	0.41	1.62	0.00	0.00	0.00
4	6.49	4.33	0.00	0.77	4.15	0.00	0.00	0.00
4	3.15	2.02	0.00	0.32	1.94	0.00	0.00	0.00
Mean	12.75	23.76	0.30	4.68	4.98	1.88	0.28	0.02
s	17.82	33.79	0.60	8.36	4.94	3.75	0.57	0.05

(a) Site 1 = Southwest Ocean Disposal Site; Site 2 = North Reference Site; Site 3 = South Beach Site; Site 4 = Half Moon Bay Site.

(b) Other crustaceans were mostly crangon shrimp.

(c) Mostly sand dollars.

TABLE 2 . Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Southwest Ocean Disposal Site in April 1992

Taxa	Individual Trawls										Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4		Trawl 5		kg/ha	#/ha
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha		
English sole	4.4	30.3	14.2	93.8	3.8	22.7	4.4	40.5	4.9	34.9	6.3	44.4
Pacific tomcod	22.7	484.5	34.7	1003.8	4.5	136.2	14.1	453.1	0.9	25.0	15.4	420.5
Surf smelt	1.8	TNC (a)	13.1	TNC	3.0	TNC	12.1	TNC	0.2	129.7	6.1	NMC (b)
Gadids (unidentified)	0.0	0.0	0.0	0.0	<0.1	15.1	0.0	0.0	0.0	0.0	<0.1	3.0
C-O sole	0.0	0.0	0.0	0.0	0.0	0.0	<0.1	8.1	0.0	0.0	<0.1	1.6

(a) TNC = Too numerous to count.

(b) NMC = No mean calculated.

TABLE 3 . Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the North Reference Site in April 1992

Taxa	Individual Trawls										Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4		Trawl 5		kg/ha	#/ha
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha		
English sole	1.7	10.7	31.1	171.9	0.0	0.0	1.5	14.0	10.0	70.4	8.9	53.4
Pacific tomcod	1.6	64.0	137.8	2062.5	2.2	130.9	7.3	392.0	15.3	551.6	32.8	640.2
Surf smelt	0.6	287.9	6.6	2781.3	0.4	212.7	1.2	616.0	0.6	363.9	1.9	852.3
Starry flounder	0.0	0.0	0.0	0.0	0.0	0.0	1.8	7.0	0.0	0.0	0.4	1.4

TABLE 4. Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the South Beach Site in April 1992

Taxa	Individual Trawls								Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4		kg/ha	#/ha
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha		
Starry flounder	0.0	0.0	0.1	5.3	0.0	0.0	0.9	847.6	0.3	213.2
Surf smelt	0.0	0.0	0.0	0.0	0.4	255.9	0.0	0.0	0.1	64.0

TABLE 5. Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Hall Moon Bay Site in April 1992

[illegible]

TABLE 6 . Trawl Catch Data Summary (Excluding Dungeness Crab Catch) from the June 1992 Epibenthic Survey Near Grays Harbor, Washington (Results are presented in kg/ha.)

Site (a)	Biological Catch Wt.	Debris Wt.	Commercially Important Fish	Other Fish	Crustaceans (b)	Echinoderms	Mollusks	Other Invertebrates
1	38.29	0.00	5.99	12.00	1.25	0.00	2.86	2.57
1	52.77	0.00	14.35	10.38	1.08	0.08	1.54	8.08 (c)
1	89.56	0.00	31.39	32.22	0.67	5.56	1.67	14.00
1	164.80	0.00	32.03	24.00	0.00	2.20	5.40	4.70
1	111.00	13.33	26.58	91.67	0.67	8.33	2.33	0.00
Mean	91.28	2.67	22.07	34.05	0.73	3.23	2.76	5.87
s	50.22	5.96	11.45	33.43	0.48	3.64	1.57	5.42
2	10.13	122.50	0.00	1.38	0.06	0.00	8.75	0.00
2	133.82	0.00	10.55	0.36	0.00	0.00	2.18	0.00
2	36.86	3.33	2.93	1.00	0.29	0.00	0.67	39.05 (c)
2	57.27	13.08	7.54	23.08	0.88	0.00	2.50	33.08 (c)
2	50.63	0.00	11.19	4.06	0.63	0.00	0.78	8.75 (c)
Mean	57.74	27.78	6.44	5.98	0.37	0.00	2.98	16.18
s	46.21	53.22	4.86	9.66	0.38	0.00	3.33	18.62
3	5.06	0.00	0.21	0.24	0.00	0.00	0.03	0.00
3	45.81	0.04	21.88	15.00	0.04	2.00	0.81	0.08
3	6.00	0.00	16.06	23.75	0.31	2.06	2.00	0.00
3	49.75	0.00	20.34	11.88	0.03	2.13	1.44	0.00
Mean	26.66	0.01	14.62	12.72	0.10	1.55	1.07	0.02
s	24.45	0.02	9.92	9.72	0.14	1.03	0.85	0.04
4	6.81	230.77	0.42	5.92	0.46	0.00	0.00	0.00
4	26.22	413.33	0.00	2.22	1.06	0.00	0.00	0.00
4	16.44	287.11	0.00	10.00	4.44	2.00	0.00	0.00
4	14.75	22.60	0.05	11.00	3.7	0.00	0.00	0.00
Mean	16.06	238.45	0.12	7.29	2.42	0.50	0.00	0.00
s	7.97	162.89	0.20	4.03	1.95	1.00	0.00	0.00

(a) Site 1 = Southwest Ocean Disposal Site; Site 2 = North Reference Site; Site 3 = South Beach Site; Site 4 = Half Moon Bay Site.

(b) Other crustaceans were mostly crangon shrimp.

(c) Mostly jellyfish.

TABLE 7. Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Southwest Ocean Disposal Site in June 1992

Taxa	Individual Trawls										Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4		Trawl 5			
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha		
English sole	0.0	0.0	3.7	30.8	26.2	166.7	32.0	480.0	22.8	133.3	16.9	162.2
Pacific tomcod	0.3	57.1	0.3	46.2	1.4	277.8	0.0	0.0	1.8	183.3	0.8	112.9
Surf smelt	5.7	1592.9	10.4	2715.4	3.8	766.7	0.0	0.0	2.0	466.7	4.4	1108.3

TABLE 8. Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the North Reference Site in June 1992

Taxa	Individual Trawls										Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4		Trawl 5			
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha
English sole	0.0	0.0	1.1	9.1	0.7	9.5	4.1	23.1	0.0	0.0	1.2	8.3
Pacific tomcod	0.0	0.0	<0.1	18.2	0.3	71.4	0.7	176.9	8.1	543.8	1.8	162.1
Surf smelt	0.0	0.0	0.2	TNC (a)	2.0	504.8	2.8	550.0	3.1	862.5	1.6	479.3 (b)

(a) TNC = Too numerous to count.

(b) Site mean for surf smelt (#/ha) was calculated using trawls 1, 3, 4, 5 only.

TABLE 9. Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the South Beach Site in June (199:)

Taxa	Individual Trawls								Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4			
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha
English sole	0.0	0.0	0.7	15.4	1.9	43.8	2.5	87.5	1.3	36.7
Surf smelt	0.2	41.2	13.5	TNC (a)	8.1	TNC	11.9	TNC	8.4	NMC (b)
Pacific tomcod	<0.1	5.9	7.7	253.9	5.9	243.8	5.9	456.3	4.9	239.9
Lingcod	0.0	0.0	0.0	7.7	0.0	6.3	0.0	0.0	0.0	3.5
C-O sole	0.0	0.0	0.0	0.0	<0.1	6.3	0.0	12.5	<0.1	4.7

(a) TNC = Too numerous to count.

(b) NMC = No mean calculated.

TABLE 10. Densities of Commercially Important Fish Taxa Captured in Beam Trawls at the Half Moon Bay Site in June 1992

Taxa	Individual Trawls								Site Mean	
	Trawl 1		Trawl 2		Trawl 3		Trawl 4			
	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha	kg/ha	#/ha
English sole	0.2	15.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.8
Starry flounder	0.3	7.7	0.0	0.0	0.0	0.0	0.0	0.0	0.1	1.9
Lingcod	0.0	0.0	0.0	0.0	0.0	0.0	0.1	10.0	0.0	2.5

TABLE 11. Grain Size Analysis (as Percentage Retained, Dry Weight) of Sediment Grab Samples Collected in April 1992 from the Grays Harbor Area

Screen Number	Sieve Size (mm)	South Beach Site					
		Station Designations (a)					
		<u>SJ-3</u>	<u>SJ-4</u>	<u>SJ-5</u>	<u>SJ-6</u>	<u>SJ-7</u>	<u>SJ-9</u>
#6	>3.34	0.00	1.13	0.46	0.00	0.00	2.49
#10	2.00-3.34	0.00	0.37	0.00	0.00	0.48	1.28
#18	1.00-2.00	0.26	0.14	0.35	0.12	0.52	0.82
#35	0.50-1.00	0.12	0.11	0.22	0.35	1.55	0.42
#60	0.25-0.50	1.15	1.62	2.53	2.47	5.44	3.81
#120	0.125-0.250	84.46	81.22	83.56	87.18	75.91	78.35
#230	0.063-0.125	14.02	15.42	12.88	9.87	16.10	12.83

Screen Number	Sieve Size (mm)	Half Moon Bay Site					
		Station Designations (a)					
		<u>HM-1</u>	<u>HM-2</u>	<u>HM-2(b)</u>	<u>HM-3</u>	<u>HM-4</u>	<u>HM-5</u>
#6	>3.34	0.00	0.00	0.00	0.00	0.00	0.00
#10	2.00-3.34	0.00	0.00	0.00	0.00	0.00	0.00
#18	1.00-2.00	0.00	0.00	0.00	0.78	0.00	0.00
#35	0.50-1.00	0.02	0.01	0.02	2.11	0.03	0.13
#60	0.25-0.50	4.46	0.74	0.68	9.05	0.64	3.40
#120	0.125-0.250	92.55	94.03	94.65	83.45	95.08	94.69
#230	0.063-0.125	2.97	5.22	4.66	4.62	4.25	1.78

(a) Station designations provided by USACE, Seattle District.

(b) Duplicate analysis of HM-2 sediment sample.

TABLE 12. Hydrographic Data from Sites Near Grays Harbor, Washington, Collected for the Dungeness Crab Survey Between April 21 and 24, 1992

<u>Site #</u>	<u>Date</u>	<u>Depth (m)</u>	<u>Salinity (ppt)</u>	<u>Temperature (°C)</u>
4	April 21	1.0	28.5	12.3
4	April 21	2.0	28.5	12.3
4	April 21	3.0	28.5	12.3
4	April 21	4.0	28.5	12.3
4	April 21	5.0	28.5	12.3
4	April 21	6.2	28.5	12.3
2	April 23	1.0	29.0	12.2
2	April 23	2.0	29.0	12.2
2	April 23	3.0	29.1	12.1
2	April 23	4.0	29.6	11.8
2	April 23	5.0	29.9	11.9
2	April 23	10.0	30.8	11.8
2	April 23	15.0(a)	31.5	11.8
5	April 24	1.0	27.4	12.7
5	April 24	2.0	27.4	12.7
5	April 24	3.0	27.5	12.7
5	April 24	4.0	27.5	12.7
5	April 24	5.0	27.5	12.6
5	April 24	10.0	27.5	12.6

(a) Depth of measurement limited by cable length on CTD unit.
Total depth was 37 m at Site 2.

TABLE 13. Hydrographic Data from Sites Near Grays Harbor, Washington, Collected for the Dungeness Crab Survey Between June 15 and 16, 1992

<u>Site #</u>	<u>Date</u>	<u>Depth (m)</u>	<u>Salinity (ppt)</u>	<u>Temperature (°C)</u>
2	June 15	0.0	28.8	15.0
2	June 15	1.0	28.8	15.0
2	June 15	2.0	28.8	15.0
2	June 15	3.0	28.8	14.9
2	June 15	4.0	28.8	14.7
2	June 15	5.0	29.0	14.4
2	June 15	10.0	30.1	13.4
2	June 15	15.0	30.9	13.2
2	June 15	20.0	32.3	10.5
2	June 15	25.0(a)	32.8	9.2
1	June 16	0.0	30.3	13.3
1	June 16	1.0	30.3	13.2
1	June 16	2.0	30.4	13.2
1	June 16	3.0	30.8	13.0
1	June 16	4.0	30.9	13.1
1	June 16	5.0	31.2	13.6
1	June 16	10.0	31.6	12.3
1	June 16	15.0	32.3	10.1
1	June 16	20.0	32.7	9.3
1	June 16	25.0(a)	32.6	9.4
4	June 16	1.0	30.4	13.7
4	June 16	2.0	30.4	13.6
4	June 16	3.0	30.5	13.5
4	June 16	4.0	30.4	13.5
4	June 16	5.0	30.5	13.4
4	June 16	10.0	30.5	13.4
4	June 16	15.0	30.5	13.4

(a) Depth of measurement limited by cable length on CTD unit. Total Depth was 29.3 m at Site 2, and 38.7 m at Site 1.

FIGURE 1a. Grain Size Analysis of Sediment Samples Collected from the South Beach Site near Grays Harbor, Washington, in April 1992 (Station designations were provided by USACE, Seattle District.)

FIGURE 1b. Grain Size Analysis of Sediment Samples Collected from the South Beach Site near Grays Harbor, Washington, in April 1992 (Station designations were provided by USACE, Seattle District.)

FIGURE 1c. Grain Size Analysis of Sediment Samples from the Half Moon Bay Site near Grays Harbor, Washington, in April 1992
(Station designations were provided by USACE, Seattle District.)

FIGURE 1d. Grain Size Analysis of Sediment Samples from the Half Moon Bay Site near Grays Harbor, Washington, in April 1992 (Station designations were provided by USACE, Seattle District.)

APPENDIX B

EPIBENTHIC SURVEYS IN 1990-1992 AT TWO PROPOSED BERM SITES NEAR GRAYS HARBOR, WASHINGTON

SUMMARY

To mitigate for shoreline erosion at the base of South Jetty at Grays Harbor Entrance, the U.S. Army Corps of Engineers, Seattle District, has proposed construction of sediment berms at South Beach and Half Moon Bay, off Point Chehalis. Sediments for the berms would be dredged from the Outer Bar and Entrance Reaches of Grays Harbor navigation channel.

Four epibenthic surveys were conducted between 1990 and 1992 to characterize the biological resources at the proposed berm sites. This work was completed in conjunction with biological monitoring at the Southwest Ocean Disposal Site. Survey efforts included epibenthic trawls with a plumb-staff beam trawl, grain-size analysis of surface sediments, and assessment of juvenile razor clam resources. Dungeness crab were identified by age class, enumerated, and measured (carapace width). Trawl catches of fish were grouped into round fish and flatfish, and commercially important fish were identified to species in 1990 and 1992. Because of the nature of the data set (i.e., limited number of trawls and high between-catch variability), a qualitative description of findings was appropriate.

In comparison to the Southwest Ocean Disposal Site and reference areas, no extraordinary aggregations of Dungeness crab, commercially important fish, or razor clams were found at either berm site. Epibenthic biota at the berm sites were typical for the Washington coast, with a few distinctive features. In general, Young-of-the-Year (YOY) Dungeness crab were more abundant outside Grays Harbor than in the harbor entrance (Half Moon Bay and the north side of the entrance channel). Older crab (1⁺ and adults) were found at higher densities in the harbor entrance than at sites outside Grays Harbor.

Surface sediments at the South Beach Site were dominated by fine sand. This site was also characterized by sand dollars and small gravel, both with a patchy distribution. The most abundant fish caught in trawls at South Beach were surf smelt.

Half Moon Bay also had fine sand as the dominant surface sediment layer. This site had the highest density of 1⁺ crab and the lowest density of YOY

crab found during the surveys. Woody debris was abundant in the trawls at Half Moon Bay. Very few commercially important fish were found at this site.

1.0 INTRODUCTION

The shoreline has been eroding at an alarming rate from Point Chehalis at the entrance to Grays Harbor (Figure B.1). In recent years, the beach has retreated more than 100 meters at Westport Light State Park on the western side of Point Chehalis. On the eastern side of the point, Half Moon Bay has formed and grown since the construction of the South Jetty in 1940. A continuation of these trends could lead to a breach of Point Chehalis at the base of South Jetty. In fact, waves from a severe winter storm in the winter of 1990 broke over the land at the base of South Jetty on Point Chehalis. A breach of Point Chehalis could result in a significant loss of land area and would affect entrance-channel maintenance, local navigation, and stability of the South Jetty.

The U.S. Army Corps of Engineers, Seattle District (USACE), proposed construction of sediment berms at South Beach and Half Moon Bay (Figure B.1) to slow erosional processes and prevent a breach at the base of the South Jetty. Sediment berms could mitigate these processes in two ways: 1) by providing a sediment supply for accretion on the shores and nearshore sacrificial material, and 2) by tripping larger waves further offshore from the beaches. The South Beach Site is located outside Grays Harbor approximately 1.7 nm (1.5 miles) south-southeast of Point Chehalis, between the 9.1- and 13.7-m (30- and 46-ft) depth contours. The Half Moon Bay Site is located along the south shore of Grays Harbor Entrance, at the base of South Jetty between the 4.9- and 7.3-m (16- and 24-ft) depth contours. Sediments for berm construction would be dredged from Grays Harbor Outer Bar Reach and Entrance Reach of the federal navigation channel.

Epibenthic surveys at the two sediment berm sites were conducted in 1990, 1991, and 1992, in conjunction with monitoring at the Southwest Ocean Disposal Site. Detailed survey results were reported for 1990 in Higgins and Pearson (1991) and for 1991 in Antrim et al. (1992). Survey reports for April and June of 1992 were letter reports to USACE and are included in Appendix A. During these surveys, beam trawls and sediment grab samples were taken for identification of Dungeness crab (June 1990, June 1991, April and June 1992),

FIGURE B.1. Grays Harbor Entrance Area and Locations of Proposed Berm Sites

commercially important fish (June 1990, April and June 1992), and razor clam resources (April 1992) at the berm sites. In addition, grain-size analyses of sediment grab samples were completed to characterize the surface sediments. This report summarizes findings from the four surveys at berm sites conducted by MSL. Trawl catches from berm sites are compared with catches at two offshore sites, the Southwest Ocean Disposal Site and North Reference area. Epibenthic biota at the berm sites were typical for the Washington coast, with a few distinctive features.

2.0 METHODS

In June of 1990, 1991, and 1992, and April 1992, epibenthic surveys at the berm sites were conducted in conjunction with biological monitoring at the Southwest Ocean Disposal Site. Trawls were taken using the methodology described in the main body of this report (Section 2.0). Survey efforts were augmented in 1992 to address concerns raised by regulatory agencies, primarily the Washington Department of Fisheries (WDF). The expanded scope of work in 1992 included identification of commercially important fish from trawl catches (April and June 1992) and sampling for juvenile razor clams at the South Beach Site (April 1992). The purpose of razor clam sampling was to determine if the density of juvenile razor clams was high at the South Beach Site. A reference site, WDF razor clam site "I", was sampled for comparison with South Beach. Sediment samples for juvenile razor clam analysis were taken with a Smith-McIntyre grab sampler, as recommended by WDF (personal communication, Doug Simons, WDF, Shellfish Program). The Smith-McIntyre grab effectively sampled the upper 3 to 4 cm of sediments, where juvenile razor clams would be found. No survey for razor clams was completed in June 1992 because plans to use the South Beach Site for berm construction were cancelled.

Characterization of surface sediments at the berm sites was completed in 1991 and 1992. A modified Van Veen grab sampler (0.1 m²) was used to collect surface sediments (upper 2 cm) from stations along the perimeter of and within the two berm sites. Grain-size analysis was conducted to determine percent composition of gravel, sand, silt, and clay in sediment samples using Method D-422 (ASTM 1972).

In 1991, two trawls were completed on the north side of Grays Harbor entrance as a reference site for comparison with Half Moon Bay. Bottom soundings conducted prior to trawling showed that the North Entrance area was characterized by irregular bottom topography, including 1.2- to 1.5-m (4- to 5-ft) high sand waves. These conditions restricted trawls to relatively smooth portions of this area.

The data were well suited to general observations on epifauna and bottom characteristics at the berm sites. Although statistical comparisons of trawl catches could have been made between berm sites and other sites surveyed,

these analyses were not completed for a number of reasons. The sampling effort at the berm sites was relatively limited and differed between years. For example, three surveys were completed at Half Moon Bay in 1991 and 1992, while four surveys were completed at the South Beach Site between 1990 and 1992. One survey at the South Beach Site included only two trawls (1990), whereas all other surveys included four trawls. Moreover, the variability in density estimates between individual trawls was relatively large, providing standard deviations (s) equal to or greater than means. As a result, this report provides a qualitative description rather than a quantitative analysis of the findings.

3.0 RESULTS

Trawl catches of Dungeness crab at the two berm sites are summarized in Table 3.1. Because each berm can be considered a unique project and study site, the following sections discuss the individual berm sites separately.

3.1 South Beach Berm Site

In surveys between 1990 and 1992, there was considerable variability in catch of Young-of-the-Year (YOY) Dungeness crab at the South Beach Site, yet catches of older Dungeness crab were relatively consistent. The mean density of YOY Dungeness crab ranged from 2 to 4,266 YOY crab/ha. For older age-classes of Dungeness crab, mean densities ranged from 0 to 118 1⁺ crab/ha and 8 to 37 adult crab/ha at this site. In June 1990, no 1⁺ crab were found during the entire survey that also included trawls at offshore sites (Southwest Ocean Disposal Site and the North Reference area). Dungeness crab distributions at the South Beach Site were more similar to offshore sites than Half Moon Bay.

Non-crab trawl catches at the South Beach Site were similar to catches at other areas sampled for these surveys. One unique biological feature was the abundance of sand dollars (Dendraster excentricus) at the South Beach Site. At some stations, surface sediment grabs were not obtainable because a dense cover of sand dollars prevented penetration by the grab sampler. Commercially-important fish taxa identified in 1992 included surf smelt, starry flounder, English and C-O sole, Pacific tomcod, and lingcod. Surf smelt were the most abundant fish caught in the trawls.

No razor clams were found in 12 grab samples taken in April 1992 at the South Beach Site between the 9.1-m and 13.7-m (30-ft and 45-ft) contours. The WDF Site "I" was also devoid of juvenile razor clams in April 1992. Because razor clam larval settlement typically occurs in May through July on Washington beaches (Lassuy and Simons 1989), the April 1992 survey probably preceded larval settlement.

Surface sediments were consistently dominated by fine sand ($\geq 80\%$ by weight). Some trawls, however, contained large amounts of small gravel.

TABLE 3.1. Summary of Dungeness Crab Densities (number/ha) from Beam Trawl Surveys at South Beach and Half Moon Bay Berm Sites, June 1990 - June 1992

Date	YOY Crab Density (number/ha)			
	South Beach		Half Moon Bay	
	Mean	s	Mean	s
June 1990	3,275	1,902	ND ^(a)	ND
June 1991	4,266	4,207	2	4
April 1992	216	206	333	537
June 1992	2	3	63	73

Date	1 ⁺ Crab Density (number/ha)			
	South Beach		Half Moon Bay	
	Mean	s	Mean	s
June 1990	0	0	ND	ND
June 1991	3	3	164	57
April 1992	118	221	37	51
June 1992	6	11	876	81

Date	Adult Crab Density (number/ha)			
	South Beach		Half Moon Bay	
	Mean	s	Mean	s
June 1990	31	9	ND	ND
June 1991	19	13	31	16
April 1992	37	12	45	18
June 1992	8	15	79	47

(a) ND = No Data.

These findings indicate that the bottom is relatively homogeneous sand with patches or bands of sand dollars and gravel throughout the area.

3.2 Half Moon Bay Berm Site

Trawls at Half Moon Bay were completed in 1991 and 1992 only. The abundance of Dungeness crab from each age-class was comparable to that found at other sites surveyed by MSL. An exception was 1⁺ crab, which were generally more abundant at Half Moon Bay. In 1991 and 1992, their mean density ranging from 37 to 876 1⁺ crab/ha at Half Moon Bay and from 0 to 118 1⁺ crab/ha at all other sites. The mean density of YOY Dungeness crab ranged from 2 to 333 YOY crab/ha, with the highest density found in April 1992. The mean density of adult crab at Half Moon Bay ranged from 31 to 79 adults/ha. In June 1991, megalopal larvae were seen floating in dense concentrations near the water surface. While megalopal larvae constituted 71% of the total Dungeness crab catch at this site in 1991, these larvae in trawl catches could have been collected from the water column as well as from the sediment surface.

Because sampling at the North Entrance area was limited to two trawls in 1991, comparisons of this data with catches at other sites have little scientific rigor. In general, this area had more YOY and adult Dungeness crab and fewer megalopae and 1⁺ Dungeness crab than Half Moon Bay.

Trawl catches of commercially important fish were lower at Half Moon Bay than at other sites surveyed in 1992. Catches were limited to a few individuals of each species and consisted of English and C-O sole, starry flounder, Pacific herring, and lingcod.

Surface sediments were dominated by fine sand over the entire area sampled at Half Moon Bay. In general, the trawl catch of debris was higher from the shallower portions of Half Moon Bay than at other sites surveyed. Debris was predominantly wood chips and small sticks.

4.0 DISCUSSION AND CONCLUSIONS

No unique biological features were identified in epibenthic surveys at the proposed berm sites. In general, the density of YOY Dungeness crab was higher outside Grays Harbor than inside the harbor entrance (Half Moon Bay and North Entrance Sites). However, sites in the harbor entrance had a better representation of 1⁺ and adult age class Dungeness crab than sites outside Grays Harbor. Surveys at the berm sites confirmed several characteristics of trawl catches discussed in the main body of this report including annual variability in YOY crab settlement and density, high inter-trawl variability in catches, and higher estuarine density of 1⁺ crab relative to areas outside the estuary.

5.0 REFERENCES

ASTM (American Society of Testing and Materials). 1972. Standard Method for Particle-Size Analysis of Soils. D-422, American Society of Testing and Materials, Philadelphia.

Antrim, L. D., V. I. Cullinan, and W. H. Pearson. 1992. Dungeness Crab Survey for the Southwest Ocean Disposal Site and Additional Sites off Grays Harbor, Washington, June 1991. PNL-7932, prepared for U.S. Army Corps of Engineers - Seattle District by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory, Richland, Washington.

Higgins, B. J., and W. H. Pearson. 1991. Dungeness Crab Survey for the Southwest Ocean Disposal Site off Grays Harbor, Washington, June 1990. PNL-5045, prepared for U.S. Army Corps of Engineers - Seattle District by Battelle/Marine Sciences Laboratory, Pacific Northwest Laboratory, Richland, Washington.

Lassuy, D. R., and D. Simons. 1989. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Northwest) -- Pacific Razor Clam. U.S. Fish and Wildlife Service Biol. Rep. 82(11.89). U.S. Army Corps of Engineers, TR-EL-82-4.

DISTRIBUTION

No. of
Copies

No. of
Copies

OFFSITE

Routing

2 DOE/Office of Scientific and
Technical Information

10 Bert Brun
U.S. Army Corps of Engineers
4735 E. Marginal Way South
Seattle, WA 98134

M. J. Graham
C. J. Hostetler
P. M. Irving
R. L. Skaggs
C. S. Sloane
P. C. Hays (last)

ONSITE

18 Pacific Northwest Laboratory

L. D. Antrim (5)
R. W. Bienert
V. I. Cullinan
R. M. Ecker
B. J. Higgins
W. H. Pearson
D. K. Shreffler
Publishing Coordination
Technical Report Files (5)

END

DATE
FILMED

4 / 30 / 93