2.

DOE/PC/91295--T4
OCT 05 1992

DE-FG22-91PC91295

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

A NOVEL, INTEGRATED TREATMENT SYSTEM FOR COAL
WASTEWATERS:

QUARTERLY REPORT:

FOR THE PERIOD June 2, 1992 - September 1, 1992.

DOE/PC/91295--T4

DE93 004928

PIs:

Prof. Henry Y. Wang
Department of Chemical Engineering
The University of Michigan
Ann Arbor, MI 48109-2136.

&

Dr. Keeran R. Srinivasan
Assoc. Research Scientist
Department of Chemical Engineering
The University of Michigan
Ann Arbor, MI 48109-2136.

Submitted To:

Pittsburgh Energy Technology Center
U. S. Department of Energy
Pittsburgh, PA.

ABSTRACT:

The aims of this study are to develop, characterize and optimize a novel treatment scheme that would be effective simultaneously against the toxic organics and heavy metals present in coal conversion wastewaters. Hec-CBDA-DT, a modified hectorite containing a mixed bilayer of a cationic (CBDA) and a diamine (DT) type surfactants, was shown to adsorb simultaneously heavy metals such as Cu^{2+} and Cd^{2+} and toxic organics such as chlorophenols. The pH dependence of the adsorption of chlorophenols on Hec-CBDA-DT suggested hydrophobic partitioning, counterion adsorption and hydrogen-bond interaction of the phenolate ions with the surface diamine groups as the possible mechanisms of sorption. Adsorption of Cu^{2+} by Hec-CBDA-DT was shown to be only slightly affected by a large excess of counterions, thus indicating that Cu^{2+} adsorbed specifically to the surface. A second type of modified clay, Hec-CBDA-Palmitic Acid (Hec-CBDA-PA) adsorbed both Cu^{2+} and Cd^{2+} strongly. However, the pH optima for adsorption were 6.0 and 8.0 respectively in the case of Cu^{2+} and Cd^{2+} . It is inferred that the mechanisms of adsorption of these two metal ions to Hec-CBDA-PA are different, specific and counterion binding respectively in the case of Cu and Cd ions.

MASTER

I. INTRODUCTION:

In our previous reports (1,2,3), we showed that, a number of clay complexes prepared by treating a smectitic clay such as hectorite sequentially with a cationic surfactant, CBDA, and then with a diamine or carboxylic acid type surfactant, were quite effective in adsorbing heavy metals such as Cu^{2+} and Cd^{2+} . A second approach was to directly attach the protonated form of the diamine to the negative sites of the clay layer to form a bilayer. More significantly, our previous work revealed that the pH of the continuous phase could be used as a "switch" to adsorb/desorb metal ions from such clay complexes. Four different complexes have been prepared and these are:

- (1) Hectorite-CBDA-Duomeen T: (Hec-CBDA-DT, the diamine type)
- (2) Hectorite-CBDA-Palmitic Acid: (Hec-CBDA-PA, the carboxylic acid type)
- (3) Montmorillonite-Duomeen T: (Mont-DT, the diamine type in which the functional moiety is directly attached to the clay surface)
- (4) Montmorillonite-CBDA-DT: (Mont-CBDA-DT, similar to Hec-CBDA-DT).

It was shown that Hec-CBDA-DT and Hec-CBDA-PA were the most effective adsorbents for Cu^{2+} ions with a pH optima of 7.0 and 6.0 respectively (3).

The main objective of our research, an integrated treatment for coal conversion wastewaters, requires a SINGLE adsorbent capable of adsorbing heavy metals reversibly and acting as a sink for the toxic organics in wastewaters. After metal ion recovery, the organic-laden adsorbent would be used as the source material in a biodegradation process for the mineralization of the adsorbed organics. In the first part of this report we have examined the adsorption characteristics of chlorinated phenols (as surrogates of hydroxy phenols in coal wastewaters) onto Hec-CBDA-DT in presence and in the absence of Cu^{2+} ions. The second part of the report deals with the effect of ionic strength on the adsorption of Cu^{2+} onto Hec-CBDA-DT at pH 7.0 to verify that ionic strength has a minimal effect. In the third part, we compare the effectiveness of Duomac-T, (DuT), a water-soluble analog of DT, when used to prepare a clay complex analogous to Hec-CBDA-DT. In the final part, the ability of Hec-CBDA-PA to strongly adsorb Cd^{2+} is demonstrated. It is shown that the optimum pH for Cd^{2+} adsorption is 8.0 in comparison to an optimal pH value of 6.0 for Cu^{2+} adsorption.

II. Adsorption of Chlorophenols:

3,5-dichlorophenol (DCP) and 2,4,5-trichlorophenol (TCP) and Hec-CBDA-DT were chosen as the adsorbates and the adsorbent respectively for this study. Table 1 is a summary of the properties of the two target molecules (4);

Table 1: Physicochemical Properties of DCP and TCP:

Property	DCP	TCP
Water Solubility (g/kg)	4.5	0.75
pKa	8.15	7.75
Absorption Max. (nm)	277, 284	297

Fig. 1 Adsorption of DCP on Hec-CBDA-DT:
pH Effect

Fig. 2 Adsorption of DCP on Hec-CBDA-DT at pH 7.0:
The Effect of Cu ions

Stock solutions of DCP (3,902 mg/L; 23.9 mM) and (504.6 mg/L; 2.56 mM) were prepared by dispersing and dissolving the respective compounds in deionized water. Adsorption studies were conducted in a batch mode at 25°C at a constant adsorbent concentration of 1 mg/ml. As in the case of metal uptake studies, pH was maintained constant at the selected pH using 20mM buffers. A stock adsorbent slurry was prepared at the pH of interest and sonicated for 10 minutes to insure complete dispersion. Controls were run concurrently to account for losses of adsorbates by processes other than sorption. An overall mass balance of greater than 90% was obtained in all cases. Initial DCP and TCP concentrations ranged respectively from 0.16 - 7.17 and 0.026 - 0.768 mM.

After 24 hours of well-mixed adsorption, phases were separated by centrifugation. DCP and TCP concentrations were measured by UV absorbance at 277 and 297 respectively. In the case of DCP, it was noted that absorbance at 284 nm was more reproducible at pH 8.0 and 9.0 and this wavelength was used at these pH. Separate calibration curves were obtained for each pH. When Cu^{2+} ions were added to the adsorption mixture, a new calibration curve in presence of Cu^{2+} ions was used to quantify adsorption data. The following experiments were undertaken:

- (1) DCP adsorption on Hec-CBDA-DT as a function of pH (Fig. 1).
- (2) The effect of Cu^{2+} ions on DCP adsorption at pH 7.0 (Fig. 2).
- (3) pH dependence of TCP adsorption to Hec-CBDA-DT (Fig. 3).
- (4) The role of Cu^{2+} on TCP adsorption at pH 7.0 (Fig. 4).

DCP Adsorption:

The pH dependence of DCP adsorption is shown in Fig. 1. At pH 3.0, DCP is essentially uncharged, while its degree of dissociation increases from 0.066 at pH 7.0 to 0.414 at pH 8.0 and to 0.88 at pH 9.0. Adsorption of DCP can be attributed to three different causes:

- (A) The neutral moiety of DCP can be expected to partition into the hydrophobic regions of Hec-CBDA-DT.
- (B) The phenolate form would bind to the protonated diamine headgroups as counterions.
- (C) Hydrogen-bonded or dipolar interactions between the phenolate anion and the uncharged diamine groups on the surface can occur.

The increase in the amount of DCP adsorption with increasing pH, as seen in Fig. 1, indicates that all the three modes are operational as the pH is increased. However, the initial part of the isotherm is identical at all pH indicating that hydrophobic partitioning is a dominant mechanism. Secondly, the height of the adsorption plateau increases only slightly with pH. Thus, sorption as counterions is also likely. The ratio of the charged vs uncharged diamine groups on the surface decreases with pH. Thus, at pH 9.0 all the three modes of binding contribute to DCP adsorption. Thus, the highest levels of adsorption are observed at this pH.

Note, however, that the the degree of dissociation of DCP has increased by a factor of 13 in the pH range of 7.0 - 9.0. But the actual enhancement in adsorption in this pH range is only slight. This would suggest that hydrogen-bonded or dipolar type interaction between the phenolate ions and the uncharged component of the surface diamine groups plays only a minor role in DCP adsorption on Hec-CBDA-DT. Further work is indicated to resolve these issues..

Fig 3: Adsorption of TCP onto Hec-CBDA-DT:
pH Effect

Fig. 4 Adsorption of TCP on Hec-CBDA-DT at pH 7.0:
The Effect of Cu ions

DCP Adsorption in presence of Cu^{2+} ions:

Cu^{2+} ions at an initial concentration of 0.4 mM were added to examine their role in DCP adsorption. At pH 3.0 Cu^{2+} had no effect on DCP adsorption. This was expected since DCP is uncharged at pH 3.0 and Cu^{2+} does not bind to Hec-CBDA-DT at pH 3.0.

The results obtained at pH 7.0 are shown in Fig. 2. At low DCP concentrations, Cu^{2+} ions block DCP adsorption as evidenced by the offset along the x-axis. The isotherm in presence of Cu^{2+} lies below the one without Cu ions. We conclude that Cu^{2+} ions form a soluble complex with DCP and this is supported by the observation that DCP in turn inhibits Cu^{2+} adsorption at the same pH. These experiments and our earlier work indicate that Cu^{2+} adsorption onto Hec-CBDA-DT involves the formation of a surface complex. This tentative conclusion is further verified by the ionic strength effect discussed in a later section of this report.

TCP Adsorption: pH and Cu^{2+} ions Effect:

Results of TCP adsorption at pH 3.0 and 9.0 in the absence of Cu ions are shown in Fig. 3. The differences are much more dramatic than in the case of DCP. TCP is almost totally in the phenolate form at pH 9.0 (the degree of dissociation of 0.95). At pH 9.0, the positive charge on the adsorbent surface is probably low. Thus, it is postulated that TCP adsorption at pH 9.0 proceeds via all the three adsorption mechanisms indicated above. However, a few differences between TCP and DCP adsorption were noted.

It is interesting that, as shown in Fig. 4, the strongest adsorption of TCP occurs at pH 7.0. At this pH, the surface will have more positive charges than at pH 9.0, and the degree of dissociation of TCP is 0.16. Therefore, counterion binding and hydrophobic sorption will be the two dominant mechanisms at pH 7.0. This probably explains why the affinity of TCP for Hec-CBDA-DT is the highest at this pH.

All the above results indicate that TCP and DCP adsorption is a complicated process involving hydrophobic bonding, dipolar interactions and counterion adsorption. The relative contributions of these mechanisms to the overall binding process will be pH dependent.

Earlier work on chlorophenol adsorption on surfactant-modified clays, i.e organo-clays, revealed no pH effect (5, 6). But, Alumina or hydroxy-aluminum-pillared clays exhibited a strong pH dependence of chlorophenol adsorption (4). Since Hec-CBDA-DT combines the hydrophobicity of a surfactant-modified clay and a net, pH dependent, positive charge of a hydroxy-aluminum-pillared clay, it is not surprising that the sorption properties of chlorophenols is a combination of different binding mechanisms.

In presence of Cu^{2+} , TCP adsorption is inhibited (Fig. 4). The extent of Cu^{2+} adsorption is also lowered. Thus, it is surmised that a soluble, Cu^{2+} -TCP complex is formed in solution which inhibits the adsorption of both components.

Finally, it is worth noting that UV absorbance of DCP and TCP is modified by Cu^{2+} ions in solution.

III. Adsorption of Cu^{2+} on Hec-CBDA-DT: Effect of Ionic Strength:

The above experiments were undertaken to study the stability of Cu^{2+} -DT interaction in presence of increasing concentrations of 1:1 electrolyte. Ionic strength was varied by adding NaCl at the

Fig. 6 Adsorption of Cu on Hec-CBDA-DuT:
pH Effect

Fig. 5 Adsorption of Cu on Hec-CBDA-DT at pH 7.0:
Ionic Strength Effect

same pH. An electrostatically-mediated adsorption process will be greatly modified by an excess of other counterions in solution. However, as shown in Fig. 5, Cu^{2+} adsorption is unaffected upto an ionic strength of 50 mM and only slightly modified at an ionic strength of 200 mM. The highest concentration of Cu^{2+} used for this experiment was 0.63 mM. These results confirm our earlier suggestions that specific interactions between Cu^{2+} and DT are responsible for the uptake of Cu^{2+} ions even at a pH (7.0), a pH at which the surface may still be positively charged (refer to the earlier sections of this report).

IV. Experiments with Hec-CBDA-Duomac-T (Hec-CBDA-DuT):

It was postulated that the effectiveness of DT in adsorbing Cu^{2+} or Cd^{2+} was related to the manner in which this active component was anchored to the CBDA-modified-hectorite. As described in our earlier report (3), DT attaches strongly to CBDA-hectorite through its hydrocarbon tail, thus exposing its diamine functionality to the solid-solution interface. Further, its insolubility in water minimizes its loss from the surface through leaching in aqueous media.

To verify the above hypothesis, a water-soluble analog of DT, Duomac-T (DuT), was obtained from Akzo Chemicals (7) and Hec-CBDA-DuT was prepared by treating CBDA-modified hectorite with an aqueous solution of Duomac-T (DuT). The preparation scheme was identical to the one used for Hec-CBDA-DT. A comparison of the two adsorbents is shown in Table 2.

Table 2. Comparison of Hec-CBDA-DT and Hec-CBDA-DuT:

Property	Hec-CBDA-DT	Hec-CBDA-DuT
% Carbon	39.3	25.1
%Nitrogen	3.45	1.93
%Hydrogen	6.91	4.35
Amino groups (mmoles/g)	0.65-0.71	0.16

The results shown in Table 2 reveal that DuT did not bind strongly to Hectorite-CBDA surface. This is reflected in the data shown in Fig. 6.

Adsorption of Cu^{2+} onto Hec-CBDA-DuT was monitored as a function of pH using an initial Cu^{2+} concentration of 0.32 mM. Control experiments were concurrently carried out to account for nonspecific effects. The results (shown in Fig. 6) are plotted in the form of a ratio of the residual to the initial Cu^{2+} concentrations as a function of pH.

It can be seen that Hec-CBDA-DuT has very little affinity for Cu^{2+} at any pH. There is no discernible difference between controls and the adsorbent. It is possible that adsorbed DuT was leached by successive (a total of 6) washings by deionized water or DuT may have displaced CBDA from the clay surface because of its positive charge.

On the basis of the above experiments and our earlier work, it may be concluded that water-insoluble, easily ionizable surfactants with a long hydrocarbon tails are the most optimal coatings that can render the resultant clays potent adsorbents of toxic organics and heavy metals.

Fig. 8 Adsorption of Cd or Cu on Hec-CBDA-PA
at the pH Optimum

Fig. 7 Adsorption of Cd on Hec-CBDA-PA:
pH Effect

V Cd^{2+} adsorption onto Hec-CBDA-PA:

In our earlier work (3), we reported on the preparation of Hec-CBDA-PA and the pH optimization of Cu^{2+} adsorption on this adsorbent. It was shown, based on a single-point isotherm, that Cu^{2+} adsorption was the highest at pH 6.0. In the present report, results from a similar optimization study for Cd^{2+} adsorption are reported. Further, complete isotherms of Cu^{2+} and Cd^{2+} adsorption at their respective optimal pH are also presented and discussed.

Cd^{2+} adsorption on Hec-CBDA-PA was monitored as a function of pH at an initial metal ion concentration of 0.18 mM. Appropriate controls were included to validate the results and these are displayed in Fig. 7.

The gradient of adsorption affinity vs pH changes dramatically near pH 7.0, and at pH 8.0 and above, the uptake of Cd ions from solution is total. It is noteworthy that the pH optima for Cu^{2+} and Cd^{2+} are respectively 6.0 and 8.0. At the higher pH, the negative surface charge of Hec-CBDA-PA is expected to be much higher than at pH 6.0. This would indicate that Cd^{2+} may be adsorbing as counterions while Cu^{2+} adsorption is probably similar to the formation of an inner sphere complex. Ionic strength studies in conjunction with spectroscopic measurements will help resolve these differences.

The full isotherms of Cu^{2+} and Cd^{2+} adsorption on Hec-CBDA-PA at pH 6.0 and 8.0 respectively are shown in Fig. 8. Adsorption of Cd^{2+} appears to be of the high affinity type and the surface sites are not fully covered even at the highest initial concentrations used. On the other hand, Cu^{2+} adsorption is seen to level off even though this isotherm is also of the high affinity type.

VI. Conclusions:

- (1) Surfactant-modified clays such as Hec-CBDA-DT have been shown to adsorb simultaneously heavy metals and chlorophenols.
- (2) Metal ions such as Cu^{2+} complex with the anionic forms of chlorophenols and this results in mutual inhibition of adsorption.
- (3) The specific nature of Cu^{2+} adsorption on Hec-CBDA-DT was inferred from a very small effect produced by an excess of counterions in solution.
- (4) The differences in pH optimum of Cu (pH 6.0) and Cd (pH 8.0) ion adsorption on Hec-CBDA-PA suggest respectively specific and electrostatic types of interactions between these two metal ions and the adsorbent.

VII. References:

- (1) Henry Y. Wang and Keeran R. Srinivasan: I Quarterly Report, DOE/PETC, Dec. 1991.
- (2) Henry Y. Wang and Keeran R. Srinivasan: II Quarterly Report, DOE/PETC, March, 1992.
- (3) Henry Y. Wang and Keeran R. Srinivasan: III Quarterly Report, DOE/PETC, June, 1992.
- (4) Richard C. Zielke and Thomas J. Pinnavaia: Clays and Clay Minerals, 36(5), 403 (1988).

- (5) M. M. Mortland, S. Shaobai and S. A. Boyd: Clays and Clay Minerals 34, 581 (1986).
- (6) Keeran R. Srinivasan and H. Scott Fogler: " Use of Inorgano-organo-Clays in the Treatment of Coal Conversion Wastewaters " Proc. 1990 International Symposium on Dioxin and Other Pollutants, held at Bayereuth, Fed. Republic of Germany, September, 1990.
- (7) Akzo Chemie America: Bulletin 85-1, "Specification and Properties of DUOMEEN Diamines and Diamine Salts", An Aramak Chemical Publication, (1985).

**DATE
FILMED
02/02/93**