EGG-MS-10535
October 1992

**Idaho
National
Engineering
Laboratory**

*Managed
by the U.S.
Department
of Energy*

INFORMAL REPORT

**Galileo's Relativity Principle, the
Concept of Pressure, and Complex
Characteristics, for the Six-Equation,
One-Pressure Model**

Henry Makowitz

*Work performed under
DOE Contract
No. DE-AC07-76ID01570*

EGG-MS-10535
OCT 13 1992

ABSTRACT

We have studied various formulations of the concept of pressure, in the context of the usual Six-Equation Model of thermal-hydraulics. A different concept of pressure, than the usual one, has been used. This new pressure concept is Galilean Invariant, and results for the One-Pressure Model with the same complex characteristic roots as the "Basic Ill-Posed Model," discussed in the literature for the cases we have investigated. We have also examined several Two-Pressure formulations and shown that two pressures are a necessary but not sufficient condition for obtaining a Well-Posed system. Several counter examples are presented. We have shown that the standard theory is *not* Galilean Invariant and suggested that the origin of Ill-Posedness is due to our closure relationships. We also question whether the current theory can satisfy conservation principles for mass, energy, and momentum.

RECEIVED

EP

INTRODUCTION

We have studied a number of formulations of the Six-Equation, One-Pressure Model^(1,2) utilizing a different concept of pressure than the usual one. Our interest was to examine the resulting characteristics of the "Basic Ill-Posed Model" discussed by Wendroff,⁽²⁾ and study their behavior as a function of our "Galilean Invariant" pressure concept.

We show that for the Six-Equation, One-Pressure Model, the complex part of the complex characteristics are Galilean Invariant and remain complex as long as a Single-Pressure Model is used and the two fluid velocities are not equal. We examine several two-pressure formulations and show that they do not result in a Well-Posed model in general. However, certain specific cases are Well-Posed. We show that the standard theory is *not* Galilean Invariant and discuss the origin of Ill-Posedness in the context of an invariant pressure and sound speed. We also question whether the current theory can satisfy fundamental conservation principles for mass, energy, and momentum

The Concept of Pressure and Galileo's Relativity Principle

L. D. Landau,⁽³⁾ in the context of a One-Pressure, Two-Fluid, Two-Velocity theory of Superfluid Dynamics, derived a pressure, P , composed of a thermodynamic component and a velocity component, given below by Equation (1),

where

E_0 = internal energy

ρ = total density

ρ_n = normal fluid density

s = specific entropy

T = temperature

μ = chemical potential

$\underline{v}_n$ = normal fluid velocity

$\underline{v}_s$ = superfluid velocity.

$$P = -E_o + T\varrho_s + \mu\varrho + \varrho_n (\underline{v}_n - \underline{v}_s)^2 . \quad (1)$$

We rewrite this as the usual pressure concept P_* , and the velocity component, via Equation (2), where ϱ_{**} is some function of densities 1 and 2,

$$P = P_* + \varrho_{**}(\underline{v}_1 - \underline{v}_2)^2 . \quad (2)$$

Landau⁽³⁾ argued that

"The usual thermodynamic definition of the pressure as the mean force acting on unit area relates to a medium at rest. In ordinary fluid dynamics, however, there is no ambiguity in the definition of pressure (dissipative processes being neglected), since we can always take a coordinate system in which the fluid volume element considered is at rest. In superfluid dynamics, however, we can eliminate only one of the two simultaneous motions by a suitable choice of the coordinate system, and so the usual definition of pressure cannot be applied.

"In a fluid entirely at rest, the definition [Equation (1)] is of course the same as the ordinary definition, since in that case $\mu\varrho + T\varrho_s - E_o = P$ by the definition of the chemical potential."

Although Landau's formulation of the pressure is specific to his Two-Fluid, Two-Velocity, Super Fluid Dynamics Model, we argue that the form of Equation (2) is quite general, and may be applicable to the Six-Equation, One-Pressure Model of thermal-hydraulics. An argument supporting this hypothesis is presented below.

Consider, as per Figure 1, three reference frames, r_0 , r_1 , and r_2 , with r_0 at rest relative to an observer, r_1 moving with constant velocity $\underline{v}_1$, and r_2 moving with constant velocity $\underline{v}_2$ relative to r_0 . If identical tanks filled with a liquid are placed in each reference frame, and a sound wave is initiated in each tank at each frame's origin, (assuming that the magnitude of the sound speed C_s is the

Figure 1. Reference frame geometry.

same in each frame), the observer in r_0 will observe that, in reference frame r_0 , the observed sound speed $C_{s(i)}$ for frame "i" is given by Equation (3),

$$C_{s(0)} = C_s \quad ; \quad (3a)$$

$$C_{s(1)} = C_s + |\underline{v}_1| \quad ; \quad (3b)$$

and

$$C_{s(2)} = C_s + |\underline{v}_2| \quad . \quad (3c)$$

The difference measured in frame r_0 of the observed sound speeds in frames r_1 and r_2 is given by Equation (4),

$$\Delta C_{s(1,2)} = C_{s(1)} - C_{s(2)} = (|\underline{v}_1| - |\underline{v}_2|) \quad . \quad (4)$$

If we place the observer in frame r_1 , then

$$C_{s(1)} = C_s \quad ; \quad (5a)$$

$$C_{s(2)} = C_s + (|\underline{v}_2| - |\underline{v}_1|) \quad ; \quad (5b)$$

and

$$\Delta C_{s(1,2)} = C_{s(1)} - C_{s(2)} = (|\underline{v}_1| - |\underline{v}_2|) \quad . \quad (5c)$$

If for simplicity we assume that the liquids in the tanks behave like an "isothermal ideal" gas, then the isothermal sound speed $C_s(\text{ISO})^{(4)}$ is given by

$$C_s(\text{ISO}) = \left(\sqrt{\frac{P}{\rho}} \right)_{\text{ISOTHERMAL}} \quad . \quad (6)$$

If we assume that the density ρ is constant and frame invariant, then, if $C_s(\text{ISO}) \equiv C_s$, we can define P_{INV} , ΔP , and C_{INV} such that

$$\Delta C_{s(1,2)} \equiv \sqrt{\frac{\Delta P}{\rho}} \quad , \quad (7)$$

where the subscript "INV" indicates a frame invariant quantity. (Refer to Appendix A for a motivation of P_{INV} and C_{INV} .)

If we consider One-Dimensional Motion (i.e., $\underline{v}_1 \parallel \underline{v}_2$), then, in frame r_0 ,

$$\Delta P_{(0)} = \rho_{(0)}(v_1 - v_2)^2 ; \quad (8a)$$

and in frame r_1 ,

$$\Delta P_{(1)} = \rho_{(1)}(v_1 - v_2)^2 , \quad (8b)$$

and since $\rho_{(1)} = \rho_{(0)} = \rho$ by definition, we see that $\Delta P_{(i)}$ is frame invariant if it is of the form given by Equation (8). This suggests that a general form for the pressure is that given by Equation (2). However, it should be noted that this definition is not unique. Although a detailed formal analysis is required for the general case of the Six-Equation Model of thermal-hydraulics, as per Reference 3 for superfluid dynamics, for now let us assume that the pressure is of the form given by Equation (2), and study the resulting characteristics.

ANALYSIS

We are interested in the characteristics of the simple system^(1,2,5) given by Equation (9),

$$\frac{\partial(\alpha \rho_v)}{\partial t} + \frac{\partial}{\partial x} (\alpha \rho_v U_v) = 0 ; \quad (9a)$$

$$\frac{\partial}{\partial t} [(1 - \alpha) \rho_l] + \frac{\partial}{\partial x} [(1 - \alpha) \rho_l U_l] = 0 ; \quad (9b)$$

$$(\alpha \rho_v) \left[\frac{\partial(U_v)}{\partial t} + (U_v) \frac{\partial}{\partial x} (U_v) \right] + (\alpha) \frac{\partial}{\partial x} (P_1) = 0 ; \quad (9c)$$

$$[(1 - \alpha) \rho_l] \left[\frac{\partial(U_l)}{\partial t} + (U_l) \frac{\partial}{\partial x} (U_l) \right] + (1 - \alpha) \frac{\partial}{\partial x} (P_2) = 0 ; \quad (9d)$$

and where the characteristic equation,⁽⁶⁾ Equation (10b),

$$\underline{A} \frac{\partial}{\partial t} (\zeta) + \underline{B} \frac{\partial}{\partial x} (\zeta) + \underline{C} = 0 , \quad (10a)$$

$$| -\tau \underline{A} + \underline{B} | = 0 \quad (10b)$$

needs to have real roots,^b for the system of Equation (10a) to be "Well-Posed."⁽⁶⁾ We have investigated the following cases:

$$\text{Case A1: } \frac{\partial}{\partial x} (q_l) = \frac{\partial}{\partial x} (q_v) = 0 ; P_1 = P_2 \equiv P. ; \zeta = \begin{bmatrix} \alpha \\ U_v \\ U_l \\ P. \end{bmatrix}$$

$$\text{Case A2: } \frac{\partial}{\partial x} (q_l) = \frac{\partial}{\partial x} (q_v) = 0 ; P = P_1 = P_2 \equiv P. + q_l U_v^2 ;$$

$$\frac{\partial}{\partial t} (U_l) \neq 0 ; \frac{\partial}{\partial x} (U_l) \neq 0 ; U_l = 0 \text{ (in "U}_l\text{" frame)} ; \zeta = \begin{bmatrix} \alpha \\ U_v \\ U_l \\ P \end{bmatrix} .$$

Case A3: $P = P_1 = P_2 \equiv P. + q_v U_v^2$; the rest is the same as A2.

Case A4: $P_1 \equiv P.$, for $U_l = 0$ Equation; $P_2 \equiv P. + q_l U_v^2$, for U_v Equation;
the rest is the same as A2, except $P_1 \neq P_2$.

Case A5: $P_1 \equiv P.$, for $U_l = 0$ Equation; $P_2 \equiv P. + q_v U_v^2$, for U_v Equation;
the rest is the same as A2, except $P_1 \neq P_2$.

$$\text{Case A6: } \frac{\partial}{\partial x} (q_l) = \frac{\partial}{\partial x} (q_v) = 0 ; P = P_1 = P_2 \equiv P. + q_v U_l^2 ;$$

$$\frac{\partial}{\partial t} (U_v) \neq 0 ; \frac{\partial}{\partial x} (U_v) \neq 0 ; U_v = 0 \text{ (in "U}_v\text{" frame)} ;$$

$$\zeta = \begin{bmatrix} \alpha \\ U_v \\ U_l \\ P \end{bmatrix} .$$

b. The reader should note that $| |$ in Equation (10b) implies a determinant.

Case A7: $P = P_1 = P_2 \equiv P_\bullet + \varrho_l U_l^2$; the rest is the same as A6 .

Case A8: $P_1 \equiv P_\bullet$, for $U_v = 0$ Equation; $P_2 \equiv P_\bullet + \varrho_l U_l^2$, for U_l Equation;
the rest is the same as A6, except $P_1 \neq P_2$.

Case A9: $P_1 \equiv P_\bullet$, for $U_v = 0$ Equation; $P_2 \equiv P_\bullet + \varrho_v U_v^2$, for U_l Equation;
the rest is the same as A6, except $P_1 \neq P_2$.

Case B1: $C_l^2 \rightarrow +\infty$ as $\frac{\partial}{\partial x} (\varrho_l) \rightarrow 0$; $P_1 = P_2 \equiv P_\bullet$;

$$\frac{\partial}{\partial x} (P_\bullet) = C_v^2 \frac{\partial}{\partial x} (\varrho_v) = C_l^2 \frac{\partial}{\partial x} (\varrho_l) ; \zeta = \begin{bmatrix} \alpha \\ \varrho_v \\ U_v \\ U_l \end{bmatrix} .$$

Case B2: $C_l^2 \rightarrow +\infty$ as $\frac{\partial}{\partial x} (\varrho_l) \rightarrow 0$; $P = P_1 = P_2 \equiv P_\bullet + \varrho_l U_v^2$;

$$\frac{\partial}{\partial x} (P) \equiv \frac{\partial(P_\bullet)}{\partial x} + \frac{\partial}{\partial x} (\varrho_l U_v^2) ; \frac{\partial}{\partial t} (U_l) \neq 0 ; \frac{\partial}{\partial x} (U_l) \neq 0 ;$$

$U_l = 0$ (in “ U_l ” frame); the rest is the same as B1, except $P \neq P_\bullet$.

Case B3: $P = P_1 = P_2 \equiv P_\bullet + \varrho_v U_l^2$; $\frac{\partial}{\partial x} (P) \equiv \frac{\partial}{\partial x} (P_\bullet) + \frac{\partial}{\partial x} (\varrho_v U_l^2)$;

$$\frac{\partial}{\partial t} (U_v) \neq 0 ; \frac{\partial}{\partial x} (U_v) \neq 0 ; U_v = 0 \text{ (in “} U_v \text{” frame)} ;$$

the rest is the same as B1, except $P \neq P_\bullet$.

Case C1: $P_1 = P_2 \equiv P_\bullet$; $\frac{\partial}{\partial x} (P_\bullet) = C_v^2 \frac{\partial}{\partial x} (\varrho_v) = C_l^2 \frac{\partial}{\partial x} (\varrho_l) ; \zeta = \begin{bmatrix} \alpha \\ \varrho_v \\ U_v \\ U_l \end{bmatrix} ;$

$$\frac{\partial}{\partial x} (U_l) \neq 0 ; \frac{\partial}{\partial t} (U_l) \neq 0 ; U_l = 0 \text{ (in “} U_l \text{” frame)} .$$

Case C2: $P_1 = P_2 \equiv P$; $P = P_\bullet + \rho_l U_V^2$;
the rest is the same as C1, except
 $P \neq P_\bullet$.

Case C3: $P_1 = P_\bullet$; $P_2 = P_\bullet + \rho_l U_V^2$;
the rest is the same as C1, except
 $P \neq P_\bullet$ and $P_1 \neq P_2$.

Case C4: $\frac{\partial}{\partial x} (P_{1.}) = C_V^2 \frac{\partial}{\partial x} (\rho_v)$;
 $\frac{\partial}{\partial x} (P_{2.}) = C_l^2 \frac{\partial}{\partial x} (\rho_l)$;
 $P_1 = P_{1.}$; $P_2 = P_{2.}$;
the rest is the same as C1, except
 $P_1 \neq P_2$; $\alpha = \text{cont.}$; and ζ redefined
in terms of ρ_v , ρ_l , U_v , U_l .

Case C5: $P_1(U_l = 0 \text{ frame}) = P_{1.}$;
 $P_2(U_v) = P_{2.} + \rho_l U_V^2$;
the rest is the same as C4 .

Case C6: $P_1(U_l = 0 \text{ frame}) = P_{1.} + \rho_l U_V^2$;
 $P_2(U_v) = P_{2.} + \rho_l U_V^2$;
the rest is the same as C4 .

Since we are only interested in the complex roots, we assume that $U_v \ll C_v$, $U_v \ll C_l$, $U_l \ll C_v$,
and $U_l \ll C_l$, as per Reference 5, and mostly ignore the real roots near $U_v \pm C_v$ and/or $U_l \pm C_l$.
To simplify the algebra we evaluate many of our cases in an instantaneous "Rest" frame (i.e.,
 $U_l = 0$ or $U_v = 0$). However, we do not assume constant velocities, that is, if

$$U_l = 0 \quad , \quad \frac{\partial}{\partial x} (U_l) \neq 0 \quad ,$$

and

$$\frac{\partial}{\partial t} (U_i) \neq 0 ,$$

and only assume a “slow” rate of change in the frame velocity. Cases A1 and B1 have the same complex roots given by Equation (11),

$$\tau = \frac{U_i + \varepsilon^2 U_v}{(1 + \varepsilon^2)} \pm \frac{i\varepsilon (U_i - U_v)}{(1 + \varepsilon^2)} \quad (11a)$$

and

$$\varepsilon^2 = \frac{(1 - \alpha) \varrho_v}{\alpha \varrho_i} , \quad (11b)$$

while Case C1 corresponds to Equation (11) with $U_i = 0$, since it is in the “ $U_i = 0$ ” frame. All One-Pressure cases where $P = P_1 = P_2 = P^* + \varrho_i U_j^2$ ($i = \iota$ or v , $j = \iota$ or v , and i may equal j) have complex roots of the form of Equation (11), where either $U_i = 0$ or $U_v = 0$, depending on our choice of “Rest” frame. Hence, when the pressure in the One-Pressure, Six-Equation Model is of the form given by Equation (2), the complex part of the characteristic roots are Galilean Invariant. We summarize all of our results for Case A in Table 1A, for Case B in Table 1B, and for Case C in Table 1C. For completeness, we show in Appendix A that the “Basic III-Posed Model” is **not** Galilean Invariant and discuss the origin of III-Posedness in the context of an invariant pressure and sound speed.

DISCUSSION

It is interesting to note that if we choose a pressure in the form of Equation (2), Galilean Invariance of the complex part of the characteristic equation's complex roots only occurs for the One-Pressure, Six-Equation Model. Additionally, for the cases that we have investigated, only the Two-Pressure Models of the form of Case C4 result in Well-Posed systems. Hence, two pressures do not automatically result in a Well-Posed system, contrary to many such observations in the literature. It should also be noted that Cases C5 and C6 do **not** result in Galilean Invariant roots, relative to Case C4, suggesting that pressures in the form of Equation (2) are incorrect for

Two-Pressure Model formulations. Finally, we observe that any frame invariant definition of the pressure, in the context of a One-Pressure Model, results in the loss of the correct Doppler shifts, since we have redefined the sound speed so that it is also frame invariant. (Refer to Appendix A for further discussion.)

Based on the study presented in the body of this paper, we can make several theoretical observations of a general nature. In Appendix A we have inferred that a connection exists between Ill-Posedness and the requirements of Galilean Invariance of the signal propagation velocity. Another way to view this point is in the context of a “Lorentz Invariant Force.”^(7,8,9) Since the Six-Equation, One-Pressure Model attempts to define a pressure and thus a force (pressure is defined as a force per unit area) that is invariant in two frames of reference—inertial or accelerating—simultaneously, and connects the signal propagation velocity (speed of sound) between these two frames, inadvertently, via the closure relationships, thus distorting the orthogonality of space and time (per Appendix A), we can ask whether mass, energy, and momentum are invariants in such a theory.

In the Special Theory of Relativity,^(7,8,9) Lorentz invariant quantities are four vectors, and are conserved in a four-dimensional space. Mass, for example, can appear to increase as a function of velocity, that is,

$$\text{mass} = \text{rest mass} \times \left[1 - \frac{U_O^2}{C^2} \right]^{-1/2}$$

for an observer at rest relative to velocity U_O . We show in Appendix B that if one imposes the condition that the scalar fields in the One-Pressure Theory are conserved in flat space, the Theory becomes Well-Posed.⁽¹⁰⁾ We show in Appendix C that the linearized One-Pressure Theory is not conservative.⁽¹¹⁾ At this time, to our knowledge, no conservation proofs exist for the quasi-linear or nonlinear One-Pressure Theory. However, a proof by Lax⁽¹²⁾ exists that the nonlinear theory is *not* bounded under special conditions. Lax also states that Energy conservation does not provide a bound for the One-Pressure Theory.

SUMMARY AND CONCLUSIONS

We have shown that the standard theory is *not* Galilean Invariant and suggested that the origin of Ill-Posedness is due to our closure relationships (refer to Appendix A). We have also questioned whether the current theory can satisfy conservation principles for mass, energy, and momentum.

We have studied various formulations of the concept of pressure, in the context of the usual Six-Equation Model^(1,2) of thermal-hydraulics. A different concept of pressure, than the usual one, has been used. This new pressure concept is Galilean Invariant, and produces the same results for the One-Pressure Model with the same complex characteristic roots as the "Basic Ill-Posed Model,"^(2,5) discussed in the literature for the cases we have investigated. We have also examined several Two-Pressure Formulations and shown that two pressures are a necessary but not sufficient condition for obtaining a Well-Posed system. Several counter examples were presented.

Table I. Summary of analysis.

A		B		C	
Case A1 (One-Pressure)	III-Posed; roots given by Equation (11)	Case B1 (One-Pressure)	III-Posed; roots given by Equation (11)	Case C1 (One-Pressure)	III-Posed; roots given by Equation (11) with $U_t = 0$
Case A2 (One-Pressure)	III-Posed; same complex roots as A1 with $U_t = 0$	Case B2 (One-Pressure)	III-Posed; same as B1 with $U_t = 0$	Case C2 (One-Pressure)	III-Posed; same as C1
Case A3 (One-Pressure)	III-Posed; same complex roots as A1 with $U_t = 0$	Case B3 (One-Pressure)	III-Posed; same as B1 with $U_v = 0$	Case C3 (Two-Pressure)	III-Posed, but reduced region versus C1
Case A4 (Two-Pressure)	III-Posed, but reduced region versus A1	—	—	Case C4 (Two-Pressure)	Well-Posed; $\tau_1 = U_v + C_v$ $\tau_2 = U_v - C_v$ $\tau_3 = + C_t$ $\tau_4 = - C_t$
Case A5 (Two-Pressure)	III-Posed, but reduced region versus A1	—	—	Case C5 (Two-Pressure)	Well-Posed; complicated expression for roots
Case A6 (One-Pressure)	III-Posed; same complex roots as A1 with $U_v = 0$	—	—	Case C6 (Two-Pressure)	Well-Posed; complicated expression for roots
Case A7 (One-Pressure)	III-Posed, same complex roots as A1 with $U_v = 0$	—	—	—	—
Case A8 (Two-Pressure)	III-Posed, but reduced region versus A1	—	—	—	—
Case A9 (Two-Pressure)	III-Posed, but reduced region versus A1	—	—	—	—

APPENDIX A

In this appendix, we show that Case C1 of the main text [i.e., the “Basic Ill-Posed Model”⁽²⁾] is not Galilean Invariant. We then motivate a Galilean Invariant Pressure concept, and discuss the origin of Ill-Posedness in the context of a Galilean Invariant sound speed.

FAILURE OF GALILEAN INVARIANCE FOR THE ILL-POSED MODEL

If we evaluate Case C1 in the lab frame for $\xi = \begin{bmatrix} P. \\ \alpha \\ U_v \\ U_l \end{bmatrix}$,

$C_l \gg U_l$, $C_v \gg U_v$, $\varepsilon^2 = 1$, and $C_l = C_v \equiv C_s$,

we obtain the following four roots:

$$\tau_{1,2} = \frac{U_l + U_v}{2} \pm C_s ; \quad (A.1.0)$$

$$\tau_{3,4} = \frac{U_l + U_v}{2} \pm \frac{i}{2} (U_v - U_l) . \quad (A.2.0)$$

For simplicity, let us say that $U_l = -U_v$ and $|U_l| = |U_v| \equiv U$. Then, in the lab frame,

$$\tau_{1,2} = \pm C_s \quad (A.3.1)$$

and

$$\tau_{3,4} = \pm iU . \quad (A.3.2)$$

In the $U_v = 0$ frame, $U_l = -2U$, and

$$\tau_{1,2} = -U \pm C_s \quad (A.4.1)$$

and

$$\tau_{3,4} = -U \pm iU \quad . \quad (A.4.2)$$

In the $U_x = 0$ frame, $U_y = 2U$, and

$$\tau_{1,2} = U \pm C_s \quad (A.5.1)$$

and

$$\tau_{3,4} = U \pm iU \quad . \quad (A.5.2)$$

Hence, only the complex part of the complex roots is Galilean Invariant. Both the real sonic roots and the real part of the complex roots are not Galilean Invariant. (Q.E.D.)

MOTIVATION FOR A GALILEAN INVARIANT PRESSURE CONCEPT FOR IDENTICAL SYSTEMS

Let us begin with Figure 1 and Equations (1) through (8) of our section entitled "The Concept of Pressure and Galileo's Relativity Principle" in the main text. If we define C_{INV} as

$$C_{INV} \equiv \sqrt{\frac{P_{INV}}{\rho_{INV}}} \quad , \quad (A.6.0)$$

then

$$C_{INV}^2 = \frac{P_s}{\rho_{INV}} + \frac{\Delta P}{\rho_{INV}} \quad , \quad (A.7.1)$$

or

$$C_{INV}^2 = C_s^2 + (v_1 - v_2)^2 \quad , \quad (A.7.2)$$

or

$$C_{INV} = C_s \sqrt{1 + \left(\frac{\Delta V}{C_s}\right)^2} \quad , \quad (A.7.3)$$

where C_* is the local sound speed in the observer's frame of reference, and we have considered identical systems as per Figure 1. Since $\Delta V = (v_1 - v_2)$ is Galilean Invariant and C_* is the same in all three inertial frames (i.e., identical systems per Figure 1). Equation (A.7.3) is Galilean Invariant. An alternative to the definition, Equations (A.6.0) and (A.7.3) can be the following; we can argue that

$$C_{*(1)} = C_{*(2)} = C_{*(3)} \quad (\text{A.8.0})$$

and just ignore ΔP and ΔC . However, we prefer the form given by Equations (A.6.0) and (A.7.3).

THE ORIGIN OF ILL-POSEDNESS

If we construct a Galilean Invariant, Pressure, P_{INV} , and Sound Speed, C_{INV} , we no longer are in a Euclidean space-time geometry and space and time are not orthogonal for such a system [i.e., Albert Einstein, "The Special Theory of Relativity,"⁽⁷⁾ 1905]. Galilean Invariance of the signal propagation velocity for our "Basic Ill-Posed Model" will result in a complex formulation [i.e., Minkowski Space⁽⁸⁾ for the Theory of Special Relativity].

We illustrate the reason for the origin of complex characteristics below. Let us consider two time lines, as viewed in the laboratory frame, of sound signals C_1 and C_2 associated with reference frames R_2 and R_1 for identical systems, in a Euclidean space where space (x) and time (t) are orthogonal, as per Figure A-1.

If we define

$$t'_i \equiv C_i t_i \quad , \quad (\text{A.9.0})$$

then by Figure A.1, since $C_1 = C_2 = C_{INV}$ and hence $\Delta t_* = 0$ (because of Galilean Invariance of P_{INV} and C_{INV}),

$$\Delta L_1^2 + \Delta t_1'^2 = \Delta L_2^2 + \Delta t_2'^2 \quad ; \quad (\text{A.9.1})$$

$$C_1 = (C_s \pm v_1) \quad ; \quad (\text{A.9.2})$$

$$C_2 = (C_s \pm v_2) ; \quad (A.9.3)$$

$$\Delta L_1 = \Delta L_2 \equiv \Delta L ; \quad (A.9.4)$$

$$\Delta t_1 = \Delta t_2 \equiv \Delta t ; \quad (A.9.5)$$

resulting with

$$(C_s \pm v_1)^2 - (C_s \pm v_2)^2 = 0 . \quad (A.9.6)$$

The simplest case to consider is $v_1 = -v_2$ with $|v_1| = |v_2| \equiv v$. This results with

$$v_1 = v_2 \quad (A.9.7)$$

or

$$\sqrt{1} = \sqrt{-1} \quad (A.9.8)$$

or

$$\pm 1 = i . \quad (A.9.9)$$

The usual closure relationships for the "Basic Ill-Posed Model" viewed in the laboratory frame

$$\frac{\partial P}{\partial t} = C_1^2 \frac{\partial}{\partial t} q_1 = C_2^2 \frac{\partial}{\partial t} q_2 \quad (A.10.0)$$

and

$$\frac{\partial P}{\partial x} = C_1^2 \frac{\partial}{\partial x} q_1 = C_2^2 \frac{\partial}{\partial x} q_2 , \quad (A.10.1)$$

inadvertently connect C_1 and C_2 , and hence the orthogonality of space and time.

Figure A-1. Galilean Invariance of P&C (pressure and sound speed), concept of time (lab frame).

APPENDIX B

We have studied a number of formulations of the Six-Equation, One-Pressure Model^(1,2) with the objective of obtaining a Well-Posed equation system.⁽⁶⁾ Although several approaches are known of rendering the “Basic Ill-Posed Model,”⁽²⁾ Well-Posed,^(2,14,15,16,17) such as adding surface tension,⁽¹⁴⁾ viscosity,^(16,17) or a second pressure,⁽¹⁵⁾ we question the necessity of these additions, and investigate the Ill-Posedness problem from a different perspective.

In this Appendix we show that the “Basic Ill-Posed,” Six-Equation, One-Pressure Model becomes Well-Posed when additional “Scalar Field” conservation equations are added to the existing equation set.

Let us consider cases B1 and C1 of the main text. Both result in an Ill-Posed problem with roots given by Equation (11) (of the main text). If we argue that

$$\frac{\partial}{\partial x} (P.) = C_v^2 \frac{\partial}{\partial x} (q_v) = C_t^2 \frac{\partial}{\partial x} (q_t) \quad (\text{B.1.0})$$

but

$$\frac{\partial}{\partial t} (P.) \neq C_v^2 \frac{\partial}{\partial t} (q_v) \neq C_t^2 \frac{\partial}{\partial t} (q_t) \quad (\text{B.2.0})$$

Closure requirements demand an additional equation.

We considered the additional Scalar Field conservation equations, Equations (B.3.0) and (B.4.0),

$$\frac{\partial}{\partial t} (\alpha) + \frac{\partial}{\partial x} (\alpha U_v) = 0 \quad (\text{B.3.0})$$

$$\frac{\partial}{\partial t} (q_v) + \frac{\partial}{\partial x} (q_v U_v) = 0 \quad (\text{B.4.0})$$

Analysis of Case C1, consisting of Equation (9) of the main text and Equation (B.3.0) with

$$\zeta = \begin{bmatrix} \alpha \\ \varrho_v \\ \varrho_l \\ U_v \\ U_l \end{bmatrix},$$

results in a Well-Posed system with the characteristic roots $\tau = 0$, $\tau = U_l$ and $\tau = U_v$. Analysis of Case B1, also consisting of Equations (9) and (B.3.0) with

$$\zeta = \begin{bmatrix} \alpha \\ \varrho_v \\ \varrho_l \\ U_v \\ U_l \end{bmatrix},$$

results in a Well-Posed system with the same characteristic roots as Case C1.

Additionally, if equality is assumed for Equation (B.2.0), and Equation (B.3.0) or (B.4.0) is substituted into Equation (9a) or (9b) of the main text, as a constraint, Well-Posedness results.

APPENDIX C

In this Appendix we show that, for the linearized case, mass, energy, and momentum are not conserved for an Ill-Posed system such as the Six-Equation, One-Pressure Model.

Let us begin with a Partial Differential Equation system of the form

$$\underline{A} \zeta_t + \underline{B} \zeta_x + \underline{C} = \pm \underline{D} \zeta_{xx} \pm \underline{E} \zeta_{tt} , \quad (\text{C.1.0})$$

where $\underline{A}$, $\underline{B}$, $\underline{C}$, $\underline{D}$, and $\underline{E}$ are $(NM) \times (NM)$ matrices and ζ is a column vector of (NM) elements. Subscripts in ζ indicate partial differentiation with respect to the subscript variable. For example,

$$\zeta_{tt} \equiv \frac{\partial^2 \zeta}{\partial t^2} \text{ and } \zeta_x \equiv \frac{\partial \zeta}{\partial x} .$$

"N" indicates the number of equation sets and "M" the number of equations in the set.

We are interested in the equation set that corresponds to the two-compressible-fluid, one-pressure-"averaged"^(1,2) model which is "consistent" with Equation (C.1.0) when $\underline{D} = \underline{E} = 0$, $M = 3$, and $N = 2$. This equation set requires "Equations of State" which we will assume are of the form

$$(P_i)_x = (e_i)_x c_i^2 , \quad (\text{C.2.0})$$

where

$$c_i^2 = (P_i)_{e_i} , \quad (\text{C.3.0})$$

and constant entropy is assumed for fluid "i". In this system we set⁽²⁾

$$P_1 = P_2 \equiv P . \quad (\text{C.4.0})$$

For our purposes we will use the equation set corresponding to Wendroff's⁽²⁾ "Basic Ill-Posed Model." From this point onward we will call this the "averaged" model.

Let's return to Equation (C.1.0) with $\underline{D} = \underline{E} = 0$. This corresponds to the form consistent with our "averaged model." The notion that this system is Ill-Posed, when the characteristics^(6,13) of this system are complex, is discussed in the literature.^(2, 5,12) Since it is known that under certain conditions the "averaged model" is Ill-Posed,⁽²⁾ we would like to understand what this means in terms of physical principles. Several excellent discussions on the relationship between "stability" and "Ill-Posedness" can be found in the literature.^(13,14,15) Let us, for simplicity, utilize a system where $N = 1$, $M = 2$, for the purposes of our discussion. This system is also discussed in Reference 6, and we will utilize the sign convention adapted there.

$$\rho_t + \rho u_x + u \rho_x = 0 \quad (C.5.1)$$

$$\rho u_t + \rho u u_x + c^2 \rho_x = 0 \quad (C.5.2)$$

The characteristic equation of this system is given by

$$\begin{vmatrix} 1 & (u - \tau) \\ (u - \tau) & c^2 \end{vmatrix} = 0 \quad (C.6.0)$$

where a linearized analysis is utilized with

$$\zeta = \zeta_0 \exp[-ik(x - \tau)] \quad (C.7.0)$$

This is equivalent to a local linear stability analysis⁽¹⁸⁾ for the case that $k \rightarrow \infty$ and $\omega \rightarrow \infty$ ($\lambda \rightarrow 0$), where

$$\zeta = \zeta_0 + \zeta' \exp(ikx - i\omega t) \quad (C.8.0)$$

and

$$\begin{vmatrix} 1 & \left(u - \frac{\omega}{k}\right) \\ \left(u - \frac{\omega}{k}\right) & c^2 \end{vmatrix} = 0 \quad (C.9.0)$$

is the dispersion equation under the assumption that (ω/k) has a finite limit. The roots of Equation (C.6.0) are

$$\tau_1 = (u + c) \quad (C.9.1)$$

$$\tau_2 = (u - c) \quad (C.9.2)$$

and the roots of (C.9.0) are (assuming ω -complex, k -real)

$$\omega_1 = (u + c) k \quad (C.10.0)$$

$$\omega_2 = (u - c) k \quad (C.10.1)$$

or (assuming k -complex, ω -real)

$$k_1 = \omega/(u + c) \quad (C.10.2)$$

$$k_2 = \omega/(u - c) \quad (C.10.3)$$

The condition for hyperbolicity for this system is that⁽⁶⁾

$$c^2 \geq 0. \quad (C.11.0)$$

If we assume c^2 is less than zero, we have an "Ill-Posed" problem. Let us examine what this means. Let's assume that

$$c = \beta_0 + i\beta_1 \quad (C.12.0)$$

and that $\beta_0 = 0$. If we study Equation (C.5) for adiabatic conditions, with constant initial energy, momentum, and mass, then Equation (C.12.0) implies lack of conservation of mass, energy, and momentum, for the situation of no sources or sinks, when a small perturbation occurs to the system Equation (C.5.0). This occurs even though the resulting equations with $c^2 = (i\beta_1)^2$ are "conservative" in a given unit cell volume because $\omega \equiv 2\pi c/\lambda$ which implies unbounded growth in time, that is,

for all

$$\exp \left(+ \frac{2\pi c t}{\lambda} \right) \text{ wavelengths.}$$

The reader should note that we have utilized a local linear stability analysis⁽¹⁸⁾ for a set of nonlinear equations. Nonlinear terms may saturate this unbounded growth. Hence, we have only shown that a serious problem may exist (from a theoretical point of view) for the “averaged model” and that an in-depth analysis of the nonlinear terms is required for all wavelengths (refer to Reference 18 for details). It should be kept in mind that while linear stability analysis can predict the onset of instabilities in nonlinear systems, the linearized equations are only valid for short times after the instability begins and are incapable of predicting system behavior once the oscillations have grown to a point for the nonlinear terms to be significant. However, Lax⁽¹²⁾ states that the nonlinear “averaged model” equations are not bounded and that total energy conservation arguments do not provide such bounds.

REFERENCES

1. M. Ishii, *Thermo-Fluid Dynamic Theory of Two-Phase Flow*, Direction des Etudes et Recherches d'Electricite de France, Eyrolles, 1975.
2. B. Wendroff, *Two-Fluid Models: A Critical Survey*, LA-UR-79-291, 1979.
3. L. D. Landau and E. M. Lifshits, *Fluid Mechanics, Chapter 16*, Pergamon Press, 1959.
4. P. Halliday and R. Resnick, *Physics*, John Wiley and Sons, Inc., 1966.
5. H. B. Stewart, *J. Comp. Phys.*, 33, 259, 1979.
6. R. Courant and D. Hilbert, *Methods of Mathematical Physics*, Interscience Publishers, 1962.
7. A. Einstein ("The Special Theory of Relativity"), "Zur Elektrodynamik bewegter Körper (On The Electrodynamics of Moving Bodies)," *Annalen der Physik*, 17, 1905, p. 891.
8. H. Minkowski, *A Translation of an Address at the 80th Assembly of German Natural Scientists and Physicians at Cologne, September 21, 1908, In the Principle of Relativity*, Dover Pubs. Inc., 1952, pp. 75–91.
9. C. Møller, *The Theory of Relativity*, Oxford: Clarendon Press, 1952.
10. H. Makowitz, *Transactions of the American Nuclear Society*, 64, 1991, pp. 316–318.
11. H. Makowitz, Internal INEL correspondence (unpublished), December 1990.
12. P. D. Lax, *Communications on Pure and Applied Mathematics*, 33, 3, 1980, pp. 395–397.
13. R. D. Richtmyer and K. W. Morton, *Difference Methods for Initial-Value Problems*, 2nd Edition, Interscience Publishers, 1967.

14. J. D. Ramshaw and J. A. Trapp, *Nuclear Science and Engineering*, 66, 1978, pp. 93–102.
15. V. H. Ransom and D. C. Hicks, *Journal of Computational Physics*, 53, 1984, pp. 124–151.
16. M. Arai, *Nuclear Science and Engineering*, 74, pp. 77–83, 1980.
17. F. H. Harlow and A. A. Amsden, *J. Comp. Phys.*, 17, pp. 19–52, 1975.
18. N. A. Krall and A. W. Trivelpiece, *Principles of Plasma Physics*, McGraw-Hill, Inc., 1973.

END

DATE
FILMED

2 / 22 / 93