Illinois State Water Survey

ATMOSPHERIC SCIENCES DIVISION

SWS Contract Report 497

STUDY OF ATMOSPHERIC POLLUTION SCAVENGING

Twenty-Fourth Progress Report
Grant Number DE - FG02 - 88ER60635

August 1990

Author:

Allen L. Williams
Office of Precipitation Quality

Sponsored by:
United States Department of Energy
Office of Health and Environmental Research
Washington, DC

Richard G. Semonin
Principal Investigator

ENR

Illinois Department of Energy and Natural Resources

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

SWS Contract Report 497

STUDY OF ATMOSPHERIC POLLUTION SCAVENGING

Twenty-Fourth Progress Report
Grant Number DE - FG02 - 88ER60635

August 1990

Author:

Allen L. Williams

**Office of Precipitation Quality
Illinois State Water Survey
2204 Griffith Drive
Champaign, Illinois 61820-7495**

Sponsored by:

**United States Department of Energy
Office of Health and Environmental Research
Washington, DC**

**Richard G. Semonin
Principal Investigator**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Contents

Apportionment of Multiple Aerosol Size Distributions Modes
Using Factor Analysis Techniques.....1

Implications of the Formation of Cloud Condensation Nuclei
from Gaseous Precursors.....19

Study of Atmospheric Pollution Scavenging
Ten Year Review (1980-1990) of DOE Research.....27

*2 conf papers
cycled separately
ch*

**Study of Atmospheric Pollution Scavenging
Ten Year Review (1980-1990) of DOE Research**

**by Allen Williams
Illinois State Water Survey
Champaign, IL**

**Research supported by the Office of Health and Environmental
Research of the United States Department of Energy**

I. Introduction

Atmospheric scavenging research conducted by the Illinois State Water Survey under contract with the Department of Energy has been a significant factor in the historical development of the field of precipitation scavenging. Emphasis of the work during the 1980's became focused on the problem of acid rain problem with the Survey being chosen as the Central Analytical Laboratory for sample analysis of the National Atmospheric Deposition Program National Trends Network (NADP/NTN). The DOE research was responsible for laying the groundwork from the standpoint of sampling and chemical analysis that has now become routine features of NADP/NTN. A significant aspect of the research has been the participation by the Water Survey in the MAP3S precipitation sampling network which is totally supported by DOE, is the longest continuous precipitation sampling network in existence, and maintains an event sampling protocol.

The following review consists of a short description of each of the papers appearing in the Study of Atmospheric Scavenging progress reports starting with the Eighteenth Progress Report in 1980 to the Twenty-Third Progress Report in 1989. In addition a listing of the significant publications and interviews associated with the program are given in the bibliography.

II. Review of Eighteenth Progress Report, (Contract Number EY-76-S-02-1199) Report Number COO-1199-60, SWS Contract Report 235, July 1980

A. Ambient Aerosol Measurements at Champaign, IL, G. J. Stensland and J. D. Bartlett

The sampling and extraction procedures for all the aerosol filter samples is described and three types of data are presented. First, a comparison of Whatman versus Nuclepore filters for about 70 samples is presented. Second, a comparison of Nuclepore versus Fluoropore filters for six samples is presented. The final section presents the results for one year of daily aerosol sampling at the Willard Airport site.

B. Nitrate Measurements in Precipitation for the Study of Trends and Patterns, G. J. Stensland

Spatial patterns and the time trends of nitrate in precipitation are examined. The data are from studies carried out in the United States in the mid 1950's and the late 1970's. For the temporal trends, data from the northeastern United States are emphasized. An annual

cycle of the nitrate concentrations in precipitation for Champaign, Illinois is compared to ambient aerosol data for the same location.

C. SCORE-78 Case Study for July 2, 1978, G. J. Stensland

The concentrations of the various ion concentrations in a single precipitation event over an approximately 50 square mile area is examined. Contours of the ammonium, sulfate, nitrate, calcium, magnesium, sodium, potassium, chloride, pH, and conductivity are presented. Sequential data at one of the sites is also discussed.

D. Multielement Analyses of Selected Samples from SCORE-1979, D. F. Gatz and G. J. Stensland

This preliminary report describes a case study to determine if the SCORE-1979 rain chemistry data falls into the same factor analysis groupings of soluble and insoluble crustal elements, soluble metals and sulfate, and insoluble metals as was found in a METROMEX analysis. The rain sampling network and sample analysis procedures are describe.

E. An Urban Influence on Deposition of Sulfate and Soluble Metals in Summer Rains, D. F. Gatz

Observations suggest that rainfall deposits at least some locally-emitted sulfur at short distances downwind of the sources which causes enhanced deposition and variability in the affected areas. The variation in airborne sulfate concentrations is roughly the same as that observed in either concentration or deposition of sulfate in rain. Thus, there is no need to invoke in-cloud scavenging of sulfur dioxide to explain enhanced sulfur in rain near the city, since there is ample aerosol sulfate and an available scavenging mechanism (nucleation) to account for it. The observation of consistently similar deposition patterns for sulfate and soluble pollutant metals places some interesting constraints on possible scavenging mechanisms and suggests that more detailed study of this group of materials would be fruitful for understanding the processes involved.

F. Associations and Mesoscale Spatial Relationships Among Rainwater Constituents, D. F. Gatz

Factor analysis of multielement deposition of both soluble and insoluble materials in single storms over networks of approximately 2000 km² near St. Louis have revealed that four main groupings of deposition patterns occur.

- 1) Soluble soil-derived elements,
- 2) Insoluble soil-derived elements,
- 3) Soluble pollutants, and
- 4) Insoluble pollutants.

G. Spatial Variability of Rain Water Impurities in Mesoscale Events, D. Q. Naiman and D. F. Gatz

A summary of an investigation of the spatial variability of precipitation impurities in the METROMEX network of precipitation samplers is presented. Reasons why precipitation impurity variations are important are discussed and on some typical results presented. It is shown that taking account of spatial correlation improves the accuracy of estimating summer rainfall and deposition from single events using open collectors (corrected for dry deposition contributions) over 2000 km² in the St. Louis area.

H. A Precipitation Chemistry Data Management System - Summary, R. K. Stahlhut and V. C. Bowersox

An automated data management system for precipitation chemistry is described. The system has two principle features: (1) a computerized data base to store the information in an easily and selectively accessible form, and (2) a library of computer programs to process and check the data for consistency and errors and to organize and report the data in formats amenable to the data user and to the data analyst. Together these features combine to provide a simple and efficient means of managing large precipitation chemistry data sets.

I. Analytical Chemistry Laboratory, M. E. Peden

The expansion of the Analytical Chemistry Laboratory with the installation of an automated ion chromatograph and associated equipment is described. In addition, the Analytical Chemistry participation in the Fourth-Environmental Protection Agency Precipitation Reference Sample Interlaboratory Comparison is discussed.

J. Ambient Air Filter Extraction Procedures, J. Bartlett and G. J. Stensland

The development of an extraction technique for the analysis of aerosol filter samples is described. The

cation exchange capacity of the soil particles can influence the ionic concentrations of Ca, Mg, Na, and K in the extraction medium and complicate the determination of the amount of soluble ionic material. The extraction method developed accounts for the exchangeable ions to better simulate the soluble ions scavenged by precipitation.

III. **Review of Nineteenth Progress Report, (Contract Number DE-AC02-76EV01199) Report Number C00-1199-60, SWS Contract Report 252, May 19**

A. **Summary of Sulfate and pH for SCORE-79, G. J. Stensland**

The Summer Chemistry Of Rain Experiment for 1979 (SCORE-79) had the objectives of assessing the surface rainfall chemistry variability. All available data from each site for the 1979 field experiment were included in the analysis. The sulfate values varied between 0.5 and 15 mg/L or by more than a factor of 30. The range of median values for the data set was 1.7 to 4.7 mg/L or a factor of 2.8. The range of pH values was 3.3 to 7.4 for all data, but for heavier rainfalls the range was 3.4 to 7.1. The median pH for all data was about 4.3.

B. **Reproducibility of Aerosol Samples Collected on Nuclepore Filters Under Inverted Funnel Rain Shields, D. F. Gatz**

An experiment is reported in which samples were collected simultaneously on identically exposed filters using inverted funnel rain shields. Good agreement was found for Si, S, K, Ca, Fe, and TSP and poorer agreement for Ti, Mn, Zn, Br, and Pb. However, statistical analyses reveals that the agreement is clearly less than desired. Some possible explanations for the discrepancies are offered, but additional experiments are needed to clarify the problem for addressing spatial variability problems.

C. **Comparison of Two Sampling Methods for Medium-Volume Aerosol Sampling, D. F. Gatz**

A comparison was carried out between an inverted funnel aerosol collection method and an isokinetic sampler free to move with the wind. The data shows that the isokinetic sampler collects more aerosol for the 11 elements analyzed and TSP than the inverted funnel. The isokinetic/funnel ratio ranged from 1.05 for S to 1.32 for Si and Ti. This finding must now be placed in the context of its impact on scavenging ratio calculations.

- D. Comparison of Total Versus Soluble Data for Ambient Aerosol Filter Samples, G. J. Stensland and D. F. Gatz

An attempt was made to determine the soluble/insoluble ratio for Ca, K, and S of daily aerosol samples. The total concentration was found by ion-excited X-ray fluorescence while the soluble contribution was determined by flame atomic absorption and automated colorimetric methods. The results are inconclusive and a number of possible problems need to be explored before quantitative values will be available.

- E. Investigation of Multivariate Statistical Methods for Estimating Background Lithium in Precipitation Tracer Experiments, D. F. Gatz

Statistical methods to estimate background lithium in tracer experiments were used to help predict the deposition patterns. The principle components regression did not yield a result useful to estimate scavenging efficiency. The primary useful result is that the background concentration of any tracer must be carefully assessed in designing future experiments, and the methods development points the direction that such assessments must take.

- F. Design of a Feasibility Study to Assess the Impact of Precipitation Quality on Soil Water Quality, V. C. Bowersox

To study the impact of precipitation soil water quality, the chemical composition of the rain would be compared to the ground water chemical composition. A field lysimeter device is discussed by which the soil water can be collected at different depths. The results would depend on the soil chemistry eg soil exchange capacity and pH, which would also be assayed at the sampling site.

- G. A New Search for Old Evidence to Identify Trends in the Chemistry of Precipitation in the Continental United States, V. C. Bowersox

In order to more clearly identify the trends in precipitation chemistry it is desirable to have a records that pre-date the current high levels of SO_x and NO_x . Data is being collected for the early decades of this century and evaluated according to sample location, sampling protocol, laboratory and analytical techniques, and contamination sources.

- H. Seasonal Precipitation concentrations and Depositions for North American from the CANSAP/NADP Networks, R. G. Semonin

Using combined data sets from the NADP and CANSAP networks, descriptive maps of calcium, ammonium, nitrate, sulfate and pH were prepared. Seasonal and annual distributions are shown for the data period July 1978 through September 1980.

- I. A New Sequential Rainwater Sampler, G. J. Stensland

The evaluation of a newly designed sequential sampler for precipitation chemistry was carried out in a field experiment in 1980. The results are quite promising although some minor changes will be made as a result of the evaluation prior to construction of additional units.

IV. Review of Twentieth Progress Report, (Contract Number DE-AC02-76EV01199) Report Number C00-1199-64, SWS Contract Report 347, September 1984

- A. Acid Rain Data Collection, Handling, Analysis and Interpretation, R. G. Semonin

Some recent data are shown for pH to illustrate its great spatial variability in the United States. These data show that the lowest values exist in the Northeast while the Great Plains show the highest pH values. The ratios of hydrogen, ammonium, and calcium ions to the sum of sulfate and nitrate reveal that maximum values occur in the Northeast, central Great Plains, and the intermountain basin, respectively. Finally, a brief account of reinterpreting the older data show that the previously perceived rapidly worsening precipitation acidity problem is not true. When account is taken of different sampling procedures, analytical methods, and, most importantly, meteorological sampling conditions, the early chemistry was proven to be inadequate as an initiating point to determine a trend in the United States.

- B. A Comparison of Four Methods of Computing Precipitation pH Averages, G. J. Stensland and V. C. Bowersox

The question of the calculation and presentation of pH data is one of importance to the determination of acidic precipitation trends owing to the fact that different numbers can be obtained even from the same data set. Examples are shown for the calculation of various climatological values of pH. These range from the

precipitation-weighted average pH calculated from the weighted average conservative ion concentrations. Since all of these methods have appeared in one or more publications dealing with the spatial distribution and the trend of precipitation acidity across the U.S., it is important to consider which of these methods is most appropriate for an assessment of the acid rain issue. In summary, this study shows there is no "correct" answer to provide the best value of pH. However, a clearly stated purpose for which the average or central tendency value is needed will help to determine which method will give the most representative value.

C. Regional Characterization of Rain Acidity Utilizing Gran's Plot Titrations, M. E. Peden and J. M. Lockard

In addition to the major anions (sulfate and nitrate) contributing to the acidity of precipitation, it is now recognized that organic acids also contribute to the total acidity. Some preliminary work is reported on the regional characterization of rain acidity through the use of Grans plot titrations. It was found that the ratio of strong to weak acids decreases from east to west as a result of both decreased strong acidities and increased ammonium concentrations. Due to sampling problems, no evidence of a significant contribution of organic acids was found in samples from any of the sites used in the study. Considerable additional work is needed in this area.

D. The Chemistry and Meteorology of Extreme pH Departures in Precipitation on the Regional Scale, K. G. Doty and R. G. Semonin

To explore this possibility that airborne surficial material may have biased the precipitation chemistry of the mid-1950's statistical methods were employed on NADP network data sets selected from recent years. A case was selected from 1981 with relatively high pH values over the Northeast, and an additional case was chosen with relatively low pH values in the same area. Using these data, a meteorological explanation was then sought related to suspended airborne material. Trajectory analyses in one case in the spring of 1981 provided evidence for the potential transport of crustal dust aerosol from the southern Great Plains and incorporated in precipitation at sites in the Midwest. The more acid case of July 1981 revealed high concentrations of most ions in comparison with adjacent weeks. These high concentrations appeared to be related to the smaller average sample volume across the Midwest and Northeast and the relatively long dry period that was observed

prior to the main rainfall event. Additional case studies will be pursued to assist in understanding the natural variability of precipitation acidity.

E. Dry Bucket Analysis of the NADP Network, R. G. Semonin

The MAP3S and NADP networks both use a sampler with a collector for wet-only samples and a collector exposed to dry deposition during non-precipitating periods. The analysis of the dry bucket samples collected from the NADP are presented. Lack of agreement over a dry deposition measurement method has resulted in a lack of data for a full assessment of the impact of acidic deposition. The dry bucket data may be questioned by the "purist." This is a first attempt to describe the regional dry deposition using readily available data. While more work is needed, these results show between station consistency within large areas of the eastern half of the United States. The absolute magnitude of the dry deposition values may be questioned, but the data do give a first glimpse of the areal distribution that is expected with implementation of more sophisticated equipment in the future.

F. Metal Solubility in Atmospheric Deposition, D. F. Gatz and L. C. Chu

The assessment of the effects of acidic deposition can not be independently achieved without due consideration given to other coincident and potentially important precipitation chemistry. A set of atmospheric deposition samples in wet, dry, and bulk precipitation were analyzed for metal solubility. The samples were analyzed for zinc, aluminum, copper, iron, cadmium, and lead. The percent soluble metals show zinc is about equal to cadmium and both are greater than copper, which is greater than lead. In general, the wet-only samples showed more solubility than bulk and dry which were nearly equal. These results have important implications for precipitation sampling for metal analysis as well as for the understanding metal chemistry in precipitation, airborne metal scavenging processes, metals as tracers for sources of acidity, and the effects of acidic precipitation on ecosystems.

G. Source Apportionment of Rain Water Impurities in Central Illinois, D. F. Gatz

The identity and relative contributions of various sources of impurities in precipitation are needed to understand the acidic precipitation phenomenon. Factor analysis was used to identify, and chemical element balance calculations were performed to apportion,

possible sources of impurities in event precipitation samples from central Illinois collected over a two year period. Although factor analyses could group elements or ions that occurred together for chemical, meteorological, or microphysical reasons, as well as that of having a common source, the following likely sources were identified: crustal dust, sea and/or road salt, and possibly strong acids. In addition, a gaseous precursor factor was identified. The smallest contribution appears to be sea and/or road salt with the greatest contribution in the acids and the gaseous precursors. The contribution of crustal dust is sizable and certainly warrants further investigation. Extensions of this work should include plotting the factor scores against the date of the sample collection as a means to relate the source term to meteorological conditions.

H. A Few Preliminary Precipitation Chemistry Statistics, R. G. Semonin

The NADP network data were used to examine the correlation coefficients between the various ions that are commonly analyzed. The relationships between these various ions are described and maps are presented indicating areas of relatively high correlations between specific ions. These maps indicate that the Northeast United States experiences very high correlations between the acid-forming ions, and high correlations of acid neutralizing materials over the interior continent. The coastal regions show correlations which are equal or greater than either the acid or base ions between sea salt components.

I. Results of the World Meteorological Organization (WMO) Interlaboratory Comparison of Reference Precipitation Samples, M. E. Peden

The research results presented in this report are given with full confidence in the quality of the analytical values that form the basis for interpretation and subsequent conclusions concerning acidic deposition. To sustain high quality analytical methodology, the laboratory continues to participate in various interlaboratory comparisons for low concentration water samples. The most recent World Meteorological Organization comparison results are presented in chapter 10. Throughout the past four years, the laboratory results have consistently yielded mean percent deviations of 3 to 5% from the test sample input. These activities will be continued to insure the highest level of accuracy for reported numbers from precipitation samples to insure the user that the data are meaningful and can be

interpreted and reported with confidence.

- J. Microcomputer Based Data Acquisition and Reduction System for Ion Chromatography, S. R. Bachman and R. K. Stahlhut

The dependability of analyses is only one essential part of the laboratory performance. The other part is the rapidity with which samples can be analyzed and made available to the research community. A microcomputer based data acquisition and reduction system for ion chromatography is described. A microcomputer is connected on-line to an ion chromatograph forming an analytical system of proven accuracy and efficiency for data acquisition and reduction of precipitation sample ion determinations. The most important aspect of this automation is that the analyst time devoted to data reduction tasks has been reduced by an estimated 25%. Another way of viewing this automation is that the through-put in the laboratory has been increased by a comparable percent.

- K. Determination of Organic Acids in Bondville, Illinois Precipitation Samples by Ion chromatography Exclusion (Ice), S. R. Bachman, J. M. Lockard and M. E. Peden

Acidic precipitation samples are generally thought to be composed of a mixture of the strong acids sulfuric and nitric. The potential contribution of other components, such as weak acids that are only partially disassociated, can not be overlooked. Some preliminary work on the determination of organic acids to the chemistry of rainfall can be quite varied. The weak to strong acid ratio for Illinois precipitation is similar to that of sites in the northeastern United States. Based on the data accumulated thus far for central Illinois, the findings are in agreement with those reported for the Northeast U.S. that strong acids control the pH of rainfall, but weak acids are important to the total chemistry of precipitation. Since this preliminary data base is limited to samples collected during the spring months, however, additional work needs to be carried out to examine seasonal differences in weak acid concentrations, and weak to strong acid relationships.

- L. Feasibility Study to Assess the Impact of Precipitation Quality on Soil Water Quality, V. C. Bowersox

The acidity of precipitation over the eastern half of the cash grain belt across Iowa, Illinois, Indiana, Ohio, and north and south of those states is of the same value as the much publicized Adirondack lake area of New York state. One of the early concerns in assessment of the

acidic deposition problem was the short or long-term damage to soil productivity. Because of this concern and the fact that Illinois represents much of the corn belt (from a climatological point of view), a small effort was devoted to estimating the effects of precipitation quality on soil water quality. The results from this preliminary and limited effort are discussed. With the data available using lysimeters to sample soil water, a few results indicate that the precipitation pH is clearly less than the soil water pH for the soils of central Illinois and that neutralization of the precipitation occurs in the top most layer of the soil for even the most acidic rainfalls observed (pH of 3.82). These data also indicate that a weak relationship exists between soil water pH and free hydrogen ion deposition with the soil water pH slightly depressed with increased hydrogen ion input. However, much additional research effort is needed to clarify this relationship and to further estimate the impact of acidic deposition on the soils of the Midwest.

- M. Precipitation, Chemical Depositions and Water Resource Issues, R. G. Semonin

Drawing upon the extensive literature on the various presumed effects of acidic deposition on water resources, some of the knowns and unknowns concerning this aspect of a total assessment are described. The material presented was collected together for presentation before an international conference, but the material bears repeating in this Progress Report as it addresses a topic of some national interest. The final conclusion of this is that little is known concerning the effects of acidic deposition on the chemical quality of surface and ground water supplies.

- N. A Review of Particle Tracers of Atmospheric Process, D. F. Gatz

A review of particle tracers of atmospheric processes is presented in Chapter 15. This paper was presented at the Department of Energy Workshop on Atmospheric Tracers held in the spring of 1984, but is relevant to the entire subject of acidic deposition assessment since it reviews one of the possible research techniques available to investigators.

- O. Tracer Applications for the Study of Precipitation Processes, R. G. Semonin

Application of tracers to the study of precipitation processes with specific application to the acidic

deposition question is discussed. The precipitation process is described as beginning with the availability of water vapor and condensation nuclei, and then moves through the requirement for motions in the atmosphere to produce clouds and precipitation. There is a clear need for tracers in the study of all of these atmospheric attributes leading to precipitation production. However, the presently available suite of gaseous and particle tracers each have inadequacies which inhibit their direct application to resolving acidic precipitation research problems. There is a need for the development of a new tracer which can be freely used in the atmosphere and will have a direct bearing on the conversion and scavenging of acidic deposition precursors.

V. **Review of Twenty-First Progress Report, (Contract Number DE-AC02-76EV01199) Report Number C00-1199-65, SWS Contract Report 426, July 1987**

A. **Wet Deposition Chemistry, R. G. Semonin**

Interest in precipitation chemistry dates back to the 18th century, but only since 1977 has the U.S. established a national network for monitoring at a level sufficient to describe the chemical climate. The siting criteria, quality assurance, and standardized equipment and analytical methodology are addressed to insure the highest quality data for interpretative research. The national distribution of concentration and deposition for calcium, ammonium, chloride, nitrate, sulfate, and pH are shown in a series of maps. The maps show high concentrations of calcium and ammonium in the Great Plains, high concentrations of nitrate, sulfate, and hydrogen ion in the eastern states, and high concentrations of chloride along the coastal states. The primary conclusions are 1) the greatest deposition is directly associated with the largest emission regions, and 2) dry deposits are largely unknown but clearly must be measured before the biochemical and geochemical cycles will be fully understood.

B. **Intra-Annual Variations in the Concentrations of Major Inorganic Ions in the Precipitation of the United States, V. C. Bowersox and G. J. Stensland**

This study examines the intra-annual variations of inorganic ion concentrations in precipitation across the U.S. These variations were summarized on contour maps showing the spatial distribution of ratios of warm period to cold period weighted average and median ion concentrations. The data suggests that precipitation

concentrations followed a rather well defined pattern of intra-annual variations with summer and winter extremes. For sodium and chloride, the Northern Plains and nearly all of the East had warm/cold period ratios <1.0 . Some east coast sites even had ratios <0.5 , probably owing to high wintertime airborne sea-salt concentrations. For ions other than sodium and chloride there were generally higher average warm (growing) season than cold (dormant) season concentrations over much of the country.

C. Estimation of Bias in the Average Wet Deposition of Calcium, Sulfate, and Sodium from National Atmospheric Deposition Program/National Trends Network (NADP/NTN) Data, G. J. Stensland

Estimates were made of the percent bias in the wet deposition loading of Ca, SO_4 , and Na from the NADP/NTN data set. These estimates were based on the data of record from mid-1978 through 1986 and are thus applicable to long-term averages. The calculations of an overall bias were based on data for the contamination from field handling of dry buckets, laboratory handling and analysis, and the buckets themselves. Sources of possible contamination that were not addressed included (a) field handling to remove a liquid aliquot of sample for field chemistry measurements and (b) chemical degradation of the liquid sample from time of collection to time of analysis. No attempt was made to address the bias due to samples undercatch or overcatch of precipitation. Infiltration of the lid to bucket seal by aerosol Ca, SO_4 , and Na occurred over large spatially contiguous areas, and contamination from the lid (gasket) was typically the largest single source of bias.

D. Spatial Relationships Between Acid Rain, Air Quality, and Visibility Data, G. J. Stensland

Visibility and acid rain are two different phenomena which both have direct relationships to air quality. The objective of this paper is to explore the relationships between all three data sets, with emphasis on the situations for the eastern U.S. This paper presents interpretation of the most current precipitation chemistry data from the National Atmospheric Deposition Program/National Trends Network (NADP/NTN), which started in the eastern U.S. in 1978 with less than 20 sites and continues today with over 200 sites spread across the entire U.S. The high site density and relatively long data record make the precipitation sulfate data particularly important. Correspondence is found between precipitation sulfate and air quality sulfate for areas

to the east. In coastal areas the spatial pattern for visibility is not consistent with the precipitation sulfate pattern, presumably due to the effect of relative humidity.

E. Variations in Precipitation Composition and Deposition with Synoptic Weather Type, K. G. Doty and D. F. Gatz

As one portion of a broader analysis of five years of event precipitation chemistry data for central Illinois, we examined variations in ion composition and deposition with synoptic weather type. Variations of H, NO₃, and SO₄ deposition and concentration with synoptic weather type were very consistent in all four seasons. Deposition frequencies were strongly a function of the sample volume distributions, which was in turn largely determined by the frequency of the various synoptic types, while concentrations were relatively constant over all synoptic types. The highest concentrations occurred in spring and summer, the lowest in fall and winter. The results suggest that historical depositions of certain ions in precipitation might be estimated from precipitation records, combined with appropriate annual mean concentrations of the major ions. The appropriate concentrations might be inferred from airborne concentration measurements, if available, or relevant emissions data.

F. Back Trajectory Climatologies for Sulfate and Nitrate Wet Deposition Fluxes at a Central Illinois Site, K. G. Doty and V. C. Bowersox

Precipitation samples have been routinely collected with automated wet-only collectors since September 1979 at a field site in rural east-central Illinois. Event record data were combined with the chemical concentration measurements of samples from the collectors to compute wet-only fluxes of sulfate and nitrate. Back trajectory calculations for these flux classes utilizing the ISWS trajectory model are presented. The analysis involves approximately 250 precipitation chemistry samples, and the calculation of about 1100 back trajectories each 72 hours in length.

G. Interlaboratory Comparisons of Reference Precipitation Samples, M. E. Peden and J. M. Lockard

The analytical chemistry laboratory has continued its participation in the interlaboratory comparisons of precipitation chemical analyses sponsored by the World Meteorological Organization (WMO). In the past three

years our involvement in international precipitation chemistry intercomparisons has expanded to include participation in studies conducted by the Canadian Long Range Transport of Atmospheric Pollutants (LRTAP) Program and the European Monitoring and Evaluation Program (EMEP) carried out by the Norwegian Institute for Air Research. These programs provide valuable data on the comparability of chemical determinations made at geographically disperse laboratories as well as documentation of the analytical procedures that are being utilized for precipitation chemistry measurements. This data base is an important source of information when comparing chemical results from different monitoring programs. Summaries of these programs and results from our laboratory are presented.

H. Quality Assurance of Precipitation Chemistry Analyses Using Ion Balance Calculations, J. M. Lockard

The National Atmospheric Deposition Program (NADP) was initiated in 1978 to assess spatial and temporal trends in precipitation chemistry within the United States. From the beginning of the program, the Illinois State Water Survey (ISWS) has been responsible for the chemical analyses of precipitation and dry deposition samples generated from the monitoring network. A rigorous internal quality assurance program has been developed to ensure that the laboratory measurements are of the highest quality. As a part of this program, each sample analysis is verified by the use of an ion balance calculation. The ion balance data have been used in part to develop objective criteria for the reanalysis of samples on a routine basis. A closer examination of these data has revealed patterns in the ion ratios that are a function of sample pH, site location, and collection season. These patterns are discussed in terms of how to most effectively utilize ion balance data to verify the accuracy of laboratory measurements.

I. A Comparison of Ion Chromatography and Automated Colorimetry for the Determination of Major Anions in Precipitation, S. R. Bachman

The two analytical methods of automated colorimetry and ion chromatography have been compared with respect to comparability of analytical results and physical features. Detection limits are somewhat improved with the ion chromatograph, although a bias for one method over the other was not detected with regard to sample concentrations. A wide range of analyte values was examined and correlations between the two methods are excellent.

- J. Long-Term Ionic Stability of Precipitation Samples, M. E. Peden and L. M. Skowron

The long-term stability of precipitation samples has been investigated by measurement of samples stored in a refrigerator for six years. Of the eight constituents measured, Ca, Mg, Cl, SO_4 , and NO_3 revealed reanalysis results within 10% of their original values. Evaporative losses during storage were not evident. Hydrogen ion concentrations increased by nearly 40%, on average, with a concomitant increase in solution conductivity. Ammonium results did not reveal a significant change in mean concentration, although the data were highly variable compared to the other analyte differences that were measured. No distinct patterns of long-term stability related to geographic regions or collection seasons were evident in the study.

- K. Synthetic Rainwater Reference Samples, L. M. Skowron and M. E. Peden

The objective of this study was to develop internally formulated reference materials that could be used in conjunction with NBS Standard Reference Materials, which cannot be used on a routine basis to monitor laboratory performance because of their high cost. Results presented indicate that a concentrated stock solution containing all of the major constituents in wet deposition can be easily prepared using commonly available salts and mineral acids. Dilutions of this single stock solution to the concentration levels characteristic of wet deposition have resulted in stable formulations that can be used for routine quality control checks, as well as for internal blind audit samples. Five months of data indicate that sample integrity is maintained without the addition of any chemical preservatives or refrigerated storage. Ongoing studies are focusing on the long term stability of the stock concentrate as well as the dilutions to develop statistical guidelines for using data obtained from these solution analyses to ensure traceability to the NBS Standard Reference Material.

- L. Preliminary Data for Daily Ambient Aerosol Measurements at the Bondville Research Site, G. J. Stensland

Results from the chemical analysis of daily aerosol filter samples are presented. It is found that:

- The teflon filters are significantly more efficient in collecting SO_4 and K particulate matter than are $0.8 \mu\text{m}$ polycarbonate filters.

- The Na blank levels are unacceptably large for the aerosol data.
- SO_4 , Ca, and Mg are maximum in summer and minimum in winter; K is maximum in fall and minimum in winter and spring; and Na is maximum in winter.

M. Ratios of Ion Concentrations in Precipitation to Aerosols at the Bondville Site, G. J. Stensland and V. C. Bowersox

Gases and aerosols are incorporated into precipitation by scavenging processes which occur within and below clouds. In this chapter we present aerosol and precipitation chemistry data for SO_4 , Ca, Mg, K, and Na. These data are summarized by season from measurements made at the Bondville Field Site, operated under this contract. Of the five species, only SO_4 has both particle and gaseous species to consider.

- From this preliminary report we conclude that scavenging ratios for Na and K may be affected by problems related to sampling, handling, and extraction methods, which make the values unrepresentative. Values cannot be directly compared to those for SO_4 , Ca, and Mg.
- Further sensitivity and statistical analyses are required to see if differences in the scavenging ratios for SO_4 and Ca or for SO_4 and Mg can be used to predict the gaseous versus particulate contributions to the total SO_4 in precipitation. The present results indicate that particulate removal is more important than gaseous removal for sulfur, especially in the summer.

N. A Computer Program for Calculating State Averages from Point Values, J. Su and G. J. Stensland

We have developed a computer program to calculate a single representative statistic for each state, given a variable that is measured at a limited number of sites across the United States. Our program is called AREA for ARea Estimating Algorithm. The AREA program is a useful program that will calculate area-weighted statistics on a state-by-state basis for variables that change with geographical locations. The current implementation of AREA can be easily altered to suite the user's needs, being, for example, general enough that with only a few minor changes it can be incorporated as a subroutine of another program.

- O. The Description and Comparison of a Prototype Trajectory Model with the ARL-ATAD Model, K. G. Doty

Concerns surrounding air and precipitation quality have led to frequent use of trajectory models in the past decade. An example of a relatively simple diagnostic model is the Air Resources Laboratories Atmospheric Transport and Dispersion Model (ATAD). The ATAD model was designed to calculate large numbers of boundary layer trajectories from several origins in an economical manner. The results of trajectory analyses for five years of event precipitation chemistry data for a site in Central Illinois are shown. The analyses led to the development of a new trajectory model (ISWS) which is described and compared to ATAD.

- P. Screening Criteria for NADP Dry Bucket Data, D. F. Gatz, V. C. Bowersox, and J. Su

It is quite clear from the results presented that "dry" fluxes of H ion are substantially decreased if samples containing water are eliminated from the data set, and some evidence that the same may be true for nitrate. Similarly, the dry fluxes of K, NH_4 , and PO_4 , while relatively insensitive to elimination of samples containing water, are much reduced when samples having one or more types of visual contamination are eliminated. The dry flux of Ca actually increased as the screening level increased. The dry fluxes of the remaining ions examined--Na, Mg, Cl, and SO_4 --showed no significant differences between screening levels. The present analyses suggest that additional future work on the current data set should include quantification of the individual effects of each potential difficulty (e.g., water in the sample, or visual contamination), and particularly, each different form of contamination. In addition, the samples might be grouped regionally (at least east vs west) to see whether there are regional differences in the way that potential contaminants affect the measurement of dry fluxes using buckets.

- Q. An Annotated Description of the Atmospheric Chemistry Sampling Station at Bondville, Illinois, S. R. Dossett

The maps, figures and text included describe the area surrounding the Bondville Field station, detail the history of the research activities, and briefly describe the research currently supported at the site. DOE contract funds make up a considerable portion of the

budget upon which the Bondville site depends. Data collected at the Bondville site offer a mix of precipitation and air chemistry information with meteorological measurements. As such, the site offers a good opportunity for the research community to study in detail the interrelated nature of the climatology, wet deposition, dry deposition, and aerosol components of the atmosphere.

VI. Review of Twenty-Second Progress Report, (Grant Number DE-FG02-88ER60635) Report Number C00-1199-66, SWS Contract Report 451, August 1988

A. A Description of the Sequential Chemistry of Rain Experiment-1988, A Component of 3CPO, V. C. Bowersox

SECORE-88, or Sequential Chemistry of Rain Experiment-1988, operated a smaller network of four wet-only sequential samplers and 13 raingage sites. Sites were separated by about 1.1 kilometers and rainfall sequences were collected every .05 cm. The common anions and cations were measured at three locations and at one of these locations, the Bondville site, samples from one sequential sampler were collected exclusively for aqueous H_2C analyses. SECORE-88 stations were nested within the larger PRECP-VI array and measurements were made on a higher time resolution than at PRECP-VI stations.

The objectives of SECORE-88 were:

- (1) to measure the wet fall chemistry of convective raincells and to quantitatively describe the variability of ion concentrations in time and space for convective cells;
- (2) to examine the variations in chemistry both along and lateral to the motion of convective raincells and to relate these variations to the dominant scavenging processes;
- (3) to look for differences in the wet fluxes of species with gaseous precursors (HNO_3), gaseous and small particle precursors (SO_2 + aerosol SO_4^{2-}), large particle precursors (Ca^{2+} from soils and road dust), and others;
- (4) to apply statistical techniques of source identification; and
- (5) to provide data to the 3CPO community for use in collaborative research with other investigators.

B. Procedures for the Analysis of Aqueous Hydrogen Peroxide in Rain Samples Collected During SECORE-88, S. R. Bachman and M. E. Peden

One of the measurement objectives of SECORE-88 was to obtain data on the H_2O_2 content of rain samples from convective showers. An efficient method, relatively free of interference, and with sufficient sensitivity to detect aqueous peroxide in rain samples, was sought. Below is a description of the techniques employed.

Samples were collected in 60 mL high density polyethylene bottles or 50 mL polypropylene centrifuge tubes, depending upon the type of automatic sequential sampler employed. Both the bottles and tubes and the collection surfaces of the automatic sampler were cleaned with deionized water. system blank samples, prepared by collecting water that had been poured over the collection surfaces and then allowed to accumulate in pre-cleaned bottles or tubes, were measured to assess the potential for contamination from the sample collection and handling procedures.

Upon cessation of a rain event, samples (rain and field blanks) were recovered from the field and transported to the lab and placed in a 4° C refrigerator. Within 24 hours, the samples designated for H_2O_2 analysis were numbered, weighed, and prepared for analysis. Samples were not filtered prior to analysis. Sample preparation involved addition of a fixing reagent, whereby a 2 mL aliquot of sample was removed from the collection bottle and added to an equal amount of a freshly prepared derivatizing (fixative) solution (see reagents). Analysis followed within one hour. Details of the fluorescence procedure for the determination of H_2O_2 are included in Kelly *et al.*, (1985) and Lazrus *et al.*, (1985).

C. Preliminary Data for SECORE-88 Case 1: June 8, 1988, V. C. Bowersox and M. E. Peden

On 8 June 1988 the 13 raingages and four sequential chemistry samplers in the SECORE-88 network all recorded the first precipitation event to occur after the network was activated on 20 May 1988. Rain fell from a stratiform deck associated with the passage of a mesoscale low that tracked across southern Illinois.

Rainfall amounts were light and relatively uniform across the network, with an average amount of 0.32 ± 0.04 inches and a range of 0.25 inches to 0.38 inches. Rain began at about 1600L and continued without interruption for 3 hours 10 minutes to 3 hours 20 minutes. Averaged over the entire event, the rainfall rates were about .09 - .11 inches/hour. during the first 5 minutes of the event, the rainfall rate was .16 - .20/hour, and for the duration of the event, the rates remained uniform and steady at .08 - .10 inches/hour.

At the outset of rain, the relative humidity at the surface measured 62% and it rose rapidly to 88% during the first 45 minutes. Thereafter as the rain rates decreased, the rise in surface relative humidity slowed until it reached 94%. Given the light rainfall rates and corresponding small drop sizes, the potential effects of evaporation on raindrop sizes and concentrations must be considered.

Another change which must be considered is the surface (10 meter) wind direction. For the first 30 minutes of the event, winds were out of 327°. At that point the wind direction began to back toward a more northerly direction until it reached $17^\circ \pm 4^\circ$, where it remained until the end of the event. Figure 1 displays the aqueous peroxide concentrations in samples collected at the Bondville field site during the 8 June event. The drop in concentrations from about 1 mg/L (-56μ equiv/L) to about 0.3 mg/L corresponds to the time period during which the wind direction shifted. One hypothesis, which must be tested, is that fresh SO_2 from Champaign may have accounted for the drop in H_2O_2 . There is no apparent rise in SO_4^{2-} concentrations to support this hypothesis, however (see Figure 2).

D. Research Summary Report to the National Acid Precipitation Assessment Program -- Task Group III Activities, V. C. Bowersox and R. G. Semonin

Studies of the scavenging of atmospheric pollutants that affect the acidity of precipitation have been done in pursuit of the goals of Task Group III: to investigate meteorological and chemical influences on acidic deposition. This effort has focused on the analysis of existing wet deposition data sets, such as NADP/NTN and MAP3S data, that have been collected in response to NAPAP needs to quantify spatial and temporal patterns and trends in wet deposition. These data sets have been used to identify the important chemical contributions to the ionic balance in precipitation samples, then to examine the influence of meteorological factors and source regions on concentrations of these constituents. Results of this work can provide insight into the important processes that modelers must consider to accurately estimate acidic deposition.

1. Except for coastal areas where marine salts are an important influence on the chemistry of precipitation, the pH of precipitation in the 31 eastern states of the U.S. can be calculated from the chemical balance of four inorganic anions and cations (SO_4^{2-} , NO_3^- , NH_4^+ , Ca^{2+}) in equilibrium with H^+ and HCO_3^- .
2. A second order empirical model of the measured pH versus the calculated pH can account for the weak inorganic acids, (formic and acetic) in

precipitation in the eastern U.S.

3. Soil aerosols from dust storms can be transported long distances (750-1500 km) and then deposited by precipitation, markedly affecting the concentrations of Ca^{2+} and other ions.
4. Highest SO_4^{2-} and NO_3^- fluxes from precipitation occur in the spring and summer and are associated with convective showers (often thunderstorms) in air that, relative to low flux cases, has had more passes and/or longer residence times over high emissions areas.
5. Near point and areas sources of urban pollution, chemical concentrations in precipitation are demonstrably increased, relative to concentrations at regional NADP/NTN sites.
6. Screening the dryfall bucket data from the NADP network results in physically consistent changes in the measured dry sedimentation fluxes, though independent studies are needed to test the validity and representativeness of these fluxes.

E. A Modified Successive Corrections Method that Meets Statistical Goals, J. Su and G. J. Stensland

The problem of determining the best placement of isopleth lines is of great importance for acid rain research. For a set of environmental data unevenly distributed in space and time, objective analysis is a method for estimating the regular grid point values from which isopleth lines can be drawn. There are many objective analysis schemes, including Thiessen polygons, polynomial fitting, variational method, smoothing isopleth lines, optimal interpolation, Kriging, and successive corrections.

Kriging and statistical methods in general are of great interest to acid rain research. Kriging calculates the interpolation weights by minimizing the standard error of interpolation and provides the uncertainty associated with the interpolated values. This is a powerful tool when comparing two sets of atmospheric chemistry data that vary in space and time. a major drawback of Kriging is the amount of computer resources that it requires. Utilizing the National Atmospheric Deposition Program/National Trends Network (NADP/NTN) data set for 1986 on a VAX 11/750, Kriging took 8.6 times longer to run than a successive corrections method and required three times more core memory. As the size of the data set becomes larger, these differences grow. Successive corrections schemes, on the other hand, are widely used in numerical weather modeling precisely because they require more modest computer resources. Unfortunately, the available successive

corrections methods do not incorporate the uncertainty of interpolation, nor can they calculate an optimal set of interpolation weights. It is our aim to implement error estimation for a successive corrections method and to formalize procedures for finding an optimal set of successive corrections parameters, thus maintaining the best of both Kriging and the successive corrections methods.

F. Dry Deposition Flux Estimates for Large-Particle Elements, from Dry-Bucket Measurements, D. F. Gatz, V. C. Bowersox and J. Su

There were two main purposes for this work. The first purpose was to explore methods for screening the data set, by:

1. Comparing distributions of dry deposition rates in dry "uncontaminated" samples with those in samples containing various single "contaminants" (including water), and
2. Determining the effects of various screening criteria on the statistical distributions of dry deposition rates measured by the NADP dry bucket samplers.

The second main purpose was to derive mean dry deposition fluxes for Ca, Mg, K, and Na for the eastern and western U.S., and for "dusty" and "clean" seasons.

This paper is based on analytical data for 2170 dry-bucket samples collected at original National Atmospheric Deposition Program (NADP) sites in what is now the combined National Atmospheric Deposition Program/National Trends Network (NADP/NTN). Samples were collected between 1978 and 1987, but the period of record varies between sites. Data from a small subset of the samples were examined previously by Hicks (1985), who found high frequencies of contaminated samples at many of the sampling sites.

G. Spatial Relationships Between Acid Rain, Air Quality, and Visibility Data, G. J. Stensland

Visibility and acid rain are two different phenomena which both have direct relationships to air quality. The objective of this study (Stensland, 1987) was to explore the relationships between all three data sets, with emphasis on the situations for the eastern U.S.

H. Synthetic Precipitation Reference Samples, M. E. Peden

The need for stable precipitation reference samples was described in the Twenty-First Progress report to the U.S. Department of Energy (Skowron and Peden, 1987). Although the National Bureau of Standards (NBS) has recently developed a Standard Reference Material (SR) for precipitation chemistry studies, the NBS solutions (SR 2694) do not include certified concentration values for nitrate, chloride, or ammonium. SRM's are also relatively expensive, so that their cost becomes prohibitive when they are used for several parameters by several chemists on a daily basis. Another drawback is that the certified concentrations of the parameters in SRM 2694 do not adequately reflect those actually found in wet-only deposition. NBS has also reported problems associated with the long-term stability of ammonium ion and pH in low ionic strength solutions (Koch, 1986).

Because of these limitations, our laboratory developed its own internal formulations with the following characteristics:

1. NBS traceable bias and precision
2. Long-term stability
3. Low cost
4. Analyze concentrations approximating the range found in natural precipitation

Preliminary data from the analysis of two different dilutions of the stock synthetic solution in 1987 indicated that, even at very low concentrations, these solutions were chemically stable for at least five months when stored at room temperature in high density polyethylene bottles.

The recent incorporation of these solutions into new and existing quality assurance programs has emphasized the need for low cost reference materials for precipitation chemistry studies. These solutions have been proven to be chemically stable when stored at room temperature and are formulated to provide the user with a choice of analyte concentrations depending on the dilution used when preparing samples from the concentrated stock solution. The reagents used in preparing the stock solution are currently available in most water chemistry laboratories or are readily available from chemical supply vendors.

VII. Review of Twenty-Third Progress Report, (Grant Number DE-FG02-88ER60635) Report Number C00-1199-67, SWS Contract Report 468, August 1989

- A. The Removal of Atmospheric SO_2 by Dry Deposited Road Dust, A. L. Williams
- B. Sensitivity Analyses of Area Weighted Average Algorithm, J. Su and G. J. Stensland

The computer program was developed on the VAX 11/750 using standard Fortran 77. An UNIX version and an IBM PC version of the program are also available. In this chapter we will discuss the algorithm as applied to calculating deposition to states. We digitized simplified state boundaries for all 48 contiguous states of the USA using the boundaries found in the graphics package distributed by the National Center for Atmospheric Research (NCAR, 1981). This step was necessary to minimize the computational time required by the algorithm so that it could be run on a PC. In this simplified scheme, each state is defined by an average of 30 points. A very small deviation in state area, less than one percent, was introduced by using the simplified boundaries.

The program itself consists of two parts. The first part reads in the data record for each site and calculate descriptive statistics for the variable of interest. This statistic along with the spatial coordinates of the data collection site are then piped into a spatial analysis routine. The spatial analysis subprogram interpolates the values of the environmental variable at the sites to the corners of each grid square. Any spatial analysis algorithm will suffice; some examples include Laplacian smoothing splines, potimum interpolation, and Kriging. We implemented an objective analysis scheme developed by Barnes (Barnes, 1964) and modified by Achtemeier (Achtemeier, et al., 1977). The second part of the algorithm reads in the digitized state boundaries and integrate the spatial field over each state to solve for the state area weighted average. The algorithm was described in detail in the last progress report.

C. A First Look at Data to Examine the Stability of Major Inorganic Ions in Rain Samples in Central Illinois, V. C. Bowersox and M. E. Peden

At the recent meeting of Multistate Atmospheric Power Production Pollution Study (MAP3S) scientists in Bethesda, MD, in April of this year, workshop sessions were held to highlight major accomplishments of MAP3S research over the last 10+ years, as well as important research needs of the future. Under research needs, one topic that received extensive discussion was the question of sample stability. Are event or daily samples chemically stable? Are the changes in concentrations of the major inorganic ions small over the daily (or event) sampling period used at MAP3S sites? Are chemical changes effectively arrested by refrigeration? Apart from the addition of TCM (a preservative for dissolved SO_2) to a sample aliquot used only for SIV) analyses, the only routine sample preservation procedure employed in the MAP3S network is refrigeration. What happens to the chemistry of a sample left in the sampler at ambient temperatures in the summer? The consensus of the group of researchers was that we do not know enough about the short term changes in the chemistry of samples held for 24-48 hours with no added preservative.

A study was designed to examine the stability of the chemistry of samples from a summer rain event. Major inorganic cations and anions were the focus of this study, since these all are reported by the networks presently in operation in North America. In a previous study, Peden and Skowron (1978) had found that bulk deposition samples exhibited large changes in Ca and Mg when left unfiltered at about 25C. The present study, however, focused on wet-only deposition samples, which contain much less dry deposited soil than bulk samples, because exposure is limited to the occurrence of precipitation.

D. Lead and Cadmium Loadings to the Great Lakes from Precipitation, D. F. Gatz, V. C. Bowersox and J. Su

We present the first analysis of precipitation chemistry data from the Great Lakes Atmospheric Deposition (GLAD) network of wet-only precipitation samplers. Lake loadings were calculated from 1- or 2-yr (1982-1983) volume-weighted mean ion concentrations and 30-yr mean annual precipitation amounts, using an objective analysis method to interpolate measurements to a 40 x 40 km square grid over the Great Lakes region. The results show that atmospheric loadings of Cd, Pb, and other cations to the Great Lakes are in many cases considerably smaller than those estimated from the bulk precipitation measurements of the early 1970s. For Pb, some portion of the differences can be accounted for by the decrease in emissions that resulted from reductions in the Pb content of gasoline. Valid dry deposition in the bulk samplers also accounts for a portion of the differences. However, for the most extreme differences, contamination of the bulk collectors by resuspended local surface dust is likely to be a major cause. Good agreement was noted between the precipitation-only loadings derived from the GLAD data and those recently given by Strachan and Eisenreich (1987).

VIII. Bibliography of Papers, Reports, and Interviews associated with the Study of Atmospheric Pollution Scavenging research over the ten year period 1980-1989

Brown, L., Stensland, G.J., Klein, J., and R. Middleton, 1989: Atmospheric deposition of ⁷Be and ¹⁰Be. Geochimica et Cosmochimica Acta, 53:135-142.

Gatz, D.F., Bowersox, V.C., and J. Su, 1988: "Screening criteria for NADP dry-bucket data," In: Proceedings 81st Annual APCA Meeting (paper 88-101.6), Air Pollution Control Association, Pittsburgh, PA, 16 pp.

Gatz, D.F., Bowersox, V.C., and J. Su, 1989: Lead and cadmium loadings to the Great Lakes from precipitation. J. Great Lakes Res., 15(2):246-264.

Stensland, G.J. and A.L. Williams, 1989: Dry deposited road dust as a major sink of atmospheric sulfur dioxide. EOS.

Gatz, D.F., Bowersox, V.C. and Su, J., 1988: "Dry deposition flux estimates for large-particle elements from dry-bucket measurements," presented at 1988 NADP Technical Committee Meeting, Champaign, IL, 24-27 October.

Gatz, D.F., Bowersox, V.C., and J. Su, 1988: "Dry deposition of large particles, based on NADP dry bucket measurements," presented at 1988 Rocky Mountain Conference, Denver, CO, 31 July - 5 August.

Gatz, D.F., Bowersox, V.C., and J. Su, 1988: Screening criteria for NADP dry bucket data," presented at Annual APCA Meeting, Dallas, TX, 20-24 June.

Gatz, D.F., Barnard, W.R., Bowersox, V.C., Stensland, G.J., and J. Su, 1988: "Base cations in precipitation: distributions and sources," presented at 9th Rocky Mountain ACS Regional Meeting, Las Vegas, NV, 27-30 March. 8-10 March.

Stensland, G.J., 1988: "Precipitation chemistry at MLO O-ring bias," presented at NOAA-GMCC Annual Meeting, Boulder, CO, 8-10 March. Su, J.,

Stensland, G.J., and V.C. Bowersox, 1988: "Spatial analysis of 1985 NADP/NTN sulfate and sodium data," presented at a mini-symposium, The Spatial Analysis of Wet Deposition Data, Champaign, IL, 27-28 October.

Bowersox, V.C., 1989: "What we have learned about the differences in daily or event samples and weekly NADP/NTN samples," presented at NADP Technical Committee Meeting, Provincetown, MA, 25 October.

Bowersox, V.C., 1989: "Wet deposition chemistry in the U.S. today- spatial patterns and temporal trends," presented at regional conference, Atmospheric Deposition in Pennsylvania: A Critical Assessment. University Park, PA, 11 September.

Bowersox, V.C., 1989: "Recent trends in the chemistry of atmospheric deposition in the eastern U.S.," presented at Acid Rain Workshop at 28th International Geological Congress, Washington, D.C., 15 July.

Bowersox, V.C., 1989: "Ecological concerns of acid rain," presented at Centel Corporation Executive Seminar, St. Charles, IL, 30 January.

- Bowersox, V.C., 1988: "Studies of pollutant scavenging," presented at NAPAP Task Group III Peer Review Meeting, Richmond, VA, 5 May.
- Bowersox, V.C., 1988: "Research at the Illinois MAP3S site," presented at MAP3S Review Meeting, Gaithersburg, MD, 26 January.
- Bowersox, V.C., 1989: "Acid Rain, an Update of Recent Research Findings," radio inter-view, WILL, Champaign, IL, 12 January
- Bowersox, V.C., 1988: "The acid rain problem," radio interview, WILL, Champaign, IL, 20 July.
- Bowersox, V.C., 1988: "SECORE-88 Field Project to Study Acid Rain," TV interview, WCIA, Champaign, IL, 7 July.
- Bowersox, V.C., 1988: "The 3CPO field project at the Water Survey," radio interview, WDWS, Champaign, IL, 8 June. Jan 88-
- Williams, A.L. and G.J. Stensland 1989: "The scavenging of sulfur dioxide by dry deposited road dust," In: Proceedings of the 1989 EPA/AWMA International Symposium on Measurement of Toxic and Related Air Pollutants in Raleigh, NC, May 1-5, 1989.
- Stensland, G.J., Krupa, S.V., and V. C. Bowersox, 1988: "Acid precipitation patterns and trends: a technical amplification of NAPAP's findings, In: Proceedings of APCA International Conference, Acidic Precipitation, a Technical Amplification of NAPAP's Findings, Pittsburgh, PA, January, pp 32-47.
- Bowersox, V.C., 1988: "Study of atmospheric pollution scavenging," In: Research Summaries for Task Group III, NAPAP, 722 Jackson Place, NW, Washington, DC. 20503.
- Semonin, R.G., Bowersox, V.C., Stensland, G.J., Peden, M.E., Su, J. and A.L. Williams, 1989: Twenty-Third Progress Report to the U.S. DOE, Office of Health and Environmental Research (Grant #DE-FG02-88ER60635) Washington, DC., 43 pp.
- Semonin, R.G., Bowersox, V.C., Stensland, G.J., Peden, M.E., Gatz, D.F., Bachman, S.R., Bachman, S.R., and J. Su, 1988: Pollution Scavenging. Twenty-Second Progress Report to the U.S. DOE, Office of Health and Environmental Research (Grant #DE-FG02-88ER60635) Washington, DC, 49 pp
- Gatz, D.F., and D.Q. Naiman, 1980: Spatial variability of rainwater impurities in mesoscale events. Paper presented at the symposium on Intermediate Range Atmospheric Transport Processes and Technology Assessment, Gatlinburg, TN, 1-3 October 1980.

Gatz, D.F., and D.Q. Naiman, 1981: Statistical analysis of NADP data and implications for network design. Paper presented at the ASTM Symposium and Workshop on Sampling and Analysis of Rain, Philadelphia, PA., 7 October 1981.

Gatz, D.F., B.K. Warner, and L.-C. Chu, 1982: Solubility of metal ions in rainwater. Paper presented at the American Chemical Society National Meeting, Las Vegas, NV, April.

Bridgman, H.A., and D.F. Gatz, 1982: Mesoscale spatial variability of sulfate in air and rain near St. Louis. Paper presented at the 4th International Conference on Precipitation Scavenging, Dry Deposition and Resuspension, Santa Monica, CA, 29 November- 3 December, 1982.

Gatz, D.F., B.K. Warner, and L.-C. Chu, 1982: Solubility of metal ions in rain water, Paper presented at the National Atmospheric Deposition Program Technical Committee Meeting, St. Louis, MO, 3-5 November.

Gatz, D.F., 1983: Source apportionment of rain water impurities in central Illinois, Paper presented at the Annual Meeting of the Air Pollution Control Association, Atlanta, GA, June.

Gatz, D.F., 1983: Source apportionment of rain water impurities in central Illinois, Paper presented at the Commission on Atmospheric Chemistry and Global Pollution International Symposium on Tropospheric Chemistry, Oxford, England, 29 August-3 September.

Gatz, D.F., B.K. Warner, and L.-C. Chu, 1983: Heavy metals: solubility in atmospheric deposition. Paper presented at the International Conference on Heavy Metals in the Environment, Heidelberg, West Germany, September.

Gatz, D.F., 1983: Source apportionment of rain water impurities in central Illinois, Paper presented at the National Atmospheric Deposition Program Technical Committee Meeting, Knoxville, TN, 6-10 November.

Gatz, D.F., 1984: A review of particle tracers of atmospheric processes, Keynote paper presented at the Department of Energy Workshop on Atmospheric Tracers, Santa Fe, NM, 22 May.

Gatz, D.F., W.R. Barnard, G.J. Stensland, and R.G. Semonin, 1986: Sources of cations in U.S. precipitation: implications of NADP/NTN data. Paper presented at the NADP Technical Committee Meeting, Atlanta, Georgia, 29 October.

Gatz, D.F., V.C. Bowersox, and J. Su, 1987: Screening criteria for NADP dry bucket data. Paper presented at the 1987 NADP Technical Committee Meeting, Minneapolis, MN, 2-5 November.

Semonin, R.G., 1981: Acid Rain in North America, talk, International Conference on Acid Deposition, Chicago, May 19, 1981.

Semonin, R.G., 1981: Implications of Recent Maps of Acid Rain to Long Range Transport, talk, U.S./ Canada Memorandum of Intent Review Panel, Washington, DC., September 21, 1981.

Bowersox, V.C., 1980: "Precipitation Chemistry Studies at the Illinois State Water Survey," talk, Coordinating Research Council-Air Pollution Research Advisory Committee, Champaign, IL., January 1980.

Bowersox, V.C., 1980: "Relative collection Efficiency of the NADP Wet/Dry Collector and a Weighing Bucket Raingage," presentation, NADP Technical Committee Meeting, St. Louis, MO, October 1980.

Bowersox, V.C., 1981: Seasonal Patterns of Sulfate and Nitrate in Precipitation in the United States," presented paper, 74th APCA Annual Meeting, Philadelphia, PA, 22 June 1981.

Bowersox, V.C., 1981: "Quality Assurance Program." presentation, site review team for DOE Contract E-AC02-76EV01199, Champaign, IL, 23 June 1981.

Bowersox, V.C., 1981: "Chemistry of Precipitation - IL vs PA," presentation, site review team for DOE Contract DE-AC0276EV01199, Champaign, IL, 23 June 1981.

Bowersox, V.C., 1981: "Intercomparison of the Chemistry of Precipitation from Daily (Event) Versus Weekly Sampling Protocols," presentation, National Atmospheric Deposition Program Technical Committee, St. Louis, MO, 2 November 1981.

Bowersox, V.C., 1982: "Feasibility Study to Assess the Impact of Precipitation Quality on Soil Water Quality," presentation, site review team for DOE Contract DE-AC02-76EV01199, Champaign, IL, 23 September 1982.

Bowersox, V.C., 1982: "Preliminary Report on CANSAP/NADP Intercomparison Study," presentation, NADP Technical Committee Meeting, St. Louis, MO., 4 November 1982.

Bowersox, V.C., 1982: "Characterization of the Inorganic Chemistry of Precipitation in North America," presented paper, 4th International conference on Precipitation Scavenging, Dry Deposition, and Resuspension, Santa Monica, CA., 2 December 1982.

Bowersox, V.C., 1983: "Characterization of the Wet Deposition Chemistry Over the United States from Recent Measurements," presented paper, 6th Midwest Water chemistry Workshop, Urbana, IL, 20 October 1983.

Bowersox, V.C., 1984: "Another Look at the Acid Rain Trend in the Eastern U.S.," talk, East Central Illinois Section of the American Society of Civil Engineers, Champaign, IL, 17 April 1984.

Bowersox, V.C., 1985: "Seasonal Variations in the Chemistry of Precipitation of the U.S." presented paper, 78th APCA Annual Meeting, Detroit, MI, 20 June 1985.

Bowersox, V.C., 1985: "Studies of Atmospheric," presentation, National Acid Precipitation Assessment Program review of Task Groups A-D, Boulder, CO, 11 September 1985.

Bowersox, V.C., 1985: "Comparison of Isopleth Maps Resulting from Objective and Subjective Analyses of NADP/NTN Data - Man Versus Machine." presented poster, National Atmospheric Deposition Program Technical Committee, Fort Collins, CO, 9 October 1985.

Bowersox, V.C., 1985: "Comparing Precipitation pH Averages." presentation, site review team for DOE Contract DE-AC02-76EV01199, Champaign, IL, 16 October 1985.

Bowersox, V.C., 1985: "Seasonal Variations in Wet Deposition in the U.S." presentation, site review team for DOE Contract DE-AC02-76EV01199, Champaign, IL, 17 October 1985.

Bowersox, V.C., 1985: "Precipitation Chemistry Patterns and Trends in the West." presentation, site review team for DOE Contract DE-AC02-76EV01199, Champaign, IL, 17 October 1985.

Bowersox, V.C., 1986: "Event and Weekly Precipitation Chemistry Samples - Are There Biases?" presentation, MAP3S/CORE Workshop, Champaign, IL, 16 April 1986.

Bowersox, V.C., 1986: "Standard Procedures for Wet Deposition Sampling - RADM Verification," presentation, Regional Model Evaluation Workshop on Quality Assurance, Toronto, Canada, 11 June 1986.

Bowersox, V.C., 1986: "Reconciling Sampling Protocols for the RADM Evaluation Field Study," presentation, RADM Evaluation Workshop, Toronto, Canada, 14 October 1986.

Bowersox, V.C., 1987: "Patterns and Trends of Sodium in Precipitation," presented poster, CACGP Symposium, Peterborough, Ontario, Canada, 27 August 1987.

Bowersox, V.C., 1987: "Gradient in Precipitation pH Across the Mid-Continental United States," presented paper, CACGP Symposium, Peterborough, Ontario, Canada, 28 August 1987.

Bowersox, V.C., 1987: "Atmospheric Chemistry at the Bondville Site -- A Local Climatology," presentation, Workshop of Cloud Chemistry, Cloud Physics Organization, Champaign, IL, 30 September 1987.

Bowersox, V.C., 1987: "pH Gradient across the Upper Mid-West," presented talk, NADP Technical Committee Meeting, Minneapolis, MN, 4 November 1987.

Bowersox, V.C., 1987: "Gradient in Precipitation pH Dry Bucket Sample Data," presented poster, NADP Technical Committee Meeting, Minneapolis, MN, 4 November 1987.

Miller, M.V., D.F. Gatz, G.J. Stensland, and A.C.D. Leslie, 1980: Aerosol Sources in Rural Central Illinois. Presented at American Society of Agronomy - Crop Science Society of America - Soil Science Society of America, Detroit, MI, 30 Nov. - 5 Dec.

Stensland, G.J. and R.G. Semonin, 1980: Network Operation-Siting. Presented at the NOAA Global Monitoring for Climate Change Annual Meeting, Special Session on Precipitation Chemistry, Boulder, CO, 6 March.

Stensland, G.J., 1981: Characterization of Acid at Hilo, Hawaii. Presented at the NOAA Global Monitoring for Climate Change Annual Meeting, Boulder, Co, 3-4 March.

Stensland, G.J., 1981: Characteristics and interpretation of the Ion Concentrations of Sequential Rain Samples from Hilo, Hawaii. Presented at IAMAP Meeting, Hamburg, F.R.G., Aug 17-28.

Stensland, G.J., 1982: Major Ion Correlations and Scavenging Ratios for Nine Convective Rain Shower Cases. Presented at the 4th International Conference on Precipitation Scavenging, Dry Deposition, and Resuspension, Santa Monica, CA, Nov. 29 - Dec. 3.

Stensland, G.J., 1984: A Critical Comparison of Data from Precipitation Chemistry Monitoring Networks - Past and Present. A refresher course presented at the 77th Annual meeting of the Air Pollution Control Assoc., San Francisco, CA, June 26.

Stensland, G.J., 1984: A Comparison of Methods of Computing Precipitation pH Averages. Paper presented at the Technical Committee Meeting of NADP (IR-7), Asheville, NC, Nov. 1.

Stensland, G.J., 1985: Potential for Bias and Outliers in NADP/NTN Data Due to the Sample container O-Ring. NADP Technical Committee - Abstracts of Papers. NREL, CSU, Fort Collins, CO.

Stensland, G.J., 1985: Preliminary Results of Beryllium-10 Measurements for Mauna Loa Observatory, HI, Rain Samples. Paper presented at the NOAA-GMCC Annual Meeting, Boulder, CO, March 14.

Gatz, D.F., W.R. Barnard, G.J. Stensland, and R.G. Semonin, 1986: Sources of Cations in Precipitation Chemistry: A Brief Review of the Issues. Water, Air, Soil Pollut., 30:245-251.

Stensland, G.J., V.C. Bowersox, and M.E. Peden, 1987: Patterns and Trends of Sodium in Precipitation. Abstract in NADP Technical Committee Meeting - Abstracts of Papers, NREL, CSU, Fort Collins, CO.

Semonin, R.G., 1982: North American Precipitation Chemistry, talk, AIAA 20th Aerospace Sciences Meeting, Orlando, FL, January 13, 1982.

Semonin, R.G., 1982: Is Acid Rain a Recent Problem?, talk, Interstate Coal Task Force Meeting, Indianapolis, IN, February 10, 1982.

Semonin, R.G., 1982: North American Precipitation Chemistry, invited talk, Edison Electric Institute Environmental Managers Meeting, New Orleans, LA, May 21, 1982.

Semonin, R.G., 1982: Precipitation Chemistry Sampling Requirements, talk, Precipitation Chemistry Workshop, Rensselaerville, NY, August 10, 1982.

Semonin, R.G., 1982: Characterizing North American Precipitation Chemistry, invited talk, National Symposium on Acid Rain, Pittsburgh, PA, October 7, 1982.

Semonin, R.G., 1982: Acid Rain: Is It a Recent Problem?, invited lecture, Sigma XI, Illinois State University, Normal, December 9, 1982.

Semonin, R.G., 1983: Characterization of North American Precipitation, paper, American Meteorological Society, conference on Climate/Energy Interactions, New Orleans, LA, January 11, 1983.

Semonin, R.G., 1983: Acid Rain Issues, invited talk, Illinois State Bar Association, Normal, April 28, 1983.

Semonin, R.G., 1983: Monitoring and Chemical Characteristics of Precipitation Quality in the U.S., talk, 12th Annual Illinois ENR Conference, Urbana, September 14, 1983.

Semonin, R.G., 1983: The Quality of Acidic Deposition Data, invited talk, Conference on Air Pollution and Forest Productivity, Washington, October 4, 1983.

Semonin, R.G., 1984: Tracer Applications to Precipitation Processes, talk at U.S. Dept. of Energy Workshop on Atmospheric Tracers, Santa Fe, NM, May 22, 1984.

Semonin, R.G., 1984: Precipitation Chemistry Research at the Water Survey,, talk at U.S. Dept. of Energy Program Managers Meeting, Washington, D.C., July 11, 1984.

Semonin, R.G., 1984: The Trend of Acid Rain in the Northeast U.S., talk as panel member at American Institute of chemical Engineers, St. Louis, MO, September 13, 1984.

Semonin, R.G., 1984: Acidic Deposition and Its Trends, talk at American Meteorological Society Chapter Meeting, Indianapolis, IN, October 25, 1984.

Semonin, R.G., 1984: NADP Dry Bucket chemistry, talk at National Atmospheric Deposition Program Technical Committee meeting, Asheville, NC, October 30, 1984.

Semonin, R.G., 1984: Assessment of Acid Rain Trends, talk, U.S. Dept. of Energy Division Managers meeting, Washington, D., November 8, 1984.

Semonin, R.G., 1984: Acid Rain and Its Trend in the U.S., talk, Southern Illinois University Coal Research Center Seminar, Carbondale, IL, November 29, 1984.

Semonin, R.G., 1985: Acid Rain Sampling, Analysis and Interpretation, talk, American Chemical Society, Peoria, IL, March 19, 1985.

Semonin, R.G., 1985: Precipitation Chemistry and Its Acidity, talk, University of Louisville Water Research Center Seminar, Louisville, KY, April 18, 1985.

Semonin, R.G., 1985: Wet Deposition Chemistry, invited talk, American Chemical Society Symposium, Arlington, VA, June 17, 1985.

Semonin, R.G., 1985: Acid Precipitation: Origins and Distribution in the United States, keynote talk, American Water Works Assoc. Symposium, Washington, DC, June 23, 1985.

Semonin, R.G., 1985: Rain Sampling, Analysis and Precipitation Scavenging Research at the Water Survey, talk, DOE Program Managers Meeting, Washington, DC, September 17, 1985.

Semonin, R.G., 1985: The Changing Acid Rain Picture, talk, University of Georgia Agricultural Experiment Station, Griffin, GA, September 19, 1985.

Semonin, R.G., 1985: Overview of Acid Rain Research, talk, DOE Review Meeting, Champaign, IL, October 17, 1985.

Peden, M.E., 1981: Analytical Protocols for the Sampling and Analysis of Atmospheric Deposition Samples, talk, ASTM Symposium on Sampling and Analysis of Rain, Philadelphia, October 13, 1981.

Peden, M.E., 1985: Theory Versus Practice in the Electrometric Determination of pH in precipitation, paper presented at the 27th Rocky Mountain Conference, Denver, CO, July, 1985.

Peden, M.E., 1987: Comparison of the Limit of Detection and Limit of Quantitation, paper presented at the 1987 Pacific Conference on Chemistry and Spectroscopy, Irvine, CA, October, 1987.

Semonin, R.G., 1982: Acid Rain Issues, radio interview, WILL Focus 580, Urbana January 26, 1982.

Semonin, R.G., 1982: Long-Range Transport and Acid Rain, talk, Edison Electric Institute News Media Conference, Washington, DC, April 19, 1982.

Semonin, R.G., 1982: National Acid Rain Issues, 1-hour radio discussion show, KNUS, Denver, Co. July 7, 1982.

Semonin, R.G., 1982: The Acid Rain Issue, 1-hour radio talk show, WDWS, Champaign, IL, July 22, 1982. Semonin, R.G., 1983: Effects from Acid Rain, radio tape interview, CBC radio, Toronto, Canada, January 3, 1983.

Semonin, R.G., 1983: Acid Rain Issues, 319NANADec 87 Illinois Press television tape, discussion panel, WILL-TV Urbana, March 16, 1983.

Semonin, R.G., 1983: The National Picture of Acidic Wet Deposition, media presentation, Chicago, IL, June 2, 1983.

Semonin, R.G., 1984: Acid Rain Economic Issues, Illinois Press television tape, panel discussion, Urbana, April 30, 1984.

Semonin, R.G., 1985: Acid Rain Research at the University of Illinois, TV interview, WILL-TV, June 7, 1985.

Bowersox, V.C., 1986: "Sampling for Radionuclides in Illinois Rainfall," newspaper interview, Copley News Service, Chicago, IL, 30 May 1986.

Bowersox, V.C., 1986: "Radionuclides from Chernobyl in Illinois Rain," news paper interview, Peoria Journal Star, Peoria, IL, 30 May 1986.

Bowersox, V.C., 1984: "What is Acid Rain?" radio interview, WBEZ National Public Radio, 3 January 1984.

Semonin, R.G., 1982: North American Acidic Precipitation Issues, testimony, U.S. Senate Energy and Natural Resources Committee Hearings on Acid Precipitation and the Use of Fossil Fuels, Washington, DC, August 19, 1982.

Semonin, R.G., 1984: North American Acidic Precipitation Issues, testimonies, U.S. House of Representatives Committee on Energy and Commerce Hearings, Chicago, March 5, 1984.

Semonin, R.G., J.D. Bartlett, V.C. Bowersox, D.F. Gatz, D.Q. Naiman, M.E. Peden, R.K. Stahlhut, and G.J. Stensland, 1980: —, 18th Progress Report, Contract No. EY-76-S-02-1199, U.S. Dept. of Energy, COO-1199-60, Illinois State Water Survey, Champaign, IL 61820, 81 pp.

Stensland, G.J., 1980: Acid Rain - Is It a National Problem? Proceedings of the 19th Ann. Conf. of Association of Conservation Engineers, Springfield, IL, 28 Sept. - 2 Oct., 1980. pp. 198-206.

Bowersox, V.C. and G.J. Stensland, 1981: Seasonal Pattern of Sulfate and Nitrate in Precipitation in the United States. Prepring 81-6.1, Proceedings of the 74th Ann. Meeting of the Air Pollut. Control Assoc., Philadelphia, PA, June 21-26, pp.

Hansen, A., G.M. Hidy, and G.J. Stensland, 1981: Examination of the Basis for Trend Interpretation of Historical Rain Chemistry in the Eastern United States. Report by Environmental Research and Technology, Westlake Village, CA, ERT Doc. No. P-A097R (earlier versions published in Aug. 1980 and Feb. 1981).

Stensland, G.J., 1981: Nitrate Measurements in Precipitation for the Study of Trends Patterns, and Scavenging Processes. Proceedings, U.S. EPA-ESRL Workshop on the Formation and Fate of Atmospheric Nitrates, Research Triangle Park, NC, Oct. 22-23, EPA-600.9-81-025, 96-117.

Shannon, J., J.C. Doran, T. Horst, D. Pack, C. Spicer, and G.J. Stensland, 1981: Air Quality Effects of Coal Technologies: Research Needs. Working Group Report in Environmental Effects of Coal Tech.: Research Needs, MITRE Corp., MTR-79W159-03 for the Fed. Interagency Comm. on Health and Env. Effects of Energy Technologies.

Gatz, D.F., G.J. Stensland, and M.V. Miller, 1982: Sources of Airborne Calcium in Rural Central Illinois. ACS Symposium Series No. 167, Atmospheric Aerosol: Source/Air Quality Relationships, 303-325.

Semonin, R.G., V.C. Bowersox, D.F. Gatz, M.E. Peden, and G.J. Stensland, 1981: Study of Atmospheric Pollution Scavenging, Dept. of Energy, COO-1199-63, Illinois State Water Survey, Champaign, IL. 126 pp.

Stensland, G.J., 1981: Acid Rain Issues. Proceedings of National Coil Coaters Association Fall Tech. Meeting, Chicago, IL, Sept. 30-Oct. 2-4.

Barrie, L., J. Hales, K. Anlauf, J. Wilson, A. Wiebe, D. Whelpdale, G.J. Stensland, and P. Summers, 1982: Monitoring and Deposition of Acid and Alkaline Materials - Preliminary Data Interpretation. U.S. Canada Mem. of Intent on Transboundary Air Pollution, Work Group 2 Rpt. 2F-1, 4-1 to 4-80.

Stensland, G.J., 1983: Working Group 2-Chairman's Summary. In: Proceedings: Advisory Workshop on Methods for Comparing Precipitation Chemistry Data at Rensselaerville, NY, Aug. 10-12, 1982. Report UAPSP-100, Southern Company Services, Birmingham, AL, pp. 2-16 to 1-23.

Stensland, G.J., 1983: Deposition Monitoring. Chapter 8 In: The Acidic Deposition Phenomenon and Its Effects, Critical Assessment Review Papers, Vol. 1, Atmospheric Sciences, Public Review Draft. EPAP-600/8-83-016A.

Stensland, G.J., 1983: Acid Rain. Encyclopedia of Environmental Science and Engineering. [J.R. Pfafflin and E.N. Ziegler, Eds.], Vol. 1, Second Edition, Gordon and Breach Science Publishers, New York, 1-9.

Dolske, D.A. and G.J. Stensland, 1983: A Comparison of Ambient Airborne Sulfate and Concentrations Determined by Several Different Filtration Techniques. Proceedings of the 3rd National Symposium on Recent Advances in Pollutant Monitoring of Ambient Air and Stationary Sources, Raleigh, NC, May 3-6, 10 pp.

Stensland, G.J., 1984: Deposition Monitoring, Chapter In: The Acidic Deposition Phenomenon and Its Effects, Critical Assessment Review Papers, Vol. I, Atmospheric Sciences, Final Report. EPA-600/8-83-016AF, pp. 8-1 to 8-17 and 8-31 to 8-71.

Miller, J.M., G.J. Stensland, and R.G. Semonin, 1984: The Chemistry of Precipitation for the Island of Hawaii. NOAA Tech. Memorandum ERL ARL-133, Silver Spring, MD, 33 pp.

Stensland, G.J., Whelpdale, D.M., and G. Oehlert, 1986: Precipitation Chemistry, In: Acid Deposition Long-Term Trends, National Academy Press, Washington, D.C., pp 128-199.

Gatz, D.F., 1980: An urban influence on deposition of sulfate and soluble metals in summer rains. In: Shriner, D.S., C.R. Richmond, and S.E. Lingberg, Eds., Atmospheric Sulfur Deposition, Environmental Impact, and Health Effects, Ann Arbor Science Publishers, Ann Arbor, MI, pp. 245-261.

Gatz, D.F., and D.Q. Naiman, 1980: Spatial variability of rain water impurities in mesoscale events. Proceedings, Symposium on Intermediate Range Atmospheric Transport Processes and Technology Assessment, Gatlinburg, TN, 1-3 Oct 1980, pp 327-334. Available as CONF-801064 from NTIS, Springfield, VA.

Gatz, D.F., G.J. Stensland, M.V. Miller, and A.C.D. Leslie, 1981: Sources of airborne calcium in rural central Illinois. In: Macias, E.S. and P.K. Hopke, Eds, Atmospheric Aerosol: Source/Air Quality Relationships, ACS Symposium Series 167, American Chemical Society, Washington, D.C., pp. 303-325.

Davis, B.L., L.R. Johnson, R.K. Stevens, D.F. Gatz, and G.J. Stensland, 1982: Observation of sulfate compounds on filter substrates by means of x-ray diffraction. In: Shryer, D.R., Ed., Heterogeneous Atmospheric Chemistry. Geophysical Monograph 26, American Geophysical Union, Washington, DC., pp. 149-156.

Gatz, D.F., B.K. Warner, and L.-C. Chu, 1983: Solubility of metal ions in rain water. In: Teasley, J., and B.B. Hicks, Eds., Deposition Both Wet and Dry, Ann Arbor Science Publishers, Ann Arbor, MI, pp. 133-151.

Gatz, D.F., B.K. Warner and L.-C. Chu, 1983: Metal solubility in atmospheric deposition. Proceedings, International Conference on Heavy Metals in the Environment, Heidelberg, West Germany, 6-9 September 1983, Volume 1, pp. 183-186.

Gatz, D.F., 1983: Source apportionment of rain water impurities in central Illinois. Paper 83-28.3, Proceedings of the 76th Annual Meeting of the Air Pollution Control Association, Pittsburgh, PA.

Gatz, D.F., L.-C. Chu, 1986: Metal solubility in atmospheric deposition. In: Nriagu, J.O. and C.I. Davidson, Eds., Metals in the Air, Advances in Environmental Science and Technology Series, John Wiley, NY, pp. 391-408.

Gatz, D.F., 1986: A review of particle tracers of atmospheric processes. In: Heiken, J., Ed., Atmospheric Tracers, Noyes Publishing Co., Park Ridge, New Jersey.

Bowersox, V.C., 1986: "Collection, Analysis, and Screening of Wet Deposition Samples." In: Proceedings Methods for Acidic Deposition Measurement, Envirotech, Seattle, WA. pp. 3-19 to 3-20.

Peden, M.E., 1983: Sampling, Analytical, and Quality Assurance Protocols for the National Atmospheric Deposition Program. Sampling and Analysis of rain, ASTM STP 823, S.A. Campbell, Ed., pp. 72-83.

Semonin, R.G., 1982: Seasonal Precipitation concentrations and Depositions for North America from the CANSAP/NADP Networks, Reprint 82-03343, AIAA 20th Aerospace Sciences Meeting, 6 pp.

Semonin, R.G., 1981: Surface weather conditions. METROMEX: A Review and Summary, AMS Meteorological Monograph, 18, 17-40.

Semonin, R.G., 1981: Cloud characteristics. METROMEX: A Review and Summary, AMS Meteorological Monograph, 18, 63-73.

Semonin, R.G., 1982: Acid Rain: Is It a Recent Problem? Proceedings of the Illinois Mining Institute, Urbana, IL, pp 19-31.

Semonin, R.G., 1982: Long Range Transport and Acid Rain. Proceedings Ninth Annual Illinois Energy Conference, Chicago, IL pp 43-45. CAST Task Force on Acid Rain, 1983: Acid Deposition in Relation to Agriculture, Forestry, and Aquatic Biology. Council for Agriculture and Science and Technology Report No. 100, 54 pp.

Semonin, R.G., 1983: Acid rain data collection, handling, analysis and interpretation. Proceedings of Air Pollution and the Productivity of the Forest, Isaac Walton League, Washington, 11-48.

Semonin, R.G., 1984: Monitoring and chemical characteristics of precipitation quality in the United States. Proceedings of the 12th ENR Conference, Springfield, IL, 182-218.

Doty, K.G. and R.G. Semonin, 1984: A Case Study of the Effects of Local and Distant Sources of Alkaline Materials on Precipitation in the Midwest and Plains. Paper 84-20.3, 77th Annual APCA Meeting, San Francisco, CA, 21 pp.

Semonin, R.G., 1984: Trace Applications for the Study of Precipitation Processes, In: Proceedings of the Atmospheric Tracers and Tracer Application Workshop, Santa Fe, NM, 25 pp.

Miller, J.M. and G.J. Stensland, and R.G. Semonin, 1984: The Chemistry of Precipitation for the Island of Hawaii During the HAMEC Project, NOAA Tech. Memo ER1 ARL-133, Silver Spring, MD, 33 pp.

Semonin, R.G., 1985: Acid Precipitation: Origins and Distribution in the United States, In: Proc. AWWA Symposium on Acid Rain, American Water Works Assoc., Denver, CO, pp. 3-29.

Semonin, R.G., 1985: Has the Acidity of Precipitation in the United States Changed Since 1955? UARG Report, Washington, DC, 9 pp. Semonin, R.G., 1985: Wet and Dry Deposition in the United States, UARG Report, Washington, DC, 34 pp.

Semonin, R.G., 1986: Wet Deposition Chemistry, In: Materials Degradation Caused by Acid Rain, ACS Symposium Series 318, American Chemical Society, Washington, DC, pp. 23-41.

Keller, B.J., M.E. Peden, and A. Rattonetti, 1984: Graphite-Furnace Atomic Absorption Method for Trace Level Determination of Total mercury. Anal. Chem., 56:2617-2618.

Doty, K.G., 1986: "Spatial patterns of NADP Sites in the Eastern U.S. for Specific Ions Using Cluster Analysis. In: Transactions of the Air Pollution Control Association's Second International Conference on the Meteorology of Acidic Deposition, Albany, New York, March 17-20, 1986, pp. 365-380.

Doty, K.G. and R.G. Semonin, 1987: Two Case Studies of the Transport of dust Storm Aerosol and the Potential Incorporation into Precipitation. J. Clim. Appl. Met., 26: 763-773.

Bachman, S.R. and M.E. Peden, 1987: Determination of Organic Acid Anions in Precipitation by Ion Exclusion Chromatography. Water, Air, Soil Pollut., 33: 191-198.

Peden, M.E. and J. Rothert, 1987: Acid Rain: A Historical Look, Standardization News, April, pp. 42-45.

Brennan, C.J. and M.E. Peden, 1987: Theory and Practice in the Electrometric Determination of pH in Precipitation. Atmos. Environ., 21: 4, 901-907.

Bowersox, V.C. and R.G. de Pena, 1980: Analysis of Precipitation Chemistry at a Central Pennsylvania Site. J. Geophys. Res., 85: 5614-5620.

Pena, J.A., de Pena, R.G., Bowersox, V.C., and J.F. Takas, 1982: SO₂ Content in Precipitation and Its Relationship with Surface Concentrations of SO₂ in the Air. Atmos. Environ., 16: 1711-1715.

Semonin, R.G. and V.C. Bowersox, 1983: "Characterization of the Inorganic Chemistry of the Precipitation of North America," In: Precipitation Scavenging, Dry Deposition, and Resuspension, Vol. 1, Pruppacher, Semonin, and Slinn, Eds., Elsevier, New York, NY., pp. 191-201.

Stensland, G.J. and V.C. Bowersox, 1984: "Quality Assurance in Acid Precipitation Monitoring Through the Use of Ion Balance Calculations." In: Quality Assurance in Air Pollution Measurements, Johnson and Penkala, Eds., Air Pollution Control Association, Pittsburgh, PA, pp. 425-443.

Bigelow, D.S., Still, M.E. and V.C. Bowersox, 1984: "Quality Assurance Considerations for Multiple-Network Data Comparison." In: Quality Assurance in Air Pollution Measurements, Johnson and Penkala, Eds., Air Pollution control Association, Pittsburgh, PA, pp. 468-487.

Bowersox, V.C., Stensland, G.J. and M.E. Peden, 1987: "Evaluation of Bias in Wet Deposition Loading from the National Atmospheric Deposition Program/National Trends Network (NADP/NTN)." In: Proceedings 80th Annual APCA Meeting (paper 87-91.6), Air Pollution Control Association, Pittsburgh, PA. 13 pp.

Gatz, D.F., 1980: Associations and spatial relationships among rainwater constituents. J. Geophys. Res., 85, 5588-5598.

Ochs, H.T., and D.F. Gatz, 1980: Water solubility of atmospheric aerosols. Atmos. Environ., 14, 615-616.

Gatz, D.F., J. Bartlett, and J.J. Bassett, 1981: Metal pollutants in agricultural soils and the St. Louis urban rainfall anomaly. Wat., Air, Soil Pollut., 15, 61-75.

Gatz, D.F., 1984: Source apportionment of rainwater impurities in central Illinois, Atmos. Environ., 18, 1895-1904.

Gatz, D.F., W.R. Barnard, and G.J. Stensland, 1986: The role of alkaline materials in precipitation chemistry: A brief review of the issues. Wat., Air, Soil Pollut., 30, 245-251.

Evans, L.S., G.R. Henrey, G.J. Stensland, D.W. Johnson, and A.J. Francis, 1981: Acidic Precipitation: Considerations for an Air Quality Standard. In: Proceedings of Air Pollut. Control Assoc. Speciality Meeting - the Proposed SO₂ and Particulate Standards, Atlanta, GA, 16-18 Sept., 1980 and Water, Air, Soil Pollut., 16: 469-509.

Stensland, G.J. and R.G. Semonin, 1982: Another Interpretation of the pH Trend in the United States/ Bull. Amer. Meteorol. Soc., 63: 1277-1284.

Davis, P.L., L.R. Johnson, R.K. Stevenes, D.F. Gatz, and G.J. Stensland, 1982: Observation of Sulfate Compounds on Filter Substrate by means of X-Ray Diffraction. In: Heterogeneous Atmospheric Chemistry, Geophysical Monograph Series, 26: AGU, 149-156.

Stensland, G.J., L. Brown, J. Klein, and R. Middleton, 1983: Beryllium-10 in rain, Abstract in EOS, Transactions, AGU, 64: 18, 283.

Stensland, G.J. and R.G. Semonin, 1983: Response to "Comments on Another Interpretation of the pH Trend in the United States," by J.P. Bruce. Bull. Amer. Meteorol. Soc., 64: 1082-1083.

Stensland, G.J. and R.G. Semonin, 1984: Response to "Comments on Another Interpretation of the pH Trend in the United States." by T.J. Butler, C.V. Cogbill, and G.E. Likens. Bull. Amer. Meteorol. Soc., 65: 640-643.

Stensland, G.J. and V.C. Bowersox, 1984: A Comparison of Methods of Computing Precipitation Averages. Preprint 84-19.1, Proceedings of the 77th Annual Meeting of the Air Pollut. Control Assoc., San Francisco, CA, June 24-29, 16 pp. Semonin, R.G. and G.J. Stensland, 1984: Acid Rain Trends? Weatherwise, 37-5, 250-251.

Stensland, G.J., V.C. Bowersox, : Quality Assurance in Acid Precipitation Monitoring Through the Use of Ion Balance Calculations. Transactions (peer reviewed) of the Air Pollution Control Association Specialty Meeting "Quality Assurance in Air Pollution Measurements." Boulder, CO, Oct. 15-18, 1984. pp 425-443.

Bowersox, V.C. and G.J. Stensland, 1985: Seasonal Variations in the Chemistry of Precipitation of the United States. Paper 85-6A.2 I₁: Proceedings of the 78th Annual meeting of the Air Pollut. Control Assoc., Detroit, MI, June 17-21, 1985. 18 pp.

Summers, P.W., G.J. Stensland, and V.C. Bowersox, 1986: Acidic Deposition in Eastern North America. Paper presented at the International Symposium on Acidic Precipitation, Muskoka, Ontario, Sept. 15-20, 1985. Water, Air, Soil Pollut. 31: 528-535. SWS Reprint 738.

Stensland, G.J., 1987: Spatial Relationships Between Acid Rain, Air Quality and Visibility Data. Paper 87-40A.6 in the Proceedings of the 80th Annual Air Pollut. Control Assoc. Meeting, New York, NY, 16 pp. SWS Reprint 841.

END

**DATE
FILMED
416192**