DOE/ER/13368--9

DE93 007266

ENZYMOLGY OF ACETONE-BUTANOL-ISOPROPANOL FORMATION

Progress Report

January 1, 1992 - December 31, 1992

Jiann-Shin Chen

Virginia Polytechnic Institute and State University

Blacksburg, Virginia 24061

RECEIVED

FEB 10 1993

OSTI

January 1993

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY

UNDER GRANT NUMBER DE-FG05-85ER13368

MASTER

EP

I. Research Objectives

This research project is concerned with a process through which anaerobic bacteria convert various carbohydrates into useful chemicals, including acetone, butanol, and isopropanol, which are traditionally called solvents. The long-term goal of the project is to understand the fundamental properties of the process so that it would be possible to prevent degeneration of producing strains, to allow positive control of the solventogenic switch, and to regulate the product pattern according to market needs. Our approach is to elucidate first the molecular properties of solvent-producing enzymes and then to apply the information gained from the enzymological study to investigate control mechanisms for the solvent-producing pathways and the expression of solvent-production genes.

Our research primarily involves Clostridium beijerinckii, which is a species containing strains with the rare ability to produce the useful chemical isopropanol, in addition to acetone, butanol and ethanol. In related studies, we also included another solvent-producing organism, Bacillus macerans, which is a facultative anaerobe and produces a high level of acetone and ethanol under anaerobic conditions. Because B. macerans does not produce butyric acid or butanol, it provides a simpler system for the study of the acetoacetyl-CoA-reacting enzymes and also an organism outside the clostridial group for a mechanistic study of the solventogenic switch.

The objectives for the report period were

- (1) To purify and characterize distinct forms of alcohol dehydrogenases.
- (2) To purify and characterize acetoacetyl-CoA-reacting enzymes.
- (3) To study the organization of solvent-production genes.

II. Summary of Scientific Progress..

(1) Alcohol dehydrogenases (ADHs).

(a) Primary/secondary ADH. Primary/secondary ADH has been purified from two strains of isopropanol-producing C.

beijerinckii and extensively characterized. Two important properties (coenzyme specificity and quaternary structure) of the ADH were further examined with the purified enzyme. The primary/secondary ADH is practically an NADP(H)-dependent enzyme; however, an NAD⁺-linked activity has now been detected with a much more sensitive assay, which uses tetrazolium dye to reoxidize NADH. The result provides another example that the specificity of an enzyme for NAD(H) or NADP(H) is not absolute, although it can be very selective toward one of them.

The primary/secondary ADHs purified from C. beijerinckii and Thermoanaerobium brockii share a 75% identity in their amino acid sequences and a similar subunit molecular weight of about 38,000. Nevertheless, their native molecular weights differ significantly. We thoroughly re-examined the native molecular weight of the two purified ADHs from C. beijerinckii and the extensively purified ADH from T. brockii under identical conditions. The previously observed difference in molecular weights was confirmed: 100,000 for the C. beijerinckii enzyme and 150,000 for the T. brockii enzyme. The apparent difference in the quaternary structure between the ADHs from the two organisms is intriguing and requires further investigation.

(b) Primary ADHs. C. beijerinckii NRRL B592 produces acetone, butanol and ethanol but not isopropanol, and only primary ADH activities are detected in this organism. During the purification of primary ADH from C. beijerinckii NRRL B592, two types of ADH with distinct coenzyme specificity and stability were clearly separated by chromatography on either Reactive Red 120- or Cibacron Blue 3GA-agarose. One of them was NADPH-dependent (the P-ADH), whereas the other could use either NADH or NADPH as the coenzyme (the D/P-ADH). The P-ADH was more thermostable and was not inactivated by O₂. The D/P-ADH, which is the predominant ADH in solvent-producing cells, has been purified to a state that it contained only ADH. The D/P-ADH had a native molecular weight of 80,000. The seemingly homogeneous D/P-ADH consistently gave two equally intense bands (Coomassie blue stained; mol. wt. 40,000 and 43,500) after electrophoresis

on polyacrylamide gel in the presence of SDS. Furthermore, three protein bands that all showed NADH- and NADPH-linked ADH activities were observed when the seemingly homogeneous D/P-ADH was electrophoresed under non-denaturing conditions. Results from subunit analysis and peptide mapping of the three electrophoretically separated forms of D/P-ADH suggest that the ADH in the upper, middle, and lower bands had, respectively, the following subunit composition: α_2 , $\alpha\beta$, and β_2 , with α and β designating subunits with molecular weights of 40,000 and 43,500, respectively. The presence of both homo- and hetero-dimers of ADH in a prokaryote is uncommon. Also, it indicates that C. beijerinckii NRRL B592 contains at least four distinct primary ADHs. The role of individual forms of ADH in the solvent-producing process will need to be defined.

(2) Acetoacetyl-CoA-reacting enzymes.

(a) 3-Hydroxybutyryl-CoA dehydrogenase (3HBDH).

3HBDH, which catalyzes the first committed reaction toward butanol production, was purified from C. beijerinckii NRRL B593 during the previous reporting period. Characterization of the enzyme was completed during 1992, and the results have been published. This portion of the study was mainly funded by Neste Oy and the State of Virginia, but it is an integral part of our effort to define the enzymology of acetone-butanol-isopropanol formation.

(b) Acetoacetyl-CoA:acetate/butyrate CoA-transferase (CoA-transferase). During 1992, we have purified to homogeneity CoA-transferase from C. beijerinckii NRRL B593. CoA-transferase catalyzes the first specific reaction toward the formation of acetone and isopropanol. The CoA-transferase of solvent-producing clostridia is an unstable enzyme, and it requires high concentrations of ammonium sulfate (0.5-0.75 M) and glycerol (15-20% vol/vol) in 25 mM MOPS [3-(N-morpholino)-propanesulfonic acid] buffer at pH 7.0 to preserve the activity during the purification. The purified C. beijerinckii CoA-transferase had a native molecular weight of 85,000, and two types of subunits with

apparent molecular weights in the range of 28,000 and 23,000. The apparent K_m values of the C. beijerinckii CoA-transferase for acetate and butyrate were significantly lower than those of the C. acetobutylicum CoA-transferase. Whether or not the difference is physiologically significant will need to be examined.

(3) Organization of solvent-production genes.

Only a limited survey of the genomic region containing the structural gene (adh) for the primary/secondary ADH has been conducted with 12 strains of isopropanol-producing C. beijerinckii. In 8 strains, an EcoRI fragment, ranging in size from 2.8 to 4 kilobases, were found to hybridize to a probe specific for the N-terminal region of the ADH. In the other 4 strains, two EcoRI fragments, 2- and 2.5-kilobases in length, showed homology to the probe. The 12 strains fell into four classes based on the number and size of their adh-containing EcoRI fragments. The next step is to see if the putative transcription regulator gene (stc), which we observed in C. beijerinckii NRRL B593 directly upstream to the adh gene, occurs near the adh gene. The stc gene is potentially related to the onset of the solventogenic switch.

III. Publications.

A. Journal article:

1. Colby, G. D., and J.-S. Chen. 1992. Purification and properties of 3-hydroxybutyryl-coenzyme A dehydrogenase from Clostridium beijerinckii ("Clostridium butylicum") NRRL B593. Appl. Environ. Microbiol. 58:3297-3302.

2. Ismaiel, A. A., C.-X. Zhu, G. D. Colby, and J.-S. Chen. Purification and characterization of primary/secondary alcohol dehydrogenase from two strains of Clostridium beijerinckii. (to be submitted)

B. Book Chapter:

1. Chen, J.-S. 1993. Properties of acid- and solvent-forming enzymes of clostridia. In D. R. Woods (ed.), The clostridia and biotechnology. Butterworth-Heinemann, Stoneham, Mass. (in press)

C. Abstracts:

1. Colby, G. D. and J.-S. Chen. 1992. Purification and characterization of 3-hydroxybutyryl-CoA dehydrogenase from Clostridium beijerinckii strain NRRL B593. Abstr. 92nd General Meeting Am. Soc. Microbiol., K-76, p.269.

2. Rifaat, M. and J.-S. Chen. 1992. A transcriptional regulatory gene upstream from the adh gene of Clostridium beijerinckii NRRL B593: cloning and sequence analysis. Abstr. 92nd General Meeting Am. Soc. Microbiol., O-66, p.320.

3. Ismaiel, A., R.-T. Yan, and J.-S. Chen. 1993. Multiple primary alcohol dehydrogenases in Clostridium beijerinckii NRRL B592. Abstr. 93rd General Meeting Am. Soc. Microbiol. (submitted)

IV. Appendix

Reprints or preprints of the listed articles and abstracts are included as Appendices #1 to #5.

Reprints + Preprints removed

END

**DATE
FILMED**

3 / 17 / 93