Classical Theory of Continuous Spin and Hidden Gauge Invariance

D. Zoller

DOE/ER/40272--126

DE92 008777

*Physics Department
University of Florida
Gainesville, Fl 32611*

Abstract: We present a classical higher derivative point particle theory whose quantization gives Wigner's continuous spin representation of the Poincaré group. Although the theory is not reparametrization invariant in the usual sense, it does possess a hidden gauge invariance that provides a non-local representation of the reparametrization group. The Hamiltonian of the theory does not vanish and its value is the continuous spin parameter. The theory presented here represents the simplest example of a wide class of higher derivative theories possessing a hidden gauge invariance.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Wigner's classification of the unitary representations of the Poincaré group¹ constitutes one of the most basic organizational principles of particle physics. Wigner found two classes of massless representations—one with a scale and one without. Ordinary massless particles such as the photon are described in a scale free fashion by the invariant value of the helicity. The other massless representations are described by the invariant length of the Pauli-Lubanski vector. These are the continuous spin representations² so called because there are a continuum of polarization states available. Here we shall show that the higher derivative theory defined by the length of the acceleration is a classical theory of continuous spin.

Our motivation is twofold. On the one hand, knowledge of the classical theory may provide a better understanding of the role of the continuous spin representation in nature. Do continuous spin particles violate causality? How does it couple to gravity? On the other hand, the higher derivative theory defined by the length of the acceleration is the simplest example of a wide variety of theories possessing a hidden reparametrization invariance. The ordinary massless representations of the Poincaré group such as the photon can be obtained from a classical higher derivative action^{3,4} that possesses both the hidden reparametrization invariance and the ordinary reparametrization invariance. The Lagrangian is the product of the arc length and the curvature. The hidden gauge invariance represents an alternative to local supersymmetry for describing spinning particles. The string Lagrangian defined by the product of the element of area and curvature (distinct from the Polyakov term⁵ which is quadratic in the curvature) also possesses a hidden gauge invariance⁶ and is a good candidate for a quantum mechanically sensible bosonic string theory. This string theory is an example of a higher derivative field theory possessing a hidden gauge invariance and one can contemplate other higher derivative field theories possessing a hidden gauge invariance.

That there is a connection between Wigner's continuous spin representation and point

particle theories with one higher derivative than usual⁷ can be seen by examining the canonical variables. Ostrogradsky⁸ first formulated higher derivative theories in Hamiltonian form and the generalized canonical coordinates are

$$\begin{aligned} q' &= \dot{q} \\ p' &= \frac{\partial L}{\partial \ddot{q}} \\ p &= \frac{\partial L}{\partial \dot{q}} - \frac{d}{dt} \left(\frac{\partial L}{\partial \ddot{q}} \right) \end{aligned}$$

where the basic dynamical variable is q and q, p and q', p' are the canonically conjugate phase space variables. The phase space is doubled as is appropriate for a fourth order differential equation. Wigner phrased his continuous spin wave equation² in terms of a field $\Psi = \Psi(x, x')$ where x is the position four vector and x' is a translationally invariant four vector. This wave function corresponds precisely to the Schrodinger wave function of the higher derivative point particle theory as the configuration space is x, x' . In terms of the usual Schrodinger representation of the canonically conjugate variables X, P and X', P' , Wigner's continuous spin wave equation is

$$P^2 \Psi = 0$$

$$P \cdot P' \Psi = 0$$

$$P'^2 \Psi = \Psi$$

$$P \cdot X' \Psi = E \Psi$$

where the real number E is called the continuous spin parameter.

There is, however, an obstacle to obtaining this theory from a reparametrization invariant action. Noether's theorem applied to the local symmetry of reparametrization invariance implies the Hamiltonian vanishes and $p'q' = 0$. Since $P'X' = 0$ is not one of the continuous spin equations, no reparametrization invariant theory could give rise to

the continuous spin wave equation. This fact leads one to contemplate constrained higher derivative theories that are not reparametrization invariant.

Constraints occur generically in higher derivative theories if the Lagrangian is homogeneous of degree one in the accelerations. The simplest constrained theory is the length of the acceleration

$$L = \sqrt{\ddot{x}^2}$$

where $\dot{x}$ means the derivative of x with respect to the evolution parameter t and the Minkowski metric has signature $+++ -$. Here the coupling constant with dimensions of inverse length has been set to one. The Hamiltonian which follows from Noether's theorem applied to t translation symmetry

$$H = P \cdot X' + P' \cdot \ddot{X} - L$$

reduces to

$$H = P \cdot X'.$$

The complete set of first class constraints are

$$P'^2 = 1$$

$$P \cdot P' = 0$$

$$P^2 = 0.$$

The first constraint is primary and occurs because the Lagrangian is homogeneous of degree one much like the mass shell condition of the arc length theory. The second constraint is secondary and the third is tertiary. The quantization of this theory immediately gives Wigner's continuous spin wave equation. The three constraints promoted to conditions on the Schrodinger wave function $\Psi(x, x')$ provide the first three of Wigner's equations. The last is the Schrodinger wave equation restricted to the energy eigenvalue E . The value

of the Casimirs of the Poincaré group can be calculated classically from the constraints. The value the length W of the Pauli-Lubanski vector turns out to be H where the Pauli-Lubanski vector is

$$W^\mu = \epsilon^{\mu\nu\alpha\beta} P_\nu M_{\alpha\beta}.$$

Note that the angular momenta are those following from Noether's theorem

$$M^{\alpha\beta} = X^{[\alpha} P^{\beta]} + X'^{[\alpha} P'^{\beta]}.$$

The theory defined by the length of the acceleration when restricted to a particular energy E therefore provides an irreducible representation of the Poincaré group with zero mass and continuous spin parameter E . We shall now examine some features of the classical theory.

The general solution of the classical equation can be easily found in the Hamiltonian form the theory. The Hamiltonian is not unique but ambiguous up to an arbitrary function $\lambda(t)$ of the evolution parameter times the primary constraint

$$H = P \cdot X' + \lambda(P'^2 - 1).$$

Eliminating X' from the equation of motion for X and X' gives

$$\ddot{X}^\mu = 2\lambda P'^\mu.$$

The equation of motion for P'

$$\dot{P}'^\mu = -P^\mu$$

is solved by

$$P'^\mu = P_o'^\mu - P^\mu t.$$

The acceleration therefore is

$$\ddot{X}^\mu = 2\lambda(P_o'^\mu - P^\mu t)$$

and integrating twice gives the general solution

$$X^\mu = X_o^\mu + X'^\mu_o t + \int^t \int 2\lambda(P'^\mu_o - P^\mu t) dt$$

The continuous spin parameter of this solution is $H = P \cdot X'_o$ and its four-momentum is P^μ .

The acceleration theory of continuous spin allows us to address the issue of the causality of the classical solution. Recall that the arc length theory with the wrong sign with respect to our choice of metric

$$L = m\sqrt{\dot{x}^2}$$

gives a tachyonic representation of the Poincaré group and provides the basis for the claim that tachyons are acausal. The four velocity of the trajectory

$$\frac{dx^\mu}{dx^o} = \frac{P^\mu}{P^o}$$

is a space like gauge invariant observable and hence the theory is acausal. The four velocity of the continuous spin theory expressed in terms of phase space variables

$$\frac{dx^\mu}{dx^o} = \frac{X'^\mu}{X'^o}$$

is gauge variant and hence not an observable. This gauge variance can be thought of as a manifestation of the well known non-localizability of massless particles. The famous comment of Newton and Wigner⁹ that massless particles of helicity one and greater do not have a position operator was proven and extended to continuous spin particles by Wightman¹⁰. The continuous spin theory therefore escapes the causality objection which plagues the tachyon theory.

We shall now attempt to discover what gauge group is responsible for the first class constraints of the acceleration theory of continuous spin. Ordinarily the Lagrangian is discovered by demanding it be invariant under a known gauge transformation. The constraints that appear in the Hamiltonian form of the theory can be understood in a general

way by applying Noether's theorem to the infinite dimensional gauge invariance. The acceleration theory, however, presents us with the converse situation where the Lagrangian was guessed by the desire to produce certain Hamiltonian constraints. The problem now is to pass from the first class constraints in the Hamiltonian formulation to uncover the gauge transformations in the Lagrangian formulation.

Let us first study the problem in the context of the simple arc length theory

$$L = m\sqrt{-\dot{x}^2}.$$

The constraint

$$P^2 + m^2 = 0$$

generates the phase space motion

$$\delta X = [P^2 + m^2, X] = 2P$$

$$\delta P = [P^2 + m^2, P] = 0.$$

Although the phase space motion involves no arbitrary function, the arbitrary function characteristic of a gauge transformation enters through the arbitrary function in the Hamiltonian. To eliminate the phase space variable P in favor of the Lagrangian variable $\dot{X}$ we use the equation of motion

$$\dot{X} = [H, X] = 2\lambda P$$

$$H = \lambda(P^2 + m^2)$$

to get

$$\delta X = \frac{\dot{X}}{2\lambda} = u\dot{X}$$

where $u = \frac{1}{2\lambda}$. These are the infinitesimal gauge transformations of the reparameterization group which have the Lie algebra

$$\delta_u \delta_v X - \delta_v \delta_u X = \delta_{[u,v]} X$$

$$[u, v] = u'v - v'u.$$

The procedure can be repeated for the acceleration theory. The primary constraint of the acceleration theory

$$P'^2 - 1 = 0$$

generates the phase space motion

$$\delta^1 X = 0$$

$$\delta^1 X' = 2P'$$

$$\delta^1 P = 0$$

$$\delta^1 P' = 0.$$

To eliminate X' and P' in favor of the derivatives of X we use the equations of motion

$$X' = \dot{X}$$

$$\ddot{X} = 2\lambda P'$$

to get

$$\delta^1 \dot{X} = \frac{\ddot{X}}{\lambda} = u \ddot{X}$$

where $u = \frac{1}{\lambda}$. Integration gives the infinitesimal gauge transformation

$$\delta^1 X = \int^t u \ddot{X}.$$

These infinitesimal gauge transformations also provide a representation of the Lie algebra of the reparametrization group

$$\delta_u^1 \delta_v^1 X - \delta_v^1 \delta_u^1 X = \delta_{[u, v]}^1 X.$$

This representation of the reparametrization group is qualitatively different from the ordinary representation in that the transformations are non-local due to the integral over the

evolution parameter. Thus it is the reparametrization group, albeit in a new guise, which underlies the acceleration theory of continuous spin.

The Lagrangian is indeed invariant up to a surface term under this non-local gauge transformation. The variation of the Lagrangian is

$$\delta^1 L = \frac{\ddot{x} \delta^1 \dot{x}}{\sqrt{\ddot{x}^2}} = \frac{\ddot{x} (u \ddot{x})}{\sqrt{\ddot{x}^2}} = (u \sqrt{\ddot{x}^2}).$$

The gauge transformations associated with the secondary and tertiary constraints can be found in a similar fashion. They are

$$\delta^2 X = \frac{u}{2} \ddot{X} = \frac{1}{2} \delta^1 \dot{X}$$

$$\delta^3 X = -(u \ddot{X}) = -\delta^1 \ddot{X}$$

where their relation to δ^1 is natural as these constraints are generated by consistency (i.e., by taking time derivatives). However, δ^2 and δ^3 do not give invariances of the Lagrangian up to surface terms. It is perhaps possible to introduce auxillary variables to promote them to invariances up to surface terms although we shall not pursue this question further here.

The finite version of the gauge transformation $\delta^1 X$ is simply

$$X \rightarrow \int^t \dot{X} \circ F$$

where F is the finite version of u . The action

$$S = \int \sqrt{\ddot{x}^2} dt$$

is invariant under this gauge transformation in the same way the usual arc length action is. Unlike the arc length theory, however, solutions are not mapped into solutions. The difficulty is that the boundary conditions of the variational principle are not respected by gauge transformations. If x is fixed at end points t_1 and t_2 , then the transformed x is not

because of the non-local nature of the gauge transformations. The hidden reparametrization invariance is thus broken by boundary conditions.

The acceleration theory of continuous spin represents a new kind of model possessing a non-local reparametrization invariance which would have been excluded on the basis of ordinary reparametrization invariance. The model provides further insight into Wigner's continuous spin wave equation allowing a discussion of causality in the context of the classical theory. The hidden gauge symmetry is much like local supersymmetry⁴ in providing a model of spinning particles. Like supersymmetry, it may also have application in the context of other models. There are indications⁶ that the string version of this hidden gauge symmetry may provide a quantum mechanically sensible bosonic string theory. The gauge symmetry also restricts the form of terms that can appear in the Lagrangian. For example, the point particle analog of the cosmological constant term, the arc length, is prevented from appearing in the Lagrangian. It might be interesting to construct higher derivative field theories of gravity possessing a hidden gauge invariance in connection with the cosmological constant problem.

ACKNOWLEDGEMENTS

I would like to thank M. Awada, J. Klauder, C. Thorn and R. Woodard for suggestions and stimulating conversations.

REFERENCES

1. E. Wigner, *Ann. Math.*, **40** , 149 (1939); V. Bargmann and E. Wigner, *Proc. N.A.S.*, **34** , 211 (1948).
2. E. Wigner, *Z. Physik*, **124**, 665 (1947); also in *Theoretical Physics* (International Atomic Energy Agency, Vienna, 1963), p. 59.
3. M. Plyushchay, *Mod. Phys. Lett. A* **4** , 837 (1989).
4. D. Zoller, *Phys. Rev. Lett.* **65**, 2236 (1990).
5. A. Polyakov, *Nucl. Phys. B* **268**, 406 (1986).
6. D. Zoller, preprint University of Florida, Gainesville UFIFT in preparation.
7. D. Zoller, preprint University of Florida, Gainesville UFIFT 89-24.
8. E. Whittaker, *A Treatise on the Analytic Dynamics of Particles and Rigid Bodies* (Cambridge University Press, 1988) p. 265.
9. T. Newton and E. Wigner, *Rev. Mod. Phys.* **21** , 400 (1949).
10. A. Wightman, *Rev. Mod. Phys.* **34** , 845 (1962).

END

DATE
FILMED

4/9/92

3