Title: A POTPOURRI OF RESULTS IN QCD FROM LARGE LATTICE
SIMULATIONS ON THE CM5

Author(s): TANMOY BHATTACHARYA AND RAJAN GUPTA, T-8, MS B285
LOS ALAMOS NATIONAL LABORATORY
LOS ALAMOS, NM 87545

Submitted to: TO BE SUBMITTED TO PROCEEDINGS OF LATTICE '93
HELD IN DALLAS, TX, OCTOBER 11-15, 1993.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Los Alamos
NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Form No. 836 R5
ST 2629 10/91

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED 270

A pot-pourri of results in QCD from large lattice simulations on the CM5

Tanmoy Bhattacharya and Rajan Gupta ^a

^aT-8 Group, MS B285, Los Alamos National Laboratory, Los Alamos, New Mexico 87545 U. S. A.

We present a status report on simulations being done on $32^3 \times 64$ lattices at $\beta = 6.0$ using quenched Wilson fermions. Results for the spectrum, decay constants, the kaon B-parameter B_K , and semi-leptonic form factors are given based on the current statistical sample of 28 configurations. These "Grand Challenge" calculations are being done on the CM5 at Los Alamos in collaboration with G. Kilcup, S. Sharpe and P. Tamayo. We end with a brief statement of code performance.

1. Lattice parameters.

All the results described in these two talks have been obtained using the following lattice parameters. The 32^4 gauge lattices were generated at $\beta = 6.0$ using the combination 10 over-relaxed (OR) sweeps followed by 1 Metropolis sweep. We have stored lattices every 2500 OR sweeps. Quark propagators, using the simple Wilson action, have been calculated after doubling the lattices in the time direction, *i.e.* ($32^3 \times 32 \rightarrow 32^3 \times 64$). This doubling was done in response to the initial hardware/operational constraints of the CM5 at LANL. Currently we are generating $32^3 \times 64$. Periodic boundary conditions are used in all 4 directions, both during lattice update and propagator calculation. Quark propagators have been calculated using two kinds of extended sources - Wuppertal and Wall - at $\kappa = 0.135$ (C), 0.153 (S), 0.155 (U_1), 0.1558 (U_2), and 0.1563 (U_3). These quark masses correspond to pseudoscalar mesons of mass 2800, 980, 700, 550 and 440 MeV respectively where we have used $1/a = 2.3 GeV$ for the lattice scale. The three light quarks allow us to extrapolate the data to the physical u and d quarks, while the C and S κ values are selected to be close to the physical charm and strange quark masses. The C (S) quark mass is slightly lower (higher) than the physical value. So far our statistical sample consists of 28 configurations, each separated by 5000 OR sweeps. The goal is to analyze 100 configurations so the present analysis should be considered preliminary. Also, Sharpe [4] and Bernard and Golterman [3] have pointed

out that in the quenched approximation η' loops give rise to unphysical terms in the chiral expansion for observables. These effects have not been included in the analysis when extrapolating quantities to the chiral limit.

We find that the data for almost all observables show a significant curvature as a function of the quark mass. So we make both a linear and a quadratic fit in each case and quote final numbers from the fit with lower χ^2 . More details about the fits are given in the individual sections.

2. Spectrum.

We have analyzed three types of hadron correlators distinguished by the type of source/sink used to generate quark propagators. These are (i) wall source and point sink (WL), (ii) Wuppertal source and point sink (SL), and Wuppertal source and sink (SS). With these three cases we are able to make consistency checks since the effective mass $m_{eff}(t)$ converges to the asymptotic value from below for WL correlators and from above for SL and SS correlators in all hadron channels. Fig. 1 illustrates, using the pion $U_1 U_1$ buffer, noteworthy features of the convergence of $m_{eff}(t)$. In the case of WL correlators, the approach of $m_{eff}(t)$ to its asymptotic value is monotonic but quite slow and one needs to fit for $t > 15$. The statistical errors in $m_{eff}(t)$ are smallest of the three cases. The SS correlators reach the asymptotic value at $t \approx 8$ and the time-slice to time-slice fluctuations are of the order of the statistical errors. The SL correlators for the

Figure 1. Comparison of the convergence of $m_{eff}(t)$ for three types of pion ($U_1 U_1$) correlators. SL data has been shifted by +0.05.

pion show correlated fluctuations in $m_{eff}(t)$, wiggles with a period of about 12 time slices, which may be a feature of the Wuppertal source. Another possibility, which we will check on the new set of $32^3 \times 64$ lattices, is that it is due to doubling the lattice in the time direction. In spite of these differences in $m_{eff}(t)$, the bottom line is that all three kinds of correlators give consistent (within 1σ) results for masses for all mesons and the error estimates are similar too. We give the final results for masses in lattice units in Table 1.

To extrapolate the data to the chiral limit in the light quark we fit as a function of m_π^2/m_ρ^2 and m_q , where the latter is defined either as $\log(1 + 0.5(1/\kappa - 1/\kappa_c))$ or $0.5(1/\kappa - 1/\kappa_c)$, or determined non-perturbatively as discussed in Ref. [1]. Using a quadratic extrapolation of m_π^2 data versus m_q (as it fits all the points) we get

$$\kappa_c = 0.15721(7). \quad (1)$$

This value is higher than that given in Ref. [1], but consistent with QCDPAX result 0.15717(3). Similarly, using a quadratic fit, the extrapolated value of m_ρ is (see Fig. 2)

$$m_\rho(m_q = 0) = 0.339(3) \quad (2)$$

Figure 2. Quadratic extrapolation of Δ , nucleon and ρ data to the chiral limit.

which gives a lattice scale of

$$a^{-1}(m_\rho) = 2.27(2) \text{ GeV}. \quad (3)$$

We shall, for simplicity, use $1/a = 2.3 \text{ GeV}$.

Wuppertal source quark propagators allow one to construct hadron correlators with non-zero momenta. In Fig. 3 we show the lowest four modes for the pion. The signal in $\vec{p} = (1, 0, 0)$ and $(1, 1, 0)$ correlators is quite good. Thus, as we show later, using 32^3 lattices at $\beta = 6.0$ leads to a significantly improved signal in the calculation of phenomenologically interesting quantities like semi-leptonic form factors and B_K .

2.1. Baryons.

In our previous work[1] we had found that wall and Wuppertal buffers give results that are significantly different. In Fig. 4 we show a typical example of the current status. We find that on the larger lattices WL and SL correlators give consistent results. The plateau in $m_{eff}(t)$ is not very clean and any systematic difference between the two cases is an artifact of the fit range and/or the statistical errors. The numbers given in Table 1 for the case of degenerate quarks are the mean of SL and WL results. Extrapolating the

Figure 3. Effective mass plot for pion (U_1U_1) at different momenta.

data to the chiral limit (Fig. 2) gives

$$\begin{aligned} m_N a &= 0.477(84) \\ m_\Delta a &= 0.61(15), \end{aligned} \quad (4)$$

however, before comparing these numbers to experimental data the reader should note that we have not included the results of Labrenz and Sharpe[6] who predict additional non-analytic terms in the chiral behavior of quenched QCD.

3. Decay constants.

Meson decay constants can be calculated using SL and SS correlators. We use the generic notation f_{PS} for pseudoscalar mesons and f_V for vector meson. There are many different ways of combining these correlators to extract f_{PS} and f_V , some of which are described in Ref. [1]. We find that all methods give essentially identical results, so we quote the mean value in Table 2. For the error estimate we give the largest individual error. We also find that the value of f_{PS} is independent of the pion's momentum. These data are obtained using local currents with two different renormalization factors. Columns 2 and 4 are using the Lepage-Mackenzie mean-field improved scheme[5] while columns 1 and 3 use the naive

Table 1

Masses in lattice units

	m_π	m_ρ	Nucleon	Δ
CC	1.220(2)	1.233(02)		
CS	0.858(2)	0.884(03)		
CU_1	0.819(3)	0.848(04)		
CU_2	0.804(4)	0.835(05)		
CU_3	0.796(5)	0.828(06)		
SS	0.427(2)	0.510(03)	0.790(15)	0.851(19)
SU_1	0.368(2)	0.468(04)		
SU_2	0.343(2)	0.451(05)		
SU_3	0.327(2)	0.441(06)		
U_1U_1	0.301(2)	0.424(05)	0.639(14)	0.715(22)
U_2U_2	0.239(2)	0.390(09)	0.570(13)	0.659(30)
U_3U_3	0.191(2)	0.366(15)	0.525(16)	0.623(33)

Table 2

f_π and f_ρ

	f_π^{naive}	$f_\pi^{\text{imp.}}$	f_ρ^{naive}	$f_\rho^{\text{imp.}}$
CC	0.126(3)	0.205(6)	0.097(2)	0.20(1)
CS	0.103(3)	0.137(4)	0.121(3)	0.20(1)
CU_1	0.094(3)	0.122(4)	0.116(3)	0.19(1)
CU_2	0.090(3)	0.116(4)	0.115(4)	0.19(1)
CU_3	0.088(4)	0.113(5)	0.114(4)	0.18(1)
SS	0.090(3)	0.098(3)	0.228(6)	0.31(1)
SU_1	0.083(3)	0.089(3)	0.241(7)	0.32(1)
SU_2	0.080(3)	0.084(3)	0.246(7)	0.33(1)
SU_3	0.078(3)	0.081(3)	0.248(8)	0.33(1)
U_1U_1	0.077(3)	0.080(3)	0.261(7)	0.34(1)
U_2U_2	0.071(3)	0.072(3)	0.276(8)	0.35(1)
U_3U_3	0.066(4)	0.067(4)	0.285(9)	0.36(1)

$\sqrt{2\kappa}$ renormalization with $Z_\psi = 0.57$ and the 1-loop perturbative value for Z_A . In both cases we use a boosted value for the coupling, $g^2 = 1.75$. The results are displayed in Fig. 5.

As shown in Fig. 5 the data for the case when only one of the quark's mass is varied are qualitatively different from when both quark masses are taken to be the same. Therefore, to get f_π we extrapolate to the chiral limit using the degenerate combinations SS , U_1U_1 , U_2U_2 and U_3U_3 . Using the mean-field improved normalizations we

Figure 4. Effective mass plot for nucleon with WL and SL correlators.

get

$$\begin{aligned} f_{\pi}a &= 0.044(5) && \text{linear} \\ f_{\pi}a &= 0.063(18) && \text{quadratic.} \end{aligned} \quad (5)$$

The data are much better fit by the quadratic form, though the parameters have larger errors. The fact that the type of fit makes such a big difference underscores the fact that one needs more precise data before the phenomenologically interesting parameters f_K and f_D can be extracted. Data for f_V^{-1} show considerable dependence on the renormalization constant (naive versus improved). This issue needs more study.

4. B_K .

The major problem in the calculation of the kaon B -parameter with Wilson fermions is the bad chiral behavior induced by the mixing of the $\Delta S = 2$ 4-fermion operator with wrong chirality operators. This mixing arises as a result of the r -term in the action and has been calculated to 1-loop in perturbation theory. In Ref. [2] it was shown that the lattice artifacts completely overwhelm the signal when using the 1-loop improved operator, however, by calculating the matrix el-

Figure 5. Meson decay constants f_{PS} and f_V^{-1} as a function of m_{π}^2/m_{ρ}^2 . Data for f_V^{-1} has been shifted by +0.25 for clarity.

ement at different values of momentum transfer one can remove most of these. The main limitation of that calculation was that the spatial lattice size was small (16^3), consequently the momentum gap ($2\pi/16$) was large, and the signal in the non-zero momentum correlators was marginal. Both these limitations have been addressed on the present set of lattices and as a result the statistical quality of the data is markedly improved.

In Fig. 6 we show the behavior of the lattice B_K , evaluated at momentum transfer $p = 2\pi/32$, as a function of the time slice at which the operator is inserted. The data shown are for two different kaon sources, *i.e.* $\bar{\psi}_s \gamma_5 \psi_d$ and $\bar{\psi}_s \gamma_4 \gamma_5 \psi_d$ as they converge from opposite direction. The statistical errors are small and the two fits agree within errors. The results for $p = 2\pi/32 \times (1, 1, 0)$ and $p = 2\pi/32 \times (1, 1, 1)$ are also reliable. Full analysis will be presented elsewhere Ref. [7].

In Table 3 we present preliminary results for the B -parameters using perturbatively improved operators evaluated with a boosted $g^2 = 1.75$ and $\mu a = 1.0$. In column 1 (2) we give the raw number for zero (one) unit of momentum trans-

Figure 6. B_K , evaluated at momentum transfer $\vec{p} = (1,0,0)$, as a function of t at which the operator is inserted. The two data sets correspond to the two kinds of operators used to produce the kaons.

fer. The final values of B_K after removing the leading bad chiral contributions by the method of momentum subtraction (Ref. [2]) are given in column 3. In columns 4 and 5 we give the results for the two left-right operators O_7 and O_8 (defined in Ref. [2]).

The data in columns 1 and 2 show that the statistical errors have been reduced to the level of a few percent. The reason that the errors in the final B -parameters are large for heavy quarks is mainly because $E(p) \approx m$, while for light quarks the statistical fluctuations are large.

The numbers extrapolated to the kaon mass are given in the last two rows in Table 3. We use two methods for this: in the first we keep S mass fixed and extrapolate in the light quark mass using S , U_1 , U_2 , U_3 . In the second, the two quarks are taken to be degenerate. Both methods give similar results for all three operators. This suggests that the $SU(3)$ breaking does not give a large contribution to these B -parameters. Also, there is little evidence of unphysical contributions from η' loops which are expected to be significant near the chiral limit in the quenched approximation[8].

Table 3

B_K , B_7 and B_8

	B_K			B_7	B_8
	($p=0$)	($p=1$)	subtr.		
CC	0.95(3)	0.95(4)	1.32(37)	0.97(3)	0.97(3)
CS	0.83(2)	0.84(3)	1.09(21)	0.94(3)	0.95(3)
CU_1	0.79(2)	0.80(2)	0.94(19)	0.92(3)	0.94(3)
CU_2	0.77(2)	0.77(2)	0.76(20)	0.91(3)	0.93(3)
CU_3	0.76(2)	0.75(3)	0.50(24)	0.89(3)	0.92(3)
SS	0.50(1)	0.53(1)	0.82(7)	0.87(2)	0.91(2)
SU_1	0.36(1)	0.41(2)	0.72(6)	0.84(1)	0.89(1)
SU_2	0.29(1)	0.34(2)	0.66(7)	0.83(1)	0.88(1)
SU_3	0.23(1)	0.28(2)	0.60(8)	0.82(1)	0.87(1)
U_1U_1	0.13(1)	0.21(2)	0.62(6)	0.81(1)	0.86(1)
U_2U_2	-.26(2)	-.08(3)	0.48(8)	0.76(1)	0.82(1)
U_3U_3	-.78(3)	-.42(4)	0.37(11)	0.72(1)	0.78(1)
Ext 1				0.48	0.80
Ext 2				0.43	0.80

The values of B_K , B_7 and B_8 are significantly lower than those found in Ref. [2]. We believe that this is due to using a larger lattice and a smaller $\vec{p}$. Very accurate and reliable results for B_K , obtained using staggered fermions, have been presented at this conference by S. Sharpe[8]. Wilson results are consistent with these but have larger systematic errors as a result of mixing with operators having the wrong chiral behavior. By improving the quality of Wilson fermion results one can, by comparison with staggered results, check our understanding of lattice artifacts.

5. Semileptonic Meson Form Factors.

We have calculated the semileptonic pseudoscalar form factors of the D meson, both for $D \rightarrow K e \nu$ and $D \rightarrow \pi e \nu$. In calculating these form factors we have held the D -meson at zero spatial momentum, and varied the momentum of the other meson from 0 to $\pi/8$. This provides a large enough range (both positive and negative) in the invariant mass of the leptonic subsystem ($-Q^2$) to test the pole dominance hypotheses. To control systematic effects, we have done the calculation using three different transcriptions (called 'local', 'extended' and 'conserved') of the current.

Table 4

The meson form factors extrapolated to $Q^2 = 0$

	$f_+^K(0)$	$f_+^\pi(0)$	$f_0^K(0)$	$f_0^\pi(0)$
U_1	0.686(44)	0.638(46)	0.645(32)	0.535(30)
U_2	0.690(51)	0.602(61)	0.639(35)	0.468(35)
U_3	0.691(61)	0.569(89)	0.633(42)	0.415(48)

The details of the method are given in Ref. [9] and a complete analysis for the present set of lattices will be presented in Ref. [10].

The renormalization of all three currents is carried out using the Lepage-Mackenzie[5] mean field improved perturbation theory. We find that the values obtained using the three currents are very close. (See Fig. 7 for an example of consistency between different currents.) We notice that the statistical errors are very small when the hadrons carry zero or one unit of momentum, and some systematic difference between the local and the non-local currents become visible. The errors are much larger for $\vec{p} = (1, 1, 1)$ probably because we do not average over all possible equivalent cases.

Except when $\vec{p} = (1, 1, 1)$, the data are consistent (see Fig. 7 for an example) with the hypothesis that the form factor is dominated by the nearest pole with the right quantum numbers (D_s^* and D^* respectively for the K and π decay). Assuming pole dominance we can extract $f(Q^2 = 0)$, and the results are summarized in Table 4. To within statistical errors the numbers are independent of the light quark mass for f^K and there is a slight decrease in f^π . For this reason we have not at this stage extrapolated the data to the chiral limit, or to the physical values of charm and strange quark masses. Our final numbers are very close to the phenomenological values for f_+^π and f_+^K .

6. Code Performance on the CM5

Our CM5 code uses the SIMD programming environment. All computationally intensive portions of the code have been written in CDPEAC and modularized. Calling these library of low level codes is not a significant overhead due to the performance of the Sparc chip, while having the algorithm control statements in CM For-

Figure 7. Comparison of $1/f_0$ obtained using the three vector currents in the case of the lightest U quark. The fits show the agreement with pole dominance. m_{D_s} has been held fixed at its lattice value when doing the fit.

tran provide flexibility, and adding/changing algorithms and physics has been easy. Our present gauge update code runs at 25 megaflops/node and the Wilson propagator generation sustains 35 megaflops/node. These timings include all I/O and setup overhead. For the present set of physics objectives we need 1800 node-hours to generate and process one configuration.

REFERENCES

1. R. Gupta *et al.*, *Phys. Rev.* **D46** (1992) 3130.
2. R. Gupta *et al.*, *Phys. Rev.* **D47** (1993) 5113.
3. C. Bernard and M. Golterman, *Phys. Rev.* **D46** (1992) 853.
4. S. Sharpe, *Phys. Rev.* **D41** (1990) 1990.
5. P. Lepage and P. Mackenzie, *Phys. Rev.* **D48** (1993) 2250.
6. J. Labrenz, S. Sharpe, these proceedings.
7. T. Bhattacharya *et al.*, in preparation.
8. S. Sharpe, these proceedings.
9. R. Gupta *et al.*, hep-lat/9310007.
10. T. Bhattacharya *et al.*, in preparation.

DATE

FILMED

3 / 7 / 94

END