Methods for Testing Transport Models: Third Year Annual Report

Clifford Singer and Dennis Cox
Departments of Nuclear Engineering and Statistics
University of Illinois at Urbana-Champaign

January 5, 1993

Abstract

This report documents progress to date under our three year contract for developing "Methods for Testing Transport Models." The work described includes (1) choice of best methods for producing "code emulators" for analysis of very large global energy confinement databases, (2) recent applications of stratified regressions for treating individual measurement errors as well as calibration/modeling errors randomly distributed across various tokamaks, (3) Bayesian methods for utilizing prior information due to previous empirical and/or theoretical analyses, (4) extension of code emulator methodology to profile data, (5) application of nonlinear least squares estimators to simulation of profile data, (6) development of more sophisticated statistical methods for handling profile data, (7) acquisition of a much larger experimental database, and (8) extensive exploratory simulation work on a large variety of discharges using recently improved models for transport theories and boundary conditions. From all of this work, it has been possible to define a complete methodology for testing new sets of reference transport models against much larger multi-institutional databases.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

1 Introduction

Under grant DE-FG02-88ER53269, we have been funded primarily for the purpose of developing methods for testing theoretical plasma transport models against data from experiments on tokamaks. We have accomplished this task by developing a fairly complete set of statistical, plasma simulation, and database handling methods. This primary purpose of this report is to describe these developments.

There are growing bodies of excellent work on statistical methods for empirical description of tokamak data, transport theory, and transport simulation and transport analysis. What distinguishes our work is an intimate connection between statistics, theory, and time-dependent transport modeling. We began this work with the hypothesis that such a coordinated approach would be a useful complement to detailed studies by various other researchers on various aspects of this problem. We now have results which amply illustrate the utility of developing statistical methods in parallel with time-dependent transport modeling using appropriate transport theories. A second purpose of this report is to describe these results.

We have also made substantial progress in gaining access to a useful database for testing theory-based transport tokamak transport models. The methods we have developed are specifically tailored to use of large, published or publicly posted databases collected from many sources. We have developed and begun automating the construction of the “code emulators” described below, so our methods will be adequate to cover the largest possible range of tokamak operating parameters with the greatest possible accuracy. We have developed methods appropriate to the use of the sometimes incomplete data in publicly posted databases so that our results can be reproduced and extended by researchers at any institution, without access to privileged data held at other institutions. A third purpose of this report is to describe what we have learned from working with different types of experimental data from a wide variety of sources.

While our works on statistical methods and transport simulation are actually intimately interrelated, it is convenient here to divide them into two separate discussions. Thus, the first part of this report deals primarily with statistical methods. The second part deals primarily with transport simulation. Illustrations of the utility of developing statistical methods in parallel

with time-dependent transport modeling using appropriate transport theories are included throughout these sections.

2 Code Emulation and Statistical Analysis

There is a roughly comparable amount of information available to us in global confinement and plasma profile databases. The global confinement databases obviously provide less information per discharge, but they include over an order of magnitude more discharges over a very wide operating parameter range. Ability to reproduce global energy confinement results is also readily communicated in the form of simple residuals plots. Plasma profile data, on the other hand contains a qualitatively different type of information not available in global confinement databases. For these reasons, methods for using both types of data are described in each of the following subsections.

As described below, we have found it essential to do full time-dependent simulations to analyze some of the discharges in the data available to us. However, it is computationally impractical to repeat this for several a priori uncertain transport model parameters for each of the hundreds or thousands of discharges in the available databases. To retain a consistent methodology and make full use of the data which is readily available, it is thus necessary to develop more efficient methods for applying transport simulation methods to this problem. Our solution has been to develop “code emulators” which adequately approximate the expected value of the transport code outputs of interest, as a function of various input parameters. Therefore, methods for developing code emulators are first described for each type of data. Methods for actually testing transport models against experimental data are then described.

It is also commonly the case that we wish to utilize prior information from previous empirical and/or theoretical analyses. Bayesian methods are appropriate for this, and examples of their application to use of global confinement and profile data are also described.

2.1 Global Confinement Data

2.1.1 Code Emulators

Our work has been very substantially facilitated by the development of “code emulators” for global energy confinement results. To produce a code emulator, it is assumed that $y_{iC} = \ln \tau_{EiC}$ deviate randomly from a function $Y(\mathbf{s}_i)$ of a set $\mathbf{s}_i$ of code input parameters:

$$y_{iC} = Y(\mathbf{s}_i) + \epsilon_{iC} \quad (1)$$

Here τ_{EiC} is the computer codes’ prediction of the global energy confinement time for the i th tokamak discharge. For our applications to date, $Y(\mathbf{s}_i)$ has been a linear function of a priori uncertain theory model parameters and of the logarithms of parameters varied for each tokamak in the database (e.g. heating power toroidal current, line-average electron density, and toroidal magnetic field). ϵ_{iC} allows for deviations of the computer results from the best (in our case linear) regression. These deviations may be highly correlated for similar code input parameters but essentially uncorrelated for very dissimilar code input parameters. ϵ_{iC} also allows for completely uncorrelated random errors in a given computation (e.g. due to use of Monte Carlo computational algorithms.) Appendix A gives more details of this statistical model. Cox, Sacks, Park and Djemil [1, 2, 3] and have estimated regression coefficients and variance parameters by maximizing the likelihood that the computer results were sampled from such a distribution. Analysis of the results indicated that it is practical to construct code emulators accurate enough to make it unnecessary to simulate all of the discharges and in a large database individually.

Cox et al. went on to investigate a number of methods for estimating a priori uncertain simulation parameters [1, 2]. They did this in two contexts. The first context involved “toy model” experiments, where parameters in a given function were estimated from a limited set of evaluations of that function. This study showed that the optimal method for parameter estimation can depend sensitively on the problem at hand (c.f. Table 1 on p. 11 of attached Appendix A). This suggested that development of statistical methods for analyzing transport theories is best done using the simulations based on these theories themselves, rather than on simpler model systems. Park’s

thesis with Cox also examined various methods for applying code emulator methodology to the problem of estimating a priori uncertain parameters used in modelling tokamak confinement data with the BALDUR transport simulation code [2].

2.1.2 Stratified Nonlinear Regression

One of the postulates we have been investigating concerns multiple sources of random deviation between reported experimental data and results of models. We noted that refined calibration and modeling of experimental measurements reaches a point of diminishing returns when resulting uncertainties are comparable to those due to random individual measurement errors (and vice versa). Thus, both individual random measurement errors and “calibration/modeling errors” common to blocks of measurements might need to be included in an adequate statistical analysis. Djemil, Singer, and Cox have recently produced some improvements of their examples of stratified regressions which allow for such multiple sources of variation ([4], c.f. Appendix B). The results of the log-linear regressions in this work are consistent with the supposition that deviations from a best empirical fit which are to all of the data from a given tokamak are at least as large a source of variance as individual random measurement errors. The nonlinear Lowess fits in this work indicate that log-linear empirical regressions leave systematic residuals. (An example for dependence of global energy confinement time on density is shown in Fig. 4 on the last page of attached Appendix B.) Since we had already found similar results with a theory-based transport simulation in good qualitative agreement with experimental data [5, 6, 7], this strongly suggests that our program of stratified calibration of non-linear theory-based transport simulation models may ultimately prove more fruitful than use of more statistically complex empirically based descriptions.

A complete description of our best presently identified method for dealing with stratified sources of variance when calibrating theory-based simulations of large global energy confinement data bases is given in Appendix C.

2.1.3 Bayesian Methods

We have found some applications of Bayesian statistical methods which appear to fit naturally with both numerical analysis and scientific communications problems encountered in this field. Appendix C gives one example. In Appendix C, we discuss a Levenburg-Marquardt type numerical algorithm for obtaining maximum likelihood estimates of statistical and transport theory model parameters. This algorithm starts with highly informative prior distributions for transport theory parameters and measurement error variance estimates. It uses the resulting estimates to as a starting point for maximum likelihood estimates of these parameters on successively less informative prior distributions, until the limit of completely uninformative priors.

The approach just outlined does much more than define a robust numerical algorithm for maximization over potentially complicated multi-dimensional parameter spaces. It will also produce graphs of theory parameter estimates as a function of assumed prior accuracy of the theory. This alleviates a scientific communication problem by allowing the theorists themselves the ultimate scientific judgement about the meaning of the results, rather than forcing the data analyst to make a less informed decision about the accuracy resulting from complex theoretical derivations of transport theories. Similarly, the experimentalist can be provided results which can be interpreted in terms of their entire experience with measurement accuracy, in addition to the “uninformative prior limit” based only on the particular data set used for the statistical analysis reported.

We have already reported less general examples of the kind of analysis just described. For example, Park’s thesis gives maximum likelihood estimates and confidence regions for theory model parameters using a particular set of semi-informative prior distributions based on conversations by Singer with the authors of different transport theories ([2]; c.f. Fig. 5 on p. 19 of attached Appendix D). Djemil et al. also computed estimates for a variance ratio for particular semi-informative and uninformative priors ([4], Appendix B).

2.2 Profile Data

2.2.1 Code Emulators for Profiles

We have examined two approaches to construction of code emulators for temperature and density profiles. The first approach uses a tensor product of quadratic functions of transport model parameters with splines for the radial variable. Results are described in Fig. 1 on p. 9 of attached Appendix D. Our second approach extends the random function approach already used for global energy confinement emulators. Our results suggest that the spline method will be sufficient for simulations based on the twenty-two reference discharges and modest extensions thereof described below. The random function approach is being further investigated in case it is needed for simulating the majority of the entire “world profile database” maintained at the University of Texas at Austin.

2.2.2 Nonlinear Least Squares

One important finding from our work on nonlinear least squares estimates of transport model parameters is shown in Fig. 2 on p. 10 of attached Appendix D. The results shown provide by far the clearest demonstration we have seen of why sawtooth modelling and treatment of sawtooth phasing can be critical for estimation of theory-based transport model parameters. The particular result shown in this figure is that nonlinear least squares estimate of a multiplier of nominal theory-based turbulence saturation levels for a JET case was straightforward with correct sawtooth phasing information but impossible without taking some account of sawtooth phasing.

We have also automated the process of finding simple nonlinear least squares fits estimates for a priori uncertain parameters in theory-based simulations of tokamak profile data. Fig. 4 and Section 3.1.5 on pp. 12-17 of attached Appendix D give an example of the results of a nonlinear least squares result for two theory model parameters (correspondingly physically to turbulence saturation levels for drift/ η_i and resistive ballooning modes.) These results were produced using a Unix c-shell script and associated coding which automatically modifies transport code input files, executes transport code runs, edits and interpolates output files, computes statistics, and finds least squares minimization allowing for random variations in transport code

output due to use of Monte Carlo numerical algorithms. This script and associated coding has been imbedded in continuously updated L^AT_EX documentation and distributed to interested co-workers [8].

2.3 Improved Statistical Methods for Profile Data

We have also defined a statistical model which allows for radially correlated covariance in the deviation between transport simulations and experimental data. Some details of this work are given in Section B of Appendix D.

3 Experimental Data Bases

We have also recently obtained and explored a much larger experimental data base. This includes the following: First, we have visited and interviewed experimentalists to obtain background information and time-dependent data needed for the careful transport simulation of the 22 reference discharges from seven tokamaks as listed in Table 1 on p. 7 of attached Appendix D, as well as a number of other cases. We have also worked intensively with a large L-mode global confinement data base and held extensive discussions with authors of the ITER H-mode data base [9] concerning any potential problems which might arise in modeling its contents with our time- and divertor-configuration-dependent boundary conditions model [10]. In addition, we have identified a number of discharges in the “world profile database” maintained at the University of Texas for use in our studies of statistical and computational methods for using large profile data bases to systematically calibrate theory-based predictive transport simulation models.

4 Transport Simulations and Improved Theory-based Models

Finally, we have done extensive exploratory work on simulating a variety of tokamak profile data using our recent survey [11] of improved transport theories. Results of this work are given in Appendix E. Although this kind of work could be used to support a random coefficients model (c.f. [12]),

it should be emphasized that this is *not* our intent. Rather, the process of fitting a variety of different discharges separately with newer transport theories has been used to guide development of a new class of theory-based simulation models which we expect may be able provide a unified framework for reproducing experimental results from a much wider variety of tokamak discharges. An example of such a model is given in Section A.2 on pp.28-30 of attached Appendix D.

5 Summary

It should be clear from this report that we have made substantial progress on a variety of the problems which need to be solved in order to proceed with calibration and testing of theoretical transport models against large multi-institutional experimental data bases. This should allow us and interested co-workers from other institutions to proceed with this substantial but potentially very productive effort, along the general outlines further described in the attached Appendix D.

References

- [1] Cox, D., Park, J.S., Sacks, J., and Singer, C.E. (1992), "Tuning Complex Computer Code to Data," *Proceedings on the 23rd Symposium of the Interface of Computer Science and Statistics*, Interface Foundation, Fairfax Station, VA, 266 – 271.
- [2] J. S. Park, "Tuning Complex Computer Codes to Data and Optimal Designs," University of Illinois at Urbana-Champaign Department of Statistics Ph.D. Thesis (1991).
- [3] T. Djemil, "Methods for Testing Transport Models," University of Illinois at Urbana-Champaign Department of Nuclear Engineering Ph.D. Thesis (1992).
- [4] T. Djemil, C. E. Singer, and D. Cox, "A Bayesian Analysis of Multisite Tokamak Confinement Data," submitted to *Technometrics*.
- [5] E. Ghanem, *Theoretical Transport Model for Tokamaks*, University of Illinois at Urbana-Champaign Department of Nuclear Engineering Ph.D. Thesis (1991).
- [6] E. Ghanem, C. E. Singer, G. Bateman, and D. Stotler, *Nucl. Fusion* **30** (1990) 1595.
- [7] J. Kinsey, G. Bateman, T. Djemil, E. Ghanem, and C. Singer, "Theory-Based Transport Simulation of Tokamaks" *Proc. 19th European Conf. on Controlled Fusion and Plasma Physics* (Innsbruck, 29 June–3 July, 1992).
- [8] C. E. Singer, "Automated Variance Calculations for Optimizing BALDUR Transport Code Fits to Experimental Data: Version 0.1," University of Illinois personal communication to G. Bateman, G. Murphy, and G. L. Singer (1992).
- [9] J. P. Christiansen, J. G. Cordey, K. Thomsen, A. Tanga, J. Deboo, D. Schissel, T. Taylor, O. Kardaun, F. Wagner, F. Ryter, S. M. Kaye, Y. Miura, N. Suzuki, M. Mori, T. Matsuda, H. Tamai, T. Takizuka, S.-I. Itoh, and K. Itoh, "A Global Energy Confinement H-mode Database

for ITER," ITER Report ITER-IL-PH-4-1-1 (JET Joint Undertaking, Abingon, UK).

- [10] C. E. Singer, G. Bateman, and D. D. Stotler, "Boundary Conditions for OH, L, and H-mode Simulations," Princeton University Plasma Physics Report PPPL-2527 (1988).
- [11] C. E. Singer and G. Bateman, "Predictive Tokamak Transport Simulations," University of Illinois Fusion Studies Laboratory Report FLS-333 (1991).
- [12] K. S. Riedel, "Advanced Statistics for Tokamak Colinearity and Tokamak to Tokamak Variation," Courant Institute of Mathematical Science: Magneto-Fluid Dynamics Division Report DOE/ER/53233-98 & MF-118 (1989).

*preprints received;
proof paper
cycled
separately.*

**DATE
FILMED**

2 / 24 / 93