BNL--48166

DE93 006392

A CLIMATIC GUIDE FOR NORTH CENTRAL OKLAHOMA

Robert M. Brown
Department of Applied Science
Brookhaven National Laboratory
Upton, NY 11973

June 1991

UNDER CONTRACT NO. DE-AC02-76CH00016 WITH
THE UNITED STATES DEPARTMENT OF ENERGY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

for

North Central Oklahoma

Table of Content

	Page
Introduction.	1-4
	5
1. General map.	
Specific locations.	6
Latitude, longitude, elevation.	
2. Insolation.	7-12
Monthly average values.	13
Annual daily values.	14
Map of ave. annual values.	15
Duration of sunshine.	16
Heating degree days-based on 65 F.	17
Cooling degree days-based on 65 F.	
3. Temperature.	
Data listing.	18
Average max. mean and min.	19-31
Plots of location temperatures.	32
Average plot of all locations.	33
Map of ave. annual temperatures.	
4. Precipitation.	34
Data listing.	35-51
Plots of location precipitation.	52
Average plot of all locations.	53
Map of ave. annual precipitation.	
5. Cloudiness.	54
Plots of Ok. City and Tulsa.	
6. Thunderstorm days.	55
Data listing.	56
Plot of location thunderstorms.	57
Average plot of all locations.	58
Map of ave. annual thunderstorms.	
7. Snowfall.	59
Data listing.	60
Plot of location snowfalls.	61
Average plots of all locations.	62
Map of ave. annual snowfalls.	
8. Wind.	63
Annual wind direction frequencies.	64
January wind directions.	65
July wind directions.	66
Annual wind speed frequencies.	
9. Fog.	67
Annual fog days.	
10. Drought.	68
Percentage months of drought.	69
11. Moisture content.	
	70
References	

A Climatic Guide for North Central Oklahoma

Introduction

This guide provides some climatological data pertaining to central and north central Oklahoma. The information was derived from standard reference material to reflect what general surface meteorological characteristics exist in that region. It is intended to assist those individuals involved in the implementation of the first ARM site in that locale. A similar guide already exists for the region involved in Kansas entitled, "One Regional ARM Guide for Climatic Evaluation". The Oklahoma-Kansas area was selected as the first site from the process reported in the "Identification, Recommendation and Justification of Potential Locales for ARM Sites".

Oklahoma, like Kansas, has extreme climatic variations, is centrally located, is compatible with other programs, has good airfields and accommodations to plan and operate an ARM site for continuous use and for special field campaigns.

The map on page 5 shows the area in Oklahoma that will be involved in the first ARM measurements. It gives the names of the cities, towns and the counties and their boundaries. Seventeen (17) meteorological reporting locations in Oklahoma were used in this guide to provide the results presented. The map on page 6 shows their locations, latitudes, longitudes and elevations.

1. General Features.

Most of Oklahoma has rolling plains which slope upward from east to west. There are hilly sections associated with the Wichita Mountains in the southwest, the Arbuckle Mountains in the south-central, the Ozarks in the northeast and the Ouachita Mountains in the southeast. The lowest part of the State (300 ft.) occurs in the southeastern corner while the highest (5000 ft.) is in the northwestern Panhandle. The State lies in the drainage basin of the Mississippi River where the Arkansas drains the northern sections and the Red River drains the south. There are a number of man made lakes to control flooding and to provide recreational areas; the largest are Texoma on the Red River and the Eufaula Reservoir on the Canadian River.

The climate is continental with warm air moving northward from the Gulf of Mexico, cold air moving southward from Canada and downslope winds moving eastward from the Rocky Mountains. This combination of air movement throughout Oklahoma and Kansas causes some of the most severe weather in the world. Summers are long and occasionally very hot whereas winters are shorter and cold but less so than more northern Plain States. Long periods of extreme cold are rare. Humidity and cloudiness are generally greater and precipitation heavier in the south and eastern sections as compared to the north and western sections of the State. Droughts do occur during hot summers accompanied by moderately high winds. The drought of the early to mid 1930's is an example of this condition.

Excessively heavy rains can occur. Amounts of 10 inches or more have been recorded in a 24-hour period in several areas of

the State. Thunderstorms are frequent and are often the cause of damaging tornadoes and hail. A "low level jet stream", identified in earlier experimental studies (Bonner, 1968), flows from the southeast corner of New Mexico through parts of Texas, Oklahoma, Kansas and into Missouri and Iowa. It has a speed of from 11 to 14 meters/sec. at an altitude of from 700 to 900 meters above the surface. The location of this jet stream coincides with some of the most severe weather recorded in those states and is a possible area for further investigation.

2. Insolation.

All of the solar radiation data available in past climatological records show a near linear increase of amounts from the northeast to the southwest in Oklahoma. The maps on pages 7 to 12 show the average monthly isopleths of solar radiation during the year and the map on page 13 shows the average annual isopleths. These maps show the definite increase from northeast to southwest. The map on page 14 shows the average annual solar radiation through the 17 locations studied. As noted, there is some interpolation on many of these maps so some caution is required in their analysis. The values range from over 350 to 450 Langleys/day. Oklahoma and Kansas, as shown on the map on page 15, has between 2800 to 3300 hours of duration of sunshine. Only New Mexico, Arizona and southern California have more sunshine hours.

Annual values of heating and cooling degree days are shown on pages 16 and 17. They are provided in this guide as an indication of the air conditioning and heatings loads to be expected in that region.

3. Temperature.

The table on page 18 gives the numerical listing of the temperatures in north central Oklahoma. The average temperatures of the various locations is given at the bottom of that page. The plots of the temperatures are shown on pages 19 through 31. The average temperatures of the locations is shown on page 32. The map on page 33 shows the average mean temperature isotherms in that region. There is a definite decrease in surface temperature from the southeastern to the northwestern sections, however, the summer average maximum temperature in northern central Oklahoma can be the highest in the State as given in the Oklahoma chapter of "The Climate of the States".

4. Precipitation.

The locations studied in this guide show a bimodal characteristic in annual precipitation amounts. A maximum occurs in May and another in September. This is shown in each of the plots from page 35 to 51. It is also evident in the plot of their averages shown on page 52. The map on page 53 shows the characteristic decrease of precipitation in Oklahoma from east to west of from 38 to 23 inches per year. The pattern and amounts agree well with those given for Kansas.

5. Cloudiness.

Clear skies are more frequent in the western part of the State than in the east, where clear and cloudy conditions are about equal. Sunshine records indicate an annual average of near 70% of possible sunshine in Oklahoma City as compared to 60% for Tulsa. Summer has the greatest period of sunshine and winter the least. The plots on page 54 show the clear, partly cloudy and cloudy monthly values for Oklahoma City and Tulsa.

6. Thunderstorms.

Oklahoma, like Kansas, experiences between 45 and 60 thunderstorm events each year. The plot on page 56 shows the thunderstorm days for six (6) locations in north central Oklahoma. As can be seen, most of them occur between April and October, peaking in May and June. There is a slight increase in August. The average plot of the 6 locations is shown on page 57. The map on page 58 shows the near linear decrease of thunderstorm activity from the northeast to the southwestern sections. It is well known that thunderstorms in the Oklahoma and Kansas region can be severe at times and that tornadoes and hail can do considerable damage. The guide on Kansas shows a plot of the risk of hail activity with elevation which applies to Oklahoma as well. A map showing the incidence and paths of tornadoes is also given in that guide and will not be repeated in this one. In "The Climate of the States" 1600 tornadoes were reported in the 100 year period 1875 to 1975 or about 16 tornadoes per year for the Oklahoma-Kansas region.

7. Snowfall.

The annual amount of snowfall in Oklahoma tends to increase from the southeast toward the northwest sections as shown on page 62. The average annual snowfall has been approximately 2 inches in the southeast to over 20 inches in the western sections of the Panhandle. Strong winds with heavy snowfalls can cause deep drifting which restricts visibility and traffic.

8. Wind direction and speed.

The annual percentage frequencies of wind directions are shown on page 63. The map shows a predominance of south and north winds. The map on page 64 gives the average directions during January which indicate an almost even north or south frequency while the map on page 65 indicates more southerly winds than northerly winds during July.

The average annual wind speeds shown on page 66 indicate an almost symmetrical distribution throughout the year. The north-south wind direction sectors dominate somewhat however. As can be noted, wind speeds average over 5 meters/sec most of the time.

9. Fog.

The map on page 67 shows the mean annual number of days with fog in the Oklahoma-Kansas region. As can be seen from the map, from 10 to 25 days a year can be expected to decrease visibility and have an effect on other atmospheric measurements.

10. Drought.

The map on page 68 provides some information on the severity of extreme drought conditions in the Oklahoma-Kansas region. It can be seen that a maximum occurs in the western sections of Kansas. A drought condition will have a profound effect on the type of vegetation measurements from overhead satellite recordings, a note to be remembered.

11. Moisture content.

The Oklahoma-Kansas region has, on the average, about as much moisture available from the surface to 325 mbar as the eastern central United States. The map on page 69 shows this feature.

Oklahoma

Location	Lat.	Long.	Elev.(ft)
1. Mutual	36:14	99:10	1865
2. Alva	36:48	98:39	1374
3. Cherokee	36:46	98:21	1180
4. Helena	36:32	98:16	1350
5. Jefferson	36:43	97:48	1070
6. Enid	36:25	97:52	1245
7. Hennessey	36:07	97:54	1170
8. Kingfisher	35:43	97:48	1070
9. Marshall	36:09	97:38	1045
0. Newkirk	36:53	97:04	1135
1. Ponca City	36:44	97:06	999
2. Perry	36:17	97:18	1025
3. Guthrie	35:53	97:27	980
4. Oklahoma City	35:24	97:36	1285
5. Ralston	36:30	96:44	825
6. Pawnee	36:20	96:48	840
7. Tulsa	36:11	95:54	668
* Lamont	36:41	97:29	1005

January

February

March

April

May

June

September

October

November

December

Average Daily Global Solar Radiation on a Horizontal Surface (MJ/m^2)

ANNUAL

Annual Mean Daily Insolation for the United States
1950-1964

Mean annual duration of sunshine, in hundreds of hours, based on black-bulb sunshine recorders at 125 stations, 1931-1960, and 50 for fewer years. (After U.S. Weather Bureau National Atlas map.)

Heating Degree Days

ANNUAL

Cooling Degree Days

ANNUAL

North Central Oklahoma Data Temperature (F)													
Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Mutual													
Ave.Max	47	52	61	72	80	90	96	95	86	75	59	50	72
Mean	34	39	47	58	67	77	82.5	81	72.5	61	46.5	37.5	58.5
Ave.Min	21	26	33	44	54	64	69	67	59	47	34	25	45
Alva													
Ave.Max	46	52	61	72	81	91	97	95	86	76	60	50	72
Mean	34	39.5	47.5	59	68.5	78	83.5	82	73	62	47.5	38	59.5
Ave.Min	22	27	34	46	56	65	70	69	60	48	35	26	47
Cherokee													
Ave.Max	47	53	62	74	82	92	97	96	87	77	60	51	73
Mean	34.5	40	48	60	68.5	79	83.5	82.5	73.5	62.5	47	38.5	59.5
Ave.Min	22	27	34	46	55	66	70	69	60	48	34	26	46
Jefferson													
Ave.Max	46	52	61	73	82	92	97	95	87	76	60	50	73
Mean	34	39.5	48	59.5	69	79	84	82	74	62.5	48	38.5	60
Ave.Min	22	27	35	46	56	66	71	69	61	49	36	27	47
Enid													
Ave.Max	48	52	62	72	80	90	95	95	87	75	61	50	72
Mean	36.5	40	49.5	59.5	68	78	82.5	82	74	62.5	49	39	60
Ave.Min	25	28	37	47	56	66	70	69	61	50	37	28	48
Hennessey													
Ave.Max	47	53	62	73	81	91	97	95	87	76	60	50	73
Mean	35.5	41	49	60	69	78.5	84	82	74	63	48.5	39	60.5
Ave.Min	24	29	36	47	57	66	71	69	61	50	37	28	48
Kingfisher													
Ave.Max	48	54	63	74	82	91	97	96	87	77	61	52	73
Mean	36	41.5	49.5	61	69.5	78.5	84	82.5	74	63	49	40	60.5
Ave.Min	24	29	36	48	57	66	71	69	61	49	37	28	48
Newkirk													
Ave.Max	44	50	60	72	80	89	94	93	85	74	58	48	71
Mean	33.5	38.5	47.5	59.5	68.5	77.5	82.5	81	73	62	47	37.5	59
Ave.Min	23	27	35	47	57	66	71	69	61	50	36	27	47
Ponca City													
Ave.Max	43	49	58	70	79	88	94	93	83	73	58	47	70
Mean	32.5	38	46.5	58.5	68	77	82.5	81	72	61	47	36.5	58.5
Ave.Min	22	27	35	47	57	66	71	69	61	49	36	26	47
Perry													
Ave.Max	48	54	63	74	81	90	95	95	87	77	61	52	73
Mean	36.5	41.5	50	61.5	69	78.5	83	82.5	74.5	63.5	49.5	40.5	61
Ave.Min	25	29	37	49	57	67	71	70	62	50	38	29	49
Guthrie													
Ave.Max	48	54	63	74	81	89	95	95	87	76	62	52	73
Mean	36	41.5	50	61.5	69	77.5	83	82	74.5	63	49.5	40	60.5
Ave.Min	24	29	37	49	57	66	71	69	62	50	37	28	48
Ok. City													
Ave.Max	47	52	60	71	78	88	92	93	86	75	60	51	71
Mean	37	41.5	48.5	60	68	77.5	81.5	81.5	74	63.5	48.5	43.5	65.5
Ave.Min	27	31	37	49	58	67	71	70	62	52	37	36	60
Tulsa													
Ave.Max	48	52	60	71	79	88	93	93	86	75	60	51	71
Mean	38	41.5	48.5	60.5	68.5	77.5	82	81.5	74	63	49	41	60.5
Ave.Min	28	31	37	50	58	67	71	70	62	51	38	31	50
Averages													
Ave.Max	46.7	52.2	61.2	72.5	80.5	89.9	95.3	94.5	86.2	75.5	60.0	50.3	72.1
Mean	35.2	40.2	48.4	59.9	68.5	78.0	83.0	81.8	73.6	62.5	48.2	39.2	60.3
Ave.Min	23.8	28.2	35.6	47.3	56.5	66.0	70.6	69.1	61.0	49.5	36.3	28.1	48.5

Temperature—Mutual Ok.

Temperature—Alva Ok.

Temperature—Cherokee Ok.

Temperature—Jefferson Ok.

Temperature—Enid Ok.

Temperature—Hennessey Ok.

Temperature—Kingfisher Ok.

Temperature—Newkirk Ok.

Temperature—Ponca City Ok.

Temperature—Perry Ok.

Temperature—Gutherie Ok.

Temperature—Oklahoma City Ok.

Temperature – Tulsa Ok.

Average Temperature

North Central Oklahoma

ANNUAL MEAN TEMPERATURE

North Central Oklahoma Data
Precipitation (inches)

Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Mutual	0.51	0.93	1.58	2.45	4.32	3.17	2.56	2.20	2.48	1.52	1.15	0.66	23.52
Alva	0.56	0.87	1.62	2.43	4.06	3.80	2.59	2.89	2.47	1.57	1.20	0.81	24.87
Cherokee	0.69	0.92	1.93	2.55	3.85	3.99	2.76	2.58	2.67	1.82	1.28	0.87	25.91
Helena	0.71	1.00	1.92	2.57	4.34	3.95	3.08	2.61	2.87	2.12	1.54	0.94	28.05
Jefferson	0.70	0.97	1.93	2.77	3.92	3.98	3.92	3.25	3.13	2.55	1.92	1.03	30.07
Enid	0.91	1.16	1.89	2.78	5.01	4.12	3.18	3.36	3.21	2.80	1.78	1.03	31.23
Hennessey	0.71	1.16	1.86	2.38	5.32	3.90	2.51	2.69	3.39	2.11	1.63	0.99	28.65
Kingfisher	0.83	1.13	1.76	2.42	4.94	3.76	2.57	2.39	3.60	2.44	1.53	1.13	28.50
Marshall	0.76	1.16	1.99	2.38	5.25	4.00	2.59	2.75	3.51	2.60	1.63	1.14	29.76
Newkirk	0.80	1.10	1.98	2.95	4.72	4.59	3.55	3.50	3.54	2.77	1.94	1.22	32.72
Ponca Cit	0.91	1.22	2.10	2.90	4.49	4.17	4.10	3.36	3.84	2.60	2.05	1.27	33.01
Perry	0.87	1.32	2.36	2.70	5.28	4.13	3.53	3.33	3.74	2.63	1.80	1.20	32.89
Guthrie	0.91	1.26	2.01	2.60	5.42	3.96	2.84	2.38	3.98	2.66	1.80	1.20	31.02
Ok. City	0.96	1.29	2.07	2.91	5.50	3.87	3.04	2.40	3.41	2.71	1.53	1.20	30.89
Ralston	1.00	1.30	2.52	2.97	4.72	4.39	3.49	2.92	3.86	2.69	1.95	1.36	33.17
Pawnee	1.01	1.31	2.48	2.97	4.84	4.02	3.13	3.01	4.37	2.72	1.88	1.25	32.99
Tulsa	1.35	1.74	3.14	4.15	5.14	4.57	3.51	3.01	4.37	3.41	2.56	1.82	38.77
Average	0.83	1.17	2.07	2.76	4.77	4.02	3.11	2.86	3.44	2.45	1.72	1.12	30.35

Precipitation – Mutual Ok.

Precipitation—Alva Ok.

Precipitation – Cherokee Ok.

Precipitation – Helena Ok.

Precipitation—Jefferson Ok.

Precipitation—Enid Ok.

Precipitation—Hennessey Ok.

Precipitation—Kingfisher Ok.

Precipitation—Marshall Ok.

North Central Oklahoma

Precipitation—Newkirk Ok.

Precipitation—Ponca City Ok.

Precipitation—Perry Ok.

Precipitation – Guthrie Ok.

Precipitation—Oklahoma City Ok.

Precipitation—Ralston Ok.

Precipitation—Pawnee Ok.

Precipitation—Tulsa Ok.

Average Precipitation

North Central Oklahoma

AVERAGE ANNUAL PRECIPITATION

Oklahoma Cloudiness

Oklahoma City

Oklahoma Cloudiness

Tulsa

North Central Oklahoma Data													
Thunderstorm days													
Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Ponca City	0.4	1.0	2.4	5.4	8.7	9.8	8.1	8.0	4.4	3.5	0.8	0.5	53.0
Tulsa	1.0	2.0	3.0	6.0	10.0	11.0	7.0	7.0	5.0	4.0	1.0	1.0	58.0
Enid	0.3	0.9	2.0	5.4	8.9	9.8	7.1	6.9	4.7	3.2	0.5	0.5	50.2
Seminole	0.7	1.3	2.8	5.2	9.4	7.2	7.2	4.5	2.6	2.1	0.6	0.6	44.2
Ok City	1.0	1.0	3.0	5.0	7.0	8.0	5.0	6.0	4.0	3.0	1.0	1.0	45.0
Alva	0.3	0.9	2.0	5.4	8.9	9.8	7.1	6.9	4.7	3.2	0.5	0.5	50.2
Averages	0.6	1.2	2.5	5.4	8.8	9.3	6.9	6.6	4.2	3.2	0.7	0.7	50.1

Thunderstorm Days

North Central Oklahoma

Average Thunderstorm Days

North Central Oklahoma

AVERAGE ANNUAL THUNDERSTORM DAYS

Thunderstorm Days

Location	Thunderstorm Days
1. Mutual	50.2
2. Alva	
3. Cherokee	
4. Helena	
5. Jefferson	50.2
6. Enid	
7. Hennessey	
8. Kingfisher	
9. Marshall	
10. Newkirk	
11. Ponca City	53.0
12. Perry	
13. Guthrie	
14. Oklahoma City	45.0
15. Ralston	
16. Pawnee	
17. Tulsa	58.0
* Lamont	44.2
Seminole	

North Central Oklahoma Data
Average Snowfall (inches)

Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
Ponca City	3.6	2.4	2.1	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.4	10.6
Tulsa	3.6	2.1	1.3	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.4	8.8
Enid	3.7	3.3	1.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.1	10.4
Seminole	2.0	1.8	1.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	6.3
Ok. City	3.5	2.3	1.8	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.7	9.7
Alva	3.5	4.8	3.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	1.1	3.3	16.1
Averages	3.3	2.8	1.9	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.7	10.3

Snowfall

North Central Oklahoma

Average Snowfall

North Central Oklahoma

AVERAGE ANNUAL SNOWFALL

Average annual percentage frequencies of wind direction, 1951-1960. Areas of sectors are proportional to hourly frequencies. (From SLUSSER, 1965.)

Average percentage frequencies of wind direction in January, 1951-1960. (From Slusser, 1965)

Average percentage frequencies of wind directions in: A. January, and B. July, 1951-1960. Areas of sectors are proportional to hourly frequencies. (From SLUSSER, 1965.)

Average annual wind speeds, 1951-1960. Areas of sectors are proportional to hourly speeds. (From SLUSSER, 1965.)

Mean annual number of days with fog (visibility less than 300 m), from records of 251 stations through 1960. (From COURT and GERSTON, 1966.)

Mean annual moisture content of atmosphere up to 325 mbar over 50 stations, 1946-1956, in millimeters of condensed water, or decigrams per square centimeter of earth's surface. (After REITAN, 1960a.)

References

1. World WeatherDisc. Version 1.0, Sept. 1989.
2. World Survey of Climatology. H. Landsberg, Vol. 11
Elsevier Scientific Publishing Co., 1974.
3. Climates of the States. Vol. 2. NOAA 1974.
4. Solar Radiation Energy Resources Atlas of the
United States. SERI/SP-642-1037, U. S. Dept.
of Energy. 1981.
5. Hammond's Illustrated Library World Atlas. C.S.
Hammond & Co. Inc., 1947.
6. Slusser, W.F., 1965. Wind rose maps of the United
States. Weatherwise, 18: 260-263.
7. Bonner, W.D., 1968. Climatology of the low-level
jet. Mon. Weather Rev., 99: 833-850.
8. Palmer, W.C., 1965. Meteorological drought. U.S.
Weather Bur., Res. Pap., 45: vi + 68pp.
9. Court, A. and Gerston, R.D., 1966. Fog frequency
in the United States. Geogr. Rev., 56: 543-550.
10. Reitan, C.H., 1960. Mean monthly values of
precipitable water over the United States,
1946-1956, Mon. Weather Rev., 88: 25-35.

END

**DATE
FILMED**
5/25/93