DOE/ER/40272--133

DE92 008775

A Higher Derivative View of the
Arc Length and Area Actions

D. Zoller

Physics Department MAR 11 1992
University of Florida
Gainesville, FL 32611

Abstract: Higher derivative versions of the arc length and area actions are presented. The higher derivative theories are equivalent with the corresponding lower derivative theories in absence of interactions. The extra degrees of freedom associated with the higher derivatives are pure gauge due to a hidden reparametrization invariance these theories possess. The extra degrees of freedom are no longer pure gauge in the presence of background fields and the equivalence no longer holds. A string version of Wigner's continuous spin particle is also noted.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

The arc length and area actions arise in many investigations of particle physics and statistical mechanics. The arc length action provides a model of the simplest relativistic point particle. The area action provides a model of flux tubes in QCD and can also describe gravitons. Such actions also arise in the context of statistical mechanics. The 2-D Ising model can be reformulated by considering the boundary of a region of like spins. The energy of such a configuration is proportional to the arc length of the boundary due to the nearest neighbor nature of the interaction. Such a picture can be fully developed for the 2-D Ising model¹ and similar ideas may apply to the 3-D Ising model.

The arc length and area actions are thus of general importance and here we present equivalent higher derivative reformulations of them. The extrema of such actions give higher order differential equations and so imply extra degrees of freedom.* The free higher derivative theories possess a hidden reparametrization invariance which implies the extra degrees of freedom can be gauged away. Gauge fixing the extra degrees of freedom leaves a theory equivalent to the usual arc length or area action. If background gravitational or electromagnetic fields are introduced, the extra degrees of freedom are not pure gauge and the higher derivative theories are no longer equivalent to the corresponding lower derivative theories.

The key to the reformulation of the arc length and area actions lies in the existence of a hidden reparametrization invariance present in some higher derivative theories. There are actions that are not invariant in the usual sense but still possess a non-standard reparametrization invariance. The simplest such theory is

$$S = m \int d\tau \sqrt{\ddot{x}^2} \quad (1)$$

* These extra degrees of freedom correspond to extra dimensions much as in a Kaluza-Klein theory. The extra dimensions in a Kaluza-Klein theory are unobservable at distance scales large compared to the size of the extra dimensions but do appear at small distance scales. By contrast, the extra dimensions that arise from the free higher derivative theories discussed here are non-dynamical and unobservable at any distance scale.

where τ is an arbitrary evolution parameter and $\dot{x}$ means the derivative of x with respect to τ . The Minkowski metric has signature $- - - +$. Although this action is not invariant under the usual reparametrization

$$x(\tau) \rightarrow x(f(\tau)) \quad (2)$$

it is invariant under the non-local gauge transformation

$$x(\tau) \rightarrow \int^{\tau} d\tau \dot{x}(f(\tau)) \quad (3.)$$

The term hidden* reparametrization invariance is used as the representation of the reparametrization group defined by (3) is different from the standard one defined by (2). In terms of $\dot{x}$ this gauge transformation is local

$$\dot{x}(\tau) \rightarrow \dot{x}(f(\tau)) \quad (4)$$

The gauge invariance reduces the number of degrees of freedom of the fourth order differential equation arising from S . We shall now show that the action S gives the same theory as the arc length action.

We shall recast the theory in Hamiltonian form to easily discuss the representation of the Poincaré group of the constrained system. Ostrogradsky first formulated higher derivative theories in Hamiltonian form³ and the generalized canonical coordinates are

$$\begin{aligned} q' &= \dot{q} \\ p' &= \frac{\partial L}{\partial \ddot{q}} \\ p &= \frac{\partial L}{\partial \dot{q}} - \frac{d}{dt} \frac{\partial L}{\partial \ddot{q}} \end{aligned} \quad (5)$$

where the basic dynamical variable is q . The Hamiltonian is

$$H = pq' + p'\dot{q}' - L \quad (6)$$

* The term hidden reparametrization invariance in Ref. 2 was used also to imply the invariance was broken.

and p, q and p', q' are canonically conjugate phase space variables. The Poisson brackets are

$$\begin{aligned} [p, q] &= 1 \\ [p', q'] &= 1 \end{aligned} \tag{7.}$$

The phase space is doubled as is appropriate for a fourth order differential equation.

The Hamiltonian of the theory S is

$$H = P \cdot X' \tag{8}$$

where the dynamical variable q is now denoted as X . The constraints are

$$\begin{aligned} P'^2 &= m^2 \\ P' \cdot P &= 0 \\ P^2 &= 0. \end{aligned} \tag{9}$$

The first constraint is primary, the second secondary, and the third tertiary. The first constraint implies P' is timelike and so there is a frame where $P' = (m, 0, \dots, 0)$. Here the dimension d of spacetime is arbitrary. The second constraint then implies the time component of P is zero. The third gives that P^μ is identically zero. The three constraints are thus equivalent to the $d + 1$ constraints

$$\begin{aligned} P'^2 &= m^2 \\ P^\mu &= 0 \end{aligned} \tag{10}$$

and the Hamiltonian H vanishes on the constraint surface. The constraint $P^\mu = 0$ can be simply eliminated by fixing the gauge

$$X^\mu = 0 \tag{11}$$

and this gauge fixed theory is clearly equivalent to the arc length theory.

The Nambu area action can also be reformulated as a higher derivative theory. The higher derivative action

$$S = \int d\sigma d\tau \sqrt{(\ddot{x} \cdot \dot{x}')^2 - \dot{x}^2 \dot{x}'^2} \quad (12)$$

arises by replacing x with $\dot{x}$ in the usual area action with string tension set to one. Again S is invariant under reparametrizations of $\dot{x}$ which in terms of x is the non-local gauge transformation

$$x(\sigma, \tau) \rightarrow \int^\tau d\tau' \dot{x}(\sigma'(\sigma, \tau), \tau'(\sigma, \tau)) \quad (13)$$

To establish the equivalence of S to the Nambu area action we shall formulate it in Hamiltonian terms.

The Hamiltonian of S is

$$H = \int d\sigma P \cdot \dot{X} \quad (14)$$

Here $\dot{X}$ corresponds to q' and is not the time derivative of X . We have departed from the earlier notation to maintain the usual string notation where X' means the derivative of X with respect to σ . The canonically conjugate phase space variables are P, X and $P', \dot{X}$ and have the Poisson brackets

$$[P(\sigma), X(\sigma')] = \delta(\sigma - \sigma')$$

$$[P'(\sigma), (\sigma')] = \delta(\sigma - \sigma')$$

The constraints are

$$P'^2 + \dot{X}'^2 = 0$$

$$P' \cdot \dot{X}' = 0$$

$$P \cdot P' = 0 \quad (15)$$

$$P \cdot \dot{X}' = 0$$

$$P^2 = 0.$$

The first two constraints are primary, the second two secondary, and the third tertiary. The timelike character of P' again implies that $P^\mu = 0$. Fixing the gauge $X^\mu(\sigma) = 0$ eliminates the constraint $P^\mu = 0$ and we are left with the constraints

$$\begin{aligned} P'^2 + \dot{X}'^2 &= 0 \\ P' \cdot \dot{X}' &= 0 \end{aligned} \tag{16}$$

which are equivalent to those of the Nambu area action.

The equivalence of these higher derivative theories with the corresponding ordinary derivative theories no longer holds in the presence of background gravitational or electromagnetic interactions. Consider the point particle coupled to a background electromagnetic field whose vector potential is A_μ . The natural coupling that arises from the replacement of x by $\dot{x}$ is

$$L = m\sqrt{\ddot{x}} + A_\mu(\dot{x})\ddot{x}^\mu \tag{17}$$

The difficulty is that the primary constraint

$$(P' - A)^2 = m^2 \tag{18}$$

and the secondary constraint

$$P \cdot (P' - A) = 0 \tag{19}$$

no longer commute. The commutator is proportional to

$$P^\mu (P' - A)^\nu F_{\mu\nu} \tag{20}$$

where $F_{\mu\nu}$ is the field strength associated with A_μ . The dynamical system with interactions possesses two second class constraints instead of three first class constraints. The theory based on L is no longer equivalent to the arc length theory as it possesses more degrees

of freedom. Similar arguments apply to the arc length theory coupled to background gravitational fields and to the Nambu area action coupled to background fields.

The arclength and area actions also come in a spacelike variety in addition to the time-like variety already discussed. The higher derivative spacelike varieties arise by replacing the argument of the square root with its negative

$$L = m\sqrt{-\dot{x}^2}$$

$$L = \sqrt{\dot{x}^2\dot{x}'^2 - (\ddot{x} \cdot \dot{x}')^2} \quad (21)$$

These theories, however, are not equivalent to the corresponding lower derivative versions. The vector P^μ is no longer timelike and the argument that $P^\mu = 0$ no longer holds. The theories thus possess extra degrees of freedom and the corresponding Hamiltonians no longer vanish. The higher derivative spacelike arc length theory restricted to a particular value of the Hamiltonian gives rise to Wigner's continuous spin representation² of the Poincaré group. The area action version of this theory gives a string version of Wigner's continuous spin particle. The Hamiltonian constraints are given by (15) with the $+$ sign in the first constraint replaced by a $-$ sign.

Higher derivative theories have not been systematically studied and may reveal some surprises. In this note we have pointed out that the higher derivative versions of the arc length and area actions do not possess the extra degrees of freedom typical of higher derivative theories. The hidden reparametrization invariance in combination with the causal character of the Lagrangian implied the expected extra degrees of freedom were pure gauge. Higher derivative actions have been suggested⁴ to play a role in the string ansatz to the loop equation arising from the $\frac{1}{N}$ approximation to QCD . The hidden reparametrization invariance and the consequent reduction in degrees of freedom may be useful⁵ in this context.

REFERENCES

1. A. Polyakov, Les Houches lectures(1989).
2. D. Zoller, preprint University of Florida, Gainesville UFIFT 91-7.
3. E. Whittaker, A Treatise on the Analytic Dynamics of Paticles and Rigid Bodies (Cambridge University Press, 1988) p. 265.
4. A. Polyakov, *Nucl. Phys. B* **268**, 406 (1986).
5. D. Zoller, preprint University of Florida, Gainesville UFIFT, in preparation.

END

DATE
FILMED

4/10/92